



# Early Syphilis in the Buffalo Region, New York State


**Age-Adjusted Rate:  
Syphilis Cases per 100,000 Residents, 2014**



**Number of Cases in 2014  
Compared to 2010-2013 average**



**New Infections by Year**



76% of New Yorkers support condom availability in High Schools

85% of syphilis cases are among men who have sex with men


Syphilis increases risk of HIV transmission two to five fold

PrEP does not protect from STDs  
Condoms and screening are important


Sources: NYSDOH Bureau of STD Prevention and Epidemiology STDC@health.ny.gov and 2014 Behavioral Risk Factor Surveillance System

# Early Syphilis in the Rochester Region, New York State


**Age-Adjusted Rate:  
Syphilis Cases per 100,000 Residents, 2014**



**Number of Cases in 2014  
Compared to 2010-2013 average**



**New Infections by Year**



76% of New Yorkers support condom availability in High Schools

85% of syphilis cases are among men who have sex with men



Syphilis increases risk of HIV transmission two to five fold

PrEP does not protect from STDs  
Condoms and screening are important


Sources: NYSDOH Bureau of STD Prevention and Epidemiology STDC@health.ny.gov and 2014 Behavioral Risk Factor Surveillance System

# Early Syphilis in Central New York


**Age-Adjusted Rate:  
Syphilis Cases per 100,000 Residents, 2014**



**Number of Cases in 2014  
Compared to 2010-2013 average**



**New Infections by Year**



76% of New Yorkers support condom availability in High Schools

85% of syphilis cases are among men who have sex with men



Syphilis increases risk of HIV transmission two to five fold

PrEP does not protect from STDs  
Condoms and screening are important


Sources: NYSDOH Bureau of STD Prevention and Epidemiology STDC@health.ny.gov and 2014 Behavioral Risk Factor Surveillance System

# Early Syphilis in the Albany Region, New York State


**Age-Adjusted Rate:  
Syphilis Cases per 100,000 Residents, 2014**



**Number of Cases in 2014  
Compared to 2010-2013 average**



**New Infections by Year**



76% of New Yorkers support condom availability in High Schools

85% of syphilis cases are among men who have sex with men



Syphilis increases risk of HIV transmission two to five fold

PrEP does not protect from STDs  
Condoms and screening are important


Sources: NYSDOH Bureau of STD Prevention and Epidemiology STDC@health.ny.gov and 2014 Behavioral Risk Factor Surveillance System

# Early Syphilis in the Hudson Valley, New York State


**Age-Adjusted Rate:  
Syphilis Cases per 100,000 Residents, 2014**



**Number of Cases in 2014  
Compared to 2010-2013 average**



**New Infections by Year**



76% of New Yorkers support condom availability in High Schools


85% of syphilis cases are among men who have sex with men

Syphilis increases risk of HIV transmission two to five fold


PrEP does not protect from STDs  
Condoms and screening are important

# Primary & Secondary Syphilis in New York City (NYC)

**Primary & Secondary Syphilis Case Rates  
(per 100,000 Population)  
Overall and by Borough of Residence, 2014**




**Primary & Secondary Syphilis Cases by Sex, NYC, 2000-2014**




**United Hospital Fund (UHF) Neighborhoods  
New York City**

- NYC Boroughs
- Brooklyn
- Manhattan
- Queens
- Richmond




**P&S Syphilis Among Men in NYC  
Reported 01/01/2014 - 12/31/2014**

- Rate / per 100,000
- 2 - 29
- 30 - 85
- 86 - 171




**P&S Syphilis Among Women in NYC  
Reported 01/01/2014 - 12/31/2014**

- Rate / per 100,000
- 0
- 1
- 2 - 3





# Early Syphilis in Nassau and Suffolk Counties, New York State


**Age-Adjusted Rate:  
Syphilis Cases per 100,000 Residents, 2014**



**Number of Cases in 2014  
Compared to 2010-2013 average**



**New Infections by Year**



76% of New Yorkers support condom availability in High Schools

85% of syphilis cases are among men who have sex with men

Syphilis increases risk of HIV transmission two to five fold

PrEP does not protect from STDs  
Condoms and screening are important

Sources: NYSDOH Bureau of STD Prevention and Epidemiology STDC@health.ny.gov and 2014 Behavioral Risk Factor Surveillance System