

## R.I.C.H. Report [Regional Information on Community HIV]

New York State HIV/AIDS 2013-2022





# 4

Highlights:

34% decrease in new HIV diagnoses since 2013 46% decrease in new AIDS diagnoses since 2013

## HIV Statistics 2022 104

104,124


individuals currently living with HIV

2,318


persons newly diagnosed with HIV

2,036

deaths among individuals with HIV



The percentage of males<sup>a</sup> newly diagnosed with HIV increased 1% since 2013. (80% to 81%)



# Non-Hispanic Black 941 Hispanic 830 Non-Hispanic White 430 Asian 93 Multi Race 19 Native American 3 HI/PI<sup>b</sup> 2

New HIV diagnoses among Non-Hispanic Black
individuals have decreased by 35% since 2013 but
continue to be the most affected sub-population.

Age at HIV diagnosis	People newly diagnosed with HIV
<u>0-19</u>	78
<u>20-29</u>	786
<u>30-39</u>	728
<u>40-49</u>	353
<u>50-59</u>	228
<u>60+</u>	145

The largest group of individuals newly diagnosed with HIV every year since 2013 was 20-29 year olds.



81% of people were linked to care within 30 days of diagnosis.



58% of people newly diagnosed with HIV reached viral load suppression in 3 months.



# HIV Statistics 2022 81,

81,079

individuals currently living with HIV

Highlights:


37% decrease in new HIV diagnoses since 2013 49% decrease in new AIDS diagnoses since 2013



persons newly diagnosed with HIV

1,568

deaths among individuals with HIV




newly diagnosed with HIV increased 1% since 2013. (80% to 81%)

The number of people with male to male sexual contact (MSM) newly diagnosed with HIV has decreased by 45% since 2013, but remains the most affected subpopulation, accounting for 53% of new diagnoses in 2022.

7 14 13

Pediatric Injection Drug MSM and IDU Use (IDU)



#### Non-Hispanic Black 746

Hispanic 676

Non-Hispanic White 199

Asian 73

Multi Race 15

HI/PIb 2

Native American 2

New HIV diagnoses among Non-Hispanic Black individuals have decreased by 35% since 2013 but continue to be the most affected sub-population.

Age at HIV diagnosis	People newly diagnosed with HIV
<u>0-19</u>	56
<u>20-29</u>	570
<u>30-39</u>	552
<u>40-49</u>	270
<u>50-59</u>	161
<u>60+</u>	104

The largest group of individuals newly diagnosed with HIV every year since 2013 was 20-29 year olds.



80% of people were linked to care within 30 days of diagnosis.



57% of people newly diagnosed with HIV reached viral load suppression in 3 months.



#### **HIV Statistics** 2022

23,045

individuals currently living with HIV


#### Highlights:

24% decrease in new HIV diagnoses since 2013 37% decrease in new AIDS diagnoses since 2013




persons newly diagnosed with HIV 468

deaths among individuals with HIV



The percentage of males<sup>a</sup> newly diagnosed with HIV increased 2% since 2013. (80% to 82%)



\*Groups with zero diagnoses are not displayed in this chart

Non-Hispanic	White	231

Non-Hispanic Black 195

Hispanic 154

Asian 20

Multi Race 4

Native American 1

HI/PIb 0

Since 2013, the number of new HIV diagnoses amongst
Non-Hispanic White and Non-Hispanic Black
individuals have been similar, with both decreasing
(20% and 36% respectively) into 2022.

Age at HIV diagnosis	People newly diagnosed with HIV
<u>0-19</u>	22
<u>20-29</u>	216
<u>30-39</u>	176
<u>40-49</u>	83
<u>50-59</u>	67
60+	41

The largest group of individuals newly diagnosed with HIV every year since 2013 was 20-29 year olds.



82% of people were linked to care within 30 days of diagnosis.



60% of people newly diagnosed with HIV reached viral load suppression in 3 months.

Source: NYS HIV Registry, as of June 2023 aSex at Birth bHI/PI-Native Hawaiian/Pacific Islander ROS: NYS excluding NYC



### Albany

HIV Statistics

2022

2,987

individuals currently living with HIV

Non-Hispanic White 40



#### Highlights:


4% increase in new HIV diagnoses since 2013 33% decrease in new AIDS diagnoses since 2013


71

persons newly diagnosed with HIV

68

deaths among individuals with HIV





\*Groups with zero diagnoses are not displayed in this chart

1101-111spainte White 40		
Non-Hispanic Black 17		
Hispanic 8		
Asian 4		
Multi-Race 2		
Native American 0		
HI/PI <sup>b</sup> 0		

New HIV diagnoses among Non-Hispanic White individuals have increased by 21% since 2013 and continue to be the most affected sub-population.

Age at HIV diagnosis	People newly diagnosed with HIV
<u>0-19</u>	2
<u>20-29</u>	26
<u>30-39</u>	28
<u>40-49</u>	7
<u>50-59</u>	6
<u>60+</u>	2

Individuals diagnosed between ages 30-39 have been increasing since 2018 and have remainsed the highest group of all age groups.



83% of people were linked to care within 30 days of diagnosis.



61% of people newly diagnosed with HIV reached viral load suppression in 3 months.

Source: NYS HIV Registry, as of June 2023. Excludes individuals who are incarcerated aSex at Birth

bHI/PI-Native Hawaiian/Pacific Islander



# Binghamton HIV Statistics 2022



#### Highlights:

31% increase in new HIV diagnoses since 2013 58% decrease in new AIDS diagnoses since 2013

512


individuals currently living with HIV

**17** 

persons newly diagnosed with HIV


17

deaths among individuals with HIV



increased 30% since 2013.

(23% to 53%)



\*Groups with zero diagnoses are not displayed in this chart

	Non-Hispanic White 10
Non-Hispanic Bla	ck 5

Hispanic 2

Multi-Race 0

Asian 0

Native American 0

HI/PIb 0

New HIV diagnoses among Non-Hispanic White individuals have increased by 43% since 2013 and continue to be the most affected sub-population.

Age at HIV diagnosis	People newly diagnosed with HIV
<u>0-19</u>	0
<u>20-29</u>	4
<u>30-39</u>	8
<u>40-49</u>	3
<u>50-59</u>	2
<u>60+</u>	0

The largest group of individuals newly diagnosed with HIV was 30-39 year olds in 2022.



94% of people were linked to care within 30 days of diagnosis.



35% of people newly diagnosed with HIV reached viral load suppression in 3 months.

Source: NYS HIV Registry, as of June 2023. Excludes individuals who are incarcerated aSex at Birth

bHI/PI-Native Hawaiian/Pacific Islander (Includes counties of: Broome, Chenango, Tioga)



2022

#### Buffalo HIV Statistics

Highlights:

39% decrease in new HIV diagnoses since 2013 22% decrease in new AIDS diagnoses since 2013

2,923


individuals currently living with HIV


74


persons newly diagnosed with HIV

68

deaths among individuals with HIV







Age at HIV diagnosis	People newly diagnosed with HIV
<u>0-19</u>	5
<u>20-29</u>	33
<u>30-39</u>	18
<u>40-49</u>	8
<u>50-59</u>	6
<u>60+</u>	4

The largest group of individuals newly diagnosed with HIV every year since 2013 was 20-29 year olds.



91% of people were linked to care within 30 days of diagnosis.



66% of people newly diagnosed with HIV reached viral load suppression in 3 months.

New HIV diagnoses among Non-Hispanic Black individuals have decreased by 27% since 2013 but continue to be the most affected sub-population.

Source: NYS HIV Registry, as of June 2023. Excludes individuals who are incarcerated a Sex at Birth

bHI/PI-Native Hawaiian/Pacific Islander

(Includes counties of: Allegany, Cattaraugus, Chautauqua, Erie, Genesee, Niagara, Orleans, Wyoming)

#### Lower Hudson 2

#### **HIV Statistics** 2022

3,574

individuals currently living with HIV


#### Highlights:

40% decrease in new HIV diagnoses since 2013 61% decrease in new AIDS diagnoses since 2013


**82** 

persons newly diagnosed with HIV 57

deaths among individuals with HIV



The percentage of males<sup>a</sup> newly diagnosed with HIV is the same as it was in 2013. (78% to 78%)



\*Groups with zero diagnoses are not displayed in this chart

#### Non-Hispanic Black 30

Non-Hispanic White 28

Hispanic 24

Native American 0

Asian 0

Multi Race 0

HI/PIb 0

New HIV diagnoses among Non-Hispanic Black individuals have decreased by 52% since 2013 but continue to be the most affected sub-population.

Age at HIV diagnosis	People newly diagnosed with HIV
<u>0-19</u>	2
<u>20-29</u>	26
<u>30-39</u>	26
<u>40-49</u>	13
<u>50-59</u>	5
<u>60+</u>	10

The largest group of individuals newly diagnosed with HIV every year since 2013 was 20-29 year olds. However, new HIV diagnoses among 30-39 year olds have increased and matched 20-29 year olds.



76% of people were linked to care within 30 days of diagnosis.



57% of people newly diagnosed with HIV reached viral load suppression in 3 months.

Source: NYS HIV Registry, as of June 2023. Excludes individuals who are incarcerated aSex at Birth

bHI/PI-Native Hawaiian/Pacific Islander

(Includes counties of: Putnam, Rockland, Westchester)



#### Mid-Hudson 4

#### **HIV Statistics** 2022

2,253

individuals currently living with HIV


#### Highlights:


35% decrease in new HIV diagnoses since 2013 22% decrease in new AIDS diagnoses since 2013



persons newly diagnosed with HIV **50** 

deaths among individuals with HIV





\*Groups with zero diagnoses are not displayed in this chart

	Non-Hispanic White 15	
Non-Hispanic Black 10		
	Hispanic 10	
Asian 1		
HI/PI <sup>b</sup> 0		
Multi Race 0		
Native America	n 0	

Since 2013, the number of new HIV diagnoses amongst
Non-Hispanic White and Non-Hispanic Black
individuals have been similar, with both decreasing
(32% and 41% respectively) into 2022.

Age at HIV diagnosis	People newly diagnosed with HIV
<u>0-19</u>	1
<u>20-29</u>	11
<u>30-39</u>	6
<u>40-49</u>	4
<u>50-59</u>	8
<u>60+</u>	6

The largest group of individuals newly diagnosed with HIV in 2022 was 20-29 year olds.



78% of people were linked to care within 30 days of diagnosis.



50% of people newly diagnosed with HIV reached viral load suppression in 3 months.

#### Nassau-Suffolk 2

## HIV Statistics 2022 5,4

5,474 individuals currently living with HIV

169

Highlights:


persons newly diagnosed with HIV

00


24% decrease in new HIV diagnoses since 2013

36% decrease in new AIDS diagnoses since 2013

deaths among individuals with HIV



The percentage of males<sup>a</sup> newly diagnosed with HIV increased 4% since 2013. (80% to 84%)



Hispanic 74

Non-Hispanic Black 46

Non-Hispanic White 44

Asian 5

Multi Race 0

Native American 0

HI/PIb 0

New HIV diagnoses among Hispanic individuals have
increased by 25% since 2013, however, this is a sharp
increase when compared to recent years.

Age at HIV diagnosis	People newly diagnosed with HIV
<u>0-19</u>	8
<u>20-29</u>	64
<u>30-39</u>	42
<u>40-49</u>	25
<u>50-59</u>	22
<u>60+</u>	8

The largest group of individuals newly diagnosed with HIV every year since 2013 was 20-29 year olds.



78% of people were linked to care within 30 days of diagnosis.



60% of people newly diagnosed with HIV reached viral load suppression in 3 months.

Source: NYS HIV Registry, as of June 2023. Excludes individuals who are incarcerated a Sex at Birth

<sup>b</sup>HI/PI-Native Hawaiian/Pacific Islander (Includes counties of: Nassau, Suffolk)



# Rochester HIV Statistics 2022

Highlights:

5% increase in new HIV diagnoses since 2013 37% decrease in new AIDS diagnoses since 2013

2,855


individuals currently living with HIV

**79** 

persons newly diagnosed with HIV


**78** 


deaths among individuals with HIV



increased 1% since 2013.

(15% to 16%)





Age at HIV diagnosis	People newly diagnosed with HIV
<u>0-19</u>	4
<u>20-29</u>	25
<u>30-39</u>	27
<u>40-49</u>	12
<u>50-59</u>	6
<u>60+</u>	5

The largest group of individuals newly diagnosed with HIV was 30-39 year olds in 2022.



91% of people were linked to care within 30 days of diagnosis.



66% of people newly diagnosed with HIV reached viral load suppression in 3 months.

Source: NYS HIV Registry, as of June 2023. Excludes individuals who are incarcerated a Sex at Birth

bHI/PI-Native Hawaiian/Pacific Islander

(Includes counties of: Chemung, Livingston, Monroe, Ontario, Schuyler, Seneca, Steuben, Wayne, Yates)

individuals have decreased by 24% since 2013 but continue to be the most affected sub-population.

# Syracuse HIV Statistics 2022 2,1

Highlights:

11% decrease in new HIV diagnoses since 2013 14% decrease in new AIDS diagnoses since 2013

2,188


individuals currently living with HIV


**65** 

persons newly diagnosed with HIV

42

deaths among individuals with HIV





Non-Hispanic White 38

Non-Hispanic Black 18

Hispanic 7

Asian 1

Native American 1

Multi-Race 0

HI/PIb 0

New HIV diagnoses among Non-Hispanic White individuals have increased by 31% since 2013 and continue to be the most affected sub-population.

Age at HIV diagnosis	People newly diagnosed with HIV
<u>0-19</u>	0
<u>20-29</u>	22
<u>30-39</u>	18
<u>40-49</u>	9
<u>50-59</u>	10
<u>60+</u>	6

The largest group of individuals newly diagnosed with HIV every year since 2013 was 20-29 year olds.




77% of people were linked to care within 30 days of diagnosis.



65% of people newly diagnosed with HIV reached viral load suppression in 3 months.

Source: NYS HIV Registry, as of June 2023. Excludes individuals who are incarcerated "Sex at Birth





#### **Contact information:**

Data Analysis and Research Translation Division of Epidemiology, Evaluation, and Partner Services P.O. Box 2073, ESP Station Albany NY 12220-0073

Phone: (518) 474-4284

E-mail: bhae@health.ny.gov

#### Link to the general HIV/AIDS data and reports:

https://www.health.ny.gov/diseases/aids/general/statistics/