Expanded BRFSS, 2003

Allegany and Wyoming Locality

Table of Contents:	
Optional Module 1: Adult Asthma History	
•	
Optional Module 4: Cholesterol Awareness	4
Optional Module 5: Diabetes	<i>6</i>
Optional Module 9: Fruits and Vegetables	10
Optional Module 10: Health Care Coverage	11
Optional Module 12: Hypertension Awareness	
Optional Module 14: Immunization	14
Optional Module 15: Oral Health	18
Optional Module 17: Skin Cancer	20
Optional Module 19: Tobacco Consumption	22
Optional Module 23: Seat Belts	
Ontional Module 24: Weight Control	25

New York State Department of Health

Have had an Asthma Episode or Attack in the Past 12 Months, among Those with Asthma

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	28	53.1	22	46.9	16.6

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

This page left blank intentionally

This page left blank intentionally

Ever had Blood Cholesterol Checked

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	472	73.3	133	26.7	4.1
Gender					
Male	164	68.2	68	31.8	6.6
Female	308	78.7	65	21.3	4.9
Age					
18-34	64	50.6	64	49.4	9.2
35-54	195	76.6	56	23.4	5.8
≥55	206	93.4	12	6.6	4.0
Education					
≤High School	240	70.6	78	29.4	5.9
>High School	231	76.4	55	23.6	5.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent ³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health Had Blood Cholesterol Checked in the Past Five Years *Allegany and Wyoming*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	435	69.3	158	30.7	4.3
Gender					
Male	150	64.1	78	35.9	6.8
Female	285	74.8	80	25.2	5.1
Age					
18-34	59	47.2	65	52.8	9.3
35-54	177	71.3	70	28.7	6.2
≥55	193	89.7	21	10.3	4.7
Education					
≤High School	223	67.9	85	32.1	6.0
>High School	211	70.8	73	29.2	6.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Check Blood for Glucose or Sugar at Least Once per Day among Those with Diabetes *Allegany and Wyoming*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	29	55.9	19	44.1	15.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

Check Feet for Sores or Irritations at Least Once per Day among Those with Diabetes *Allegany and Wyoming*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	31	59.3	18	40.7	15.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Had Feet Checked by a Medical Professional in Past Twelve Months, among Those with Diabetes

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	34	68.0	15	32.0	14.4

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Had Dilated Eye Exam in Past 12 Months, among Those with Diabetes *Allegany and Wyoming*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	37	75.2	11	24.8	13.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent ³95% Confidence Interval

Eat Five Fruits or Vegetable Servings per Day

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	155	22.4	465	77.6	3.6
Gender					
Male	36	13.6	206	86.4	4.6
Female	119	31.8	259	68.2	5.3
Age					
18-34	25	18.9	108	81.1	7.2
35-54	55	19.4	201	80.6	5.2
≥55	74	30.2	149	69.8	6.5
Education					
≤High School	58	14.8	270	85.2	4.1
>High School	96	31.1	195	68.9	6.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Main Reason for Not Having Health Care Coverage

Demographic Groups	n¹	Changed or lost job	C.I. ³	n	Could not afford	C.I.	n	Other %	C.I.
Total	12	22.1	12.7	27	44.4	14.5	21	33.5	13.6

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

No Health Care Coverage for Past Two Years Among Uninsured Allegany and Wyoming

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	25	44.3	33	55.7	14.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent ³95% Confidence Interval

Ever Been Told Have High Blood Pressure by a Medical Professional *Allegany and Wyoming*

Demographic	1	Yes		No	CI3
Groups	n¹	% ²	n	%	C.I. ³
Total	180	25.3	434	74.7	3.7
Gender					
Male	71	26.5	168	73.5	5.9
Female	109	24.0	266	76.0	4.5
Age					
18-34	8	5.5	123	94.5	3.9
35-54	58	23.5	197	76.5	6.1
≥55	110	48.9	110	51.1	7.3
Education					
≤High School	114	30.0	212	70.0	5.5
>High School	65	19.5	222	80.5	4.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Had a Flu Shot in the Past 12 Months, Aged 65 and Older *Allegany and Wyoming*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	97	73.5	32	26.5	8.6

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent ³95% Confidence Interval

Had a Flushot in the Past 12 Months

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	229	33.3	385	66.7	4.1
Gender					
Male	73	28.2	165	71.8	6.1
Female	156	38.7	220	61.3	5.4
Age					
18-64	126	24.4	351	75.6	4.3
≥65	97	73.5	32	26.5	8.6
Education					
≤High School	113	29.5	212	70.5	5.4
>High School	115	37.5	173	62.5	6.3

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent ³95% Confidence Interval

Ever Had a Pneumonia Shot, Aged 65 and Older

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	92	70.9	36	29.1	8.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent ³95% Confidence Interval

Ever Had a Pneumonia Shot

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	166	25.3	422	74.7	3.9
Gender					
Male	53	22.2	170	77.8	5.9
Female	113	28.5	252	71.5	5.0
Age					
18-64	70	15.0	383	85.0	3.8
≥65	92	70.9	36	29.1	8.7
Education					
≤High School	97	27.5	213	72.5	5.5
>High School	68	22.7	209	77.3	5.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent ³95% Confidence Interval

Have Had Permanent Teeth Removed because of Tooth Decay or Gum Disease *Allegany and Wyoming*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	335	51.2	268	48.8	4.6
Gender					
Male	127	50.6	108	49.4	7.0
Female	208	51.9	160	48.1	5.7
Age					
18-34	30	23.0	100	77.0	7.7
35-54	131	54.7	120	45.3	6.9
≥55	171	79.6	43	20.4	6.1
Education					
≤High School	219	64.0	100	36.0	6.2
>High School	116	36.2	167	63.8	6.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Having Seen a Dental Professional in the Past 12 Months

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	374	66.7	172	33.3	4.6
Gender					
Male	130	59.3	83	40.7	7.3
Female	244	74.6	89	25.4	5.1
Age					
18-34	77	61.9	49	38.1	9.0
35-54	156	62.4	81	37.6	7.2
≥55	135	78.3	40	21.7	6.7
Education					
≤High School	172	61.4	97	38.6	6.8
>High School	201	72.2	75	27.8	6.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

Had a Severe Sunburn Lasting 12 Hours in Past 12 Months *Allegany and Wyoming*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	249	46.4	364	53.6	4.5
Gender					
Male	119	54.2	121	45.8	6.8
Female	130	37.9	243	62.1	5.6
Age					
18-34	86	69.6	43	30.4	8.2
35-54	119	47.1	135	52.9	7.0
≥55	42	20.8	180	79.2	6.1
Education					
≤High School	117	42.8	208	57.2	6.3
>High School	132	50.6	155	49.4	6.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Number of Severe Sunburns Lasting 12 or More Hours in Past 12 Months *Allegany and Wyoming*

Demographic Groups	n¹	3 or more	C.I. ³	n	1-2 %	C.I.	n	None %	C.I.
Total	110	22.7	4.1	128	22.5	3.8	364	54.8	4.6
Gender									
Male	58	30.0	6.7	55	22.6	5.6	121	47.4	6.9
Female	52	14.8	4.3	73	22.5	4.9	243	62.7	5.6
Age									
18-64	106	26.9	4.9	115	25.3	4.4	244	47.8	5.1
≥65	4	4.0	4.2	11	8.5	5.3	114	87.5	6.5
Education									
≤High School	50	20.7	5.7	58	20.0	5.0	208	59.3	6.3
>High School	60	24.9	6.1	70	25.5	5.7	155	49.6	6.6

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

^{395%} Confidence Interval

Number of Cigarettes Smoked per day among Everyday Smokers *Allegany and Wyoming*

Demographic Groups	n¹	1-10 cigarettes %	n	≥11 cigarettes %	C.I. ³
Total	34	26.1	94	73.9	8.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

Use Seatbelt when Ride or Drive in a Car (Always, Nearly Always or Never Ride in a Car)

Demographic Groups	n¹	Yes %2	n	No %	C.I. ³
Total	566	91.1	46	8.9	2.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Use Seatbelt when Ride or Drive in a Car (Always or Never ride in a Car) *Allegany and Wyoming*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	493	78.0	119	22.0	3.9
Gender					
Male	169	70.1	70	29.9	6.3
Female	324	86.6	49	13.4	3.9
Age					
18-64	371	76.4	103	23.6	4.4
≥65	116	86.9	14	13.1	6.8
Education					
≤High School	254	75.6	69	24.4	5.5
>High School	238	80.8	50	19.2	5.3

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

^{395%} Confidence Interval

Trying to Lose Weight

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	259	40.7	354	59.3	4.4
Gender					
Male	74	30.2	165	69.8	6.2
Female	185	52.0	189	48.0	5.6
Age					
18-34	54	38.6	75	61.4	8.7
35-54	115	42.7	140	57.3	6.8
≥55	86	39.4	135	60.6	7.2
Education					
≤High School	129	37.8	195	62.2	5.9
>High School	130	44.0	158	56.0	6.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent ³95% Confidence Interval

Trying to Maintain Current Weight of Those not Trying to Lose Weight *Allegany and Wyoming*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	198	53.7	152	46.3	6.0
Gender					
Male	86	50.6	78	49.4	8.4
Female	112	58.7	74	41.3	7.9
Age					
18-34	33	42.7	41	57.3	12.0
35-54	81	57.7	59	42.3	9.4
≥55	82	61.0	51	39.0	9.2
Education					
≤High School	98	49.6	95	50.4	8.1
>High School	99	59.0	57	41.0	9.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

Eating Fewer Calories or Less Fat to Lose or Maintain Weight Allegany and Wyoming

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	332	72.0	120	28.0	4.8
Gender					
Male	97	61.9	60	38.1	8.3
Female	235	80.9	60	19.1	4.8
Age					
18-34	64	75.0	22	25.0	9.8
35-54	142	70.8	53	29.2	7.5
≥55	123	71.7	42	28.3	7.9
Education					
≤High School	170	73.1	54	26.9	6.9
>High School	162	71.0	65	29.0	6.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health Using Physical Exercise to Lose or Maintain Weight Allegany and Wyoming

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	311	71.2	142	28.8	4.6
Gender					
Male	104	66.6	55	33.4	7.8
Female	207	75.3	87	24.7	5.1
Age					
18-34	74	84.9	13	15.1	7.7
35-54	136	71.4	58	28.6	7.1
≥55	97	57.8	69	42.2	8.4
Education					
≤High School	153	69.7	70	30.3	6.8
>High School	158	72.8	71	27.2	6.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

^{395%} Confidence Interval