

Expanded BRFSS, 2003

Kings Locality

Table of Contents:

Optional Module 1: Adult Asthma History 1
Optional Module 5: Diabetes..... 2
Optional Module 7: Disability/Quality of Life..... 6

Optional Module 10: Health Care Coverage 10
Optional Module 13: Injury Control - Falls..... 12
Optional Module 19: Tobacco Consumption 13
Optional Module 20: Tobacco - Media 14
Optional Module 21: Tobacco – other products 22
Optional Module 22: Tobacco – Work site ETS 23
Optional Module 23: Seat Belts..... 25
Optional Module 24: Weight Control..... 27

New York State Department of Health

Expanded BRFSS, 2003 - New York State Department of Health

Have had an Asthma Episode or Attack in the Past 12 Months, among Those with Asthma

Kings

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	36	64.6	20	35.4	14.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Check Blood for Glucose or Sugar at Least Once per Day among Those with Diabetes

Kings

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	25	50.1	24	49.9	15.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Check Feet for Sores or Irritations at Least Once per Day among Those with Diabetes

Kings

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	25	51.1	23	48.9	15.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Had Feet Checked by a Medical Professional in Past Twelve Months, among Those with Diabetes

Kings

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	34	75.0	11	25.0	14.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Had Dilated Eye Exam in Past 12 Months, among Those with Diabetes

Kings

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	32	70.8	14	29.2	15.4

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Adults with a Disability

Kings

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	108	16.3	513	83.7	3.2
Gender					
Male	38	16.4	181	83.6	5.5
Female	70	16.2	332	83.8	3.8
Age					
18-34	14	7.7	195	92.3	4.7
35-54	40	15.5	203	84.5	5.1
≥55	51	27.4	110	72.6	7.5
Education					
≤High School	54	19.4	206	80.6	5.3
>High School	54	14.0	301	86.0	4.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Adults with a Disability Requiring Help

Kings

Demographic Groups	n ¹	Disability need help % ²	C.I. ³	n	Disability no help needed %	C.I.	n	No Disability %	C.I.
Total	39	5.3	1.8	69	11.0	2.8	513	83.7	3.2
Gender									
Male	8	2.9	2.1	30	13.5	5.2	181	83.6	5.5
Female	31	7.2	2.7	39	8.9	2.9	332	83.8	3.8
Age									
18-64	29	4.5	1.8	49	9.2	2.9	457	86.2	3.4
≥65	8	8.3	5.9	19	19.4	8.8	51	72.2	10.2
Education									
≤High School	20	5.8	2.7	34	13.6	4.8	206	80.6	5.3
>High School	19	4.9	2.4	35	9.0	3.3	301	86.0	4.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

This page left intentionally blank

This page left intentionally blank

Expanded BRFSS, 2003 - New York State Department of Health

Main Reason for Not Having Health Care Coverage

Kings

Demographic Groups	n ¹	Changed or lost job % ²	C.I. ³	n	Could not afford %	C.I.	n	Other %	C.I.
Total	10	10.7	6.9	22	27.5	11.1	50	61.8	11.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

No Health Care Coverage for Past Two Years Among Uninsured

Kings

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	30	35.6	55	64.4	11.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Had a Fall in Past Three Months

Kings

Demographic Groups	n ¹	No fall % ²	C.I. ³	n	Fall with injury %	C.I.	n	Fall no injury %	C.I.
Total	218	80.8	5.8	25	7.9	3.4	26	11.3	5.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Number of Cigarettes Smoked per day among Everyday Smokers

Kings

Demographic Groups	n¹	1-10 cigarettes %²	n	≥11 cigarettes %	C.I.³
Total	37	49.5	38	50.5	13.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from Television

Kings

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	379	63.6	221	36.4	4.5
Gender					
Male	136	63.8	80	36.2	7.4
Female	243	63.5	141	36.5	5.4
Age					
18-34	145	72.6	60	27.4	6.8
35-54	151	64.0	88	36.0	7.1
≥55	80	51.8	69	48.2	9.5
Education					
≤High School	175	69.2	73	30.8	6.6
>High School	201	58.5	147	41.5	6.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from Radio

Kings

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	192	32.6	407	67.4	4.3
Gender					
Male	76	36.2	134	63.8	7.4
Female	116	29.6	273	70.4	5.1
Age					
18-34	81	44.2	123	55.8	7.9
35-54	66	25.8	168	74.2	6.2
≥55	43	26.8	111	73.2	8.2
Education					
≤High School	83	33.1	165	66.9	6.7
>High School	109	32.9	236	67.1	5.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from Billboards

Kings

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	215	39.4	381	60.6	4.7
Gender					
Male	79	40.8	131	59.2	7.8
Female	136	38.3	250	61.7	5.6
Age					
18-34	90	48.2	114	51.8	7.9
35-54	79	37.1	154	62.9	7.3
≥55	45	31.4	110	68.6	9.1
Education					
≤High School	94	39.5	156	60.5	7.0
>High School	120	39.4	222	60.6	6.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from Buses

Kings

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	155	27.8	436	72.2	4.3

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from Magazines

Kings

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	296	50.8	308	49.2	4.7
Gender					
Male	109	52.0	106	48.0	7.7
Female	187	49.8	202	50.2	5.6
Age					
18-34	99	52.1	106	47.9	7.9
35-54	111	47.8	126	52.2	7.4
≥55	83	53.2	72	46.8	9.4
Education					
≤High School	135	52.7	117	47.3	7.1
>High School	160	49.3	187	50.7	6.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from a Website

Kings

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	71	12.2	536	87.8	3.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from Movie Theater

Kings

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	86	15.0	519	85.0	3.4

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard or Seen Anti-Tobacco Advertisement that Refers to the NYS Quitline in Past 30 Days

Kings

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	250	43.3	332	56.7	4.7
Gender					
Male	95	46.5	110	53.5	7.9
Female	155	40.7	222	59.3	5.6
Age					
18-34	95	48.5	100	51.5	8.1
35-54	93	40.1	135	59.9	7.3
≥55	59	40.7	94	59.3	9.4
Education					
≤High School	117	47.9	131	52.1	7.2
>High School	131	39.5	197	60.5	6.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Used or Tried Any Smokeless Tobacco Products such as Chewing Tobacco or Snuff

Kings

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	31	5.1	576	94.9	2.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Work Indoors Most of the Time

Kings

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	291	81.4	59	18.6	5.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Smoking Policy at Place of Work among Indoor Workers***Kings***

Demographic Groups	n¹	Not allowed anywhere %²	C.I.³	n	Allowed in some or all places %	C.I.	n	No official policy %	C.I.
Total	239	81.5	5.2	35	13.7	4.6	12	4.8	2.9
Gender									
Male	82	75.7	9.0	17	16.7	7.8	7	7.6	5.5
Female	157	86.8	5.5	18	10.9	5.1	5	2.3	2.2
Age									
18-34	83	77.6	9.3	17	19.8	9.0	4	2.6	2.9
35-54	117	84.1	6.9	14	9.6	5.2	6	6.3	5.0
≥55	36	82.8	13.6	4	10.8	10.4	2	6.5	9.9
Education									
≤High School	59	70.0	11.5	16	21.8	10.3	5	8.2	7.0
>High School	177	87.2	5.2	18	9.5	4.5	7	3.3	2.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution²Weighted Percent³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Use Seatbelt when Ride or Drive in a Car (Always, Nearly Always or Never Ride in a Car)

Kings

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	543	88.6	64	11.4	3.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Use Seatbelt when Ride or Drive in a Car (Always or Never ride in a Car)

Kings

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	465	76.1	142	23.9	4.0
Gender					
Male	154	71.8	62	28.2	6.9
Female	311	79.6	80	20.4	4.6
Age					
18-64	401	75.2	126	24.8	4.4
≥65	58	79.7	15	20.3	10.3
Education					
≤High School	195	76.0	54	24.0	6.3
>High School	265	75.9	87	24.1	5.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Trying to Lose Weight

Kings

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	315	48.7	293	51.3	4.6
Gender					
Male	91	39.7	124	60.3	7.5
Female	224	56.1	169	43.9	5.6
Age					
18-34	101	47.6	105	52.4	7.8
35-54	136	56.6	103	43.4	7.3
≥55	77	40.7	79	59.3	9.0
Education					
≤High School	124	42.5	127	57.5	6.9
>High School	187	53.8	164	46.2	6.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Trying to Maintain Current Weight of Those not Trying to Lose Weight

Kings

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	171	57.8	120	42.2	6.7
Gender					
Male	69	55.9	55	44.1	10.0
Female	102	60.1	65	39.9	8.6
Age					
18-34	55	48.2	49	51.8	11.2
35-54	62	60.7	41	39.3	10.8
≥55	49	65.2	29	34.8	11.8
Education					
≤High School	74	58.0	52	42.0	9.8
>High School	95	56.6	68	43.4	9.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Eating Fewer Calories or Less Fat to Lose or Maintain Weight

Kings

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	368	80.2	105	19.8	3.9
Gender					
Male	114	77.1	41	22.9	7.1
Female	254	82.5	64	17.5	4.4
Age					
18-34	114	75.7	38	24.3	7.5
35-54	150	81.3	46	18.7	5.7
≥55	98	83.4	21	16.6	7.7
Education					
≤High School	156	82.4	37	17.6	5.8
>High School	208	78.1	67	21.9	5.4

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Using Physical Exercise to Lose or Maintain Weight

Kings

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	328	68.8	157	31.2	4.7
Gender					
Male	113	72.9	47	27.1	7.8
Female	215	65.8	110	34.2	5.8
Age					
18-34	124	81.9	32	18.1	6.2
35-54	130	66.3	68	33.7	7.6
≥55	72	58.1	53	41.9	10.3
Education					
≤High School	116	62.4	81	37.6	7.6
>High School	209	74.7	73	25.3	5.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval