Expanded BRFSS, 2003

Dutchess and Putnam Locality

Table of Contents:	
Optional Module 1: Adult Asthma History	
Optional Module 4: Cholesterol Awareness	4
Optional Module 5: Diabetes	e
Optional Module 9: Fruits and Vegetables	10
Optional Module 10: Health Care Coverage	
Optional Module 12: Hypertension Awareness	
Optional Module 13: Injury Control - Falls	
Optional Module 14: Immunization	
Optional Module 19: Tobacco Consumption	
Optional Module 20: Tobacco - Media	20
Ontional Module 23: Seat Belts.	28

New York State Department of Health

Have had an Asthma Episode or Attack in the Past 12 Months, among Those with Asthma

Demographic Groups	n¹	Yes %°	n	No %	C.I. ³
Total	26	43.6	29	56.4	14.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

This page left blank intentionally

This page left blank intentionally

Ever had Blood Cholesterol Checked

Demographic		Yes		No	
Groups	n¹	% ²	n	%	C.I. ³
Total	518	84.6	78	15.4	3.8
Gender					
Male	209	86.2	21	13.8	6.1
Female	309	83.2	57	16.8	4.5
Age					
18-34	87	69.7	35	30.3	9.8
35-54	236	88.1	29	11.9	4.9
≥55	189	92.9	14	7.1	4.2
Education					
≤High School	162	78.7	35	21.3	7.2
>High School	356	87.9	43	12.1	4.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent ³95% Confidence Interval

Had Blood Cholesterol Checked in the Past Five Years

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	475	79.5	107	20.5	4.1
Gender					
Male	194	81.9	31	18.1	6.5
Female	281	77.2	76	22.8	5.0
Age					
18-34	73	62.4	44	37.6	10.3
35-54	215	82.4	43	17.6	5.6
≥55	181	90.1	20	9.9	4.7
Education					
≤High School	151	75.1	42	24.9	7.5
>High School	324	81.9	65	18.1	4.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Check Blood for Glucose or Sugar at Least Once per Day among Those with Diabetes *Dutchess and Putnam*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	25	58.1	20	41.9	15.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Check Feet for Sores or Irritations at Least Once per Day among Those with Diabetes *Dutchess and Putnam*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	30	72.0	14	28.0	13.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

Had Feet Checked by a Medical Professional in Past Twelve Months, among Those with Diabetes

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	34	80.6	9	19.4	12.4

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Had Dilated Eye Exam in Past 12 Months, among Those with Diabetes *Dutchess and Putnam*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	35	80.2	9	19.8	13.3

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

Eat Five Fruits or Vegetable Servings per Day

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	184	28.6	420	71.4	4.0
Gender					
Male	56	21.8	177	78.2	5.6
Female	128	35.4	243	64.6	5.4
Age					
18-34	25	18.4	100	81.6	7.4
35-54	79	28.9	187	71.1	6.0
≥55	77	36.8	130	63.2	7.4
Education					
≤High School	55	27.1	146	72.9	6.9
>High School	129	29.4	274	70.6	4.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Main Reason for Not Having Health Care Coverage

Demographic		Changed or lost job Could not afford				Other			
Groups	n¹	%2	C.I. ³	n	%	C.I.	n	%	C.I.
Total	6	13.2	11.0	20	36.5	15.4	25	50.3	16.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

No Health Care Coverage for Past Two Years Among Uninsured

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	28	57.4	23	42.6	15.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent ³95% Confidence Interval

Ever Been Told Have High Blood Pressure by a Medical Professional Dutchess and Putnam

Demographic	1	Yes		No	CI3
Groups	n¹	% ²	n	%	C.I. ³
Total	154	23.6	448	76.4	3.7
Gender					
Male	79	29.5	153	70.5	6.4
Female	75	17.7	295	82.3	4.0
Age					
18-34	9	5.1	116	94.9	3.7
35-54	50	20.7	215	79.3	5.5
≥55	94	45.4	113	54.6	7.7
Education					
≤High School	62	28.6	139	71.4	7.0
>High School	92	20.8	309	79.2	4.3

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Had a Fall in Past Three Months

Demographic Groups	n¹	No fall	C.I. ³	n	Fall with injury %	C.I.	n	Fall no injury %	C.I.
Total	279	85.9	4.3	15	5.7	3.1	34	8.3	3.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

Had a Flu Shot in the Past 12 Months, Aged 65 and Older

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	65	59.0	46	41.0	10.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent ³95% Confidence Interval

Had a Flushot in the Past 12 Months

Demographic Groups	n¹	Yes %2	n	No %	C.I. ³
Total	164	25.2	434	74.8	3.9
Gender					
Male	57	23.5	173	76.5	6.0
Female	107	26.9	261	73.1	4.9
Age					
18-64	96	18.3	386	81.7	3.9
≥65	65	59.0	46	41.0	10.2
Education					
≤High School	53	25.7	145	74.3	6.9
>High School	111	25.0	289	75.0	4.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

^{395%} Confidence Interval

Ever Had a Pneumonia Shot, Aged 65 and Older

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	74	75.0	33	25.0	8.4

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent ³95% Confidence Interval

Ever Had a Pneumonia Shot

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	132	23.8	430	76.2	4.1
Gender					
Male	52	25.7	155	74.3	6.7
Female	80	22.1	275	77.9	4.7
Age					
18-64	57	13.6	393	86.4	3.8
≥65	74	75.0	33	25.0	8.4
Education					
≤High School	53	29.1	134	70.9	7.4
>High School	79	21.0	296	79.0	4.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

^{395%} Confidence Interval

Number of Cigarettes Smoked per day among Everyday Smokers *Dutchess and Putnam*

Demographic Groups	n¹	1-10 cigarettes %	n	≥11 cigarettes %	C.I. ³
Total	35	38.4	68	61.6	12.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

Heard, Read or Seen Information about the Effects of Tobacco from Television *Dutchess and Putnam*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	370	63.8	218	36.2	4.5
Gender					
Male	153	66.2	78	33.8	7.0
Female	217	61.4	140	38.6	5.5
Age					
18-34	96	78.6	28	21.4	8.2
35-54	157	57.0	105	43.0	6.8
≥55	113	58.6	84	41.4	7.7
Education					
≤High School	132	67.3	63	32.7	7.6
>High School	238	61.9	155	38.1	5.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Heard, Read or Seen Information about the Effects of Tobacco from Radio *Dutchess and Putnam*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	161	28.5	424	71.5	4.4
Gender					
Male	78	34.7	148	65.3	7.2
Female	83	22.3	276	77.7	4.6
Age					
18-34	36	32.1	87	67.9	10.2
35-54	73	26.5	184	73.5	5.8
≥55	49	27.1	151	72.9	7.2
Education					
≤High School	56	29.7	139	70.3	7.5
>High School	105	27.8	285	72.2	5.3

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Heard, Read or Seen Information about the Effects of Tobacco from Billboards *Dutchess and Putnam*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	225	39.4	362	60.6	4.7
Gender					
Male	91	43.4	133	56.6	7.6
Female	134	35.4	229	64.6	5.3
Age					
18-34	66	54.5	57	45.5	10.2
35-54	107	38.2	154	61.8	6.6
≥55	50	26.8	148	73.2	7.1
Education					
≤High School	82	41.6	113	58.4	8.1
>High School	143	38.1	249	61.9	5.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Heard, Read or Seen Information about the Effects of Tobacco from Buses *Dutchess and Putnam*

Demographic Groups	n¹	Yes %2	n	No %	C.I. ³
Total	33	6.1	556	93.9	2.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent ³95% Confidence Interval

Heard, Read or Seen Information about the Effects of Tobacco from Magazines *Dutchess and Putnam*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	325	55.4	267	44.6	4.6
Gender					
Male	125	55.8	102	44.2	7.4
Female	200	55.1	165	44.9	5.6
Age					
18-34	67	54.3	57	45.7	10.3
35-54	140	52.8	123	47.2	6.8
≥55	114	59.4	86	40.6	7.6
Education					
≤High School	115	60.9	80	39.1	8.0
>High School	210	52.5	187	47.5	5.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Heard, Read or Seen Information about the Effects of Tobacco from a Website *Dutchess and Putnam*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	52	10.1	545	89.9	3.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

Heard, Read or Seen Information about the Effects of Tobacco from Movie Theater *Dutchess and Putnam*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	66	14.1	528	85.9	3.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Heard or Seen Anti-Tobacco Advertisement that Refers to the NYS Quitline in Past 30 Days

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	208	38.5	361	61.5	4.7
Gender					
Male	84	41.6	136	58.4	7.7
Female	124	35.4	225	64.6	5.5
Age					
18-34	55	52.2	60	47.8	10.7
35-54	99	37.9	157	62.1	6.7
≥55	53	27.1	141	72.9	7.0
Education					
≤High School	77	43.9	112	56.1	8.4
>High School	131	35.5	249	64.5	5.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Use Seatbelt when Ride or Drive in a Car (Always, Nearly Always or Never Ride in a Car)

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	547	90.0	51	10.0	3.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Use Seatbelt when Ride or Drive in a Car (Always or Never ride in a Car) Dutchess and Putnam

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	495	79.9	103	20.1	4.1
Gender					
Male	172	71.4	58	28.6	7.1
Female	323	88.4	45	11.6	3.5
Age					
18-64	398	78.6	86	21.4	4.7
≥65	92	86.0	17	14.0	7.4
Education					
≤High School	155	77.0	43	23.0	6.9
>High School	340	81.6	60	18.4	5.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

^{395%} Confidence Interval