Expanded BRFSS, 2003

Oneida, Oswego

Ever had Blood Cholesterol Checked

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	515	77.9	118	22.1	3.9
Gender					
Male	187	75.6	54	24.4	6.2
Female	328	80.1	64	19.9	4.8
Age					
18-34	79	55.7	58	44.3	9.4
35-54	209	82.3	45	17.7	5.2
≥55	224	92.9	15	7.1	3.9
Education					
≤High School	223	68.8	77	31.2	6.3
>High School	290	86.5	41	13.5	4.4

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health Had Blood Cholesterol Checked in the Past Five Years Oneida, Oswego

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	476	72.8	151	27.2	4.1
Gender					
Male	174	70.6	67	29.4	6.5
Female	302	74.9	84	25.1	5.1
Age					
18-34	72	50.7	65	49.3	9.4
35-54	190	75.7	64	24.3	5.8
≥55	211	90.0	22	10.0	4.5
Education					
≤High School	206	64.0	92	36.0	6.4
>High School	268	81.2	59	18.8	4.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Adults with a Disability

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	141	17.9	499	82.1	3.1
Gender					
Male	35	12.9	207	87.1	4.5
Female	106	22.6	292	77.4	4.3
Age					
18-34	10	5.2	129	94.8	3.7
35-54	63	20.7	194	79.3	5.3
≥55	68	26.7	173	73.3	6.2
Education					
≤High School	72	19.1	230	80.9	4.7
>High School	69	16.8	267	83.2	4.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent ³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health **Eat Five Fruits or Vegetable Servings per Day** Oneida, Oswego

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	164	23.8	478	76.2	3.6
Gender					
Male	42	15.9	201	84.1	4.9
Female	122	31.4	277	68.6	5.1
Age					
18-34	32	19.4	107	80.6	6.9
35-54	51	20.9	207	79.1	5.6
≥55	79	31.0	163	69.0	6.4
Education					
≤High School	61	18.4	243	81.6	4.8
>High School	102	29.0	234	71.0	5.3

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

Ever Been Told Have High Blood Pressure by a Medical Professional *Oneida, Oswego*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	181	26.7	461	73.3	3.8
Gender					
Male	77	29.4	167	70.6	6.3
Female	104	24.1	294	75.9	4.5
Age					
18-34	5	4.4	134	95.6	4.3
35-54	58	24.2	201	75.8	6.0
≥55	117	50.4	124	49.6	7.1
Education					
≤High School	102	29.9	202	70.1	5.7
>High School	78	23.5	258	76.5	5.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Had a Fall in Past Three Months

Demographic Groups	n¹	No fall	C.I. ³		Fall with injury %	C.I.	n	Fall no injury %	C.I.
Total	301	82.7	4.4	22	5.4	2.6	42	11.8	3.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Had a Flu Shot in the Past 12 Months, Aged 65 and Older

Oneida, Oswego

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	89	61.0	58	39.0	8.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

Had a Flushot in the Past 12 Months

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	215	32.2	424	67.8	4.1
Gender					
Male	83	33.0	159	67.0	6.6
Female	132	31.5	265	68.5	5.0
Age					
18-64	125	25.4	364	74.6	4.4
≥65	89	61.0	58	39.0	8.7
Education					
≤High School	101	29.6	201	70.4	5.7
>High School	112	34.4	223	65.6	5.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

^{395%} Confidence Interval

Ever Had a Pneumonia Shot, Aged 65 and Older

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	96	66.8	47	33.2	8.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

Ever Had a Pneumonia Shot

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	160	25.1	437	74.9	3.9
Gender					
Male	67	28.5	150	71.5	6.7
Female	93	22.1	287	77.9	4.5
Age					
18-64	64	14.7	387	85.3	4.0
≥65	96	66.8	47	33.2	8.5
Education					
≤High School	85	25.8	200	74.2	5.5
>High School	73	24.1	237	75.9	5.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

^{395%} Confidence Interval

Have Had Permanent Teeth Removed because of Tooth Decay or Gum Disease *Oneida, Oswego*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	364	52.7	263	47.3	4.5
Gender					
Male	133	49.6	107	50.4	7.1
Female	231	55.6	156	44.4	5.6
Age					
18-34	29	18.9	110	81.1	6.8
35-54	137	52.2	118	47.8	6.8
≥55	197	86.3	33	13.7	4.9
Education					
≤High School	194	58.2	100	41.8	6.6
>High School	168	47.3	163	52.7	6.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

^{395%} Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Having Soon a Dontal Professional in the Pas

Having Seen a Dental Professional in the Past 12 Months *Oneida, Oswego*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	452	75.2	136	24.8	4.1
Gender					
Male	160	69.4	64	30.6	6.9
Female	292	80.8	72	19.2	4.5
Age					
18-34	94	68.3	44	31.7	8.7
35-54	195	78.5	55	21.5	5.7
≥55	160	78.1	37	21.9	7.1
Education					
≤High School	189	68.4	78	31.6	6.6
>High School	262	81.2	58	18.8	5.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Trying to Lose Weight

Demographic Groups	n¹	Yes %2	n	No %	C.I. ³
Total	299	44.8	341	55.2	4.4
Gender					
Male	81	34.4	161	65.6	6.7
Female	218	54.6	180	45.4	5.4
Age					
18-34	59	39.1	80	60.9	9.0
35-54	139	53.0	118	47.0	6.8
≥55	99	40.2	142	59.8	6.9
Education					
≤High School	130	41.6	172	58.4	6.3
>High School	169	48.0	167	52.0	6.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent ³95% Confidence Interval

Trying to Maintain Current Weight of Those not Trying to Lose Weight *Oneida, Oswego*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	213	61.8	125	38.2	6.0
Gender					
Male	91	56.1	68	43.9	8.6
Female	122	69.5	57	30.5	7.5
Age					
18-34	42	53.4	38	46.6	12.2
35-54	77	63.5	41	36.5	9.7
≥55	93	67.9	46	32.1	8.6
Education					
≤High School	105	58.8	65	41.2	8.5
>High School	108	65.4	58	34.6	8.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

^{395%} Confidence Interval

Eating Fewer Calories or Less Fat to Lose or Maintain Weight *Oneida, Oswego*

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	385	74.8	121	25.2	4.5
Gender					
Male	120	68.5	50	31.5	8.1
Female	265	79.8	71	20.2	4.8
Age					
18-34	76	71.5	25	28.5	10.6
35-54	176	81.6	38	18.4	5.8
≥55	131	69.5	57	30.5	7.5
Education					
≤High School	173	74.5	58	25.5	6.5
>High School	212	75.1	63	24.9	6.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health Using Physical Exercise to Lose or Maintain Weight Oneida, Oswego

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	366	73.6	144	26.4	4.3
Gender					
Male	133	78.9	38	21.1	6.9
Female	233	69.5	106	30.5	5.4
Age					
18-34	82	79.4	19	20.6	9.0
35-54	165	78.4	51	21.6	6.0
≥55	117	62.8	73	37.2	7.7
Education					
≤High School	151	68.1	82	31.9	6.7
>High School	215	78.3	62	21.7	5.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Ever Had a Pap Smear

Demographic Groups	n¹	Yes %2	n	No %	C.I. ³
Total	309	96.5	11	3.5	2.4

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent

³95% Confidence Interval

Had a Pap Smear in Past 3 Years

Demographic Groups	n¹	Yes	n	No %	C.I. ³
Total	267	84.8	53	15.2	4.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution ²Weighted Percent