

# **New York State Expanded Behavioral Risk Factor Surveillance System**

**Erie County**

**Final Report  
July 2008 - June 2009**

**New York State Department of Health  
December 2009**

## Contents

<b>Introduction</b>	12
Table 1. Erie County and New York State adult US Census Bureau population estimates, 2007	13
<b>General Health</b>	
Table 2a. Fair or poor general health among adults in Erie County and New York State	14
Table 2b. Fair or poor general health among adults in Erie County and New York State - Age-adjusted	15
Table 3a. Poor physical health 14 or more days within the past month among adults in Erie County and New York State	16
Table 3b. Poor physical health 14 or more days within the past month among adults in Erie County and New York State - Age-adjusted	17
Table 4a. Poor mental health 14 or more days within the past month among adults in Erie County and New York State	18
Table 4b. Poor mental health 14 or more days within the past month among adults in Erie County and New York State - Age-adjusted	19
Table 5a. Poor physical or mental health 14 or more days within the past month among adults in Erie County and New York State	20
Table 5b. Poor physical or mental health 14 or more days within the past month among adults in Erie County and New York State - Age-adjusted	21
<b>Quality Health Care</b>	
Table 6a. Health insurance among adults in Erie County and New York State	22
Table 6b. Health insurance among adults in Erie County and New York State - Age-adjusted	23
Table 7a. Health insurance among adults aged 18-64 in Erie County and New York State	24
Table 7b. Health insurance among adults aged 18-64 in Erie County and New York State - Age-adjusted	25
Table 8a. No health insurance among adults in Erie County and New York State	26

## Contents

Table 8b. No health insurance among adults in Erie County and New York State - Age-adjusted	27
Table 9a. No health insurance among adults aged 18-64 in Erie County and New York State	28
Table 9b. No health insurance among adults aged 18-64 in Erie County and New York State - Age-adjusted	29
Table 10a. Adults with regular health care provider in Erie County and New York State	30
Table 10b. Adults with regular health care provider in Erie County and New York State - Age-adjusted	31
Table 11a. Cost prevented visit to doctor within the past year among adults in Erie County and New York State	32
Table 11b. Cost prevented visit to doctor within the past year among adults in Erie County and New York State - Age-adjusted	33
Table 12a. Visited a doctor for routine checkup within the past year among adults in Erie County and New York State	34
Table 12b. Visited a doctor for routine checkup within the past year among adults in Erie County and New York State - Age-adjusted	35
Table 13a. Visited a doctor for routine checkup within the past 2 years among adults in Erie County and New York State	36
Table 13b. Visited a doctor for routine checkup within the past 2 years among adults in Erie County and New York State - Age-adjusted	37
Table 14a. Dentist visit within the past year among adults in Erie County and New York State	38
Table 14b. Dentist visit within the past year among adults in Erie County and New York State - Age-adjusted	39
Table 15a. Had teeth cleaned within the past year among adults in Erie County and New York State	40
Table 15b. Had teeth cleaned within the past year among adults in Erie County and New York State - Age-adjusted	41
Table 16a. Permanent teeth extracted due to decay or gum disease among adults in Erie County and New York State	42

## Contents

Table 16b. Permanent teeth extracted due to decay or gum disease among adults in Erie County and New York State - Age-adjusted	43
Table 17a. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older in Erie County and New York State	44
Table 17b. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older in Erie County and New York State - Age-adjusted	45
Table 18a. Flu shot within the past 12 months in Erie County and New York State	46
Table 18b. Flu shot within the past 12 months in Erie County and New York State - Age-adjusted	47
Table 19a. Flu vaccine that was sprayed in nose within the past 12 months in Erie County and New York State	48
Table 19b. Flu vaccine that was sprayed in nose within the past 12 months in Erie County and New York State - Age-adjusted	49
Table 20a. Flu shot or flu vaccine sprayed in nose within the past 12 months in Erie County and New York State	50
Table 20b. Flu shot or flu vaccine sprayed in nose within the past 12 months in Erie County and New York State - Age-adjusted	51
Table 21a. Flu shot among adults aged 65 and older in Erie County and New York State	52
Table 21b. Flu shot among adults aged 65 and older in Erie County and New York State - Age-adjusted	53
Table 22a. Pneumonia shot or pneumococcal vaccine in Erie County and New York State	54
Table 22b. Pneumonia shot or pneumococcal vaccine in Erie County and New York State - Age-adjusted	55
Table 23a. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older in Erie County and New York State	56
Table 23b. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older in Erie County and New York State - Age-adjusted	57

## Contents

### Cancer Screening

Table 24a. Ever had a mammogram among women aged 40 and older in Erie County and New York State	58
Table 24b. Ever had a mammogram among women aged 40 and older in Erie County and New York State - Age-adjusted	59
Table 25a. Had mammogram within the past 2 years among women aged 40 and older in Erie County and New York State	60
Table 25b. Had mammogram within the past 2 years among women aged 40 and older in Erie County and New York State - Age-adjusted	61
Table 26a. Had mammogram within the past 2 years among women aged 50 and older in Erie County and New York State	62
Table 26b. Had mammogram within the past 2 years among women aged 50 and older in Erie County and New York State - Age-adjusted	63
Table 27a. Ever had a Pap test among women in Erie County and New York State	64
Table 27b. Ever had a Pap test among women in Erie County and New York State - Age-adjusted	65
Table 28a. Had a Pap test within the past 3 years among women in Erie County and New York State	66
Table 28b. Had a Pap test within the past 3 years among women in Erie County and New York State - Age-adjusted	67
Table 29a. Ever had a digital rectal exam among men aged 40 and older in Erie County and New York State	68
Table 29b. Ever had a digital rectal exam among men aged 40 and older in Erie County and New York State - Age-adjusted	69
Table 30a. Had digital rectal exam within the past 2 years among men aged 40 and older in Erie County and New York State	70
Table 30b. Had digital rectal exam within the past 2 years among men aged 40 and older in Erie County and New York State - Age-adjusted	71

## Contents

Table 31a. Ever had a prostate specific antigen test among men aged 40 and older in Erie County and New York State	72
Table 31b. Ever had a prostate specific antigen test among men aged 40 and older in Erie County and New York State - Age-adjusted	73
Table 32a. Had prostate specific antigen test within the past 2 years among men aged 40 and older in Erie County and New York State	74
Table 32b. Had prostate specific antigen test within the past 2 years among men aged 40 and older in Erie County and New York State - Age-adjusted	75
Table 33a. Prostate cancer among men aged 40 and older in Erie County and New York State	76
Table 33b. Prostate cancer among men aged 40 and older in Erie County and New York State - Age-adjusted	77
Table 34a. Home blood stool test ever used among adults aged 50 and older in Erie County and New York State	78
Table 34b. Home blood stool test ever used among adults aged 50 and older in Erie County and New York State - Age-adjusted	79
Table 35a. Home blood stool test used within the past year among adults aged 50 and older in Erie County and New York State	80
Table 35b. Home blood stool test used within the past year among adults aged 50 and older in Erie County and New York State - Age-adjusted	81
Table 36a. Home blood stool test used within the past 2 years among adults aged 50 and older in Erie County and New York State	82
Table 36b. Home blood stool test used within the past 2 years among adults aged 50 and older in Erie County and New York State - Age-adjusted	83
Table 37a. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in Erie County and New York State	84
Table 37b. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in Erie County and New York State - Age-adjusted	85
Table 38a. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Erie County and New York State	86

## Contents

Table 38b. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Erie County and New York State - Age-adjusted	87
Table 39a. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Erie County and New York State	88
Table 39b. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Erie County and New York State - Age-adjusted	89

### Chronic Diseases

Table 40a. High blood pressure among adults in Erie County and New York State	90
Table 40b. High blood pressure among adults in Erie County and New York State - Age-adjusted	91
Table 41a. Blood pressure medication among adults with high blood pressure in Erie County and New York State	92
Table 41b. Blood pressure medication among adults with high blood pressure in Erie County and New York State - Age-adjusted	93
Table 42a. Ever had blood cholesterol checked among adults in Erie County and New York State	94
Table 42b. Ever had blood cholesterol checked among adults in Erie County and New York State - Age-adjusted	95
Table 43a. Blood cholesterol checked within the past 5 years among adults in Erie County and New York State	96
Table 43b. Blood cholesterol checked within the past 5 years among adults in Erie County and New York State - Age-adjusted	97
Table 44a. Cardiovascular disease among adults in Erie County and New York State	98
Table 44b. Cardiovascular disease among adults in Erie County and New York State - Age-adjusted	99
Table 45a. Coronary heart disease among adults in Erie County and New York State	100
Table 45b. Coronary heart disease among adults in Erie County and New York State - Age-adjusted	101

## Contents

Table 46a. Ever diagnosed with diabetes among adults in Erie County and New York State	102
Table 46b. Ever diagnosed with diabetes among adults in Erie County and New York State - Age-adjusted	103
Table 47a. Ever diagnosed with asthma (lifetime) among adults in Erie County and New York State	104
Table 47b. Ever diagnosed with asthma (lifetime) among adults in Erie County and New York State - Age-adjusted	105
Table 48a. Current asthma among adults in Erie County and New York State	106
Table 48b. Current asthma among adults in Erie County and New York State - Age-adjusted	107
Table 49a. Chronic joint symptoms among adults in Erie County and New York State	108
Table 49b. Chronic joint symptoms among adults in Erie County and New York State - Age-adjusted	109
Table 50a. Arthritis among adults in Erie County and New York State	110
Table 50b. Arthritis among adults in Erie County and New York State - Age-adjusted	111
Table 51a. Activity limitations because of physical, mental or emotional problems among adults in Erie County and New York State	112
Table 51b. Activity limitations because of physical, mental or emotional problems among adults in Erie County and New York State - Age-adjusted	113
Table 52a. Have health problem that requires the use of special equipment among adults in Erie County and New York State	114
Table 52b. Have health problem that requires the use of special equipment among adults in Erie County and New York State - Age-adjusted	115
Table 53a. Disability among adults in Erie County and New York State	116
Table 53b. Disability among adults in Erie County and New York State - Age-adjusted	117

## Health Behaviors

Table 54a. Current smoking among adults in Erie County and New York State	118
---	-----



## Contents

Table 54b. Current smoking among adults in Erie County and New York State - Age-adjusted	119
Table 55a. Everyday smoking among adults in Erie County and New York State	120
Table 55b. Everyday smoking among adults in Erie County and New York State - Age-adjusted	121
Table 56a. Adults living in homes in which smoking is prohibited in Erie County and New York State	122
Table 56b. Adults living in homes in which smoking is prohibited in Erie County and New York State - Age-adjusted	123
Table 57a. Binge drinking within the past month among adults in Erie County and New York State	124
Table 57b. Binge drinking within the past month among adults in Erie County and New York State - Age-adjusted	125
Table 58a. Heavy drinking within the past month among adults in Erie County and New York State	126
Table 58b. Heavy drinking within the past month among adults in Erie County and New York State - Age-adjusted	127
Table 59a. Never or rarely been asked about sexual history during routine check-up among adults in Erie County and New York State	128
Table 59b. Never or rarely been asked about sexual history during routine check-up among adults in Erie County and New York State - Age-adjusted	129
Table 60a. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults in Erie County and New York State	130
Table 60b. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults in Erie County and New York State - Age-adjusted	131
Table 61a. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults in Erie County and New York State	132
Table 61b. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults in Erie County and New York State - Age-adjusted	133

## Contents

### Overweight / Obesity

Table 62a. Overweight among adults in Erie County and New York State	134
Table 62b. Overweight among adults in Erie County and New York State - Age-adjusted	135
Table 63a. Obesity among adults in Erie County and New York State	136
Table 63b. Obesity among adults in Erie County and New York State - Age-adjusted	137
Table 64a. Overweight or obesity among adults in Erie County and New York State	138
Table 64b. Overweight or obesity among adults in Erie County and New York State - Age-adjusted	139
Table 65a. Received advice about weight by a health professional among adults in Erie County and New York State	140
Table 65b. Received advice about weight by a health professional among adults in Erie County and New York State - Age-adjusted	141
Table 66a. Received advice to lose weight by a health professional among adults provided weight advice in Erie County and New York State	142
Table 66b. Received advice to lose weight by a health professional among adults provided weight advice in Erie County and New York State - Age-adjusted	143

### Physical Activity / Nutrition

Table 67a. No leisure-time physical activity among adults in Erie County and New York State	144
Table 67b. No leisure-time physical activity among adults in Erie County and New York State - Age-adjusted	145
Table 68a. Consumption of 5 or more servings of fruits and vegetables per day among adults in Erie County and New York State	146
Table 68b. Consumption of 5 or more servings of fruits and vegetables per day among adults in Erie County and New York State - Age-adjusted	147

## Contents

### Injury

Table 69a. Had a fall within the past 3 months among adults in Erie County and New York State	148
Table 69b. Had a fall within the past 3 months among adults in Erie County and New York State - Age-adjusted	149
Table 70a. Had a fall that caused injury within the past 3 months among adults in Erie County and New York State	150
Table 70b. Had a fall that caused injury within the past 3 months among adults in Erie County and New York State - Age-adjusted	151

## Introduction

The New York State Expanded Behavioral Risk Factor Surveillance System (Expanded BRFSS) is a random digit dialed telephone survey of adults (18 years of age and older) that is representative of the non-institutionalized civilian population with landline telephones living in New York State. The goal of the Expanded BRFSS is to collect county-specific data on preventive health practices, risk behaviors, injuries and preventable chronic and infectious diseases. Topics assessed by the Expanded BRFSS include tobacco use, physical inactivity, diet, use of cancer screening services and other factors linked to the leading causes of morbidity and mortality. For each topic, health indicators are based on answers to one or more questions in the survey. The survey results are weighted to more accurately represent the adult population living in each county. Weighting procedures adjust for the differential selection probabilities by race, age and gender. More detailed information on weighting procedures is available in the technical brief.

Survey data collection occurred in 58 geographic areas of New York State, including each of the 57 counties outside of New York City and New York City (5 counties) as a single area. A standard questionnaire was used in all areas. Data collection for the 2009 Expanded BRFSS took place during the period of July 2008 through June 2009. The goal was to complete about 650 interviews in each area. The final number of interviews varies because of the sampling design of the survey and differential response rates. More detailed information on the sampling design is available in the technical brief.

The New York State Department of Health contracted with RTI International, which is responsible for the data collection, sampling, weighting and creation of the statistical reports. RTI International subcontracted with Clearwater Research Inc. for data collection and additional expertise in sample design and data quality control.

This report presents both crude estimates and estimates that have been age-adjusted to the Year 2000 US Standard Population when they represent the entire adult population of an area. Estimates for specific age groups are not age-adjusted. Age adjustment is a statistical process applied to rates that allows communities with different age structures to be compared. Counties can compare their data with results from New York State as a whole and with other counties. Confidence intervals at the 95% level associated with each health indicator give an estimated range of values which is likely to include the actual prevalence of the indicator. Confidence intervals are similar to margins of error. When the confidence intervals of two estimates of the same indicator from different areas do not overlap, they may be said to be statistically significantly different, i.e., these differences are unlikely related to chance and are considered true differences. If there is any value that is included in both intervals, the two estimates are not statistically significantly different.

The survey results for each of the 69 health indicators included in the survey are typically provided by gender, age, race and ethnicity, education and income in the Expanded BRFSS county reports. The estimates for some of these groupings may be suppressed in geographic areas with low proportions of adult respondents in these groupings.

**Expanded BRFSS Report: July 2008 - June 2009  
Erie County**

**Table 1. Erie County and New York State adult US Census Bureau population estimates,<sup>1</sup> 2007**

	Erie County		New York State	
	Adult Population <sup>1</sup>	Percent	Adult Population <sup>1</sup>	Percent
<b>Total</b>	713,480	100	14,884,315	100
<b>Gender</b>				
Male	337,843	47.4	7,103,791	47.7
Female	375,637	52.6	7,780,524	52.3
<b>Age (years)</b>				
18-34	201,824	28.3	4,497,479	30.2
35-44	121,131	17.0	2,844,115	19.1
45-54	142,436	20.0	2,854,316	19.2
55-64	106,618	14.9	2,142,000	14.4
≥ 65	141,471	19.8	2,546,405	17.1
<b>Race/Ethnicity</b>				
White Non-Hispanic	586,553	82.2	9,343,884	62.8
Black Non-Hispanic	85,873	12.0	2,156,941	14.5
Hispanic	22,336	3.1	2,258,717	15.2
Other Non-Hispanic	18,718	2.6	1,124,773	7.6

<sup>1</sup> Source: National Center for Health Statistics. Postcensal estimates of the resident population of the United States for July 1, 2000-July 1, 2007, by year, county, age, bridged race, Hispanic origin, and sex (Vintage 2007). Prepared under a collaborative arrangement with the U.S. Census Bureau; released August 7, 2008. Available from: [http://www.cdc.gov/nchs/nvss/bridged\\_race.htm](http://www.cdc.gov/nchs/nvss/bridged_race.htm) as of November 25, 2009.

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 2a. Fair or poor general health among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	661	116,400	16.3	(12.9 - 19.7)	38,124	2,531,400	17.1	(15.1 - 19.0)
<b>Gender</b>								
Male	239	57,500	17.0	(11.1 - 23.0)	14,599	1,069,700	15.1	(12.4 - 17.8)
Female	422	58,800	15.7	(12.0 - 19.3)	23,525	1,461,700	18.8	(16.1 - 21.6)
<b>Age (years)</b>								
18-34	*	*	*	*	3,940	535,800	12.0	(7.2 - 16.8)
35-44	*	*	*	*	4,985	289,200	10.0	(6.8 - 13.3)
45-54	149	28,700	20.2	(11.1 - 29.2)	7,819	507,800	17.9	(14.0 - 21.8)
55-64	*	*	*	*	8,757	424,700	19.9	(16.0 - 23.9)
≥ 65	224	42,800	30.4	(23.3 - 37.5)	12,623	773,900	30.5	(26.8 - 34.2)
<b>Race/Ethnicity</b>								
White Non-Hispanic	587	82,700	14.1	(11.1 - 17.1)	35,811	1,226,600	13.2	(11.7 - 14.6)
Black Non-Hispanic	*	*	*	*	739	457,300	21.3	(14.8 - 27.9)
Hispanic	*	*	*	*	861	684,100	30.4	(22.2 - 38.6)
Other Non-Hispanic	*	*	*	*	713	163,400	14.5	(5.1 - 24.0)
<b>Education</b>								
≤ High School	265	69,400	24.6	(17.8 - 31.3)	15,682	1,466,500	26.3	(22.6 - 30.0)
Some College	176	26,400	13.7	(7.5 - 19.9)	9,904	508,500	14.3	(10.9 - 17.8)
≥ College Degree	218	20,200	8.5	(4.9 - 12.2)	12,457	493,500	8.8	(6.5 - 11.1)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,633	1,269,200	36.3	(30.1 - 42.4)
\$25,000-\$49,999	163	27,100	16.1	(9.7 - 22.4)	9,649	447,600	15.4	(11.9 - 18.9)
\$50,000-\$74,999	*	*	*	*	5,769	237,800	12.1	(8.0 - 16.2)
≥ \$75,000	136	17,200	8.2	(3.0 - 13.4)	8,562	218,300	4.9	(3.6 - 6.1)

<sup>1</sup> Defined as reporting fair or poor general health

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have fair or poor health (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 2b. Fair or poor general health among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	661	14.8	(11.5 - 18.1)	38,124	16.7	(14.8 - 18.6)
<b>Gender</b>						
Male	239	16.1	(10.4 - 21.8)	14,599	15.3	(12.6 - 18.0)
Female	422	13.7	(10.2 - 17.2)	23,525	18.0	(15.2 - 20.7)
<b>Race/Ethnicity</b>						
White Non-Hispanic	587	12.9	(10.0 - 15.9)	35,811	12.9	(11.4 - 14.4)
Black Non-Hispanic	*	*	*	739	20.8	(14.2 - 27.3)
Hispanic	*	*	*	861	29.9	(21.8 - 37.9)
Other Non-Hispanic	*	*	*	713	13.8	(4.8 - 22.8)
<b>Education</b>						
≤ High School	265	23.1	(16.4 - 29.9)	15,682	25.8	(22.1 - 29.5)
Some College	176	12.4	(6.5 - 18.2)	9,904	14.1	(10.7 - 17.6)
≥ College Degree	218	7.3	(4.0 - 10.6)	12,457	8.5	(6.3 - 10.8)
<b>Income</b>						
≤ \$24,999	*	*	*	9,633	35.9	(29.7 - 42.0)
\$25,000-\$49,999	163	15.5	(9.0 - 22.0)	9,649	15.0	(11.6 - 18.5)
\$50,000-\$74,999	*	*	*	5,769	11.8	(7.7 - 16.0)
≥ \$75,000	136	7.5	(2.8 - 12.2)	8,562	4.7	(3.5 - 6.0)

<sup>1</sup> Defined as reporting fair or poor general health

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Erie County**

**Table 3a. Poor physical health 14 or more days within the past month among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	643	86,600	12.5	(9.4 - 15.6)	37,327	1,472,900	10.1	(8.7 - 11.6)
<b>Gender</b>								
Male	233	24,000	7.3	(3.7 - 10.9)	14,325	604,200	8.7	(7.0 - 10.5)
Female	410	62,600	17.1	(12.4 - 21.8)	23,002	868,600	11.4	(9.2 - 13.6)
<b>Age (years)</b>								
18-34	*	*	*	*	3,891	268,100	6.1	(2.7 - 9.5)
35-44	*	*	*	*	4,948	232,100	8.1	(5.2 - 11.1)
45-54	147	15,400	11.0	(5.3 - 16.8)	7,764	350,900	12.5	(9.4 - 15.6)
55-64	131	24,900	25.0	(16.0 - 33.9)	8,621	271,200	13.0	(10.3 - 15.8)
≥ 65	217	25,400	18.6	(12.5 - 24.7)	12,103	350,600	14.5	(12.1 - 16.9)
<b>Race/Ethnicity</b>								
White Non-Hispanic	572	62,000	10.8	(8.0 - 13.6)	35,073	929,700	10.2	(8.8 - 11.6)
Black Non-Hispanic	*	*	*	*	724	208,200	9.9	(4.8 - 14.9)
Hispanic	*	*	*	*	835	250,300	11.5	(6.5 - 16.4)
Other Non-Hispanic	*	*	*	*	695	84,600	7.6	(1.3 - 13.9)
<b>Education</b>								
≤ High School	255	48,900	17.9	(12.0 - 23.8)	15,178	692,800	12.9	(10.6 - 15.3)
Some College	173	17,000	9.1	(4.6 - 13.6)	9,751	362,500	10.3	(7.4 - 13.2)
≥ College Degree	214	20,300	8.7	(3.8 - 13.6)	12,327	359,300	6.4	(4.8 - 8.1)
<b>Income</b>								
≤ \$24,999	175	38,200	25.4	(16.5 - 34.4)	9,294	625,600	18.4	(13.6 - 23.3)
\$25,000-\$49,999	159	12,800	7.8	(3.8 - 11.8)	9,495	279,100	9.8	(7.3 - 12.3)
\$50,000-\$74,999	*	*	*	*	5,733	186,400	9.5	(5.8 - 13.2)
≥ \$75,000	134	14,000	6.8	(1.4 - 12.3)	8,510	219,100	4.9	(3.6 - 6.2)

<sup>1</sup> Defined as 14 or more days within the past month when physical health was not good

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults with poor physical health 14 or more days within the past month (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 3b. Poor physical health 14 or more days within the past month among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	643	11.4	(8.3 - 14.5)	37,327	9.9	(8.4 - 11.4)
<b>Gender</b>						
Male	233	6.8	(3.3 - 10.3)	14,325	8.7	(6.9 - 10.5)
Female	410	15.5	(10.7 - 20.4)	23,002	11.0	(8.7 - 13.2)
<b>Race/Ethnicity</b>						
White Non-Hispanic	572	9.7	(6.9 - 12.5)	35,073	9.9	(8.5 - 11.3)
Black Non-Hispanic	*	*	*	724	9.7	(4.6 - 14.9)
Hispanic	*	*	*	835	11.3	(6.3 - 16.2)
Other Non-Hispanic	*	*	*	695	7.6	(1.2 - 14.0)
<b>Education</b>						
≤ High School	255	17.0	(11.0 - 23.1)	15,178	12.7	(10.2 - 15.1)
Some College	173	8.1	(3.7 - 12.6)	9,751	10.1	(7.2 - 12.9)
≥ College Degree	214	7.6	(2.9 - 12.4)	12,327	6.3	(4.7 - 8.0)
<b>Income</b>						
≤ \$24,999	175	24.6	(15.2 - 34.0)	9,294	18.4	(13.5 - 23.4)
\$25,000-\$49,999	159	6.4	(3.0 - 9.9)	9,495	9.4	(7.0 - 11.9)
\$50,000-\$74,999	*	*	*	5,733	9.0	(5.3 - 12.6)
≥ \$75,000	134	6.8	(1.2 - 12.4)	8,510	4.9	(3.5 - 6.2)

<sup>1</sup> Defined as 14 or more days within the past month when physical health was not good

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 4a. Poor mental health 14 or more days within the past month among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	650	69,300	9.9	(7.3 - 12.5)	37,528	1,473,200	10.1	(8.6 - 11.5)
<b>Gender</b>								
Male	234	29,000	8.7	(4.9 - 12.6)	14,407	555,200	7.9	(6.3 - 9.6)
Female	416	40,300	11.0	(7.4 - 14.5)	23,121	918,000	12.0	(9.7 - 14.3)
<b>Age (years)</b>								
18-34	*	*	*	*	3,889	496,700	11.2	(7.4 - 15.1)
35-44	91	19,000	16.1	(8.2 - 24.0)	4,947	254,100	8.9	(6.7 - 11.1)
45-54	147	13,400	9.6	(4.6 - 14.6)	7,749	347,200	12.3	(9.0 - 15.6)
55-64	136	18,700	17.7	(9.4 - 25.9)	8,623	211,300	10.1	(7.9 - 12.3)
≥ 65	217	10,600	7.9	(3.8 - 12.0)	12,320	163,900	6.7	(4.7 - 8.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	580	58,500	10.1	(7.3 - 12.9)	35,266	899,200	9.8	(8.4 - 11.2)
Black Non-Hispanic	*	*	*	*	722	227,800	10.9	(5.4 - 16.4)
Hispanic	*	*	*	*	845	263,700	11.8	(6.8 - 16.8)
Other Non-Hispanic	*	*	*	*	695	82,500	7.4	(1.4 - 13.5)
<b>Education</b>								
≤ High School	253	31,700	11.8	(7.1 - 16.5)	15,335	665,000	12.2	(10.0 - 14.5)
Some College	176	20,000	10.4	(4.8 - 16.0)	9,771	404,500	11.5	(8.3 - 14.8)
≥ College Degree	218	17,500	7.4	(4.0 - 10.8)	12,342	346,600	6.2	(4.5 - 7.9)
<b>Income</b>								
≤ \$24,999	175	26,000	17.6	(10.0 - 25.2)	9,398	611,500	17.9	(13.0 - 22.8)
\$25,000-\$49,999	160	13,600	8.2	(3.0 - 13.5)	9,547	260,300	9.0	(6.7 - 11.4)
\$50,000-\$74,999	*	*	*	*	5,726	187,300	9.6	(6.0 - 13.1)
≥ \$75,000	136	13,800	6.6	(2.6 - 10.6)	8,520	276,500	6.2	(4.6 - 7.7)

<sup>1</sup> Defined as 14 or more days within the past month when mental health was not good

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults with poor mental health 14 or more days within the past month (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 4b. Poor mental health 14 or more days within the past month among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	650	9.9	(7.1 - 12.7)	37,528	10.2	(8.7 - 11.7)
<b>Gender</b>						
Male	234	8.8	(4.8 - 12.8)	14,407	7.9	(6.3 - 9.6)
Female	416	10.9	(7.1 - 14.8)	23,121	12.2	(9.8 - 14.6)
<b>Race/Ethnicity</b>						
White Non-Hispanic	580	10.3	(7.3 - 13.4)	35,266	10.0	(8.5 - 11.4)
Black Non-Hispanic	*	*	*	722	11.1	(5.5 - 16.7)
Hispanic	*	*	*	845	11.5	(6.5 - 16.4)
Other Non-Hispanic	*	*	*	695	7.6	(1.5 - 13.7)
<b>Education</b>						
≤ High School	253	11.6	(6.8 - 16.4)	15,335	12.3	(10.0 - 14.6)
Some College	176	10.8	(4.6 - 17.0)	9,771	11.8	(8.5 - 15.1)
≥ College Degree	218	7.5	(3.8 - 11.1)	12,342	6.3	(4.6 - 8.0)
<b>Income</b>						
≤ \$24,999	175	16.9	(9.1 - 24.7)	9,398	18.2	(13.2 - 23.1)
\$25,000-\$49,999	160	8.4	(2.6 - 14.2)	9,547	9.4	(6.9 - 11.9)
\$50,000-\$74,999	*	*	*	5,726	9.5	(5.9 - 13.1)
≥ \$75,000	136	7.5	(2.9 - 12.1)	8,520	6.2	(4.6 - 7.7)

<sup>1</sup> Defined as 14 or more days within the past month when mental health was not good

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 5a. Poor physical or mental health 14 or more days within the past month among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	639	126,300	18.3	(14.7 - 21.9)	37,005	2,385,900	16.5	(14.8 - 18.2)
<b>Gender</b>								
Male	230	40,900	12.6	(8.0 - 17.1)	14,231	937,700	13.6	(11.4 - 15.8)
Female	409	85,400	23.4	(18.1 - 28.7)	22,774	1,448,200	19.1	(16.5 - 21.7)
<b>Age (years)</b>								
18-34	*	*	*	*	3,859	602,600	13.8	(9.7 - 17.9)
35-44	90	20,200	17.3	(9.1 - 25.4)	4,924	388,300	13.7	(10.4 - 17.0)
45-54	148	22,300	15.8	(9.2 - 22.5)	7,722	551,000	19.6	(15.8 - 23.4)
55-64	130	33,100	33.3	(23.6 - 43.0)	8,545	395,500	19.2	(16.1 - 22.3)
≥ 65	213	31,500	23.9	(17.1 - 30.6)	11,955	448,500	18.8	(15.9 - 21.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	568	97,800	17.1	(13.5 - 20.6)	34,778	1,485,700	16.3	(14.8 - 17.8)
Black Non-Hispanic	*	*	*	*	715	343,700	16.5	(10.3 - 22.7)
Hispanic	*	*	*	*	832	412,900	19.0	(12.6 - 25.3)
Other Non-Hispanic	*	*	*	*	680	143,500	13.0	(5.0 - 21.0)
<b>Education</b>								
≤ High School	250	63,900	23.8	(17.2 - 30.5)	15,018	1,077,200	20.3	(17.3 - 23.2)
Some College	174	28,500	15.1	(8.9 - 21.4)	9,672	615,500	17.6	(14.1 - 21.1)
≥ College Degree	214	33,600	14.4	(8.7 - 20.1)	12,244	634,200	11.4	(9.2 - 13.6)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,199	924,000	27.3	(21.9 - 32.8)
\$25,000-\$49,999	158	23,100	14.0	(7.8 - 20.2)	9,425	456,000	16.1	(13.0 - 19.2)
\$50,000-\$74,999	91	10,700	10.9	(3.8 - 17.9)	5,696	312,000	16.0	(11.4 - 20.5)
≥ \$75,000	134	24,300	11.9	(5.5 - 18.2)	8,475	447,600	10.0	(8.1 - 12.0)

<sup>1</sup> Defined as 14 or more days within the past month when physical or mental health prevented usual activities, such as self-care, work or recreation

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults with poor physical or mental health 14 or more days within the past month (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 5b. Poor physical or mental health 14 or more days within the past month among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	639	17.5	(13.7 - 21.2)	37,005	16.4	(14.7 - 18.2)
<b>Gender</b>						
Male	230	12.2	(7.6 - 16.8)	14,231	13.6	(11.4 - 15.8)
Female	409	22.1	(16.5 - 27.7)	22,774	19.0	(16.3 - 21.6)
<b>Race/Ethnicity</b>						
White Non-Hispanic	568	16.5	(12.7 - 20.2)	34,778	16.3	(14.7 - 17.8)
Black Non-Hispanic	*	*	*	715	16.7	(10.3 - 23.0)
Hispanic	*	*	*	832	18.5	(12.2 - 24.8)
Other Non-Hispanic	*	*	*	680	13.1	(5.0 - 21.3)
<b>Education</b>						
≤ High School	250	22.9	(16.0 - 29.8)	15,018	20.1	(17.1 - 23.1)
Some College	174	14.6	(7.9 - 21.3)	9,672	17.6	(14.1 - 21.1)
≥ College Degree	214	13.8	(8.0 - 19.6)	12,244	11.4	(9.2 - 13.7)
<b>Income</b>						
≤ \$24,999	*	*	*	9,199	27.6	(22.1 - 33.1)
\$25,000-\$49,999	158	13.3	(6.8 - 19.7)	9,425	16.1	(12.9 - 19.3)
\$50,000-\$74,999	91	10.6	(3.3 - 17.8)	5,696	15.6	(11.0 - 20.1)
≥ \$75,000	134	12.4	(5.7 - 19.1)	8,475	10.0	(8.0 - 12.0)

<sup>1</sup> Defined as 14 or more days within the past month when physical or mental health prevented usual activities, such as self-care, work or recreation

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 6a. Health insurance among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	661	666,000	93.5	(90.4 - 96.6)	38,119	12,854,200	86.6	(84.2 - 89.0)
<b>Gender</b>								
Male	238	314,200	93.3	(88.3 - 98.3)	14,597	5,881,100	83.0	(78.9 - 87.1)
Female	423	351,800	93.7	(89.8 - 97.5)	23,522	6,973,100	89.9	(87.2 - 92.6)
<b>Age (years)</b>								
18-34	59	178,700	88.6	(79.1 - 98.0)	3,910	3,302,100	74.3	(67.8 - 80.7)
35-44	92	111,000	91.6	(84.9 - 98.3)	4,986	2,560,300	88.9	(84.7 - 93.1)
45-54	149	137,000	96.1	(93.3 - 99.0)	7,831	2,530,100	88.9	(84.6 - 93.2)
55-64	137	101,400	95.1	(90.6 - 99.6)	8,764	1,986,700	92.8	(90.2 - 95.5)
≥ 65	224	138,000	98.3	(95.9 - 100)	12,628	2,475,000	97.8	(96.7 - 98.8)
<b>Race/Ethnicity</b>								
White Non-Hispanic	587	548,900	93.7	(90.3 - 97.2)	35,811	8,591,200	92.2	(90.6 - 93.8)
Black Non-Hispanic	*	*	*	*	736	1,781,600	82.7	(75.3 - 90.1)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	264	253,300	90.0	(83.7 - 96.3)	15,673	4,448,900	79.7	(74.9 - 84.5)
Some College	176	177,300	92.0	(85.7 - 98.3)	9,906	3,004,600	85.0	(80.1 - 89.8)
≥ College Degree	218	233,800	98.8	(97.6 - 99.9)	12,459	5,361,700	95.3	(93.3 - 97.2)
<b>Income</b>								
≤ \$24,999	180	137,500	89.7	(83.3 - 96.0)	9,640	2,437,200	69.7	(62.5 - 76.8)
\$25,000-\$49,999	163	156,600	92.8	(86.4 - 99.2)	9,662	2,384,000	82.2	(76.1 - 88.3)
\$50,000-\$74,999	91	95,600	97.2	(92.8 - 100)	5,772	1,855,700	94.0	(91.2 - 96.7)
≥ \$75,000	136	198,900	95.0	(87.9 - 100)	8,567	4,388,400	97.5	(96.6 - 98.5)

<sup>1</sup> Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have health insurance (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 6b. Health insurance among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	661	93.1	(89.7 - 96.4)	38,119	86.7	(84.4 - 89.1)
<b>Gender</b>						
Male	238	93.2	(88.1 - 98.3)	14,597	83.6	(79.7 - 87.5)
Female	423	92.9	(88.5 - 97.4)	23,522	89.6	(86.8 - 92.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	587	93.2	(89.4 - 97.0)	35,811	92.1	(90.5 - 93.7)
Black Non-Hispanic	*	*	*	736	82.5	(75.0 - 90.0)
Hispanic	*	*	*	860	68.7	(59.0 - 78.4)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	264	89.2	(82.4 - 96.1)	15,673	80.1	(75.4 - 84.7)
Some College	176	91.3	(84.1 - 98.5)	9,906	84.6	(79.7 - 89.4)
≥ College Degree	218	98.7	(97.4 - 100)	12,459	95.4	(93.6 - 97.2)
<b>Income</b>						
≤ \$24,999	180	88.9	(81.9 - 95.8)	9,640	70.2	(63.2 - 77.1)
\$25,000-\$49,999	163	92.3	(85.1 - 99.6)	9,662	81.9	(76.0 - 87.9)
\$50,000-\$74,999	91	97.0	(91.9 - 100)	5,772	94.0	(91.3 - 96.7)
≥ \$75,000	136	94.8	(87.6 - 100)	8,567	97.5	(96.6 - 98.5)

<sup>1</sup> Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 7a. Health insurance among adults aged 18-64<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	432	524,600	92.3	(88.4 - 96.1)	25,165	10,251,300	84.2	(81.3 - 87.1)
<b>Gender</b>								
Male	173	259,300	92.7	(86.8 - 98.6)	10,078	4,839,900	80.3	(75.5 - 85.0)
Female	259	265,300	91.9	(86.9 - 96.8)	15,087	5,411,400	88.0	(84.7 - 91.3)
<b>Age (years)</b>								
18-34	59	178,700	88.6	(79.1 - 98.0)	3,910	3,302,100	74.3	(67.8 - 80.7)
35-44	91	109,900	91.5	(84.8 - 98.3)	4,981	2,556,600	88.9	(84.7 - 93.1)
45-54	148	136,600	96.1	(93.3 - 99.0)	7,791	2,472,900	88.7	(84.3 - 93.0)
55-64	134	99,400	95.0	(90.4 - 99.6)	8,483	1,919,700	92.8	(90.1 - 95.5)
≥ 65	-	-	-	-	-	-	-	-
<b>Race/Ethnicity</b>								
White Non-Hispanic	369	418,400	92.4	(88.0 - 96.9)	23,354	6,582,200	90.3	(88.3 - 92.4)
Black Non-Hispanic	*	*	*	*	567	1,464,100	80.2	(71.6 - 88.7)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	141	174,400	86.5	(77.8 - 95.2)	9,037	3,191,500	74.6	(68.6 - 80.6)
Some College	129	151,600	91.7	(84.5 - 98.8)	7,031	2,540,400	82.9	(77.4 - 88.4)
≥ College Degree	161	198,300	98.5	(97.2 - 99.9)	9,068	4,502,400	94.5	(92.2 - 96.8)
<b>Income</b>								
≤ \$24,999	88	87,900	85.4	(75.9 - 94.9)	4,918	1,707,200	62.6	(54.1 - 71.2)
\$25,000-\$49,999	109	118,700	91.8	(83.8 - 99.9)	6,180	1,783,000	77.7	(70.3 - 85.1)
\$50,000-\$74,999	75	84,800	96.9	(92.0 - 100)	4,554	1,553,400	92.9	(89.7 - 96.1)
≥ \$75,000	122	188,100	94.7	(87.3 - 100)	7,324	3,987,600	97.3	(96.3 - 98.3)

<sup>1</sup> Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 18-64 who have health insurance (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 7b. Health insurance among adults aged 18-64<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	432	92.0	(88.0 - 96.0)	25,165	84.3	(81.5 - 87.2)
<b>Gender</b>						
Male	173	92.5	(86.5 - 98.6)	10,078	80.5	(75.9 - 85.1)
Female	259	91.5	(86.2 - 96.9)	15,087	87.9	(84.6 - 91.2)
<b>Race/Ethnicity</b>						
White Non-Hispanic	369	92.1	(87.5 - 96.8)	23,354	90.3	(88.3 - 92.3)
Black Non-Hispanic	*	*	*	567	80.0	(71.3 - 88.6)
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	141	86.0	(77.0 - 95.0)	9,037	75.1	(69.3 - 80.9)
Some College	129	91.0	(83.0 - 99.1)	7,031	82.6	(77.1 - 88.1)
≥ College Degree	161	98.5	(97.0 - 100)	9,068	94.6	(92.5 - 96.8)
<b>Income</b>						
≤ \$24,999	88	84.9	(74.9 - 94.9)	4,918	63.2	(54.9 - 71.6)
\$25,000-\$49,999	109	91.5	(82.6 - 100)	6,180	77.5	(70.2 - 84.7)
\$50,000-\$74,999	75	96.7	(91.1 - 100)	4,554	93.0	(89.8 - 96.2)
≥ \$75,000	122	94.6	(87.0 - 100)	7,324	97.3	(96.3 - 98.3)

<sup>1</sup> Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 8a. No health insurance among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	661	46,400	6.5	(3.4 - 9.6)	38,119	1,989,300	13.4	(11.0 - 15.8)
<b>Gender</b>								
Male	238	22,600	6.7	(1.7 - 11.7)	14,597	1,207,400	17.0	(12.9 - 21.1)
Female	423	23,800	6.3	(2.5 - 10.2)	23,522	781,900	10.1	(7.4 - 12.8)
<b>Age (years)</b>								
18-34	*	*	*	*	3,910	1,144,100	25.7	(19.3 - 32.2)
35-44	*	*	*	*	4,986	319,200	11.1	(6.9 - 15.3)
45-54	*	*	*	*	7,831	316,500	11.1	(6.8 - 15.4)
55-64	*	*	*	*	8,764	153,300	7.2	(4.5 - 9.8)
≥ 65	*	*	*	*	12,628	56,200	2.2	(1.2 - 3.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	587	36,600	6.3	(2.8 - 9.7)	35,811	727,100	7.8	(6.2 - 9.4)
Black Non-Hispanic	*	*	*	*	736	372,800	17.3	(9.9 - 24.7)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	264	28,100	10.0	(3.7 - 16.3)	15,673	1,134,400	20.3	(15.5 - 25.1)
Some College	*	*	*	*	9,906	531,400	15.0	(10.2 - 19.9)
≥ College Degree	*	*	*	*	12,459	266,300	4.7	(2.8 - 6.7)
<b>Income</b>								
≤ \$24,999	180	15,900	10.3	(4.0 - 16.7)	9,640	1,061,200	30.3	(23.2 - 37.5)
\$25,000-\$49,999	*	*	*	*	9,662	516,700	17.8	(11.7 - 23.9)
\$50,000-\$74,999	*	*	*	*	5,772	119,100	6.0	(3.3 - 8.8)
≥ \$75,000	*	*	*	*	8,567	111,100	2.5	(1.5 - 3.4)

<sup>1</sup> Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have no health insurance (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 8b. No health insurance among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	661	6.9	(3.6 - 10.3)	38,119	13.3	(10.9 - 15.6)
<b>Gender</b>						
Male	238	6.8	(1.7 - 11.9)	14,597	16.4	(12.5 - 20.3)
Female	423	7.1	(2.6 - 11.5)	23,522	10.4	(7.7 - 13.2)
<b>Race/Ethnicity</b>						
White Non-Hispanic	587	6.8	(3.0 - 10.6)	35,811	7.9	(6.3 - 9.5)
Black Non-Hispanic	*	*	*	736	17.5	(10.0 - 25.0)
Hispanic	*	*	*	860	31.3	(21.6 - 41.0)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	264	10.8	(3.9 - 17.6)	15,673	19.9	(15.3 - 24.6)
Some College	*	*	*	9,906	15.4	(10.6 - 20.3)
≥ College Degree	*	*	*	12,459	4.6	(2.8 - 6.4)
<b>Income</b>						
≤ \$24,999	180	11.1	(4.2 - 18.1)	9,640	29.8	(22.9 - 36.8)
\$25,000-\$49,999	*	*	*	9,662	18.1	(12.1 - 24.0)
\$50,000-\$74,999	*	*	*	5,772	6.0	(3.3 - 8.7)
≥ \$75,000	*	*	*	8,567	2.5	(1.5 - 3.4)

<sup>1</sup> Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 9a. No health insurance among adults aged 18-64<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	432	44,000	7.7	(3.9 - 11.6)	25,165	1,928,400	15.8	(12.9 - 18.7)
<b>Gender</b>								
Male	173	20,500	7.3	(1.4 - 13.2)	10,078	1,190,200	19.7	(15.0 - 24.5)
Female	259	23,500	8.1	(3.2 - 13.1)	15,087	738,200	12.0	(8.7 - 15.3)
<b>Age (years)</b>								
18-34	*	*	*	*	3,910	1,144,100	25.7	(19.3 - 32.2)
35-44	*	*	*	*	4,981	319,200	11.1	(6.9 - 15.3)
45-54	*	*	*	*	7,791	316,100	11.3	(7.0 - 15.7)
55-64	*	*	*	*	8,483	148,900	7.2	(4.5 - 9.9)
≥ 65	-	-	-	-	-	-	-	-
<b>Race/Ethnicity</b>								
White Non-Hispanic	369	34,200	7.6	(3.1 - 12.0)	23,354	703,100	9.7	(7.6 - 11.7)
Black Non-Hispanic	*	*	*	*	567	361,800	19.8	(11.3 - 28.4)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	141	27,200	13.5	(4.8 - 22.2)	9,037	1,087,700	25.4	(19.4 - 31.4)
Some College	*	*	*	*	7,031	522,800	17.1	(11.6 - 22.6)
≥ College Degree	*	*	*	*	9,068	261,200	5.5	(3.2 - 7.8)
<b>Income</b>								
≤ \$24,999	88	15,000	14.6	(5.1 - 24.1)	4,918	1,018,000	37.4	(28.8 - 45.9)
\$25,000-\$49,999	*	*	*	*	6,180	511,100	22.3	(14.9 - 29.7)
\$50,000-\$74,999	*	*	*	*	4,554	118,800	7.1	(3.9 - 10.3)
≥ \$75,000	*	*	*	*	7,324	110,700	2.7	(1.7 - 3.7)

<sup>1</sup> Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 18-64 who have no health insurance (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 9b. No health insurance among adults aged 18-64<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	432	8.0	(4.0 - 12.0)	25,165	15.7	(12.8 - 18.5)
<b>Gender</b>						
Male	173	7.5	(1.4 - 13.5)	10,078	19.5	(14.9 - 24.1)
Female	259	8.5	(3.1 - 13.8)	15,087	12.1	(8.8 - 15.4)
<b>Race/Ethnicity</b>						
White Non-Hispanic	369	7.9	(3.2 - 12.5)	23,354	9.7	(7.7 - 11.7)
Black Non-Hispanic	*	*	*	567	20.0	(11.4 - 28.7)
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	141	14.0	(5.0 - 23.0)	9,037	24.9	(19.1 - 30.7)
Some College	*	*	*	7,031	17.4	(11.9 - 22.9)
≥ College Degree	*	*	*	9,068	5.4	(3.2 - 7.5)
<b>Income</b>						
≤ \$24,999	88	15.1	(5.1 - 25.1)	4,918	36.8	(28.4 - 45.1)
\$25,000-\$49,999	*	*	*	6,180	22.5	(15.3 - 29.8)
\$50,000-\$74,999	*	*	*	4,554	7.0	(3.8 - 10.2)
≥ \$75,000	*	*	*	7,324	2.7	(1.7 - 3.7)

<sup>1</sup> Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 10a. Adults with regular health care provider<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	660	627,900	88.7	(84.3 - 93.0)	38,135	12,324,300	83.0	(80.4 - 85.6)
<b>Gender</b>								
Male	238	278,500	83.6	(75.7 - 91.6)	14,593	5,494,300	77.6	(73.2 - 82.0)
Female	422	349,500	93.2	(89.4 - 96.9)	23,542	6,830,000	87.8	(85.1 - 90.5)
<b>Age (years)</b>								
18-34	*	*	*	*	3,929	2,931,600	65.8	(59.2 - 72.5)
35-44	92	109,700	90.5	(84.4 - 96.6)	4,988	2,471,000	86.1	(81.9 - 90.3)
45-54	149	132,200	92.8	(88.3 - 97.4)	7,823	2,529,300	88.6	(84.6 - 92.7)
55-64	137	100,600	94.3	(88.2 - 100)	8,770	2,002,000	93.6	(91.7 - 95.4)
≥ 65	224	134,100	95.1	(91.6 - 98.6)	12,625	2,390,400	94.1	(92.2 - 96.0)
<b>Race/Ethnicity</b>								
White Non-Hispanic	586	530,100	91.2	(87.5 - 94.9)	35,821	8,227,700	88.2	(86.4 - 89.9)
Black Non-Hispanic	*	*	*	*	740	1,753,200	81.3	(72.6 - 89.9)
Hispanic	*	*	*	*	860	1,453,100	64.7	(54.7 - 74.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	264	235,800	83.6	(75.3 - 92.0)	15,692	4,249,200	75.9	(70.9 - 80.8)
Some College	176	169,500	88.0	(79.9 - 96.0)	9,904	2,935,000	82.8	(77.2 - 88.4)
≥ College Degree	217	221,100	95.3	(90.9 - 99.7)	12,456	5,046,100	89.8	(87.1 - 92.6)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,644	2,554,200	72.7	(65.9 - 79.6)
\$25,000-\$49,999	163	148,600	88.1	(79.5 - 96.7)	9,655	2,243,100	77.3	(70.0 - 84.6)
\$50,000-\$74,999	91	91,200	92.8	(86.8 - 98.8)	5,771	1,767,500	89.5	(85.2 - 93.9)
≥ \$75,000	135	187,800	91.8	(84.7 - 98.9)	8,560	4,122,800	91.9	(90.2 - 93.7)

<sup>1</sup> Defined as having one or more persons respondent thinks of as a personal doctor or health care provider

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have at least one person they think of as their personal doctor or health care provider (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 10b. Adults with regular health care provider<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	660	88.1	(83.5 - 92.7)	38,135	83.0	(80.4 - 85.5)
<b>Gender</b>						
Male	238	83.0	(74.7 - 91.3)	14,593	78.1	(73.9 - 82.3)
Female	422	92.7	(88.5 - 96.9)	23,542	87.4	(84.6 - 90.1)
<b>Race/Ethnicity</b>						
White Non-Hispanic	586	90.7	(86.8 - 94.6)	35,821	88.0	(86.2 - 89.7)
Black Non-Hispanic	*	*	*	740	81.4	(73.0 - 89.8)
Hispanic	*	*	*	860	65.2	(55.5 - 75.0)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	264	82.4	(73.3 - 91.5)	15,692	75.8	(71.0 - 80.6)
Some College	176	87.3	(78.9 - 95.8)	9,904	82.9	(77.5 - 88.3)
≥ College Degree	217	95.2	(90.8 - 99.7)	12,456	89.8	(87.1 - 92.4)
<b>Income</b>						
≤ \$24,999	*	*	*	9,644	72.9	(66.1 - 79.6)
\$25,000-\$49,999	163	87.5	(78.5 - 96.4)	9,655	77.4	(70.5 - 84.3)
\$50,000-\$74,999	91	92.5	(86.0 - 99.1)	5,771	89.6	(85.3 - 94.0)
≥ \$75,000	135	91.6	(84.3 - 98.9)	8,560	91.6	(89.8 - 93.4)

<sup>1</sup> Defined as having one or more persons respondent thinks of as a personal doctor or health care provider

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 11a. Cost prevented visit to doctor within the past year among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	661	83,500	11.7	(8.1 - 15.4)	38,143	2,034,800	13.7	(11.6 - 15.8)
<b>Gender</b>								
Male	238	36,400	10.8	(5.0 - 16.6)	14,607	903,300	12.7	(9.5 - 16.0)
Female	423	47,100	12.5	(7.9 - 17.2)	23,536	1,131,500	14.6	(11.8 - 17.4)
<b>Age (years)</b>								
18-34	*	*	*	*	3,939	947,300	21.2	(15.6 - 26.9)
35-44	92	20,700	17.1	(8.8 - 25.4)	4,987	347,900	12.1	(8.0 - 16.2)
45-54	149	17,000	11.9	(5.6 - 18.3)	7,825	382,200	13.5	(9.9 - 17.1)
55-64	137	11,700	11.0	(4.8 - 17.2)	8,766	236,800	11.1	(8.0 - 14.2)
≥ 65	*	*	*	*	12,626	120,600	4.8	(2.5 - 7.0)
<b>Race/Ethnicity</b>								
White Non-Hispanic	588	75,700	12.9	(8.7 - 17.1)	35,833	984,600	10.6	(8.9 - 12.2)
Black Non-Hispanic	*	*	*	*	738	280,000	13.1	(7.2 - 19.0)
Hispanic	*	*	*	*	859	597,900	26.5	(17.3 - 35.7)
Other Non-Hispanic	*	*	*	*	713	172,300	15.3	(5.7 - 24.9)
<b>Education</b>								
≤ High School	264	47,900	17.0	(9.7 - 24.4)	15,696	996,600	17.8	(13.8 - 21.9)
Some College	176	24,300	12.6	(5.4 - 19.7)	9,911	512,400	14.5	(10.4 - 18.6)
≥ College Degree	218	9,800	4.1	(1.6 - 6.7)	12,453	466,900	8.3	(6.0 - 10.6)
<b>Income</b>								
≤ \$24,999	179	26,600	17.6	(9.0 - 26.1)	9,639	854,500	24.3	(18.4 - 30.3)
\$25,000-\$49,999	163	29,100	17.3	(8.8 - 25.7)	9,656	532,900	18.4	(12.5 - 24.3)
\$50,000-\$74,999	*	*	*	*	5,776	202,700	10.3	(6.0 - 14.5)
≥ \$75,000	*	*	*	*	8,561	192,000	4.3	(2.9 - 5.6)

<sup>1</sup> Defined as needing to see a doctor within the past year but not being able to because of cost

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who did not see a doctor because of cost within the past year (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 11b. Cost prevented visit to doctor within the past year among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	661	12.5	(8.5 - 16.6)	38,143	13.8	(11.7 - 15.9)
<b>Gender</b>						
Male	238	11.0	(5.0 - 17.1)	14,607	12.5	(9.4 - 15.6)
Female	423	13.9	(8.5 - 19.3)	23,536	15.0	(12.1 - 17.9)
<b>Race/Ethnicity</b>						
White Non-Hispanic	588	13.9	(9.3 - 18.5)	35,833	10.8	(9.2 - 12.5)
Black Non-Hispanic	*	*	*	738	13.2	(7.2 - 19.3)
Hispanic	*	*	*	859	26.1	(17.2 - 35.1)
Other Non-Hispanic	*	*	*	713	14.8	(5.5 - 24.2)
<b>Education</b>						
≤ High School	264	18.2	(10.3 - 26.2)	15,696	17.8	(13.8 - 21.7)
Some College	176	14.4	(6.1 - 22.8)	9,911	15.3	(11.0 - 19.5)
≥ College Degree	218	4.3	(1.5 - 7.1)	12,453	8.2	(6.0 - 10.5)
<b>Income</b>						
≤ \$24,999	179	19.7	(9.9 - 29.6)	9,639	24.5	(18.6 - 30.4)
\$25,000-\$49,999	163	18.5	(9.2 - 27.8)	9,656	18.7	(12.9 - 24.6)
\$50,000-\$74,999	*	*	*	5,776	10.4	(6.1 - 14.7)
≥ \$75,000	*	*	*	8,561	4.4	(3.1 - 5.8)

<sup>1</sup> Defined as needing to see a doctor within the past year but not being able to because of cost

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 12a. Visited a doctor for routine checkup within the past year among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	660	530,900	74.5	(69.5 - 79.5)	38,026	10,809,300	72.9	(70.6 - 75.3)
<b>Gender</b>								
Male	239	228,700	67.7	(59.3 - 76.1)	14,592	4,738,400	66.9	(63.0 - 70.9)
Female	421	302,200	80.6	(75.1 - 86.1)	23,434	6,070,900	78.5	(75.9 - 81.0)
<b>Age (years)</b>								
18-34	*	*	*	*	3,921	2,747,400	61.7	(55.6 - 67.9)
35-44	*	*	*	*	4,973	1,883,100	65.5	(60.2 - 70.7)
45-54	149	110,700	77.7	(69.6 - 85.9)	7,808	2,174,100	76.6	(73.2 - 80.0)
55-64	136	83,100	78.2	(69.3 - 87.0)	8,751	1,728,600	81.4	(78.7 - 84.2)
≥ 65	224	132,600	94.0	(90.4 - 97.6)	12,573	2,276,000	90.0	(87.5 - 92.5)
<b>Race/Ethnicity</b>								
White Non-Hispanic	586	424,900	72.5	(67.1 - 78.0)	35,721	6,746,200	72.4	(70.6 - 74.2)
Black Non-Hispanic	*	*	*	*	736	1,688,900	78.8	(70.4 - 87.3)
Hispanic	*	*	*	*	857	1,550,900	69.3	(60.0 - 78.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	264	214,800	76.1	(68.3 - 83.9)	15,640	4,150,600	74.7	(70.7 - 78.7)
Some College	*	*	*	*	9,880	2,524,000	71.3	(66.1 - 76.5)
≥ College Degree	218	177,400	74.9	(66.7 - 83.2)	12,428	4,042,900	71.9	(68.5 - 75.3)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,616	2,469,200	70.4	(64.2 - 76.6)
\$25,000-\$49,999	*	*	*	*	9,631	2,068,400	71.3	(65.3 - 77.3)
\$50,000-\$74,999	*	*	*	*	5,765	1,466,000	74.8	(69.6 - 80.0)
≥ \$75,000	*	*	*	*	8,547	3,176,500	70.7	(67.3 - 74.2)

<sup>1</sup> Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past year

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have visited a doctor for a routine checkup within the past year (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 12b. Visited a doctor for routine checkup within the past year among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	660	73.3	(68.0 - 78.6)	38,026	72.7	(70.3 - 75.0)
<b>Gender</b>						
Male	239	67.4	(58.7 - 76.0)	14,592	67.1	(63.3 - 70.9)
Female	421	78.6	(72.4 - 84.9)	23,434	77.7	(75.1 - 80.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	586	71.0	(65.1 - 76.8)	35,721	71.9	(70.1 - 73.7)
Black Non-Hispanic	*	*	*	736	79.3	(71.3 - 87.4)
Hispanic	*	*	*	857	69.2	(60.1 - 78.2)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	264	75.2	(66.9 - 83.4)	15,640	74.4	(70.4 - 78.4)
Some College	*	*	*	9,880	70.9	(65.9 - 76.0)
≥ College Degree	218	74.3	(65.7 - 83.0)	12,428	71.7	(68.3 - 75.0)
<b>Income</b>						
≤ \$24,999	*	*	*	9,616	70.2	(64.1 - 76.2)
\$25,000-\$49,999	*	*	*	9,631	71.1	(65.3 - 76.9)
\$50,000-\$74,999	*	*	*	5,765	74.5	(69.2 - 79.8)
≥ \$75,000	*	*	*	8,547	70.3	(66.9 - 73.8)

<sup>1</sup> Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past year

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 13a. Visited a doctor for routine checkup within the past 2 years among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	660	619,700	86.9	(83.1 - 90.8)	38,026	12,726,600	85.9	(83.9 - 87.9)
<b>Gender</b>								
Male	239	283,600	84.0	(77.4 - 90.5)	14,592	5,791,900	81.8	(78.2 - 85.4)
Female	421	336,100	89.6	(85.3 - 94.0)	23,434	6,934,700	89.6	(87.9 - 91.3)
<b>Age (years)</b>								
18-34	*	*	*	*	3,921	3,535,400	79.4	(73.9 - 85.0)
35-44	92	94,500	78.0	(69.0 - 87.0)	4,973	2,355,600	81.9	(78.0 - 85.7)
45-54	149	130,300	91.5	(85.4 - 97.6)	7,808	2,507,900	88.4	(85.6 - 91.2)
55-64	136	97,800	92.0	(87.3 - 96.7)	8,751	1,924,800	90.7	(88.6 - 92.7)
≥ 65	224	134,500	95.3	(92.0 - 98.7)	12,573	2,402,900	95.0	(93.5 - 96.5)
<b>Race/Ethnicity</b>								
White Non-Hispanic	586	505,500	86.3	(82.0 - 90.6)	35,721	8,070,000	86.6	(85.4 - 87.9)
Black Non-Hispanic	*	*	*	*	736	1,914,500	89.4	(82.3 - 96.5)
Hispanic	*	*	*	*	857	1,832,800	81.9	(73.7 - 90.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	264	242,300	85.9	(79.2 - 92.6)	15,640	4,713,700	84.8	(81.2 - 88.4)
Some College	175	162,800	84.7	(77.7 - 91.6)	9,880	2,990,500	84.5	(79.6 - 89.4)
≥ College Degree	218	213,000	90.0	(83.6 - 96.3)	12,428	4,928,300	87.6	(85.3 - 89.9)
<b>Income</b>								
≤ \$24,999	179	139,400	91.1	(85.7 - 96.5)	9,616	2,911,700	83.0	(77.4 - 88.6)
\$25,000-\$49,999	163	145,300	86.1	(78.8 - 93.4)	9,631	2,375,500	81.9	(76.1 - 87.7)
\$50,000-\$74,999	91	86,700	88.2	(80.8 - 95.6)	5,765	1,733,500	88.4	(84.9 - 91.9)
≥ \$75,000	136	172,600	82.5	(72.7 - 92.2)	8,547	3,907,000	87.0	(84.8 - 89.2)

<sup>1</sup> Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past 2 years

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have visited a doctor for a routine checkup within the past 2 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Erie County**

**Table 13b. Visited a doctor for routine checkup within the past 2 years among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	660	86.0	(81.9 - 90.2)	38,026	85.7	(83.7 - 87.6)
<b>Gender</b>						
Male	239	83.2	(76.4 - 90.0)	14,592	81.9	(78.4 - 85.3)
Female	421	88.6	(83.6 - 93.5)	23,434	89.1	(87.4 - 90.9)
<b>Race/Ethnicity</b>						
White Non-Hispanic	586	85.1	(80.5 - 89.7)	35,721	86.2	(85.0 - 87.5)
Black Non-Hispanic	*	*	*	736	89.6	(83.0 - 96.3)
Hispanic	*	*	*	857	81.8	(73.8 - 89.8)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	264	85.1	(78.0 - 92.1)	15,640	84.6	(81.1 - 88.1)
Some College	175	82.9	(75.0 - 90.8)	9,880	84.4	(79.8 - 89.0)
≥ College Degree	218	89.5	(83.0 - 96.1)	12,428	87.3	(85.0 - 89.6)
<b>Income</b>						
≤ \$24,999	179	90.5	(84.5 - 96.5)	9,616	83.2	(77.8 - 88.5)
\$25,000-\$49,999	163	84.9	(76.9 - 93.0)	9,631	81.9	(76.3 - 87.4)
\$50,000-\$74,999	91	86.6	(77.9 - 95.3)	5,765	88.1	(84.4 - 91.8)
≥ \$75,000	136	82.3	(72.5 - 92.1)	8,547	86.4	(84.2 - 88.7)

<sup>1</sup> Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past 2 years

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 14a. Dentist visit within the past year among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	659	470,900	66.3	(60.8 - 71.7)	38,026	10,540,400	71.1	(68.6 - 73.6)
<b>Gender</b>								
Male	239	210,200	62.2	(53.2 - 71.2)	14,573	4,845,800	68.4	(64.5 - 72.2)
Female	420	260,700	70.0	(63.7 - 76.2)	23,453	5,694,600	73.6	(70.4 - 76.8)
<b>Age (years)</b>								
18-34	*	*	*	*	3,924	2,789,700	62.6	(56.3 - 69.0)
35-44	92	92,600	76.4	(67.4 - 85.5)	4,979	2,181,000	75.8	(70.9 - 80.7)
45-54	149	107,500	75.5	(66.0 - 85.0)	7,814	2,149,300	75.8	(71.9 - 79.7)
55-64	136	82,200	77.3	(68.9 - 85.7)	8,742	1,690,500	79.2	(75.9 - 82.6)
≥ 65	224	92,200	65.3	(58.1 - 72.5)	12,567	1,729,800	68.5	(64.7 - 72.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	586	393,100	67.1	(61.5 - 72.7)	35,721	6,959,900	74.7	(72.7 - 76.6)
Black Non-Hispanic	*	*	*	*	735	1,405,800	66.2	(57.9 - 74.5)
Hispanic	*	*	*	*	862	1,425,600	63.1	(53.7 - 72.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	263	162,600	58.1	(48.9 - 67.4)	15,613	3,620,700	65.1	(60.7 - 69.5)
Some College	*	*	*	*	9,880	2,504,700	70.8	(66.1 - 75.5)
≥ College Degree	218	184,600	78.0	(69.8 - 86.1)	12,453	4,402,400	78.2	(74.6 - 81.8)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,594	1,972,200	56.2	(49.8 - 62.6)
\$25,000-\$49,999	*	*	*	*	9,649	1,896,000	65.8	(59.5 - 72.1)
\$50,000-\$74,999	91	83,100	84.6	(75.7 - 93.5)	5,763	1,482,400	75.2	(69.4 - 81.0)
≥ \$75,000	*	*	*	*	8,564	3,707,900	82.4	(79.5 - 85.3)

<sup>1</sup> Defined as having visited a dentist or a dental clinic for any reason within the past year

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have visited a dentist or a dental clinic for any reason within the past year (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 14b. Dentist visit within the past year among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	659	65.8	(60.0 - 71.5)	38,026	71.1	(68.6 - 73.5)
<b>Gender</b>						
Male	239	61.7	(52.5 - 71.0)	14,573	68.3	(64.6 - 72.1)
Female	420	69.4	(62.5 - 76.4)	23,453	73.6	(70.4 - 76.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	586	66.3	(60.3 - 72.2)	35,721	74.4	(72.4 - 76.3)
Black Non-Hispanic	*	*	*	735	66.6	(58.3 - 74.9)
Hispanic	*	*	*	862	63.6	(54.4 - 72.9)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	263	58.3	(48.5 - 68.1)	15,613	65.3	(60.9 - 69.6)
Some College	*	*	*	9,880	70.4	(65.7 - 75.1)
≥ College Degree	218	77.0	(68.2 - 85.8)	12,453	78.1	(74.6 - 81.7)
<b>Income</b>						
≤ \$24,999	*	*	*	9,594	56.4	(50.1 - 62.8)
\$25,000-\$49,999	*	*	*	9,649	65.8	(59.6 - 72.0)
\$50,000-\$74,999	*	*	*	5,763	75.0	(69.1 - 80.8)
≥ \$75,000	*	*	*	8,564	82.1	(79.1 - 85.0)

<sup>1</sup> Defined as having visited a dentist or a dental clinic for any reason within the past year

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 15a. Had teeth cleaned within the past year among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	588	451,100	69.2	(63.6 - 74.8)	34,070	9,985,900	71.7	(69.1 - 74.3)
<b>Gender</b>								
Male	215	211,500	68.6	(59.4 - 77.8)	13,173	4,616,400	69.0	(65.0 - 73.1)
Female	373	239,600	69.7	(62.9 - 76.4)	20,897	5,369,500	74.2	(70.8 - 77.5)
<b>Age (years)</b>								
18-34	*	*	*	*	3,860	2,746,400	63.0	(56.5 - 69.5)
35-44	*	*	*	*	4,875	2,112,100	75.5	(70.7 - 80.3)
45-54	142	100,800	74.7	(65.2 - 84.1)	7,438	2,022,600	73.3	(69.1 - 77.4)
55-64	128	77,300	77.6	(68.8 - 86.3)	7,906	1,560,900	77.7	(73.8 - 81.6)
≥ 65	171	81,500	76.8	(69.2 - 84.3)	9,991	1,543,900	77.1	(73.7 - 80.5)
<b>Race/Ethnicity</b>								
White Non-Hispanic	520	386,000	72.5	(67.0 - 78.0)	31,982	6,601,500	75.4	(73.4 - 77.4)
Black Non-Hispanic	*	*	*	*	680	1,227,200	61.6	(52.1 - 71.0)
Hispanic	*	*	*	*	789	1,421,300	66.6	(57.1 - 76.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	*	*	*	*	12,740	3,388,600	67.7	(63.1 - 72.3)
Some College	162	121,800	67.7	(57.9 - 77.6)	9,145	2,286,800	68.5	(62.9 - 74.1)
≥ College Degree	210	182,400	78.6	(70.8 - 86.5)	12,128	4,298,500	78.3	(74.8 - 81.7)
<b>Income</b>								
≤ \$24,999	*	*	*	*	7,525	1,861,800	60.3	(53.5 - 67.1)
\$25,000-\$49,999	*	*	*	*	8,709	1,681,700	62.6	(55.4 - 69.8)
\$50,000-\$74,999	89	80,400	83.3	(74.1 - 92.4)	5,534	1,447,800	76.3	(70.7 - 81.8)
≥ \$75,000	*	*	*	*	8,448	3,617,500	80.8	(77.8 - 83.8)

<sup>1</sup> Defined as having had teeth cleaned by a dentist or dental hygienist within the past year

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have had their teeth cleaned by a dentist or dental hygienist within the past year (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 15b. Had teeth cleaned within the past year among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	588	68.0	(62.0 - 74.0)	34,070	71.7	(69.1 - 74.2)
<b>Gender</b>						
Male	215	67.9	(58.4 - 77.4)	13,173	69.2	(65.3 - 73.1)
Female	373	68.1	(60.6 - 75.5)	20,897	73.9	(70.5 - 77.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	520	71.1	(65.1 - 77.0)	31,982	75.1	(73.0 - 77.1)
Black Non-Hispanic	*	*	*	680	62.0	(52.7 - 71.4)
Hispanic	*	*	*	789	66.9	(57.5 - 76.2)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	12,740	67.7	(63.2 - 72.3)
Some College	*	*	*	9,145	68.1	(62.6 - 73.6)
≥ College Degree	210	77.4	(68.8 - 86.0)	12,128	78.3	(74.9 - 81.7)
<b>Income</b>						
≤ \$24,999	*	*	*	7,525	60.2	(53.4 - 67.0)
\$25,000-\$49,999	*	*	*	8,709	62.7	(55.7 - 69.7)
\$50,000-\$74,999	*	*	*	5,534	76.4	(70.9 - 81.9)
≥ \$75,000	*	*	*	8,448	80.4	(77.4 - 83.4)

<sup>1</sup> Defined as having had teeth cleaned by a dentist or dental hygienist within the past year

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 16a. Permanent teeth extracted due to decay or gum disease among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	655	362,200	51.1	(45.6 - 56.5)	37,652	7,426,500	50.4	(47.8 - 52.9)
<b>Gender</b>								
Male	238	162,200	48.1	(39.4 - 56.8)	14,457	3,367,000	47.8	(43.9 - 51.6)
Female	417	200,000	53.7	(47.2 - 60.3)	23,195	4,059,500	52.7	(49.5 - 56.0)
<b>Age (years)</b>								
18-34	59	35,000	17.3	(7.6 - 27.0)	3,921	1,121,600	25.4	(19.7 - 31.0)
35-44	*	*	*	*	4,964	1,250,000	43.6	(37.9 - 49.2)
45-54	148	81,500	57.7	(48.4 - 66.9)	7,780	1,595,300	56.1	(51.4 - 60.7)
55-64	135	71,400	67.9	(58.5 - 77.3)	8,648	1,382,800	65.5	(61.6 - 69.4)
≥ 65	221	119,200	85.2	(80.4 - 90.1)	12,339	2,076,900	83.0	(80.6 - 85.4)
<b>Race/Ethnicity</b>								
White Non-Hispanic	582	286,000	49.0	(43.5 - 54.6)	35,364	4,425,700	47.9	(46.0 - 49.9)
Black Non-Hispanic	*	*	*	*	727	1,244,700	58.2	(48.9 - 67.4)
Hispanic	*	*	*	*	853	1,220,500	54.2	(44.9 - 63.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	260	166,800	59.8	(50.0 - 69.5)	15,427	3,436,000	61.9	(57.2 - 66.6)
Some College	*	*	*	*	9,798	1,726,200	49.3	(44.3 - 54.2)
≥ College Degree	218	92,500	39.1	(30.7 - 47.5)	12,351	2,242,600	40.1	(36.5 - 43.6)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,483	2,171,900	62.3	(55.8 - 68.8)
\$25,000-\$49,999	*	*	*	*	9,552	1,577,400	54.7	(48.3 - 61.0)
\$50,000-\$74,999	*	*	*	*	5,744	1,000,700	51.5	(45.6 - 57.4)
≥ \$75,000	*	*	*	*	8,516	1,738,200	38.8	(35.3 - 42.2)

<sup>1</sup> Defined as having had at least one permanent tooth extracted because of tooth decay or gum disease

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have had at least one permanent tooth extracted because of tooth decay or gum disease (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 16b. Permanent teeth extracted due to decay or gum disease among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	655	48.4	(42.7 - 54.1)	37,652	49.4	(46.9 - 51.9)
<b>Gender</b>						
Male	238	47.3	(38.4 - 56.2)	14,457	47.9	(44.1 - 51.7)
Female	417	49.4	(42.2 - 56.5)	23,195	50.7	(47.4 - 54.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	582	46.3	(40.6 - 52.0)	35,364	46.5	(44.5 - 48.5)
Black Non-Hispanic	*	*	*	727	57.7	(48.6 - 66.9)
Hispanic	*	*	*	853	53.8	(44.6 - 62.9)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	15,427	61.3	(56.7 - 66.0)
Some College	*	*	*	9,798	47.9	(43.0 - 52.9)
≥ College Degree	218	36.8	(27.8 - 45.8)	12,351	39.0	(35.5 - 42.5)
<b>Income</b>						
≤ \$24,999	*	*	*	9,483	61.6	(55.2 - 68.1)
\$25,000-\$49,999	*	*	*	9,552	54.2	(48.0 - 60.3)
\$50,000-\$74,999	*	*	*	5,744	49.9	(44.0 - 55.9)
≥ \$75,000	*	*	*	8,516	37.7	(34.2 - 41.1)

<sup>1</sup> Defined as having had at least one permanent tooth extracted because of tooth decay or gum disease

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 17a. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	221	34,500	24.7	(18.3 - 31.1)	12,339	491,200	19.6	(16.0 - 23.3)
<b>Gender</b>								
Male	*	*	*	*	4,352	167,200	16.4	(11.8 - 20.9)
Female	157	18,800	22.4	(15.3 - 29.6)	7,987	324,000	21.8	(16.6 - 27.0)
<b>Age (years)</b>								
≥ 65	221	34,500	24.7	(18.3 - 31.1)	12,339	491,200	19.6	(16.0 - 23.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	212	33,400	25.6	(19.0 - 32.3)	11,885	322,600	16.8	(14.3 - 19.2)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	121	26,200	33.1	(23.6 - 42.6)	6,358	314,100	25.3	(19.9 - 30.7)
Some College	*	*	*	*	2,736	90,600	20.2	(12.1 - 28.2)
≥ College Degree	*	*	*	*	3,214	80,500	10.1	(3.6 - 16.5)
<b>Income</b>								
≤ \$24,999	91	10,200	20.7	(11.9 - 29.5)	4,558	199,600	26.9	(19.3 - 34.4)
\$25,000-\$49,999	*	*	*	*	3,365	123,900	21.4	(13.1 - 29.6)
\$50,000-\$74,999	*	*	*	*	1,179	21,400	7.3	(3.8 - 10.8)
≥ \$75,000	*	*	*	*	1,187	9,600	2.5	(1.0 - 4.0)

<sup>1</sup> Defined as having all permanent teeth extracted because of tooth decay or gum disease among adults aged 65 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 65 and older who have had all permanent teeth extracted because of tooth decay or gum disease (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 17b. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	221	25.1	(18.3 - 31.9)	12,339	19.4	(15.7 - 23.2)
<b>Gender</b>						
Male	*	*	*	4,352	16.5	(11.5 - 21.4)
Female	157	22.4	(15.2 - 29.5)	7,987	22.1	(16.6 - 27.7)
<b>Race/Ethnicity</b>						
White Non-Hispanic	212	25.9	(18.9 - 32.9)	11,885	16.6	(13.9 - 19.3)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	6,358	24.9	(19.3 - 30.5)
Some College	*	*	*	2,736	21.0	(12.1 - 29.9)
≥ College Degree	*	*	*	3,214	10.0	(3.6 - 16.3)
<b>Income</b>						
≤ \$24,999	91	20.1	(11.3 - 28.9)	4,558	27.3	(19.1 - 35.5)
\$25,000-\$49,999	*	*	*	3,365	21.0	(12.9 - 29.1)
\$50,000-\$74,999	*	*	*	1,179	7.6	(3.6 - 11.5)
≥ \$75,000	*	*	*	1,187	2.2	(0.9 - 3.5)

<sup>1</sup> Defined as having all permanent teeth extracted because of tooth decay or gum disease among adults aged 65 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 18a. Flu shot within the past 12 months<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	657	333,700	47.5	(42.2 - 52.8)	38,079	6,255,900	42.3	(39.9 - 44.8)
<b>Gender</b>								
Male	237	151,300	46.1	(37.4 - 54.8)	14,579	2,798,900	39.6	(35.8 - 43.5)
Female	420	182,400	48.7	(42.3 - 55.1)	23,500	3,456,900	44.8	(41.6 - 47.9)
<b>Age (years)</b>								
18-49	208	131,300	35.2	(26.5 - 43.8)	12,505	2,685,300	31.1	(27.3 - 34.9)
50-64	228	97,000	50.9	(43.3 - 58.5)	12,972	1,662,900	46.0	(42.6 - 49.5)
≥ 65	221	105,400	75.9	(69.4 - 82.4)	12,602	1,907,700	75.2	(72.0 - 78.4)
<b>Race/Ethnicity</b>								
White Non-Hispanic	583	263,300	45.7	(40.4 - 51.1)	35,770	4,029,000	43.3	(41.4 - 45.3)
Black Non-Hispanic	*	*	*	*	738	907,000	42.1	(33.1 - 51.1)
Hispanic	*	*	*	*	860	882,900	39.6	(30.4 - 48.9)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	261	144,900	53.0	(44.2 - 61.7)	15,663	2,457,800	44.3	(39.8 - 48.8)
Some College	176	78,300	40.6	(31.0 - 50.3)	9,898	1,377,700	38.9	(34.0 - 43.9)
≥ College Degree	217	109,500	46.6	(37.8 - 55.3)	12,436	2,404,400	43.0	(39.5 - 46.4)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,635	1,656,800	47.6	(41.2 - 54.0)
\$25,000-\$49,999	*	*	*	*	9,647	1,200,600	41.4	(35.6 - 47.3)
\$50,000-\$74,999	*	*	*	*	5,763	792,300	40.2	(34.3 - 46.2)
≥ \$75,000	*	*	*	*	8,548	1,776,100	39.6	(36.3 - 42.8)

<sup>1</sup> Defined as having had a flu shot within the past 12 months

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have had a flu shot within the past 12 months (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 18b. Flu shot within the past 12 months<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	657	45.6	(40.0 - 51.2)	38,079	41.7	(39.2 - 44.1)
<b>Gender</b>						
Male	237	45.9	(36.9 - 54.9)	14,579	40.0	(36.2 - 43.8)
Female	420	45.3	(38.3 - 52.3)	23,500	43.1	(40.0 - 46.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	583	43.8	(38.2 - 49.5)	35,770	42.4	(40.4 - 44.3)
Black Non-Hispanic	*	*	*	738	42.3	(33.3 - 51.3)
Hispanic	*	*	*	860	39.0	(29.9 - 48.1)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	261	51.5	(42.2 - 60.8)	15,663	43.7	(39.1 - 48.2)
Some College	176	38.4	(28.4 - 48.3)	9,898	37.8	(33.0 - 42.7)
≥ College Degree	217	44.8	(35.3 - 54.2)	12,436	42.5	(39.1 - 46.0)
<b>Income</b>						
≤ \$24,999	*	*	*	9,635	47.3	(41.0 - 53.7)
\$25,000-\$49,999	*	*	*	9,647	40.3	(34.5 - 46.0)
\$50,000-\$74,999	*	*	*	5,763	39.4	(33.4 - 45.4)
≥ \$75,000	*	*	*	8,548	39.1	(35.8 - 42.4)

<sup>1</sup> Defined as having had a flu shot within the past 12 months

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 19a. Flu vaccine that was sprayed in nose within the past 12 months<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	*	*	*	*	38,106	248,300	1.7	(0.6 - 2.8)
<b>Gender</b>								
Male	*	*	*	*	14,586	148,000	2.1	(0.0 - 4.2)
Female	*	*	*	*	23,520	100,300	1.3	(0.5 - 2.1)
<b>Age (years)</b>								
18-49	*	*	*	*	12,535	184,900	2.1	(0.3 - 3.9)
50-64	*	*	*	*	12,968	36,000	1.0	(0.2 - 1.8)
≥ 65	*	*	*	*	12,603	27,400	1.1	(0.4 - 1.8)
<b>Race/Ethnicity</b>								
White Non-Hispanic	*	*	*	*	35,796	66,000	0.7	(0.4 - 1.0)
Black Non-Hispanic	*	*	*	*	740	38,500	1.8	(0.0 - 4.1)
Hispanic	*	*	*	*	860	133,700	5.9	(0.0 - 12.3)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	*	*	*	*	15,679	172,500	3.1	(0.3 - 5.9)
Some College	*	*	*	*	9,897	43,500	1.2	(0.4 - 2.0)
≥ College Degree	*	*	*	*	12,449	31,600	0.6	(0.3 - 0.8)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,636	138,400	3.9	(0.0 - 8.1)
\$25,000-\$49,999	*	*	*	*	9,648	41,500	1.4	(0.0 - 3.1)
\$50,000-\$74,999	*	*	*	*	5,766	13,300	0.7	(0.2 - 1.2)
≥ \$75,000	*	*	*	*	8,561	35,300	0.8	(0.3 - 1.3)

<sup>1</sup> Defined as having had flu vaccine sprayed in nose within the past 12 months

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have had flu vaccine sprayed in nose within the past 12 months (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 19b. Flu vaccine that was sprayed in nose within the past 12 months<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	*	*	*	38,106	1.7	(0.6 - 2.7)
<b>Gender</b>						
Male	*	*	*	14,586	2.0	(0.1 - 3.9)
Female	*	*	*	23,520	1.4	(0.5 - 2.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	*	*	*	35,796	0.7	(0.4 - 1.0)
Black Non-Hispanic	*	*	*	740	1.9	(0.0 - 4.4)
Hispanic	*	*	*	860	5.9	(0.0 - 11.9)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	15,679	3.1	(0.4 - 5.8)
Some College	*	*	*	9,897	1.2	(0.4 - 2.0)
≥ College Degree	*	*	*	12,449	0.6	(0.3 - 0.8)
<b>Income</b>						
≤ \$24,999	*	*	*	9,636	3.8	(0.0 - 7.7)
\$25,000-\$49,999	*	*	*	9,648	1.5	(0.0 - 3.4)
\$50,000-\$74,999	*	*	*	5,766	0.7	(0.1 - 1.2)
≥ \$75,000	*	*	*	8,561	0.8	(0.3 - 1.3)

<sup>1</sup> Defined as having had flu vaccine sprayed in nose within the past 12 months

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Erie County**

**Table 20a. Flu shot or flu vaccine sprayed in nose within the past 12 months<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	657	334,300	47.6	(42.3 - 52.8)	38,059	6,302,300	42.7	(40.2 - 45.1)
<b>Gender</b>								
Male	237	151,900	46.3	(37.6 - 55.0)	14,568	2,822,700	40.0	(36.1 - 43.9)
Female	420	182,400	48.7	(42.3 - 55.1)	23,491	3,479,500	45.1	(42.0 - 48.2)
<b>Age (years)</b>								
18-49	208	131,800	35.3	(26.7 - 43.9)	12,501	2,726,400	31.6	(27.8 - 35.4)
50-64	228	97,000	50.9	(43.3 - 58.5)	12,964	1,664,200	46.1	(42.7 - 49.5)
≥ 65	221	105,400	75.9	(69.4 - 82.4)	12,594	1,911,700	75.4	(72.2 - 78.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	583	263,800	45.8	(40.5 - 51.2)	35,750	4,048,700	43.6	(41.7 - 45.5)
Black Non-Hispanic	*	*	*	*	738	910,000	42.2	(33.2 - 51.2)
Hispanic	*	*	*	*	860	903,300	40.6	(31.3 - 49.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	261	144,900	53.0	(44.2 - 61.7)	15,651	2,480,100	44.7	(40.2 - 49.3)
Some College	176	78,300	40.6	(31.0 - 50.3)	9,895	1,387,300	39.2	(34.3 - 44.1)
≥ College Degree	217	110,100	46.8	(38.0 - 55.6)	12,431	2,418,300	43.2	(39.8 - 46.7)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,626	1,675,100	48.2	(41.8 - 54.6)
\$25,000-\$49,999	*	*	*	*	9,643	1,208,300	41.7	(35.8 - 47.6)
\$50,000-\$74,999	*	*	*	*	5,761	795,600	40.4	(34.5 - 46.3)
≥ \$75,000	*	*	*	*	8,548	1,791,200	39.9	(36.6 - 43.2)

<sup>1</sup> Defined as having had a flu shot or flu vaccine sprayed in nose within the past 12 months

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have had a flu shot or flu vaccine sprayed in nose within the past 12 months (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 20b. Flu shot or flu vaccine sprayed in nose within the past 12 months<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	657	45.7	(40.1 - 51.3)	38,059	42.0	(39.6 - 44.5)
<b>Gender</b>						
Male	237	46.1	(37.2 - 55.1)	14,568	40.4	(36.6 - 44.2)
Female	420	45.3	(38.3 - 52.3)	23,491	43.5	(40.3 - 46.7)
<b>Race/Ethnicity</b>						
White Non-Hispanic	583	43.9	(38.3 - 49.6)	35,750	42.6	(40.7 - 44.6)
Black Non-Hispanic	*	*	*	738	42.4	(33.4 - 51.4)
Hispanic	*	*	*	860	40.0	(30.9 - 49.1)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	261	51.5	(42.2 - 60.8)	15,651	44.1	(39.6 - 48.6)
Some College	176	38.4	(28.4 - 48.3)	9,895	38.1	(33.2 - 43.0)
≥ College Degree	217	45.0	(35.6 - 54.5)	12,431	42.8	(39.4 - 46.3)
<b>Income</b>						
≤ \$24,999	*	*	*	9,626	48.0	(41.6 - 54.3)
\$25,000-\$49,999	*	*	*	9,643	40.5	(34.8 - 46.3)
\$50,000-\$74,999	*	*	*	5,761	39.6	(33.6 - 45.6)
≥ \$75,000	*	*	*	8,548	39.4	(36.1 - 42.7)

<sup>1</sup> Defined as having had a flu shot or flu vaccine sprayed in nose within the past 12 months

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 21a. Flu shot among adults aged 65 and older<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	221	105,400	75.9	(69.4 - 82.4)	12,602	1,907,700	75.2	(72.0 - 78.4)
<b>Gender</b>								
Male	*	*	*	*	4,417	772,900	74.6	(69.3 - 79.8)
Female	158	64,900	76.8	(69.5 - 84.2)	8,185	1,134,800	75.6	(71.6 - 79.6)
<b>Age (years)</b>								
≥ 65	221	105,400	75.9	(69.4 - 82.4)	12,602	1,907,700	75.2	(72.0 - 78.4)
<b>Race/Ethnicity</b>								
White Non-Hispanic	212	98,300	76.0	(69.4 - 82.5)	12,137	1,487,100	76.0	(73.5 - 78.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	120	59,400	75.3	(66.4 - 84.2)	6,518	930,800	74.0	(68.9 - 79.2)
Some College	*	*	*	*	2,790	317,800	69.7	(62.7 - 76.7)
≥ College Degree	*	*	*	*	3,261	654,000	80.5	(76.4 - 84.6)
<b>Income</b>								
≤ \$24,999	*	*	*	*	4,665	545,300	72.4	(65.6 - 79.3)
\$25,000-\$49,999	*	*	*	*	3,422	430,600	72.8	(67.0 - 78.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	1,198	301,000	77.5	(70.9 - 84.2)

<sup>1</sup> Defined as having had a flu shot during the past 12 months among adults aged 65 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 65 and older who have had a flu shot within the past 12 months (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 21b. Flu shot among adults aged 65 and older<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	221	75.7	(68.8 - 82.5)	12,602	75.0	(71.5 - 78.5)
<b>Gender</b>						
Male	*	*	*	4,417	74.4	(68.7 - 80.1)
Female	158	76.8	(69.5 - 84.2)	8,185	75.5	(71.3 - 79.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	212	76.0	(69.2 - 82.9)	12,137	75.9	(73.0 - 78.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	120	75.0	(65.6 - 84.5)	6,518	73.3	(67.5 - 79.1)
Some College	*	*	*	2,790	70.0	(62.7 - 77.2)
≥ College Degree	*	*	*	3,261	80.7	(76.2 - 85.1)
<b>Income</b>						
≤ \$24,999	*	*	*	4,665	72.6	(65.1 - 80.1)
\$25,000-\$49,999	*	*	*	3,422	71.4	(65.0 - 77.9)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	1,198	78.0	(70.8 - 85.1)

<sup>1</sup> Defined as having had a flu shot during the past 12 months among adults aged 65 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 22a. Pneumonia shot or pneumococcal vaccine<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	605	215,400	33.4	(28.3 - 38.5)	35,303	3,456,800	26.1	(23.9 - 28.3)
<b>Gender</b>								
Male	205	99,700	34.2	(25.2 - 43.2)	13,052	1,542,200	25.2	(21.5 - 28.8)
Female	400	115,700	32.7	(27.2 - 38.3)	22,251	1,914,600	26.9	(24.2 - 29.6)
<b>Age (years)</b>								
18-49	182	66,900	19.9	(11.9 - 27.9)	10,943	1,097,000	14.6	(11.3 - 17.9)
50-64	210	46,500	26.7	(19.6 - 33.8)	12,179	758,000	23.0	(19.8 - 26.2)
≥ 65	213	102,000	75.9	(69.1 - 82.6)	12,181	1,601,700	65.7	(61.7 - 69.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	535	180,500	34.4	(29.1 - 39.8)	33,225	2,358,700	28.7	(26.9 - 30.5)
Black Non-Hispanic	*	*	*	*	685	584,400	28.4	(19.5 - 37.3)
Hispanic	*	*	*	*	769	408,000	19.5	(12.6 - 26.4)
Other Non-Hispanic	*	*	*	*	624	105,600	12.2	(2.3 - 22.2)
<b>Education</b>								
≤ High School	247	106,800	41.2	(32.3 - 50.1)	14,700	1,535,300	30.4	(26.4 - 34.3)
Some College	160	55,800	32.3	(22.4 - 42.1)	9,151	836,200	27.2	(21.9 - 32.5)
≥ College Degree	196	52,000	24.6	(17.5 - 31.7)	11,375	1,070,700	21.4	(18.6 - 24.1)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,082	1,050,900	33.1	(27.6 - 38.6)
\$25,000-\$49,999	155	49,000	31.1	(21.6 - 40.6)	9,033	797,500	30.3	(24.1 - 36.6)
\$50,000-\$74,999	80	11,300	13.3	(6.1 - 20.5)	5,320	378,800	21.8	(17.4 - 26.1)
≥ \$75,000	*	*	*	*	7,773	648,700	16.3	(14.1 - 18.5)

<sup>1</sup> Defined as having ever had a pneumonia shot or pneumococcal vaccine

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have ever had a pneumonia shot or pneumococcal vaccine (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 22b. Pneumonia shot or pneumococcal vaccine<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	605	30.7	(25.5 - 35.9)	35,303	25.4	(23.2 - 27.6)
<b>Gender</b>						
Male	205	33.7	(24.6 - 42.9)	13,052	25.8	(22.3 - 29.4)
Female	400	28.2	(22.6 - 33.9)	22,251	25.1	(22.4 - 27.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	535	31.9	(26.3 - 37.4)	33,225	27.7	(25.9 - 29.5)
Black Non-Hispanic	*	*	*	685	27.9	(19.1 - 36.7)
Hispanic	*	*	*	769	19.6	(12.8 - 26.5)
Other Non-Hispanic	*	*	*	624	11.3	(2.0 - 20.7)
<b>Education</b>						
≤ High School	247	38.2	(29.0 - 47.4)	14,700	29.5	(25.6 - 33.5)
Some College	*	*	*	9,151	26.2	(21.0 - 31.3)
≥ College Degree	196	22.5	(15.2 - 29.8)	11,375	21.0	(18.3 - 23.8)
<b>Income</b>						
≤ \$24,999	*	*	*	9,082	32.3	(26.9 - 37.8)
\$25,000-\$49,999	155	28.2	(18.5 - 37.8)	9,033	29.0	(23.0 - 35.0)
\$50,000-\$74,999	80	12.3	(4.8 - 19.7)	5,320	21.2	(16.8 - 25.7)
≥ \$75,000	*	*	*	7,773	16.2	(13.9 - 18.5)

<sup>1</sup> Defined as having ever had a pneumonia shot or pneumococcal vaccine

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 23a. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	213	102,000	75.9	(69.1 - 82.6)	12,181	1,601,700	65.7	(61.7 - 69.7)
<b>Gender</b>								
Male	*	*	*	*	4,180	606,400	62.1	(55.5 - 68.7)
Female	155	65,000	78.3	(70.7 - 85.9)	8,001	995,300	68.0	(63.1 - 73.0)
<b>Age (years)</b>								
≥ 65	213	102,000	75.9	(69.1 - 82.6)	12,181	1,601,700	65.7	(61.7 - 69.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	204	93,900	75.2	(68.2 - 82.3)	11,735	1,281,300	68.2	(64.9 - 71.5)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	119	59,700	76.8	(68.2 - 85.5)	6,345	806,200	65.6	(59.5 - 71.7)
Some College	*	*	*	*	2,694	289,500	66.8	(60.1 - 73.4)
≥ College Degree	*	*	*	*	3,111	501,100	65.6	(58.5 - 72.7)
<b>Income</b>								
≤ \$24,999	*	*	*	*	4,533	497,100	67.7	(59.9 - 75.5)
\$25,000-\$49,999	*	*	*	*	3,324	344,300	61.5	(53.3 - 69.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	1,138	236,300	64.5	(55.2 - 73.8)

<sup>1</sup> Defined as having ever had a pneumonia shot or pneumococcal vaccine among adults aged 65 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 65 and older who have ever had a pneumonia shot or pneumococcal vaccine (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 23b. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	213	75.4	(68.3 - 82.5)	12,181	64.7	(60.4 - 69.1)
<b>Gender</b>						
Male	*	*	*	4,180	61.5	(54.3 - 68.6)
Female	155	78.2	(70.6 - 85.9)	8,001	67.6	(62.4 - 72.9)
<b>Race/Ethnicity</b>						
White Non-Hispanic	204	74.6	(67.2 - 82.1)	11,735	67.1	(63.2 - 70.9)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	119	76.7	(67.7 - 85.7)	6,345	64.2	(57.4 - 71.0)
Some College	*	*	*	2,694	65.8	(58.4 - 73.3)
≥ College Degree	*	*	*	3,111	65.3	(57.7 - 72.9)
<b>Income</b>						
≤ \$24,999	*	*	*	4,533	66.0	(57.0 - 74.9)
\$25,000-\$49,999	*	*	*	3,324	60.3	(51.9 - 68.6)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	*	*

<sup>1</sup> Defined as having ever had a pneumonia shot or pneumococcal vaccine among adults aged 65 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 24a. Ever had a mammogram among women aged 40 and older<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	361	232,400	93.9	(90.7 - 97.0)	19,491	4,328,800	91.8	(89.7 - 94.0)
<b>Gender</b>								
Male	-	-	-	-	-	-	-	-
Female	361	232,400	93.9	(90.7 - 97.0)	19,491	4,328,800	91.8	(89.7 - 94.0)
<b>Age (years)</b>								
40-49	61	56,500	89.7	(80.8 - 98.7)	3,851	1,287,500	85.0	(79.7 - 90.3)
50-64	141	97,000	96.8	(93.4 - 100)	7,452	1,641,700	96.0	(94.1 - 97.9)
≥ 65	159	78,900	93.5	(88.7 - 98.2)	8,188	1,399,600	94.1	(90.5 - 97.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	316	183,500	92.7	(88.9 - 96.6)	18,548	2,976,100	93.9	(92.7 - 95.1)
Black Non-Hispanic	*	*	*	*	342	746,600	93.8	(87.3 - 100)
Hispanic	*	*	*	*	333	447,600	91.8	(84.6 - 99.0)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	151	93,700	94.6	(90.2 - 99.0)	8,266	1,698,700	91.0	(87.4 - 94.5)
Some College	105	64,700	90.7	(83.5 - 98.0)	5,322	1,055,500	93.6	(90.2 - 97.0)
≥ College Degree	102	72,400	95.8	(90.9 - 100)	5,872	1,560,700	91.6	(87.5 - 95.6)
<b>Income</b>								
≤ \$24,999	119	61,700	93.0	(86.8 - 99.2)	5,691	1,023,400	93.5	(89.7 - 97.4)
\$25,000-\$49,999	81	49,600	96.0	(90.3 - 100)	4,857	860,300	89.5	(83.6 - 95.5)
\$50,000-\$74,999	*	*	*	*	2,648	547,100	94.6	(92.2 - 96.9)
≥ \$75,000	54	47,100	91.1	(81.4 - 100)	3,582	1,185,100	92.2	(88.2 - 96.1)

<sup>1</sup> Defined as having ever had a mammogram among women aged 40 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of women aged 40 and older who have ever had a mammogram (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 24b. Ever had a mammogram among women aged 40 and older<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	361	93.3	(89.5 - 97.2)	19,491	91.1	(88.7 - 93.6)
<b>Gender</b>						
Male	-	-	-	-	-	-
Female	361	93.3	(89.5 - 97.2)	19,491	91.1	(88.7 - 93.6)
<b>Race/Ethnicity</b>						
White Non-Hispanic	316	91.9	(87.2 - 96.7)	18,548	93.3	(92.0 - 94.7)
Black Non-Hispanic	*	*	*	342	93.1	(85.7 - 100)
Hispanic	*	*	*	333	91.0	(82.8 - 99.2)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	151	94.6	(89.8 - 99.3)	8,266	90.2	(86.1 - 94.2)
Some College	105	89.9	(81.3 - 98.6)	5,322	92.9	(88.9 - 96.9)
≥ College Degree	102	94.8	(87.9 - 100)	5,872	90.9	(86.6 - 95.3)
<b>Income</b>						
≤ \$24,999	119	92.3	(85.0 - 99.5)	5,691	93.0	(88.5 - 97.4)
\$25,000-\$49,999	81	96.0	(90.2 - 100)	4,857	88.8	(82.5 - 95.1)
\$50,000-\$74,999	*	*	*	2,648	94.3	(91.7 - 96.8)
≥ \$75,000	*	*	*	3,582	91.2	(86.8 - 95.6)

<sup>1</sup> Defined as having ever had a mammogram among women aged 40 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 25a. Had mammogram within the past 2 years among women aged 40 and older<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	358	213,300	87.0	(83.0 - 90.9)	19,376	3,760,000	80.3	(77.8 - 82.9)
<b>Gender</b>								
Male	-	-	-	-	-	-	-	-
Female	358	213,300	87.0	(83.0 - 90.9)	19,376	3,760,000	80.3	(77.8 - 82.9)
<b>Age (years)</b>								
40-49	59	52,000	85.2	(75.2 - 95.1)	3,842	1,122,700	74.4	(68.5 - 80.2)
50-64	140	91,600	91.7	(86.7 - 96.7)	7,428	1,487,900	87.2	(84.5 - 89.8)
≥ 65	159	69,700	82.6	(76.1 - 89.2)	8,106	1,149,400	78.5	(74.2 - 82.9)
<b>Race/Ethnicity</b>								
White Non-Hispanic	313	164,500	84.1	(79.3 - 88.8)	18,438	2,531,500	80.2	(78.3 - 82.2)
Black Non-Hispanic	*	*	*	*	339	667,100	85.6	(77.5 - 93.8)
Hispanic	*	*	*	*	332	419,700	86.1	(77.5 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	151	86,400	87.3	(81.5 - 93.1)	8,204	1,486,900	80.5	(76.4 - 84.6)
Some College	104	57,900	81.5	(72.8 - 90.2)	5,297	929,400	83.0	(78.8 - 87.1)
≥ College Degree	100	67,500	91.5	(85.3 - 97.8)	5,845	1,334,800	78.6	(73.9 - 83.2)
<b>Income</b>								
≤ \$24,999	118	53,300	80.7	(72.4 - 89.1)	5,650	882,300	80.8	(75.9 - 85.7)
\$25,000-\$49,999	81	46,700	90.3	(82.8 - 97.8)	4,846	732,300	76.3	(69.7 - 82.9)
\$50,000-\$74,999	*	*	*	*	2,641	494,200	85.6	(81.9 - 89.4)
≥ \$75,000	*	*	*	*	3,572	1,025,900	80.0	(75.2 - 84.8)

<sup>1</sup> Defined as having had a mammogram within the past 2 years among women aged 40 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of women aged 40 and older who have had a mammogram within the past 2 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 25b. Had mammogram within the past 2 years among women aged 40 and older<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	358	86.9	(82.4 - 91.3)	19,376	79.7	(76.9 - 82.5)
<b>Gender</b>						
Male	-	-	-	-	-	-
Female	358	86.9	(82.4 - 91.3)	19,376	79.7	(76.9 - 82.5)
<b>Race/Ethnicity</b>						
White Non-Hispanic	313	83.8	(78.4 - 89.2)	18,438	79.7	(77.6 - 81.8)
Black Non-Hispanic	*	*	*	339	85.3	(76.5 - 94.2)
Hispanic	*	*	*	332	85.2	(75.6 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	151	88.4	(82.7 - 94.2)	8,204	80.1	(75.6 - 84.7)
Some College	104	80.7	(70.9 - 90.6)	5,297	82.3	(77.7 - 86.9)
≥ College Degree	100	90.7	(82.7 - 98.6)	5,845	77.8	(72.9 - 82.8)
<b>Income</b>						
≤ \$24,999	118	80.8	(72.0 - 89.7)	5,650	80.8	(75.4 - 86.2)
\$25,000-\$49,999	81	89.5	(81.2 - 97.8)	4,846	75.7	(68.7 - 82.6)
\$50,000-\$74,999	*	*	*	2,641	85.5	(81.6 - 89.4)
≥ \$75,000	*	*	*	3,572	79.0	(73.8 - 84.2)

<sup>1</sup> Defined as having had a mammogram within the past 2 years among women aged 40 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 26a. Had mammogram within the past 2 years among women aged 50 and older<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	299	161,300	87.6	(83.5 - 91.7)	15,534	2,637,300	83.2	(80.7 - 85.7)
<b>Gender</b>								
Male	-	-	-	-	-	-	-	-
Female	299	161,300	87.6	(83.5 - 91.7)	15,534	2,637,300	83.2	(80.7 - 85.7)
<b>Age (years)</b>								
50-64	140	91,600	91.7	(86.7 - 96.7)	7,428	1,487,900	87.2	(84.5 - 89.8)
≥ 65	159	69,700	82.6	(76.1 - 89.2)	8,106	1,149,400	78.5	(74.2 - 82.9)
<b>Race/Ethnicity</b>								
White Non-Hispanic	263	127,300	85.2	(80.3 - 90.1)	14,887	1,907,000	82.3	(80.3 - 84.4)
Black Non-Hispanic	*	*	*	*	249	385,600	89.9	(83.2 - 96.7)
Hispanic	*	*	*	*	220	286,700	93.9	(89.1 - 98.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	133	65,400	85.1	(78.5 - 91.8)	7,026	1,094,500	83.3	(80.0 - 86.6)
Some College	86	45,500	84.4	(75.8 - 93.0)	4,124	646,700	87.3	(84.2 - 90.3)
≥ College Degree	77	48,900	94.1	(88.9 - 99.3)	4,356	890,500	80.6	(75.1 - 86.1)
<b>Income</b>								
≤ \$24,999	109	47,300	80.4	(71.7 - 89.0)	4,991	649,100	81.0	(76.8 - 85.2)
\$25,000-\$49,999	65	35,000	93.6	(85.6 - 100)	3,913	533,300	78.6	(70.9 - 86.2)
\$50,000-\$74,999	*	*	*	*	1,943	359,300	89.6	(86.3 - 93.0)
≥ \$75,000	*	*	*	*	2,303	601,300	84.6	(79.3 - 89.9)

<sup>1</sup> Defined as having had a mammogram within the past 2 years among women aged 50 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of women aged 50 and older who have had a mammogram within the past 2 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 26b. Had mammogram within the past 2 years among women aged 50 and older<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	299	87.9	(83.8 - 92.0)	15,534	83.1	(80.3 - 85.8)
<b>Gender</b>						
Male	-	-	-	-	-	-
Female	299	87.9	(83.8 - 92.0)	15,534	83.1	(80.3 - 85.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	263	85.4	(80.6 - 90.3)	14,887	82.1	(79.9 - 84.3)
Black Non-Hispanic	*	*	*	249	90.4	(83.6 - 97.2)
Hispanic	*	*	*	220	94.1	(89.3 - 99.0)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	133	85.8	(79.4 - 92.3)	7,026	83.6	(80.1 - 87.0)
Some College	86	84.4	(75.6 - 93.2)	4,124	87.3	(84.1 - 90.5)
≥ College Degree	77	94.1	(88.6 - 99.5)	4,356	79.9	(73.9 - 86.0)
<b>Income</b>						
≤ \$24,999	109	81.2	(72.8 - 89.6)	4,991	81.3	(76.9 - 85.6)
\$25,000-\$49,999	65	92.7	(83.7 - 100)	3,913	78.1	(69.9 - 86.4)
\$50,000-\$74,999	*	*	*	1,943	90.2	(87.1 - 93.4)
≥ \$75,000	*	*	*	2,303	83.9	(77.9 - 89.9)

<sup>1</sup> Defined as having had a mammogram within the past 2 years among women aged 50 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 27a. Ever had a Pap test among women<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	418	354,400	95.3	(91.2 - 99.3)	23,381	7,045,300	92.1	(90.0 - 94.3)
<b>Gender</b>								
Male	-	-	-	-	-	-	-	-
Female	418	354,400	95.3	(91.2 - 99.3)	23,381	7,045,300	92.1	(90.0 - 94.3)
<b>Age (years)</b>								
18-34	*	*	*	*	2,377	1,896,600	85.2	(79.3 - 91.0)
35-44	*	*	*	*	3,023	1,301,500	92.6	(87.7 - 97.5)
45-54	93	71,800	99.2	(97.6 - 100)	4,657	1,420,300	98.3	(97.0 - 99.6)
55-64	84	54,800	100	(100 - 100)	5,187	1,071,200	97.2	(95.0 - 99.3)
≥ 65	158	79,600	94.9	(90.3 - 99.4)	8,137	1,355,700	92.3	(88.5 - 96.1)
<b>Race/Ethnicity</b>								
White Non-Hispanic	363	286,000	97.8	(95.8 - 99.8)	21,976	4,439,700	94.2	(92.1 - 96.3)
Black Non-Hispanic	*	*	*	*	481	1,231,600	91.2	(84.0 - 98.4)
Hispanic	*	*	*	*	533	984,500	90.1	(85.3 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	165	120,400	94.1	(89.2 - 99.1)	9,526	2,540,500	88.3	(83.8 - 92.7)
Some College	117	97,500	97.1	(92.9 - 100)	6,462	1,782,200	93.5	(90.4 - 96.6)
≥ College Degree	134	135,500	94.9	(85.8 - 100)	7,345	2,643,200	95.0	(92.2 - 97.9)
<b>Income</b>								
≤ \$24,999	127	74,800	96.7	(92.5 - 100)	6,526	1,717,200	93.7	(90.9 - 96.6)
\$25,000-\$49,999	100	87,200	94.3	(88.2 - 100)	5,787	1,378,400	91.0	(85.6 - 96.4)
\$50,000-\$74,999	56	58,500	100	(100 - 100)	3,310	933,900	96.0	(91.8 - 100)
≥ \$75,000	66	86,500	100	(100 - 100)	4,570	1,943,300	95.4	(92.6 - 98.2)

<sup>1</sup> Defined as having ever had a Pap test among women

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of women who have ever had a Pap test (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 27b. Ever had a Pap test among women<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	418	94.7	(89.8 - 99.7)	23,381	91.8	(89.6 - 94.0)
<b>Gender</b>						
Male	-	-	-	-	-	-
Female	418	94.7	(89.8 - 99.7)	23,381	91.8	(89.6 - 94.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	363	98.0	(96.0 - 100)	21,976	94.0	(91.8 - 96.2)
Black Non-Hispanic	*	*	*	481	90.8	(83.5 - 98.1)
Hispanic	*	*	*	533	89.7	(84.6 - 94.7)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	165	94.0	(88.3 - 99.6)	9,526	87.8	(83.2 - 92.4)
Some College	117	96.9	(91.8 - 100)	6,462	93.1	(89.7 - 96.4)
≥ College Degree	*	*	*	7,345	94.9	(92.0 - 97.7)
<b>Income</b>						
≤ \$24,999	127	97.2	(93.7 - 100)	6,526	93.3	(90.1 - 96.5)
\$25,000-\$49,999	100	94.0	(87.2 - 100)	5,787	91.1	(85.7 - 96.5)
\$50,000-\$74,999	56	100	(100 - 100)	3,310	96.0	(91.9 - 100)
≥ \$75,000	66	100	(100 - 100)	4,570	95.0	(91.9 - 98.1)

<sup>1</sup> Defined as having ever had a Pap test among women

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 28a. Had a Pap test within the past 3 years among women<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	412	308,700	83.8	(78.9 - 88.7)	23,109	6,205,600	82.2	(79.8 - 84.6)
<b>Gender</b>								
Male	-	-	-	-	-	-	-	-
Female	412	308,700	83.8	(78.9 - 88.7)	23,109	6,205,600	82.2	(79.8 - 84.6)
<b>Age (years)</b>								
18-34	*	*	*	*	2,371	1,818,400	81.8	(75.5 - 88.1)
35-44	*	*	*	*	3,012	1,220,300	86.9	(81.8 - 92.0)
45-54	92	66,500	92.8	(87.1 - 98.5)	4,640	1,311,500	92.0	(89.8 - 94.3)
55-64	83	45,200	83.1	(74.8 - 91.4)	5,152	924,300	84.5	(81.0 - 88.0)
≥ 65	155	56,500	68.6	(60.1 - 77.1)	7,934	931,100	66.4	(61.8 - 71.0)
<b>Race/Ethnicity</b>								
White Non-Hispanic	357	246,500	85.3	(81.3 - 89.4)	21,720	3,807,100	81.5	(79.2 - 83.8)
Black Non-Hispanic	*	*	*	*	473	1,142,800	86.7	(79.1 - 94.3)
Hispanic	*	*	*	*	527	894,900	83.6	(76.3 - 91.0)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	162	98,700	78.1	(70.8 - 85.5)	9,384	2,132,600	76.0	(71.1 - 81.0)
Some College	116	86,000	86.0	(79.1 - 92.9)	6,401	1,594,300	84.0	(80.4 - 87.5)
≥ College Degree	132	122,900	87.2	(77.2 - 97.1)	7,281	2,406,000	86.9	(83.7 - 90.1)
<b>Income</b>								
≤ \$24,999	125	55,200	72.7	(63.5 - 81.9)	6,441	1,453,900	82.0	(77.9 - 86.0)
\$25,000-\$49,999	100	76,700	83.0	(74.2 - 91.7)	5,740	1,182,000	78.3	(72.6 - 83.9)
\$50,000-\$74,999	56	56,400	96.3	(92.1 - 100)	3,296	825,000	85.0	(78.0 - 91.9)
≥ \$75,000	65	79,400	92.5	(85.8 - 99.2)	4,555	1,810,200	89.0	(85.7 - 92.2)

<sup>1</sup> Defined as having had a Pap test within the past 3 years among women

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of women who have had a Pap test within the past 3 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 28b. Had a Pap test within the past 3 years among women<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	412	84.9	(79.4 - 90.5)	23,109	82.7	(80.2 - 85.1)
<b>Gender</b>						
Male	-	-	-	-	-	-
Female	412	84.9	(79.4 - 90.5)	23,109	82.7	(80.2 - 85.1)
<b>Race/Ethnicity</b>						
White Non-Hispanic	357	87.3	(83.2 - 91.4)	21,720	82.3	(80.0 - 84.6)
Black Non-Hispanic	*	*	*	473	87.0	(79.3 - 94.6)
Hispanic	*	*	*	527	83.4	(75.9 - 90.9)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	162	80.8	(73.3 - 88.2)	9,384	76.4	(71.3 - 81.5)
Some College	116	87.5	(80.4 - 94.5)	6,401	84.2	(80.5 - 88.0)
≥ College Degree	*	*	*	7,281	87.5	(84.3 - 90.6)
<b>Income</b>						
≤ \$24,999	125	75.7	(66.8 - 84.6)	6,441	82.7	(78.5 - 86.9)
\$25,000-\$49,999	100	84.0	(74.7 - 93.3)	5,740	79.2	(73.5 - 84.8)
\$50,000-\$74,999	56	97.3	(94.2 - 100)	3,296	85.7	(78.7 - 92.8)
≥ \$75,000	65	92.6	(85.4 - 99.7)	4,555	88.9	(85.5 - 92.3)

<sup>1</sup> Defined as having had a Pap test within the past 3 years among women

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 29a. Ever had a digital rectal exam among men aged 40 and older<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	195	159,500	75.4	(67.0 - 83.9)	12,033	3,176,500	73.8	(69.7 - 77.8)
<b>Gender</b>								
Male	195	159,500	75.4	(67.0 - 83.9)	12,033	3,176,500	73.8	(69.7 - 77.8)
Female	-	-	-	-	-	-	-	-
<b>Age (years)</b>								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	3,098	922,800	66.8	(59.9 - 73.8)
55-64	*	*	*	*	3,504	865,600	87.1	(82.5 - 91.7)
≥ 65	63	48,700	87.8	(78.5 - 97.0)	4,348	953,900	92.9	(91.0 - 94.9)
<b>Race/Ethnicity</b>								
White Non-Hispanic	180	142,900	80.5	(74.0 - 87.0)	11,411	2,452,100	80.3	(77.7 - 82.9)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	*	*	*	*	5,013	931,600	64.1	(57.3 - 71.0)
Some College	*	*	*	*	2,695	674,200	75.5	(67.8 - 83.2)
≥ College Degree	*	*	*	*	4,302	1,568,000	80.5	(74.0 - 87.0)
<b>Income</b>								
≤ \$24,999	*	*	*	*	2,573	573,300	65.9	(56.0 - 75.9)
\$25,000-\$49,999	*	*	*	*	3,168	530,100	74.6	(67.2 - 82.1)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,309	1,351,100	78.0	(72.6 - 83.5)

<sup>1</sup> Defined as having ever had a digital rectal exam among men aged 40 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of men aged 40 and older who have ever had a digital rectal exam (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 29b. Ever had a digital rectal exam among men aged 40 and older<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	195	73.3	(64.1 - 82.6)	12,033	73.7	(69.6 - 77.8)
<b>Gender</b>						
Male	195	73.3	(64.1 - 82.6)	12,033	73.7	(69.6 - 77.8)
Female	-	-	-	-	-	-
<b>Race/Ethnicity</b>						
White Non-Hispanic	180	78.6	(71.4 - 85.8)	11,411	79.9	(77.2 - 82.5)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	5,013	64.7	(57.8 - 71.7)
Some College	*	*	*	2,695	75.0	(67.1 - 82.9)
≥ College Degree	*	*	*	4,302	80.3	(73.8 - 86.8)
<b>Income</b>						
≤ \$24,999	*	*	*	2,573	66.9	(56.9 - 76.9)
\$25,000-\$49,999	*	*	*	3,168	74.8	(67.3 - 82.2)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,309	77.5	(72.0 - 83.0)

<sup>1</sup> Defined as having ever had a digital rectal exam among men aged 40 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 30a. Had digital rectal exam within the past 2 years among men aged 40 and older<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	193	115,400	55.0	(46.4 - 63.7)	11,923	2,380,300	55.8	(51.8 - 59.8)
<b>Gender</b>								
Male	193	115,400	55.0	(46.4 - 63.7)	11,923	2,380,300	55.8	(51.8 - 59.8)
Female	-	-	-	-	-	-	-	-
<b>Age (years)</b>								
40-44	*	*	*	*	1,078	246,800	27.5	(18.5 - 36.5)
45-54	*	*	*	*	3,074	650,000	47.6	(40.8 - 54.4)
55-64	*	*	*	*	3,467	673,500	68.1	(62.4 - 73.8)
≥ 65	*	*	*	*	4,304	809,900	79.7	(75.0 - 84.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	178	109,000	62.0	(54.1 - 69.9)	11,305	1,875,600	62.2	(59.2 - 65.2)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	*	*	*	*	4,963	718,700	49.6	(43.2 - 56.1)
Some College	*	*	*	*	2,665	511,500	58.5	(50.8 - 66.3)
≥ College Degree	*	*	*	*	4,277	1,148,400	59.4	(52.8 - 66.0)
<b>Income</b>								
≤ \$24,999	*	*	*	*	2,553	388,000	45.0	(35.0 - 55.0)
\$25,000-\$49,999	*	*	*	*	3,139	422,400	60.5	(53.0 - 68.0)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,295	978,300	56.9	(51.1 - 62.7)

<sup>1</sup> Defined as having had a digital rectal exam within the past 2 years among men aged 40 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of men aged 40 and older who have had a digital rectal exam within the past 2 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 30b. Had digital rectal exam within the past 2 years among men aged 40 and older<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	193	53.3	(44.4 - 62.3)	11,923	55.9	(51.7 - 60.0)
<b>Gender</b>						
Male	193	53.3	(44.4 - 62.3)	11,923	55.9	(51.7 - 60.0)
Female	-	-	-	-	-	-
<b>Race/Ethnicity</b>						
White Non-Hispanic	178	60.2	(51.9 - 68.5)	11,305	62.0	(58.9 - 65.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	4,963	50.5	(43.7 - 57.3)
Some College	*	*	*	2,665	58.3	(50.4 - 66.2)
≥ College Degree	*	*	*	4,277	59.2	(52.4 - 65.9)
<b>Income</b>						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,139	60.8	(53.1 - 68.4)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,295	56.5	(50.5 - 62.4)

<sup>1</sup> Defined as having had a digital rectal exam within the past 2 years among men aged 40 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 31a. Ever had a prostate specific antigen test among men aged 40 and older<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	187	126,900	62.1	(53.3 - 70.8)	11,557	2,822,600	67.6	(63.9 - 71.2)
<b>Gender</b>								
Male	187	126,900	62.1	(53.3 - 70.8)	11,557	2,822,600	67.6	(63.9 - 71.2)
Female	-	-	-	-	-	-	-	-
<b>Age (years)</b>								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,938	760,600	57.4	(50.6 - 64.3)
55-64	*	*	*	*	3,373	811,400	83.4	(78.8 - 88.1)
≥ 65	61	45,100	83.5	(73.6 - 93.5)	4,215	905,400	90.4	(87.1 - 93.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	172	108,900	63.8	(55.6 - 72.1)	10,950	1,965,400	67.0	(63.9 - 70.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	*	*	*	*	4,775	829,200	59.0	(52.4 - 65.6)
Some College	*	*	*	*	2,584	535,000	61.9	(53.7 - 70.2)
≥ College Degree	*	*	*	*	4,176	1,455,700	76.8	(72.5 - 81.1)
<b>Income</b>								
≤ \$24,999	*	*	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	*	3,048	444,100	64.1	(56.0 - 72.3)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,188	1,211,100	72.3	(67.6 - 77.0)

<sup>1</sup> Defined as having ever had a prostate specific antigen test among men aged 40 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of men aged 40 and older who have ever had a prostate specific antigen test (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 31b. Ever had a prostate specific antigen test among men aged 40 and older<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	187	61.5	(52.6 - 70.5)	11,557	67.4	(63.7 - 71.1)
<b>Gender</b>						
Male	187	61.5	(52.6 - 70.5)	11,557	67.4	(63.7 - 71.1)
Female	-	-	-	-	-	-
<b>Race/Ethnicity</b>						
White Non-Hispanic	172	62.3	(53.8 - 70.9)	10,950	66.6	(63.4 - 69.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	4,775	59.7	(52.9 - 66.4)
Some College	*	*	*	2,584	61.4	(53.0 - 69.9)
≥ College Degree	*	*	*	4,176	76.3	(71.9 - 80.7)
<b>Income</b>						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,048	64.2	(56.1 - 72.4)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,188	71.5	(66.6 - 76.3)

<sup>1</sup> Defined as having ever had a prostate specific antigen test among men aged 40 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 32a. Had prostate specific antigen test within the past 2 years among men aged 40 and older<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	186	113,700	55.9	(47.2 - 64.6)	11,437	2,493,800	59.9	(56.0 - 63.9)
<b>Gender</b>								
Male	186	113,700	55.9	(47.2 - 64.6)	11,437	2,493,800	59.9	(56.0 - 63.9)
Female	-	-	-	-	-	-	-	-
<b>Age (years)</b>								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,917	659,400	49.9	(42.8 - 57.0)
55-64	*	*	*	*	3,326	719,700	74.5	(69.4 - 79.7)
≥ 65	*	*	*	*	4,166	830,000	83.3	(78.4 - 88.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	172	99,900	58.6	(50.3 - 66.9)	10,836	1,762,200	60.4	(57.3 - 63.5)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	*	*	*	*	4,727	734,200	52.4	(45.7 - 59.1)
Some College	*	*	*	*	2,558	483,800	56.3	(48.2 - 64.4)
≥ College Degree	*	*	*	*	4,133	1,273,200	67.4	(61.6 - 73.3)
<b>Income</b>								
≤ \$24,999	*	*	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	*	3,021	410,800	59.6	(51.5 - 67.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,170	1,080,100	64.6	(59.1 - 70.0)

<sup>1</sup> Defined as having had a prostate specific antigen test within the past 2 years among men aged 40 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of men aged 40 and older who have had a prostate specific antigen test within the past 2 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 32b. Had prostate specific antigen test within the past 2 years among men aged 40 and older<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	186	55.5	(46.5 - 64.4)	11,437	59.8	(55.8 - 63.9)
<b>Gender</b>						
Male	186	55.5	(46.5 - 64.4)	11,437	59.8	(55.8 - 63.9)
Female	-	-	-	-	-	-
<b>Race/Ethnicity</b>						
White Non-Hispanic	172	56.9	(48.4 - 65.4)	10,836	60.1	(56.9 - 63.4)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	4,727	52.9	(45.9 - 59.9)
Some College	*	*	*	2,558	56.0	(47.7 - 64.2)
≥ College Degree	*	*	*	4,133	67.1	(61.2 - 73.1)
<b>Income</b>						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,021	59.8	(51.7 - 67.9)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,170	63.9	(58.3 - 69.6)

<sup>1</sup> Defined as having had a prostate specific antigen test within the past 2 years among men aged 40 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Erie County**

**Table 33a. Prostate cancer among men aged 40 and older<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	197	8,700	4.1	(1.6 - 6.7)	12,133	178,200	4.1	(3.0 - 5.2)
<b>Gender</b>								
Male	197	8,700	4.1	(1.6 - 6.7)	12,133	178,200	4.1	(3.0 - 5.2)
Female	-	-	-	-	-	-	-	-
<b>Age (years)</b>								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	3,120	6,100	0.4	(0.0 - 0.9)
55-64	*	*	*	*	3,532	31,600	3.1	(1.5 - 4.8)
≥ 65	*	*	*	*	4,388	140,400	13.6	(9.7 - 17.4)
<b>Race/Ethnicity</b>								
White Non-Hispanic	182	8,700	4.9	(1.9 - 7.9)	11,502	143,700	4.7	(3.5 - 5.8)
Black Non-Hispanic	*	*	*	*	184	23,500	5.1	(0.0 - 11.0)
Hispanic	*	*	*	*	197	7,700	1.6	(0.0 - 3.7)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	*	*	*	*	5,065	46,300	3.1	(2.2 - 4.1)
Some College	*	*	*	*	2,709	39,200	4.4	(1.3 - 7.5)
≥ College Degree	*	*	*	*	4,336	92,500	4.7	(3.0 - 6.5)
<b>Income</b>								
≤ \$24,999	*	*	*	*	2,597	35,800	4.1	(2.2 - 6.0)
\$25,000-\$49,999	*	*	*	*	3,196	29,600	4.1	(2.9 - 5.4)
\$50,000-\$74,999	*	*	*	*	2,007	20,300	2.8	(1.2 - 4.4)
≥ \$75,000	*	*	*	*	3,322	56,500	3.2	(1.7 - 4.8)

<sup>1</sup> Defined as having ever been told by a health professional that respondent had prostate cancer among men aged 40 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of men aged 40 and older who have ever been told they had prostate cancer (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 33b. Prostate cancer among men aged 40 and older<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	197	4.1	(1.5 - 6.7)	12,133	4.5	(3.2 - 5.7)
<b>Gender</b>						
Male	197	4.1	(1.5 - 6.7)	12,133	4.5	(3.2 - 5.7)
Female	-	-	-	-	-	-
<b>Race/Ethnicity</b>						
White Non-Hispanic	182	4.9	(1.9 - 8.0)	11,502	5.1	(3.8 - 6.4)
Black Non-Hispanic	*	*	*	184	5.7	(0.0 - 12.8)
Hispanic	*	*	*	197	1.8	(0.0 - 4.4)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	5,065	3.3	(2.3 - 4.3)
Some College	*	*	*	2,709	5.0	(1.1 - 9.0)
≥ College Degree	*	*	*	4,336	5.2	(3.1 - 7.2)
<b>Income</b>						
≤ \$24,999	*	*	*	2,597	4.5	(2.2 - 6.7)
\$25,000-\$49,999	*	*	*	3,196	4.3	(3.0 - 5.6)
\$50,000-\$74,999	*	*	*	2,007	3.0	(1.1 - 5.0)
≥ \$75,000	*	*	*	3,322	3.5	(1.6 - 5.4)

<sup>1</sup> Defined as having ever been told by a health professional that respondent had prostate cancer among men aged 40 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 34a. Home blood stool test ever used among adults aged 50 and older<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	447	132,800	40.5	(35.2 - 45.7)	25,322	2,288,500	38.0	(35.5 - 40.4)
<b>Gender</b>								
Male	147	46,400	32.4	(23.7 - 41.0)	9,571	1,081,900	38.9	(35.1 - 42.6)
Female	300	86,400	46.7	(40.3 - 53.1)	15,751	1,206,600	37.2	(34.0 - 40.4)
<b>Age (years)</b>								
50-64	226	70,800	37.6	(30.2 - 44.9)	12,834	1,192,000	34.0	(30.8 - 37.2)
≥ 65	219	61,600	44.5	(37.2 - 51.9)	12,472	1,093,400	43.6	(39.8 - 47.4)
<b>Race/Ethnicity</b>								
White Non-Hispanic	399	110,200	40.2	(34.9 - 45.6)	24,213	1,754,500	39.1	(37.1 - 41.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	198	53,800	37.6	(30.1 - 45.2)	11,146	823,500	34.9	(30.5 - 39.2)
Some College	*	*	*	*	6,261	480,300	37.0	(32.6 - 41.4)
≥ College Degree	129	39,700	41.5	(32.1 - 51.0)	7,851	973,700	41.6	(37.8 - 45.3)
<b>Income</b>								
≤ \$24,999	146	38,100	41.4	(32.0 - 50.8)	7,222	505,000	35.4	(29.7 - 41.2)
\$25,000-\$49,999	*	*	*	*	6,609	467,600	40.3	(35.2 - 45.4)
\$50,000-\$74,999	*	*	*	*	3,500	313,000	38.2	(32.6 - 43.9)
≥ \$75,000	*	*	*	*	4,647	657,800	37.3	(33.0 - 41.6)

<sup>1</sup> Defined as having ever used a home blood stool test among adults aged 50 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 50 and older who have ever used a home blood stool test (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 34b. Home blood stool test ever used among adults aged 50 and older<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	447	39.6	(34.3 - 44.9)	25,322	38.1	(35.5 - 40.8)
<b>Gender</b>						
Male	147	32.0	(23.5 - 40.5)	9,571	39.4	(35.3 - 43.5)
Female	300	46.1	(39.6 - 52.6)	15,751	36.9	(33.5 - 40.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	399	39.4	(34.0 - 44.9)	24,213	38.9	(36.7 - 41.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	198	37.1	(29.3 - 44.8)	11,146	35.1	(30.3 - 39.9)
Some College	*	*	*	6,261	36.6	(32.0 - 41.3)
≥ College Degree	129	40.6	(31.1 - 50.0)	7,851	41.9	(37.9 - 46.0)
<b>Income</b>						
≤ \$24,999	146	40.1	(30.6 - 49.7)	7,222	35.9	(29.5 - 42.4)
\$25,000-\$49,999	*	*	*	6,609	40.2	(34.8 - 45.6)
\$50,000-\$74,999	*	*	*	3,500	38.1	(31.9 - 44.3)
≥ \$75,000	*	*	*	4,647	37.5	(32.9 - 42.2)

<sup>1</sup> Defined as having ever used a home blood stool test among adults aged 50 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 35a. Home blood stool test used within the past year among adults aged 50 and older<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	444	42,300	13.0	(9.5 - 16.4)	25,074	863,300	14.4	(12.6 - 16.2)
<b>Gender</b>								
Male	146	16,300	11.5	(6.3 - 16.6)	9,508	444,400	16.0	(13.2 - 18.9)
Female	298	26,000	14.1	(9.4 - 18.8)	15,566	418,900	13.1	(10.8 - 15.4)
<b>Age (years)</b>								
50-64	225	19,600	10.4	(6.2 - 14.7)	12,759	454,400	13.0	(10.6 - 15.5)
≥ 65	217	22,700	16.6	(10.8 - 22.3)	12,299	406,400	16.4	(13.7 - 19.0)
<b>Race/Ethnicity</b>								
White Non-Hispanic	396	34,700	12.8	(9.1 - 16.4)	23,971	634,300	14.3	(12.8 - 15.7)
Black Non-Hispanic	*	*	*	*	396	136,000	18.6	(9.4 - 27.8)
Hispanic	*	*	*	*	349	53,400	9.9	(2.5 - 17.2)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	195	14,700	10.4	(5.6 - 15.3)	11,026	288,600	12.3	(9.4 - 15.3)
Some College	117	15,500	17.5	(9.6 - 25.4)	6,208	212,500	16.4	(12.5 - 20.4)
≥ College Degree	129	12,100	12.7	(6.8 - 18.6)	7,778	352,400	15.2	(12.5 - 17.9)
<b>Income</b>								
≤ \$24,999	145	16,800	18.4	(10.6 - 26.2)	7,157	196,300	13.9	(10.1 - 17.6)
\$25,000-\$49,999	109	11,300	14.0	(6.8 - 21.2)	6,562	221,500	19.2	(14.1 - 24.3)
\$50,000-\$74,999	*	*	*	*	3,469	111,900	13.7	(10.1 - 17.4)
≥ \$75,000	*	*	*	*	4,623	205,100	11.7	(8.8 - 14.6)

<sup>1</sup> Defined as having used a home blood stool test within the past year among adults aged 50 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 50 and older who have used a home blood stool test within the past year (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 35b. Home blood stool test used within the past year among adults aged 50 and older<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	444	13.0	(9.4 - 16.5)	25,074	14.6	(12.6 - 16.7)
<b>Gender</b>						
Male	146	11.9	(6.5 - 17.3)	9,508	16.4	(13.2 - 19.6)
Female	298	13.9	(9.2 - 18.6)	15,566	12.9	(10.4 - 15.4)
<b>Race/Ethnicity</b>						
White Non-Hispanic	396	12.9	(9.1 - 16.7)	23,971	14.2	(12.7 - 15.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	349	10.2	(1.7 - 18.7)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	195	10.3	(5.3 - 15.3)	11,026	12.7	(9.2 - 16.1)
Some College	117	17.5	(9.3 - 25.6)	6,208	16.6	(12.3 - 20.9)
≥ College Degree	129	13.0	(6.9 - 19.0)	7,778	15.3	(12.4 - 18.3)
<b>Income</b>						
≤ \$24,999	145	18.5	(10.3 - 26.6)	7,157	14.3	(9.8 - 18.7)
\$25,000-\$49,999	109	14.1	(6.7 - 21.5)	6,562	19.3	(13.8 - 24.8)
\$50,000-\$74,999	*	*	*	3,469	13.8	(9.8 - 17.7)
≥ \$75,000	*	*	*	4,623	11.8	(8.6 - 15.0)

<sup>1</sup> Defined as having used a home blood stool test within the past year among adults aged 50 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 36a. Home blood stool test used within the past 2 years among adults aged 50 and older<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	444	62,300	19.1	(15.0 - 23.1)	25,074	1,218,000	20.4	(18.3 - 22.5)
<b>Gender</b>								
Male	146	24,200	17.0	(10.7 - 23.3)	9,508	593,100	21.4	(18.3 - 24.5)
Female	298	38,100	20.7	(15.4 - 25.9)	15,566	624,900	19.5	(16.6 - 22.4)
<b>Age (years)</b>								
50-64	225	30,000	16.0	(10.8 - 21.1)	12,759	629,900	18.1	(15.4 - 20.7)
≥ 65	217	32,200	23.5	(17.1 - 29.9)	12,299	585,500	23.6	(20.2 - 27.0)
<b>Race/Ethnicity</b>								
White Non-Hispanic	396	52,200	19.2	(14.9 - 23.5)	23,971	877,500	19.7	(18.2 - 21.3)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	195	23,900	16.9	(11.2 - 22.6)	11,026	454,600	19.4	(15.6 - 23.3)
Some College	117	17,900	20.3	(12.0 - 28.5)	6,208	262,200	20.3	(16.3 - 24.2)
≥ College Degree	129	20,500	21.5	(13.6 - 29.3)	7,778	491,400	21.2	(18.1 - 24.2)
<b>Income</b>								
≤ \$24,999	145	24,300	26.6	(18.1 - 35.1)	7,157	263,900	18.6	(14.6 - 22.7)
\$25,000-\$49,999	109	14,900	18.4	(10.2 - 26.6)	6,562	285,900	24.8	(19.6 - 30.0)
\$50,000-\$74,999	53	7,000	16.4	(6.9 - 25.8)	3,469	167,500	20.6	(15.6 - 25.5)
≥ \$75,000	*	*	*	*	4,623	294,900	16.8	(13.7 - 20.0)

<sup>1</sup> Defined as having used a home blood stool test within the past 2 years among adults aged 50 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 50 and older who have used a home blood stool test within the past 2 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 36b. Home blood stool test used within the past 2 years among adults aged 50 and older<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	444	18.8	(14.7 - 22.9)	25,074	20.5	(18.2 - 22.8)
<b>Gender</b>						
Male	146	17.2	(10.8 - 23.6)	9,508	21.7	(18.3 - 25.1)
Female	298	20.2	(15.0 - 25.5)	15,566	19.4	(16.3 - 22.5)
<b>Race/Ethnicity</b>						
White Non-Hispanic	396	19.0	(14.7 - 23.4)	23,971	19.7	(18.0 - 21.3)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	195	16.5	(10.8 - 22.3)	11,026	19.5	(15.4 - 23.7)
Some College	117	20.0	(11.5 - 28.5)	6,208	20.4	(16.1 - 24.7)
≥ College Degree	129	21.4	(13.6 - 29.3)	7,778	21.4	(18.1 - 24.7)
<b>Income</b>						
≤ \$24,999	145	26.1	(17.4 - 34.9)	7,157	18.9	(14.2 - 23.6)
\$25,000-\$49,999	109	18.4	(10.0 - 26.8)	6,562	24.8	(19.3 - 30.3)
\$50,000-\$74,999	53	16.3	(6.9 - 25.7)	3,469	20.6	(15.3 - 25.9)
≥ \$75,000	*	*	*	4,623	16.9	(13.4 - 20.4)

<sup>1</sup> Defined as having used a home blood stool test within the past 2 years among adults aged 50 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Erie County**

**Table 37a. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	450	225,200	68.4	(63.2 - 73.6)	25,357	4,019,100	66.6	(64.1 - 69.1)
<b>Gender</b>								
Male	147	95,300	66.4	(57.4 - 75.4)	9,582	1,894,100	68.0	(64.2 - 71.7)
Female	303	129,900	69.9	(63.9 - 75.9)	15,775	2,125,000	65.5	(62.1 - 68.8)
<b>Age (years)</b>								
50-54	*	*	*	*	4,179	718,900	50.6	(44.6 - 56.5)
55-64	136	77,800	73.7	(64.3 - 83.1)	8,695	1,493,500	71.0	(67.3 - 74.8)
≥ 65	222	97,600	70.0	(63.0 - 77.0)	12,466	1,802,000	72.1	(68.6 - 75.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	402	193,800	70.5	(65.4 - 75.6)	24,248	3,093,900	68.8	(66.8 - 70.8)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	199	93,800	65.5	(57.2 - 73.7)	11,121	1,445,100	61.2	(56.7 - 65.8)
Some College	119	58,800	66.2	(56.5 - 76.0)	6,287	835,300	64.3	(59.7 - 69.0)
≥ College Degree	129	72,500	75.8	(66.8 - 84.9)	7,884	1,732,400	73.7	(70.0 - 77.4)
<b>Income</b>								
≤ \$24,999	148	56,000	60.5	(50.7 - 70.4)	7,220	892,600	62.7	(56.8 - 68.6)
\$25,000-\$49,999	*	*	*	*	6,619	697,300	60.1	(54.6 - 65.7)
\$50,000-\$74,999	*	*	*	*	3,514	613,700	74.2	(68.9 - 79.6)
≥ \$75,000	*	*	*	*	4,665	1,294,400	73.3	(69.2 - 77.5)

<sup>1</sup> Defined as having ever had a sigmoidoscopy or colonoscopy among adults aged 50 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 50 and older who have ever had a sigmoidoscopy or colonoscopy (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 37b. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	450	67.9	(62.5 - 73.2)	25,357	66.6	(63.9 - 69.3)
<b>Gender</b>						
Male	147	66.1	(57.1 - 75.1)	9,582	68.5	(64.5 - 72.5)
Female	303	69.4	(63.3 - 75.6)	15,775	64.8	(61.2 - 68.4)
<b>Race/Ethnicity</b>						
White Non-Hispanic	402	70.2	(64.8 - 75.5)	24,248	68.7	(66.5 - 70.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	199	64.8	(56.3 - 73.4)	11,121	61.3	(56.4 - 66.2)
Some College	*	*	*	6,287	63.8	(58.8 - 68.8)
≥ College Degree	129	75.9	(66.8 - 85.0)	7,884	74.0	(70.2 - 77.9)
<b>Income</b>						
≤ \$24,999	*	*	*	7,220	63.0	(56.7 - 69.3)
\$25,000-\$49,999	*	*	*	6,619	59.7	(53.8 - 65.6)
\$50,000-\$74,999	*	*	*	3,514	74.4	(68.3 - 80.5)
≥ \$75,000	*	*	*	4,665	73.6	(69.3 - 78.0)

<sup>1</sup> Defined as having ever had a sigmoidoscopy or colonoscopy among adults aged 50 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 38a. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	448	216,900	66.1	(60.9 - 71.4)	25,169	3,855,800	64.6	(62.1 - 67.1)
<b>Gender</b>								
Male	145	89,900	63.3	(54.1 - 72.5)	9,521	1,835,400	66.4	(62.6 - 70.1)
Female	303	127,000	68.3	(62.2 - 74.3)	15,648	2,020,400	63.1	(59.8 - 66.4)
<b>Age (years)</b>								
50-54	*	*	*	*	4,169	700,500	49.3	(43.3 - 55.3)
55-64	135	74,600	70.9	(61.3 - 80.6)	8,652	1,455,100	69.4	(65.7 - 73.2)
≥ 65	221	94,200	68.0	(60.8 - 75.1)	12,331	1,695,700	69.5	(65.9 - 73.1)
<b>Race/Ethnicity</b>								
White Non-Hispanic	400	185,600	67.8	(62.5 - 73.1)	24,069	2,981,500	66.6	(64.6 - 68.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	198	92,100	64.5	(56.3 - 72.8)	11,017	1,360,400	59.0	(54.5 - 63.5)
Some College	*	*	*	*	6,251	807,100	62.4	(57.8 - 67.1)
≥ College Degree	128	68,300	72.0	(62.6 - 81.4)	7,841	1,682,500	71.7	(68.0 - 75.4)
<b>Income</b>								
≤ \$24,999	147	55,000	59.8	(49.9 - 69.6)	7,163	858,900	61.0	(55.1 - 67.0)
\$25,000-\$49,999	*	*	*	*	6,584	664,900	57.7	(52.2 - 63.2)
\$50,000-\$74,999	*	*	*	*	3,501	602,600	73.1	(67.7 - 78.5)
≥ \$75,000	*	*	*	*	4,655	1,260,400	71.5	(67.3 - 75.7)

<sup>1</sup> Defined as having had a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 50 and older who have had a sigmoidoscopy or colonoscopy within the past 10 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 38b. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	448	65.8	(60.4 - 71.2)	25,169	64.6	(61.9 - 67.3)
<b>Gender</b>						
Male	145	63.2	(54.0 - 72.4)	9,521	66.9	(62.9 - 70.9)
Female	303	68.0	(61.8 - 74.2)	15,648	62.4	(58.9 - 66.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	400	67.6	(62.2 - 73.1)	24,069	66.5	(64.3 - 68.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	198	64.0	(55.4 - 72.5)	11,017	59.0	(54.1 - 63.9)
Some College	*	*	*	6,251	61.9	(56.9 - 66.9)
≥ College Degree	128	72.4	(62.9 - 81.8)	7,841	72.1	(68.2 - 75.9)
<b>Income</b>						
≤ \$24,999	*	*	*	7,163	61.3	(54.9 - 67.6)
\$25,000-\$49,999	*	*	*	6,584	57.2	(51.4 - 63.0)
\$50,000-\$74,999	*	*	*	3,501	73.3	(67.2 - 79.4)
≥ \$75,000	*	*	*	4,655	71.8	(67.4 - 76.2)

<sup>1</sup> Defined as having had a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 39a. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	440	224,400	69.4	(64.1 - 74.6)	24,792	4,049,200	69.1	(66.7 - 71.6)
<b>Gender</b>								
Male	143	93,000	66.4	(57.2 - 75.6)	9,418	1,945,200	70.7	(67.0 - 74.3)
Female	297	131,500	71.6	(65.7 - 77.6)	15,374	2,104,100	67.8	(64.6 - 71.1)
<b>Age (years)</b>								
50-54	*	*	*	*	4,132	751,200	54.1	(48.0 - 60.1)
55-64	132	76,400	74.1	(64.4 - 83.8)	8,366	1,487,600	73.7	(70.0 - 77.4)
≥ 65	218	100,200	72.9	(66.1 - 79.6)	12,294	1,810,500	74.0	(70.6 - 77.4)
<b>Race/Ethnicity</b>								
White Non-Hispanic	394	193,100	71.3	(66.2 - 76.5)	23,726	3,091,400	70.4	(68.4 - 72.4)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	195	94,500	67.0	(58.7 - 75.3)	10,872	1,461,600	64.2	(59.8 - 68.7)
Some College	117	60,500	69.2	(59.7 - 78.8)	6,170	873,300	68.4	(63.9 - 72.8)
≥ College Degree	125	69,400	74.2	(64.8 - 83.6)	7,708	1,709,000	74.6	(70.9 - 78.3)
<b>Income</b>								
≤ \$24,999	144	60,900	67.4	(57.7 - 77.1)	7,092	923,300	66.0	(60.1 - 71.9)
\$25,000-\$49,999	*	*	*	*	6,526	730,000	64.2	(58.9 - 69.6)
\$50,000-\$74,999	*	*	*	*	3,467	621,500	76.6	(71.4 - 81.9)
≥ \$75,000	*	*	*	*	4,615	1,292,500	73.8	(69.8 - 77.9)

<sup>1</sup> Defined as having used a home blood stool test within the past year or a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 years and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 50 and older who have used a home blood stool test in the past year or sigmoidoscopy or colonoscopy within the past 10 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009  
Erie County**

**Table 39b. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	440	68.9	(63.5 - 74.3)	24,792	69.3	(66.7 - 71.9)
<b>Gender</b>						
Male	143	66.3	(57.2 - 75.5)	9,418	71.5	(67.6 - 75.4)
Female	297	71.1	(64.9 - 77.2)	15,374	67.2	(63.7 - 70.7)
<b>Race/Ethnicity</b>						
White Non-Hispanic	394	71.0	(65.6 - 76.3)	23,726	70.3	(68.1 - 72.4)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	195	66.2	(57.6 - 74.7)	10,872	64.5	(59.8 - 69.3)
Some College	117	68.5	(58.6 - 78.4)	6,170	68.2	(63.4 - 72.9)
≥ College Degree	125	74.4	(64.9 - 83.9)	7,708	75.0	(71.2 - 78.8)
<b>Income</b>						
≤ \$24,999	*	*	*	7,092	66.4	(60.2 - 72.6)
\$25,000-\$49,999	*	*	*	6,526	63.8	(58.2 - 69.5)
\$50,000-\$74,999	*	*	*	3,467	76.7	(70.6 - 82.7)
≥ \$75,000	*	*	*	4,615	74.4	(70.2 - 78.6)

<sup>1</sup> Defined as having used a home blood stool test within the past year or a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 years and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 40a. High blood pressure among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	658	226,200	31.9	(27.4 - 36.5)	38,125	3,974,800	26.8	(24.9 - 28.7)
<b>Gender</b>								
Male	239	116,300	34.4	(26.6 - 42.3)	14,611	1,960,500	27.7	(24.7 - 30.7)
Female	419	109,900	29.6	(24.7 - 34.6)	23,514	2,014,300	26.0	(23.6 - 28.3)
<b>Age (years)</b>								
18-34	*	*	*	*	3,935	312,200	7.0	(4.2 - 9.8)
35-44	92	20,900	17.2	(8.6 - 25.9)	4,984	426,200	14.8	(11.6 - 18.1)
45-54	149	49,500	34.8	(25.2 - 44.3)	7,825	862,400	30.3	(25.9 - 34.8)
55-64	*	*	*	*	8,765	890,800	41.6	(37.6 - 45.7)
≥ 65	222	85,800	61.4	(54.2 - 68.5)	12,616	1,483,200	58.5	(54.7 - 62.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	584	184,300	31.7	(27.0 - 36.4)	35,814	2,650,900	28.4	(26.9 - 30.0)
Black Non-Hispanic	*	*	*	*	736	558,500	26.2	(19.6 - 32.8)
Hispanic	*	*	*	*	863	538,100	23.8	(17.0 - 30.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	263	103,100	37.0	(29.0 - 45.0)	15,688	1,851,200	33.1	(29.4 - 36.9)
Some College	176	62,100	32.2	(23.4 - 41.0)	9,906	851,100	24.1	(20.9 - 27.3)
≥ College Degree	217	61,000	25.8	(18.9 - 32.7)	12,452	1,253,400	22.3	(19.7 - 24.9)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,640	1,153,000	32.8	(27.6 - 38.1)
\$25,000-\$49,999	162	57,500	34.2	(25.5 - 42.8)	9,657	807,800	27.9	(23.7 - 32.1)
\$50,000-\$74,999	91	21,800	22.2	(13.3 - 31.1)	5,773	507,600	25.8	(21.4 - 30.1)
≥ \$75,000	136	49,600	23.7	(14.5 - 33.0)	8,560	942,500	21.0	(18.4 - 23.6)

<sup>1</sup> Defined as having ever been told by a health professional that respondent has high blood pressure

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have ever been told by a health professional that they have high blood pressure (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 40b. High blood pressure among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	658	28.8	(24.3 - 33.4)	38,125	25.7	(23.8 - 27.5)
<b>Gender</b>						
Male	239	33.2	(25.3 - 41.2)	14,611	27.7	(24.7 - 30.7)
Female	419	24.8	(20.0 - 29.5)	23,514	23.8	(21.5 - 26.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	584	28.9	(24.1 - 33.6)	35,814	27.2	(25.6 - 28.7)
Black Non-Hispanic	*	*	*	736	24.9	(18.4 - 31.3)
Hispanic	*	*	*	863	23.3	(16.6 - 30.0)
Other Non-Hispanic	*	*	*	712	19.3	(9.5 - 29.1)
<b>Education</b>						
≤ High School	263	34.2	(26.1 - 42.4)	15,688	32.1	(28.4 - 35.8)
Some College	176	29.3	(20.6 - 38.0)	9,906	22.8	(19.7 - 25.9)
≥ College Degree	217	22.5	(15.8 - 29.3)	12,452	21.2	(18.6 - 23.8)
<b>Income</b>						
≤ \$24,999	*	*	*	9,640	31.8	(26.7 - 37.0)
\$25,000-\$49,999	162	31.0	(22.3 - 39.7)	9,657	26.9	(22.8 - 30.9)
\$50,000-\$74,999	91	20.0	(11.1 - 28.9)	5,773	24.8	(20.4 - 29.2)
≥ \$75,000	136	21.6	(12.7 - 30.5)	8,560	19.9	(17.3 - 22.5)

<sup>1</sup> Defined as having ever been told by a health professional that respondent has high blood pressure

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Erie County**

**Table 41a. Blood pressure medication among adults with high blood pressure<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	274	185,900	82.4	(75.1 - 89.6)	15,288	3,145,800	79.2	(76.0 - 82.4)
<b>Gender</b>								
Male	*	*	*	*	6,080	1,432,100	73.1	(67.6 - 78.5)
Female	182	97,500	88.7	(82.4 - 95.0)	9,208	1,713,700	85.2	(82.1 - 88.2)
<b>Age (years)</b>								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,315	645,500	74.9	(66.8 - 82.9)
55-64	*	*	*	*	4,045	782,300	87.8	(84.5 - 91.1)
≥ 65	135	78,600	92.1	(86.4 - 97.7)	7,675	1,383,400	93.4	(91.1 - 95.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	241	149,000	81.1	(73.0 - 89.1)	14,442	2,117,300	79.9	(77.5 - 82.4)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	*	*	*	*	7,354	1,482,800	80.2	(74.7 - 85.6)
Some College	*	*	*	*	3,868	632,000	74.3	(67.8 - 80.7)
≥ College Degree	*	*	*	*	4,035	1,013,200	80.9	(76.8 - 84.9)
<b>Income</b>								
≤ \$24,999	*	*	*	*	4,898	948,200	82.3	(75.7 - 88.9)
\$25,000-\$49,999	*	*	*	*	4,088	668,300	82.8	(77.6 - 88.0)
\$50,000-\$74,999	*	*	*	*	1,971	391,500	77.2	(67.7 - 86.6)
≥ \$75,000	*	*	*	*	2,397	683,700	72.5	(66.1 - 78.9)

<sup>1</sup> Defined as currently taking medicine for high blood pressure among adults with high blood pressure

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults with high blood pressure who are taking medicine for high blood pressure (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 41b. Blood pressure medication among adults with high blood pressure<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	274	80.0	(71.9 - 88.2)	15,288	78.5	(75.2 - 81.7)
<b>Gender</b>						
Male	*	*	*	6,080	73.3	(67.9 - 78.7)
Female	182	86.3	(77.9 - 94.6)	9,208	84.0	(80.7 - 87.2)
<b>Race/Ethnicity</b>						
White Non-Hispanic	241	78.2	(69.0 - 87.5)	14,442	79.0	(76.5 - 81.6)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	7,354	79.7	(74.3 - 85.2)
Some College	*	*	*	3,868	72.5	(65.7 - 79.3)
≥ College Degree	*	*	*	4,035	80.4	(76.1 - 84.6)
<b>Income</b>						
≤ \$24,999	*	*	*	4,898	81.6	(75.0 - 88.3)
\$25,000-\$49,999	*	*	*	4,088	82.1	(76.7 - 87.5)
\$50,000-\$74,999	*	*	*	1,971	76.4	(66.7 - 86.1)
≥ \$75,000	*	*	*	2,397	71.6	(64.8 - 78.3)

<sup>1</sup> Defined as currently taking medicine for high blood pressure among adults with high blood pressure

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 42a. Ever had blood cholesterol checked among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	646	556,700	82.6	(77.4 - 87.7)	37,450	11,496,600	80.3	(77.7 - 83.0)
<b>Gender</b>								
Male	230	250,900	80.0	(71.4 - 88.7)	14,389	5,288,400	77.7	(73.3 - 82.0)
Female	416	305,800	84.7	(78.7 - 90.7)	23,061	6,208,200	82.7	(79.6 - 85.8)
<b>Age (years)</b>								
18-34	*	*	*	*	3,684	2,296,400	55.3	(48.7 - 61.9)
35-44	89	99,100	84.0	(76.1 - 92.0)	4,883	2,264,200	81.6	(76.6 - 86.5)
45-54	147	134,100	95.5	(91.3 - 99.7)	7,731	2,539,400	91.0	(87.0 - 94.9)
55-64	136	103,200	97.8	(95.5 - 100)	8,718	2,014,900	95.4	(93.5 - 97.2)
≥ 65	221	132,500	97.0	(94.4 - 99.5)	12,434	2,381,700	95.9	(94.3 - 97.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	575	453,100	81.3	(75.5 - 87.2)	35,198	7,832,600	86.3	(84.8 - 87.9)
Black Non-Hispanic	*	*	*	*	720	1,588,900	77.8	(68.4 - 87.2)
Hispanic	*	*	*	*	842	1,224,600	56.8	(46.8 - 66.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	260	208,900	78.3	(69.7 - 86.9)	15,342	3,795,700	70.6	(65.6 - 75.6)
Some College	*	*	*	*	9,732	2,599,800	77.4	(71.7 - 83.2)
≥ College Degree	212	198,200	89.1	(81.9 - 96.3)	12,296	5,008,300	91.3	(88.9 - 93.6)
<b>Income</b>								
≤ \$24,999	175	131,200	90.8	(83.8 - 97.8)	9,437	2,277,500	68.4	(61.5 - 75.3)
\$25,000-\$49,999	*	*	*	*	9,492	2,053,000	71.9	(64.5 - 79.3)
\$50,000-\$74,999	*	*	*	*	5,721	1,736,100	90.0	(87.7 - 92.3)
≥ \$75,000	*	*	*	*	8,460	4,019,300	92.2	(90.8 - 93.6)

<sup>1</sup> Defined as ever having blood cholesterol checked

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have ever had blood cholesterol checked (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 42b. Ever had blood cholesterol checked among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	646	80.7	(75.2 - 86.3)	37,450	79.9	(77.3 - 82.5)
<b>Gender</b>						
Male	230	79.0	(70.2 - 87.9)	14,389	78.0	(73.9 - 82.1)
Female	416	82.2	(75.3 - 89.2)	23,061	81.6	(78.4 - 84.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	575	79.4	(73.2 - 85.7)	35,198	85.6	(84.0 - 87.2)
Black Non-Hispanic	*	*	*	720	77.5	(68.2 - 86.7)
Hispanic	*	*	*	842	57.2	(47.4 - 67.0)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	260	75.7	(66.3 - 85.0)	15,342	70.2	(65.2 - 75.1)
Some College	*	*	*	9,732	76.7	(71.1 - 82.3)
≥ College Degree	212	87.9	(79.9 - 95.9)	12,296	90.9	(88.6 - 93.2)
<b>Income</b>						
≤ \$24,999	175	89.9	(82.0 - 97.8)	9,437	68.1	(61.3 - 74.9)
\$25,000-\$49,999	*	*	*	9,492	71.6	(64.4 - 78.7)
\$50,000-\$74,999	*	*	*	5,721	89.6	(87.2 - 92.0)
≥ \$75,000	*	*	*	8,460	91.7	(90.1 - 93.2)

<sup>1</sup> Defined as ever having blood cholesterol checked

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 43a. Blood cholesterol checked within the past 5 years among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	644	533,800	79.8	(74.5 - 85.1)	37,201	11,035,200	77.8	(75.1 - 80.4)
<b>Gender</b>								
Male	229	232,400	75.3	(66.3 - 84.3)	14,307	5,020,700	74.5	(70.2 - 78.9)
Female	415	301,400	83.7	(77.6 - 89.7)	22,894	6,014,600	80.7	(77.6 - 83.8)
<b>Age (years)</b>								
18-34	*	*	*	*	3,647	2,179,800	53.3	(46.7 - 59.9)
35-44	89	94,400	80.1	(71.4 - 88.8)	4,862	2,161,200	78.0	(72.8 - 83.2)
45-54	146	128,900	92.2	(86.1 - 98.3)	7,697	2,408,600	86.6	(82.4 - 90.8)
55-64	136	101,600	96.3	(93.1 - 99.4)	8,677	1,957,900	93.3	(91.4 - 95.3)
≥ 65	221	130,800	95.7	(92.7 - 98.7)	12,318	2,327,800	94.8	(93.1 - 96.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	573	430,200	78.0	(72.0 - 84.0)	34,960	7,486,600	83.4	(81.8 - 85.0)
Black Non-Hispanic	*	*	*	*	719	1,561,700	76.5	(67.1 - 85.9)
Hispanic	*	*	*	*	838	1,184,000	55.6	(45.6 - 65.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	259	196,900	75.2	(66.3 - 84.0)	15,215	3,612,500	68.3	(63.2 - 73.3)
Some College	*	*	*	*	9,673	2,524,200	75.4	(69.8 - 81.1)
≥ College Degree	211	190,300	85.8	(77.7 - 93.9)	12,236	4,806,500	88.1	(85.5 - 90.6)
<b>Income</b>								
≤ \$24,999	175	127,900	88.6	(81.3 - 95.9)	9,350	2,214,800	66.9	(60.1 - 73.8)
\$25,000-\$49,999	*	*	*	*	9,454	1,936,600	68.8	(61.5 - 76.1)
\$50,000-\$74,999	*	*	*	*	5,701	1,640,400	85.9	(82.4 - 89.5)
≥ \$75,000	*	*	*	*	8,435	3,872,400	89.2	(87.5 - 90.9)

<sup>1</sup> Defined as having had blood cholesterol checked within the past 5 years

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have had blood cholesterol checked within the past 5 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 43b. Blood cholesterol checked within the past 5 years among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	644	77.9	(72.2 - 83.6)	37,201	77.3	(74.7 - 79.9)
<b>Gender</b>						
Male	229	74.3	(65.0 - 83.5)	14,307	74.8	(70.7 - 79.0)
Female	415	81.0	(74.0 - 88.0)	22,894	79.5	(76.3 - 82.7)
<b>Race/Ethnicity</b>						
White Non-Hispanic	573	76.0	(69.6 - 82.4)	34,960	82.6	(81.0 - 84.3)
Black Non-Hispanic	*	*	*	719	76.2	(66.9 - 85.5)
Hispanic	*	*	*	838	55.9	(46.1 - 65.8)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	259	72.5	(62.9 - 82.0)	15,215	67.8	(62.9 - 72.8)
Some College	*	*	*	9,673	74.7	(69.1 - 80.2)
≥ College Degree	211	84.5	(75.7 - 93.2)	12,236	87.5	(85.0 - 90.1)
<b>Income</b>						
≤ \$24,999	175	87.2	(78.9 - 95.5)	9,350	66.6	(59.8 - 73.4)
\$25,000-\$49,999	*	*	*	9,454	68.5	(61.4 - 75.6)
\$50,000-\$74,999	*	*	*	5,701	85.4	(81.7 - 89.0)
≥ \$75,000	*	*	*	8,435	88.5	(86.7 - 90.3)

<sup>1</sup> Defined as having had blood cholesterol checked within the past 5 years

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Erie County**

**Table 44a. Cardiovascular disease among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	655	57,900	8.2	(6.0 - 10.3)	37,810	1,135,000	7.7	(6.5 - 8.8)
<b>Gender</b>								
Male	237	30,500	9.1	(5.3 - 12.8)	14,477	664,600	9.4	(7.5 - 11.4)
Female	418	27,400	7.4	(5.0 - 9.7)	23,333	470,400	6.1	(4.8 - 7.4)
<b>Age (years)</b>								
18-34	*	*	*	*	3,930	100,800	2.3	(0.0 - 4.8)
35-44	*	*	*	*	4,971	103,400	3.6	(1.5 - 5.7)
45-54	148	7,400	5.3	(1.8 - 8.8)	7,768	131,300	4.6	(2.9 - 6.4)
55-64	133	16,200	15.5	(7.0 - 24.1)	8,699	224,200	10.6	(8.3 - 12.9)
≥ 65	223	33,000	23.5	(17.1 - 29.9)	12,442	575,300	23.0	(19.9 - 26.1)
<b>Race/Ethnicity</b>								
White Non-Hispanic	583	49,300	8.4	(6.3 - 10.6)	35,522	772,200	8.3	(7.4 - 9.2)
Black Non-Hispanic	*	*	*	*	728	140,700	6.6	(3.3 - 9.9)
Hispanic	*	*	*	*	854	163,300	7.3	(2.0 - 12.5)
Other Non-Hispanic	*	*	*	*	706	58,800	5.2	(0.9 - 9.5)
<b>Education</b>								
≤ High School	262	28,500	10.2	(6.6 - 13.8)	15,486	590,800	10.7	(8.2 - 13.2)
Some College	175	18,400	9.6	(4.3 - 14.9)	9,849	227,100	6.4	(4.7 - 8.1)
≥ College Degree	215	11,000	4.7	(2.2 - 7.2)	12,393	314,600	5.6	(4.3 - 6.9)
<b>Income</b>								
≤ \$24,999	176	22,500	15.1	(9.1 - 21.0)	9,505	448,100	12.9	(9.1 - 16.7)
\$25,000-\$49,999	161	11,400	6.8	(3.2 - 10.3)	9,571	179,300	6.2	(4.8 - 7.6)
\$50,000-\$74,999	*	*	*	*	5,751	166,900	8.5	(5.5 - 11.4)
≥ \$75,000	*	*	*	*	8,538	183,500	4.1	(2.9 - 5.3)

<sup>1</sup> Defined as having ever been told by a health professional that respondent had a heart attack or angina or a stroke

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults with cardiovascular disease (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 44b. Cardiovascular disease among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	655	7.1	(5.1 - 9.0)	37,810	7.6	(6.4 - 8.7)
<b>Gender</b>						
Male	237	8.5	(5.0 - 12.0)	14,477	9.7	(7.8 - 11.7)
Female	418	5.8	(3.9 - 7.7)	23,333	5.6	(4.3 - 6.9)
<b>Race/Ethnicity</b>						
White Non-Hispanic	583	7.4	(5.4 - 9.4)	35,522	8.2	(7.2 - 9.1)
Black Non-Hispanic	*	*	*	728	6.8	(3.2 - 10.4)
Hispanic	*	*	*	854	7.0	(2.0 - 12.0)
Other Non-Hispanic	*	*	*	706	5.5	(0.9 - 10.0)
<b>Education</b>						
≤ High School	262	8.9	(5.6 - 12.2)	15,486	10.5	(8.0 - 13.0)
Some College	175	8.5	(3.7 - 13.2)	9,849	6.4	(4.5 - 8.2)
≥ College Degree	215	4.0	(1.8 - 6.2)	12,393	5.5	(4.2 - 6.9)
<b>Income</b>						
≤ \$24,999	176	14.0	(8.1 - 19.9)	9,505	12.8	(9.0 - 16.6)
\$25,000-\$49,999	161	5.9	(2.6 - 9.1)	9,571	6.0	(4.7 - 7.4)
\$50,000-\$74,999	*	*	*	5,751	8.5	(5.4 - 11.6)
≥ \$75,000	*	*	*	8,538	4.0	(2.7 - 5.2)

<sup>1</sup> Defined as having ever been told by a health professional that respondent had a heart attack or angina or a stroke

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Erie County**

**Table 45a. Coronary heart disease among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	655	47,300	6.7	(4.7 - 8.7)	37,838	929,400	6.3	(5.3 - 7.3)
<b>Gender</b>								
Male	236	26,600	7.9	(4.3 - 11.5)	14,479	598,600	8.5	(6.6 - 10.4)
Female	419	20,700	5.6	(3.5 - 7.6)	23,359	330,800	4.3	(3.4 - 5.2)
<b>Age (years)</b>								
18-34	*	*	*	*	3,932	63,400	1.4	(0.0 - 3.6)
35-44	*	*	*	*	4,973	95,300	3.3	(1.2 - 5.4)
45-54	*	*	*	*	7,778	108,300	3.8	(2.1 - 5.6)
55-64	132	11,900	11.5	(3.1 - 19.8)	8,703	188,600	8.9	(6.7 - 11.1)
≥ 65	224	27,900	19.8	(13.7 - 25.8)	12,452	473,800	18.9	(16.1 - 21.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	583	39,500	6.8	(4.8 - 8.7)	35,552	636,700	6.9	(6.0 - 7.7)
Black Non-Hispanic	*	*	*	*	728	113,100	5.3	(2.3 - 8.4)
Hispanic	*	*	*	*	853	128,000	5.7	(1.1 - 10.3)
Other Non-Hispanic	*	*	*	*	705	51,500	4.6	(0.5 - 8.7)
<b>Education</b>								
≤ High School	261	21,500	7.7	(4.6 - 10.8)	15,485	474,500	8.6	(6.3 - 10.8)
Some College	175	16,500	8.6	(3.4 - 13.8)	9,860	193,000	5.5	(3.9 - 7.1)
≥ College Degree	216	9,300	3.9	(1.6 - 6.3)	12,411	259,700	4.6	(3.4 - 5.8)
<b>Income</b>								
≤ \$24,999	176	19,400	13.0	(7.4 - 18.5)	9,507	372,300	10.7	(7.3 - 14.1)
\$25,000-\$49,999	161	9,300	5.5	(2.3 - 8.7)	9,585	136,300	4.7	(3.8 - 5.7)
\$50,000-\$74,999	*	*	*	*	5,755	154,600	7.8	(4.9 - 10.7)
≥ \$75,000	*	*	*	*	8,541	151,800	3.4	(2.3 - 4.5)

<sup>1</sup> Defined as having ever been told by a health professional that respondent had a heart attack or angina

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults with coronary heart disease (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 45b. Coronary heart disease among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	655	5.8	(4.0 - 7.7)	37,838	6.3	(5.2 - 7.3)
<b>Gender</b>						
Male	236	7.5	(4.1 - 10.8)	14,479	8.8	(6.9 - 10.8)
Female	419	4.4	(2.7 - 6.0)	23,359	3.9	(3.0 - 4.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	583	5.9	(4.1 - 7.8)	35,552	6.8	(5.9 - 7.7)
Black Non-Hispanic	*	*	*	728	5.6	(2.2 - 9.0)
Hispanic	*	*	*	853	5.4	(1.1 - 9.6)
Other Non-Hispanic	*	*	*	705	4.8	(0.4 - 9.2)
<b>Education</b>						
≤ High School	261	6.7	(3.9 - 9.5)	15,485	8.5	(6.2 - 10.7)
Some College	175	7.7	(3.1 - 12.4)	9,860	5.5	(3.7 - 7.2)
≥ College Degree	216	3.5	(1.3 - 5.6)	12,411	4.7	(3.4 - 5.9)
<b>Income</b>						
≤ \$24,999	176	12.0	(6.6 - 17.5)	9,507	10.7	(7.2 - 14.1)
\$25,000-\$49,999	161	4.8	(1.9 - 7.7)	9,585	4.7	(3.6 - 5.7)
\$50,000-\$74,999	*	*	*	5,755	8.0	(4.8 - 11.1)
≥ \$75,000	*	*	*	8,541	3.3	(2.1 - 4.5)

<sup>1</sup> Defined as having ever been told by a health professional that respondent had a heart attack or angina

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 46a. Ever diagnosed with diabetes among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	661	80,300	11.3	(8.3 - 14.2)	38,174	1,382,700	9.3	(8.0 - 10.5)
<b>Gender</b>								
Male	239	40,100	11.9	(6.7 - 17.0)	14,626	530,400	7.5	(6.2 - 8.8)
Female	422	40,200	10.7	(7.5 - 13.9)	23,548	852,300	11.0	(8.9 - 13.0)
<b>Age (years)</b>								
18-34	*	*	*	*	3,941	150,200	3.4	(1.0 - 5.8)
35-44	*	*	*	*	4,987	165,100	5.7	(3.6 - 7.9)
45-54	149	14,400	10.1	(1.9 - 18.4)	7,830	279,000	9.8	(6.7 - 12.8)
55-64	137	22,200	20.8	(11.7 - 30.0)	8,771	332,900	15.5	(12.3 - 18.8)
≥ 65	225	30,100	21.3	(15.2 - 27.3)	12,645	455,600	17.9	(15.0 - 20.8)
<b>Race/Ethnicity</b>								
White Non-Hispanic	587	56,200	9.6	(7.2 - 12.0)	35,862	809,300	8.7	(7.8 - 9.5)
Black Non-Hispanic	*	*	*	*	739	261,000	12.1	(6.6 - 17.6)
Hispanic	*	*	*	*	862	227,300	10.1	(5.4 - 14.8)
Other Non-Hispanic	*	*	*	*	711	85,100	7.6	(2.3 - 12.8)
<b>Education</b>								
≤ High School	265	40,700	14.4	(8.8 - 20.0)	15,709	730,000	13.0	(10.5 - 15.6)
Some College	175	20,900	10.9	(5.3 - 16.5)	9,918	303,500	8.6	(5.9 - 11.2)
≥ College Degree	218	18,100	7.6	(4.0 - 11.3)	12,464	337,500	6.0	(4.8 - 7.2)
<b>Income</b>								
≤ \$24,999	180	27,800	18.2	(9.7 - 26.6)	9,652	543,400	15.5	(11.4 - 19.5)
\$25,000-\$49,999	163	19,900	11.8	(6.6 - 17.0)	9,667	291,000	10.0	(7.5 - 12.6)
\$50,000-\$74,999	*	*	*	*	5,777	151,000	7.6	(5.0 - 10.3)
≥ \$75,000	135	14,600	7.0	(1.9 - 12.1)	8,565	220,800	4.9	(3.8 - 6.1)

<sup>1</sup> Defined as having ever been told by a doctor that respondent had diabetes, excluding pre-diabetes and women with diabetes only when pregnant

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have diabetes (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 46b. Ever diagnosed with diabetes among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	661	10.5	(7.6 - 13.4)	38,174	9.0	(7.8 - 10.3)
<b>Gender</b>						
Male	239	11.1	(6.3 - 16.0)	14,626	7.5	(6.1 - 8.9)
Female	422	9.9	(6.5 - 13.3)	23,548	10.4	(8.4 - 12.4)
<b>Race/Ethnicity</b>						
White Non-Hispanic	587	8.9	(6.4 - 11.3)	35,862	8.4	(7.5 - 9.2)
Black Non-Hispanic	*	*	*	739	11.4	(6.0 - 16.7)
Hispanic	*	*	*	862	10.1	(5.4 - 14.9)
Other Non-Hispanic	*	*	*	711	7.7	(2.3 - 13.0)
<b>Education</b>						
≤ High School	265	13.7	(8.0 - 19.5)	15,709	12.6	(10.0 - 15.2)
Some College	175	9.9	(4.8 - 15.1)	9,918	8.3	(5.6 - 10.9)
≥ College Degree	218	7.1	(3.5 - 10.8)	12,464	5.9	(4.7 - 7.2)
<b>Income</b>						
≤ \$24,999	180	17.3	(8.7 - 25.8)	9,652	15.2	(11.1 - 19.2)
\$25,000-\$49,999	163	11.2	(5.8 - 16.7)	9,667	9.7	(7.1 - 12.2)
\$50,000-\$74,999	*	*	*	5,777	7.6	(4.9 - 10.2)
≥ \$75,000	135	6.5	(2.0 - 11.1)	8,565	4.7	(3.6 - 5.9)

<sup>1</sup> Defined as having ever been told by a doctor that respondent had diabetes, excluding pre-diabetes and women with diabetes only when pregnant

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 47a. Ever diagnosed with asthma (lifetime) among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	662	93,900	13.2	(9.7 - 16.6)	38,096	2,225,100	15.0	(13.2 - 16.8)
<b>Gender</b>								
Male	239	18,000	5.3	(1.9 - 8.7)	14,601	875,800	12.4	(10.1 - 14.7)
Female	423	75,900	20.2	(14.8 - 25.7)	23,495	1,349,300	17.4	(14.6 - 20.1)
<b>Age (years)</b>								
18-34	59	35,700	17.7	(7.8 - 27.5)	3,931	825,100	18.5	(13.8 - 23.3)
35-44	*	*	*	*	4,972	431,700	15.1	(11.2 - 19.1)
45-54	149	19,800	13.9	(8.1 - 19.7)	7,819	365,900	12.8	(9.7 - 16.0)
55-64	137	12,000	11.3	(5.9 - 16.6)	8,756	316,700	14.8	(11.9 - 17.7)
≥ 65	225	14,900	10.5	(5.7 - 15.4)	12,618	285,700	11.2	(8.6 - 13.8)
<b>Race/Ethnicity</b>								
White Non-Hispanic	588	69,200	11.8	(8.3 - 15.3)	35,783	1,365,700	14.6	(13.0 - 16.3)
Black Non-Hispanic	*	*	*	*	737	367,000	17.2	(10.2 - 24.2)
Hispanic	*	*	*	*	863	317,100	14.1	(9.0 - 19.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	265	39,700	14.0	(8.3 - 19.8)	15,675	831,600	14.9	(12.0 - 17.8)
Some College	176	23,400	12.1	(5.4 - 18.9)	9,893	587,500	16.6	(13.1 - 20.1)
≥ College Degree	218	29,900	12.6	(7.1 - 18.2)	12,444	739,000	13.2	(10.4 - 15.9)
<b>Income</b>								
≤ \$24,999	180	28,300	18.4	(10.0 - 26.8)	9,643	708,900	20.2	(14.9 - 25.5)
\$25,000-\$49,999	163	17,500	10.4	(3.8 - 17.0)	9,641	389,800	13.4	(10.0 - 16.9)
\$50,000-\$74,999	*	*	*	*	5,762	257,800	13.2	(10.0 - 16.5)
≥ \$75,000	136	24,700	11.8	(5.7 - 17.9)	8,562	577,900	12.8	(10.4 - 15.3)

<sup>1</sup> Defined as adults having ever been told by a health professional that they had asthma

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have lifetime asthma (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 47b. Ever diagnosed with asthma (lifetime) among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	662	13.4	(9.7 - 17.2)	38,096	15.2	(13.3 - 17.0)
<b>Gender</b>						
Male	239	5.2	(1.8 - 8.7)	14,601	12.6	(10.3 - 14.9)
Female	423	20.8	(14.7 - 26.9)	23,495	17.5	(14.8 - 20.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	588	12.2	(8.3 - 16.0)	35,783	14.9	(13.2 - 16.6)
Black Non-Hispanic	*	*	*	737	17.1	(10.1 - 24.1)
Hispanic	*	*	*	863	14.5	(9.3 - 19.7)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	265	14.3	(8.1 - 20.5)	15,675	15.1	(12.1 - 18.0)
Some College	176	12.7	(5.3 - 20.0)	9,893	16.9	(13.4 - 20.4)
≥ College Degree	218	12.8	(6.7 - 18.9)	12,444	13.3	(10.6 - 16.0)
<b>Income</b>						
≤ \$24,999	180	18.8	(9.7 - 27.9)	9,643	20.3	(15.0 - 25.6)
\$25,000-\$49,999	163	10.7	(3.6 - 17.9)	9,641	13.7	(10.2 - 17.2)
\$50,000-\$74,999	*	*	*	5,762	13.5	(10.0 - 17.0)
≥ \$75,000	136	12.1	(5.6 - 18.6)	8,562	13.0	(10.6 - 15.4)

<sup>1</sup> Defined as adults having ever been told by a health professional that they had asthma

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Erie County**

**Table 48a. Current asthma among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	658	55,100	7.8	(5.4 - 10.2)	37,933	1,415,000	9.6	(8.1 - 11.0)
<b>Gender</b>								
Male	*	*	*	*	14,546	440,700	6.3	(4.8 - 7.7)
Female	420	48,100	12.9	(8.7 - 17.1)	23,387	974,300	12.6	(10.2 - 15.0)
<b>Age (years)</b>								
18-34	*	*	*	*	3,909	467,200	10.5	(6.8 - 14.3)
35-44	*	*	*	*	4,947	246,800	8.7	(5.8 - 11.6)
45-54	148	14,900	10.5	(5.3 - 15.7)	7,787	240,000	8.4	(6.1 - 10.8)
55-64	136	8,400	7.9	(3.2 - 12.7)	8,726	231,100	10.8	(8.1 - 13.5)
≥ 65	224	12,700	9.0	(4.4 - 13.6)	12,564	229,900	9.1	(6.5 - 11.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	584	38,400	6.6	(4.3 - 8.9)	35,629	888,400	9.6	(8.2 - 11.0)
Black Non-Hispanic	*	*	*	*	736	246,900	11.6	(5.7 - 17.5)
Hispanic	*	*	*	*	859	186,800	8.3	(4.4 - 12.2)
Other Non-Hispanic	*	*	*	*	709	93,000	8.3	(1.4 - 15.2)
<b>Education</b>								
≤ High School	262	28,400	10.1	(5.8 - 14.5)	15,594	595,100	10.7	(8.2 - 13.2)
Some College	*	*	*	*	9,861	322,000	9.1	(7.0 - 11.3)
≥ College Degree	218	18,600	7.8	(3.5 - 12.2)	12,394	430,900	7.7	(5.8 - 9.7)
<b>Income</b>								
≤ \$24,999	178	20,600	13.5	(7.1 - 19.9)	9,600	489,200	14.0	(9.7 - 18.3)
\$25,000-\$49,999	162	7,600	4.6	(1.4 - 7.7)	9,590	268,400	9.3	(6.2 - 12.4)
\$50,000-\$74,999	*	*	*	*	5,741	143,400	7.4	(5.1 - 9.6)
≥ \$75,000	136	15,000	7.2	(2.3 - 12.0)	8,541	339,200	7.6	(5.8 - 9.4)

<sup>1</sup> Defined as adults having ever been told by a health professional that they had asthma and still have asthma

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who currently have asthma (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Erie County**

**Table 48b. Current asthma among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	658	7.7	(5.1 - 10.3)	37,933	9.7	(8.2 - 11.2)
<b>Gender</b>						
Male	*	*	*	14,546	6.4	(4.9 - 7.9)
Female	420	13.0	(8.4 - 17.6)	23,387	12.7	(10.2 - 15.1)
<b>Race/Ethnicity</b>						
White Non-Hispanic	584	6.6	(4.1 - 9.1)	35,629	9.7	(8.3 - 11.2)
Black Non-Hispanic	*	*	*	736	11.5	(5.6 - 17.3)
Hispanic	*	*	*	859	8.6	(4.5 - 12.6)
Other Non-Hispanic	*	*	*	709	8.3	(1.3 - 15.3)
<b>Education</b>						
≤ High School	262	10.0	(5.3 - 14.7)	15,594	10.8	(8.3 - 13.3)
Some College	*	*	*	9,861	9.4	(7.2 - 11.6)
≥ College Degree	218	8.1	(3.2 - 12.9)	12,394	7.8	(5.8 - 9.7)
<b>Income</b>						
≤ \$24,999	178	13.7	(6.7 - 20.6)	9,600	14.1	(9.7 - 18.4)
\$25,000-\$49,999	162	4.5	(1.2 - 7.8)	9,590	9.5	(6.3 - 12.7)
\$50,000-\$74,999	*	*	*	5,741	7.5	(5.0 - 10.1)
≥ \$75,000	136	7.4	(2.1 - 12.7)	8,541	7.7	(5.9 - 9.5)

<sup>1</sup> Defined as adults having ever been told by a health professional that they had asthma and still have asthma

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 49a. Chronic joint symptoms among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	661	359,100	50.4	(45.0 - 55.8)	38,099	6,377,100	42.9	(40.5 - 45.3)
<b>Gender</b>								
Male	238	164,600	48.8	(40.0 - 57.6)	14,595	3,033,600	42.8	(39.0 - 46.6)
Female	423	194,500	51.8	(45.4 - 58.2)	23,504	3,343,500	43.0	(40.0 - 46.0)
<b>Age (years)</b>								
18-34	*	*	*	*	3,933	1,261,300	28.3	(22.6 - 34.0)
35-44	*	*	*	*	4,984	1,110,900	38.6	(33.1 - 44.0)
45-54	149	76,900	54.0	(44.5 - 63.5)	7,817	1,456,500	51.2	(46.4 - 55.9)
55-64	136	63,500	59.9	(50.0 - 69.9)	8,751	1,167,400	54.6	(50.4 - 58.8)
≥ 65	225	87,700	62.0	(54.8 - 69.1)	12,614	1,381,000	54.3	(50.5 - 58.1)
<b>Race/Ethnicity</b>								
White Non-Hispanic	587	293,200	50.0	(44.5 - 55.6)	35,785	4,267,600	45.8	(43.8 - 47.7)
Black Non-Hispanic	*	*	*	*	740	890,500	41.3	(32.5 - 50.0)
Hispanic	*	*	*	*	860	827,500	36.7	(27.7 - 45.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	264	154,600	54.9	(45.8 - 63.9)	15,676	2,450,000	43.8	(39.4 - 48.1)
Some College	*	*	*	*	9,900	1,598,300	45.1	(40.1 - 50.1)
≥ College Degree	218	106,900	45.1	(36.4 - 53.8)	12,440	2,313,100	41.1	(37.8 - 44.5)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,641	1,483,000	42.2	(36.3 - 48.1)
\$25,000-\$49,999	*	*	*	*	9,647	1,332,100	45.9	(39.8 - 52.1)
\$50,000-\$74,999	*	*	*	*	5,768	894,200	45.3	(39.5 - 51.1)
≥ \$75,000	*	*	*	*	8,549	1,874,200	41.7	(38.4 - 45.1)

<sup>1</sup> Defined as adults having symptoms of pain, aching or stiffness in or around a joint during the past 30 days and began more than 3 months ago

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have chronic joint symptoms (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 49b. Chronic joint symptoms among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	661	48.9	(43.2 - 54.6)	38,099	42.1	(39.7 - 44.5)
<b>Gender</b>						
Male	238	48.5	(39.5 - 57.6)	14,595	42.6	(38.8 - 46.3)
Female	423	49.3	(42.2 - 56.4)	23,504	41.6	(38.6 - 44.7)
<b>Race/Ethnicity</b>						
White Non-Hispanic	587	48.6	(42.7 - 54.5)	35,785	44.7	(42.7 - 46.6)
Black Non-Hispanic	*	*	*	740	40.2	(31.5 - 48.9)
Hispanic	*	*	*	860	36.6	(27.8 - 45.4)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	264	53.8	(44.2 - 63.4)	15,676	43.1	(38.8 - 47.5)
Some College	*	*	*	9,900	43.9	(39.0 - 48.8)
≥ College Degree	218	43.4	(34.1 - 52.8)	12,440	40.3	(37.0 - 43.7)
<b>Income</b>						
≤ \$24,999	*	*	*	9,641	41.6	(35.7 - 47.4)
\$25,000-\$49,999	*	*	*	9,647	45.0	(39.0 - 51.0)
\$50,000-\$74,999	*	*	*	5,768	44.3	(38.4 - 50.2)
≥ \$75,000	*	*	*	8,549	40.8	(37.4 - 44.2)

<sup>1</sup> Defined as adults having symptoms of pain, aching or stiffness in or around a joint during the past 30 days and began more than 3 months ago

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 50a. Arthritis among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	658	231,700	32.6	(28.2 - 37.1)	38,013	4,102,800	27.7	(25.9 - 29.6)
<b>Gender</b>								
Male	237	90,200	26.9	(20.2 - 33.6)	14,555	1,642,700	23.3	(20.7 - 25.8)
Female	421	141,500	37.8	(32.0 - 43.6)	23,458	2,460,100	31.8	(29.2 - 34.4)
<b>Age (years)</b>								
18-34	*	*	*	*	3,927	312,100	7.0	(4.9 - 9.1)
35-44	92	23,200	19.2	(10.9 - 27.5)	4,972	412,500	14.5	(11.3 - 17.6)
45-54	148	49,700	35.2	(25.3 - 45.1)	7,789	869,100	30.7	(26.6 - 34.8)
55-64	*	*	*	*	8,729	999,300	47.0	(42.9 - 51.1)
≥ 65	222	90,700	65.2	(58.2 - 72.3)	12,596	1,509,800	59.6	(55.8 - 63.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	584	194,000	33.3	(28.6 - 37.9)	35,710	2,967,700	31.9	(30.3 - 33.6)
Black Non-Hispanic	*	*	*	*	737	588,100	27.3	(20.2 - 34.4)
Hispanic	*	*	*	*	856	380,700	17.1	(11.8 - 22.5)
Other Non-Hispanic	*	*	*	*	710	166,300	14.8	(7.8 - 21.8)
<b>Education</b>								
≤ High School	263	112,100	39.9	(31.8 - 48.1)	15,652	1,742,700	31.3	(27.8 - 34.8)
Some College	175	60,100	31.4	(22.9 - 39.8)	9,874	902,000	25.5	(22.3 - 28.6)
≥ College Degree	218	59,200	25.0	(18.6 - 31.4)	12,407	1,437,800	25.6	(22.9 - 28.4)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,618	1,112,500	32.0	(27.1 - 36.9)
\$25,000-\$49,999	163	69,000	40.9	(31.2 - 50.6)	9,624	821,900	28.4	(24.2 - 32.6)
\$50,000-\$74,999	*	*	*	*	5,753	542,500	27.5	(23.2 - 31.9)
≥ \$75,000	135	32,100	15.4	(9.6 - 21.3)	8,536	1,012,800	22.6	(19.8 - 25.3)

<sup>1</sup> Defined as adults having ever been told by a health professional that they have some form of arthritis, rheumatoid arthritis, gout, lupus or fibromyalgia (excluding back or neck)

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have ever been told by a health professional they have some form of arthritis (excluding back or neck) (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 50b. Arthritis among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	658	29.8	(25.3 - 34.3)	38,013	26.6	(24.8 - 28.5)
<b>Gender</b>						
Male	237	25.7	(19.1 - 32.3)	14,555	23.5	(20.9 - 26.1)
Female	421	33.5	(27.4 - 39.5)	23,458	29.5	(27.0 - 32.1)
<b>Race/Ethnicity</b>						
White Non-Hispanic	584	30.5	(25.8 - 35.2)	35,710	30.4	(28.7 - 32.0)
Black Non-Hispanic	*	*	*	737	26.5	(19.3 - 33.6)
Hispanic	*	*	*	856	17.2	(11.8 - 22.6)
Other Non-Hispanic	*	*	*	710	14.9	(7.7 - 22.0)
<b>Education</b>						
≤ High School	263	37.3	(29.0 - 45.5)	15,652	30.4	(26.9 - 33.8)
Some College	175	29.1	(20.4 - 37.7)	9,874	24.1	(21.0 - 27.1)
≥ College Degree	218	22.1	(15.8 - 28.3)	12,407	24.7	(21.9 - 27.5)
<b>Income</b>						
≤ \$24,999	*	*	*	9,618	31.2	(26.3 - 36.1)
\$25,000-\$49,999	*	*	*	9,624	27.3	(23.2 - 31.3)
\$50,000-\$74,999	*	*	*	5,753	26.3	(21.9 - 30.7)
≥ \$75,000	135	14.6	(8.8 - 20.5)	8,536	21.6	(18.8 - 24.3)

<sup>1</sup> Defined as adults having ever been told by a health professional that they have some form of arthritis, rheumatoid arthritis, gout, lupus or fibromyalgia (excluding back or neck)

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 51a. Activity limitations because of physical, mental or emotional problems among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	661	164,800	23.1	(19.1 - 27.2)	37,998	3,053,000	20.6	(18.7 - 22.5)
<b>Gender</b>								
Male	238	78,200	23.3	(16.3 - 30.2)	14,566	1,318,400	18.6	(15.7 - 21.5)
Female	423	86,600	23.1	(18.5 - 27.6)	23,432	1,734,600	22.4	(19.8 - 25.0)
<b>Age (years)</b>								
18-34	*	*	*	*	3,929	580,000	13.0	(8.2 - 17.9)
35-44	92	16,100	13.3	(5.6 - 21.0)	4,973	418,600	14.5	(11.0 - 18.1)
45-54	149	33,100	23.3	(15.7 - 30.9)	7,812	620,200	21.7	(18.3 - 25.2)
55-64	137	44,100	41.4	(31.6 - 51.2)	8,739	651,600	30.5	(26.5 - 34.6)
≥ 65	224	44,600	31.9	(24.9 - 38.8)	12,545	782,700	31.1	(27.7 - 34.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	587	142,000	24.3	(19.8 - 28.7)	35,688	2,037,200	21.9	(20.3 - 23.5)
Black Non-Hispanic	*	*	*	*	735	429,700	20.1	(13.5 - 26.7)
Hispanic	*	*	*	*	863	448,200	19.8	(11.9 - 27.8)
Other Non-Hispanic	*	*	*	*	712	137,900	12.3	(5.2 - 19.3)
<b>Education</b>								
≤ High School	265	87,600	31.0	(23.5 - 38.5)	15,626	1,359,600	24.4	(20.6 - 28.1)
Some College	175	39,600	20.7	(12.7 - 28.7)	9,875	700,800	19.8	(16.5 - 23.2)
≥ College Degree	218	36,400	15.4	(10.3 - 20.4)	12,415	924,800	16.5	(14.2 - 18.7)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,603	1,055,800	30.1	(24.7 - 35.5)
\$25,000-\$49,999	163	37,900	22.5	(15.4 - 29.5)	9,628	630,900	21.8	(16.6 - 27.0)
\$50,000-\$74,999	91	18,300	18.6	(9.4 - 27.8)	5,760	408,100	20.7	(16.1 - 25.2)
≥ \$75,000	136	23,100	11.0	(5.0 - 17.0)	8,544	583,300	13.0	(11.1 - 14.9)

<sup>1</sup> Defined as activity limitations because of physical, mental or emotional problems

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who are limited in any activities because of physical, mental or emotional problems (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 51b. Activity limitations because of physical, mental or emotional problems among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	661	21.4	(17.3 - 25.5)	37,998	20.1	(18.2 - 22.0)
<b>Gender</b>						
Male	238	22.3	(15.3 - 29.3)	14,566	18.6	(15.8 - 21.4)
Female	423	20.6	(16.0 - 25.1)	23,432	21.5	(18.9 - 24.1)
<b>Race/Ethnicity</b>						
White Non-Hispanic	587	22.6	(18.0 - 27.2)	35,688	21.3	(19.7 - 22.9)
Black Non-Hispanic	*	*	*	735	19.9	(13.2 - 26.5)
Hispanic	*	*	*	863	19.1	(11.5 - 26.8)
Other Non-Hispanic	*	*	*	712	12.4	(5.3 - 19.6)
<b>Education</b>						
≤ High School	265	29.9	(22.1 - 37.6)	15,626	23.8	(20.1 - 27.4)
Some College	175	19.4	(11.1 - 27.6)	9,875	19.3	(16.0 - 22.6)
≥ College Degree	218	13.2	(8.4 - 17.9)	12,415	16.1	(13.9 - 18.3)
<b>Income</b>						
≤ \$24,999	*	*	*	9,603	29.7	(24.3 - 35.2)
\$25,000-\$49,999	163	21.1	(14.0 - 28.2)	9,628	21.2	(16.1 - 26.3)
\$50,000-\$74,999	91	17.8	(8.2 - 27.5)	5,760	20.1	(15.6 - 24.6)
≥ \$75,000	136	9.9	(4.1 - 15.6)	8,544	12.7	(10.8 - 14.6)

<sup>1</sup> Defined as activity limitations because of physical, mental or emotional problems

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 52a. Have health problem that requires the use of special equipment among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	662	75,700	10.6	(8.0 - 13.2)	38,171	1,169,400	7.9	(6.9 - 8.9)
<b>Gender</b>								
Male	239	32,200	9.5	(5.3 - 13.8)	14,617	460,700	6.5	(5.3 - 7.7)
Female	423	43,500	11.6	(8.4 - 14.8)	23,554	708,700	9.1	(7.6 - 10.7)
<b>Age (years)</b>								
18-34	*	*	*	*	3,940	68,800	1.5	(0.7 - 2.4)
35-44	*	*	*	*	4,989	122,500	4.3	(2.4 - 6.1)
45-54	149	17,000	11.9	(6.4 - 17.5)	7,836	182,500	6.4	(4.7 - 8.1)
55-64	137	19,300	18.1	(11.1 - 25.1)	8,773	219,600	10.3	(7.8 - 12.8)
≥ 65	225	29,200	20.7	(14.3 - 27.0)	12,633	576,100	22.7	(19.0 - 26.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	588	59,800	10.2	(7.4 - 13.0)	35,853	802,000	8.6	(7.6 - 9.6)
Black Non-Hispanic	*	*	*	*	739	235,800	10.9	(6.4 - 15.5)
Hispanic	*	*	*	*	864	104,100	4.6	(2.1 - 7.1)
Other Non-Hispanic	*	*	*	*	715	27,600	2.5	(0.8 - 4.1)
<b>Education</b>								
≤ High School	265	41,200	14.6	(9.4 - 19.8)	15,711	604,200	10.8	(8.7 - 12.8)
Some College	176	16,800	8.7	(4.5 - 12.9)	9,914	260,000	7.3	(5.5 - 9.1)
≥ College Degree	218	17,700	7.5	(4.1 - 10.9)	12,463	298,900	5.3	(4.1 - 6.5)
<b>Income</b>								
≤ \$24,999	180	31,000	20.2	(12.8 - 27.6)	9,656	430,500	12.2	(9.6 - 14.8)
\$25,000-\$49,999	163	16,200	9.6	(5.1 - 14.1)	9,663	238,900	8.2	(5.9 - 10.5)
\$50,000-\$74,999	*	*	*	*	5,775	103,000	5.2	(3.6 - 6.8)
≥ \$75,000	136	9,600	4.6	(1.6 - 7.6)	8,566	136,000	3.0	(2.1 - 4.0)

<sup>1</sup> Defined as having any health problem that requires the use of special equipment such as a cane, a wheelchair, a special bed, or a special telephone

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have any health problem that requires the use of special equipment (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 52b. Have health problem that requires the use of special equipment among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	662	9.6	(7.0 - 12.2)	38,171	7.6	(6.6 - 8.6)
<b>Gender</b>						
Male	239	9.4	(5.1 - 13.8)	14,617	6.8	(5.5 - 8.0)
Female	423	9.8	(6.9 - 12.7)	23,554	8.4	(6.9 - 9.9)
<b>Race/Ethnicity</b>						
White Non-Hispanic	588	9.1	(6.4 - 11.9)	35,853	8.3	(7.3 - 9.3)
Black Non-Hispanic	*	*	*	739	10.5	(6.1 - 15.0)
Hispanic	*	*	*	864	4.5	(2.1 - 6.9)
Other Non-Hispanic	*	*	*	715	2.2	(0.7 - 3.8)
<b>Education</b>						
≤ High School	265	13.7	(8.4 - 19.0)	15,711	10.3	(8.3 - 12.3)
Some College	176	8.1	(3.8 - 12.5)	9,914	7.1	(5.2 - 8.9)
≥ College Degree	218	6.2	(3.3 - 9.2)	12,463	5.3	(4.1 - 6.6)
<b>Income</b>						
≤ \$24,999	180	19.6	(11.8 - 27.4)	9,656	11.9	(9.4 - 14.5)
\$25,000-\$49,999	163	8.6	(4.4 - 12.8)	9,663	7.7	(5.5 - 10.0)
\$50,000-\$74,999	*	*	*	5,775	5.1	(3.4 - 6.8)
≥ \$75,000	136	4.2	(1.3 - 7.1)	8,566	2.9	(2.0 - 3.9)

<sup>1</sup> Defined as having any health problem that requires the use of special equipment such as a cane, a wheelchair, a special bed, or a special telephone

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 53a. Disability among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	661	183,700	25.8	(21.6 - 30.0)	38,016	3,394,200	22.9	(20.9 - 24.9)
<b>Gender</b>								
Male	238	91,000	27.1	(19.8 - 34.3)	14,568	1,469,700	20.7	(17.8 - 23.7)
Female	423	92,700	24.7	(20.0 - 29.3)	23,448	1,924,600	24.8	(22.1 - 27.5)
<b>Age (years)</b>								
18-34	*	*	*	*	3,931	590,300	13.3	(8.4 - 18.1)
35-44	92	17,300	14.3	(6.4 - 22.1)	4,974	447,900	15.6	(11.9 - 19.3)
45-54	149	35,900	25.2	(17.4 - 33.0)	7,811	643,900	22.6	(19.1 - 26.0)
55-64	137	46,400	43.5	(33.7 - 53.4)	8,740	702,100	33.0	(28.9 - 37.0)
≥ 65	224	57,300	40.9	(33.6 - 48.3)	12,560	1,010,000	39.9	(36.1 - 43.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	587	157,500	26.9	(22.3 - 31.5)	35,703	2,253,400	24.2	(22.6 - 25.9)
Black Non-Hispanic	*	*	*	*	736	511,700	23.8	(16.8 - 30.8)
Hispanic	*	*	*	*	864	484,300	21.4	(13.4 - 29.5)
Other Non-Hispanic	*	*	*	*	713	144,800	12.9	(5.8 - 19.9)
<b>Education</b>								
≤ High School	265	96,100	34.0	(26.3 - 41.8)	15,638	1,544,200	27.6	(23.7 - 31.4)
Some College	175	43,400	22.7	(14.5 - 30.8)	9,876	755,200	21.4	(17.9 - 24.8)
≥ College Degree	218	43,000	18.2	(12.7 - 23.6)	12,419	1,021,300	18.2	(15.9 - 20.5)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,613	1,179,100	33.6	(28.0 - 39.2)
\$25,000-\$49,999	163	45,000	26.7	(19.0 - 34.3)	9,629	699,000	24.2	(18.9 - 29.4)
\$50,000-\$74,999	91	18,900	19.3	(10.0 - 28.5)	5,757	430,100	21.8	(17.2 - 26.4)
≥ \$75,000	136	27,300	13.1	(6.7 - 19.4)	8,546	632,700	14.1	(12.1 - 16.1)

<sup>1</sup> Defined as having activity limitations due to physical, mental or emotional problems or have health problems that require the use of special equipment

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults with a disability (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 53b. Disability among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	661	23.8	(19.6 - 28.1)	38,016	22.3	(20.4 - 24.3)
<b>Gender</b>						
Male	238	26.2	(18.9 - 33.5)	14,568	20.9	(18.0 - 23.8)
Female	423	21.8	(17.1 - 26.4)	23,448	23.6	(21.0 - 26.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	587	25.0	(20.3 - 29.7)	35,703	23.6	(21.9 - 25.3)
Black Non-Hispanic	*	*	*	736	23.5	(16.5 - 30.5)
Hispanic	*	*	*	864	20.6	(12.9 - 28.3)
Other Non-Hispanic	*	*	*	713	12.9	(5.8 - 20.1)
<b>Education</b>						
≤ High School	265	32.7	(24.7 - 40.7)	15,638	26.8	(23.0 - 30.6)
Some College	175	21.2	(12.8 - 29.6)	9,876	20.9	(17.4 - 24.3)
≥ College Degree	218	15.7	(10.6 - 20.9)	12,419	17.9	(15.5 - 20.3)
<b>Income</b>						
≤ \$24,999	*	*	*	9,613	33.2	(27.6 - 38.8)
\$25,000-\$49,999	163	25.0	(17.3 - 32.7)	9,629	23.5	(18.3 - 28.6)
\$50,000-\$74,999	91	18.4	(8.6 - 28.1)	5,757	21.2	(16.6 - 25.8)
≥ \$75,000	136	11.7	(5.7 - 17.8)	8,546	13.8	(11.7 - 15.8)

<sup>1</sup> Defined as having activity limitations due to physical, mental or emotional problems or have health problems that require the use of special equipment

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 54a. Current smoking among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	662	182,000	25.5	(20.3 - 30.8)	38,029	2,483,500	16.8	(15.1 - 18.6)
<b>Gender</b>								
Male	239	98,300	29.1	(20.3 - 38.0)	14,559	1,356,400	19.3	(16.3 - 22.2)
Female	423	83,700	22.3	(16.4 - 28.1)	23,470	1,127,100	14.6	(12.7 - 16.6)
<b>Age (years)</b>								
18-34	*	*	*	*	3,929	993,500	22.3	(17.5 - 27.1)
35-44	*	*	*	*	4,977	421,100	14.8	(11.6 - 17.9)
45-54	149	29,300	20.6	(13.3 - 27.9)	7,807	508,800	18.3	(15.4 - 21.3)
55-64	137	27,900	26.2	(17.0 - 35.4)	8,744	314,500	14.7	(12.1 - 17.3)
≥ 65	225	9,300	6.6	(2.5 - 10.6)	12,572	245,600	9.7	(7.2 - 12.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	588	134,600	23.0	(17.8 - 28.1)	35,729	1,615,500	17.3	(15.9 - 18.8)
Black Non-Hispanic	*	*	*	*	735	284,100	13.6	(8.9 - 18.2)
Hispanic	*	*	*	*	858	350,900	15.8	(9.6 - 21.9)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	265	97,200	34.4	(25.1 - 43.8)	15,654	1,237,300	22.4	(19.1 - 25.6)
Some College	176	50,800	26.4	(16.8 - 35.9)	9,875	670,600	19.1	(15.5 - 22.8)
≥ College Degree	218	33,600	14.2	(7.2 - 21.2)	12,421	563,300	10.0	(7.5 - 12.5)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,626	835,100	24.1	(19.1 - 29.1)
\$25,000-\$49,999	163	44,000	26.1	(16.8 - 35.4)	9,629	558,100	19.4	(15.6 - 23.2)
\$50,000-\$74,999	*	*	*	*	5,756	338,100	17.3	(13.0 - 21.7)
≥ \$75,000	*	*	*	*	8,549	468,700	10.4	(8.6 - 12.3)

<sup>1</sup> Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day or some days

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adult current smokers (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 54b. Current smoking among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	662	26.7	(21.0 - 32.3)	38,029	17.0	(15.3 - 18.8)
<b>Gender</b>						
Male	239	29.6	(20.3 - 38.9)	14,559	19.2	(16.3 - 22.1)
Female	423	24.0	(17.4 - 30.6)	23,470	15.0	(13.0 - 17.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	588	24.2	(18.6 - 29.9)	35,729	17.7	(16.2 - 19.2)
Black Non-Hispanic	*	*	*	735	13.5	(8.8 - 18.2)
Hispanic	*	*	*	858	15.9	(9.8 - 22.0)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	265	36.4	(26.5 - 46.4)	15,654	22.8	(19.5 - 26.0)
Some College	*	*	*	9,875	19.4	(15.8 - 22.9)
≥ College Degree	218	14.8	(6.8 - 22.8)	12,421	10.0	(7.6 - 12.5)
<b>Income</b>						
≤ \$24,999	*	*	*	9,626	24.1	(19.2 - 29.1)
\$25,000-\$49,999	*	*	*	9,629	19.9	(16.0 - 23.7)
\$50,000-\$74,999	*	*	*	5,756	17.8	(13.3 - 22.3)
≥ \$75,000	*	*	*	8,549	10.6	(8.6 - 12.5)

<sup>1</sup> Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day or some days

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 55a. Everyday smoking among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	662	116,600	16.3	(12.4 - 20.2)	38,029	1,626,800	11.0	(9.8 - 12.2)
<b>Gender</b>								
Male	239	56,200	16.6	(10.6 - 22.6)	14,559	848,400	12.0	(10.0 - 14.1)
Female	423	60,400	16.1	(11.0 - 21.2)	23,470	778,400	10.1	(8.7 - 11.5)
<b>Age (years)</b>								
18-34	*	*	*	*	3,929	559,800	12.6	(9.6 - 15.6)
35-44	92	23,300	19.3	(10.5 - 28.0)	4,977	315,200	11.0	(8.4 - 13.7)
45-54	149	22,400	15.7	(9.0 - 22.4)	7,807	377,000	13.6	(11.1 - 16.0)
55-64	137	20,200	19.0	(10.9 - 27.0)	8,744	203,000	9.5	(7.5 - 11.5)
≥ 65	225	8,300	5.8	(1.9 - 9.8)	12,572	171,700	6.8	(4.8 - 8.8)
<b>Race/Ethnicity</b>								
White Non-Hispanic	588	95,900	16.3	(12.1 - 20.6)	35,729	1,158,100	12.4	(11.2 - 13.7)
Black Non-Hispanic	*	*	*	*	735	197,700	9.4	(5.6 - 13.3)
Hispanic	*	*	*	*	858	176,200	7.9	(4.2 - 11.6)
Other Non-Hispanic	*	*	*	*	707	94,900	8.5	(3.3 - 13.7)
<b>Education</b>								
≤ High School	265	68,100	24.1	(16.5 - 31.7)	15,654	881,900	15.9	(13.4 - 18.5)
Some College	176	32,700	16.9	(9.6 - 24.3)	9,875	410,200	11.7	(9.6 - 13.8)
≥ College Degree	218	15,500	6.6	(2.8 - 10.4)	12,421	323,500	5.8	(4.3 - 7.2)
<b>Income</b>								
≤ \$24,999	180	37,700	24.6	(14.7 - 34.4)	9,626	452,800	13.1	(10.3 - 15.8)
\$25,000-\$49,999	163	34,600	20.5	(12.1 - 28.9)	9,629	425,900	14.8	(11.9 - 17.7)
\$50,000-\$74,999	*	*	*	*	5,756	222,700	11.4	(8.5 - 14.4)
≥ \$75,000	136	17,200	8.2	(3.2 - 13.2)	8,549	320,200	7.1	(5.5 - 8.8)

<sup>1</sup> Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have smoked at least 100 cigarettes in lifetime and currently smoking everyday (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 55b. Everyday smoking among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	662	17.0	(12.8 - 21.3)	38,029	11.3	(10.0 - 12.5)
<b>Gender</b>						
Male	239	16.2	(10.2 - 22.2)	14,559	12.1	(10.1 - 14.1)
Female	423	17.8	(11.9 - 23.7)	23,470	10.5	(9.0 - 12.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	588	17.4	(12.6 - 22.1)	35,729	12.8	(11.5 - 14.1)
Black Non-Hispanic	*	*	*	735	9.4	(5.5 - 13.4)
Hispanic	*	*	*	858	8.0	(4.3 - 11.7)
Other Non-Hispanic	*	*	*	707	8.5	(3.2 - 13.8)
<b>Education</b>						
≤ High School	265	25.5	(17.3 - 33.7)	15,654	16.4	(13.8 - 19.0)
Some College	176	18.2	(10.0 - 26.5)	9,875	12.1	(9.9 - 14.3)
≥ College Degree	218	6.6	(2.4 - 10.7)	12,421	5.8	(4.3 - 7.3)
<b>Income</b>						
≤ \$24,999	*	*	*	9,626	13.4	(10.6 - 16.2)
\$25,000-\$49,999	163	21.2	(12.1 - 30.4)	9,629	15.3	(12.3 - 18.3)
\$50,000-\$74,999	*	*	*	5,756	11.7	(8.7 - 14.8)
≥ \$75,000	136	8.7	(3.1 - 14.3)	8,549	7.2	(5.5 - 8.9)

<sup>1</sup> Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 56a. Adults living in homes in which smoking is prohibited<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	658	521,200	73.3	(68.4 - 78.2)	38,045	11,997,100	80.8	(79.0 - 82.6)
<b>Gender</b>								
Male	238	232,800	69.0	(60.8 - 77.3)	14,577	5,634,400	79.4	(76.6 - 82.2)
Female	420	288,400	77.1	(71.5 - 82.7)	23,468	6,362,800	82.1	(79.9 - 84.3)
<b>Age (years)</b>								
18-34	*	*	*	*	3,934	3,655,200	82.1	(77.8 - 86.4)
35-44	92	92,800	76.6	(67.4 - 85.9)	4,985	2,479,700	86.1	(82.5 - 89.7)
45-54	149	99,800	70.1	(61.3 - 78.9)	7,808	2,194,400	77.2	(73.5 - 80.8)
55-64	137	81,200	76.2	(68.0 - 84.4)	8,753	1,732,500	81.0	(78.3 - 83.7)
≥ 65	221	107,900	77.4	(71.1 - 83.6)	12,565	1,935,400	76.5	(73.4 - 79.5)
<b>Race/Ethnicity</b>								
White Non-Hispanic	584	435,400	74.5	(69.7 - 79.3)	35,737	7,426,900	79.8	(78.3 - 81.2)
Black Non-Hispanic	*	*	*	*	738	1,692,100	78.6	(71.9 - 85.3)
Hispanic	*	*	*	*	863	2,003,900	88.7	(84.1 - 93.3)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	265	179,500	63.5	(54.6 - 72.4)	15,647	4,307,200	77.1	(74.0 - 80.2)
Some College	174	140,900	73.5	(64.6 - 82.3)	9,883	2,795,400	78.9	(75.0 - 82.8)
≥ College Degree	218	200,900	84.8	(78.5 - 91.1)	12,437	4,817,300	85.7	(83.3 - 88.2)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,608	2,627,700	74.9	(70.0 - 79.7)
\$25,000-\$49,999	163	119,200	70.6	(61.4 - 79.9)	9,640	2,314,500	79.8	(76.0 - 83.5)
\$50,000-\$74,999	*	*	*	*	5,766	1,551,300	78.7	(74.0 - 83.4)
≥ \$75,000	136	184,200	88.0	(81.3 - 94.7)	8,556	3,883,200	86.5	(84.3 - 88.7)

<sup>1</sup> Defined as having a rule that prohibits smoking anywhere inside the home

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who live in a home with a rule that prohibits smoking anywhere inside the home (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 56b. Adults living in homes in which smoking is prohibited<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	658	73.1	(67.8 - 78.3)	38,045	80.9	(79.2 - 82.7)
<b>Gender</b>						
Male	238	68.8	(60.3 - 77.3)	14,577	79.4	(76.7 - 82.2)
Female	420	76.9	(70.7 - 83.2)	23,468	82.3	(80.0 - 84.5)
<b>Race/Ethnicity</b>						
White Non-Hispanic	584	74.3	(69.1 - 79.4)	35,737	79.9	(78.4 - 81.4)
Black Non-Hispanic	*	*	*	738	78.6	(71.8 - 85.4)
Hispanic	*	*	*	863	88.9	(84.4 - 93.4)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	265	61.9	(52.3 - 71.4)	15,647	76.9	(73.8 - 80.0)
Some College	174	73.8	(64.5 - 83.2)	9,883	79.1	(75.2 - 83.0)
≥ College Degree	218	85.1	(78.6 - 91.6)	12,437	86.0	(83.5 - 88.5)
<b>Income</b>						
≤ \$24,999	*	*	*	9,608	74.9	(70.1 - 79.7)
\$25,000-\$49,999	163	71.2	(61.3 - 81.0)	9,640	79.8	(76.1 - 83.6)
\$50,000-\$74,999	*	*	*	5,766	78.6	(73.8 - 83.3)
≥ \$75,000	136	87.5	(80.5 - 94.5)	8,556	86.6	(84.3 - 88.9)

<sup>1</sup> Defined as having a rule that prohibits smoking anywhere inside the home

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 57a. Binge drinking within the past month among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	656	164,100	23.1	(18.2 - 28.1)	37,745	2,602,000	17.7	(15.7 - 19.8)
<b>Gender</b>								
Male	236	107,600	32.1	(23.6 - 40.5)	14,420	1,712,400	24.5	(20.9 - 28.1)
Female	420	56,500	15.1	(10.0 - 20.2)	23,325	889,600	11.6	(9.5 - 13.7)
<b>Age (years)</b>								
18-34	*	*	*	*	3,895	1,254,300	28.5	(22.8 - 34.2)
35-44	*	*	*	*	4,934	584,800	20.6	(16.5 - 24.7)
45-54	148	38,000	26.8	(18.2 - 35.5)	7,738	467,600	16.6	(13.3 - 19.9)
55-64	134	18,000	17.2	(9.3 - 25.1)	8,663	207,200	9.8	(7.9 - 11.8)
≥ 65	*	*	*	*	12,515	88,100	3.5	(2.5 - 4.5)
<b>Race/Ethnicity</b>								
White Non-Hispanic	582	148,600	25.5	(20.0 - 31.0)	35,453	1,761,900	19.2	(17.5 - 20.8)
Black Non-Hispanic	*	*	*	*	728	247,000	11.6	(6.3 - 17.0)
Hispanic	*	*	*	*	854	460,600	20.5	(11.3 - 29.8)
Other Non-Hispanic	*	*	*	*	710	132,500	12.1	(3.4 - 20.7)
<b>Education</b>								
≤ High School	262	64,500	23.1	(15.0 - 31.1)	15,491	945,800	17.4	(13.2 - 21.6)
Some College	*	*	*	*	9,819	636,600	18.1	(14.6 - 21.5)
≥ College Degree	215	42,800	18.2	(11.1 - 25.4)	12,358	1,018,700	18.2	(15.3 - 21.1)
<b>Income</b>								
≤ \$24,999	179	21,200	13.9	(5.7 - 22.1)	9,557	469,100	13.5	(8.5 - 18.5)
\$25,000-\$49,999	160	32,600	19.5	(11.3 - 27.7)	9,575	534,300	18.5	(13.0 - 24.0)
\$50,000-\$74,999	*	*	*	*	5,726	378,200	19.2	(14.1 - 24.3)
≥ \$75,000	*	*	*	*	8,501	1,011,400	22.6	(19.7 - 25.6)

<sup>1</sup> Defined as men having 5 or more drinks or women having 4 or more drinks on 1 or more occasion within the past month

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adult binge drinkers within the past month (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 57b. Binge drinking within the past month among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	656	24.1	(18.9 - 29.3)	37,745	18.1	(16.1 - 20.2)
<b>Gender</b>						
Male	236	32.4	(23.8 - 41.1)	14,420	24.5	(21.0 - 27.9)
Female	420	16.6	(10.7 - 22.4)	23,325	12.4	(10.2 - 14.5)
<b>Race/Ethnicity</b>						
White Non-Hispanic	582	26.6	(20.7 - 32.4)	35,453	19.8	(18.1 - 21.6)
Black Non-Hispanic	*	*	*	728	11.8	(6.4 - 17.1)
Hispanic	*	*	*	854	20.2	(11.3 - 29.1)
Other Non-Hispanic	*	*	*	710	12.1	(3.4 - 20.8)
<b>Education</b>						
≤ High School	262	23.9	(15.2 - 32.5)	15,491	17.5	(13.5 - 21.5)
Some College	*	*	*	9,819	18.6	(15.1 - 22.0)
≥ College Degree	215	19.1	(11.4 - 26.8)	12,358	18.8	(15.9 - 21.7)
<b>Income</b>						
≤ \$24,999	179	15.2	(6.0 - 24.4)	9,557	13.4	(8.6 - 18.2)
\$25,000-\$49,999	160	19.2	(10.7 - 27.7)	9,575	18.9	(13.5 - 24.2)
\$50,000-\$74,999	*	*	*	5,726	19.9	(14.8 - 25.1)
≥ \$75,000	*	*	*	8,501	23.2	(20.2 - 26.1)

<sup>1</sup> Defined as men having 5 or more drinks or women having 4 or more drinks on 1 or more occasion within the past month

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 58a. Heavy drinking within the past month among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	654	48,800	6.9	(4.2 - 9.5)	37,553	718,700	4.9	(4.2 - 5.6)
<b>Gender</b>								
Male	235	23,700	7.1	(2.9 - 11.3)	14,330	333,300	4.8	(3.9 - 5.7)
Female	419	25,100	6.7	(3.4 - 10.1)	23,223	385,400	5.0	(4.0 - 6.1)
<b>Age (years)</b>								
18-34	*	*	*	*	3,866	224,900	5.1	(3.5 - 6.8)
35-44	*	*	*	*	4,924	137,200	4.8	(3.0 - 6.7)
45-54	148	18,800	13.3	(6.5 - 20.1)	7,715	144,600	5.2	(4.1 - 6.2)
55-64	*	*	*	*	8,624	117,400	5.6	(3.9 - 7.3)
≥ 65	*	*	*	*	12,424	94,600	3.8	(2.7 - 4.9)
<b>Race/Ethnicity</b>								
White Non-Hispanic	580	46,600	8.0	(4.9 - 11.2)	35,279	572,400	6.3	(5.4 - 7.1)
Black Non-Hispanic	*	*	*	*	723	58,100	2.8	(0.6 - 4.9)
Hispanic	*	*	*	*	848	60,900	2.7	(0.5 - 4.9)
Other Non-Hispanic	*	*	*	*	703	27,200	2.5	(0.8 - 4.2)
<b>Education</b>								
≤ High School	263	17,800	6.3	(2.8 - 9.9)	15,411	228,100	4.2	(3.2 - 5.2)
Some College	*	*	*	*	9,759	163,700	4.7	(3.6 - 5.8)
≥ College Degree	215	17,500	7.5	(2.7 - 12.3)	12,306	326,800	5.9	(4.5 - 7.3)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,512	104,200	3.0	(1.8 - 4.1)
\$25,000-\$49,999	160	13,100	7.9	(2.6 - 13.1)	9,537	158,700	5.5	(3.9 - 7.2)
\$50,000-\$74,999	*	*	*	*	5,708	95,400	4.9	(3.1 - 6.7)
≥ \$75,000	135	22,000	10.5	(3.7 - 17.4)	8,481	313,900	7.0	(5.4 - 8.6)

<sup>1</sup> Defined as adult men averaging more than 2 alcoholic drinks per day and adult women averaging more than 1 alcoholic drink per day within the past month

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adult heavy drinkers within the past month (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 58b. Heavy drinking within the past month among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	654	7.1	(4.3 - 10.0)	37,553	5.0	(4.3 - 5.8)
<b>Gender</b>						
Male	235	7.0	(2.8 - 11.2)	14,330	4.9	(3.9 - 5.9)
Female	419	7.2	(3.4 - 11.1)	23,223	5.1	(4.0 - 6.2)
<b>Race/Ethnicity</b>						
White Non-Hispanic	580	8.3	(4.9 - 11.7)	35,279	6.4	(5.5 - 7.2)
Black Non-Hispanic	*	*	*	723	2.9	(0.6 - 5.1)
Hispanic	*	*	*	848	2.8	(0.5 - 5.1)
Other Non-Hispanic	*	*	*	703	2.5	(0.8 - 4.2)
<b>Education</b>						
≤ High School	263	6.1	(2.6 - 9.6)	15,411	4.3	(3.2 - 5.3)
Some College	*	*	*	9,759	4.8	(3.6 - 5.9)
≥ College Degree	215	7.8	(2.5 - 13.2)	12,306	6.0	(4.5 - 7.4)
<b>Income</b>						
≤ \$24,999	*	*	*	9,512	3.0	(1.9 - 4.1)
\$25,000-\$49,999	160	7.6	(2.5 - 12.8)	9,537	5.8	(4.0 - 7.5)
\$50,000-\$74,999	*	*	*	5,708	5.0	(3.1 - 7.0)
≥ \$75,000	135	10.9	(3.6 - 18.2)	8,481	7.0	(5.4 - 8.7)

<sup>1</sup> Defined as adult men averaging more than 2 alcoholic drinks per day and adult women averaging more than 1 alcoholic drink per day within the past month

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 59a. Never or rarely been asked about sexual history during routine check-up among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	609	423,500	65.1	(59.3 - 71.0)	35,193	8,480,600	61.9	(59.2 - 64.5)
<b>Gender</b>								
Male	213	217,400	71.5	(61.7 - 81.3)	13,299	4,357,400	67.7	(63.6 - 71.7)
Female	396	206,100	59.5	(52.7 - 66.4)	21,894	4,123,300	56.8	(53.3 - 60.2)
<b>Age (years)</b>								
18-34	*	*	*	*	3,507	1,804,600	44.3	(37.8 - 50.8)
35-44	*	*	*	*	4,531	1,635,300	62.2	(56.6 - 67.8)
45-54	140	95,700	71.4	(63.0 - 79.9)	7,215	1,715,600	65.4	(60.5 - 70.3)
55-64	124	78,300	82.0	(73.7 - 90.4)	8,168	1,416,200	70.7	(66.6 - 74.8)
≥ 65	212	110,700	83.4	(78.0 - 88.8)	11,772	1,908,900	80.3	(76.9 - 83.8)
<b>Race/Ethnicity</b>								
White Non-Hispanic	544	378,700	70.0	(64.2 - 75.8)	33,090	5,940,300	68.5	(66.3 - 70.7)
Black Non-Hispanic	*	*	*	*	683	831,800	40.8	(31.8 - 49.9)
Hispanic	*	*	*	*	784	1,071,700	53.4	(43.6 - 63.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	240	164,000	64.9	(54.9 - 74.9)	14,264	3,001,200	60.1	(55.4 - 64.9)
Some College	*	*	*	*	9,156	1,972,800	60.3	(54.9 - 65.7)
≥ College Degree	199	137,000	64.6	(55.1 - 74.2)	11,706	3,476,000	65.0	(61.2 - 68.7)
<b>Income</b>								
≤ \$24,999	*	*	*	*	8,772	1,529,400	49.3	(42.6 - 56.0)
\$25,000-\$49,999	*	*	*	*	8,932	1,705,000	64.4	(58.0 - 70.7)
\$50,000-\$74,999	*	*	*	*	5,433	1,217,000	64.5	(58.5 - 70.5)
≥ \$75,000	*	*	*	*	8,108	2,934,300	68.2	(64.7 - 71.7)

<sup>1</sup> Defined as respondents who never or rarely (hardly ever) were asked about their sexual history, such as about sexual partners and sexual practices, during a routine check-up at a doctor's office or clinic

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who were never or rarely asked about their sexual history during a routine check-up at a doctor's office or clinic (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 59b. Never or rarely been asked about sexual history during routine check-up among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	609	62.6	(56.3 - 68.9)	35,193	61.4	(58.7 - 64.0)
<b>Gender</b>						
Male	*	*	*	13,299	67.9	(64.0 - 71.9)
Female	396	56.3	(48.6 - 63.9)	21,894	55.5	(52.0 - 59.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	544	67.8	(61.6 - 74.1)	33,090	67.8	(65.6 - 70.1)
Black Non-Hispanic	*	*	*	683	41.1	(32.0 - 50.2)
Hispanic	*	*	*	784	53.0	(43.3 - 62.6)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	14,264	59.5	(54.7 - 64.3)
Some College	*	*	*	9,156	59.7	(54.4 - 65.0)
≥ College Degree	*	*	*	11,706	64.5	(60.8 - 68.2)
<b>Income</b>						
≤ \$24,999	*	*	*	8,772	48.7	(42.0 - 55.4)
\$25,000-\$49,999	*	*	*	8,932	63.7	(57.5 - 70.0)
\$50,000-\$74,999	*	*	*	5,433	64.2	(58.2 - 70.2)
≥ \$75,000	*	*	*	8,108	67.5	(64.0 - 71.0)

<sup>1</sup> Defined as respondents who never or rarely (hardly ever) were asked about their sexual history, such as about sexual partners and sexual practices, during a routine check-up at a doctor's office or clinic

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 60a. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	414	399,900	79.9	(74.8 - 84.9)	22,788	7,011,100	77.9	(75.2 - 80.5)
<b>Gender</b>								
Male	155	210,300	85.6	(79.1 - 92.1)	9,034	3,618,500	81.0	(77.2 - 84.8)
Female	259	189,600	74.4	(67.0 - 81.8)	13,754	3,392,600	74.8	(71.2 - 78.4)
<b>Age (years)</b>								
18-34	*	*	*	*	3,213	2,362,500	70.8	(65.1 - 76.6)
35-44	*	*	*	*	3,513	1,313,400	71.7	(65.6 - 77.7)
45-54	99	73,900	75.3	(65.9 - 84.8)	4,975	1,290,600	80.3	(75.9 - 84.8)
55-64	*	*	*	*	4,955	932,200	87.2	(84.1 - 90.3)
≥ 65	121	72,400	95.0	(90.3 - 99.7)	6,132	1,112,400	95.9	(94.4 - 97.5)
<b>Race/Ethnicity</b>								
White Non-Hispanic	374	330,900	79.6	(74.2 - 85.0)	21,540	4,875,200	78.3	(75.6 - 80.9)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	491	1,002,400	84.8	(77.8 - 91.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	149	150,600	84.3	(76.8 - 91.8)	8,879	2,581,700	82.1	(77.9 - 86.3)
Some College	*	*	*	*	5,953	1,599,500	77.5	(72.6 - 82.3)
≥ College Degree	148	141,200	79.0	(70.6 - 87.4)	7,928	2,824,800	75.6	(71.8 - 79.5)
<b>Income</b>								
≤ \$24,999	*	*	*	*	5,491	1,602,900	78.7	(71.9 - 85.6)
\$25,000-\$49,999	115	102,400	80.4	(71.6 - 89.2)	5,912	1,335,900	78.8	(73.8 - 83.9)
\$50,000-\$74,999	*	*	*	*	3,717	987,500	76.6	(70.4 - 82.8)
≥ \$75,000	99	135,900	80.7	(70.9 - 90.5)	5,771	2,321,600	77.6	(74.0 - 81.2)

<sup>1</sup> Defined as respondent estimating there were hardly any (0-1 out of 10) or a few (2-3 out of 10) persons their age in their community who have had a sexually transmitted disease such as gonorrhea, chlamydia, herpes or syphilis

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who believed hardly any or a few people their age in their community have had a sexually transmitted disease (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 60b. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	414	79.0	(73.6 - 84.5)	22,788	77.3	(74.7 - 79.9)
<b>Gender</b>						
Male	155	85.6	(79.0 - 92.2)	9,034	81.1	(77.4 - 84.8)
Female	259	72.7	(64.6 - 80.9)	13,754	73.6	(70.0 - 77.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	374	78.4	(72.5 - 84.3)	21,540	77.5	(74.9 - 80.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	491	84.6	(77.6 - 91.5)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	149	82.2	(73.6 - 90.9)	8,879	81.6	(77.4 - 85.7)
Some College	*	*	*	5,953	76.8	(72.0 - 81.6)
≥ College Degree	148	79.1	(70.2 - 87.9)	7,928	75.1	(71.3 - 79.0)
<b>Income</b>						
≤ \$24,999	*	*	*	5,491	77.9	(71.0 - 84.7)
\$25,000-\$49,999	115	79.6	(70.0 - 89.2)	5,912	77.9	(72.8 - 83.0)
\$50,000-\$74,999	*	*	*	3,717	76.1	(70.0 - 82.3)
≥ \$75,000	*	*	*	5,771	77.0	(73.4 - 80.6)

<sup>1</sup> Defined as respondent estimating there were hardly any (0-1 out of 10) or a few (2-3 out of 10) persons their age in their community who have had a sexually transmitted disease such as gonorrhea, chlamydia, herpes or syphilis

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Erie County**

**Table 61a. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	630	615,200	89.8	(86.8 - 92.8)	36,287	12,738,300	89.7	(88.2 - 91.3)
<b>Gender</b>								
Male	227	287,500	88.8	(83.4 - 94.1)	13,893	6,104,600	89.4	(87.2 - 91.5)
Female	403	327,700	90.7	(87.7 - 93.7)	22,394	6,633,700	90.1	(87.9 - 92.2)
<b>Age (years)</b>								
18-34	58	190,500	95.9	(90.3 - 100)	3,850	4,007,400	91.3	(87.5 - 95.1)
35-44	90	109,200	92.9	(87.5 - 98.3)	4,875	2,578,600	93.2	(90.4 - 96.0)
45-54	147	125,200	89.2	(80.8 - 97.7)	7,607	2,416,800	90.5	(87.9 - 93.2)
55-64	129	91,900	91.5	(86.8 - 96.2)	8,411	1,876,500	92.2	(90.0 - 94.5)
≥ 65	206	98,400	76.8	(70.3 - 83.2)	11,544	1,858,900	79.7	(76.6 - 82.8)
<b>Race/Ethnicity</b>								
White Non-Hispanic	559	507,200	90.1	(87.1 - 93.0)	34,102	8,050,700	89.4	(87.7 - 91.0)
Black Non-Hispanic	*	*	*	*	702	1,822,200	91.9	(88.2 - 95.6)
Hispanic	*	*	*	*	820	1,967,200	89.9	(85.2 - 94.7)
Other Non-Hispanic	*	*	*	*	663	898,200	88.6	(79.5 - 97.6)
<b>Education</b>								
≤ High School	250	227,700	84.5	(78.1 - 91.0)	14,507	4,385,300	84.6	(81.9 - 87.2)
Some College	167	167,400	90.7	(86.3 - 95.1)	9,571	3,094,300	90.0	(86.7 - 93.4)
≥ College Degree	211	219,200	95.2	(92.6 - 97.8)	12,142	5,228,800	95.4	(94.0 - 96.7)
<b>Income</b>								
≤ \$24,999	168	120,500	82.7	(74.0 - 91.4)	9,021	2,833,500	86.5	(82.6 - 90.3)
\$25,000-\$49,999	157	153,600	93.4	(89.6 - 97.1)	9,261	2,529,900	89.5	(86.5 - 92.5)
\$50,000-\$74,999	90	93,700	96.5	(92.8 - 100)	5,644	1,740,000	91.2	(87.4 - 95.1)
≥ \$75,000	135	198,300	95.4	(91.9 - 98.8)	8,397	4,184,300	94.4	(92.5 - 96.3)

<sup>1</sup> Defined as believing it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who believed it is at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 61b. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	630	90.6	(87.6 - 93.6)	36,287	89.8	(88.3 - 91.4)
<b>Gender</b>						
Male	227	88.7	(83.3 - 94.1)	13,893	89.1	(86.9 - 91.3)
Female	403	92.3	(89.3 - 95.2)	22,394	90.5	(88.3 - 92.7)
<b>Race/Ethnicity</b>						
White Non-Hispanic	559	90.8	(87.8 - 93.8)	34,102	89.5	(87.8 - 91.2)
Black Non-Hispanic	*	*	*	702	91.7	(87.8 - 95.6)
Hispanic	*	*	*	820	89.6	(84.7 - 94.5)
Other Non-Hispanic	*	*	*	663	89.0	(80.5 - 97.6)
<b>Education</b>						
≤ High School	250	85.5	(78.9 - 92.1)	14,507	84.6	(81.8 - 87.4)
Some College	167	91.3	(86.9 - 95.8)	9,571	90.2	(86.9 - 93.5)
≥ College Degree	211	95.6	(92.9 - 98.2)	12,142	95.4	(94.0 - 96.7)
<b>Income</b>						
≤ \$24,999	168	83.9	(75.2 - 92.7)	9,021	86.5	(82.6 - 90.4)
\$25,000-\$49,999	157	93.6	(89.8 - 97.4)	9,261	89.6	(86.5 - 92.6)
\$50,000-\$74,999	90	97.2	(94.3 - 100)	5,644	91.1	(86.9 - 95.3)
≥ \$75,000	135	95.1	(91.4 - 98.9)	8,397	94.5	(92.6 - 96.4)

<sup>1</sup> Defined as believing it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 62a. Overweight among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	636	233,300	34.1	(28.9 - 39.2)	36,710	5,213,000	36.4	(34.0 - 38.8)
<b>Gender</b>								
Male	234	136,800	41.2	(32.4 - 49.9)	14,502	3,073,200	43.7	(39.9 - 47.6)
Female	402	96,500	27.4	(22.1 - 32.7)	22,208	2,139,800	29.3	(26.4 - 32.2)
<b>Age (years)</b>								
18-34	*	*	*	*	3,789	1,295,200	30.1	(24.7 - 35.6)
35-44	*	*	*	*	4,785	1,052,000	37.5	(31.7 - 43.3)
45-54	143	47,100	34.1	(25.1 - 43.1)	7,495	1,053,300	38.8	(34.1 - 43.4)
55-64	133	41,300	39.8	(29.8 - 49.7)	8,356	798,100	38.8	(34.8 - 42.7)
≥ 65	218	50,700	37.3	(30.0 - 44.6)	12,285	1,014,400	41.5	(37.6 - 45.4)
<b>Race/Ethnicity</b>								
White Non-Hispanic	565	195,300	34.7	(29.5 - 40.0)	34,501	3,282,200	36.4	(34.4 - 38.3)
Black Non-Hispanic	*	*	*	*	708	740,000	36.7	(28.3 - 45.1)
Hispanic	*	*	*	*	813	781,500	35.6	(27.1 - 44.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	256	100,100	36.8	(27.9 - 45.8)	15,079	1,889,700	35.4	(31.2 - 39.5)
Some College	171	60,000	31.9	(22.4 - 41.3)	9,515	1,314,500	38.7	(33.8 - 43.7)
≥ College Degree	207	73,200	32.8	(24.8 - 40.8)	12,062	1,996,900	36.3	(32.8 - 39.7)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,346	1,170,900	34.0	(28.2 - 39.8)
\$25,000-\$49,999	*	*	*	*	9,368	1,136,000	40.4	(34.6 - 46.2)
\$50,000-\$74,999	*	*	*	*	5,615	738,700	38.3	(32.3 - 44.3)
≥ \$75,000	130	48,600	24.3	(15.8 - 32.7)	8,337	1,659,100	37.5	(34.1 - 40.9)

<sup>1</sup> Defined as having a body mass index (BMI) more than 24.9 but less than 30.0. BMI is calculated as weight in kilograms divided by the square of height in meters.

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who are overweight (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 62b. Overweight among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	636	33.4	(28.0 - 38.8)	36,710	36.2	(33.8 - 38.6)
<b>Gender</b>						
Male	234	40.6	(31.6 - 49.6)	14,502	43.9	(40.1 - 47.6)
Female	402	26.5	(20.8 - 32.2)	22,208	28.9	(25.9 - 31.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	565	33.9	(28.4 - 39.3)	34,501	36.0	(34.0 - 38.0)
Black Non-Hispanic	*	*	*	708	37.0	(28.5 - 45.5)
Hispanic	*	*	*	813	36.0	(27.5 - 44.4)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	256	37.0	(27.4 - 46.5)	15,079	35.2	(31.1 - 39.4)
Some College	171	31.4	(21.5 - 41.3)	9,515	38.3	(33.3 - 43.2)
≥ College Degree	207	31.0	(22.7 - 39.2)	12,062	36.2	(32.7 - 39.8)
<b>Income</b>						
≤ \$24,999	*	*	*	9,346	33.8	(28.0 - 39.5)
\$25,000-\$49,999	*	*	*	9,368	40.6	(34.8 - 46.4)
\$50,000-\$74,999	*	*	*	5,615	38.6	(32.5 - 44.7)
≥ \$75,000	130	22.7	(14.4 - 31.0)	8,337	37.2	(33.7 - 40.6)

<sup>1</sup> Defined as having a body mass index (BMI) more than 24.9 but less than 30.0. BMI is calculated as weight in kilograms divided by the square of height in meters.

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Erie County**

**Table 63a. Obesity among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	636	195,100	28.5	(23.7 - 33.3)	36,710	3,327,400	23.2	(21.2 - 25.3)
<b>Gender</b>								
Male	234	96,700	29.1	(21.0 - 37.2)	14,502	1,505,800	21.4	(19.0 - 23.9)
Female	402	98,400	27.9	(22.5 - 33.3)	22,208	1,821,600	25.0	(21.8 - 28.1)
<b>Age (years)</b>								
18-34	*	*	*	*	3,789	730,600	17.0	(12.4 - 21.6)
35-44	*	*	*	*	4,785	662,200	23.6	(19.4 - 27.9)
45-54	143	49,800	36.1	(26.3 - 45.8)	7,495	816,800	30.1	(25.3 - 34.8)
55-64	133	33,900	32.6	(22.9 - 42.4)	8,356	563,900	27.4	(23.7 - 31.1)
≥ 65	218	37,500	27.6	(20.9 - 34.3)	12,285	554,000	22.7	(19.2 - 26.1)
<b>Race/Ethnicity</b>								
White Non-Hispanic	565	148,700	26.5	(21.6 - 31.3)	34,501	2,062,700	22.9	(21.2 - 24.6)
Black Non-Hispanic	*	*	*	*	708	703,100	34.9	(26.1 - 43.7)
Hispanic	*	*	*	*	813	456,400	20.8	(14.3 - 27.2)
Other Non-Hispanic	*	*	*	*	688	105,200	9.6	(4.0 - 15.3)
<b>Education</b>								
≤ High School	256	80,600	29.7	(21.5 - 37.8)	15,079	1,561,600	29.2	(25.3 - 33.2)
Some College	171	58,700	31.2	(22.0 - 40.3)	9,515	792,800	23.4	(19.4 - 27.3)
≥ College Degree	207	55,300	24.7	(17.0 - 32.5)	12,062	915,000	16.6	(14.4 - 18.8)
<b>Income</b>								
≤ \$24,999	179	44,900	29.5	(19.7 - 39.3)	9,346	974,800	28.3	(22.6 - 34.0)
\$25,000-\$49,999	157	41,000	24.8	(17.1 - 32.6)	9,368	584,800	20.8	(17.1 - 24.5)
\$50,000-\$74,999	*	*	*	*	5,615	475,300	24.6	(20.0 - 29.3)
≥ \$75,000	*	*	*	*	8,337	867,700	19.6	(17.0 - 22.3)

<sup>1</sup> Defined as having a body mass index (BMI) of 30.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who are obese (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 63b. Obesity among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	636	28.0	(22.9 - 33.1)	36,710	23.1	(21.0 - 25.1)
<b>Gender</b>						
Male	234	29.6	(21.2 - 38.1)	14,502	21.5	(19.0 - 24.0)
Female	402	26.5	(20.7 - 32.3)	22,208	24.6	(21.5 - 27.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	565	26.3	(21.1 - 31.5)	34,501	22.8	(21.0 - 24.5)
Black Non-Hispanic	*	*	*	708	34.8	(25.9 - 43.6)
Hispanic	*	*	*	813	20.5	(14.1 - 26.9)
Other Non-Hispanic	*	*	*	688	9.5	(3.8 - 15.2)
<b>Education</b>						
≤ High School	256	29.2	(20.6 - 37.7)	15,079	29.4	(25.4 - 33.3)
Some College	171	30.4	(21.0 - 39.8)	9,515	23.1	(19.2 - 27.1)
≥ College Degree	207	24.8	(16.1 - 33.4)	12,062	16.4	(14.2 - 18.5)
<b>Income</b>						
≤ \$24,999	*	*	*	9,346	28.3	(22.6 - 34.1)
\$25,000-\$49,999	157	24.3	(16.1 - 32.4)	9,368	21.0	(17.2 - 24.8)
\$50,000-\$74,999	*	*	*	5,615	24.0	(19.4 - 28.7)
≥ \$75,000	*	*	*	8,337	19.3	(16.7 - 21.9)

<sup>1</sup> Defined as having a body mass index (BMI) of 30.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 64a. Overweight or obesity among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	636	428,400	62.6	(57.1 - 68.0)	36,710	8,540,500	59.6	(57.0 - 62.2)
<b>Gender</b>								
Male	234	233,500	70.3	(62.0 - 78.5)	14,502	4,579,100	65.2	(61.0 - 69.4)
Female	402	194,900	55.3	(48.6 - 62.0)	22,208	3,961,400	54.3	(51.0 - 57.6)
<b>Age (years)</b>								
18-34	*	*	*	*	3,789	2,025,800	47.1	(40.8 - 53.5)
35-44	*	*	*	*	4,785	1,714,200	61.1	(55.6 - 66.6)
45-54	143	97,000	70.2	(61.4 - 79.0)	7,495	1,870,100	68.8	(64.6 - 73.1)
55-64	133	75,200	72.4	(63.3 - 81.6)	8,356	1,362,000	66.2	(61.9 - 70.4)
≥ 65	218	88,200	64.9	(57.7 - 72.0)	12,285	1,568,400	64.1	(60.6 - 67.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	565	344,000	61.2	(55.5 - 66.9)	34,501	5,344,900	59.3	(57.2 - 61.3)
Black Non-Hispanic	*	*	*	*	708	1,443,100	71.6	(62.5 - 80.7)
Hispanic	*	*	*	*	813	1,237,900	56.4	(46.5 - 66.2)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	256	180,700	66.5	(57.8 - 75.2)	15,079	3,451,400	64.6	(59.8 - 69.4)
Some College	*	*	*	*	9,515	2,107,300	62.1	(57.0 - 67.2)
≥ College Degree	207	128,500	57.5	(48.3 - 66.7)	12,062	2,911,900	52.9	(49.2 - 56.6)
<b>Income</b>								
≤ \$24,999	179	108,500	71.2	(61.3 - 81.2)	9,346	2,145,700	62.3	(55.7 - 68.9)
\$25,000-\$49,999	*	*	*	*	9,368	1,720,700	61.2	(54.5 - 67.9)
\$50,000-\$74,999	*	*	*	*	5,615	1,214,000	62.9	(57.1 - 68.8)
≥ \$75,000	*	*	*	*	8,337	2,526,700	57.1	(53.6 - 60.6)

<sup>1</sup> Defined as having a body mass index (BMI) of 25.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who are overweight or obese (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 64b. Overweight or obesity among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	636	61.4	(55.5 - 67.2)	36,710	59.3	(56.8 - 61.9)
<b>Gender</b>						
Male	234	70.2	(61.8 - 78.6)	14,502	65.4	(61.3 - 69.4)
Female	402	53.0	(45.5 - 60.5)	22,208	53.5	(50.1 - 56.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	565	60.1	(54.0 - 66.2)	34,501	58.8	(56.7 - 60.8)
Black Non-Hispanic	*	*	*	708	71.8	(63.0 - 80.6)
Hispanic	*	*	*	813	56.5	(46.8 - 66.1)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	256	66.1	(56.9 - 75.4)	15,079	64.6	(59.9 - 69.3)
Some College	*	*	*	9,515	61.4	(56.4 - 66.5)
≥ College Degree	207	55.7	(45.7 - 65.7)	12,062	52.6	(48.9 - 56.3)
<b>Income</b>						
≤ \$24,999	*	*	*	9,346	62.1	(55.6 - 68.6)
\$25,000-\$49,999	*	*	*	9,368	61.6	(55.1 - 68.1)
\$50,000-\$74,999	*	*	*	5,615	62.6	(56.7 - 68.6)
≥ \$75,000	*	*	*	8,337	56.5	(52.9 - 60.0)

<sup>1</sup> Defined as having a body mass index (BMI) of 25.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 65a. Received advice about weight by a health professional among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	635	186,700	27.3	(22.7 - 31.9)	36,729	3,929,700	27.4	(25.1 - 29.7)
<b>Gender</b>								
Male	234	82,200	24.8	(17.4 - 32.1)	14,499	1,654,800	23.6	(20.6 - 26.6)
Female	401	104,500	29.7	(24.1 - 35.3)	22,230	2,274,900	31.0	(27.7 - 34.3)
<b>Age (years)</b>								
18-34	*	*	*	*	3,796	895,500	20.7	(15.3 - 26.2)
35-44	*	*	*	*	4,779	722,700	25.8	(20.9 - 30.6)
45-54	142	50,400	36.6	(27.6 - 45.7)	7,490	910,900	33.5	(29.0 - 38.1)
55-64	133	35,500	34.1	(24.4 - 43.9)	8,351	741,100	36.1	(32.0 - 40.3)
≥ 65	218	37,200	27.3	(20.7 - 33.9)	12,313	659,500	26.9	(23.0 - 30.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	565	152,100	27.1	(22.3 - 31.8)	34,502	2,386,200	26.5	(24.6 - 28.4)
Black Non-Hispanic	*	*	*	*	709	615,700	30.4	(22.0 - 38.8)
Hispanic	*	*	*	*	830	555,400	25.1	(17.8 - 32.4)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	255	61,000	22.5	(16.0 - 29.0)	15,116	1,469,700	27.5	(23.6 - 31.4)
Some College	171	62,300	33.1	(23.2 - 43.0)	9,515	946,800	27.8	(23.2 - 32.4)
≥ College Degree	207	62,900	28.1	(20.6 - 35.7)	12,041	1,445,500	26.2	(23.0 - 29.4)
<b>Income</b>								
≤ \$24,999	179	40,500	26.6	(18.2 - 35.0)	9,360	1,079,400	31.4	(25.3 - 37.5)
\$25,000-\$49,999	*	*	*	*	9,365	696,900	24.6	(19.8 - 29.4)
\$50,000-\$74,999	89	22,400	23.6	(13.7 - 33.5)	5,603	507,100	26.3	(21.6 - 31.0)
≥ \$75,000	130	48,600	24.3	(15.4 - 33.1)	8,334	1,207,100	27.3	(24.2 - 30.5)

<sup>1</sup> Defined as being advised by a doctor, nurse or other health professional about their weight (lose, gain or maintain)

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults provided advice by a health professional about their weight (lose, gain or maintain) (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 65b. Received advice about weight by a health professional among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	635	26.6	(21.7 - 31.4)	36,729	27.1	(24.8 - 29.4)
<b>Gender</b>						
Male	234	24.8	(17.1 - 32.6)	14,499	23.4	(20.5 - 26.3)
Female	401	28.3	(22.2 - 34.3)	22,230	30.6	(27.3 - 34.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	565	26.3	(21.3 - 31.4)	34,502	26.1	(24.2 - 28.0)
Black Non-Hispanic	*	*	*	709	30.4	(21.9 - 38.9)
Hispanic	*	*	*	830	24.9	(17.7 - 32.2)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	255	21.7	(14.9 - 28.5)	15,116	27.3	(23.4 - 31.2)
Some College	*	*	*	9,515	27.4	(22.9 - 32.0)
≥ College Degree	207	27.6	(19.3 - 36.0)	12,041	25.9	(22.7 - 29.0)
<b>Income</b>						
≤ \$24,999	179	25.2	(16.5 - 33.8)	9,360	31.0	(24.9 - 37.0)
\$25,000-\$49,999	*	*	*	9,365	24.9	(20.0 - 29.8)
\$50,000-\$74,999	89	21.7	(11.9 - 31.6)	5,603	25.9	(21.2 - 30.6)
≥ \$75,000	130	24.0	(14.6 - 33.3)	8,334	26.9	(23.7 - 30.0)

<sup>1</sup> Defined as being advised by a doctor, nurse or other health professional about their weight (lose, gain or maintain)

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 66a. Received advice to lose weight by a health professional among adults provided weight advice<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	200	153,100	82.0	(73.4 - 90.6)	9,724	3,412,900	86.8	(83.9 - 89.8)
<b>Gender</b>								
Male	*	*	*	*	3,807	1,468,700	88.8	(85.7 - 91.8)
Female	136	87,400	83.7	(73.8 - 93.5)	5,917	1,944,200	85.5	(80.8 - 90.1)
<b>Age (years)</b>								
18-34	*	*	*	*	681	779,700	87.1	(80.6 - 93.6)
35-44	*	*	*	*	1,215	622,100	86.1	(76.9 - 95.2)
45-54	*	*	*	*	2,238	813,300	89.3	(84.4 - 94.2)
55-64	*	*	*	*	2,697	645,500	87.1	(81.2 - 93.0)
≥ 65	63	30,600	82.3	(72.9 - 91.7)	2,893	552,300	83.7	(76.9 - 90.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	*	*	*	*	9,082	2,076,700	87.0	(84.5 - 89.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	251	483,600	87.1	(78.5 - 95.7)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	70	52,700	86.4	(78.0 - 94.8)	3,962	1,239,700	84.3	(78.6 - 90.1)
Some College	*	*	*	*	2,633	814,800	86.1	(80.1 - 92.0)
≥ College Degree	72	56,300	89.5	(81.2 - 97.8)	3,115	1,290,900	89.3	(85.3 - 93.3)
<b>Income</b>								
≤ \$24,999	*	*	*	*	2,539	922,100	85.4	(78.7 - 92.1)
\$25,000-\$49,999	*	*	*	*	2,482	594,100	85.2	(78.4 - 92.1)
\$50,000-\$74,999	*	*	*	*	1,521	446,100	88.0	(83.2 - 92.7)
≥ \$75,000	*	*	*	*	2,270	1,073,200	88.9	(83.7 - 94.1)

<sup>1</sup> Defined as provided advice by a health professional about their weight and advised to lose weight

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults provided advice by a health professional about their weight and advised to lose weight (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 66b. Received advice to lose weight by a health professional among adults provided weight advice<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	200	82.0	(72.6 - 91.4)	9,724	86.9	(83.8 - 89.9)
<b>Gender</b>						
Male	*	*	*	3,807	88.7	(85.7 - 91.8)
Female	*	*	*	5,917	85.5	(80.7 - 90.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	*	*	*	9,082	87.0	(84.4 - 89.6)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	251	87.6	(79.3 - 95.8)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	70	88.1	(80.4 - 95.9)	3,962	84.6	(78.7 - 90.5)
Some College	*	*	*	2,633	85.6	(79.5 - 91.7)
≥ College Degree	72	88.9	(79.4 - 98.4)	3,115	89.4	(85.4 - 93.3)
<b>Income</b>						
≤ \$24,999	*	*	*	2,539	85.7	(79.0 - 92.3)
\$25,000-\$49,999	*	*	*	2,482	85.3	(78.4 - 92.3)
\$50,000-\$74,999	*	*	*	1,521	88.0	(83.1 - 92.8)
≥ \$75,000	*	*	*	2,270	88.7	(83.1 - 94.2)

<sup>1</sup> Defined as provided advice by a health professional about their weight and advised to lose weight

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 67a. No leisure-time physical activity among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	660	144,000	20.3	(16.3 - 24.2)	38,128	3,541,700	23.9	(21.6 - 26.1)
<b>Gender</b>								
Male	237	50,400	15.0	(9.4 - 20.7)	14,602	1,482,500	20.9	(17.8 - 24.0)
Female	423	93,600	24.9	(19.5 - 30.4)	23,526	2,059,200	26.6	(23.4 - 29.7)
<b>Age (years)</b>								
18-34	59	32,800	16.3	(6.8 - 25.8)	3,938	982,900	22.0	(16.6 - 27.5)
35-44	92	10,200	8.4	(3.5 - 13.4)	4,988	556,000	19.3	(15.0 - 23.6)
45-54	149	37,700	26.5	(16.6 - 36.3)	7,827	759,000	26.8	(21.7 - 31.9)
55-64	137	21,900	20.5	(13.4 - 27.7)	8,765	464,900	21.7	(18.2 - 25.2)
≥ 65	223	41,400	29.8	(23.2 - 36.5)	12,610	778,800	30.7	(27.3 - 34.2)
<b>Race/Ethnicity</b>								
White Non-Hispanic	587	114,600	19.6	(15.6 - 23.6)	35,816	1,892,800	20.3	(18.6 - 22.0)
Black Non-Hispanic	*	*	*	*	739	690,000	32.0	(23.4 - 40.6)
Hispanic	*	*	*	*	860	600,800	26.7	(19.1 - 34.4)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	263	73,400	26.2	(19.1 - 33.3)	15,683	1,810,200	32.5	(28.4 - 36.5)
Some College	176	44,800	23.3	(14.7 - 31.8)	9,906	814,200	23.0	(18.6 - 27.3)
≥ College Degree	218	25,200	10.6	(6.2 - 15.1)	12,458	841,000	14.9	(11.9 - 18.0)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,640	1,332,200	38.0	(31.7 - 44.3)
\$25,000-\$49,999	163	43,800	26.0	(16.5 - 35.5)	9,658	648,000	22.3	(18.3 - 26.3)
\$50,000-\$74,999	*	*	*	*	5,773	430,700	21.8	(16.6 - 27.0)
≥ \$75,000	136	17,700	8.4	(4.1 - 12.7)	8,565	560,800	12.5	(10.3 - 14.6)

<sup>1</sup> Defined as having not participated in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have not participated in any physical activities or exercises (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 67b. No leisure-time physical activity among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	660	19.2	(15.1 - 23.3)	38,128	23.7	(21.5 - 25.9)
<b>Gender</b>						
Male	237	14.2	(8.8 - 19.6)	14,602	20.9	(17.8 - 23.9)
Female	423	23.7	(17.7 - 29.6)	23,526	26.3	(23.1 - 29.5)
<b>Race/Ethnicity</b>						
White Non-Hispanic	587	18.5	(14.3 - 22.7)	35,816	20.0	(18.3 - 21.7)
Black Non-Hispanic	*	*	*	739	32.0	(23.3 - 40.6)
Hispanic	*	*	*	860	26.9	(19.2 - 34.5)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	263	25.2	(17.9 - 32.5)	15,683	32.3	(28.3 - 36.4)
Some College	176	23.1	(13.8 - 32.3)	9,906	22.8	(18.5 - 27.2)
≥ College Degree	218	9.4	(5.0 - 13.8)	12,458	14.9	(11.8 - 17.9)
<b>Income</b>						
≤ \$24,999	*	*	*	9,640	37.9	(31.6 - 44.2)
\$25,000-\$49,999	*	*	*	9,658	22.0	(18.0 - 26.0)
\$50,000-\$74,999	*	*	*	5,773	21.6	(16.3 - 26.8)
≥ \$75,000	136	7.5	(3.6 - 11.3)	8,565	12.6	(10.3 - 14.9)

<sup>1</sup> Defined as having not participated in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 68a. Consumption of 5 or more servings of fruits and vegetables per day among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	662	187,500	26.3	(21.6 - 30.9)	38,203	4,023,400	27.0	(24.9 - 29.2)
<b>Gender</b>								
Male	239	64,800	19.2	(12.6 - 25.8)	14,634	1,603,900	22.6	(19.4 - 25.8)
Female	423	122,700	32.7	(26.4 - 38.9)	23,569	2,419,500	31.1	(28.2 - 34.0)
<b>Age (years)</b>								
18-34	*	*	*	*	3,940	1,261,700	28.3	(22.7 - 33.9)
35-44	92	24,100	19.9	(11.0 - 28.8)	4,989	688,000	23.9	(19.5 - 28.3)
45-54	149	36,700	25.8	(17.9 - 33.7)	7,837	746,800	26.2	(22.3 - 30.1)
55-64	137	30,100	28.2	(18.3 - 38.1)	8,777	579,100	27.0	(23.6 - 30.5)
≥ 65	225	41,200	29.1	(22.4 - 35.8)	12,660	747,800	29.4	(26.0 - 32.8)
<b>Race/Ethnicity</b>								
White Non-Hispanic	588	151,300	25.8	(21.0 - 30.6)	35,884	2,538,000	27.2	(25.4 - 28.9)
Black Non-Hispanic	*	*	*	*	740	572,400	26.5	(19.2 - 33.9)
Hispanic	*	*	*	*	864	672,200	29.8	(21.0 - 38.5)
Other Non-Hispanic	*	*	*	*	715	240,800	21.4	(12.0 - 30.8)
<b>Education</b>								
≤ High School	265	60,100	21.3	(14.1 - 28.5)	15,726	1,309,800	23.4	(19.3 - 27.4)
Some College	176	58,900	30.6	(21.5 - 39.7)	9,924	964,300	27.2	(23.0 - 31.3)
≥ College Degree	218	68,400	28.9	(20.8 - 37.0)	12,471	1,747,500	31.0	(27.9 - 34.1)
<b>Income</b>								
≤ \$24,999	180	35,900	23.4	(15.4 - 31.4)	9,667	831,500	23.7	(18.6 - 28.7)
\$25,000-\$49,999	163	37,100	22.0	(13.5 - 30.4)	9,669	917,200	31.6	(25.6 - 37.6)
\$50,000-\$74,999	*	*	*	*	5,777	501,000	25.3	(21.4 - 29.3)
≥ \$75,000	136	50,200	24.0	(14.4 - 33.6)	8,569	1,260,100	28.0	(25.0 - 31.0)

<sup>1</sup> Defined as having consumed 5 or more servings of fruits and vegetables in an average day

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who consume 5 or more servings of fruits and vegetables per day (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 68b. Consumption of 5 or more servings of fruits and vegetables per day among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	662	26.3	(21.3 - 31.3)	38,203	27.1	(24.9 - 29.2)
<b>Gender</b>						
Male	239	18.9	(12.3 - 25.6)	14,634	22.7	(19.6 - 25.9)
Female	423	33.0	(25.9 - 40.0)	23,569	31.0	(28.1 - 34.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	588	25.7	(20.6 - 30.8)	35,884	27.1	(25.4 - 28.9)
Black Non-Hispanic	*	*	*	740	26.8	(19.4 - 34.2)
Hispanic	*	*	*	864	29.5	(20.9 - 38.1)
Other Non-Hispanic	*	*	*	715	21.8	(12.2 - 31.5)
<b>Education</b>						
≤ High School	265	21.5	(13.8 - 29.3)	15,726	23.3	(19.3 - 27.3)
Some College	176	29.7	(20.1 - 39.3)	9,924	27.1	(22.9 - 31.2)
≥ College Degree	218	29.2	(20.2 - 38.1)	12,471	31.2	(28.0 - 34.3)
<b>Income</b>						
≤ \$24,999	180	24.0	(15.1 - 33.0)	9,667	23.9	(18.8 - 28.9)
\$25,000-\$49,999	163	22.4	(13.1 - 31.8)	9,669	32.0	(26.0 - 38.0)
\$50,000-\$74,999	*	*	*	5,777	25.0	(21.0 - 28.9)
≥ \$75,000	136	23.3	(13.7 - 33.0)	8,569	28.0	(24.9 - 31.1)

<sup>1</sup> Defined as having consumed 5 or more servings of fruits and vegetables in an average day

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Erie County**

**Table 69a. Had a fall within the past 3 months among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	504	59,800	15.5	(11.9 - 19.1)	28,783	1,244,800	17.0	(14.9 - 19.0)
<b>Gender</b>								
Male	168	19,300	11.1	(6.3 - 15.9)	10,966	489,400	14.4	(11.9 - 16.8)
Female	336	40,500	19.1	(14.0 - 24.1)	17,817	755,300	19.2	(16.1 - 22.3)
<b>Age (years)</b>								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	148	17,300	12.1	(7.1 - 17.2)	7,741	445,500	16.0	(12.0 - 20.0)
55-64	133	29,800	28.8	(19.4 - 38.2)	8,454	336,000	16.5	(13.4 - 19.6)
≥ 65	223	12,700	9.1	(5.3 - 12.8)	12,588	463,300	18.4	(15.2 - 21.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	451	43,800	13.7	(10.4 - 17.0)	27,456	869,000	16.6	(15.1 - 18.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	424	76,100	11.0	(4.5 - 17.4)
Other Non-Hispanic	*	*	*	*	445	44,300	10.0	(3.1 - 16.8)
<b>Education</b>								
≤ High School	214	23,700	14.3	(9.2 - 19.3)	12,417	493,000	17.3	(13.3 - 21.3)
Some College	134	15,700	15.5	(9.1 - 22.0)	7,232	299,400	18.7	(14.6 - 22.7)
≥ College Degree	153	19,800	17.0	(9.6 - 24.3)	9,086	441,500	15.4	(12.9 - 17.9)
<b>Income</b>								
≤ \$24,999	160	22,600	20.8	(13.4 - 28.2)	7,824	423,900	25.0	(18.5 - 31.6)
\$25,000-\$49,999	122	12,500	13.7	(7.4 - 20.1)	7,445	223,600	16.0	(12.8 - 19.2)
\$50,000-\$74,999	*	*	*	*	4,141	151,200	14.6	(10.8 - 18.4)
≥ \$75,000	85	9,500	11.0	(3.8 - 18.2)	5,874	287,100	12.5	(10.0 - 15.1)

<sup>1</sup> Defined as respondent unintentionally coming to rest on the ground or another lower level one or more times within the past 3 months

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have had a fall within the past 3 months (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 69b. Had a fall within the past 3 months among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	504	15.0	(11.5 - 18.6)	28,783	17.3	(15.0 - 19.5)
<b>Gender</b>						
Male	168	10.4	(5.9 - 14.8)	10,966	15.0	(12.1 - 17.8)
Female	336	19.2	(14.1 - 24.4)	17,817	19.4	(16.0 - 22.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	451	13.4	(10.1 - 16.7)	27,456	16.9	(15.2 - 18.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	424	10.7	(4.2 - 17.1)
Other Non-Hispanic	*	*	*	445	10.0	(2.9 - 17.0)
<b>Education</b>						
≤ High School	214	13.9	(8.8 - 18.9)	12,417	17.4	(13.0 - 21.7)
Some College	134	15.6	(9.0 - 22.2)	7,232	19.0	(14.7 - 23.2)
≥ College Degree	153	16.0	(8.9 - 23.1)	9,086	16.0	(13.1 - 18.8)
<b>Income</b>						
≤ \$24,999	160	20.5	(13.0 - 28.0)	7,824	25.3	(18.3 - 32.4)
\$25,000-\$49,999	122	13.4	(7.0 - 19.7)	7,445	15.9	(12.6 - 19.2)
\$50,000-\$74,999	*	*	*	4,141	14.7	(10.6 - 18.8)
≥ \$75,000	85	10.4	(3.5 - 17.2)	5,874	12.9	(9.9 - 15.8)

<sup>1</sup> Defined as respondent unintentionally coming to rest on the ground or another lower level one or more times within the past 3 months

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Erie County**

**Table 70a. Had a fall that caused injury within the past 3 months among adults<sup>1</sup> in Erie County and New York State**

	Erie County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	504	20,400	5.3	(3.3 - 7.3)	28,760	335,700	4.6	(3.7 - 5.5)
<b>Gender</b>								
Male	*	*	*	*	10,957	120,000	3.5	(2.5 - 4.6)
Female	336	15,200	7.2	(4.2 - 10.1)	17,803	215,600	5.5	(4.1 - 6.9)
<b>Age (years)</b>								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	7,737	115,100	4.1	(2.9 - 5.4)
55-64	133	12,100	11.7	(6.0 - 17.4)	8,449	90,300	4.4	(2.7 - 6.1)
≥ 65	*	*	*	*	12,574	130,300	5.2	(3.4 - 6.9)
<b>Race/Ethnicity</b>								
White Non-Hispanic	451	15,400	4.8	(2.7 - 6.9)	27,433	270,800	5.2	(4.2 - 6.1)
Black Non-Hispanic	*	*	*	*	458	47,900	4.9	(0.9 - 9.0)
Hispanic	*	*	*	*	424	12,400	1.8	(0.2 - 3.3)
Other Non-Hispanic	*	*	*	*	445	4,600	1.0	(0.4 - 1.6)
<b>Education</b>								
≤ High School	214	10,100	6.1	(2.8 - 9.3)	12,404	148,900	5.2	(3.6 - 6.8)
Some College	*	*	*	*	7,228	85,400	5.3	(3.1 - 7.5)
≥ College Degree	*	*	*	*	9,081	101,300	3.5	(2.4 - 4.6)
<b>Income</b>								
≤ \$24,999	160	9,600	8.9	(4.2 - 13.5)	7,812	90,200	5.3	(3.6 - 7.1)
\$25,000-\$49,999	*	*	*	*	7,442	93,200	6.7	(3.9 - 9.4)
\$50,000-\$74,999	*	*	*	*	4,139	37,000	3.6	(1.8 - 5.3)
≥ \$75,000	*	*	*	*	5,872	54,700	2.4	(1.5 - 3.3)

<sup>1</sup> Defined as as respondent unintentionally coming to rest on the ground or another lower level and caused respondent to limit regular activities for at least a day or to go see a doctor one or more times within the past 3 months

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have had a fall that caused an injury within the past 3 months (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Erie County**

**Table 70b. Had a fall that caused injury within the past 3 months among adults<sup>1</sup> in Erie County and New York State - Age-Adjusted<sup>2</sup>**

	Erie County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	504	5.1	(3.1 - 7.0)	28,760	4.6	(3.6 - 5.5)
<b>Gender</b>						
Male	*	*	*	10,957	3.6	(2.4 - 4.7)
Female	336	7.3	(4.2 - 10.3)	17,803	5.5	(4.1 - 6.9)
<b>Race/Ethnicity</b>						
White Non-Hispanic	451	4.7	(2.6 - 6.7)	27,433	5.2	(4.2 - 6.2)
Black Non-Hispanic	*	*	*	458	4.7	(0.7 - 8.7)
Hispanic	*	*	*	424	1.8	(0.1 - 3.4)
Other Non-Hispanic	*	*	*	445	1.0	(0.4 - 1.6)
<b>Education</b>						
≤ High School	214	6.0	(2.7 - 9.3)	12,404	5.2	(3.5 - 6.8)
Some College	*	*	*	7,228	5.3	(3.1 - 7.5)
≥ College Degree	*	*	*	9,081	3.6	(2.4 - 4.7)
<b>Income</b>						
≤ \$24,999	160	8.4	(3.8 - 12.9)	7,812	5.1	(3.3 - 7.0)
\$25,000-\$49,999	*	*	*	7,442	6.6	(3.9 - 9.2)
\$50,000-\$74,999	*	*	*	4,139	3.7	(1.7 - 5.8)
≥ \$75,000	*	*	*	5,872	2.4	(1.4 - 3.3)

<sup>1</sup> Defined as as respondent unintentionally coming to rest on the ground or another lower level and caused respondent to limit regular activities for at least a day or to go see a doctor one or more times within the past 3 months

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)