

New York State Expanded Behavioral Risk Factor Surveillance System

New York City

**Final Report
July 2008 - June 2009**

**New York State Department of Health
December 2009**

Contents

Introduction	12
Table 1. New York City and New York State adult US Census Bureau population estimates, 2007	13
General Health	
Table 2a. Fair or poor general health among adults in New York City and New York State	14
Table 2b. Fair or poor general health among adults in New York City and New York State - Age-adjusted	15
Table 3a. Poor physical health 14 or more days within the past month among adults in New York City and New York State	16
Table 3b. Poor physical health 14 or more days within the past month among adults in New York City and New York State - Age-adjusted	17
Table 4a. Poor mental health 14 or more days within the past month among adults in New York City and New York State	18
Table 4b. Poor mental health 14 or more days within the past month among adults in New York City and New York State - Age-adjusted	19
Table 5a. Poor physical or mental health 14 or more days within the past month among adults in New York City and New York State	20
Table 5b. Poor physical or mental health 14 or more days within the past month among adults in New York City and New York State - Age-adjusted	21
Quality Health Care	
Table 6a. Health insurance among adults in New York City and New York State	22
Table 6b. Health insurance among adults in New York City and New York State - Age-adjusted	23
Table 7a. Health insurance among adults aged 18-64 in New York City and New York State	24
Table 7b. Health insurance among adults aged 18-64 in New York City and New York State - Age-adjusted	25

Contents

Table 8a. No health insurance among adults in New York City and New York State	26
Table 8b. No health insurance among adults in New York City and New York State - Age-adjusted	27
Table 9a. No health insurance among adults aged 18-64 in New York City and New York State	28
Table 9b. No health insurance among adults aged 18-64 in New York City and New York State - Age-adjusted	29
Table 10a. Adults with regular health care provider in New York City and New York State	30
Table 10b. Adults with regular health care provider in New York City and New York State - Age-adjusted	31
Table 11a. Cost prevented visit to doctor within the past year among adults in New York City and New York State	32
Table 11b. Cost prevented visit to doctor within the past year among adults in New York City and New York State - Age-adjusted	33
Table 12a. Visited a doctor for routine checkup within the past year among adults in New York City and New York State	34
Table 12b. Visited a doctor for routine checkup within the past year among adults in New York City and New York State - Age-adjusted	35
Table 13a. Visited a doctor for routine checkup within the past 2 years among adults in New York City and New York State	36
Table 13b. Visited a doctor for routine checkup within the past 2 years among adults in New York City and New York State - Age-adjusted	37
Table 14a. Dentist visit within the past year among adults in New York City and New York State	38
Table 14b. Dentist visit within the past year among adults in New York City and New York State - Age-adjusted	39
Table 15a. Had teeth cleaned within the past year among adults in New York City and New York State	40
Table 15b. Had teeth cleaned within the past year among adults in New York City and New York State - Age-adjusted	41

Contents

Table 16a. Permanent teeth extracted due to decay or gum disease among adults in New York City and New York State	42
Table 16b. Permanent teeth extracted due to decay or gum disease among adults in New York City and New York State - Age-adjusted	43
Table 17a. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older in New York City and New York State	44
Table 17b. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older in New York City and New York State - Age-adjusted	45
Table 18a. Flu shot within the past 12 months in New York City and New York State	46
Table 18b. Flu shot within the past 12 months in New York City and New York State - Age-adjusted	47
Table 19a. Flu vaccine that was sprayed in nose within the past 12 months in New York City and New York State	48
Table 19b. Flu vaccine that was sprayed in nose within the past 12 months in New York City and New York State - Age-adjusted	49
Table 20a. Flu shot or flu vaccine sprayed in nose within the past 12 months in New York City and New York State	50
Table 20b. Flu shot or flu vaccine sprayed in nose within the past 12 months in New York City and New York State - Age-adjusted	51
Table 21a. Flu shot among adults aged 65 and older in New York City and New York State	52
Table 21b. Flu shot among adults aged 65 and older in New York City and New York State - Age-adjusted	53
Table 22a. Pneumonia shot or pneumococcal vaccine in New York City and New York State	54
Table 22b. Pneumonia shot or pneumococcal vaccine in New York City and New York State - Age-adjusted	55
Table 23a. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older in New York City and New York State	56
Table 23b. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older in New York City and New York State - Age-adjusted	57

Contents

Cancer Screening

Table 24a. Ever had a mammogram among women aged 40 and older in New York City and New York State	58
Table 24b. Ever had a mammogram among women aged 40 and older in New York City and New York State - Age-adjusted	59
Table 25a. Had mammogram within the past 2 years among women aged 40 and older in New York City and New York State	60
Table 25b. Had mammogram within the past 2 years among women aged 40 and older in New York City and New York State - Age-adjusted	61
Table 26a. Had mammogram within the past 2 years among women aged 50 and older in New York City and New York State	62
Table 26b. Had mammogram within the past 2 years among women aged 50 and older in New York City and New York State - Age-adjusted	63
Table 27a. Ever had a Pap test among women in New York City and New York State	64
Table 27b. Ever had a Pap test among women in New York City and New York State - Age-adjusted	65
Table 28a. Had a Pap test within the past 3 years among women in New York City and New York State	66
Table 28b. Had a Pap test within the past 3 years among women in New York City and New York State - Age-adjusted	67
Table 29a. Ever had a digital rectal exam among men aged 40 and older in New York City and New York State	68
Table 29b. Ever had a digital rectal exam among men aged 40 and older in New York City and New York State - Age-adjusted	69
Table 30a. Had digital rectal exam within the past 2 years among men aged 40 and older in New York City and New York State	70
Table 30b. Had digital rectal exam within the past 2 years among men aged 40 and older in New York City and New York State - Age-adjusted	71

Contents

Table 31a. Ever had a prostate specific antigen test among men aged 40 and older in New York City and New York State	72
Table 31b. Ever had a prostate specific antigen test among men aged 40 and older in New York City and New York State - Age-adjusted	73
Table 32a. Had prostate specific antigen test within the past 2 years among men aged 40 and older in New York City and New York State	74
Table 32b. Had prostate specific antigen test within the past 2 years among men aged 40 and older in New York City and New York State - Age-adjusted	75
Table 33a. Prostate cancer among men aged 40 and older in New York City and New York State	76
Table 33b. Prostate cancer among men aged 40 and older in New York City and New York State - Age-adjusted	77
Table 34a. Home blood stool test ever used among adults aged 50 and older in New York City and New York State	78
Table 34b. Home blood stool test ever used among adults aged 50 and older in New York City and New York State - Age-adjusted	79
Table 35a. Home blood stool test used within the past year among adults aged 50 and older in New York City and New York State	80
Table 35b. Home blood stool test used within the past year among adults aged 50 and older in New York City and New York State - Age-adjusted	81
Table 36a. Home blood stool test used within the past 2 years among adults aged 50 and older in New York City and New York State	82
Table 36b. Home blood stool test used within the past 2 years among adults aged 50 and older in New York City and New York State - Age-adjusted	83
Table 37a. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in New York City and New York State	84
Table 37b. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in New York City and New York State - Age-adjusted	85
Table 38a. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in New York City and New York State	86

Contents

Table 38b. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in New York City and New York State - Age-adjusted	87
Table 39a. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in New York City and New York State	88
Table 39b. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in New York City and New York State - Age-adjusted	89
Chronic Diseases	
Table 40a. High blood pressure among adults in New York City and New York State	90
Table 40b. High blood pressure among adults in New York City and New York State - Age-adjusted	91
Table 41a. Blood pressure medication among adults with high blood pressure in New York City and New York State	92
Table 41b. Blood pressure medication among adults with high blood pressure in New York City and New York State - Age-adjusted	93
Table 42a. Ever had blood cholesterol checked among adults in New York City and New York State	94
Table 42b. Ever had blood cholesterol checked among adults in New York City and New York State - Age-adjusted	95
Table 43a. Blood cholesterol checked within the past 5 years among adults in New York City and New York State	96
Table 43b. Blood cholesterol checked within the past 5 years among adults in New York City and New York State - Age-adjusted	97
Table 44a. Cardiovascular disease among adults in New York City and New York State	98
Table 44b. Cardiovascular disease among adults in New York City and New York State - Age-adjusted	99
Table 45a. Coronary heart disease among adults in New York City and New York State	100

Contents

Table 45b. Coronary heart disease among adults in New York City and New York State - Age-adjusted	101
Table 46a. Ever diagnosed with diabetes among adults in New York City and New York State	102
Table 46b. Ever diagnosed with diabetes among adults in New York City and New York State - Age-adjusted	103
Table 47a. Ever diagnosed with asthma (lifetime) among adults in New York City and New York State	104
Table 47b. Ever diagnosed with asthma (lifetime) among adults in New York City and New York State - Age-adjusted	105
Table 48a. Current asthma among adults in New York City and New York State	106
Table 48b. Current asthma among adults in New York City and New York State - Age-adjusted	107
Table 49a. Chronic joint symptoms among adults in New York City and New York State	108
Table 49b. Chronic joint symptoms among adults in New York City and New York State - Age-adjusted	109
Table 50a. Arthritis among adults in New York City and New York State	110
Table 50b. Arthritis among adults in New York City and New York State - Age-adjusted	111
Table 51a. Activity limitations because of physical, mental or emotional problems among adults in New York City and New York State	112
Table 51b. Activity limitations because of physical, mental or emotional problems among adults in New York City and New York State - Age-adjusted	113
Table 52a. Have health problem that requires the use of special equipment among adults in New York City and New York State	114
Table 52b. Have health problem that requires the use of special equipment among adults in New York City and New York State - Age-adjusted	115
Table 53a. Disability among adults in New York City and New York State	116
Table 53b. Disability among adults in New York City and New York State - Age-adjusted	117

Contents

Health Behaviors

Table 54a. Current smoking among adults in New York City and New York State	118
Table 54b. Current smoking among adults in New York City and New York State - Age-adjusted	119
Table 55a. Everyday smoking among adults in New York City and New York State	120
Table 55b. Everyday smoking among adults in New York City and New York State - Age-adjusted	121
Table 56a. Adults living in homes in which smoking is prohibited in New York City and New York State	122
Table 56b. Adults living in homes in which smoking is prohibited in New York City and New York State - Age-adjusted	123
Table 57a. Binge drinking within the past month among adults in New York City and New York State	124
Table 57b. Binge drinking within the past month among adults in New York City and New York State - Age-adjusted	125
Table 58a. Heavy drinking within the past month among adults in New York City and New York State	126
Table 58b. Heavy drinking within the past month among adults in New York City and New York State - Age-adjusted	127
Table 59a. Never or rarely been asked about sexual history during routine check-up among adults in New York City and New York State	128
Table 59b. Never or rarely been asked about sexual history during routine check-up among adults in New York City and New York State - Age-adjusted	129
Table 60a. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults in New York City and New York State	130
Table 60b. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults in New York City and New York State - Age-adjusted	131

Contents

Table 61a. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults in New York City and New York State	132
Table 61b. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults in New York City and New York State - Age-adjusted	133

Overweight / Obesity

Table 62a. Overweight among adults in New York City and New York State	134
Table 62b. Overweight among adults in New York City and New York State - Age-adjusted	135
Table 63a. Obesity among adults in New York City and New York State	136
Table 63b. Obesity among adults in New York City and New York State - Age-adjusted	137
Table 64a. Overweight or obesity among adults in New York City and New York State	138
Table 64b. Overweight or obesity among adults in New York City and New York State - Age-adjusted	139
Table 65a. Received advice about weight by a health professional among adults in New York City and New York State	140
Table 65b. Received advice about weight by a health professional among adults in New York City and New York State - Age-adjusted	141
Table 66a. Received advice to lose weight by a health professional among adults provided weight advice in New York City and New York State	142
Table 66b. Received advice to lose weight by a health professional among adults provided weight advice in New York City and New York State - Age-adjusted	143

Physical Activity / Nutrition

Table 67a. No leisure-time physical activity among adults in New York City and New York State	144
Table 67b. No leisure-time physical activity among adults in New York City and New York State - Age-adjusted	145

Contents

Table 68a. Consumption of 5 or more servings of fruits and vegetables per day among adults in New York City and New York State	146
Table 68b. Consumption of 5 or more servings of fruits and vegetables per day among adults in New York City and New York State - Age-adjusted	147

Injury

Table 69a. Had a fall within the past 3 months among adults in New York City and New York State	148
Table 69b. Had a fall within the past 3 months among adults in New York City and New York State - Age-adjusted	149
Table 70a. Had a fall that caused injury within the past 3 months among adults in New York City and New York State	150
Table 70b. Had a fall that caused injury within the past 3 months among adults in New York City and New York State - Age-adjusted	151

Introduction

The New York State Expanded Behavioral Risk Factor Surveillance System (Expanded BRFSS) is a random digit dialed telephone survey of adults (18 years of age and older) that is representative of the non-institutionalized civilian population with landline telephones living in New York State. The goal of the Expanded BRFSS is to collect county-specific data on preventive health practices, risk behaviors, injuries and preventable chronic and infectious diseases. Topics assessed by the Expanded BRFSS include tobacco use, physical inactivity, diet, use of cancer screening services and other factors linked to the leading causes of morbidity and mortality. For each topic, health indicators are based on answers to one or more questions in the survey. The survey results are weighted to more accurately represent the adult population living in each county. Weighting procedures adjust for the differential selection probabilities by race, age and gender. More detailed information on weighting procedures is available in the technical brief.

Survey data collection occurred in 58 geographic areas of New York State, including each of the 57 counties outside of New York City and New York City (5 counties) as a single area. A standard questionnaire was used in all areas. Data collection for the 2009 Expanded BRFSS took place during the period of July 2008 through June 2009. The goal was to complete about 650 interviews in each area. The final number of interviews varies because of the sampling design of the survey and differential response rates. More detailed information on the sampling design is available in the technical brief.

The New York State Department of Health contracted with RTI International, which is responsible for the data collection, sampling, weighting and creation of the statistical reports. RTI International subcontracted with Clearwater Research Inc. for data collection and additional expertise in sample design and data quality control.

This report presents both crude estimates and estimates that have been age-adjusted to the Year 2000 US Standard Population when they represent the entire adult population of an area. Estimates for specific age groups are not age-adjusted. Age adjustment is a statistical process applied to rates that allows communities with different age structures to be compared. Counties can compare their data with results from New York State as a whole and with other counties. Confidence intervals at the 95% level associated with each health indicator give an estimated range of values which is likely to include the actual prevalence of the indicator. Confidence intervals are similar to margins of error. When the confidence intervals of two estimates of the same indicator from different areas do not overlap, they may be said to be statistically significantly different, i.e., these differences are unlikely related to chance and are considered true differences. If there is any value that is included in both intervals, the two estimates are not statistically significantly different.

The survey results for each of the 69 health indicators included in the survey are typically provided by gender, age, race and ethnicity, education and income in the Expanded BRFSS county reports. The estimates for some of these groupings may be suppressed in geographic areas with low proportions of adult respondents in these groupings.

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 1. New York City and New York State adult US Census Bureau population estimates,¹ 2007

	New York City		New York State	
	Adult Population ¹	Percent	Adult Population ¹	Percent
Total	6,376,612	100	14,884,315	100
Gender				
Male	2,980,737	46.7	7,103,791	47.7
Female	3,395,875	53.3	7,780,524	52.3
Age (years)				
18-34	2,054,400	32.2	4,497,479	30.2
35-44	1,319,174	20.7	2,844,115	19.1
45-54	1,142,743	17.9	2,854,316	19.2
55-64	847,250	13.3	2,142,000	14.4
≥ 65	1,013,045	15.9	2,546,405	17.1
Race/Ethnicity				
White Non-Hispanic	2,439,285	38.3	9,343,884	62.8
Black Non-Hispanic	1,485,573	23.3	2,156,941	14.5
Hispanic	1,640,571	25.7	2,258,717	15.2
Other Non-Hispanic	811,183	12.7	1,124,773	7.6

¹ Source: National Center for Health Statistics. Postcensal estimates of the resident population of the United States for July 1, 2000-July 1, 2007, by year, county, age, bridged race, Hispanic origin, and sex (Vintage 2007). Prepared under a collaborative arrangement with the U.S. Census Bureau; released August 7, 2008. Available from: http://www.cdc.gov/nchs/nvss/bridged_race.htm as of November 25, 2009.

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 2a. Fair or poor general health among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	645	1,345,100	21.2	(16.7 - 25.6)	38,124	2,531,400	17.1	(15.1 - 19.0)
Gender								
Male	250	539,000	18.2	(11.9 - 24.4)	14,599	1,069,700	15.1	(12.4 - 17.8)
Female	395	806,100	23.8	(17.7 - 30.0)	23,525	1,461,700	18.8	(16.1 - 21.6)
Age (years)								
18-34	*	*	*	*	3,940	535,800	12.0	(7.2 - 16.8)
35-44	102	140,800	10.7	(3.8 - 17.5)	4,985	289,200	10.0	(6.8 - 13.3)
45-54	123	270,400	23.9	(14.6 - 33.3)	7,819	507,800	17.9	(14.0 - 21.8)
55-64	158	188,300	22.5	(13.0 - 31.9)	8,757	424,700	19.9	(16.0 - 23.9)
≥ 65	187	379,700	37.7	(28.8 - 46.6)	12,623	773,900	30.5	(26.8 - 34.2)
Race/Ethnicity								
White Non-Hispanic	393	344,300	14.2	(8.9 - 19.4)	35,811	1,226,600	13.2	(11.7 - 14.6)
Black Non-Hispanic	93	315,100	21.4	(12.1 - 30.7)	739	457,300	21.3	(14.8 - 27.9)
Hispanic	*	*	*	*	861	684,100	30.4	(22.2 - 38.6)
Other Non-Hispanic	*	*	*	*	713	163,400	14.5	(5.1 - 24.0)
Education								
≤ High School	228	801,300	30.3	(22.6 - 38.1)	15,682	1,466,500	26.3	(22.6 - 30.0)
Some College	118	228,400	18.1	(8.6 - 27.5)	9,904	508,500	14.3	(10.9 - 17.8)
≥ College Degree	295	255,200	10.8	(5.5 - 16.1)	12,457	493,500	8.8	(6.5 - 11.1)
Income								
≤ \$24,999	*	*	*	*	9,633	1,269,200	36.3	(30.1 - 42.4)
\$25,000-\$49,999	106	182,700	15.7	(7.4 - 24.1)	9,649	447,600	15.4	(11.9 - 18.9)
\$50,000-\$74,999	*	*	*	*	5,769	237,800	12.1	(8.0 - 16.2)
≥ \$75,000	*	*	*	*	8,562	218,300	4.9	(3.6 - 6.1)

¹ Defined as reporting fair or poor general health

² Sample size from survey

³ Estimated number of adults who have fair or poor health (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 2b. Fair or poor general health among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	645	21.3	(16.9 - 25.7)	38,124	16.7	(14.8 - 18.6)
Gender						
Male	250	18.9	(12.7 - 25.2)	14,599	15.3	(12.6 - 18.0)
Female	395	23.3	(17.1 - 29.4)	23,525	18.0	(15.2 - 20.7)
Race/Ethnicity						
White Non-Hispanic	393	14.9	(9.5 - 20.2)	35,811	12.9	(11.4 - 14.4)
Black Non-Hispanic	93	21.2	(11.9 - 30.5)	739	20.8	(14.2 - 27.3)
Hispanic	*	*	*	861	29.9	(21.8 - 37.9)
Other Non-Hispanic	*	*	*	713	13.8	(4.8 - 22.8)
Education						
≤ High School	228	30.1	(22.5 - 37.8)	15,682	25.8	(22.1 - 29.5)
Some College	118	18.5	(9.0 - 28.1)	9,904	14.1	(10.7 - 17.6)
≥ College Degree	295	10.9	(5.7 - 16.1)	12,457	8.5	(6.3 - 10.8)
Income						
≤ \$24,999	*	*	*	9,633	35.9	(29.7 - 42.0)
\$25,000-\$49,999	106	15.7	(7.3 - 24.1)	9,649	15.0	(11.6 - 18.5)
\$50,000-\$74,999	*	*	*	5,769	11.8	(7.7 - 16.0)
≥ \$75,000	*	*	*	8,562	4.7	(3.5 - 6.0)

¹ Defined as reporting fair or poor general health

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 3a. Poor physical health 14 or more days within the past month among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	624	621,200	10.0	(6.8 - 13.3)	37,327	1,472,900	10.1	(8.7 - 11.6)
Gender								
Male	241	244,800	8.5	(4.5 - 12.5)	14,325	604,200	8.7	(7.0 - 10.5)
Female	383	376,400	11.4	(6.4 - 16.4)	23,002	868,600	11.4	(9.2 - 13.6)
Age (years)								
18-34	*	*	*	*	3,891	268,100	6.1	(2.7 - 9.5)
35-44	*	*	*	*	4,948	232,100	8.1	(5.2 - 11.1)
45-54	121	160,600	14.4	(6.9 - 22.0)	7,764	350,900	12.5	(9.4 - 15.6)
55-64	155	100,200	12.3	(5.9 - 18.7)	8,621	271,200	13.0	(10.3 - 15.8)
≥ 65	176	105,900	11.2	(5.5 - 16.8)	12,103	350,600	14.5	(12.1 - 16.9)
Race/Ethnicity								
White Non-Hispanic	377	233,400	10.0	(4.8 - 15.1)	35,073	929,700	10.2	(8.8 - 11.6)
Black Non-Hispanic	92	131,500	9.0	(1.9 - 16.2)	724	208,200	9.9	(4.8 - 14.9)
Hispanic	111	191,600	12.1	(5.4 - 18.9)	835	250,300	11.5	(6.5 - 16.4)
Other Non-Hispanic	*	*	*	*	695	84,600	7.6	(1.3 - 13.9)
Education								
≤ High School	213	266,700	10.7	(5.8 - 15.5)	15,178	692,800	12.9	(10.6 - 15.3)
Some College	118	138,100	10.9	(3.3 - 18.6)	9,751	362,500	10.3	(7.4 - 13.2)
≥ College Degree	290	160,900	6.9	(3.2 - 10.6)	12,327	359,300	6.4	(4.8 - 8.1)
Income								
≤ \$24,999	156	328,100	17.2	(8.7 - 25.8)	9,294	625,600	18.4	(13.6 - 23.3)
\$25,000-\$49,999	104	94,700	8.3	(2.6 - 14.1)	9,495	279,100	9.8	(7.3 - 12.3)
\$50,000-\$74,999	*	*	*	*	5,733	186,400	9.5	(5.8 - 13.2)
≥ \$75,000	*	*	*	*	8,510	219,100	4.9	(3.6 - 6.2)

¹ Defined as 14 or more days within the past month when physical health was not good

² Sample size from survey

³ Estimated number of adults with poor physical health 14 or more days within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 3b. Poor physical health 14 or more days within the past month among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	624	10.2	(6.8 - 13.5)	37,327	9.9	(8.4 - 11.4)
Gender						
Male	241	8.8	(4.6 - 12.9)	14,325	8.7	(6.9 - 10.5)
Female	383	11.4	(6.4 - 16.4)	23,002	11.0	(8.7 - 13.2)
Race/Ethnicity						
White Non-Hispanic	377	10.1	(4.9 - 15.3)	35,073	9.9	(8.5 - 11.3)
Black Non-Hispanic	92	9.2	(1.9 - 16.6)	724	9.7	(4.6 - 14.9)
Hispanic	111	12.3	(5.5 - 19.1)	835	11.3	(6.3 - 16.2)
Other Non-Hispanic	*	*	*	695	7.6	(1.2 - 14.0)
Education						
≤ High School	213	10.8	(5.8 - 15.7)	15,178	12.7	(10.2 - 15.1)
Some College	118	11.2	(3.5 - 18.9)	9,751	10.1	(7.2 - 12.9)
≥ College Degree	290	7.0	(3.2 - 10.7)	12,327	6.3	(4.7 - 8.0)
Income						
≤ \$24,999	156	17.8	(9.1 - 26.5)	9,294	18.4	(13.5 - 23.4)
\$25,000-\$49,999	104	8.3	(2.6 - 14.0)	9,495	9.4	(7.0 - 11.9)
\$50,000-\$74,999	*	*	*	5,733	9.0	(5.3 - 12.6)
≥ \$75,000	*	*	*	8,510	4.9	(3.5 - 6.2)

¹ Defined as 14 or more days within the past month when physical health was not good

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 4a. Poor mental health 14 or more days within the past month among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	628	571,900	9.2	(5.9 - 12.4)	37,528	1,473,200	10.1	(8.6 - 11.5)
Gender								
Male	245	189,200	6.5	(2.8 - 10.2)	14,407	555,200	7.9	(6.3 - 9.6)
Female	383	382,700	11.5	(6.4 - 16.6)	23,121	918,000	12.0	(9.7 - 14.3)
Age (years)								
18-34	*	*	*	*	3,889	496,700	11.2	(7.4 - 15.1)
35-44	*	*	*	*	4,947	254,100	8.9	(6.7 - 11.1)
45-54	122	143,400	12.7	(4.8 - 20.6)	7,749	347,200	12.3	(9.0 - 15.6)
55-64	153	77,000	9.4	(4.3 - 14.6)	8,623	211,300	10.1	(7.9 - 12.3)
≥ 65	179	78,600	8.3	(3.3 - 13.3)	12,320	163,900	6.7	(4.7 - 8.7)
Race/Ethnicity								
White Non-Hispanic	383	196,400	8.3	(3.3 - 13.2)	35,266	899,200	9.8	(8.4 - 11.2)
Black Non-Hispanic	*	*	*	*	722	227,800	10.9	(5.4 - 16.4)
Hispanic	113	185,100	11.4	(4.7 - 18.1)	845	263,700	11.8	(6.8 - 16.8)
Other Non-Hispanic	*	*	*	*	695	82,500	7.4	(1.4 - 13.5)
Education								
≤ High School	218	245,900	9.6	(5.3 - 14.0)	15,335	665,000	12.2	(10.0 - 14.5)
Some College	116	153,400	12.2	(3.8 - 20.5)	9,771	404,500	11.5	(8.3 - 14.8)
≥ College Degree	290	117,100	5.0	(1.3 - 8.7)	12,342	346,600	6.2	(4.5 - 7.9)
Income								
≤ \$24,999	161	336,200	17.5	(9.0 - 26.1)	9,398	611,500	17.9	(13.0 - 22.8)
\$25,000-\$49,999	*	*	*	*	9,547	260,300	9.0	(6.7 - 11.4)
\$50,000-\$74,999	*	*	*	*	5,726	187,300	9.6	(6.0 - 13.1)
≥ \$75,000	*	*	*	*	8,520	276,500	6.2	(4.6 - 7.7)

¹ Defined as 14 or more days within the past month when mental health was not good

² Sample size from survey

³ Estimated number of adults with poor mental health 14 or more days within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 4b. Poor mental health 14 or more days within the past month among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	628	9.2	(5.9 - 12.4)	37,528	10.2	(8.7 - 11.7)
Gender						
Male	245	6.5	(2.9 - 10.2)	14,407	7.9	(6.3 - 9.6)
Female	383	11.5	(6.3 - 16.6)	23,121	12.2	(9.8 - 14.6)
Race/Ethnicity						
White Non-Hispanic	383	8.3	(3.3 - 13.3)	35,266	10.0	(8.5 - 11.4)
Black Non-Hispanic	*	*	*	722	11.1	(5.5 - 16.7)
Hispanic	113	11.2	(4.6 - 17.8)	845	11.5	(6.5 - 16.4)
Other Non-Hispanic	*	*	*	695	7.6	(1.5 - 13.7)
Education						
≤ High School	218	9.6	(5.2 - 14.0)	15,335	12.3	(10.0 - 14.6)
Some College	116	12.2	(3.9 - 20.6)	9,771	11.8	(8.5 - 15.1)
≥ College Degree	290	5.0	(1.3 - 8.7)	12,342	6.3	(4.6 - 8.0)
Income						
≤ \$24,999	161	17.5	(9.0 - 26.1)	9,398	18.2	(13.2 - 23.1)
\$25,000-\$49,999	*	*	*	9,547	9.4	(6.9 - 11.9)
\$50,000-\$74,999	*	*	*	5,726	9.5	(5.9 - 13.1)
≥ \$75,000	*	*	*	8,520	6.2	(4.6 - 7.7)

¹ Defined as 14 or more days within the past month when mental health was not good

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

**Table 5a. Poor physical or mental health 14 or more days within the past month among adults¹
in New York City and New York State**

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	616	900,300	14.6	(10.8 - 18.5)	37,005	2,385,900	16.5	(14.8 - 18.2)
Gender								
Male	239	327,800	11.5	(6.7 - 16.3)	14,231	937,700	13.6	(11.4 - 15.8)
Female	377	572,500	17.4	(11.7 - 23.1)	22,774	1,448,200	19.1	(16.5 - 21.7)
Age (years)								
18-34	*	*	*	*	3,859	602,600	13.8	(9.7 - 17.9)
35-44	99	133,200	10.3	(3.6 - 17.0)	4,924	388,300	13.7	(10.4 - 17.0)
45-54	121	228,700	20.6	(11.5 - 29.6)	7,722	551,000	19.6	(15.8 - 23.4)
55-64	151	146,900	18.3	(10.8 - 25.7)	8,545	395,500	19.2	(16.1 - 22.3)
≥ 65	174	158,500	16.9	(9.9 - 23.8)	11,955	448,500	18.8	(15.9 - 21.7)
Race/Ethnicity								
White Non-Hispanic	374	297,600	12.7	(7.4 - 18.0)	34,778	1,485,700	16.3	(14.8 - 17.8)
Black Non-Hispanic	89	199,000	13.9	(5.4 - 22.4)	715	343,700	16.5	(10.3 - 22.7)
Hispanic	110	301,600	19.2	(10.6 - 27.8)	832	412,900	19.0	(12.6 - 25.3)
Other Non-Hispanic	*	*	*	*	680	143,500	13.0	(5.0 - 21.0)
Education								
≤ High School	209	390,400	15.7	(10.0 - 21.5)	15,018	1,077,200	20.3	(17.3 - 23.2)
Some College	116	196,700	15.6	(6.8 - 24.4)	9,672	615,500	17.6	(14.1 - 21.1)
≥ College Degree	288	257,800	11.0	(6.0 - 16.1)	12,244	634,200	11.4	(9.2 - 13.6)
Income								
≤ \$24,999	155	468,300	24.6	(15.2 - 34.1)	9,199	924,000	27.3	(21.9 - 32.8)
\$25,000-\$49,999	103	102,800	9.1	(3.2 - 15.0)	9,425	456,000	16.1	(13.0 - 19.2)
\$50,000-\$74,999	*	*	*	*	5,696	312,000	16.0	(11.4 - 20.5)
≥ \$75,000	187	116,300	7.6	(2.7 - 12.5)	8,475	447,600	10.0	(8.1 - 12.0)

¹ Defined as 14 or more days within the past month when physical or mental health prevented usual activities, such as self-care, work or recreation

² Sample size from survey

³ Estimated number of adults with poor physical or mental health 14 or more days within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 5b. Poor physical or mental health 14 or more days within the past month among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	616	14.7	(10.9 - 18.6)	37,005	16.4	(14.7 - 18.2)
Gender						
Male	239	11.7	(6.8 - 16.5)	14,231	13.6	(11.4 - 15.8)
Female	377	17.4	(11.6 - 23.1)	22,774	19.0	(16.3 - 21.6)
Race/Ethnicity						
White Non-Hispanic	374	12.8	(7.4 - 18.2)	34,778	16.3	(14.7 - 17.8)
Black Non-Hispanic	89	14.2	(5.5 - 22.9)	715	16.7	(10.3 - 23.0)
Hispanic	110	19.1	(10.5 - 27.6)	832	18.5	(12.2 - 24.8)
Other Non-Hispanic	*	*	*	680	13.1	(5.0 - 21.3)
Education						
≤ High School	209	15.8	(9.9 - 21.7)	15,018	20.1	(17.1 - 23.1)
Some College	116	15.7	(6.9 - 24.5)	9,672	17.6	(14.1 - 21.1)
≥ College Degree	288	11.1	(6.0 - 16.2)	12,244	11.4	(9.2 - 13.7)
Income						
≤ \$24,999	155	25.1	(15.6 - 34.7)	9,199	27.6	(22.1 - 33.1)
\$25,000-\$49,999	103	9.1	(3.2 - 15.0)	9,425	16.1	(12.9 - 19.3)
\$50,000-\$74,999	*	*	*	5,696	15.6	(11.0 - 20.1)
≥ \$75,000	187	7.7	(2.7 - 12.7)	8,475	10.0	(8.0 - 12.0)

¹ Defined as 14 or more days within the past month when physical or mental health prevented usual activities, such as self-care, work or recreation

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 6a. Health insurance among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	647	5,214,300	82.0	(76.5 - 87.4)	38,119	12,854,200	86.6	(84.2 - 89.0)
Gender								
Male	251	2,260,900	75.9	(66.7 - 85.0)	14,597	5,881,100	83.0	(78.9 - 87.1)
Female	396	2,953,400	87.4	(81.4 - 93.3)	23,522	6,973,100	89.9	(87.2 - 92.6)
Age (years)								
18-34	*	*	*	*	3,910	3,302,100	74.3	(67.8 - 80.7)
35-44	102	1,153,000	87.4	(78.5 - 96.3)	4,986	2,560,300	88.9	(84.7 - 93.1)
45-54	*	*	*	*	7,831	2,530,100	88.9	(84.6 - 93.2)
55-64	160	793,700	93.7	(87.4 - 99.9)	8,764	1,986,700	92.8	(90.2 - 95.5)
≥ 65	187	971,700	96.9	(94.5 - 99.4)	12,628	2,475,000	97.8	(96.7 - 98.8)
Race/Ethnicity								
White Non-Hispanic	394	2,232,600	91.7	(85.9 - 97.6)	35,811	8,591,200	92.2	(90.6 - 93.8)
Black Non-Hispanic	*	*	*	*	736	1,781,600	82.7	(75.3 - 90.1)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	231	1,989,400	74.8	(65.2 - 84.4)	15,673	4,448,900	79.7	(74.9 - 84.5)
Some College	*	*	*	*	9,906	3,004,600	85.0	(80.1 - 89.8)
≥ College Degree	295	2,241,900	94.8	(90.4 - 99.2)	12,459	5,361,700	95.3	(93.3 - 97.2)
Income								
≤ \$24,999	*	*	*	*	9,640	2,437,200	69.7	(62.5 - 76.8)
\$25,000-\$49,999	*	*	*	*	9,662	2,384,000	82.2	(76.1 - 88.3)
\$50,000-\$74,999	77	663,500	93.3	(86.2 - 100)	5,772	1,855,700	94.0	(91.2 - 96.7)
≥ \$75,000	189	1,502,900	97.8	(95.7 - 100)	8,567	4,388,400	97.5	(96.6 - 98.5)

¹ Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

² Sample size from survey

³ Estimated number of adults who have health insurance (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 6b. Health insurance among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	647	82.5	(77.2 - 87.8)	38,119	86.7	(84.4 - 89.1)
Gender						
Male	251	77.1	(68.4 - 85.8)	14,597	83.6	(79.7 - 87.5)
Female	396	87.3	(81.2 - 93.3)	23,522	89.6	(86.8 - 92.3)
Race/Ethnicity						
White Non-Hispanic	394	91.9	(86.1 - 97.7)	35,811	92.1	(90.5 - 93.7)
Black Non-Hispanic	*	*	*	736	82.5	(75.0 - 90.0)
Hispanic	*	*	*	860	68.7	(59.0 - 78.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	231	76.0	(66.7 - 85.2)	15,673	80.1	(75.4 - 84.7)
Some College	*	*	*	9,906	84.6	(79.7 - 89.4)
≥ College Degree	295	95.1	(90.9 - 99.2)	12,459	95.4	(93.6 - 97.2)
Income						
≤ \$24,999	*	*	*	9,640	70.2	(63.2 - 77.1)
\$25,000-\$49,999	*	*	*	9,662	81.9	(76.0 - 87.9)
\$50,000-\$74,999	77	93.5	(86.6 - 100)	5,772	94.0	(91.3 - 96.7)
≥ \$75,000	189	97.9	(95.8 - 100)	8,567	97.5	(96.6 - 98.5)

¹ Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 7a. Health insurance among adults aged 18-64¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	448	4,169,500	78.9	(72.5 - 85.3)	25,165	10,251,300	84.2	(81.3 - 87.1)
Gender								
Male	*	*	*	*	10,078	4,839,900	80.3	(75.5 - 85.0)
Female	256	2,306,800	85.4	(78.1 - 92.7)	15,087	5,411,400	88.0	(84.7 - 91.3)
Age (years)								
18-34	*	*	*	*	3,910	3,302,100	74.3	(67.8 - 80.7)
35-44	102	1,153,000	87.4	(78.5 - 96.3)	4,981	2,556,600	88.9	(84.7 - 93.1)
45-54	*	*	*	*	7,791	2,472,900	88.7	(84.3 - 93.0)
55-64	152	767,000	93.7	(87.3 - 100)	8,483	1,919,700	92.8	(90.1 - 95.5)
≥ 65	-	-	-	-	-	-	-	-
Race/Ethnicity								
White Non-Hispanic	247	1,653,900	89.5	(81.9 - 97.1)	23,354	6,582,200	90.3	(88.3 - 92.4)
Black Non-Hispanic	*	*	*	*	567	1,464,100	80.2	(71.6 - 88.7)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	9,037	3,191,500	74.6	(68.6 - 80.6)
Some College	*	*	*	*	7,031	2,540,400	82.9	(77.4 - 88.4)
≥ College Degree	218	1,906,800	94.1	(89.0 - 99.2)	9,068	4,502,400	94.5	(92.2 - 96.8)
Income								
≤ \$24,999	*	*	*	*	4,918	1,707,200	62.6	(54.1 - 71.2)
\$25,000-\$49,999	*	*	*	*	6,180	1,783,000	77.7	(70.3 - 85.1)
\$50,000-\$74,999	55	552,200	92.1	(83.6 - 100)	4,554	1,553,400	92.9	(89.7 - 96.1)
≥ \$75,000	162	1,356,300	97.6	(95.2 - 100)	7,324	3,987,600	97.3	(96.3 - 98.3)

¹ Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

² Sample size from survey

³ Estimated number of adults aged 18-64 who have health insurance (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 7b. Health insurance among adults aged 18-64¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	448	79.3	(73.0 - 85.5)	25,165	84.3	(81.5 - 87.2)
Gender						
Male	*	*	*	10,078	80.5	(75.9 - 85.1)
Female	256	85.4	(78.1 - 92.7)	15,087	87.9	(84.6 - 91.2)
Race/Ethnicity						
White Non-Hispanic	247	89.4	(81.8 - 97.1)	23,354	90.3	(88.3 - 92.3)
Black Non-Hispanic	*	*	*	567	80.0	(71.3 - 88.6)
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	9,037	75.1	(69.3 - 80.9)
Some College	*	*	*	7,031	82.6	(77.1 - 88.1)
≥ College Degree	218	94.3	(89.4 - 99.1)	9,068	94.6	(92.5 - 96.8)
Income						
≤ \$24,999	*	*	*	4,918	63.2	(54.9 - 71.6)
\$25,000-\$49,999	*	*	*	6,180	77.5	(70.2 - 84.7)
\$50,000-\$74,999	55	92.2	(83.8 - 100)	4,554	93.0	(89.8 - 96.2)
≥ \$75,000	162	97.6	(95.2 - 100)	7,324	97.3	(96.3 - 98.3)

¹ Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 8a. No health insurance among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	647	1,146,100	18.0	(12.6 - 23.5)	38,119	1,989,300	13.4	(11.0 - 15.8)
Gender								
Male	251	719,800	24.1	(15.0 - 33.3)	14,597	1,207,400	17.0	(12.9 - 21.1)
Female	396	426,300	12.6	(6.7 - 18.6)	23,522	781,900	10.1	(7.4 - 12.8)
Age (years)								
18-34	*	*	*	*	3,910	1,144,100	25.7	(19.3 - 32.2)
35-44	102	166,200	12.6	(3.7 - 21.5)	4,986	319,200	11.1	(6.9 - 15.3)
45-54	*	*	*	*	7,831	316,500	11.1	(6.8 - 15.4)
55-64	*	*	*	*	8,764	153,300	7.2	(4.5 - 9.8)
≥ 65	*	*	*	*	12,628	56,200	2.2	(1.2 - 3.3)
Race/Ethnicity								
White Non-Hispanic	394	201,200	8.3	(2.4 - 14.1)	35,811	727,100	7.8	(6.2 - 9.4)
Black Non-Hispanic	*	*	*	*	736	372,800	17.3	(9.9 - 24.7)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	231	670,100	25.2	(15.6 - 34.8)	15,673	1,134,400	20.3	(15.5 - 25.1)
Some College	*	*	*	*	9,906	531,400	15.0	(10.2 - 19.9)
≥ College Degree	295	122,500	5.2	(0.8 - 9.6)	12,459	266,300	4.7	(2.8 - 6.7)
Income								
≤ \$24,999	*	*	*	*	9,640	1,061,200	30.3	(23.2 - 37.5)
\$25,000-\$49,999	*	*	*	*	9,662	516,700	17.8	(11.7 - 23.9)
\$50,000-\$74,999	*	*	*	*	5,772	119,100	6.0	(3.3 - 8.8)
≥ \$75,000	*	*	*	*	8,567	111,100	2.5	(1.5 - 3.4)

¹ Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

² Sample size from survey

³ Estimated number of adults who have no health insurance (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 8b. No health insurance among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	647	17.5	(12.2 - 22.8)	38,119	13.3	(10.9 - 15.6)
Gender						
Male	251	22.9	(14.2 - 31.6)	14,597	16.4	(12.5 - 20.3)
Female	396	12.7	(6.7 - 18.8)	23,522	10.4	(7.7 - 13.2)
Race/Ethnicity						
White Non-Hispanic	394	8.1	(2.3 - 13.9)	35,811	7.9	(6.3 - 9.5)
Black Non-Hispanic	*	*	*	736	17.5	(10.0 - 25.0)
Hispanic	*	*	*	860	31.3	(21.6 - 41.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	231	24.0	(14.8 - 33.3)	15,673	19.9	(15.3 - 24.6)
Some College	*	*	*	9,906	15.4	(10.6 - 20.3)
≥ College Degree	295	4.9	(0.8 - 9.1)	12,459	4.6	(2.8 - 6.4)
Income						
≤ \$24,999	*	*	*	9,640	29.8	(22.9 - 36.8)
\$25,000-\$49,999	*	*	*	9,662	18.1	(12.1 - 24.0)
\$50,000-\$74,999	*	*	*	5,772	6.0	(3.3 - 8.7)
≥ \$75,000	*	*	*	8,567	2.5	(1.5 - 3.4)

¹ Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 9a. No health insurance among adults aged 18-64¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	448	1,113,100	21.1	(14.7 - 27.5)	25,165	1,928,400	15.8	(12.9 - 18.7)
Gender								
Male	*	*	*	*	10,078	1,190,200	19.7	(15.0 - 24.5)
Female	256	393,300	14.6	(7.3 - 21.9)	15,087	738,200	12.0	(8.7 - 15.3)
Age (years)								
18-34	*	*	*	*	3,910	1,144,100	25.7	(19.3 - 32.2)
35-44	102	166,200	12.6	(3.7 - 21.5)	4,981	319,200	11.1	(6.9 - 15.3)
45-54	*	*	*	*	7,791	316,100	11.3	(7.0 - 15.7)
55-64	*	*	*	*	8,483	148,900	7.2	(4.5 - 9.9)
≥ 65	-	-	-	-	-	-	-	-
Race/Ethnicity								
White Non-Hispanic	247	194,100	10.5	(2.9 - 18.1)	23,354	703,100	9.7	(7.6 - 11.7)
Black Non-Hispanic	*	*	*	*	567	361,800	19.8	(11.3 - 28.4)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	9,037	1,087,700	25.4	(19.4 - 31.4)
Some College	*	*	*	*	7,031	522,800	17.1	(11.6 - 22.6)
≥ College Degree	218	120,100	5.9	(0.8 - 11.0)	9,068	261,200	5.5	(3.2 - 7.8)
Income								
≤ \$24,999	*	*	*	*	4,918	1,018,000	37.4	(28.8 - 45.9)
\$25,000-\$49,999	*	*	*	*	6,180	511,100	22.3	(14.9 - 29.7)
\$50,000-\$74,999	*	*	*	*	4,554	118,800	7.1	(3.9 - 10.3)
≥ \$75,000	*	*	*	*	7,324	110,700	2.7	(1.7 - 3.7)

¹ Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

² Sample size from survey

³ Estimated number of adults aged 18-64 who have no health insurance (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 9b. No health insurance among adults aged 18-64¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	448	20.7	(14.5 - 27.0)	25,165	15.7	(12.8 - 18.5)
Gender						
Male	*	*	*	10,078	19.5	(14.9 - 24.1)
Female	256	14.6	(7.3 - 21.9)	15,087	12.1	(8.8 - 15.4)
Race/Ethnicity						
White Non-Hispanic	247	10.6	(2.9 - 18.2)	23,354	9.7	(7.7 - 11.7)
Black Non-Hispanic	*	*	*	567	20.0	(11.4 - 28.7)
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	9,037	24.9	(19.1 - 30.7)
Some College	*	*	*	7,031	17.4	(11.9 - 22.9)
≥ College Degree	218	5.7	(0.9 - 10.6)	9,068	5.4	(3.2 - 7.5)
Income						
≤ \$24,999	*	*	*	4,918	36.8	(28.4 - 45.1)
\$25,000-\$49,999	*	*	*	6,180	22.5	(15.3 - 29.8)
\$50,000-\$74,999	*	*	*	4,554	7.0	(3.8 - 10.2)
≥ \$75,000	*	*	*	7,324	2.7	(1.7 - 3.7)

¹ Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 10a. Adults with regular health care provider¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	648	4,895,900	76.9	(71.2 - 82.6)	38,135	12,324,300	83.0	(80.4 - 85.6)
Gender								
Male	250	2,061,400	69.4	(59.7 - 79.0)	14,593	5,494,300	77.6	(73.2 - 82.0)
Female	398	2,834,500	83.5	(77.6 - 89.4)	23,542	6,830,000	87.8	(85.1 - 90.5)
Age (years)								
18-34	*	*	*	*	3,929	2,931,600	65.8	(59.2 - 72.5)
35-44	101	1,088,600	83.1	(74.1 - 92.0)	4,988	2,471,000	86.1	(81.9 - 90.3)
45-54	124	967,500	84.7	(75.1 - 94.3)	7,823	2,529,300	88.6	(84.6 - 92.7)
55-64	160	796,800	94.0	(89.7 - 98.4)	8,770	2,002,000	93.6	(91.7 - 95.4)
≥ 65	188	923,000	91.1	(86.4 - 95.8)	12,625	2,390,400	94.1	(92.2 - 96.0)
Race/Ethnicity								
White Non-Hispanic	395	2,070,900	84.9	(78.8 - 91.0)	35,821	8,227,700	88.2	(86.4 - 89.9)
Black Non-Hispanic	*	*	*	*	740	1,753,200	81.3	(72.6 - 89.9)
Hispanic	*	*	*	*	860	1,453,100	64.7	(54.7 - 74.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	232	1,838,900	68.9	(59.2 - 78.5)	15,692	4,249,200	75.9	(70.9 - 80.8)
Some College	*	*	*	*	9,904	2,935,000	82.8	(77.2 - 88.4)
≥ College Degree	294	2,074,600	88.1	(81.8 - 94.3)	12,456	5,046,100	89.8	(87.1 - 92.6)
Income								
≤ \$24,999	*	*	*	*	9,644	2,554,200	72.7	(65.9 - 79.6)
\$25,000-\$49,999	*	*	*	*	9,655	2,243,100	77.3	(70.0 - 84.6)
\$50,000-\$74,999	*	*	*	*	5,771	1,767,500	89.5	(85.2 - 93.9)
≥ \$75,000	188	1,425,400	93.3	(89.0 - 97.6)	8,560	4,122,800	91.9	(90.2 - 93.7)

¹ Defined as having one or more persons respondent thinks of as a personal doctor or health care provider

² Sample size from survey

³ Estimated number of adults who have at least one person they think of as their personal doctor or health care provider (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 10b. Adults with regular health care provider¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	648	77.5	(72.0 - 83.0)	38,135	83.0	(80.4 - 85.5)
Gender						
Male	250	70.8	(61.6 - 80.1)	14,593	78.1	(73.9 - 82.3)
Female	398	83.3	(77.3 - 89.2)	23,542	87.4	(84.6 - 90.1)
Race/Ethnicity						
White Non-Hispanic	395	85.2	(79.3 - 91.1)	35,821	88.0	(86.2 - 89.7)
Black Non-Hispanic	*	*	*	740	81.4	(73.0 - 89.8)
Hispanic	*	*	*	860	65.2	(55.5 - 75.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	232	69.7	(60.3 - 79.1)	15,692	75.8	(71.0 - 80.6)
Some College	*	*	*	9,904	82.9	(77.5 - 88.3)
≥ College Degree	294	88.2	(82.2 - 94.3)	12,456	89.8	(87.1 - 92.4)
Income						
≤ \$24,999	*	*	*	9,644	72.9	(66.1 - 79.6)
\$25,000-\$49,999	*	*	*	9,655	77.4	(70.5 - 84.3)
\$50,000-\$74,999	*	*	*	5,771	89.6	(85.3 - 94.0)
≥ \$75,000	188	93.2	(88.8 - 97.6)	8,560	91.6	(89.8 - 93.4)

¹ Defined as having one or more persons respondent thinks of as a personal doctor or health care provider

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 11a. Cost prevented visit to doctor within the past year among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	646	1,047,100	16.5	(11.7 - 21.3)	38,143	2,034,800	13.7	(11.6 - 15.8)
Gender								
Male	249	462,100	15.5	(8.2 - 22.9)	14,607	903,300	12.7	(9.5 - 16.0)
Female	397	585,100	17.3	(11.1 - 23.5)	23,536	1,131,500	14.6	(11.8 - 17.4)
Age (years)								
18-34	*	*	*	*	3,939	947,300	21.2	(15.6 - 26.9)
35-44	*	*	*	*	4,987	347,900	12.1	(8.0 - 16.2)
45-54	123	184,800	16.4	(7.8 - 25.1)	7,825	382,200	13.5	(9.9 - 17.1)
55-64	160	114,400	13.5	(6.1 - 20.9)	8,766	236,800	11.1	(8.0 - 14.2)
≥ 65	*	*	*	*	12,626	120,600	4.8	(2.5 - 7.0)
Race/Ethnicity								
White Non-Hispanic	393	321,300	13.2	(7.5 - 19.0)	35,833	984,600	10.6	(8.9 - 12.2)
Black Non-Hispanic	93	169,800	11.6	(3.3 - 19.8)	738	280,000	13.1	(7.2 - 19.0)
Hispanic	*	*	*	*	859	597,900	26.5	(17.3 - 35.7)
Other Non-Hispanic	*	*	*	*	713	172,300	15.3	(5.7 - 24.9)
Education								
≤ High School	231	519,000	19.5	(11.3 - 27.7)	15,696	996,600	17.8	(13.8 - 21.9)
Some College	*	*	*	*	9,911	512,400	14.5	(10.4 - 18.6)
≥ College Degree	294	247,600	10.5	(5.2 - 15.8)	12,453	466,900	8.3	(6.0 - 10.6)
Income								
≤ \$24,999	*	*	*	*	9,639	854,500	24.3	(18.4 - 30.3)
\$25,000-\$49,999	*	*	*	*	9,656	532,900	18.4	(12.5 - 24.3)
\$50,000-\$74,999	*	*	*	*	5,776	202,700	10.3	(6.0 - 14.5)
≥ \$75,000	189	78,800	5.1	(1.7 - 8.5)	8,561	192,000	4.3	(2.9 - 5.6)

¹ Defined as needing to see a doctor within the past year but not being able to because of cost

² Sample size from survey

³ Estimated number of adults who did not see a doctor because of cost within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 11b. Cost prevented visit to doctor within the past year among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	646	16.2	(11.5 - 20.9)	38,143	13.8	(11.7 - 15.9)
Gender						
Male	249	15.0	(8.0 - 22.1)	14,607	12.5	(9.4 - 15.6)
Female	397	17.3	(11.1 - 23.6)	23,536	15.0	(12.1 - 17.9)
Race/Ethnicity						
White Non-Hispanic	393	13.0	(7.2 - 18.7)	35,833	10.8	(9.2 - 12.5)
Black Non-Hispanic	93	11.8	(3.3 - 20.2)	738	13.2	(7.2 - 19.3)
Hispanic	*	*	*	859	26.1	(17.2 - 35.1)
Other Non-Hispanic	*	*	*	713	14.8	(5.5 - 24.2)
Education						
≤ High School	231	18.8	(10.9 - 26.7)	15,696	17.8	(13.8 - 21.7)
Some College	*	*	*	9,911	15.3	(11.0 - 19.5)
≥ College Degree	294	10.4	(5.1 - 15.6)	12,453	8.2	(6.0 - 10.5)
Income						
≤ \$24,999	*	*	*	9,639	24.5	(18.6 - 30.4)
\$25,000-\$49,999	*	*	*	9,656	18.7	(12.9 - 24.6)
\$50,000-\$74,999	*	*	*	5,776	10.4	(6.1 - 14.7)
≥ \$75,000	189	5.2	(1.7 - 8.6)	8,561	4.4	(3.1 - 5.8)

¹ Defined as needing to see a doctor within the past year but not being able to because of cost

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 12a. Visited a doctor for routine checkup within the past year among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	644	4,725,400	74.5	(69.2 - 79.8)	38,026	10,809,300	72.9	(70.6 - 75.3)
Gender								
Male	250	1,969,100	66.4	(57.4 - 75.4)	14,592	4,738,400	66.9	(63.0 - 70.9)
Female	394	2,756,400	81.7	(76.2 - 87.3)	23,434	6,070,900	78.5	(75.9 - 81.0)
Age (years)								
18-34	*	*	*	*	3,921	2,747,400	61.7	(55.6 - 67.9)
35-44	*	*	*	*	4,973	1,883,100	65.5	(60.2 - 70.7)
45-54	123	954,100	84.5	(76.9 - 92.1)	7,808	2,174,100	76.6	(73.2 - 80.0)
55-64	158	713,000	85.7	(79.5 - 91.9)	8,751	1,728,600	81.4	(78.7 - 84.2)
≥ 65	186	907,600	90.4	(84.3 - 96.5)	12,573	2,276,000	90.0	(87.5 - 92.5)
Race/Ethnicity								
White Non-Hispanic	394	1,843,100	75.7	(69.7 - 81.7)	35,721	6,746,200	72.4	(70.6 - 74.2)
Black Non-Hispanic	*	*	*	*	736	1,688,900	78.8	(70.4 - 87.3)
Hispanic	*	*	*	*	857	1,550,900	69.3	(60.0 - 78.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	227	2,066,700	78.5	(70.3 - 86.7)	15,640	4,150,600	74.7	(70.7 - 78.7)
Some College	*	*	*	*	9,880	2,524,000	71.3	(66.1 - 76.5)
≥ College Degree	295	1,687,800	71.4	(63.7 - 79.1)	12,428	4,042,900	71.9	(68.5 - 75.3)
Income								
≤ \$24,999	*	*	*	*	9,616	2,469,200	70.4	(64.2 - 76.6)
\$25,000-\$49,999	*	*	*	*	9,631	2,068,400	71.3	(65.3 - 77.3)
\$50,000-\$74,999	*	*	*	*	5,765	1,466,000	74.8	(69.6 - 80.0)
≥ \$75,000	189	1,050,000	68.4	(59.2 - 77.5)	8,547	3,176,500	70.7	(67.3 - 74.2)

¹ Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past year

² Sample size from survey

³ Estimated number of adults who have visited a doctor for a routine checkup within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 12b. Visited a doctor for routine checkup within the past year among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	644	74.9	(69.8 - 80.1)	38,026	72.7	(70.3 - 75.0)
Gender						
Male	250	67.5	(58.8 - 76.1)	14,592	67.1	(63.3 - 70.9)
Female	394	81.5	(75.9 - 87.1)	23,434	77.7	(75.1 - 80.3)
Race/Ethnicity						
White Non-Hispanic	394	75.9	(70.0 - 81.9)	35,721	71.9	(70.1 - 73.7)
Black Non-Hispanic	*	*	*	736	79.3	(71.3 - 87.4)
Hispanic	*	*	*	857	69.2	(60.1 - 78.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	227	78.7	(70.7 - 86.7)	15,640	74.4	(70.4 - 78.4)
Some College	*	*	*	9,880	70.9	(65.9 - 76.0)
≥ College Degree	295	71.8	(64.3 - 79.4)	12,428	71.7	(68.3 - 75.0)
Income						
≤ \$24,999	*	*	*	9,616	70.2	(64.1 - 76.2)
\$25,000-\$49,999	*	*	*	9,631	71.1	(65.3 - 76.9)
\$50,000-\$74,999	*	*	*	5,765	74.5	(69.2 - 79.8)
≥ \$75,000	189	68.6	(59.5 - 77.7)	8,547	70.3	(66.9 - 73.8)

¹ Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past year

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 13a. Visited a doctor for routine checkup within the past 2 years among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	644	5,525,800	87.2	(82.7 - 91.7)	38,026	12,726,600	85.9	(83.9 - 87.9)
Gender								
Male	250	2,429,700	81.9	(73.5 - 90.3)	14,592	5,791,900	81.8	(78.2 - 85.4)
Female	394	3,096,100	91.8	(88.2 - 95.4)	23,434	6,934,700	89.6	(87.9 - 91.3)
Age (years)								
18-34	*	*	*	*	3,921	3,535,400	79.4	(73.9 - 85.0)
35-44	102	1,119,000	84.8	(77.0 - 92.6)	4,973	2,355,600	81.9	(78.0 - 85.7)
45-54	123	1,029,600	91.2	(84.6 - 97.8)	7,808	2,507,900	88.4	(85.6 - 91.2)
55-64	158	773,500	93.0	(88.2 - 97.7)	8,751	1,924,800	90.7	(88.6 - 92.7)
≥ 65	186	952,900	94.9	(91.2 - 98.6)	12,573	2,402,900	95.0	(93.5 - 96.5)
Race/Ethnicity								
White Non-Hispanic	394	2,208,600	90.7	(86.9 - 94.5)	35,721	8,070,000	86.6	(85.4 - 87.9)
Black Non-Hispanic	*	*	*	*	736	1,914,500	89.4	(82.3 - 96.5)
Hispanic	*	*	*	*	857	1,832,800	81.9	(73.7 - 90.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	227	2,285,200	86.8	(79.5 - 94.1)	15,640	4,713,700	84.8	(81.2 - 88.4)
Some College	*	*	*	*	9,880	2,990,500	84.5	(79.6 - 89.4)
≥ College Degree	295	2,138,600	90.5	(85.3 - 95.6)	12,428	4,928,300	87.6	(85.3 - 89.9)
Income								
≤ \$24,999	168	1,663,100	84.1	(74.4 - 93.9)	9,616	2,911,700	83.0	(77.4 - 88.6)
\$25,000-\$49,999	*	*	*	*	9,631	2,375,500	81.9	(76.1 - 87.7)
\$50,000-\$74,999	76	644,300	92.4	(83.6 - 100)	5,765	1,733,500	88.4	(84.9 - 91.9)
≥ \$75,000	189	1,377,500	89.7	(84.2 - 95.1)	8,547	3,907,000	87.0	(84.8 - 89.2)

¹ Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past 2 years

² Sample size from survey

³ Estimated number of adults who have visited a doctor for a routine checkup within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 13b. Visited a doctor for routine checkup within the past 2 years among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	644	87.4	(83.1 - 91.7)	38,026	85.7	(83.7 - 87.6)
Gender						
Male	250	82.6	(74.6 - 90.5)	14,592	81.9	(78.4 - 85.3)
Female	394	91.6	(87.9 - 95.3)	23,434	89.1	(87.4 - 90.9)
Race/Ethnicity						
White Non-Hispanic	394	90.7	(86.9 - 94.5)	35,721	86.2	(85.0 - 87.5)
Black Non-Hispanic	93	90.2	(80.7 - 99.6)	736	89.6	(83.0 - 96.3)
Hispanic	*	*	*	857	81.8	(73.8 - 89.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	227	86.9	(79.8 - 93.9)	15,640	84.6	(81.1 - 88.1)
Some College	*	*	*	9,880	84.4	(79.8 - 89.0)
≥ College Degree	295	90.5	(85.4 - 95.5)	12,428	87.3	(85.0 - 89.6)
Income						
≤ \$24,999	168	84.7	(75.3 - 94.1)	9,616	83.2	(77.8 - 88.5)
\$25,000-\$49,999	*	*	*	9,631	81.9	(76.3 - 87.4)
\$50,000-\$74,999	76	92.1	(83.0 - 100)	5,765	88.1	(84.4 - 91.8)
≥ \$75,000	189	89.5	(83.9 - 95.1)	8,547	86.4	(84.2 - 88.7)

¹ Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past 2 years

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 14a. Dentist visit within the past year among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	647	4,352,900	68.6	(63.1 - 74.1)	38,026	10,540,400	71.1	(68.6 - 73.6)
Gender								
Male	251	1,969,700	66.1	(57.3 - 74.8)	14,573	4,845,800	68.4	(64.5 - 72.2)
Female	396	2,383,300	70.8	(63.9 - 77.8)	23,453	5,694,600	73.6	(70.4 - 76.8)
Age (years)								
18-34	*	*	*	*	3,924	2,789,700	62.6	(56.3 - 69.0)
35-44	*	*	*	*	4,979	2,181,000	75.8	(70.9 - 80.7)
45-54	123	859,300	76.3	(67.2 - 85.4)	7,814	2,149,300	75.8	(71.9 - 79.7)
55-64	160	673,300	79.5	(71.5 - 87.4)	8,742	1,690,500	79.2	(75.9 - 82.6)
≥ 65	187	629,300	63.0	(53.8 - 72.2)	12,567	1,729,800	68.5	(64.7 - 72.3)
Race/Ethnicity								
White Non-Hispanic	395	1,815,800	74.4	(67.7 - 81.2)	35,721	6,959,900	74.7	(72.7 - 76.6)
Black Non-Hispanic	*	*	*	*	735	1,405,800	66.2	(57.9 - 74.5)
Hispanic	*	*	*	*	862	1,425,600	63.1	(53.7 - 72.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	230	1,784,700	67.6	(58.6 - 76.6)	15,613	3,620,700	65.1	(60.7 - 69.5)
Some College	*	*	*	*	9,880	2,504,700	70.8	(66.1 - 75.5)
≥ College Degree	295	1,700,800	71.9	(63.9 - 80.0)	12,453	4,402,400	78.2	(74.6 - 81.8)
Income								
≤ \$24,999	*	*	*	*	9,594	1,972,200	56.2	(49.8 - 62.6)
\$25,000-\$49,999	*	*	*	*	9,649	1,896,000	65.8	(59.5 - 72.1)
\$50,000-\$74,999	*	*	*	*	5,763	1,482,400	75.2	(69.4 - 81.0)
≥ \$75,000	189	1,262,000	82.2	(74.5 - 89.8)	8,564	3,707,900	82.4	(79.5 - 85.3)

¹ Defined as having visited a dentist or a dental clinic for any reason within the past year

² Sample size from survey

³ Estimated number of adults who have visited a dentist or a dental clinic for any reason within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 14b. Dentist visit within the past year among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	647	68.9	(63.4 - 74.3)	38,026	71.1	(68.6 - 73.5)
Gender						
Male	251	66.4	(58.0 - 74.8)	14,573	68.3	(64.6 - 72.1)
Female	396	71.0	(64.0 - 78.1)	23,453	73.6	(70.4 - 76.8)
Race/Ethnicity						
White Non-Hispanic	395	74.3	(67.6 - 81.0)	35,721	74.4	(72.4 - 76.3)
Black Non-Hispanic	*	*	*	735	66.6	(58.3 - 74.9)
Hispanic	*	*	*	862	63.6	(54.4 - 72.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	230	68.0	(59.2 - 76.8)	15,613	65.3	(60.9 - 69.6)
Some College	*	*	*	9,880	70.4	(65.7 - 75.1)
≥ College Degree	295	72.2	(64.3 - 80.1)	12,453	78.1	(74.6 - 81.7)
Income						
≤ \$24,999	*	*	*	9,594	56.4	(50.1 - 62.8)
\$25,000-\$49,999	*	*	*	9,649	65.8	(59.6 - 72.0)
\$50,000-\$74,999	*	*	*	5,763	75.0	(69.1 - 80.8)
≥ \$75,000	189	82.0	(74.4 - 89.7)	8,564	82.1	(79.1 - 85.0)

¹ Defined as having visited a dentist or a dental clinic for any reason within the past year

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 15a. Had teeth cleaned within the past year among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	594	4,092,300	69.3	(63.5 - 75.2)	34,070	9,985,900	71.7	(69.1 - 74.3)
Gender								
Male	233	1,889,200	67.8	(58.5 - 77.0)	13,173	4,616,400	69.0	(65.0 - 73.1)
Female	361	2,203,200	70.7	(63.3 - 78.2)	20,897	5,369,500	74.2	(70.8 - 77.5)
Age (years)								
18-34	*	*	*	*	3,860	2,746,400	63.0	(56.5 - 69.5)
35-44	*	*	*	*	4,875	2,112,100	75.5	(70.7 - 80.3)
45-54	119	796,000	72.1	(62.3 - 81.9)	7,438	2,022,600	73.3	(69.1 - 77.4)
55-64	155	614,000	75.6	(66.5 - 84.8)	7,906	1,560,900	77.7	(73.8 - 81.6)
≥ 65	148	539,100	74.6	(65.7 - 83.5)	9,991	1,543,900	77.1	(73.7 - 80.5)
Race/Ethnicity								
White Non-Hispanic	364	1,723,400	75.9	(68.8 - 83.0)	31,982	6,601,500	75.4	(73.4 - 77.4)
Black Non-Hispanic	*	*	*	*	680	1,227,200	61.6	(52.1 - 71.0)
Hispanic	*	*	*	*	789	1,421,300	66.6	(57.1 - 76.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	198	1,702,800	71.3	(62.0 - 80.7)	12,740	3,388,600	67.7	(63.1 - 72.3)
Some College	*	*	*	*	9,145	2,286,800	68.5	(62.9 - 74.1)
≥ College Degree	285	1,656,500	72.7	(64.8 - 80.5)	12,128	4,298,500	78.3	(74.8 - 81.7)
Income								
≤ \$24,999	*	*	*	*	7,525	1,861,800	60.3	(53.5 - 67.1)
\$25,000-\$49,999	*	*	*	*	8,709	1,681,700	62.6	(55.4 - 69.8)
\$50,000-\$74,999	*	*	*	*	5,534	1,447,800	76.3	(70.7 - 81.8)
≥ \$75,000	189	1,237,100	80.5	(72.7 - 88.3)	8,448	3,617,500	80.8	(77.8 - 83.8)

¹ Defined as having had teeth cleaned by a dentist or dental hygienist within the past year

² Sample size from survey

³ Estimated number of adults who have had their teeth cleaned by a dentist or dental hygienist within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 15b. Had teeth cleaned within the past year among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	594	69.6	(63.8 - 75.4)	34,070	71.7	(69.1 - 74.2)
Gender						
Male	233	68.2	(59.4 - 77.1)	13,173	69.2	(65.3 - 73.1)
Female	361	70.8	(63.3 - 78.3)	20,897	73.9	(70.5 - 77.3)
Race/Ethnicity						
White Non-Hispanic	364	76.1	(69.0 - 83.1)	31,982	75.1	(73.0 - 77.1)
Black Non-Hispanic	*	*	*	680	62.0	(52.7 - 71.4)
Hispanic	*	*	*	789	66.9	(57.5 - 76.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	198	71.6	(62.4 - 80.7)	12,740	67.7	(63.2 - 72.3)
Some College	*	*	*	9,145	68.1	(62.6 - 73.6)
≥ College Degree	285	73.1	(65.4 - 80.8)	12,128	78.3	(74.9 - 81.7)
Income						
≤ \$24,999	*	*	*	7,525	60.2	(53.4 - 67.0)
\$25,000-\$49,999	*	*	*	8,709	62.7	(55.7 - 69.7)
\$50,000-\$74,999	*	*	*	5,534	76.4	(70.9 - 81.9)
≥ \$75,000	189	80.5	(72.7 - 88.2)	8,448	80.4	(77.4 - 83.4)

¹ Defined as having had teeth cleaned by a dentist or dental hygienist within the past year

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 16a. Permanent teeth extracted due to decay or gum disease among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	643	3,323,300	52.5	(46.8 - 58.2)	37,652	7,426,500	50.4	(47.8 - 52.9)
Gender								
Male	248	1,406,500	47.5	(38.6 - 56.3)	14,457	3,367,000	47.8	(43.9 - 51.6)
Female	395	1,916,800	57.0	(49.8 - 64.2)	23,195	4,059,500	52.7	(49.5 - 56.0)
Age (years)								
18-34	*	*	*	*	3,921	1,121,600	25.4	(19.7 - 31.0)
35-44	*	*	*	*	4,964	1,250,000	43.6	(37.9 - 49.2)
45-54	*	*	*	*	7,780	1,595,300	56.1	(51.4 - 60.7)
55-64	158	525,900	62.9	(53.5 - 72.2)	8,648	1,382,800	65.5	(61.6 - 69.4)
≥ 65	186	874,300	87.2	(81.7 - 92.6)	12,339	2,076,900	83.0	(80.6 - 85.4)
Race/Ethnicity								
White Non-Hispanic	391	1,076,800	44.8	(38.2 - 51.5)	35,364	4,425,700	47.9	(46.0 - 49.9)
Black Non-Hispanic	*	*	*	*	727	1,244,700	58.2	(48.9 - 67.4)
Hispanic	*	*	*	*	853	1,220,500	54.2	(44.9 - 63.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	230	1,657,600	62.4	(52.9 - 71.9)	15,427	3,436,000	61.9	(57.2 - 66.6)
Some College	*	*	*	*	9,798	1,726,200	49.3	(44.3 - 54.2)
≥ College Degree	294	1,016,600	43.1	(35.2 - 51.1)	12,351	2,242,600	40.1	(36.5 - 43.6)
Income								
≤ \$24,999	*	*	*	*	9,483	2,171,900	62.3	(55.8 - 68.8)
\$25,000-\$49,999	*	*	*	*	9,552	1,577,400	54.7	(48.3 - 61.0)
\$50,000-\$74,999	*	*	*	*	5,744	1,000,700	51.5	(45.6 - 57.4)
≥ \$75,000	189	639,800	41.6	(32.6 - 50.7)	8,516	1,738,200	38.8	(35.3 - 42.2)

¹ Defined as having had at least one permanent tooth extracted because of tooth decay or gum disease

² Sample size from survey

³ Estimated number of adults who have had at least one permanent tooth extracted because of tooth decay or gum disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 16b. Permanent teeth extracted due to decay or gum disease among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	643	53.0	(47.4 - 58.6)	37,652	49.4	(46.9 - 51.9)
Gender						
Male	248	49.2	(40.6 - 57.9)	14,457	47.9	(44.1 - 51.7)
Female	395	56.3	(49.1 - 63.6)	23,195	50.7	(47.4 - 54.0)
Race/Ethnicity						
White Non-Hispanic	391	45.5	(38.9 - 52.2)	35,364	46.5	(44.5 - 48.5)
Black Non-Hispanic	*	*	*	727	57.7	(48.6 - 66.9)
Hispanic	*	*	*	853	53.8	(44.6 - 62.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	230	63.1	(53.7 - 72.4)	15,427	61.3	(56.7 - 66.0)
Some College	*	*	*	9,798	47.9	(43.0 - 52.9)
≥ College Degree	294	43.4	(35.5 - 51.3)	12,351	39.0	(35.5 - 42.5)
Income						
≤ \$24,999	*	*	*	9,483	61.6	(55.2 - 68.1)
\$25,000-\$49,999	*	*	*	9,552	54.2	(48.0 - 60.3)
\$50,000-\$74,999	*	*	*	5,744	49.9	(44.0 - 55.9)
≥ \$75,000	189	41.6	(32.5 - 50.7)	8,516	37.7	(34.2 - 41.1)

¹ Defined as having had at least one permanent tooth extracted because of tooth decay or gum disease

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 17a. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	186	253,600	25.3	(16.6 - 34.0)	12,339	491,200	19.6	(16.0 - 23.3)
Gender								
Male	*	*	*	*	4,352	167,200	16.4	(11.8 - 20.9)
Female	*	*	*	*	7,987	324,000	21.8	(16.6 - 27.0)
Age (years)								
≥ 65	186	253,600	25.3	(16.6 - 34.0)	12,339	491,200	19.6	(16.0 - 23.3)
Race/Ethnicity								
White Non-Hispanic	137	102,700	18.8	(10.8 - 26.8)	11,885	322,600	16.8	(14.3 - 19.2)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	6,358	314,100	25.3	(19.9 - 30.7)
Some College	*	*	*	*	2,736	90,600	20.2	(12.1 - 28.2)
≥ College Degree	*	*	*	*	3,214	80,500	10.1	(3.6 - 16.5)
Income								
≤ \$24,999	*	*	*	*	4,558	199,600	26.9	(19.3 - 34.4)
\$25,000-\$49,999	*	*	*	*	3,365	123,900	21.4	(13.1 - 29.6)
\$50,000-\$74,999	*	*	*	*	1,179	21,400	7.3	(3.8 - 10.8)
≥ \$75,000	*	*	*	*	1,187	9,600	2.5	(1.0 - 4.0)

¹ Defined as having all permanent teeth extracted because of tooth decay or gum disease among adults aged 65 and older

² Sample size from survey

³ Estimated number of adults aged 65 and older who have had all permanent teeth extracted because of tooth decay or gum disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 17b. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	186	24.3	(15.8 - 32.8)	12,339	19.4	(15.7 - 23.2)
Gender						
Male	*	*	*	4,352	16.5	(11.5 - 21.4)
Female	*	*	*	7,987	22.1	(16.6 - 27.7)
Race/Ethnicity						
White Non-Hispanic	137	18.3	(10.1 - 26.5)	11,885	16.6	(13.9 - 19.3)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	6,358	24.9	(19.3 - 30.5)
Some College	*	*	*	2,736	21.0	(12.1 - 29.9)
≥ College Degree	*	*	*	3,214	10.0	(3.6 - 16.3)
Income						
≤ \$24,999	*	*	*	4,558	27.3	(19.1 - 35.5)
\$25,000-\$49,999	*	*	*	3,365	21.0	(12.9 - 29.1)
\$50,000-\$74,999	*	*	*	1,179	7.6	(3.6 - 11.5)
≥ \$75,000	*	*	*	1,187	2.2	(0.9 - 3.5)

¹ Defined as having all permanent teeth extracted because of tooth decay or gum disease among adults aged 65 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 18a. Flu shot within the past 12 months¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	646	2,662,000	42.2	(36.6 - 47.7)	38,079	6,255,900	42.3	(39.9 - 44.8)
Gender								
Male	250	1,191,700	40.2	(31.3 - 49.0)	14,579	2,798,900	39.6	(35.8 - 43.5)
Female	396	1,470,300	44.0	(37.0 - 50.9)	23,500	3,456,900	44.8	(41.6 - 47.9)
Age (years)								
18-49	226	1,254,100	32.5	(24.4 - 40.6)	12,505	2,685,300	31.1	(27.3 - 34.9)
50-64	232	658,400	45.8	(37.7 - 53.9)	12,972	1,662,900	46.0	(42.6 - 49.5)
≥ 65	188	749,600	74.0	(66.3 - 81.7)	12,602	1,907,700	75.2	(72.0 - 78.4)
Race/Ethnicity								
White Non-Hispanic	394	1,034,100	42.6	(36.0 - 49.2)	35,770	4,029,000	43.3	(41.4 - 45.3)
Black Non-Hispanic	*	*	*	*	738	907,000	42.1	(33.1 - 51.1)
Hispanic	*	*	*	*	860	882,900	39.6	(30.4 - 48.9)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	230	1,200,400	45.5	(36.3 - 54.7)	15,663	2,457,800	44.3	(39.8 - 48.8)
Some College	*	*	*	*	9,898	1,377,700	38.9	(34.0 - 43.9)
≥ College Degree	294	942,700	40.4	(32.7 - 48.2)	12,436	2,404,400	43.0	(39.5 - 46.4)
Income								
≤ \$24,999	*	*	*	*	9,635	1,656,800	47.6	(41.2 - 54.0)
\$25,000-\$49,999	*	*	*	*	9,647	1,200,600	41.4	(35.6 - 47.3)
\$50,000-\$74,999	*	*	*	*	5,763	792,300	40.2	(34.3 - 46.2)
≥ \$75,000	189	581,400	37.8	(29.3 - 46.4)	8,548	1,776,100	39.6	(36.3 - 42.8)

¹ Defined as having had a flu shot within the past 12 months

² Sample size from survey

³ Estimated number of adults who have had a flu shot within the past 12 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 18b. Flu shot within the past 12 months¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	646	42.3	(36.8 - 47.8)	38,079	41.7	(39.2 - 44.1)
Gender						
Male	250	41.4	(32.9 - 50.0)	14,579	40.0	(36.2 - 43.8)
Female	396	43.1	(36.1 - 50.0)	23,500	43.1	(40.0 - 46.3)
Race/Ethnicity						
White Non-Hispanic	394	43.0	(36.4 - 49.7)	35,770	42.4	(40.4 - 44.3)
Black Non-Hispanic	*	*	*	738	42.3	(33.3 - 51.3)
Hispanic	*	*	*	860	39.0	(29.9 - 48.1)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	230	45.3	(36.3 - 54.4)	15,663	43.7	(39.1 - 48.2)
Some College	*	*	*	9,898	37.8	(33.0 - 42.7)
≥ College Degree	294	40.9	(33.2 - 48.7)	12,436	42.5	(39.1 - 46.0)
Income						
≤ \$24,999	*	*	*	9,635	47.3	(41.0 - 53.7)
\$25,000-\$49,999	*	*	*	9,647	40.3	(34.5 - 46.0)
\$50,000-\$74,999	*	*	*	5,763	39.4	(33.4 - 45.4)
≥ \$75,000	189	38.2	(29.6 - 46.8)	8,548	39.1	(35.8 - 42.4)

¹ Defined as having had a flu shot within the past 12 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 19a. Flu vaccine that was sprayed in nose within the past 12 months¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	647	151,300	2.4	(0.0 - 4.8)	38,106	248,300	1.7	(0.6 - 2.8)
Gender								
Male	*	*	*	*	14,586	148,000	2.1	(0.0 - 4.2)
Female	*	*	*	*	23,520	100,300	1.3	(0.5 - 2.1)
Age (years)								
18-49	*	*	*	*	12,535	184,900	2.1	(0.3 - 3.9)
50-64	*	*	*	*	12,968	36,000	1.0	(0.2 - 1.8)
≥ 65	*	*	*	*	12,603	27,400	1.1	(0.4 - 1.8)
Race/Ethnicity								
White Non-Hispanic	*	*	*	*	35,796	66,000	0.7	(0.4 - 1.0)
Black Non-Hispanic	*	*	*	*	740	38,500	1.8	(0.0 - 4.1)
Hispanic	*	*	*	*	860	133,700	5.9	(0.0 - 12.3)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,679	172,500	3.1	(0.3 - 5.9)
Some College	*	*	*	*	9,897	43,500	1.2	(0.4 - 2.0)
≥ College Degree	*	*	*	*	12,449	31,600	0.6	(0.3 - 0.8)
Income								
≤ \$24,999	*	*	*	*	9,636	138,400	3.9	(0.0 - 8.1)
\$25,000-\$49,999	*	*	*	*	9,648	41,500	1.4	(0.0 - 3.1)
\$50,000-\$74,999	*	*	*	*	5,766	13,300	0.7	(0.2 - 1.2)
≥ \$75,000	*	*	*	*	8,561	35,300	0.8	(0.3 - 1.3)

¹ Defined as having had flu vaccine sprayed in nose within the past 12 months

² Sample size from survey

³ Estimated number of adults who have had flu vaccine sprayed in nose within the past 12 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 19b. Flu vaccine that was sprayed in nose within the past 12 months¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	647	2.3	(0.0 - 4.7)	38,106	1.7	(0.6 - 2.7)
Gender						
Male	*	*	*	14,586	2.0	(0.1 - 3.9)
Female	*	*	*	23,520	1.4	(0.5 - 2.3)
Race/Ethnicity						
White Non-Hispanic	*	*	*	35,796	0.7	(0.4 - 1.0)
Black Non-Hispanic	*	*	*	740	1.9	(0.0 - 4.4)
Hispanic	*	*	*	860	5.9	(0.0 - 11.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	15,679	3.1	(0.4 - 5.8)
Some College	*	*	*	9,897	1.2	(0.4 - 2.0)
≥ College Degree	*	*	*	12,449	0.6	(0.3 - 0.8)
Income						
≤ \$24,999	*	*	*	9,636	3.8	(0.0 - 7.7)
\$25,000-\$49,999	*	*	*	9,648	1.5	(0.0 - 3.4)
\$50,000-\$74,999	*	*	*	5,766	0.7	(0.1 - 1.2)
≥ \$75,000	*	*	*	8,561	0.8	(0.3 - 1.3)

¹ Defined as having had flu vaccine sprayed in nose within the past 12 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 20a. Flu shot or flu vaccine sprayed in nose within the past 12 months¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	646	2,674,200	42.4	(36.8 - 47.9)	38,059	6,302,300	42.7	(40.2 - 45.1)
Gender								
Male	250	1,191,700	40.2	(31.3 - 49.0)	14,568	2,822,700	40.0	(36.1 - 43.9)
Female	396	1,482,500	44.3	(37.4 - 51.3)	23,491	3,479,500	45.1	(42.0 - 48.2)
Age (years)								
18-49	226	1,266,200	32.8	(24.7 - 40.9)	12,501	2,726,400	31.6	(27.8 - 35.4)
50-64	232	658,400	45.8	(37.7 - 53.9)	12,964	1,664,200	46.1	(42.7 - 49.5)
≥ 65	188	749,600	74.0	(66.3 - 81.7)	12,594	1,911,700	75.4	(72.2 - 78.6)
Race/Ethnicity								
White Non-Hispanic	394	1,034,100	42.6	(36.0 - 49.2)	35,750	4,048,700	43.6	(41.7 - 45.5)
Black Non-Hispanic	*	*	*	*	738	910,000	42.2	(33.2 - 51.2)
Hispanic	*	*	*	*	860	903,300	40.6	(31.3 - 49.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	230	1,212,500	46.0	(36.8 - 55.1)	15,651	2,480,100	44.7	(40.2 - 49.3)
Some College	*	*	*	*	9,895	1,387,300	39.2	(34.3 - 44.1)
≥ College Degree	294	942,700	40.4	(32.7 - 48.2)	12,431	2,418,300	43.2	(39.8 - 46.7)
Income								
≤ \$24,999	*	*	*	*	9,626	1,675,100	48.2	(41.8 - 54.6)
\$25,000-\$49,999	*	*	*	*	9,643	1,208,300	41.7	(35.8 - 47.6)
\$50,000-\$74,999	*	*	*	*	5,761	795,600	40.4	(34.5 - 46.3)
≥ \$75,000	189	581,400	37.8	(29.3 - 46.4)	8,548	1,791,200	39.9	(36.6 - 43.2)

¹ Defined as having had a flu shot or flu vaccine sprayed in nose within the past 12 months

² Sample size from survey

³ Estimated number of adults who have had a flu shot or flu vaccine sprayed in nose within the past 12 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 20b. Flu shot or flu vaccine sprayed in nose within the past 12 months¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	646	42.5	(37.0 - 48.0)	38,059	42.0	(39.6 - 44.5)
Gender						
Male	250	41.4	(32.9 - 50.0)	14,568	40.4	(36.6 - 44.2)
Female	396	43.5	(36.5 - 50.5)	23,491	43.5	(40.3 - 46.7)
Race/Ethnicity						
White Non-Hispanic	394	43.0	(36.4 - 49.7)	35,750	42.6	(40.7 - 44.6)
Black Non-Hispanic	*	*	*	738	42.4	(33.4 - 51.4)
Hispanic	*	*	*	860	40.0	(30.9 - 49.1)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	230	45.8	(36.8 - 54.9)	15,651	44.1	(39.6 - 48.6)
Some College	*	*	*	9,895	38.1	(33.2 - 43.0)
≥ College Degree	294	40.9	(33.2 - 48.7)	12,431	42.8	(39.4 - 46.3)
Income						
≤ \$24,999	*	*	*	9,626	48.0	(41.6 - 54.3)
\$25,000-\$49,999	*	*	*	9,643	40.5	(34.8 - 46.3)
\$50,000-\$74,999	*	*	*	5,761	39.6	(33.6 - 45.6)
≥ \$75,000	189	38.2	(29.6 - 46.8)	8,548	39.4	(36.1 - 42.7)

¹ Defined as having had a flu shot or flu vaccine sprayed in nose within the past 12 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 21a. Flu shot among adults aged 65 and older¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	188	749,600	74.0	(66.3 - 81.7)	12,602	1,907,700	75.2	(72.0 - 78.4)
Gender								
Male	*	*	*	*	4,417	772,900	74.6	(69.3 - 79.8)
Female	129	460,600	74.9	(65.5 - 84.4)	8,185	1,134,800	75.6	(71.6 - 79.6)
Age (years)								
≥ 65	188	749,600	74.0	(66.3 - 81.7)	12,602	1,907,700	75.2	(72.0 - 78.4)
Race/Ethnicity								
White Non-Hispanic	139	416,000	74.7	(66.5 - 82.9)	12,137	1,487,100	76.0	(73.5 - 78.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	6,518	930,800	74.0	(68.9 - 79.2)
Some College	*	*	*	*	2,790	317,800	69.7	(62.7 - 76.7)
≥ College Degree	72	266,300	83.5	(74.2 - 92.8)	3,261	654,000	80.5	(76.4 - 84.6)
Income								
≤ \$24,999	*	*	*	*	4,665	545,300	72.4	(65.6 - 79.3)
\$25,000-\$49,999	*	*	*	*	3,422	430,600	72.8	(67.0 - 78.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	1,198	301,000	77.5	(70.9 - 84.2)

¹ Defined as having had a flu shot during the past 12 months among adults aged 65 and older

² Sample size from survey

³ Estimated number of adults aged 65 and older who have had a flu shot within the past 12 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 21b. Flu shot among adults aged 65 and older¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	188	73.8	(65.8 - 81.7)	12,602	75.0	(71.5 - 78.5)
Gender						
Male	*	*	*	4,417	74.4	(68.7 - 80.1)
Female	129	74.9	(65.4 - 84.3)	8,185	75.5	(71.3 - 79.8)
Race/Ethnicity						
White Non-Hispanic	139	74.5	(65.9 - 83.1)	12,137	75.9	(73.0 - 78.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	6,518	73.3	(67.5 - 79.1)
Some College	*	*	*	2,790	70.0	(62.7 - 77.2)
≥ College Degree	72	83.5	(74.1 - 93.0)	3,261	80.7	(76.2 - 85.1)
Income						
≤ \$24,999	*	*	*	4,665	72.6	(65.1 - 80.1)
\$25,000-\$49,999	*	*	*	3,422	71.4	(65.0 - 77.9)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	1,198	78.0	(70.8 - 85.1)

¹ Defined as having had a flu shot during the past 12 months among adults aged 65 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 22a. Pneumonia shot or pneumococcal vaccine¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	589	1,427,200	24.9	(20.0 - 29.9)	35,303	3,456,800	26.1	(23.9 - 28.3)
Gender								
Male	221	624,400	24.1	(15.8 - 32.4)	13,052	1,542,200	25.2	(21.5 - 28.8)
Female	368	802,800	25.6	(19.7 - 31.5)	22,251	1,914,600	26.9	(24.2 - 29.6)
Age (years)								
18-49	199	542,800	15.7	(8.7 - 22.6)	10,943	1,097,000	14.6	(11.3 - 17.9)
50-64	211	328,500	25.4	(17.6 - 33.2)	12,179	758,000	23.0	(19.8 - 26.2)
≥ 65	179	555,900	56.9	(47.5 - 66.3)	12,181	1,601,700	65.7	(61.7 - 69.7)
Race/Ethnicity								
White Non-Hispanic	357	604,900	28.4	(22.2 - 34.7)	33,225	2,358,700	28.7	(26.9 - 30.5)
Black Non-Hispanic	*	*	*	*	685	584,400	28.4	(19.5 - 37.3)
Hispanic	108	315,400	20.2	(11.2 - 29.2)	769	408,000	19.5	(12.6 - 26.4)
Other Non-Hispanic	*	*	*	*	624	105,600	12.2	(2.3 - 22.2)
Education								
≤ High School	215	678,900	28.1	(20.2 - 36.1)	14,700	1,535,300	30.4	(26.4 - 34.3)
Some College	*	*	*	*	9,151	836,200	27.2	(21.9 - 32.5)
≥ College Degree	265	404,500	18.9	(12.9 - 24.9)	11,375	1,070,700	21.4	(18.6 - 24.1)
Income								
≤ \$24,999	155	519,100	29.0	(19.9 - 38.2)	9,082	1,050,900	33.1	(27.6 - 38.6)
\$25,000-\$49,999	*	*	*	*	9,033	797,500	30.3	(24.1 - 36.6)
\$50,000-\$74,999	*	*	*	*	5,320	378,800	21.8	(17.4 - 26.1)
≥ \$75,000	171	176,400	12.8	(7.7 - 17.9)	7,773	648,700	16.3	(14.1 - 18.5)

¹ Defined as having ever had a pneumonia shot or pneumococcal vaccine

² Sample size from survey

³ Estimated number of adults who have ever had a pneumonia shot or pneumococcal vaccine (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 22b. Pneumonia shot or pneumococcal vaccine¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	589	24.9	(20.0 - 29.7)	35,303	25.4	(23.2 - 27.6)
Gender						
Male	221	24.9	(17.0 - 32.9)	13,052	25.8	(22.3 - 29.4)
Female	368	24.8	(19.0 - 30.7)	22,251	25.1	(22.4 - 27.8)
Race/Ethnicity						
White Non-Hispanic	357	28.3	(22.1 - 34.5)	33,225	27.7	(25.9 - 29.5)
Black Non-Hispanic	*	*	*	685	27.9	(19.1 - 36.7)
Hispanic	108	20.5	(11.6 - 29.5)	769	19.6	(12.8 - 26.5)
Other Non-Hispanic	*	*	*	624	11.3	(2.0 - 20.7)
Education						
≤ High School	215	28.0	(20.2 - 35.8)	14,700	29.5	(25.6 - 33.5)
Some College	*	*	*	9,151	26.2	(21.0 - 31.3)
≥ College Degree	265	19.3	(13.4 - 25.3)	11,375	21.0	(18.3 - 23.8)
Income						
≤ \$24,999	155	29.2	(20.1 - 38.3)	9,082	32.3	(26.9 - 37.8)
\$25,000-\$49,999	*	*	*	9,033	29.0	(23.0 - 35.0)
\$50,000-\$74,999	*	*	*	5,320	21.2	(16.8 - 25.7)
≥ \$75,000	171	13.2	(7.8 - 18.7)	7,773	16.2	(13.9 - 18.5)

¹ Defined as having ever had a pneumonia shot or pneumococcal vaccine

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 23a. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	179	555,900	56.9	(47.5 - 66.3)	12,181	1,601,700	65.7	(61.7 - 69.7)
Gender								
Male	*	*	*	*	4,180	606,400	62.1	(55.5 - 68.7)
Female	*	*	*	*	8,001	995,300	68.0	(63.1 - 73.0)
Age (years)								
≥ 65	179	555,900	56.9	(47.5 - 66.3)	12,181	1,601,700	65.7	(61.7 - 69.7)
Race/Ethnicity								
White Non-Hispanic	*	*	*	*	11,735	1,281,300	68.2	(64.9 - 71.5)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	6,345	806,200	65.6	(59.5 - 71.7)
Some College	*	*	*	*	2,694	289,500	66.8	(60.1 - 73.4)
≥ College Degree	*	*	*	*	3,111	501,100	65.6	(58.5 - 72.7)
Income								
≤ \$24,999	*	*	*	*	4,533	497,100	67.7	(59.9 - 75.5)
\$25,000-\$49,999	*	*	*	*	3,324	344,300	61.5	(53.3 - 69.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	1,138	236,300	64.5	(55.2 - 73.8)

¹ Defined as having ever had a pneumonia shot or pneumococcal vaccine among adults aged 65 and older

² Sample size from survey

³ Estimated number of adults aged 65 and older who have ever had a pneumonia shot or pneumococcal vaccine (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 23b. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	179	56.2	(46.5 - 65.8)	12,181	64.7	(60.4 - 69.1)
Gender						
Male	*	*	*	4,180	61.5	(54.3 - 68.6)
Female	*	*	*	8,001	67.6	(62.4 - 72.9)
Race/Ethnicity						
White Non-Hispanic	*	*	*	11,735	67.1	(63.2 - 70.9)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	6,345	64.2	(57.4 - 71.0)
Some College	*	*	*	2,694	65.8	(58.4 - 73.3)
≥ College Degree	*	*	*	3,111	65.3	(57.7 - 72.9)
Income						
≤ \$24,999	*	*	*	4,533	66.0	(57.0 - 74.9)
\$25,000-\$49,999	*	*	*	3,324	60.3	(51.9 - 68.6)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	*	*

¹ Defined as having ever had a pneumonia shot or pneumococcal vaccine among adults aged 65 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 24a. Ever had a mammogram among women aged 40 and older¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	306	1,674,600	88.5	(83.2 - 93.7)	19,491	4,328,800	91.8	(89.7 - 94.0)
Gender								
Male	-	-	-	-	-	-	-	-
Female	306	1,674,600	88.5	(83.2 - 93.7)	19,491	4,328,800	91.8	(89.7 - 94.0)
Age (years)								
40-49	*	*	*	*	3,851	1,287,500	85.0	(79.7 - 90.3)
50-64	115	634,800	96.6	(92.0 - 100)	7,452	1,641,700	96.0	(94.1 - 97.9)
≥ 65	127	550,000	91.3	(82.8 - 99.9)	8,188	1,399,600	94.1	(90.5 - 97.6)
Race/Ethnicity								
White Non-Hispanic	192	694,100	92.5	(88.1 - 96.8)	18,548	2,976,100	93.9	(92.7 - 95.1)
Black Non-Hispanic	*	*	*	*	342	746,600	93.8	(87.3 - 100)
Hispanic	*	*	*	*	333	447,600	91.8	(84.6 - 99.0)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	123	810,000	89.1	(81.9 - 96.3)	8,266	1,698,700	91.0	(87.4 - 94.5)
Some College	*	*	*	*	5,322	1,055,500	93.6	(90.2 - 97.0)
≥ College Degree	*	*	*	*	5,872	1,560,700	91.6	(87.5 - 95.6)
Income								
≤ \$24,999	83	538,600	93.9	(86.7 - 100)	5,691	1,023,400	93.5	(89.7 - 97.4)
\$25,000-\$49,999	*	*	*	*	4,857	860,300	89.5	(83.6 - 95.5)
\$50,000-\$74,999	*	*	*	*	2,648	547,100	94.6	(92.2 - 96.9)
≥ \$75,000	*	*	*	*	3,582	1,185,100	92.2	(88.2 - 96.1)

¹ Defined as having ever had a mammogram among women aged 40 and older

² Sample size from survey

³ Estimated number of women aged 40 and older who have ever had a mammogram (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 24b. Ever had a mammogram among women aged 40 and older¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	306	87.6	(81.9 - 93.2)	19,491	91.1	(88.7 - 93.6)
Gender						
Male	-	-	-	-	-	-
Female	306	87.6	(81.9 - 93.2)	19,491	91.1	(88.7 - 93.6)
Race/Ethnicity						
White Non-Hispanic	192	91.9	(87.2 - 96.6)	18,548	93.3	(92.0 - 94.7)
Black Non-Hispanic	*	*	*	342	93.1	(85.7 - 100)
Hispanic	*	*	*	333	91.0	(82.8 - 99.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	123	88.1	(80.2 - 96.0)	8,266	90.2	(86.1 - 94.2)
Some College	*	*	*	5,322	92.9	(88.9 - 96.9)
≥ College Degree	*	*	*	5,872	90.9	(86.6 - 95.3)
Income						
≤ \$24,999	83	93.2	(85.1 - 100)	5,691	93.0	(88.5 - 97.4)
\$25,000-\$49,999	*	*	*	4,857	88.8	(82.5 - 95.1)
\$50,000-\$74,999	*	*	*	2,648	94.3	(91.7 - 96.8)
≥ \$75,000	*	*	*	3,582	91.2	(86.8 - 95.6)

¹ Defined as having ever had a mammogram among women aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 25a. Had mammogram within the past 2 years among women aged 40 and older¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	304	1,454,700	77.6	(71.4 - 83.7)	19,376	3,760,000	80.3	(77.8 - 82.9)
Gender								
Male	-	-	-	-	-	-	-	-
Female	304	1,454,700	77.6	(71.4 - 83.7)	19,376	3,760,000	80.3	(77.8 - 82.9)
Age (years)								
40-49	*	*	*	*	3,842	1,122,700	74.4	(68.5 - 80.2)
50-64	115	592,000	90.1	(83.8 - 96.4)	7,428	1,487,900	87.2	(84.5 - 89.8)
≥ 65	*	*	*	*	8,106	1,149,400	78.5	(74.2 - 82.9)
Race/Ethnicity								
White Non-Hispanic	191	549,600	73.6	(66.2 - 80.9)	18,438	2,531,500	80.2	(78.3 - 82.2)
Black Non-Hispanic	*	*	*	*	339	667,100	85.6	(77.5 - 93.8)
Hispanic	*	*	*	*	332	419,700	86.1	(77.5 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	122	736,700	82.3	(74.2 - 90.5)	8,204	1,486,900	80.5	(76.4 - 84.6)
Some College	*	*	*	*	5,297	929,400	83.0	(78.8 - 87.1)
≥ College Degree	*	*	*	*	5,845	1,334,800	78.6	(73.9 - 83.2)
Income								
≤ \$24,999	83	495,300	86.4	(77.7 - 95.1)	5,650	882,300	80.8	(75.9 - 85.7)
\$25,000-\$49,999	*	*	*	*	4,846	732,300	76.3	(69.7 - 82.9)
\$50,000-\$74,999	*	*	*	*	2,641	494,200	85.6	(81.9 - 89.4)
≥ \$75,000	*	*	*	*	3,572	1,025,900	80.0	(75.2 - 84.8)

¹ Defined as having had a mammogram within the past 2 years among women aged 40 and older

² Sample size from survey

³ Estimated number of women aged 40 and older who have had a mammogram within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 25b. Had mammogram within the past 2 years among women aged 40 and older¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	304	76.7	(70.2 - 83.2)	19,376	79.7	(76.9 - 82.5)
Gender						
Male	-	-	-	-	-	-
Female	304	76.7	(70.2 - 83.2)	19,376	79.7	(76.9 - 82.5)
Race/Ethnicity						
White Non-Hispanic	191	72.6	(65.0 - 80.3)	18,438	79.7	(77.6 - 81.8)
Black Non-Hispanic	*	*	*	339	85.3	(76.5 - 94.2)
Hispanic	*	*	*	332	85.2	(75.6 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	122	81.6	(73.0 - 90.3)	8,204	80.1	(75.6 - 84.7)
Some College	*	*	*	5,297	82.3	(77.7 - 86.9)
≥ College Degree	*	*	*	5,845	77.8	(72.9 - 82.8)
Income						
≤ \$24,999	83	86.0	(76.7 - 95.2)	5,650	80.8	(75.4 - 86.2)
\$25,000-\$49,999	*	*	*	4,846	75.7	(68.7 - 82.6)
\$50,000-\$74,999	*	*	*	2,641	85.5	(81.6 - 89.4)
≥ \$75,000	*	*	*	3,572	79.0	(73.8 - 84.2)

¹ Defined as having had a mammogram within the past 2 years among women aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 26a. Had mammogram within the past 2 years among women aged 50 and older¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	240	1,026,300	82.7	(76.5 - 88.8)	15,534	2,637,300	83.2	(80.7 - 85.7)
Gender								
Male	-	-	-	-	-	-	-	-
Female	240	1,026,300	82.7	(76.5 - 88.8)	15,534	2,637,300	83.2	(80.7 - 85.7)
Age (years)								
50-64	115	592,000	90.1	(83.8 - 96.4)	7,428	1,487,900	87.2	(84.5 - 89.8)
≥ 65	*	*	*	*	8,106	1,149,400	78.5	(74.2 - 82.9)
Race/Ethnicity								
White Non-Hispanic	163	467,200	78.4	(71.2 - 85.6)	14,887	1,907,000	82.3	(80.3 - 84.4)
Black Non-Hispanic	*	*	*	*	249	385,600	89.9	(83.2 - 96.7)
Hispanic	*	*	*	*	220	286,700	93.9	(89.1 - 98.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	97	512,800	88.1	(81.4 - 94.7)	7,026	1,094,500	83.3	(80.0 - 86.6)
Some College	*	*	*	*	4,124	646,700	87.3	(84.2 - 90.3)
≥ College Degree	*	*	*	*	4,356	890,500	80.6	(75.1 - 86.1)
Income								
≤ \$24,999	68	339,900	87.9	(80.3 - 95.4)	4,991	649,100	81.0	(76.8 - 85.2)
\$25,000-\$49,999	*	*	*	*	3,913	533,300	78.6	(70.9 - 86.2)
\$50,000-\$74,999	*	*	*	*	1,943	359,300	89.6	(86.3 - 93.0)
≥ \$75,000	*	*	*	*	2,303	601,300	84.6	(79.3 - 89.9)

¹ Defined as having had a mammogram within the past 2 years among women aged 50 and older

² Sample size from survey

³ Estimated number of women aged 50 and older who have had a mammogram within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 26b. Had mammogram within the past 2 years among women aged 50 and older¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	240	82.4	(76.1 - 88.7)	15,534	83.1	(80.3 - 85.8)
Gender						
Male	-	-	-	-	-	-
Female	240	82.4	(76.1 - 88.7)	15,534	83.1	(80.3 - 85.8)
Race/Ethnicity						
White Non-Hispanic	163	78.2	(70.9 - 85.5)	14,887	82.1	(79.9 - 84.3)
Black Non-Hispanic	*	*	*	249	90.4	(83.6 - 97.2)
Hispanic	*	*	*	220	94.1	(89.3 - 99.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	97	88.2	(81.6 - 94.8)	7,026	83.6	(80.1 - 87.0)
Some College	*	*	*	4,124	87.3	(84.1 - 90.5)
≥ College Degree	*	*	*	4,356	79.9	(73.9 - 86.0)
Income						
≤ \$24,999	68	87.8	(80.3 - 95.4)	4,991	81.3	(76.9 - 85.6)
\$25,000-\$49,999	*	*	*	3,913	78.1	(69.9 - 86.4)
\$50,000-\$74,999	*	*	*	1,943	90.2	(87.1 - 93.4)
≥ \$75,000	*	*	*	2,303	83.9	(77.9 - 89.9)

¹ Defined as having had a mammogram within the past 2 years among women aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 27a. Ever had a Pap test among women¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	390	2,943,100	89.3	(84.7 - 94.0)	23,381	7,045,300	92.1	(90.0 - 94.3)
Gender								
Male	-	-	-	-	-	-	-	-
Female	390	2,943,100	89.3	(84.7 - 94.0)	23,381	7,045,300	92.1	(90.0 - 94.3)
Age (years)								
18-34	*	*	*	*	2,377	1,896,600	85.2	(79.3 - 91.0)
35-44	*	*	*	*	3,023	1,301,500	92.6	(87.7 - 97.5)
45-54	68	573,400	98.1	(95.1 - 100)	4,657	1,420,300	98.3	(97.0 - 99.6)
55-64	88	421,700	96.0	(91.0 - 100)	5,187	1,071,200	97.2	(95.0 - 99.3)
≥ 65	126	517,500	88.0	(78.8 - 97.2)	8,137	1,355,700	92.3	(88.5 - 96.1)
Race/Ethnicity								
White Non-Hispanic	230	1,055,500	87.7	(80.0 - 95.5)	21,976	4,439,700	94.2	(92.1 - 96.3)
Black Non-Hispanic	65	890,200	89.8	(80.1 - 99.4)	481	1,231,600	91.2	(84.0 - 98.4)
Hispanic	69	736,200	95.4	(91.2 - 99.6)	533	984,500	90.1	(85.3 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	151	1,254,600	86.0	(77.7 - 94.4)	9,526	2,540,500	88.3	(83.8 - 92.7)
Some College	74	611,200	91.2	(83.3 - 99.1)	6,462	1,782,200	93.5	(90.4 - 96.6)
≥ College Degree	162	1,008,700	91.9	(85.1 - 98.8)	7,345	2,643,200	95.0	(92.2 - 97.9)
Income								
≤ \$24,999	105	961,400	95.4	(91.0 - 99.8)	6,526	1,717,200	93.7	(90.9 - 96.6)
\$25,000-\$49,999	*	*	*	*	5,787	1,378,400	91.0	(85.6 - 96.4)
\$50,000-\$74,999	*	*	*	*	3,310	933,900	96.0	(91.8 - 100)
≥ \$75,000	93	619,800	91.7	(83.9 - 99.5)	4,570	1,943,300	95.4	(92.6 - 98.2)

¹ Defined as having ever had a Pap test among women

² Sample size from survey

³ Estimated number of women who have ever had a Pap test (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 27b. Ever had a Pap test among women¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	390	89.2	(84.5 - 94.0)	23,381	91.8	(89.6 - 94.0)
Gender						
Male	-	-	-	-	-	-
Female	390	89.2	(84.5 - 94.0)	23,381	91.8	(89.6 - 94.0)
Race/Ethnicity						
White Non-Hispanic	230	87.4	(79.4 - 95.4)	21,976	94.0	(91.8 - 96.2)
Black Non-Hispanic	65	89.6	(79.9 - 99.3)	481	90.8	(83.5 - 98.1)
Hispanic	69	95.6	(91.5 - 99.7)	533	89.7	(84.6 - 94.7)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	151	86.0	(77.7 - 94.4)	9,526	87.8	(83.2 - 92.4)
Some College	74	90.8	(82.3 - 99.2)	6,462	93.1	(89.7 - 96.4)
≥ College Degree	162	91.8	(84.9 - 98.7)	7,345	94.9	(92.0 - 97.7)
Income						
≤ \$24,999	105	95.2	(90.4 - 100)	6,526	93.3	(90.1 - 96.5)
\$25,000-\$49,999	*	*	*	5,787	91.1	(85.7 - 96.5)
\$50,000-\$74,999	*	*	*	3,310	96.0	(91.9 - 100)
≥ \$75,000	93	91.0	(82.5 - 99.5)	4,570	95.0	(91.9 - 98.1)

¹ Defined as having ever had a Pap test among women

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 28a. Had a Pap test within the past 3 years among women¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	378	2,662,000	82.6	(77.3 - 87.8)	23,109	6,205,600	82.2	(79.8 - 84.6)
Gender								
Male	-	-	-	-	-	-	-	-
Female	378	2,662,000	82.6	(77.3 - 87.8)	23,109	6,205,600	82.2	(79.8 - 84.6)
Age (years)								
18-34	*	*	*	*	2,371	1,818,400	81.8	(75.5 - 88.1)
35-44	*	*	*	*	3,012	1,220,300	86.9	(81.8 - 92.0)
45-54	67	543,100	95.6	(90.9 - 100)	4,640	1,311,500	92.0	(89.8 - 94.3)
55-64	86	379,200	87.3	(79.5 - 95.1)	5,152	924,300	84.5	(81.0 - 88.0)
≥ 65	*	*	*	*	7,934	931,100	66.4	(61.8 - 71.0)
Race/Ethnicity								
White Non-Hispanic	224	906,500	76.4	(68.1 - 84.7)	21,720	3,807,100	81.5	(79.2 - 83.8)
Black Non-Hispanic	*	*	*	*	473	1,142,800	86.7	(79.1 - 94.3)
Hispanic	66	670,700	89.5	(80.5 - 98.6)	527	894,900	83.6	(76.3 - 91.0)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	143	1,104,700	78.9	(69.4 - 88.4)	9,384	2,132,600	76.0	(71.1 - 81.0)
Some College	73	576,100	86.3	(77.4 - 95.1)	6,401	1,594,300	84.0	(80.4 - 87.5)
≥ College Degree	160	917,400	84.0	(76.3 - 91.6)	7,281	2,406,000	86.9	(83.7 - 90.1)
Income								
≤ \$24,999	98	868,900	90.8	(85.2 - 96.5)	6,441	1,453,900	82.0	(77.9 - 86.0)
\$25,000-\$49,999	*	*	*	*	5,740	1,182,000	78.3	(72.6 - 83.9)
\$50,000-\$74,999	*	*	*	*	3,296	825,000	85.0	(78.0 - 91.9)
≥ \$75,000	93	571,200	84.5	(75.6 - 93.4)	4,555	1,810,200	89.0	(85.7 - 92.2)

¹ Defined as having had a Pap test within the past 3 years among women

² Sample size from survey

³ Estimated number of women who have had a Pap test within the past 3 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 28b. Had a Pap test within the past 3 years among women¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	378	82.7	(77.4 - 88.1)	23,109	82.7	(80.2 - 85.1)
Gender						
Male	-	-	-	-	-	-
Female	378	82.7	(77.4 - 88.1)	23,109	82.7	(80.2 - 85.1)
Race/Ethnicity						
White Non-Hispanic	224	76.5	(68.1 - 85.0)	21,720	82.3	(80.0 - 84.6)
Black Non-Hispanic	*	*	*	473	87.0	(79.3 - 94.6)
Hispanic	66	89.7	(80.7 - 98.8)	527	83.4	(75.9 - 90.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	143	79.2	(69.7 - 88.7)	9,384	76.4	(71.3 - 81.5)
Some College	73	86.0	(76.8 - 95.3)	6,401	84.2	(80.5 - 88.0)
≥ College Degree	160	84.2	(76.6 - 91.9)	7,281	87.5	(84.3 - 90.6)
Income						
≤ \$24,999	98	90.9	(85.0 - 96.7)	6,441	82.7	(78.5 - 86.9)
\$25,000-\$49,999	*	*	*	5,740	79.2	(73.5 - 84.8)
\$50,000-\$74,999	*	*	*	3,296	85.7	(78.7 - 92.8)
≥ \$75,000	93	84.1	(74.7 - 93.5)	4,555	88.9	(85.5 - 92.3)

¹ Defined as having had a Pap test within the past 3 years among women

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 29a. Ever had a digital rectal exam among men aged 40 and older¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	209	1,181,700	66.8	(57.6 - 76.0)	12,033	3,176,500	73.8	(69.7 - 77.8)
Gender								
Male	209	1,181,700	66.8	(57.6 - 76.0)	12,033	3,176,500	73.8	(69.7 - 77.8)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	3,098	922,800	66.8	(59.9 - 73.8)
55-64	*	*	*	*	3,504	865,600	87.1	(82.5 - 91.7)
≥ 65	59	382,900	96.2	(92.1 - 100)	4,348	953,900	92.9	(91.0 - 94.9)
Race/Ethnicity								
White Non-Hispanic	145	681,400	79.5	(71.2 - 87.7)	11,411	2,452,100	80.3	(77.7 - 82.9)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	5,013	931,600	64.1	(57.3 - 71.0)
Some College	*	*	*	*	2,695	674,200	75.5	(67.8 - 83.2)
≥ College Degree	*	*	*	*	4,302	1,568,000	80.5	(74.0 - 87.0)
Income								
≤ \$24,999	*	*	*	*	2,573	573,300	65.9	(56.0 - 75.9)
\$25,000-\$49,999	*	*	*	*	3,168	530,100	74.6	(67.2 - 82.1)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,309	1,351,100	78.0	(72.6 - 83.5)

¹ Defined as having ever had a digital rectal exam among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have ever had a digital rectal exam (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 29b. Ever had a digital rectal exam among men aged 40 and older¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	209	68.3	(59.2 - 77.3)	12,033	73.7	(69.6 - 77.8)
Gender						
Male	209	68.3	(59.2 - 77.3)	12,033	73.7	(69.6 - 77.8)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	145	80.3	(72.3 - 88.3)	11,411	79.9	(77.2 - 82.5)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	5,013	64.7	(57.8 - 71.7)
Some College	*	*	*	2,695	75.0	(67.1 - 82.9)
≥ College Degree	*	*	*	4,302	80.3	(73.8 - 86.8)
Income						
≤ \$24,999	*	*	*	2,573	66.9	(56.9 - 76.9)
\$25,000-\$49,999	*	*	*	3,168	74.8	(67.3 - 82.2)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,309	77.5	(72.0 - 83.0)

¹ Defined as having ever had a digital rectal exam among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 30a. Had digital rectal exam within the past 2 years among men aged 40 and older¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	207	875,600	50.0	(41.0 - 58.9)	11,923	2,380,300	55.8	(51.8 - 59.8)
Gender								
Male	207	875,600	50.0	(41.0 - 58.9)	11,923	2,380,300	55.8	(51.8 - 59.8)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	1,078	246,800	27.5	(18.5 - 36.5)
45-54	*	*	*	*	3,074	650,000	47.6	(40.8 - 54.4)
55-64	*	*	*	*	3,467	673,500	68.1	(62.4 - 73.8)
≥ 65	*	*	*	*	4,304	809,900	79.7	(75.0 - 84.3)
Race/Ethnicity								
White Non-Hispanic	143	523,100	62.1	(52.6 - 71.7)	11,305	1,875,600	62.2	(59.2 - 65.2)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	4,963	718,700	49.6	(43.2 - 56.1)
Some College	*	*	*	*	2,665	511,500	58.5	(50.8 - 66.3)
≥ College Degree	*	*	*	*	4,277	1,148,400	59.4	(52.8 - 66.0)
Income								
≤ \$24,999	*	*	*	*	2,553	388,000	45.0	(35.0 - 55.0)
\$25,000-\$49,999	*	*	*	*	3,139	422,400	60.5	(53.0 - 68.0)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,295	978,300	56.9	(51.1 - 62.7)

¹ Defined as having had a digital rectal exam within the past 2 years among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have had a digital rectal exam within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 30b. Had digital rectal exam within the past 2 years among men aged 40 and older¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	207	51.6	(42.6 - 60.6)	11,923	55.9	(51.7 - 60.0)
Gender						
Male	207	51.6	(42.6 - 60.6)	11,923	55.9	(51.7 - 60.0)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	143	63.4	(54.0 - 72.8)	11,305	62.0	(58.9 - 65.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	4,963	50.5	(43.7 - 57.3)
Some College	*	*	*	2,665	58.3	(50.4 - 66.2)
≥ College Degree	*	*	*	4,277	59.2	(52.4 - 65.9)
Income						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,139	60.8	(53.1 - 68.4)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,295	56.5	(50.5 - 62.4)

¹ Defined as having had a digital rectal exam within the past 2 years among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 31a. Ever had a prostate specific antigen test among men aged 40 and older¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	202	1,216,900	69.7	(61.5 - 77.9)	11,557	2,822,600	67.6	(63.9 - 71.2)
Gender								
Male	202	1,216,900	69.7	(61.5 - 77.9)	11,557	2,822,600	67.6	(63.9 - 71.2)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,938	760,600	57.4	(50.6 - 64.3)
55-64	*	*	*	*	3,373	811,400	83.4	(78.8 - 88.1)
≥ 65	57	356,000	90.9	(83.1 - 98.8)	4,215	905,400	90.4	(87.1 - 93.6)
Race/Ethnicity								
White Non-Hispanic	138	554,400	66.9	(57.1 - 76.6)	10,950	1,965,400	67.0	(63.9 - 70.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	4,775	829,200	59.0	(52.4 - 65.6)
Some College	*	*	*	*	2,584	535,000	61.9	(53.7 - 70.2)
≥ College Degree	110	693,100	81.6	(73.1 - 90.2)	4,176	1,455,700	76.8	(72.5 - 81.1)
Income								
≤ \$24,999	*	*	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	*	3,048	444,100	64.1	(56.0 - 72.3)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,188	1,211,100	72.3	(67.6 - 77.0)

¹ Defined as having ever had a prostate specific antigen test among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have ever had a prostate specific antigen test (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 31b. Ever had a prostate specific antigen test among men aged 40 and older¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	202	70.6	(62.6 - 78.7)	11,557	67.4	(63.7 - 71.1)
Gender						
Male	202	70.6	(62.6 - 78.7)	11,557	67.4	(63.7 - 71.1)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	138	68.2	(58.6 - 77.8)	10,950	66.6	(63.4 - 69.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	4,775	59.7	(52.9 - 66.4)
Some College	*	*	*	2,584	61.4	(53.0 - 69.9)
≥ College Degree	110	82.1	(73.7 - 90.5)	4,176	76.3	(71.9 - 80.7)
Income						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,048	64.2	(56.1 - 72.4)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,188	71.5	(66.6 - 76.3)

¹ Defined as having ever had a prostate specific antigen test among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 32a. Had prostate specific antigen test within the past 2 years among men aged 40 and older¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	202	1,076,200	61.6	(52.7 - 70.6)	11,437	2,493,800	59.9	(56.0 - 63.9)
Gender								
Male	202	1,076,200	61.6	(52.7 - 70.6)	11,437	2,493,800	59.9	(56.0 - 63.9)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,917	659,400	49.9	(42.8 - 57.0)
55-64	*	*	*	*	3,326	719,700	74.5	(69.4 - 79.7)
≥ 65	*	*	*	*	4,166	830,000	83.3	(78.4 - 88.3)
Race/Ethnicity								
White Non-Hispanic	138	521,800	62.9	(53.2 - 72.7)	10,836	1,762,200	60.4	(57.3 - 63.5)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	4,727	734,200	52.4	(45.7 - 59.1)
Some College	*	*	*	*	2,558	483,800	56.3	(48.2 - 64.4)
≥ College Degree	*	*	*	*	4,133	1,273,200	67.4	(61.6 - 73.3)
Income								
≤ \$24,999	*	*	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	*	3,021	410,800	59.6	(51.5 - 67.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,170	1,080,100	64.6	(59.1 - 70.0)

¹ Defined as having had a prostate specific antigen test within the past 2 years among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have had a prostate specific antigen test within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 32b. Had prostate specific antigen test within the past 2 years among men aged 40 and older¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	202	62.5	(53.7 - 71.4)	11,437	59.8	(55.8 - 63.9)
Gender						
Male	202	62.5	(53.7 - 71.4)	11,437	59.8	(55.8 - 63.9)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	138	64.3	(54.6 - 74.0)	10,836	60.1	(56.9 - 63.4)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	4,727	52.9	(45.9 - 59.9)
Some College	*	*	*	2,558	56.0	(47.7 - 64.2)
≥ College Degree	*	*	*	4,133	67.1	(61.2 - 73.1)
Income						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,021	59.8	(51.7 - 67.9)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,170	63.9	(58.3 - 69.6)

¹ Defined as having had a prostate specific antigen test within the past 2 years among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 33a. Prostate cancer among men aged 40 and older¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	210	60,900	3.4	(1.0 - 5.8)	12,133	178,200	4.1	(3.0 - 5.2)
Gender								
Male	210	60,900	3.4	(1.0 - 5.8)	12,133	178,200	4.1	(3.0 - 5.2)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	3,120	6,100	0.4	(0.0 - 0.9)
55-64	*	*	*	*	3,532	31,600	3.1	(1.5 - 4.8)
≥ 65	59	53,600	13.5	(4.0 - 22.9)	4,388	140,400	13.6	(9.7 - 17.4)
Race/Ethnicity								
White Non-Hispanic	*	*	*	*	11,502	143,700	4.7	(3.5 - 5.8)
Black Non-Hispanic	*	*	*	*	184	23,500	5.1	(0.0 - 11.0)
Hispanic	*	*	*	*	197	7,700	1.6	(0.0 - 3.7)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	5,065	46,300	3.1	(2.2 - 4.1)
Some College	*	*	*	*	2,709	39,200	4.4	(1.3 - 7.5)
≥ College Degree	*	*	*	*	4,336	92,500	4.7	(3.0 - 6.5)
Income								
≤ \$24,999	*	*	*	*	2,597	35,800	4.1	(2.2 - 6.0)
\$25,000-\$49,999	*	*	*	*	3,196	29,600	4.1	(2.9 - 5.4)
\$50,000-\$74,999	*	*	*	*	2,007	20,300	2.8	(1.2 - 4.4)
≥ \$75,000	*	*	*	*	3,322	56,500	3.2	(1.7 - 4.8)

¹ Defined as having ever been told by a health professional that respondent had prostate cancer among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have ever been told they had prostate cancer (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 33b. Prostate cancer among men aged 40 and older¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	210	4.0	(1.2 - 6.8)	12,133	4.5	(3.2 - 5.7)
Gender						
Male	210	4.0	(1.2 - 6.8)	12,133	4.5	(3.2 - 5.7)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	*	*	*	11,502	5.1	(3.8 - 6.4)
Black Non-Hispanic	*	*	*	184	5.7	(0.0 - 12.8)
Hispanic	*	*	*	197	1.8	(0.0 - 4.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	5,065	3.3	(2.3 - 4.3)
Some College	*	*	*	2,709	5.0	(1.1 - 9.0)
≥ College Degree	*	*	*	4,336	5.2	(3.1 - 7.2)
Income						
≤ \$24,999	*	*	*	2,597	4.5	(2.2 - 6.7)
\$25,000-\$49,999	*	*	*	3,196	4.3	(3.0 - 5.6)
\$50,000-\$74,999	*	*	*	2,007	3.0	(1.1 - 5.0)
≥ \$75,000	*	*	*	3,322	3.5	(1.6 - 5.4)

¹ Defined as having ever been told by a health professional that respondent had prostate cancer among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 34a. Home blood stool test ever used among adults aged 50 and older¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	409	796,700	33.8	(27.9 - 39.7)	25,322	2,288,500	38.0	(35.5 - 40.4)
Gender								
Male	162	396,700	36.9	(27.8 - 46.1)	9,571	1,081,900	38.9	(35.1 - 42.6)
Female	247	400,000	31.2	(23.5 - 38.9)	15,751	1,206,600	37.2	(34.0 - 40.4)
Age (years)								
50-64	223	446,600	32.9	(25.2 - 40.6)	12,834	1,192,000	34.0	(30.8 - 37.2)
≥ 65	186	350,000	35.0	(25.8 - 44.2)	12,472	1,093,400	43.6	(39.8 - 47.4)
Race/Ethnicity								
White Non-Hispanic	291	398,700	32.3	(26.0 - 38.7)	24,213	1,754,500	39.1	(37.1 - 41.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	145	295,600	29.0	(19.3 - 38.8)	11,146	823,500	34.9	(30.5 - 39.2)
Some College	*	*	*	*	6,261	480,300	37.0	(32.6 - 41.4)
≥ College Degree	190	373,700	40.0	(31.2 - 48.8)	7,851	973,700	41.6	(37.8 - 45.3)
Income								
≤ \$24,999	*	*	*	*	7,222	505,000	35.4	(29.7 - 41.2)
\$25,000-\$49,999	*	*	*	*	6,609	467,600	40.3	(35.2 - 45.4)
\$50,000-\$74,999	*	*	*	*	3,500	313,000	38.2	(32.6 - 43.9)
≥ \$75,000	*	*	*	*	4,647	657,800	37.3	(33.0 - 41.6)

¹ Defined as having ever used a home blood stool test among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have ever used a home blood stool test (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 34b. Home blood stool test ever used among adults aged 50 and older¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	409	34.4	(28.3 - 40.5)	25,322	38.1	(35.5 - 40.8)
Gender						
Male	162	37.5	(28.1 - 46.9)	9,571	39.4	(35.3 - 43.5)
Female	247	31.4	(23.6 - 39.2)	15,751	36.9	(33.5 - 40.3)
Race/Ethnicity						
White Non-Hispanic	291	32.2	(25.7 - 38.7)	24,213	38.9	(36.7 - 41.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	11,146	35.1	(30.3 - 39.9)
Some College	*	*	*	6,261	36.6	(32.0 - 41.3)
≥ College Degree	190	40.5	(31.5 - 49.4)	7,851	41.9	(37.9 - 46.0)
Income						
≤ \$24,999	*	*	*	7,222	35.9	(29.5 - 42.4)
\$25,000-\$49,999	*	*	*	6,609	40.2	(34.8 - 45.6)
\$50,000-\$74,999	*	*	*	3,500	38.1	(31.9 - 44.3)
≥ \$75,000	*	*	*	4,647	37.5	(32.9 - 42.2)

¹ Defined as having ever used a home blood stool test among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 35a. Home blood stool test used within the past year among adults aged 50 and older¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	405	311,200	13.3	(9.0 - 17.7)	25,074	863,300	14.4	(12.6 - 16.2)
Gender								
Male	161	158,700	14.9	(8.1 - 21.7)	9,508	444,400	16.0	(13.2 - 18.9)
Female	244	152,500	12.0	(6.4 - 17.6)	15,566	418,900	13.1	(10.8 - 15.4)
Age (years)								
50-64	221	182,800	13.6	(7.6 - 19.6)	12,759	454,400	13.0	(10.6 - 15.5)
≥ 65	184	128,400	13.0	(6.7 - 19.2)	12,299	406,400	16.4	(13.7 - 19.0)
Race/Ethnicity								
White Non-Hispanic	288	139,300	11.4	(7.0 - 15.9)	23,971	634,300	14.3	(12.8 - 15.7)
Black Non-Hispanic	*	*	*	*	396	136,000	18.6	(9.4 - 27.8)
Hispanic	*	*	*	*	349	53,400	9.9	(2.5 - 17.2)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	143	93,400	9.3	(2.7 - 15.9)	11,026	288,600	12.3	(9.4 - 15.3)
Some College	*	*	*	*	6,208	212,500	16.4	(12.5 - 20.4)
≥ College Degree	188	132,300	14.3	(8.1 - 20.6)	7,778	352,400	15.2	(12.5 - 17.9)
Income								
≤ \$24,999	110	75,500	10.5	(3.5 - 17.5)	7,157	196,300	13.9	(10.1 - 17.6)
\$25,000-\$49,999	*	*	*	*	6,562	221,500	19.2	(14.1 - 24.3)
\$50,000-\$74,999	*	*	*	*	3,469	111,900	13.7	(10.1 - 17.4)
≥ \$75,000	*	*	*	*	4,623	205,100	11.7	(8.8 - 14.6)

¹ Defined as having used a home blood stool test within the past year among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have used a home blood stool test within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 35b. Home blood stool test used within the past year among adults aged 50 and older¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	405	13.8	(9.2 - 18.4)	25,074	14.6	(12.6 - 16.7)
Gender						
Male	161	15.5	(8.3 - 22.8)	9,508	16.4	(13.2 - 19.6)
Female	244	12.1	(6.3 - 17.8)	15,566	12.9	(10.4 - 15.4)
Race/Ethnicity						
White Non-Hispanic	288	11.3	(6.8 - 15.9)	23,971	14.2	(12.7 - 15.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	349	10.2	(1.7 - 18.7)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	143	10.0	(2.8 - 17.3)	11,026	12.7	(9.2 - 16.1)
Some College	*	*	*	6,208	16.6	(12.3 - 20.9)
≥ College Degree	188	14.6	(8.1 - 21.1)	7,778	15.3	(12.4 - 18.3)
Income						
≤ \$24,999	110	11.6	(3.5 - 19.7)	7,157	14.3	(9.8 - 18.7)
\$25,000-\$49,999	*	*	*	6,562	19.3	(13.8 - 24.8)
\$50,000-\$74,999	*	*	*	3,469	13.8	(9.8 - 17.7)
≥ \$75,000	*	*	*	4,623	11.8	(8.6 - 15.0)

¹ Defined as having used a home blood stool test within the past year among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 36a. Home blood stool test used within the past 2 years among adults aged 50 and older¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	405	445,300	19.1	(13.9 - 24.2)	25,074	1,218,000	20.4	(18.3 - 22.5)
Gender								
Male	161	207,900	19.5	(12.1 - 26.9)	9,508	593,100	21.4	(18.3 - 24.5)
Female	244	237,400	18.7	(11.6 - 25.8)	15,566	624,900	19.5	(16.6 - 22.4)
Age (years)								
50-64	221	245,100	18.2	(11.7 - 24.8)	12,759	629,900	18.1	(15.4 - 20.7)
≥ 65	184	200,200	20.2	(12.0 - 28.4)	12,299	585,500	23.6	(20.2 - 27.0)
Race/Ethnicity								
White Non-Hispanic	288	183,100	15.0	(10.2 - 19.9)	23,971	877,500	19.7	(18.2 - 21.3)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	143	170,400	16.9	(8.2 - 25.6)	11,026	454,600	19.4	(15.6 - 23.3)
Some College	*	*	*	*	6,208	262,200	20.3	(16.3 - 24.2)
≥ College Degree	188	184,600	20.0	(12.9 - 27.1)	7,778	491,400	21.2	(18.1 - 24.2)
Income								
≤ \$24,999	110	100,100	14.0	(6.5 - 21.5)	7,157	263,900	18.6	(14.6 - 22.7)
\$25,000-\$49,999	*	*	*	*	6,562	285,900	24.8	(19.6 - 30.0)
\$50,000-\$74,999	*	*	*	*	3,469	167,500	20.6	(15.6 - 25.5)
≥ \$75,000	115	87,800	13.6	(5.8 - 21.3)	4,623	294,900	16.8	(13.7 - 20.0)

¹ Defined as having used a home blood stool test within the past 2 years among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have used a home blood stool test within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

**Table 36b. Home blood stool test used within the past 2 years among adults aged 50 and older¹
in New York City and New York State - Age-Adjusted²**

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	405	19.4	(14.1 - 24.7)	25,074	20.5	(18.2 - 22.8)
Gender						
Male	161	19.9	(12.2 - 27.7)	9,508	21.7	(18.3 - 25.1)
Female	244	18.9	(11.7 - 26.1)	15,566	19.4	(16.3 - 22.5)
Race/Ethnicity						
White Non-Hispanic	288	15.0	(10.1 - 20.0)	23,971	19.7	(18.0 - 21.3)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	143	17.2	(8.3 - 26.2)	11,026	19.5	(15.4 - 23.7)
Some College	*	*	*	6,208	20.4	(16.1 - 24.7)
≥ College Degree	188	20.4	(13.1 - 27.7)	7,778	21.4	(18.1 - 24.7)
Income						
≤ \$24,999	110	14.9	(6.4 - 23.3)	7,157	18.9	(14.2 - 23.6)
\$25,000-\$49,999	*	*	*	6,562	24.8	(19.3 - 30.3)
\$50,000-\$74,999	*	*	*	3,469	20.6	(15.3 - 25.9)
≥ \$75,000	115	14.0	(5.9 - 22.2)	4,623	16.9	(13.4 - 20.4)

¹ Defined as having used a home blood stool test within the past 2 years among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 37a. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	410	1,509,200	63.9	(57.9 - 70.0)	25,357	4,019,100	66.6	(64.1 - 69.1)
Gender								
Male	163	731,300	67.8	(58.8 - 76.9)	9,582	1,894,100	68.0	(64.2 - 71.7)
Female	247	778,000	60.7	(52.6 - 68.7)	15,775	2,125,000	65.5	(62.1 - 68.8)
Age (years)								
50-54	*	*	*	*	4,179	718,900	50.6	(44.6 - 56.5)
55-64	158	576,700	70.4	(61.5 - 79.2)	8,695	1,493,500	71.0	(67.3 - 74.8)
≥ 65	184	673,900	67.9	(59.2 - 76.5)	12,466	1,802,000	72.1	(68.6 - 75.7)
Race/Ethnicity								
White Non-Hispanic	292	823,100	66.6	(60.3 - 72.8)	24,248	3,093,900	68.8	(66.8 - 70.8)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	11,121	1,445,100	61.2	(56.7 - 65.8)
Some College	*	*	*	*	6,287	835,300	64.3	(59.7 - 69.0)
≥ College Degree	191	664,100	70.8	(62.1 - 79.4)	7,884	1,732,400	73.7	(70.0 - 77.4)
Income								
≤ \$24,999	*	*	*	*	7,220	892,600	62.7	(56.8 - 68.6)
\$25,000-\$49,999	*	*	*	*	6,619	697,300	60.1	(54.6 - 65.7)
\$50,000-\$74,999	*	*	*	*	3,514	613,700	74.2	(68.9 - 79.6)
≥ \$75,000	*	*	*	*	4,665	1,294,400	73.3	(69.2 - 77.5)

¹ Defined as having ever had a sigmoidoscopy or colonoscopy among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have ever had a sigmoidoscopy or colonoscopy (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 37b. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	410	64.2	(58.1 - 70.3)	25,357	66.6	(63.9 - 69.3)
Gender						
Male	163	68.2	(59.2 - 77.3)	9,582	68.5	(64.5 - 72.5)
Female	247	60.2	(52.1 - 68.4)	15,775	64.8	(61.2 - 68.4)
Race/Ethnicity						
White Non-Hispanic	292	66.3	(59.8 - 72.9)	24,248	68.7	(66.5 - 70.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	11,121	61.3	(56.4 - 66.2)
Some College	*	*	*	6,287	63.8	(58.8 - 68.8)
≥ College Degree	191	71.7	(63.2 - 80.3)	7,884	74.0	(70.2 - 77.9)
Income						
≤ \$24,999	*	*	*	7,220	63.0	(56.7 - 69.3)
\$25,000-\$49,999	*	*	*	6,619	59.7	(53.8 - 65.6)
\$50,000-\$74,999	*	*	*	3,514	74.4	(68.3 - 80.5)
≥ \$75,000	*	*	*	4,665	73.6	(69.3 - 78.0)

¹ Defined as having ever had a sigmoidoscopy or colonoscopy among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 38a. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	407	1,436,100	62.0	(56.0 - 68.1)	25,169	3,855,800	64.6	(62.1 - 67.1)
Gender								
Male	162	709,000	66.5	(57.3 - 75.7)	9,521	1,835,400	66.4	(62.6 - 70.1)
Female	245	727,200	58.2	(50.2 - 66.3)	15,648	2,020,400	63.1	(59.8 - 66.4)
Age (years)								
50-54	*	*	*	*	4,169	700,500	49.3	(43.3 - 55.3)
55-64	158	566,100	69.1	(60.1 - 78.0)	8,652	1,455,100	69.4	(65.7 - 73.2)
≥ 65	181	616,900	65.1	(56.3 - 73.9)	12,331	1,695,700	69.5	(65.9 - 73.1)
Race/Ethnicity								
White Non-Hispanic	292	795,300	64.3	(57.9 - 70.7)	24,069	2,981,500	66.6	(64.6 - 68.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	11,017	1,360,400	59.0	(54.5 - 63.5)
Some College	*	*	*	*	6,251	807,100	62.4	(57.8 - 67.1)
≥ College Degree	191	644,600	68.7	(60.0 - 77.4)	7,841	1,682,500	71.7	(68.0 - 75.4)
Income								
≤ \$24,999	*	*	*	*	7,163	858,900	61.0	(55.1 - 67.0)
\$25,000-\$49,999	*	*	*	*	6,584	664,900	57.7	(52.2 - 63.2)
\$50,000-\$74,999	*	*	*	*	3,501	602,600	73.1	(67.7 - 78.5)
≥ \$75,000	*	*	*	*	4,655	1,260,400	71.5	(67.3 - 75.7)

¹ Defined as having had a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have had a sigmoidoscopy or colonoscopy within the past 10 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 38b. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	407	62.2	(56.1 - 68.4)	25,169	64.6	(61.9 - 67.3)
Gender						
Male	162	66.9	(57.7 - 76.0)	9,521	66.9	(62.9 - 70.9)
Female	245	57.7	(49.6 - 65.9)	15,648	62.4	(58.9 - 66.0)
Race/Ethnicity						
White Non-Hispanic	292	64.1	(57.5 - 70.8)	24,069	66.5	(64.3 - 68.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	11,017	59.0	(54.1 - 63.9)
Some College	*	*	*	6,251	61.9	(56.9 - 66.9)
≥ College Degree	191	69.7	(61.1 - 78.3)	7,841	72.1	(68.2 - 75.9)
Income						
≤ \$24,999	*	*	*	7,163	61.3	(54.9 - 67.6)
\$25,000-\$49,999	*	*	*	6,584	57.2	(51.4 - 63.0)
\$50,000-\$74,999	*	*	*	3,501	73.3	(67.2 - 79.4)
≥ \$75,000	*	*	*	4,655	71.8	(67.4 - 76.2)

¹ Defined as having had a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 39a. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	395	1,525,000	67.2	(61.3 - 73.2)	24,792	4,049,200	69.1	(66.7 - 71.6)
Gender								
Male	162	768,200	71.5	(62.7 - 80.3)	9,418	1,945,200	70.7	(67.0 - 74.3)
Female	233	756,800	63.4	(55.4 - 71.4)	15,374	2,104,100	67.8	(64.6 - 71.1)
Age (years)								
50-54	*	*	*	*	4,132	751,200	54.1	(48.0 - 60.1)
55-64	149	579,000	73.7	(64.7 - 82.6)	8,366	1,487,600	73.7	(70.0 - 77.4)
≥ 65	182	667,700	69.6	(61.2 - 77.9)	12,294	1,810,500	74.0	(70.6 - 77.4)
Race/Ethnicity								
White Non-Hispanic	281	804,700	67.2	(60.8 - 73.6)	23,726	3,091,400	70.4	(68.4 - 72.4)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	10,872	1,461,600	64.2	(59.8 - 68.7)
Some College	*	*	*	*	6,170	873,300	68.4	(63.9 - 72.8)
≥ College Degree	185	650,900	71.0	(62.4 - 79.6)	7,708	1,709,000	74.6	(70.9 - 78.3)
Income								
≤ \$24,999	*	*	*	*	7,092	923,300	66.0	(60.1 - 71.9)
\$25,000-\$49,999	*	*	*	*	6,526	730,000	64.2	(58.9 - 69.6)
\$50,000-\$74,999	*	*	*	*	3,467	621,500	76.6	(71.4 - 81.9)
≥ \$75,000	116	481,500	73.5	(63.7 - 83.4)	4,615	1,292,500	73.8	(69.8 - 77.9)

¹ Defined as having used a home blood stool test within the past year or a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 years and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have used a home blood stool test in the past year or sigmoidoscopy or colonoscopy within the past 10 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 39b. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	395	67.8	(61.8 - 73.8)	24,792	69.3	(66.7 - 71.9)
Gender						
Male	162	72.4	(63.7 - 81.0)	9,418	71.5	(67.6 - 75.4)
Female	233	63.1	(55.0 - 71.2)	15,374	67.2	(63.7 - 70.7)
Race/Ethnicity						
White Non-Hispanic	281	67.1	(60.5 - 73.7)	23,726	70.3	(68.1 - 72.4)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	10,872	64.5	(59.8 - 69.3)
Some College	*	*	*	6,170	68.2	(63.4 - 72.9)
≥ College Degree	185	72.3	(63.9 - 80.7)	7,708	75.0	(71.2 - 78.8)
Income						
≤ \$24,999	*	*	*	7,092	66.4	(60.2 - 72.6)
\$25,000-\$49,999	*	*	*	6,526	63.8	(58.2 - 69.5)
\$50,000-\$74,999	*	*	*	3,467	76.7	(70.6 - 82.7)
≥ \$75,000	116	74.5	(64.9 - 84.2)	4,615	74.4	(70.2 - 78.6)

¹ Defined as having used a home blood stool test within the past year or a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 years and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 40a. High blood pressure among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	646	1,517,400	23.9	(19.8 - 28.0)	38,125	3,974,800	26.8	(24.9 - 28.7)
Gender								
Male	251	702,500	23.6	(17.0 - 30.1)	14,611	1,960,500	27.7	(24.7 - 30.7)
Female	395	814,900	24.1	(19.0 - 29.2)	23,514	2,014,300	26.0	(23.6 - 28.3)
Age (years)								
18-34	*	*	*	*	3,935	312,200	7.0	(4.2 - 9.8)
35-44	102	128,400	9.7	(3.6 - 15.9)	4,984	426,200	14.8	(11.6 - 18.1)
45-54	*	*	*	*	7,825	862,400	30.3	(25.9 - 34.8)
55-64	160	328,200	38.7	(29.2 - 48.2)	8,765	890,800	41.6	(37.6 - 45.7)
≥ 65	186	567,600	56.4	(47.2 - 65.5)	12,616	1,483,200	58.5	(54.7 - 62.3)
Race/Ethnicity								
White Non-Hispanic	393	597,300	24.5	(19.5 - 29.6)	35,814	2,650,900	28.4	(26.9 - 30.0)
Black Non-Hispanic	93	321,700	21.8	(13.0 - 30.6)	736	558,500	26.2	(19.6 - 32.8)
Hispanic	116	439,400	26.8	(17.4 - 36.2)	863	538,100	23.8	(17.0 - 30.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	231	840,200	31.6	(24.1 - 39.1)	15,688	1,851,200	33.1	(29.4 - 36.9)
Some College	118	213,400	16.9	(9.4 - 24.4)	9,906	851,100	24.1	(20.9 - 27.3)
≥ College Degree	293	449,800	19.1	(13.3 - 24.8)	12,452	1,253,400	22.3	(19.7 - 24.9)
Income								
≤ \$24,999	169	577,900	29.2	(20.3 - 38.0)	9,640	1,153,000	32.8	(27.6 - 38.1)
\$25,000-\$49,999	106	230,200	19.8	(11.0 - 28.6)	9,657	807,800	27.9	(23.7 - 32.1)
\$50,000-\$74,999	*	*	*	*	5,773	507,600	25.8	(21.4 - 30.1)
≥ \$75,000	189	304,300	19.8	(13.0 - 26.6)	8,560	942,500	21.0	(18.4 - 23.6)

¹ Defined as having ever been told by a health professional that respondent has high blood pressure

² Sample size from survey

³ Estimated number of adults who have ever been told by a health professional that they have high blood pressure (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 40b. High blood pressure among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	646	23.8	(19.7 - 27.8)	38,125	25.7	(23.8 - 27.5)
Gender						
Male	251	24.6	(18.1 - 31.2)	14,611	27.7	(24.7 - 30.7)
Female	395	23.0	(18.0 - 28.0)	23,514	23.8	(21.5 - 26.0)
Race/Ethnicity						
White Non-Hispanic	393	24.9	(19.8 - 30.0)	35,814	27.2	(25.6 - 28.7)
Black Non-Hispanic	93	21.2	(12.6 - 29.9)	736	24.9	(18.4 - 31.3)
Hispanic	116	26.6	(17.3 - 35.9)	863	23.3	(16.6 - 30.0)
Other Non-Hispanic	*	*	*	712	19.3	(9.5 - 29.1)
Education						
≤ High School	231	31.5	(24.0 - 38.9)	15,688	32.1	(28.4 - 35.8)
Some College	118	16.6	(9.2 - 24.0)	9,906	22.8	(19.7 - 25.9)
≥ College Degree	293	19.1	(13.4 - 24.8)	12,452	21.2	(18.6 - 23.8)
Income						
≤ \$24,999	169	29.1	(20.4 - 37.8)	9,640	31.8	(26.7 - 37.0)
\$25,000-\$49,999	106	19.9	(11.1 - 28.6)	9,657	26.9	(22.8 - 30.9)
\$50,000-\$74,999	*	*	*	5,773	24.8	(20.4 - 29.2)
≥ \$75,000	189	19.9	(13.0 - 26.7)	8,560	19.9	(17.3 - 22.5)

¹ Defined as having ever been told by a health professional that respondent has high blood pressure

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 41a. Blood pressure medication among adults with high blood pressure¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	212	1,238,100	81.6	(73.7 - 89.5)	15,288	3,145,800	79.2	(76.0 - 82.4)
Gender								
Male	*	*	*	*	6,080	1,432,100	73.1	(67.6 - 78.5)
Female	134	721,300	88.5	(81.7 - 95.4)	9,208	1,713,700	85.2	(82.1 - 88.2)
Age (years)								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,315	645,500	74.9	(66.8 - 82.9)
55-64	58	296,400	90.3	(82.3 - 98.3)	4,045	782,300	87.8	(84.5 - 91.1)
≥ 65	103	530,500	93.5	(87.7 - 99.2)	7,675	1,383,400	93.4	(91.1 - 95.7)
Race/Ethnicity								
White Non-Hispanic	125	502,200	84.1	(75.0 - 93.1)	14,442	2,117,300	79.9	(77.5 - 82.4)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	7,354	1,482,800	80.2	(74.7 - 85.6)
Some College	*	*	*	*	3,868	632,000	74.3	(67.8 - 80.7)
≥ College Degree	72	399,500	88.8	(79.7 - 97.9)	4,035	1,013,200	80.9	(76.8 - 84.9)
Income								
≤ \$24,999	*	*	*	*	4,898	948,200	82.3	(75.7 - 88.9)
\$25,000-\$49,999	*	*	*	*	4,088	668,300	82.8	(77.6 - 88.0)
\$50,000-\$74,999	*	*	*	*	1,971	391,500	77.2	(67.7 - 86.6)
≥ \$75,000	*	*	*	*	2,397	683,700	72.5	(66.1 - 78.9)

¹ Defined as currently taking medicine for high blood pressure among adults with high blood pressure

² Sample size from survey

³ Estimated number of adults with high blood pressure who are taking medicine for high blood pressure (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 41b. Blood pressure medication among adults with high blood pressure¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	212	81.7	(73.9 - 89.4)	15,288	78.5	(75.2 - 81.7)
Gender						
Male	*	*	*	6,080	73.3	(67.9 - 78.7)
Female	134	88.0	(80.9 - 95.2)	9,208	84.0	(80.7 - 87.2)
Race/Ethnicity						
White Non-Hispanic	125	84.3	(75.4 - 93.3)	14,442	79.0	(76.5 - 81.6)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	7,354	79.7	(74.3 - 85.2)
Some College	*	*	*	3,868	72.5	(65.7 - 79.3)
≥ College Degree	72	89.0	(80.0 - 98.0)	4,035	80.4	(76.1 - 84.6)
Income						
≤ \$24,999	*	*	*	4,898	81.6	(75.0 - 88.3)
\$25,000-\$49,999	*	*	*	4,088	82.1	(76.7 - 87.5)
\$50,000-\$74,999	*	*	*	1,971	76.4	(66.7 - 86.1)
≥ \$75,000	*	*	*	2,397	71.6	(64.8 - 78.3)

¹ Defined as currently taking medicine for high blood pressure among adults with high blood pressure

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 42a. Ever had blood cholesterol checked among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	630	4,648,200	76.3	(70.4 - 82.1)	37,450	11,496,600	80.3	(77.7 - 83.0)
Gender								
Male	243	2,038,600	72.4	(62.6 - 82.2)	14,389	5,288,400	77.7	(73.3 - 82.0)
Female	387	2,609,600	79.5	(72.8 - 86.3)	23,061	6,208,200	82.7	(79.6 - 85.8)
Age (years)								
18-34	*	*	*	*	3,684	2,296,400	55.3	(48.7 - 61.9)
35-44	*	*	*	*	4,883	2,264,200	81.6	(76.6 - 86.5)
45-54	122	966,300	87.9	(78.3 - 97.6)	7,731	2,539,400	91.0	(87.0 - 94.9)
55-64	158	786,600	94.7	(90.4 - 99.0)	8,718	2,014,900	95.4	(93.5 - 97.2)
≥ 65	182	919,000	94.2	(90.0 - 98.4)	12,434	2,381,700	95.9	(94.3 - 97.6)
Race/Ethnicity								
White Non-Hispanic	387	2,111,200	88.6	(83.3 - 93.8)	35,198	7,832,600	86.3	(84.8 - 87.9)
Black Non-Hispanic	*	*	*	*	720	1,588,900	77.8	(68.4 - 87.2)
Hispanic	*	*	*	*	842	1,224,600	56.8	(46.8 - 66.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	225	1,688,200	65.8	(55.8 - 75.7)	15,342	3,795,700	70.6	(65.6 - 75.6)
Some College	*	*	*	*	9,732	2,599,800	77.4	(71.7 - 83.2)
≥ College Degree	287	2,122,000	92.6	(87.3 - 97.9)	12,296	5,008,300	91.3	(88.9 - 93.6)
Income								
≤ \$24,999	*	*	*	*	9,437	2,277,500	68.4	(61.5 - 75.3)
\$25,000-\$49,999	*	*	*	*	9,492	2,053,000	71.9	(64.5 - 79.3)
\$50,000-\$74,999	76	660,700	96.7	(92.3 - 100)	5,721	1,736,100	90.0	(87.7 - 92.3)
≥ \$75,000	185	1,440,700	97.5	(95.3 - 99.6)	8,460	4,019,300	92.2	(90.8 - 93.6)

¹ Defined as ever having blood cholesterol checked

² Sample size from survey

³ Estimated number of adults who have ever had blood cholesterol checked (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 42b. Ever had blood cholesterol checked among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	630	76.6	(70.9 - 82.3)	37,450	79.9	(77.3 - 82.5)
Gender						
Male	243	73.8	(64.5 - 83.2)	14,389	78.0	(73.9 - 82.1)
Female	387	78.9	(72.1 - 85.8)	23,061	81.6	(78.4 - 84.8)
Race/Ethnicity						
White Non-Hispanic	387	88.7	(83.6 - 93.8)	35,198	85.6	(84.0 - 87.2)
Black Non-Hispanic	*	*	*	720	77.5	(68.2 - 86.7)
Hispanic	*	*	*	842	57.2	(47.4 - 67.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	225	66.5	(56.8 - 76.2)	15,342	70.2	(65.2 - 75.1)
Some College	*	*	*	9,732	76.7	(71.1 - 82.3)
≥ College Degree	287	92.6	(87.5 - 97.8)	12,296	90.9	(88.6 - 93.2)
Income						
≤ \$24,999	*	*	*	9,437	68.1	(61.3 - 74.9)
\$25,000-\$49,999	*	*	*	9,492	71.6	(64.4 - 78.7)
\$50,000-\$74,999	76	97.0	(92.9 - 100)	5,721	89.6	(87.2 - 92.0)
≥ \$75,000	185	97.4	(95.2 - 99.6)	8,460	91.7	(90.1 - 93.2)

¹ Defined as ever having blood cholesterol checked

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 43a. Blood cholesterol checked within the past 5 years among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	626	4,473,300	74.2	(68.3 - 80.1)	37,201	11,035,200	77.8	(75.1 - 80.4)
Gender								
Male	240	1,915,000	69.4	(59.5 - 79.3)	14,307	5,020,700	74.5	(70.2 - 78.9)
Female	386	2,558,300	78.2	(71.5 - 85.0)	22,894	6,014,600	80.7	(77.6 - 83.8)
Age (years)								
18-34	*	*	*	*	3,647	2,179,800	53.3	(46.7 - 59.9)
35-44	*	*	*	*	4,862	2,161,200	78.0	(72.8 - 83.2)
45-54	*	*	*	*	7,697	2,408,600	86.6	(82.4 - 90.8)
55-64	157	766,800	93.5	(88.9 - 98.1)	8,677	1,957,900	93.3	(91.4 - 95.3)
≥ 65	180	889,800	93.0	(88.6 - 97.4)	12,318	2,327,800	94.8	(93.1 - 96.6)
Race/Ethnicity								
White Non-Hispanic	385	2,033,400	86.8	(81.5 - 92.2)	34,960	7,486,600	83.4	(81.8 - 85.0)
Black Non-Hispanic	*	*	*	*	719	1,561,700	76.5	(67.1 - 85.9)
Hispanic	*	*	*	*	838	1,184,000	55.6	(45.6 - 65.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,215	3,612,500	68.3	(63.2 - 73.3)
Some College	*	*	*	*	9,673	2,524,200	75.4	(69.8 - 81.1)
≥ College Degree	286	2,052,900	89.7	(83.9 - 95.5)	12,236	4,806,500	88.1	(85.5 - 90.6)
Income								
≤ \$24,999	*	*	*	*	9,350	2,214,800	66.9	(60.1 - 73.8)
\$25,000-\$49,999	*	*	*	*	9,454	1,936,600	68.8	(61.5 - 76.1)
\$50,000-\$74,999	75	613,400	91.8	(82.9 - 100)	5,701	1,640,400	85.9	(82.4 - 89.5)
≥ \$75,000	185	1,400,300	94.7	(91.6 - 97.9)	8,435	3,872,400	89.2	(87.5 - 90.9)

¹ Defined as having had blood cholesterol checked within the past 5 years

² Sample size from survey

³ Estimated number of adults who have had blood cholesterol checked within the past 5 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 43b. Blood cholesterol checked within the past 5 years among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	626	74.6	(68.8 - 80.3)	37,201	77.3	(74.7 - 79.9)
Gender						
Male	240	70.9	(61.4 - 80.3)	14,307	74.8	(70.7 - 79.0)
Female	386	77.7	(70.8 - 84.6)	22,894	79.5	(76.3 - 82.7)
Race/Ethnicity						
White Non-Hispanic	385	87.0	(81.8 - 92.3)	34,960	82.6	(81.0 - 84.3)
Black Non-Hispanic	*	*	*	719	76.2	(66.9 - 85.5)
Hispanic	*	*	*	838	55.9	(46.1 - 65.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	222	64.2	(54.4 - 74.0)	15,215	67.8	(62.9 - 72.8)
Some College	*	*	*	9,673	74.7	(69.1 - 80.2)
≥ College Degree	286	89.8	(84.1 - 95.4)	12,236	87.5	(85.0 - 90.1)
Income						
≤ \$24,999	*	*	*	9,350	66.6	(59.8 - 73.4)
\$25,000-\$49,999	*	*	*	9,454	68.5	(61.4 - 75.6)
\$50,000-\$74,999	75	92.2	(83.4 - 100)	5,701	85.4	(81.7 - 89.0)
≥ \$75,000	185	94.7	(91.4 - 97.9)	8,435	88.5	(86.7 - 90.3)

¹ Defined as having had blood cholesterol checked within the past 5 years

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 44a. Cardiovascular disease among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	640	490,000	7.8	(5.2 - 10.3)	37,810	1,135,000	7.7	(6.5 - 8.8)
Gender								
Male	248	274,000	9.3	(4.9 - 13.6)	14,477	664,600	9.4	(7.5 - 11.4)
Female	392	215,900	6.4	(3.5 - 9.3)	23,333	470,400	6.1	(4.8 - 7.4)
Age (years)								
18-34	*	*	*	*	3,930	100,800	2.3	(0.0 - 4.8)
35-44	*	*	*	*	4,971	103,400	3.6	(1.5 - 5.7)
45-54	*	*	*	*	7,768	131,300	4.6	(2.9 - 6.4)
55-64	157	78,000	9.4	(4.1 - 14.6)	8,699	224,200	10.6	(8.3 - 12.9)
≥ 65	184	225,900	22.6	(15.2 - 30.0)	12,442	575,300	23.0	(19.9 - 26.1)
Race/Ethnicity								
White Non-Hispanic	389	218,100	9.1	(5.9 - 12.2)	35,522	772,200	8.3	(7.4 - 9.2)
Black Non-Hispanic	*	*	*	*	728	140,700	6.6	(3.3 - 9.9)
Hispanic	114	141,600	8.7	(1.5 - 15.8)	854	163,300	7.3	(2.0 - 12.5)
Other Non-Hispanic	*	*	*	*	706	58,800	5.2	(0.9 - 9.5)
Education								
≤ High School	226	274,800	10.5	(5.3 - 15.7)	15,486	590,800	10.7	(8.2 - 13.2)
Some College	115	77,800	6.2	(1.7 - 10.7)	9,849	227,100	6.4	(4.7 - 8.1)
≥ College Degree	295	137,400	5.8	(3.1 - 8.5)	12,393	314,600	5.6	(4.3 - 6.9)
Income								
≤ \$24,999	166	237,300	12.0	(5.4 - 18.7)	9,505	448,100	12.9	(9.1 - 16.7)
\$25,000-\$49,999	*	*	*	*	9,571	179,300	6.2	(4.8 - 7.6)
\$50,000-\$74,999	77	79,900	11.2	(3.3 - 19.2)	5,751	166,900	8.5	(5.5 - 11.4)
≥ \$75,000	*	*	*	*	8,538	183,500	4.1	(2.9 - 5.3)

¹ Defined as having ever been told by a health professional that respondent had a heart attack or angina or a stroke

² Sample size from survey

³ Estimated number of adults with cardiovascular disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 44b. Cardiovascular disease among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	640	8.1	(5.5 - 10.7)	37,810	7.6	(6.4 - 8.7)
Gender						
Male	248	10.2	(5.7 - 14.7)	14,477	9.7	(7.8 - 11.7)
Female	392	6.3	(3.3 - 9.2)	23,333	5.6	(4.3 - 6.9)
Race/Ethnicity						
White Non-Hispanic	389	9.8	(6.3 - 13.3)	35,522	8.2	(7.2 - 9.1)
Black Non-Hispanic	*	*	*	728	6.8	(3.2 - 10.4)
Hispanic	114	8.3	(1.5 - 15.2)	854	7.0	(2.0 - 12.0)
Other Non-Hispanic	*	*	*	706	5.5	(0.9 - 10.0)
Education						
≤ High School	226	10.7	(5.6 - 15.9)	15,486	10.5	(8.0 - 13.0)
Some College	115	6.7	(1.8 - 11.7)	9,849	6.4	(4.5 - 8.2)
≥ College Degree	295	6.1	(3.3 - 9.0)	12,393	5.5	(4.2 - 6.9)
Income						
≤ \$24,999	166	12.5	(5.9 - 19.2)	9,505	12.8	(9.0 - 16.6)
\$25,000-\$49,999	*	*	*	9,571	6.0	(4.7 - 7.4)
\$50,000-\$74,999	77	11.9	(3.4 - 20.4)	5,751	8.5	(5.4 - 11.6)
≥ \$75,000	*	*	*	8,538	4.0	(2.7 - 5.2)

¹ Defined as having ever been told by a health professional that respondent had a heart attack or angina or a stroke

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 45a. Coronary heart disease among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	641	396,400	6.3	(4.0 - 8.6)	37,838	929,400	6.3	(5.3 - 7.3)
Gender								
Male	249	263,600	8.9	(4.6 - 13.2)	14,479	598,600	8.5	(6.6 - 10.4)
Female	392	132,800	3.9	(1.9 - 6.0)	23,359	330,800	4.3	(3.4 - 5.2)
Age (years)								
18-34	*	*	*	*	3,932	63,400	1.4	(0.0 - 3.6)
35-44	*	*	*	*	4,973	95,300	3.3	(1.2 - 5.4)
45-54	*	*	*	*	7,778	108,300	3.8	(2.1 - 5.6)
55-64	158	67,300	8.0	(3.0 - 13.0)	8,703	188,600	8.9	(6.7 - 11.1)
≥ 65	184	177,900	17.8	(11.0 - 24.5)	12,452	473,800	18.9	(16.1 - 21.7)
Race/Ethnicity								
White Non-Hispanic	390	182,700	7.6	(4.6 - 10.5)	35,552	636,700	6.9	(6.0 - 7.7)
Black Non-Hispanic	*	*	*	*	728	113,100	5.3	(2.3 - 8.4)
Hispanic	114	110,000	6.7	(0.5 - 13.0)	853	128,000	5.7	(1.1 - 10.3)
Other Non-Hispanic	*	*	*	*	705	51,500	4.6	(0.5 - 8.7)
Education								
≤ High School	227	216,800	8.2	(3.6 - 12.9)	15,485	474,500	8.6	(6.3 - 10.8)
Some College	*	*	*	*	9,860	193,000	5.5	(3.9 - 7.1)
≥ College Degree	295	111,400	4.7	(2.2 - 7.2)	12,411	259,700	4.6	(3.4 - 5.8)
Income								
≤ \$24,999	166	195,200	9.9	(3.9 - 15.9)	9,507	372,300	10.7	(7.3 - 14.1)
\$25,000-\$49,999	*	*	*	*	9,585	136,300	4.7	(3.8 - 5.7)
\$50,000-\$74,999	77	79,900	11.2	(3.3 - 19.2)	5,755	154,600	7.8	(4.9 - 10.7)
≥ \$75,000	*	*	*	*	8,541	151,800	3.4	(2.3 - 4.5)

¹ Defined as having ever been told by a health professional that respondent had a heart attack or angina

² Sample size from survey

³ Estimated number of adults with coronary heart disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 45b. Coronary heart disease among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	641	6.7	(4.3 - 9.0)	37,838	6.3	(5.2 - 7.3)
Gender						
Male	249	9.8	(5.4 - 14.3)	14,479	8.8	(6.9 - 10.8)
Female	392	3.9	(1.8 - 5.9)	23,359	3.9	(3.0 - 4.8)
Race/Ethnicity						
White Non-Hispanic	390	8.4	(5.1 - 11.8)	35,552	6.8	(5.9 - 7.7)
Black Non-Hispanic	*	*	*	728	5.6	(2.2 - 9.0)
Hispanic	114	6.4	(0.5 - 12.2)	853	5.4	(1.1 - 9.6)
Other Non-Hispanic	*	*	*	705	4.8	(0.4 - 9.2)
Education						
≤ High School	227	8.6	(4.0 - 13.2)	15,485	8.5	(6.2 - 10.7)
Some College	*	*	*	9,860	5.5	(3.7 - 7.2)
≥ College Degree	295	5.1	(2.4 - 7.8)	12,411	4.7	(3.4 - 5.9)
Income						
≤ \$24,999	166	10.4	(4.4 - 16.3)	9,507	10.7	(7.2 - 14.1)
\$25,000-\$49,999	*	*	*	9,585	4.7	(3.6 - 5.7)
\$50,000-\$74,999	77	11.9	(3.4 - 20.4)	5,755	8.0	(4.8 - 11.1)
≥ \$75,000	*	*	*	8,541	3.3	(2.1 - 4.5)

¹ Defined as having ever been told by a health professional that respondent had a heart attack or angina

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 46a. Ever diagnosed with diabetes among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	649	614,100	9.6	(6.8 - 12.5)	38,174	1,382,700	9.3	(8.0 - 10.5)
Gender								
Male	251	181,300	6.1	(3.3 - 8.9)	14,626	530,400	7.5	(6.2 - 8.8)
Female	398	432,900	12.7	(8.2 - 17.3)	23,548	852,300	11.0	(8.9 - 13.0)
Age (years)								
18-34	*	*	*	*	3,941	150,200	3.4	(1.0 - 5.8)
35-44	*	*	*	*	4,987	165,100	5.7	(3.6 - 7.9)
45-54	124	151,600	13.3	(5.9 - 20.6)	7,830	279,000	9.8	(6.7 - 12.8)
55-64	160	125,900	14.9	(7.1 - 22.6)	8,771	332,900	15.5	(12.3 - 18.8)
≥ 65	188	165,000	16.3	(9.4 - 23.2)	12,645	455,600	17.9	(15.0 - 20.8)
Race/Ethnicity								
White Non-Hispanic	395	187,700	7.7	(4.9 - 10.5)	35,862	809,300	8.7	(7.8 - 9.5)
Black Non-Hispanic	94	179,700	12.1	(4.3 - 19.9)	739	261,000	12.1	(6.6 - 17.6)
Hispanic	116	190,500	11.6	(5.2 - 18.1)	862	227,300	10.1	(5.4 - 14.8)
Other Non-Hispanic	*	*	*	*	711	85,100	7.6	(2.3 - 12.8)
Education								
≤ High School	232	374,500	14.0	(8.7 - 19.3)	15,709	730,000	13.0	(10.5 - 15.6)
Some College	118	110,600	8.8	(1.6 - 15.9)	9,918	303,500	8.6	(5.9 - 11.2)
≥ College Degree	295	119,800	5.1	(2.5 - 7.6)	12,464	337,500	6.0	(4.8 - 7.2)
Income								
≤ \$24,999	169	314,100	15.9	(8.8 - 22.9)	9,652	543,400	15.5	(11.4 - 19.5)
\$25,000-\$49,999	106	103,800	8.9	(2.9 - 14.9)	9,667	291,000	10.0	(7.5 - 12.6)
\$50,000-\$74,999	*	*	*	*	5,777	151,000	7.6	(5.0 - 10.3)
≥ \$75,000	189	64,300	4.2	(1.2 - 7.2)	8,565	220,800	4.9	(3.8 - 6.1)

¹ Defined as having ever been told by a doctor that respondent had diabetes, excluding pre-diabetes and women with diabetes only when pregnant

² Sample size from survey

³ Estimated number of adults who have diabetes (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 46b. Ever diagnosed with diabetes among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	649	9.7	(6.8 - 12.5)	38,174	9.0	(7.8 - 10.3)
Gender						
Male	251	6.5	(3.5 - 9.5)	14,626	7.5	(6.1 - 8.9)
Female	398	12.4	(7.9 - 17.0)	23,548	10.4	(8.4 - 12.4)
Race/Ethnicity						
White Non-Hispanic	395	7.9	(5.1 - 10.8)	35,862	8.4	(7.5 - 9.2)
Black Non-Hispanic	94	11.6	(4.0 - 19.2)	739	11.4	(6.0 - 16.7)
Hispanic	116	11.9	(5.3 - 18.4)	862	10.1	(5.4 - 14.9)
Other Non-Hispanic	*	*	*	711	7.7	(2.3 - 13.0)
Education						
≤ High School	232	13.8	(8.6 - 19.1)	15,709	12.6	(10.0 - 15.2)
Some College	118	8.8	(1.6 - 16.0)	9,918	8.3	(5.6 - 10.9)
≥ College Degree	295	5.4	(2.6 - 8.1)	12,464	5.9	(4.7 - 7.2)
Income						
≤ \$24,999	169	16.0	(8.9 - 23.0)	9,652	15.2	(11.1 - 19.2)
\$25,000-\$49,999	106	8.8	(2.8 - 14.8)	9,667	9.7	(7.1 - 12.2)
\$50,000-\$74,999	*	*	*	5,777	7.6	(4.9 - 10.2)
≥ \$75,000	189	4.3	(1.2 - 7.4)	8,565	4.7	(3.6 - 5.9)

¹ Defined as having ever been told by a doctor that respondent had diabetes, excluding pre-diabetes and women with diabetes only when pregnant

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 47a. Ever diagnosed with asthma (lifetime) among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	646	954,100	15.0	(10.9 - 19.1)	38,096	2,225,100	15.0	(13.2 - 16.8)
Gender								
Male	249	348,300	11.8	(6.6 - 17.0)	14,601	875,800	12.4	(10.1 - 14.7)
Female	397	605,800	17.9	(11.8 - 24.0)	23,495	1,349,300	17.4	(14.6 - 20.1)
Age (years)								
18-34	75	331,700	16.1	(6.3 - 26.0)	3,931	825,100	18.5	(13.8 - 23.3)
35-44	100	194,100	15.0	(6.7 - 23.4)	4,972	431,700	15.1	(11.2 - 19.1)
45-54	123	172,700	15.2	(7.5 - 22.8)	7,819	365,900	12.8	(9.7 - 16.0)
55-64	160	150,900	17.8	(10.8 - 24.8)	8,756	316,700	14.8	(11.9 - 17.7)
≥ 65	188	104,700	10.3	(4.0 - 16.7)	12,618	285,700	11.2	(8.6 - 13.8)
Race/Ethnicity								
White Non-Hispanic	394	339,800	14.0	(8.0 - 19.9)	35,783	1,365,700	14.6	(13.0 - 16.3)
Black Non-Hispanic	*	*	*	*	737	367,000	17.2	(10.2 - 24.2)
Hispanic	115	201,500	12.3	(5.9 - 18.8)	863	317,100	14.1	(9.0 - 19.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	231	366,200	13.7	(7.9 - 19.5)	15,675	831,600	14.9	(12.0 - 17.8)
Some College	118	205,600	16.3	(7.1 - 25.4)	9,893	587,500	16.6	(13.1 - 20.1)
≥ College Degree	293	317,500	13.6	(7.3 - 19.9)	12,444	739,000	13.2	(10.4 - 15.9)
Income								
≤ \$24,999	168	437,900	22.2	(12.9 - 31.5)	9,643	708,900	20.2	(14.9 - 25.5)
\$25,000-\$49,999	106	127,100	10.9	(3.0 - 18.9)	9,641	389,800	13.4	(10.0 - 16.9)
\$50,000-\$74,999	*	*	*	*	5,762	257,800	13.2	(10.0 - 16.5)
≥ \$75,000	189	183,600	11.9	(5.5 - 18.4)	8,562	577,900	12.8	(10.4 - 15.3)

¹ Defined as adults having ever been told by a health professional that they had asthma

² Sample size from survey

³ Estimated number of adults who have lifetime asthma (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 47b. Ever diagnosed with asthma (lifetime) among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	646	15.0	(10.9 - 19.1)	38,096	15.2	(13.3 - 17.0)
Gender						
Male	249	11.8	(6.7 - 17.0)	14,601	12.6	(10.3 - 14.9)
Female	397	17.7	(11.6 - 23.8)	23,495	17.5	(14.8 - 20.3)
Race/Ethnicity						
White Non-Hispanic	394	13.7	(7.8 - 19.7)	35,783	14.9	(13.2 - 16.6)
Black Non-Hispanic	93	18.2	(8.2 - 28.2)	737	17.1	(10.1 - 24.1)
Hispanic	115	12.7	(6.1 - 19.4)	863	14.5	(9.3 - 19.7)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	231	13.8	(7.9 - 19.6)	15,675	15.1	(12.1 - 18.0)
Some College	118	16.3	(7.1 - 25.5)	9,893	16.9	(13.4 - 20.4)
≥ College Degree	293	13.4	(7.2 - 19.6)	12,444	13.3	(10.6 - 16.0)
Income						
≤ \$24,999	168	22.2	(13.0 - 31.4)	9,643	20.3	(15.0 - 25.6)
\$25,000-\$49,999	106	10.8	(2.8 - 18.7)	9,641	13.7	(10.2 - 17.2)
\$50,000-\$74,999	*	*	*	5,762	13.5	(10.0 - 17.0)
≥ \$75,000	189	11.7	(5.3 - 18.1)	8,562	13.0	(10.6 - 15.4)

¹ Defined as adults having ever been told by a health professional that they had asthma

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 48a. Current asthma among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	644	586,700	9.3	(6.0 - 12.6)	37,933	1,415,000	9.6	(8.1 - 11.0)
Gender								
Male	248	161,300	5.5	(2.3 - 8.6)	14,546	440,700	6.3	(4.8 - 7.7)
Female	396	425,400	12.6	(7.2 - 17.9)	23,387	974,300	12.6	(10.2 - 15.0)
Age (years)								
18-34	*	*	*	*	3,909	467,200	10.5	(6.8 - 14.3)
35-44	*	*	*	*	4,947	246,800	8.7	(5.8 - 11.6)
45-54	123	101,000	8.9	(3.2 - 14.5)	7,787	240,000	8.4	(6.1 - 10.8)
55-64	160	123,000	14.5	(8.0 - 21.0)	8,726	231,100	10.8	(8.1 - 13.5)
≥ 65	187	92,300	9.1	(2.9 - 15.3)	12,564	229,900	9.1	(6.5 - 11.6)
Race/Ethnicity								
White Non-Hispanic	393	218,600	9.0	(3.9 - 14.0)	35,629	888,400	9.6	(8.2 - 11.0)
Black Non-Hispanic	93	185,300	12.7	(4.2 - 21.1)	736	246,900	11.6	(5.7 - 17.5)
Hispanic	114	117,000	7.2	(2.1 - 12.3)	859	186,800	8.3	(4.4 - 12.2)
Other Non-Hispanic	*	*	*	*	709	93,000	8.3	(1.4 - 15.2)
Education								
≤ High School	231	262,100	9.8	(4.8 - 14.8)	15,594	595,100	10.7	(8.2 - 13.2)
Some College	117	97,600	7.7	(2.3 - 13.2)	9,861	322,000	9.1	(7.0 - 11.3)
≥ College Degree	292	162,300	7.0	(2.5 - 11.5)	12,394	430,900	7.7	(5.8 - 9.7)
Income								
≤ \$24,999	168	279,500	14.2	(6.6 - 21.7)	9,600	489,200	14.0	(9.7 - 18.3)
\$25,000-\$49,999	106	99,600	8.6	(1.1 - 16.0)	9,590	268,400	9.3	(6.2 - 12.4)
\$50,000-\$74,999	*	*	*	*	5,741	143,400	7.4	(5.1 - 9.6)
≥ \$75,000	188	95,500	6.3	(1.5 - 11.0)	8,541	339,200	7.6	(5.8 - 9.4)

¹ Defined as adults having ever been told by a health professional that they had asthma and still have asthma

² Sample size from survey

³ Estimated number of adults who currently have asthma (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 48b. Current asthma among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	644	9.2	(5.9 - 12.5)	37,933	9.7	(8.2 - 11.2)
Gender						
Male	248	5.6	(2.4 - 8.8)	14,546	6.4	(4.9 - 7.9)
Female	396	12.3	(7.0 - 17.7)	23,387	12.7	(10.2 - 15.1)
Race/Ethnicity						
White Non-Hispanic	393	8.7	(3.7 - 13.8)	35,629	9.7	(8.3 - 11.2)
Black Non-Hispanic	93	12.5	(4.1 - 20.9)	736	11.5	(5.6 - 17.3)
Hispanic	114	7.4	(2.2 - 12.7)	859	8.6	(4.5 - 12.6)
Other Non-Hispanic	*	*	*	709	8.3	(1.3 - 15.3)
Education						
≤ High School	231	9.9	(4.8 - 14.9)	15,594	10.8	(8.3 - 13.3)
Some College	117	7.7	(2.2 - 13.2)	9,861	9.4	(7.2 - 11.6)
≥ College Degree	292	6.7	(2.3 - 11.2)	12,394	7.8	(5.8 - 9.7)
Income						
≤ \$24,999	168	14.3	(6.8 - 21.9)	9,600	14.1	(9.7 - 18.4)
\$25,000-\$49,999	106	8.4	(1.0 - 15.8)	9,590	9.5	(6.3 - 12.7)
\$50,000-\$74,999	*	*	*	5,741	7.5	(5.0 - 10.1)
≥ \$75,000	188	5.9	(1.3 - 10.5)	8,541	7.7	(5.9 - 9.5)

¹ Defined as adults having ever been told by a health professional that they had asthma and still have asthma

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 49a. Chronic joint symptoms among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	648	2,518,300	39.5	(34.2 - 44.9)	38,099	6,377,100	42.9	(40.5 - 45.3)
Gender								
Male	250	1,220,800	41.0	(32.3 - 49.8)	14,595	3,033,600	42.8	(39.0 - 46.6)
Female	398	1,297,400	38.2	(31.7 - 44.7)	23,504	3,343,500	43.0	(40.0 - 46.0)
Age (years)								
18-34	*	*	*	*	3,933	1,261,300	28.3	(22.6 - 34.0)
35-44	*	*	*	*	4,984	1,110,900	38.6	(33.1 - 44.0)
45-54	*	*	*	*	7,817	1,456,500	51.2	(46.4 - 55.9)
55-64	160	423,300	50.0	(40.2 - 59.7)	8,751	1,167,400	54.6	(50.4 - 58.8)
≥ 65	188	499,500	49.3	(40.1 - 58.5)	12,614	1,381,000	54.3	(50.5 - 58.1)
Race/Ethnicity								
White Non-Hispanic	394	1,046,400	43.0	(36.4 - 49.6)	35,785	4,267,600	45.8	(43.8 - 47.7)
Black Non-Hispanic	*	*	*	*	740	890,500	41.3	(32.5 - 50.0)
Hispanic	*	*	*	*	860	827,500	36.7	(27.7 - 45.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	232	1,026,600	38.4	(29.8 - 47.1)	15,676	2,450,000	43.8	(39.4 - 48.1)
Some College	*	*	*	*	9,900	1,598,300	45.1	(40.1 - 50.1)
≥ College Degree	294	965,600	40.9	(33.3 - 48.5)	12,440	2,313,100	41.1	(37.8 - 44.5)
Income								
≤ \$24,999	169	692,100	35.0	(25.2 - 44.7)	9,641	1,483,000	42.2	(36.3 - 48.1)
\$25,000-\$49,999	*	*	*	*	9,647	1,332,100	45.9	(39.8 - 52.1)
\$50,000-\$74,999	*	*	*	*	5,768	894,200	45.3	(39.5 - 51.1)
≥ \$75,000	188	691,400	45.2	(36.2 - 54.1)	8,549	1,874,200	41.7	(38.4 - 45.1)

¹ Defined as adults having symptoms of pain, aching or stiffness in or around a joint during the past 30 days and began more than 3 months ago

² Sample size from survey

³ Estimated number of adults who have chronic joint symptoms (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 49b. Chronic joint symptoms among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	648	39.5	(34.3 - 44.8)	38,099	42.1	(39.7 - 44.5)
Gender						
Male	250	41.5	(33.0 - 50.0)	14,595	42.6	(38.8 - 46.3)
Female	398	37.9	(31.3 - 44.4)	23,504	41.6	(38.6 - 44.7)
Race/Ethnicity						
White Non-Hispanic	394	42.9	(36.3 - 49.5)	35,785	44.7	(42.7 - 46.6)
Black Non-Hispanic	*	*	*	740	40.2	(31.5 - 48.9)
Hispanic	*	*	*	860	36.6	(27.8 - 45.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	232	38.5	(29.9 - 47.0)	15,676	43.1	(38.8 - 47.5)
Some College	*	*	*	9,900	43.9	(39.0 - 48.8)
≥ College Degree	294	41.3	(33.7 - 48.9)	12,440	40.3	(37.0 - 43.7)
Income						
≤ \$24,999	169	34.9	(25.3 - 44.6)	9,641	41.6	(35.7 - 47.4)
\$25,000-\$49,999	*	*	*	9,647	45.0	(39.0 - 51.0)
\$50,000-\$74,999	*	*	*	5,768	44.3	(38.4 - 50.2)
≥ \$75,000	188	45.3	(36.2 - 54.3)	8,549	40.8	(37.4 - 44.2)

¹ Defined as adults having symptoms of pain, aching or stiffness in or around a joint during the past 30 days and began more than 3 months ago

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 50a. Arthritis among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	642	1,467,800	23.2	(19.3 - 27.1)	38,013	4,102,800	27.7	(25.9 - 29.6)
Gender								
Male	249	545,200	18.5	(13.1 - 23.8)	14,555	1,642,700	23.3	(20.7 - 25.8)
Female	393	922,600	27.3	(21.8 - 32.8)	23,458	2,460,100	31.8	(29.2 - 34.4)
Age (years)								
18-34	*	*	*	*	3,927	312,100	7.0	(4.9 - 9.1)
35-44	101	116,000	9.0	(2.6 - 15.3)	4,972	412,500	14.5	(11.3 - 17.6)
45-54	122	317,500	28.0	(18.4 - 37.7)	7,789	869,100	30.7	(26.6 - 34.8)
55-64	157	371,700	44.4	(34.5 - 54.2)	8,729	999,300	47.0	(42.9 - 51.1)
≥ 65	187	562,700	55.7	(46.7 - 64.7)	12,596	1,509,800	59.6	(55.8 - 63.3)
Race/Ethnicity								
White Non-Hispanic	391	669,900	27.6	(22.2 - 33.0)	35,710	2,967,700	31.9	(30.3 - 33.6)
Black Non-Hispanic	*	*	*	*	737	588,100	27.3	(20.2 - 34.4)
Hispanic	113	296,600	18.5	(11.1 - 25.8)	856	380,700	17.1	(11.8 - 22.5)
Other Non-Hispanic	*	*	*	*	710	166,300	14.8	(7.8 - 21.8)
Education								
≤ High School	229	679,400	25.8	(19.1 - 32.4)	15,652	1,742,700	31.3	(27.8 - 34.8)
Some College	118	189,000	14.9	(8.3 - 21.6)	9,874	902,000	25.5	(22.3 - 28.6)
≥ College Degree	291	585,500	24.9	(18.6 - 31.2)	12,407	1,437,800	25.6	(22.9 - 28.4)
Income								
≤ \$24,999	165	492,500	25.3	(17.5 - 33.1)	9,618	1,112,500	32.0	(27.1 - 36.9)
\$25,000-\$49,999	104	210,600	18.2	(9.8 - 26.7)	9,624	821,900	28.4	(24.2 - 32.6)
\$50,000-\$74,999	*	*	*	*	5,753	542,500	27.5	(23.2 - 31.9)
≥ \$75,000	189	352,100	22.9	(15.5 - 30.4)	8,536	1,012,800	22.6	(19.8 - 25.3)

¹ Defined as adults having ever been told by a health professional that they have some form of arthritis, rheumatoid arthritis, gout, lupus or fibromyalgia (excluding back or neck)

² Sample size from survey

³ Estimated number of adults who have ever been told by a health professional they have some form of arthritis (excluding back or neck) (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 50b. Arthritis among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	642	23.3	(19.4 - 27.3)	38,013	26.6	(24.8 - 28.5)
Gender						
Male	249	19.9	(14.2 - 25.5)	14,555	23.5	(20.9 - 26.1)
Female	393	26.4	(20.9 - 31.8)	23,458	29.5	(27.0 - 32.1)
Race/Ethnicity						
White Non-Hispanic	391	27.6	(22.2 - 33.1)	35,710	30.4	(28.7 - 32.0)
Black Non-Hispanic	*	*	*	737	26.5	(19.3 - 33.6)
Hispanic	113	19.0	(11.5 - 26.5)	856	17.2	(11.8 - 22.6)
Other Non-Hispanic	*	*	*	710	14.9	(7.7 - 22.0)
Education						
≤ High School	229	25.8	(19.1 - 32.5)	15,652	30.4	(26.9 - 33.8)
Some College	118	14.4	(7.9 - 20.8)	9,874	24.1	(21.0 - 27.1)
≥ College Degree	291	25.4	(19.0 - 31.8)	12,407	24.7	(21.9 - 27.5)
Income						
≤ \$24,999	165	25.5	(17.6 - 33.4)	9,618	31.2	(26.3 - 36.1)
\$25,000-\$49,999	104	18.2	(9.8 - 26.5)	9,624	27.3	(23.2 - 31.3)
\$50,000-\$74,999	*	*	*	5,753	26.3	(21.9 - 30.7)
≥ \$75,000	189	23.1	(15.6 - 30.6)	8,536	21.6	(18.8 - 24.3)

¹ Defined as adults having ever been told by a health professional that they have some form of arthritis, rheumatoid arthritis, gout, lupus or fibromyalgia (excluding back or neck)

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 51a. Activity limitations because of physical, mental or emotional problems among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	645	1,224,700	19.3	(15.0 - 23.6)	37,998	3,053,000	20.6	(18.7 - 22.5)
Gender								
Male	250	491,200	16.5	(10.0 - 23.0)	14,566	1,318,400	18.6	(15.7 - 21.5)
Female	395	733,500	21.7	(16.0 - 27.4)	23,432	1,734,600	22.4	(19.8 - 25.0)
Age (years)								
18-34	*	*	*	*	3,929	580,000	13.0	(8.2 - 17.9)
35-44	102	160,800	12.2	(4.8 - 19.6)	4,973	418,600	14.5	(11.0 - 18.1)
45-54	124	247,300	21.6	(13.6 - 29.7)	7,812	620,200	21.7	(18.3 - 25.2)
55-64	159	276,400	32.7	(23.2 - 42.2)	8,739	651,600	30.5	(26.5 - 34.6)
≥ 65	185	267,700	27.0	(18.7 - 35.2)	12,545	782,700	31.1	(27.7 - 34.6)
Race/Ethnicity								
White Non-Hispanic	392	479,400	19.7	(14.2 - 25.2)	35,688	2,037,200	21.9	(20.3 - 23.5)
Black Non-Hispanic	93	293,600	20.0	(10.7 - 29.2)	735	429,700	20.1	(13.5 - 26.7)
Hispanic	*	*	*	*	863	448,200	19.8	(11.9 - 27.8)
Other Non-Hispanic	*	*	*	*	712	137,900	12.3	(5.2 - 19.3)
Education								
≤ High School	231	609,400	22.9	(15.3 - 30.6)	15,626	1,359,600	24.4	(20.6 - 28.1)
Some College	116	162,000	12.9	(4.7 - 21.0)	9,875	700,800	19.8	(16.5 - 23.2)
≥ College Degree	294	388,500	16.4	(11.5 - 21.4)	12,415	924,800	16.5	(14.2 - 18.7)
Income								
≤ \$24,999	169	517,300	26.1	(16.9 - 35.4)	9,603	1,055,800	30.1	(24.7 - 35.5)
\$25,000-\$49,999	*	*	*	*	9,628	630,900	21.8	(16.6 - 27.0)
\$50,000-\$74,999	*	*	*	*	5,760	408,100	20.7	(16.1 - 25.2)
≥ \$75,000	188	185,300	12.1	(7.3 - 16.8)	8,544	583,300	13.0	(11.1 - 14.9)

¹ Defined as activity limitations because of physical, mental or emotional problems

² Sample size from survey

³ Estimated number of adults who are limited in any activities because of physical, mental or emotional problems (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 51b. Activity limitations because of physical, mental or emotional problems among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	645	19.1	(14.9 - 23.4)	37,998	20.1	(18.2 - 22.0)
Gender						
Male	250	16.8	(10.4 - 23.1)	14,566	18.6	(15.8 - 21.4)
Female	395	21.2	(15.5 - 26.9)	23,432	21.5	(18.9 - 24.1)
Race/Ethnicity						
White Non-Hispanic	392	19.6	(14.1 - 25.1)	35,688	21.3	(19.7 - 22.9)
Black Non-Hispanic	93	20.0	(10.7 - 29.3)	735	19.9	(13.2 - 26.5)
Hispanic	*	*	*	863	19.1	(11.5 - 26.8)
Other Non-Hispanic	*	*	*	712	12.4	(5.3 - 19.6)
Education						
≤ High School	231	22.5	(15.1 - 29.9)	15,626	23.8	(20.1 - 27.4)
Some College	116	12.5	(4.5 - 20.6)	9,875	19.3	(16.0 - 22.6)
≥ College Degree	294	16.7	(11.7 - 21.7)	12,415	16.1	(13.9 - 18.3)
Income						
≤ \$24,999	169	26.2	(16.9 - 35.4)	9,603	29.7	(24.3 - 35.2)
\$25,000-\$49,999	*	*	*	9,628	21.2	(16.1 - 26.3)
\$50,000-\$74,999	*	*	*	5,760	20.1	(15.6 - 24.6)
≥ \$75,000	188	12.4	(7.4 - 17.3)	8,544	12.7	(10.8 - 14.6)

¹ Defined as activity limitations because of physical, mental or emotional problems

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 52a. Have health problem that requires the use of special equipment among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	647	506,800	8.0	(5.7 - 10.2)	38,171	1,169,400	7.9	(6.9 - 8.9)
Gender								
Male	249	163,200	5.5	(3.0 - 8.0)	14,617	460,700	6.5	(5.3 - 7.7)
Female	398	343,600	10.1	(6.7 - 13.6)	23,554	708,700	9.1	(7.6 - 10.7)
Age (years)								
18-34	*	*	*	*	3,940	68,800	1.5	(0.7 - 2.4)
35-44	*	*	*	*	4,989	122,500	4.3	(2.4 - 6.1)
45-54	124	68,800	6.0	(2.2 - 9.8)	7,836	182,500	6.4	(4.7 - 8.1)
55-64	159	90,700	10.8	(4.9 - 16.7)	8,773	219,600	10.3	(7.8 - 12.8)
≥ 65	187	283,000	28.0	(19.3 - 36.8)	12,633	576,100	22.7	(19.0 - 26.3)
Race/Ethnicity								
White Non-Hispanic	393	255,000	10.5	(7.1 - 13.9)	35,853	802,000	8.6	(7.6 - 9.6)
Black Non-Hispanic	94	168,200	11.3	(4.9 - 17.8)	739	235,800	10.9	(6.4 - 15.5)
Hispanic	116	75,200	4.6	(1.3 - 7.8)	864	104,100	4.6	(2.1 - 7.1)
Other Non-Hispanic	*	*	*	*	715	27,600	2.5	(0.8 - 4.1)
Education								
≤ High School	232	288,400	10.8	(6.6 - 15.0)	15,711	604,200	10.8	(8.7 - 12.8)
Some College	118	82,900	6.6	(2.0 - 11.2)	9,914	260,000	7.3	(5.5 - 9.1)
≥ College Degree	293	130,800	5.6	(2.9 - 8.2)	12,463	298,900	5.3	(4.1 - 6.5)
Income								
≤ \$24,999	169	191,800	9.7	(5.4 - 13.9)	9,656	430,500	12.2	(9.6 - 14.8)
\$25,000-\$49,999	106	93,600	8.0	(2.5 - 13.6)	9,663	238,900	8.2	(5.9 - 10.5)
\$50,000-\$74,999	*	*	*	*	5,775	103,000	5.2	(3.6 - 6.8)
≥ \$75,000	*	*	*	*	8,566	136,000	3.0	(2.1 - 4.0)

¹ Defined as having any health problem that requires the use of special equipment such as a cane, a wheelchair, a special bed, or a special telephone

² Sample size from survey

³ Estimated number of adults who have any health problem that requires the use of special equipment (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 52b. Have health problem that requires the use of special equipment among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	647	8.0	(5.8 - 10.2)	38,171	7.6	(6.6 - 8.6)
Gender						
Male	249	6.1	(3.3 - 8.8)	14,617	6.8	(5.5 - 8.0)
Female	398	9.6	(6.3 - 12.9)	23,554	8.4	(6.9 - 9.9)
Race/Ethnicity						
White Non-Hispanic	393	10.8	(7.3 - 14.2)	35,853	8.3	(7.3 - 9.3)
Black Non-Hispanic	94	11.1	(4.8 - 17.5)	739	10.5	(6.1 - 15.0)
Hispanic	116	4.5	(1.3 - 7.7)	864	4.5	(2.1 - 6.9)
Other Non-Hispanic	*	*	*	715	2.2	(0.7 - 3.8)
Education						
≤ High School	232	10.4	(6.4 - 14.4)	15,711	10.3	(8.3 - 12.3)
Some College	118	6.6	(1.9 - 11.3)	9,914	7.1	(5.2 - 8.9)
≥ College Degree	293	6.0	(3.1 - 8.8)	12,463	5.3	(4.1 - 6.6)
Income						
≤ \$24,999	169	9.8	(5.6 - 14.1)	9,656	11.9	(9.4 - 14.5)
\$25,000-\$49,999	106	7.7	(2.4 - 13.0)	9,663	7.7	(5.5 - 10.0)
\$50,000-\$74,999	*	*	*	5,775	5.1	(3.4 - 6.8)
≥ \$75,000	*	*	*	8,566	2.9	(2.0 - 3.9)

¹ Defined as having any health problem that requires the use of special equipment such as a cane, a wheelchair, a special bed, or a special telephone

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 53a. Disability among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	646	1,414,800	22.2	(17.8 - 26.7)	38,016	3,394,200	22.9	(20.9 - 24.9)
Gender								
Male	250	563,700	18.9	(12.3 - 25.6)	14,568	1,469,700	20.7	(17.8 - 23.7)
Female	396	851,100	25.1	(19.2 - 31.0)	23,448	1,924,600	24.8	(22.1 - 27.5)
Age (years)								
18-34	*	*	*	*	3,931	590,300	13.3	(8.4 - 18.1)
35-44	102	176,300	13.4	(5.7 - 21.0)	4,974	447,900	15.6	(11.9 - 19.3)
45-54	124	252,800	22.1	(14.1 - 30.2)	7,811	643,900	22.6	(19.1 - 26.0)
55-64	158	305,600	36.4	(26.7 - 46.0)	8,740	702,100	33.0	(28.9 - 37.0)
≥ 65	187	407,600	40.3	(31.2 - 49.5)	12,560	1,010,000	39.9	(36.1 - 43.7)
Race/Ethnicity								
White Non-Hispanic	392	566,800	23.3	(17.6 - 29.1)	35,703	2,253,400	24.2	(22.6 - 25.9)
Black Non-Hispanic	94	364,900	24.6	(14.7 - 34.4)	736	511,700	23.8	(16.8 - 30.8)
Hispanic	*	*	*	*	864	484,300	21.4	(13.4 - 29.5)
Other Non-Hispanic	*	*	*	*	713	144,800	12.9	(5.8 - 19.9)
Education								
≤ High School	232	718,900	26.9	(19.1 - 34.8)	15,638	1,544,200	27.6	(23.7 - 31.4)
Some College	117	185,900	14.7	(6.3 - 23.2)	9,876	755,200	21.4	(17.9 - 24.8)
≥ College Degree	293	440,500	18.7	(13.4 - 24.0)	12,419	1,021,300	18.2	(15.9 - 20.5)
Income								
≤ \$24,999	169	597,300	30.2	(20.6 - 39.8)	9,613	1,179,100	33.6	(28.0 - 39.2)
\$25,000-\$49,999	*	*	*	*	9,629	699,000	24.2	(18.9 - 29.4)
\$50,000-\$74,999	*	*	*	*	5,757	430,100	21.8	(17.2 - 26.4)
≥ \$75,000	188	209,200	13.6	(8.5 - 18.8)	8,546	632,700	14.1	(12.1 - 16.1)

¹ Defined as having activity limitations due to physical, mental or emotional problems or have health problems that require the use of special equipment

² Sample size from survey

³ Estimated number of adults with a disability (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 53b. Disability among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	646	22.1	(17.7 - 26.5)	38,016	22.3	(20.4 - 24.3)
Gender						
Male	250	19.6	(13.1 - 26.2)	14,568	20.9	(18.0 - 23.8)
Female	396	24.3	(18.4 - 30.2)	23,448	23.6	(21.0 - 26.3)
Race/Ethnicity						
White Non-Hispanic	392	23.5	(17.8 - 29.3)	35,703	23.6	(21.9 - 25.3)
Black Non-Hispanic	94	24.5	(14.7 - 34.4)	736	23.5	(16.5 - 30.5)
Hispanic	*	*	*	864	20.6	(12.9 - 28.3)
Other Non-Hispanic	*	*	*	713	12.9	(5.8 - 20.1)
Education						
≤ High School	232	26.2	(18.6 - 33.8)	15,638	26.8	(23.0 - 30.6)
Some College	117	14.7	(6.3 - 23.2)	9,876	20.9	(17.4 - 24.3)
≥ College Degree	293	19.2	(13.8 - 24.6)	12,419	17.9	(15.5 - 20.3)
Income						
≤ \$24,999	169	30.2	(20.6 - 39.7)	9,613	33.2	(27.6 - 38.8)
\$25,000-\$49,999	*	*	*	9,629	23.5	(18.3 - 28.6)
\$50,000-\$74,999	*	*	*	5,757	21.2	(16.6 - 25.8)
≥ \$75,000	188	13.9	(8.6 - 19.2)	8,546	13.8	(11.7 - 15.8)

¹ Defined as having activity limitations due to physical, mental or emotional problems or have health problems that require the use of special equipment

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 54a. Current smoking among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	643	912,700	14.5	(10.7 - 18.4)	38,029	2,483,500	16.8	(15.1 - 18.6)
Gender								
Male	248	552,400	18.8	(12.1 - 25.5)	14,559	1,356,400	19.3	(16.3 - 22.2)
Female	395	360,300	10.8	(6.6 - 15.0)	23,470	1,127,100	14.6	(12.7 - 16.6)
Age (years)								
18-34	75	388,400	18.9	(9.1 - 28.7)	3,929	993,500	22.3	(17.5 - 27.1)
35-44	*	*	*	*	4,977	421,100	14.8	(11.6 - 17.9)
45-54	121	158,200	14.7	(7.8 - 21.6)	7,807	508,800	18.3	(15.4 - 21.3)
55-64	160	120,100	14.2	(8.2 - 20.1)	8,744	314,500	14.7	(12.1 - 17.3)
≥ 65	186	135,000	13.4	(7.2 - 19.7)	12,572	245,600	9.7	(7.2 - 12.3)
Race/Ethnicity								
White Non-Hispanic	393	315,300	13.0	(8.1 - 17.9)	35,729	1,615,500	17.3	(15.9 - 18.8)
Black Non-Hispanic	92	158,500	11.1	(4.9 - 17.4)	735	284,100	13.6	(8.9 - 18.2)
Hispanic	114	250,600	15.5	(7.3 - 23.7)	858	350,900	15.8	(9.6 - 21.9)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	230	451,500	17.3	(11.0 - 23.6)	15,654	1,237,300	22.4	(19.1 - 25.6)
Some College	116	192,400	15.6	(5.9 - 25.2)	9,875	670,600	19.1	(15.5 - 22.8)
≥ College Degree	293	259,500	11.0	(5.4 - 16.7)	12,421	563,300	10.0	(7.5 - 12.5)
Income								
≤ \$24,999	168	397,100	20.6	(11.9 - 29.3)	9,626	835,100	24.1	(19.1 - 29.1)
\$25,000-\$49,999	103	150,500	13.3	(4.7 - 21.8)	9,629	558,100	19.4	(15.6 - 23.2)
\$50,000-\$74,999	*	*	*	*	5,756	338,100	17.3	(13.0 - 21.7)
≥ \$75,000	188	114,900	7.5	(3.1 - 11.9)	8,549	468,700	10.4	(8.6 - 12.3)

¹ Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day or some days

² Sample size from survey

³ Estimated number of adult current smokers (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 54b. Current smoking among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	643	14.5	(10.7 - 18.3)	38,029	17.0	(15.3 - 18.8)
Gender						
Male	248	18.8	(12.3 - 25.3)	14,559	19.2	(16.3 - 22.1)
Female	395	10.8	(6.6 - 15.0)	23,470	15.0	(13.0 - 17.0)
Race/Ethnicity						
White Non-Hispanic	393	13.0	(8.2 - 17.7)	35,729	17.7	(16.2 - 19.2)
Black Non-Hispanic	92	11.2	(4.8 - 17.5)	735	13.5	(8.8 - 18.2)
Hispanic	114	15.7	(7.6 - 23.9)	858	15.9	(9.8 - 22.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	230	17.6	(11.4 - 23.9)	15,654	22.8	(19.5 - 26.0)
Some College	116	15.3	(6.0 - 24.6)	9,875	19.4	(15.8 - 22.9)
≥ College Degree	293	10.8	(5.3 - 16.3)	12,421	10.0	(7.6 - 12.5)
Income						
≤ \$24,999	168	20.2	(11.8 - 28.6)	9,626	24.1	(19.2 - 29.1)
\$25,000-\$49,999	103	13.8	(5.1 - 22.5)	9,629	19.9	(16.0 - 23.7)
\$50,000-\$74,999	*	*	*	5,756	17.8	(13.3 - 22.3)
≥ \$75,000	188	7.6	(3.1 - 12.1)	8,549	10.6	(8.6 - 12.5)

¹ Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day or some days

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 55a. Everyday smoking among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	643	482,600	7.7	(5.2 - 10.2)	38,029	1,626,800	11.0	(9.8 - 12.2)
Gender								
Male	248	283,700	9.7	(5.3 - 14.0)	14,559	848,400	12.0	(10.0 - 14.1)
Female	395	198,900	6.0	(3.1 - 8.8)	23,470	778,400	10.1	(8.7 - 11.5)
Age (years)								
18-34	*	*	*	*	3,929	559,800	12.6	(9.6 - 15.6)
35-44	*	*	*	*	4,977	315,200	11.0	(8.4 - 13.7)
45-54	121	100,300	9.3	(3.7 - 14.9)	7,807	377,000	13.6	(11.1 - 16.0)
55-64	160	67,400	8.0	(3.3 - 12.6)	8,744	203,000	9.5	(7.5 - 11.5)
≥ 65	186	94,000	9.4	(4.4 - 14.3)	12,572	171,700	6.8	(4.8 - 8.8)
Race/Ethnicity								
White Non-Hispanic	393	201,500	8.3	(4.1 - 12.4)	35,729	1,158,100	12.4	(11.2 - 13.7)
Black Non-Hispanic	*	*	*	*	735	197,700	9.4	(5.6 - 13.3)
Hispanic	114	108,800	6.7	(2.0 - 11.5)	858	176,200	7.9	(4.2 - 11.6)
Other Non-Hispanic	*	*	*	*	707	94,900	8.5	(3.3 - 13.7)
Education								
≤ High School	230	269,200	10.3	(5.6 - 15.1)	15,654	881,900	15.9	(13.4 - 18.5)
Some College	*	*	*	*	9,875	410,200	11.7	(9.6 - 13.8)
≥ College Degree	293	138,400	5.9	(2.6 - 9.2)	12,421	323,500	5.8	(4.3 - 7.2)
Income								
≤ \$24,999	168	131,600	6.8	(2.8 - 10.9)	9,626	452,800	13.1	(10.3 - 15.8)
\$25,000-\$49,999	103	105,800	9.3	(3.4 - 15.3)	9,629	425,900	14.8	(11.9 - 17.7)
\$50,000-\$74,999	*	*	*	*	5,756	222,700	11.4	(8.5 - 14.4)
≥ \$75,000	188	91,500	6.0	(1.8 - 10.1)	8,549	320,200	7.1	(5.5 - 8.8)

¹ Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day

² Sample size from survey

³ Estimated number of adults who have smoked at least 100 cigarettes in lifetime and currently smoking everyday (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 55b. Everyday smoking among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	643	7.9	(5.4 - 10.4)	38,029	11.3	(10.0 - 12.5)
Gender						
Male	248	9.9	(5.6 - 14.2)	14,559	12.1	(10.1 - 14.1)
Female	395	6.1	(3.2 - 9.0)	23,470	10.5	(9.0 - 12.0)
Race/Ethnicity						
White Non-Hispanic	393	8.5	(4.4 - 12.6)	35,729	12.8	(11.5 - 14.1)
Black Non-Hispanic	*	*	*	735	9.4	(5.5 - 13.4)
Hispanic	114	6.8	(2.1 - 11.6)	858	8.0	(4.3 - 11.7)
Other Non-Hispanic	*	*	*	707	8.5	(3.2 - 13.8)
Education						
≤ High School	230	10.7	(6.0 - 15.4)	15,654	16.4	(13.8 - 19.0)
Some College	*	*	*	9,875	12.1	(9.9 - 14.3)
≥ College Degree	293	5.9	(2.6 - 9.3)	12,421	5.8	(4.3 - 7.3)
Income						
≤ \$24,999	168	7.1	(2.9 - 11.3)	9,626	13.4	(10.6 - 16.2)
\$25,000-\$49,999	103	9.8	(3.6 - 16.0)	9,629	15.3	(12.3 - 18.3)
\$50,000-\$74,999	*	*	*	5,756	11.7	(8.7 - 14.8)
≥ \$75,000	188	6.2	(1.9 - 10.5)	8,549	7.2	(5.5 - 8.9)

¹ Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 56a. Adults living in homes in which smoking is prohibited¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	646	5,286,900	83.1	(79.3 - 87.0)	38,045	11,997,100	80.8	(79.0 - 82.6)
Gender								
Male	251	2,445,500	82.0	(75.9 - 88.2)	14,577	5,634,400	79.4	(76.6 - 82.2)
Female	395	2,841,400	84.0	(79.3 - 88.8)	23,468	6,362,800	82.1	(79.9 - 84.3)
Age (years)								
18-34	74	1,779,100	86.9	(78.2 - 95.6)	3,934	3,655,200	82.1	(77.8 - 86.4)
35-44	102	1,164,800	88.3	(80.8 - 95.8)	4,985	2,479,700	86.1	(82.5 - 89.7)
45-54	123	870,300	76.4	(67.9 - 85.0)	7,808	2,194,400	77.2	(73.5 - 80.8)
55-64	160	710,000	83.8	(77.7 - 89.9)	8,753	1,732,500	81.0	(78.3 - 83.7)
≥ 65	187	762,600	75.6	(68.1 - 83.0)	12,565	1,935,400	76.5	(73.4 - 79.5)
Race/Ethnicity								
White Non-Hispanic	392	1,963,600	81.0	(76.0 - 86.0)	35,737	7,426,900	79.8	(78.3 - 81.2)
Black Non-Hispanic	94	1,179,400	79.4	(70.0 - 88.8)	738	1,692,100	78.6	(71.9 - 85.3)
Hispanic	116	1,509,100	92.0	(86.4 - 97.5)	863	2,003,900	88.7	(84.1 - 93.3)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	231	2,260,400	84.9	(79.1 - 90.7)	15,647	4,307,200	77.1	(74.0 - 80.2)
Some College	*	*	*	*	9,883	2,795,400	78.9	(75.0 - 82.8)
≥ College Degree	293	1,963,000	83.3	(77.7 - 88.9)	12,437	4,817,300	85.7	(83.3 - 88.2)
Income								
≤ \$24,999	169	1,590,800	80.4	(72.2 - 88.5)	9,608	2,627,700	74.9	(70.0 - 79.7)
\$25,000-\$49,999	106	1,004,100	86.3	(78.1 - 94.6)	9,640	2,314,500	79.8	(76.0 - 83.5)
\$50,000-\$74,999	*	*	*	*	5,766	1,551,300	78.7	(74.0 - 83.4)
≥ \$75,000	188	1,335,700	87.2	(82.0 - 92.4)	8,556	3,883,200	86.5	(84.3 - 88.7)

¹ Defined as having a rule that prohibits smoking anywhere inside the home

² Sample size from survey

³ Estimated number of adults who live in a home with a rule that prohibits smoking anywhere inside the home (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 56b. Adults living in homes in which smoking is prohibited¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	646	83.1	(79.3 - 86.9)	38,045	80.9	(79.2 - 82.7)
Gender						
Male	251	81.8	(75.8 - 87.9)	14,577	79.4	(76.7 - 82.2)
Female	395	84.2	(79.4 - 89.0)	23,468	82.3	(80.0 - 84.5)
Race/Ethnicity						
White Non-Hispanic	392	81.0	(76.0 - 85.9)	35,737	79.9	(78.4 - 81.4)
Black Non-Hispanic	94	78.9	(69.4 - 88.5)	738	78.6	(71.8 - 85.4)
Hispanic	116	92.3	(87.0 - 97.6)	863	88.9	(84.4 - 93.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	231	84.6	(78.9 - 90.4)	15,647	76.9	(73.8 - 80.0)
Some College	*	*	*	9,883	79.1	(75.2 - 83.0)
≥ College Degree	293	83.3	(77.6 - 88.9)	12,437	86.0	(83.5 - 88.5)
Income						
≤ \$24,999	169	80.2	(72.1 - 88.3)	9,608	74.9	(70.1 - 79.7)
\$25,000-\$49,999	106	86.1	(77.8 - 94.5)	9,640	79.8	(76.1 - 83.6)
\$50,000-\$74,999	*	*	*	5,766	78.6	(73.8 - 83.3)
≥ \$75,000	188	87.2	(81.9 - 92.5)	8,556	86.6	(84.3 - 88.9)

¹ Defined as having a rule that prohibits smoking anywhere inside the home

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 57a. Binge drinking within the past month among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	640	1,011,000	16.2	(11.5 - 20.8)	37,745	2,602,000	17.7	(15.7 - 19.8)
Gender								
Male	247	697,200	23.9	(15.7 - 32.1)	14,420	1,712,400	24.5	(20.9 - 28.1)
Female	393	313,800	9.4	(5.0 - 13.9)	23,325	889,600	11.6	(9.5 - 13.7)
Age (years)								
18-34	*	*	*	*	3,895	1,254,300	28.5	(22.8 - 34.2)
35-44	100	237,800	18.4	(9.9 - 26.8)	4,934	584,800	20.6	(16.5 - 24.7)
45-54	122	168,800	15.0	(7.2 - 22.8)	7,738	467,600	16.6	(13.3 - 19.9)
55-64	159	73,400	8.9	(4.5 - 13.4)	8,663	207,200	9.8	(7.9 - 11.8)
≥ 65	*	*	*	*	12,515	88,100	3.5	(2.5 - 4.5)
Race/Ethnicity								
White Non-Hispanic	391	449,300	18.9	(13.2 - 24.6)	35,453	1,761,900	19.2	(17.5 - 20.8)
Black Non-Hispanic	*	*	*	*	728	247,000	11.6	(6.3 - 17.0)
Hispanic	*	*	*	*	854	460,600	20.5	(11.3 - 29.8)
Other Non-Hispanic	*	*	*	*	710	132,500	12.1	(3.4 - 20.7)
Education								
≤ High School	224	413,200	16.2	(7.5 - 25.0)	15,491	945,800	17.4	(13.2 - 21.6)
Some College	117	148,900	11.8	(3.2 - 20.5)	9,819	636,600	18.1	(14.6 - 21.5)
≥ College Degree	295	448,900	19.0	(12.6 - 25.4)	12,358	1,018,700	18.2	(15.3 - 21.1)
Income								
≤ \$24,999	167	239,400	12.2	(3.5 - 21.0)	9,557	469,100	13.5	(8.5 - 18.5)
\$25,000-\$49,999	*	*	*	*	9,575	534,300	18.5	(13.0 - 24.0)
\$50,000-\$74,999	*	*	*	*	5,726	378,200	19.2	(14.1 - 24.3)
≥ \$75,000	188	334,400	21.8	(14.2 - 29.5)	8,501	1,011,400	22.6	(19.7 - 25.6)

¹ Defined as men having 5 or more drinks or women having 4 or more drinks on 1 or more occasion within the past month

² Sample size from survey

³ Estimated number of adult binge drinkers within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 57b. Binge drinking within the past month among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	640	15.9	(11.4 - 20.4)	37,745	18.1	(16.1 - 20.2)
Gender						
Male	247	23.2	(15.3 - 31.1)	14,420	24.5	(21.0 - 27.9)
Female	393	9.5	(5.0 - 14.0)	23,325	12.4	(10.2 - 14.5)
Race/Ethnicity						
White Non-Hispanic	391	18.7	(13.1 - 24.4)	35,453	19.8	(18.1 - 21.6)
Black Non-Hispanic	*	*	*	728	11.8	(6.4 - 17.1)
Hispanic	*	*	*	854	20.2	(11.3 - 29.1)
Other Non-Hispanic	*	*	*	710	12.1	(3.4 - 20.8)
Education						
≤ High School	224	15.7	(7.4 - 24.0)	15,491	17.5	(13.5 - 21.5)
Some College	117	11.6	(3.1 - 20.1)	9,819	18.6	(15.1 - 22.0)
≥ College Degree	295	18.8	(12.5 - 25.2)	12,358	18.8	(15.9 - 21.7)
Income						
≤ \$24,999	167	11.6	(3.3 - 19.9)	9,557	13.4	(8.6 - 18.2)
\$25,000-\$49,999	*	*	*	9,575	18.9	(13.5 - 24.2)
\$50,000-\$74,999	*	*	*	5,726	19.9	(14.8 - 25.1)
≥ \$75,000	188	21.5	(13.9 - 29.1)	8,501	23.2	(20.2 - 26.1)

¹ Defined as men having 5 or more drinks or women having 4 or more drinks on 1 or more occasion within the past month

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 58a. Heavy drinking within the past month among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	636	198,100	3.2	(1.8 - 4.6)	37,553	718,700	4.9	(4.2 - 5.6)
Gender								
Male	246	60,400	2.1	(0.6 - 3.6)	14,330	333,300	4.8	(3.9 - 5.7)
Female	390	137,700	4.1	(1.9 - 6.3)	23,223	385,400	5.0	(4.0 - 6.1)
Age (years)								
18-34	*	*	*	*	3,866	224,900	5.1	(3.5 - 6.8)
35-44	*	*	*	*	4,924	137,200	4.8	(3.0 - 6.7)
45-54	*	*	*	*	7,715	144,600	5.2	(4.1 - 6.2)
55-64	158	58,000	7.1	(2.9 - 11.2)	8,624	117,400	5.6	(3.9 - 7.3)
≥ 65	*	*	*	*	12,424	94,600	3.8	(2.7 - 4.9)
Race/Ethnicity								
White Non-Hispanic	387	121,300	5.1	(2.6 - 7.6)	35,279	572,400	6.3	(5.4 - 7.1)
Black Non-Hispanic	*	*	*	*	723	58,100	2.8	(0.6 - 4.9)
Hispanic	*	*	*	*	848	60,900	2.7	(0.5 - 4.9)
Other Non-Hispanic	*	*	*	*	703	27,200	2.5	(0.8 - 4.2)
Education								
≤ High School	*	*	*	*	15,411	228,100	4.2	(3.2 - 5.2)
Some College	*	*	*	*	9,759	163,700	4.7	(3.6 - 5.8)
≥ College Degree	291	136,100	5.8	(2.7 - 8.9)	12,306	326,800	5.9	(4.5 - 7.3)
Income								
≤ \$24,999	*	*	*	*	9,512	104,200	3.0	(1.8 - 4.1)
\$25,000-\$49,999	*	*	*	*	9,537	158,700	5.5	(3.9 - 7.2)
\$50,000-\$74,999	*	*	*	*	5,708	95,400	4.9	(3.1 - 6.7)
≥ \$75,000	189	103,400	6.7	(2.8 - 10.7)	8,481	313,900	7.0	(5.4 - 8.6)

¹ Defined as adult men averaging more than 2 alcoholic drinks per day and adult women averaging more than 1 alcoholic drink per day within the past month

² Sample size from survey

³ Estimated number of adult heavy drinkers within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 58b. Heavy drinking within the past month among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	636	3.2	(1.8 - 4.6)	37,553	5.0	(4.3 - 5.8)
Gender						
Male	246	2.2	(0.6 - 3.7)	14,330	4.9	(3.9 - 5.9)
Female	390	4.1	(1.8 - 6.3)	23,223	5.1	(4.0 - 6.2)
Race/Ethnicity						
White Non-Hispanic	387	5.0	(2.5 - 7.6)	35,279	6.4	(5.5 - 7.2)
Black Non-Hispanic	*	*	*	723	2.9	(0.6 - 5.1)
Hispanic	*	*	*	848	2.8	(0.5 - 5.1)
Other Non-Hispanic	*	*	*	703	2.5	(0.8 - 4.2)
Education						
≤ High School	*	*	*	15,411	4.3	(3.2 - 5.3)
Some College	*	*	*	9,759	4.8	(3.6 - 5.9)
≥ College Degree	291	5.9	(2.7 - 9.0)	12,306	6.0	(4.5 - 7.4)
Income						
≤ \$24,999	*	*	*	9,512	3.0	(1.9 - 4.1)
\$25,000-\$49,999	*	*	*	9,537	5.8	(4.0 - 7.5)
\$50,000-\$74,999	*	*	*	5,708	5.0	(3.1 - 7.0)
≥ \$75,000	189	6.6	(2.6 - 10.5)	8,481	7.0	(5.4 - 8.7)

¹ Defined as adult men averaging more than 2 alcoholic drinks per day and adult women averaging more than 1 alcoholic drink per day within the past month

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 59a. Never or rarely been asked about sexual history during routine check-up among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	599	3,246,300	55.2	(49.3 - 61.0)	35,193	8,480,600	61.9	(59.2 - 64.5)
Gender								
Male	233	1,637,300	60.6	(51.4 - 69.7)	13,299	4,357,400	67.7	(63.6 - 71.7)
Female	366	1,609,100	50.6	(43.2 - 58.0)	21,894	4,123,300	56.8	(53.3 - 60.2)
Age (years)								
18-34	*	*	*	*	3,507	1,804,600	44.3	(37.8 - 50.8)
35-44	*	*	*	*	4,531	1,635,300	62.2	(56.6 - 67.8)
45-54	*	*	*	*	7,215	1,715,600	65.4	(60.5 - 70.3)
55-64	150	489,600	61.8	(52.1 - 71.5)	8,168	1,416,200	70.7	(66.6 - 74.8)
≥ 65	173	725,600	76.9	(68.5 - 85.2)	11,772	1,908,900	80.3	(76.9 - 83.8)
Race/Ethnicity								
White Non-Hispanic	369	1,497,100	64.8	(57.3 - 72.3)	33,090	5,940,300	68.5	(66.3 - 70.7)
Black Non-Hispanic	*	*	*	*	683	831,800	40.8	(31.8 - 49.9)
Hispanic	*	*	*	*	784	1,071,700	53.4	(43.6 - 63.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	206	1,250,700	53.1	(43.4 - 62.9)	14,264	3,001,200	60.1	(55.4 - 64.9)
Some College	*	*	*	*	9,156	1,972,800	60.3	(54.9 - 65.7)
≥ College Degree	279	1,367,800	59.8	(51.6 - 68.1)	11,706	3,476,000	65.0	(61.2 - 68.7)
Income								
≤ \$24,999	*	*	*	*	8,772	1,529,400	49.3	(42.6 - 56.0)
\$25,000-\$49,999	*	*	*	*	8,932	1,705,000	64.4	(58.0 - 70.7)
\$50,000-\$74,999	*	*	*	*	5,433	1,217,000	64.5	(58.5 - 70.5)
≥ \$75,000	183	1,018,000	67.2	(58.2 - 76.1)	8,108	2,934,300	68.2	(64.7 - 71.7)

¹ Defined as respondents who never or rarely (hardly ever) were asked about their sexual history, such as about sexual partners and sexual practices, during a routine check-up at a doctor's office or clinic

² Sample size from survey

³ Estimated number of adults who were never or rarely asked about their sexual history during a routine check-up at a doctor's office or clinic (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 59b. Never or rarely been asked about sexual history during routine check-up among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	599	55.6	(49.8 - 61.4)	35,193	61.4	(58.7 - 64.0)
Gender						
Male	233	62.1	(53.3 - 70.9)	13,299	67.9	(64.0 - 71.9)
Female	366	50.1	(42.7 - 57.6)	21,894	55.5	(52.0 - 59.0)
Race/Ethnicity						
White Non-Hispanic	369	65.4	(58.0 - 72.9)	33,090	67.8	(65.6 - 70.1)
Black Non-Hispanic	*	*	*	683	41.1	(32.0 - 50.2)
Hispanic	*	*	*	784	53.0	(43.3 - 62.6)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	206	53.3	(43.7 - 63.0)	14,264	59.5	(54.7 - 64.3)
Some College	*	*	*	9,156	59.7	(54.4 - 65.0)
≥ College Degree	279	60.4	(52.3 - 68.6)	11,706	64.5	(60.8 - 68.2)
Income						
≤ \$24,999	*	*	*	8,772	48.7	(42.0 - 55.4)
\$25,000-\$49,999	*	*	*	8,932	63.7	(57.5 - 70.0)
\$50,000-\$74,999	*	*	*	5,433	64.2	(58.2 - 70.2)
≥ \$75,000	183	67.6	(58.8 - 76.5)	8,108	67.5	(64.0 - 71.0)

¹ Defined as respondents who never or rarely (hardly ever) were asked about their sexual history, such as about sexual partners and sexual practices, during a routine check-up at a doctor's office or clinic

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 60a. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	340	2,645,000	76.9	(70.3 - 83.4)	22,788	7,011,100	77.9	(75.2 - 80.5)
Gender								
Male	139	1,310,600	75.8	(66.4 - 85.2)	9,034	3,618,500	81.0	(77.2 - 84.8)
Female	201	1,334,400	77.9	(68.8 - 87.1)	13,754	3,392,600	74.8	(71.2 - 78.4)
Age (years)								
18-34	*	*	*	*	3,213	2,362,500	70.8	(65.1 - 76.6)
35-44	*	*	*	*	3,513	1,313,400	71.7	(65.6 - 77.7)
45-54	*	*	*	*	4,975	1,290,600	80.3	(75.9 - 84.8)
55-64	76	317,700	87.4	(79.2 - 95.7)	4,955	932,200	87.2	(84.1 - 90.3)
≥ 65	87	389,500	97.0	(92.9 - 100)	6,132	1,112,400	95.9	(94.4 - 97.5)
Race/Ethnicity								
White Non-Hispanic	228	1,213,000	75.0	(65.7 - 84.4)	21,540	4,875,200	78.3	(75.6 - 80.9)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	56	671,100	86.9	(77.4 - 96.5)	491	1,002,400	84.8	(77.8 - 91.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	108	1,089,700	82.0	(72.5 - 91.5)	8,879	2,581,700	82.1	(77.9 - 86.3)
Some College	*	*	*	*	5,953	1,599,500	77.5	(72.6 - 82.3)
≥ College Degree	176	1,086,400	73.8	(64.6 - 83.0)	7,928	2,824,800	75.6	(71.8 - 79.5)
Income								
≤ \$24,999	*	*	*	*	5,491	1,602,900	78.7	(71.9 - 85.6)
\$25,000-\$49,999	*	*	*	*	5,912	1,335,900	78.8	(73.8 - 83.9)
\$50,000-\$74,999	*	*	*	*	3,717	987,500	76.6	(70.4 - 82.8)
≥ \$75,000	*	*	*	*	5,771	2,321,600	77.6	(74.0 - 81.2)

¹ Defined as respondent estimating there were hardly any (0-1 out of 10) or a few (2-3 out of 10) persons their age in their community who have had a sexually transmitted disease such as gonorrhea, chlamydia, herpes or syphilis

² Sample size from survey

³ Estimated number of adults who believed hardly any or a few people their age in their community have had a sexually transmitted disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 60b. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	340	77.0	(70.5 - 83.4)	22,788	77.3	(74.7 - 79.9)
Gender						
Male	139	76.2	(67.0 - 85.4)	9,034	81.1	(77.4 - 84.8)
Female	201	77.7	(68.6 - 86.9)	13,754	73.6	(70.0 - 77.3)
Race/Ethnicity						
White Non-Hispanic	228	75.3	(66.1 - 84.5)	21,540	77.5	(74.9 - 80.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	56	86.9	(77.4 - 96.4)	491	84.6	(77.6 - 91.5)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	108	82.2	(72.9 - 91.5)	8,879	81.6	(77.4 - 85.7)
Some College	*	*	*	5,953	76.8	(72.0 - 81.6)
≥ College Degree	176	74.1	(65.0 - 83.2)	7,928	75.1	(71.3 - 79.0)
Income						
≤ \$24,999	*	*	*	5,491	77.9	(71.0 - 84.7)
\$25,000-\$49,999	*	*	*	5,912	77.9	(72.8 - 83.0)
\$50,000-\$74,999	*	*	*	3,717	76.1	(70.0 - 82.3)
≥ \$75,000	*	*	*	5,771	77.0	(73.4 - 80.6)

¹ Defined as respondent estimating there were hardly any (0-1 out of 10) or a few (2-3 out of 10) persons their age in their community who have had a sexually transmitted disease such as gonorrhea, chlamydia, herpes or syphilis

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 61a. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	610	5,298,800	88.5	(85.0 - 92.0)	36,287	12,738,300	89.7	(88.2 - 91.3)
Gender								
Male	241	2,521,900	88.5	(83.6 - 93.4)	13,893	6,104,600	89.4	(87.2 - 91.5)
Female	369	2,776,900	88.5	(83.6 - 93.4)	22,394	6,633,700	90.1	(87.9 - 92.2)
Age (years)								
18-34	74	1,829,300	89.9	(82.0 - 97.8)	3,850	4,007,400	91.3	(87.5 - 95.1)
35-44	98	1,142,600	92.0	(85.9 - 98.0)	4,875	2,578,600	93.2	(90.4 - 96.0)
45-54	115	868,000	86.5	(79.7 - 93.3)	7,607	2,416,800	90.5	(87.9 - 93.2)
55-64	150	734,700	93.5	(88.2 - 98.8)	8,411	1,876,500	92.2	(90.0 - 94.5)
≥ 65	173	724,300	78.7	(71.1 - 86.3)	11,544	1,858,900	79.7	(76.6 - 82.8)
Race/Ethnicity								
White Non-Hispanic	374	1,973,700	84.2	(78.3 - 90.2)	34,102	8,050,700	89.4	(87.7 - 91.0)
Black Non-Hispanic	85	1,253,200	93.9	(89.0 - 98.8)	702	1,822,200	91.9	(88.2 - 95.6)
Hispanic	112	1,424,200	89.3	(83.0 - 95.6)	820	1,967,200	89.9	(85.2 - 94.7)
Other Non-Hispanic	*	*	*	*	663	898,200	88.6	(79.5 - 97.6)
Education								
≤ High School	208	2,051,100	85.5	(80.2 - 90.9)	14,507	4,385,300	84.6	(81.9 - 87.2)
Some College	112	1,045,300	85.9	(76.8 - 95.1)	9,571	3,094,300	90.0	(86.7 - 93.4)
≥ College Degree	287	2,184,800	95.0	(91.9 - 98.0)	12,142	5,228,800	95.4	(94.0 - 96.7)
Income								
≤ \$24,999	154	1,607,200	88.8	(82.1 - 95.5)	9,021	2,833,500	86.5	(82.6 - 90.3)
\$25,000-\$49,999	103	1,028,500	89.6	(82.6 - 96.5)	9,261	2,529,900	89.5	(86.5 - 92.5)
\$50,000-\$74,999	*	*	*	*	5,644	1,740,000	91.2	(87.4 - 95.1)
≥ \$75,000	186	1,387,900	91.4	(86.0 - 96.8)	8,397	4,184,300	94.4	(92.5 - 96.3)

¹ Defined as believing it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community

² Sample size from survey

³ Estimated number of adults who believed it is at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 61b. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	610	88.3	(84.8 - 91.8)	36,287	89.8	(88.3 - 91.4)
Gender						
Male	241	88.0	(83.0 - 92.9)	13,893	89.1	(86.9 - 91.3)
Female	369	88.6	(83.6 - 93.6)	22,394	90.5	(88.3 - 92.7)
Race/Ethnicity						
White Non-Hispanic	374	84.0	(77.9 - 90.0)	34,102	89.5	(87.8 - 91.2)
Black Non-Hispanic	85	93.5	(88.3 - 98.8)	702	91.7	(87.8 - 95.6)
Hispanic	112	89.0	(82.5 - 95.4)	820	89.6	(84.7 - 94.5)
Other Non-Hispanic	*	*	*	663	89.0	(80.5 - 97.6)
Education						
≤ High School	208	85.2	(79.7 - 90.8)	14,507	84.6	(81.8 - 87.4)
Some College	112	85.7	(76.6 - 94.7)	9,571	90.2	(86.9 - 93.5)
≥ College Degree	287	94.9	(91.9 - 97.9)	12,142	95.4	(94.0 - 96.7)
Income						
≤ \$24,999	154	88.6	(81.8 - 95.3)	9,021	86.5	(82.6 - 90.4)
\$25,000-\$49,999	103	89.5	(82.5 - 96.4)	9,261	89.6	(86.5 - 92.6)
\$50,000-\$74,999	*	*	*	5,644	91.1	(86.9 - 95.3)
≥ \$75,000	186	91.5	(86.2 - 96.8)	8,397	94.5	(92.6 - 96.4)

¹ Defined as believing it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 62a. Overweight among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	615	2,198,800	36.0	(30.7 - 41.3)	36,710	5,213,000	36.4	(34.0 - 38.8)
Gender								
Male	248	1,251,900	42.6	(33.9 - 51.2)	14,502	3,073,200	43.7	(39.9 - 47.6)
Female	367	947,000	29.9	(23.6 - 36.3)	22,208	2,139,800	29.3	(26.4 - 32.2)
Age (years)								
18-34	*	*	*	*	3,789	1,295,200	30.1	(24.7 - 35.6)
35-44	*	*	*	*	4,785	1,052,000	37.5	(31.7 - 43.3)
45-54	*	*	*	*	7,495	1,053,300	38.8	(34.1 - 43.4)
55-64	153	293,800	35.9	(26.8 - 45.1)	8,356	798,100	38.8	(34.8 - 42.7)
≥ 65	177	424,500	44.2	(34.7 - 53.6)	12,285	1,014,400	41.5	(37.6 - 45.4)
Race/Ethnicity								
White Non-Hispanic	375	876,100	37.2	(30.5 - 43.9)	34,501	3,282,200	36.4	(34.4 - 38.3)
Black Non-Hispanic	*	*	*	*	708	740,000	36.7	(28.3 - 45.1)
Hispanic	*	*	*	*	813	781,500	35.6	(27.1 - 44.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	215	900,800	35.9	(27.4 - 44.4)	15,079	1,889,700	35.4	(31.2 - 39.5)
Some College	*	*	*	*	9,515	1,314,500	38.7	(33.8 - 43.7)
≥ College Degree	289	794,600	34.0	(26.2 - 41.7)	12,062	1,996,900	36.3	(32.8 - 39.7)
Income								
≤ \$24,999	163	660,200	33.8	(23.9 - 43.8)	9,346	1,170,900	34.0	(28.2 - 39.8)
\$25,000-\$49,999	*	*	*	*	9,368	1,136,000	40.4	(34.6 - 46.2)
\$50,000-\$74,999	*	*	*	*	5,615	738,700	38.3	(32.3 - 44.3)
≥ \$75,000	187	540,100	35.4	(26.6 - 44.1)	8,337	1,659,100	37.5	(34.1 - 40.9)

¹ Defined as having a body mass index (BMI) more than 24.9 but less than 30.0. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Sample size from survey

³ Estimated number of adults who are overweight (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 62b. Overweight among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	615	36.1	(30.8 - 41.4)	36,710	36.2	(33.8 - 38.6)
Gender						
Male	248	43.0	(34.5 - 51.4)	14,502	43.9	(40.1 - 47.6)
Female	367	29.7	(23.2 - 36.1)	22,208	28.9	(25.9 - 31.8)
Race/Ethnicity						
White Non-Hispanic	375	36.9	(30.3 - 43.5)	34,501	36.0	(34.0 - 38.0)
Black Non-Hispanic	*	*	*	708	37.0	(28.5 - 45.5)
Hispanic	*	*	*	813	36.0	(27.5 - 44.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	215	35.9	(27.5 - 44.3)	15,079	35.2	(31.1 - 39.4)
Some College	*	*	*	9,515	38.3	(33.3 - 43.2)
≥ College Degree	289	34.3	(26.5 - 42.1)	12,062	36.2	(32.7 - 39.8)
Income						
≤ \$24,999	163	33.7	(23.9 - 43.5)	9,346	33.8	(28.0 - 39.5)
\$25,000-\$49,999	*	*	*	9,368	40.6	(34.8 - 46.4)
\$50,000-\$74,999	*	*	*	5,615	38.6	(32.5 - 44.7)
≥ \$75,000	187	35.7	(26.8 - 44.6)	8,337	37.2	(33.7 - 40.6)

¹ Defined as having a body mass index (BMI) more than 24.9 but less than 30.0. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 63a. Obesity among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	615	1,302,500	21.3	(16.8 - 25.9)	36,710	3,327,400	23.2	(21.2 - 25.3)
Gender								
Male	248	427,600	14.5	(9.5 - 19.6)	14,502	1,505,800	21.4	(19.0 - 23.9)
Female	367	874,800	27.6	(20.7 - 34.6)	22,208	1,821,600	25.0	(21.8 - 28.1)
Age (years)								
18-34	71	294,800	15.0	(5.5 - 24.6)	3,789	730,600	17.0	(12.4 - 21.6)
35-44	100	240,100	18.5	(10.0 - 27.0)	4,785	662,200	23.6	(19.4 - 27.9)
45-54	*	*	*	*	7,495	816,800	30.1	(25.3 - 34.8)
55-64	153	200,000	24.5	(15.9 - 33.1)	8,356	563,900	27.4	(23.7 - 31.1)
≥ 65	177	216,600	22.5	(14.0 - 31.1)	12,285	554,000	22.7	(19.2 - 26.1)
Race/Ethnicity								
White Non-Hispanic	375	442,200	18.8	(12.8 - 24.7)	34,501	2,062,700	22.9	(21.2 - 24.6)
Black Non-Hispanic	*	*	*	*	708	703,100	34.9	(26.1 - 43.7)
Hispanic	111	315,700	19.7	(11.3 - 28.2)	813	456,400	20.8	(14.3 - 27.2)
Other Non-Hispanic	*	*	*	*	688	105,200	9.6	(4.0 - 15.3)
Education								
≤ High School	215	762,700	30.4	(22.3 - 38.5)	15,079	1,561,600	29.2	(25.3 - 33.2)
Some College	*	*	*	*	9,515	792,800	23.4	(19.4 - 27.3)
≥ College Degree	289	271,200	11.6	(7.1 - 16.0)	12,062	915,000	16.6	(14.4 - 18.8)
Income								
≤ \$24,999	163	550,100	28.2	(18.3 - 38.1)	9,346	974,800	28.3	(22.6 - 34.0)
\$25,000-\$49,999	100	139,300	12.5	(4.8 - 20.2)	9,368	584,800	20.8	(17.1 - 24.5)
\$50,000-\$74,999	*	*	*	*	5,615	475,300	24.6	(20.0 - 29.3)
≥ \$75,000	187	239,000	15.6	(9.0 - 22.2)	8,337	867,700	19.6	(17.0 - 22.3)

¹ Defined as having a body mass index (BMI) of 30.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Sample size from survey

³ Estimated number of adults who are obese (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 63b. Obesity among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	615	21.5	(16.9 - 26.1)	36,710	23.1	(21.0 - 25.1)
Gender						
Male	248	14.9	(9.9 - 20.0)	14,502	21.5	(19.0 - 24.0)
Female	367	27.6	(20.6 - 34.6)	22,208	24.6	(21.5 - 27.8)
Race/Ethnicity						
White Non-Hispanic	375	19.0	(13.0 - 25.0)	34,501	22.8	(21.0 - 24.5)
Black Non-Hispanic	*	*	*	708	34.8	(25.9 - 43.6)
Hispanic	111	19.7	(11.3 - 28.0)	813	20.5	(14.1 - 26.9)
Other Non-Hispanic	*	*	*	688	9.5	(3.8 - 15.2)
Education						
≤ High School	215	31.0	(22.8 - 39.1)	15,079	29.4	(25.4 - 33.3)
Some College	*	*	*	9,515	23.1	(19.2 - 27.1)
≥ College Degree	289	11.4	(7.0 - 15.7)	12,062	16.4	(14.2 - 18.5)
Income						
≤ \$24,999	163	28.6	(18.7 - 38.5)	9,346	28.3	(22.6 - 34.1)
\$25,000-\$49,999	100	13.3	(5.2 - 21.4)	9,368	21.0	(17.2 - 24.8)
\$50,000-\$74,999	*	*	*	5,615	24.0	(19.4 - 28.7)
≥ \$75,000	187	15.3	(8.8 - 21.7)	8,337	19.3	(16.7 - 21.9)

¹ Defined as having a body mass index (BMI) of 30.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 64a. Overweight or obesity among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	615	3,501,300	57.3	(51.5 - 63.2)	36,710	8,540,500	59.6	(57.0 - 62.2)
Gender								
Male	248	1,679,500	57.1	(47.9 - 66.3)	14,502	4,579,100	65.2	(61.0 - 69.4)
Female	367	1,821,800	57.6	(50.3 - 64.8)	22,208	3,961,400	54.3	(51.0 - 57.6)
Age (years)								
18-34	*	*	*	*	3,789	2,025,800	47.1	(40.8 - 53.5)
35-44	*	*	*	*	4,785	1,714,200	61.1	(55.6 - 66.6)
45-54	*	*	*	*	7,495	1,870,100	68.8	(64.6 - 73.1)
55-64	*	*	*	*	8,356	1,362,000	66.2	(61.9 - 70.4)
≥ 65	177	641,200	66.7	(58.3 - 75.2)	12,285	1,568,400	64.1	(60.6 - 67.7)
Race/Ethnicity								
White Non-Hispanic	375	1,318,300	56.0	(49.0 - 63.0)	34,501	5,344,900	59.3	(57.2 - 61.3)
Black Non-Hispanic	*	*	*	*	708	1,443,100	71.6	(62.5 - 80.7)
Hispanic	*	*	*	*	813	1,237,900	56.4	(46.5 - 66.2)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	215	1,663,400	66.3	(56.3 - 76.3)	15,079	3,451,400	64.6	(59.8 - 69.4)
Some College	*	*	*	*	9,515	2,107,300	62.1	(57.0 - 67.2)
≥ College Degree	289	1,065,800	45.6	(37.4 - 53.7)	12,062	2,911,900	52.9	(49.2 - 56.6)
Income								
≤ \$24,999	*	*	*	*	9,346	2,145,700	62.3	(55.7 - 68.9)
\$25,000-\$49,999	*	*	*	*	9,368	1,720,700	61.2	(54.5 - 67.9)
\$50,000-\$74,999	*	*	*	*	5,615	1,214,000	62.9	(57.1 - 68.8)
≥ \$75,000	187	779,200	51.0	(41.8 - 60.2)	8,337	2,526,700	57.1	(53.6 - 60.6)

¹ Defined as having a body mass index (BMI) of 25.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Sample size from survey

³ Estimated number of adults who are overweight or obese (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 64b. Overweight or obesity among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	615	57.6	(51.9 - 63.3)	36,710	59.3	(56.8 - 61.9)
Gender						
Male	248	57.9	(49.0 - 66.8)	14,502	65.4	(61.3 - 69.4)
Female	367	57.3	(50.0 - 64.6)	22,208	53.5	(50.1 - 56.8)
Race/Ethnicity						
White Non-Hispanic	375	55.9	(48.9 - 62.9)	34,501	58.8	(56.7 - 60.8)
Black Non-Hispanic	*	*	*	708	71.8	(63.0 - 80.6)
Hispanic	*	*	*	813	56.5	(46.8 - 66.1)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	215	66.9	(57.1 - 76.6)	15,079	64.6	(59.9 - 69.3)
Some College	*	*	*	9,515	61.4	(56.4 - 66.5)
≥ College Degree	289	45.7	(37.6 - 53.8)	12,062	52.6	(48.9 - 56.3)
Income						
≤ \$24,999	*	*	*	9,346	62.1	(55.6 - 68.6)
\$25,000-\$49,999	*	*	*	9,368	61.6	(55.1 - 68.1)
\$50,000-\$74,999	*	*	*	5,615	62.6	(56.7 - 68.6)
≥ \$75,000	187	50.9	(41.7 - 60.1)	8,337	56.5	(52.9 - 60.0)

¹ Defined as having a body mass index (BMI) of 25.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 65a. Received advice about weight by a health professional among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	619	1,827,200	29.8	(24.7 - 35.0)	36,729	3,929,700	27.4	(25.1 - 29.7)
Gender								
Male	247	653,700	22.3	(15.6 - 29.0)	14,499	1,654,800	23.6	(20.6 - 26.6)
Female	372	1,173,500	36.8	(29.5 - 44.0)	22,230	2,274,900	31.0	(27.7 - 34.3)
Age (years)								
18-34	*	*	*	*	3,796	895,500	20.7	(15.3 - 26.2)
35-44	100	311,600	24.1	(14.2 - 34.0)	4,779	722,700	25.8	(20.9 - 30.6)
45-54	*	*	*	*	7,490	910,900	33.5	(29.0 - 38.1)
55-64	152	317,200	39.0	(29.2 - 48.9)	8,351	741,100	36.1	(32.0 - 40.3)
≥ 65	180	294,400	30.4	(21.1 - 39.8)	12,313	659,500	26.9	(23.0 - 30.7)
Race/Ethnicity								
White Non-Hispanic	377	699,300	29.7	(23.1 - 36.3)	34,502	2,386,200	26.5	(24.6 - 28.4)
Black Non-Hispanic	*	*	*	*	709	615,700	30.4	(22.0 - 38.8)
Hispanic	112	410,100	25.5	(15.7 - 35.4)	830	555,400	25.1	(17.8 - 32.4)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	214	738,700	29.5	(21.4 - 37.7)	15,116	1,469,700	27.5	(23.6 - 31.4)
Some College	*	*	*	*	9,515	946,800	27.8	(23.2 - 32.4)
≥ College Degree	291	651,100	27.8	(20.6 - 34.9)	12,041	1,445,500	26.2	(23.0 - 29.4)
Income								
≤ \$24,999	*	*	*	*	9,360	1,079,400	31.4	(25.3 - 37.5)
\$25,000-\$49,999	*	*	*	*	9,365	696,900	24.6	(19.8 - 29.4)
\$50,000-\$74,999	*	*	*	*	5,603	507,100	26.3	(21.6 - 31.0)
≥ \$75,000	187	485,800	31.8	(23.5 - 40.1)	8,334	1,207,100	27.3	(24.2 - 30.5)

¹ Defined as being advised by a doctor, nurse or other health professional about their weight (lose, gain or maintain)

² Sample size from survey

³ Estimated number of adults provided advice by a health professional about their weight (lose, gain or maintain) (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 65b. Received advice about weight by a health professional among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	619	29.7	(24.7 - 34.8)	36,729	27.1	(24.8 - 29.4)
Gender						
Male	247	22.3	(15.8 - 28.8)	14,499	23.4	(20.5 - 26.3)
Female	372	36.5	(29.2 - 43.8)	22,230	30.6	(27.3 - 34.0)
Race/Ethnicity						
White Non-Hispanic	377	29.7	(23.1 - 36.3)	34,502	26.1	(24.2 - 28.0)
Black Non-Hispanic	*	*	*	709	30.4	(21.9 - 38.9)
Hispanic	112	25.3	(15.6 - 35.0)	830	24.9	(17.7 - 32.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	214	29.5	(21.4 - 37.6)	15,116	27.3	(23.4 - 31.2)
Some College	*	*	*	9,515	27.4	(22.9 - 32.0)
≥ College Degree	291	27.8	(20.7 - 34.9)	12,041	25.9	(22.7 - 29.0)
Income						
≤ \$24,999	*	*	*	9,360	31.0	(24.9 - 37.0)
\$25,000-\$49,999	*	*	*	9,365	24.9	(20.0 - 29.8)
\$50,000-\$74,999	*	*	*	5,603	25.9	(21.2 - 30.6)
≥ \$75,000	187	32.0	(23.6 - 40.4)	8,334	26.9	(23.7 - 30.0)

¹ Defined as being advised by a doctor, nurse or other health professional about their weight (lose, gain or maintain)

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 66a. Received advice to lose weight by a health professional among adults provided weight advice¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	192	1,586,100	86.8	(80.7 - 92.9)	9,724	3,412,900	86.8	(83.9 - 89.8)
Gender								
Male	72	599,800	91.8	(85.4 - 98.2)	3,807	1,468,700	88.8	(85.7 - 91.8)
Female	120	986,200	84.0	(75.3 - 92.8)	5,917	1,944,200	85.5	(80.8 - 90.1)
Age (years)								
18-34	*	*	*	*	681	779,700	87.1	(80.6 - 93.6)
35-44	*	*	*	*	1,215	622,100	86.1	(76.9 - 95.2)
45-54	*	*	*	*	2,238	813,300	89.3	(84.4 - 94.2)
55-64	*	*	*	*	2,697	645,500	87.1	(81.2 - 93.0)
≥ 65	*	*	*	*	2,893	552,300	83.7	(76.9 - 90.6)
Race/Ethnicity								
White Non-Hispanic	112	622,100	89.0	(81.5 - 96.4)	9,082	2,076,700	87.0	(84.5 - 89.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	251	483,600	87.1	(78.5 - 95.7)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	3,962	1,239,700	84.3	(78.6 - 90.1)
Some College	*	*	*	*	2,633	814,800	86.1	(80.1 - 92.0)
≥ College Degree	87	577,900	88.8	(80.2 - 97.3)	3,115	1,290,900	89.3	(85.3 - 93.3)
Income								
≤ \$24,999	*	*	*	*	2,539	922,100	85.4	(78.7 - 92.1)
\$25,000-\$49,999	*	*	*	*	2,482	594,100	85.2	(78.4 - 92.1)
\$50,000-\$74,999	*	*	*	*	1,521	446,100	88.0	(83.2 - 92.7)
≥ \$75,000	*	*	*	*	2,270	1,073,200	88.9	(83.7 - 94.1)

¹ Defined as provided advice by a health professional about their weight and advised to lose weight

² Sample size from survey

³ Estimated number of adults provided advice by a health professional about their weight and advised to lose weight (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 66b. Received advice to lose weight by a health professional among adults provided weight advice¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	192	86.9	(80.6 - 93.1)	9,724	86.9	(83.8 - 89.9)
Gender						
Male	72	92.2	(86.1 - 98.3)	3,807	88.7	(85.7 - 91.8)
Female	120	83.9	(74.9 - 92.9)	5,917	85.5	(80.7 - 90.3)
Race/Ethnicity						
White Non-Hispanic	112	89.2	(81.8 - 96.6)	9,082	87.0	(84.4 - 89.6)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	251	87.6	(79.3 - 95.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	3,962	84.6	(78.7 - 90.5)
Some College	*	*	*	2,633	85.6	(79.5 - 91.7)
≥ College Degree	87	89.1	(80.7 - 97.5)	3,115	89.4	(85.4 - 93.3)
Income						
≤ \$24,999	*	*	*	2,539	85.7	(79.0 - 92.3)
\$25,000-\$49,999	*	*	*	2,482	85.3	(78.4 - 92.3)
\$50,000-\$74,999	*	*	*	1,521	88.0	(83.1 - 92.8)
≥ \$75,000	*	*	*	2,270	88.7	(83.1 - 94.2)

¹ Defined as provided advice by a health professional about their weight and advised to lose weight

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 67a. No leisure-time physical activity among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	646	1,723,300	27.1	(22.1 - 32.2)	38,128	3,541,700	23.9	(21.6 - 26.1)
Gender								
Male	251	716,300	24.0	(16.9 - 31.2)	14,602	1,482,500	20.9	(17.8 - 24.0)
Female	395	1,006,900	29.9	(23.0 - 36.8)	23,526	2,059,200	26.6	(23.4 - 29.7)
Age (years)								
18-34	*	*	*	*	3,938	982,900	22.0	(16.6 - 27.5)
35-44	102	281,600	21.3	(12.3 - 30.4)	4,988	556,000	19.3	(15.0 - 23.6)
45-54	*	*	*	*	7,827	759,000	26.8	(21.7 - 31.9)
55-64	160	188,700	22.3	(14.0 - 30.5)	8,765	464,900	21.7	(18.2 - 25.2)
≥ 65	187	294,700	29.2	(20.9 - 37.5)	12,610	778,800	30.7	(27.3 - 34.2)
Race/Ethnicity								
White Non-Hispanic	395	496,800	20.4	(14.4 - 26.3)	35,816	1,892,800	20.3	(18.6 - 22.0)
Black Non-Hispanic	*	*	*	*	739	690,000	32.0	(23.4 - 40.6)
Hispanic	*	*	*	*	860	600,800	26.7	(19.1 - 34.4)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	230	869,100	32.9	(24.6 - 41.1)	15,683	1,810,200	32.5	(28.4 - 36.5)
Some College	*	*	*	*	9,906	814,200	23.0	(18.6 - 27.3)
≥ College Degree	294	438,400	18.6	(11.6 - 25.5)	12,458	841,000	14.9	(11.9 - 18.0)
Income								
≤ \$24,999	*	*	*	*	9,640	1,332,200	38.0	(31.7 - 44.3)
\$25,000-\$49,999	106	220,100	18.9	(9.9 - 27.9)	9,658	648,000	22.3	(18.3 - 26.3)
\$50,000-\$74,999	*	*	*	*	5,773	430,700	21.8	(16.6 - 27.0)
≥ \$75,000	189	188,500	12.3	(6.6 - 17.9)	8,565	560,800	12.5	(10.3 - 14.6)

¹ Defined as having not participated in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise

² Sample size from survey

³ Estimated number of adults who have not participated in any physical activities or exercises (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 67b. No leisure-time physical activity among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	646	27.2	(22.2 - 32.2)	38,128	23.7	(21.5 - 25.9)
Gender						
Male	251	24.1	(17.1 - 31.1)	14,602	20.9	(17.8 - 23.9)
Female	395	30.0	(23.0 - 37.0)	23,526	26.3	(23.1 - 29.5)
Race/Ethnicity						
White Non-Hispanic	395	20.3	(14.4 - 26.3)	35,816	20.0	(18.3 - 21.7)
Black Non-Hispanic	*	*	*	739	32.0	(23.3 - 40.6)
Hispanic	*	*	*	860	26.9	(19.2 - 34.5)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	230	32.8	(24.6 - 40.9)	15,683	32.3	(28.3 - 36.4)
Some College	*	*	*	9,906	22.8	(18.5 - 27.2)
≥ College Degree	294	18.7	(11.8 - 25.6)	12,458	14.9	(11.8 - 17.9)
Income						
≤ \$24,999	*	*	*	9,640	37.9	(31.6 - 44.2)
\$25,000-\$49,999	106	18.9	(9.9 - 27.9)	9,658	22.0	(18.0 - 26.0)
\$50,000-\$74,999	*	*	*	5,773	21.6	(16.3 - 26.8)
≥ \$75,000	189	12.7	(6.9 - 18.6)	8,565	12.6	(10.3 - 14.9)

¹ Defined as having not participated in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 68a. Consumption of 5 or more servings of fruits and vegetables per day among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	649	1,663,500	26.1	(21.3 - 30.9)	38,203	4,023,400	27.0	(24.9 - 29.2)
Gender								
Male	251	686,100	23.0	(15.7 - 30.3)	14,634	1,603,900	22.6	(19.4 - 25.8)
Female	398	977,400	28.8	(22.5 - 35.1)	23,569	2,419,500	31.1	(28.2 - 34.0)
Age (years)								
18-34	*	*	*	*	3,940	1,261,700	28.3	(22.7 - 33.9)
35-44	102	320,700	24.3	(15.2 - 33.5)	4,989	688,000	23.9	(19.5 - 28.3)
45-54	124	286,800	25.1	(16.0 - 34.2)	7,837	746,800	26.2	(22.3 - 30.1)
55-64	160	222,700	26.3	(18.2 - 34.3)	8,777	579,100	27.0	(23.6 - 30.5)
≥ 65	188	291,300	28.8	(20.6 - 36.9)	12,660	747,800	29.4	(26.0 - 32.8)
Race/Ethnicity								
White Non-Hispanic	395	688,700	28.2	(22.2 - 34.3)	35,884	2,538,000	27.2	(25.4 - 28.9)
Black Non-Hispanic	94	334,500	22.5	(12.5 - 32.5)	740	572,400	26.5	(19.2 - 33.9)
Hispanic	*	*	*	*	864	672,200	29.8	(21.0 - 38.5)
Other Non-Hispanic	*	*	*	*	715	240,800	21.4	(12.0 - 30.8)
Education								
≤ High School	232	670,900	25.1	(16.9 - 33.4)	15,726	1,309,800	23.4	(19.3 - 27.4)
Some College	*	*	*	*	9,924	964,300	27.2	(23.0 - 31.3)
≥ College Degree	295	683,300	28.9	(22.0 - 35.8)	12,471	1,747,500	31.0	(27.9 - 34.1)
Income								
≤ \$24,999	169	446,500	22.6	(13.9 - 31.2)	9,667	831,500	23.7	(18.6 - 28.7)
\$25,000-\$49,999	*	*	*	*	9,669	917,200	31.6	(25.6 - 37.6)
\$50,000-\$74,999	77	135,000	19.0	(10.1 - 27.9)	5,777	501,000	25.3	(21.4 - 29.3)
≥ \$75,000	189	418,300	27.2	(19.3 - 35.1)	8,569	1,260,100	28.0	(25.0 - 31.0)

¹ Defined as having consumed 5 or more servings of fruits and vegetables in an average day

² Sample size from survey

³ Estimated number of adults who consume 5 or more servings of fruits and vegetables per day (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
New York City

Table 68b. Consumption of 5 or more servings of fruits and vegetables per day among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	649	26.2	(21.4 - 31.0)	38,203	27.1	(24.9 - 29.2)
Gender						
Male	251	23.3	(16.2 - 30.4)	14,634	22.7	(19.6 - 25.9)
Female	398	28.8	(22.4 - 35.1)	23,569	31.0	(28.1 - 34.0)
Race/Ethnicity						
White Non-Hispanic	395	28.4	(22.4 - 34.5)	35,884	27.1	(25.4 - 28.9)
Black Non-Hispanic	94	22.3	(12.4 - 32.2)	740	26.8	(19.4 - 34.2)
Hispanic	*	*	*	864	29.5	(20.9 - 38.1)
Other Non-Hispanic	*	*	*	715	21.8	(12.2 - 31.5)
Education						
≤ High School	232	25.0	(16.9 - 33.1)	15,726	23.3	(19.3 - 27.3)
Some College	*	*	*	9,924	27.1	(22.9 - 31.2)
≥ College Degree	295	29.4	(22.5 - 36.3)	12,471	31.2	(28.0 - 34.3)
Income						
≤ \$24,999	169	22.9	(14.2 - 31.6)	9,667	23.9	(18.8 - 28.9)
\$25,000-\$49,999	*	*	*	9,669	32.0	(26.0 - 38.0)
\$50,000-\$74,999	77	18.8	(10.0 - 27.7)	5,777	25.0	(21.0 - 28.9)
≥ \$75,000	189	27.5	(19.5 - 35.5)	8,569	28.0	(24.9 - 31.1)

¹ Defined as having consumed 5 or more servings of fruits and vegetables in an average day

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 69a. Had a fall within the past 3 months among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	456	552,900	19.2	(14.1 - 24.2)	28,783	1,244,800	17.0	(14.9 - 19.0)
Gender								
Male	184	197,400	14.9	(8.9 - 21.0)	10,966	489,400	14.4	(11.9 - 16.8)
Female	272	355,500	22.8	(15.2 - 30.3)	17,817	755,300	19.2	(16.1 - 22.3)
Age (years)								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	120	183,900	16.8	(7.0 - 26.5)	7,741	445,500	16.0	(12.0 - 20.0)
55-64	150	143,800	18.2	(10.6 - 25.9)	8,454	336,000	16.5	(13.4 - 19.6)
≥ 65	186	225,300	22.5	(14.8 - 30.3)	12,588	463,300	18.4	(15.2 - 21.6)
Race/Ethnicity								
White Non-Hispanic	305	263,900	19.9	(14.4 - 25.4)	27,456	869,000	16.6	(15.1 - 18.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	424	76,100	11.0	(4.5 - 17.4)
Other Non-Hispanic	*	*	*	*	445	44,300	10.0	(3.1 - 16.8)
Education								
≤ High School	163	225,100	17.8	(9.0 - 26.7)	12,417	493,000	17.3	(13.3 - 21.3)
Some College	*	*	*	*	7,232	299,400	18.7	(14.6 - 22.7)
≥ College Degree	211	188,900	16.7	(10.7 - 22.7)	9,086	441,500	15.4	(12.9 - 17.9)
Income								
≤ \$24,999	*	*	*	*	7,824	423,900	25.0	(18.5 - 31.6)
\$25,000-\$49,999	72	64,000	14.4	(5.1 - 23.6)	7,445	223,600	16.0	(12.8 - 19.2)
\$50,000-\$74,999	*	*	*	*	4,141	151,200	14.6	(10.8 - 18.4)
≥ \$75,000	134	109,900	13.7	(7.0 - 20.5)	5,874	287,100	12.5	(10.0 - 15.1)

¹ Defined as respondent unintentionally coming to rest on the ground or another lower level one or more times within the past 3 months

² Sample size from survey

³ Estimated number of adults who have had a fall within the past 3 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 69b. Had a fall within the past 3 months among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	456	19.5	(14.3 - 24.7)	28,783	17.3	(15.0 - 19.5)
Gender						
Male	184	16.0	(9.4 - 22.6)	10,966	15.0	(12.1 - 17.8)
Female	272	22.8	(15.0 - 30.6)	17,817	19.4	(16.0 - 22.8)
Race/Ethnicity						
White Non-Hispanic	305	20.4	(14.7 - 26.2)	27,456	16.9	(15.2 - 18.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	424	10.7	(4.2 - 17.1)
Other Non-Hispanic	*	*	*	445	10.0	(2.9 - 17.0)
Education						
≤ High School	163	17.8	(8.8 - 26.9)	12,417	17.4	(13.0 - 21.7)
Some College	*	*	*	7,232	19.0	(14.7 - 23.2)
≥ College Degree	211	17.8	(11.3 - 24.3)	9,086	16.0	(13.1 - 18.8)
Income						
≤ \$24,999	*	*	*	7,824	25.3	(18.3 - 32.4)
\$25,000-\$49,999	72	14.0	(5.3 - 22.8)	7,445	15.9	(12.6 - 19.2)
\$50,000-\$74,999	*	*	*	4,141	14.7	(10.6 - 18.8)
≥ \$75,000	134	14.3	(7.0 - 21.6)	5,874	12.9	(9.9 - 15.8)

¹ Defined as respondent unintentionally coming to rest on the ground or another lower level one or more times within the past 3 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 70a. Had a fall that caused injury within the past 3 months among adults¹ in New York City and New York State

	New York City				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	456	132,200	4.6	(2.4 - 6.8)	28,760	335,700	4.6	(3.7 - 5.5)
Gender								
Male	*	*	*	*	10,957	120,000	3.5	(2.5 - 4.6)
Female	272	88,000	5.6	(2.3 - 9.0)	17,803	215,600	5.5	(4.1 - 6.9)
Age (years)								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	7,737	115,100	4.1	(2.9 - 5.4)
55-64	*	*	*	*	8,449	90,300	4.4	(2.7 - 6.1)
≥ 65	186	58,600	5.9	(1.6 - 10.1)	12,574	130,300	5.2	(3.4 - 6.9)
Race/Ethnicity								
White Non-Hispanic	305	90,800	6.8	(3.3 - 10.4)	27,433	270,800	5.2	(4.2 - 6.1)
Black Non-Hispanic	*	*	*	*	458	47,900	4.9	(0.9 - 9.0)
Hispanic	*	*	*	*	424	12,400	1.8	(0.2 - 3.3)
Other Non-Hispanic	*	*	*	*	445	4,600	1.0	(0.4 - 1.6)
Education								
≤ High School	163	65,000	5.1	(1.7 - 8.6)	12,404	148,900	5.2	(3.6 - 6.8)
Some College	*	*	*	*	7,228	85,400	5.3	(3.1 - 7.5)
≥ College Degree	211	38,700	3.4	(0.9 - 6.0)	9,081	101,300	3.5	(2.4 - 4.6)
Income								
≤ \$24,999	*	*	*	*	7,812	90,200	5.3	(3.6 - 7.1)
\$25,000-\$49,999	*	*	*	*	7,442	93,200	6.7	(3.9 - 9.4)
\$50,000-\$74,999	*	*	*	*	4,139	37,000	3.6	(1.8 - 5.3)
≥ \$75,000	*	*	*	*	5,872	54,700	2.4	(1.5 - 3.3)

¹ Defined as as respondent unintentionally coming to rest on the ground or another lower level and caused respondent to limit regular activities for at least a day or to go see a doctor one or more times within the past 3 months

² Sample size from survey

³ Estimated number of adults who have had a fall that caused an injury within the past 3 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
New York City**

Table 70b. Had a fall that caused injury within the past 3 months among adults¹ in New York City and New York State - Age-Adjusted²

	New York City			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	456	4.5	(2.4 - 6.6)	28,760	4.6	(3.6 - 5.5)
Gender						
Male	*	*	*	10,957	3.6	(2.4 - 4.7)
Female	272	5.6	(2.3 - 8.9)	17,803	5.5	(4.1 - 6.9)
Race/Ethnicity						
White Non-Hispanic	305	6.8	(3.3 - 10.2)	27,433	5.2	(4.2 - 6.2)
Black Non-Hispanic	*	*	*	458	4.7	(0.7 - 8.7)
Hispanic	*	*	*	424	1.8	(0.1 - 3.4)
Other Non-Hispanic	*	*	*	445	1.0	(0.4 - 1.6)
Education						
≤ High School	163	5.0	(1.6 - 8.3)	12,404	5.2	(3.5 - 6.8)
Some College	*	*	*	7,228	5.3	(3.1 - 7.5)
≥ College Degree	211	3.5	(1.0 - 6.0)	9,081	3.6	(2.4 - 4.7)
Income						
≤ \$24,999	*	*	*	7,812	5.1	(3.3 - 7.0)
\$25,000-\$49,999	*	*	*	7,442	6.6	(3.9 - 9.2)
\$50,000-\$74,999	*	*	*	4,139	3.7	(1.7 - 5.8)
≥ \$75,000	*	*	*	5,872	2.4	(1.4 - 3.3)

¹ Defined as as respondent unintentionally coming to rest on the ground or another lower level and caused respondent to limit regular activities for at least a day or to go see a doctor one or more times within the past 3 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)