

New York State Expanded Behavioral Risk Factor Surveillance System

Oswego County

**Final Report
July 2008 - June 2009**

**New York State Department of Health
December 2009**

Contents

Introduction	12
Table 1. Oswego County and New York State adult US Census Bureau population estimates, 2007	13
General Health	
Table 2a. Fair or poor general health among adults in Oswego County and New York State	14
Table 2b. Fair or poor general health among adults in Oswego County and New York State - Age-adjusted	15
Table 3a. Poor physical health 14 or more days within the past month among adults in Oswego County and New York State	16
Table 3b. Poor physical health 14 or more days within the past month among adults in Oswego County and New York State - Age-adjusted	17
Table 4a. Poor mental health 14 or more days within the past month among adults in Oswego County and New York State	18
Table 4b. Poor mental health 14 or more days within the past month among adults in Oswego County and New York State - Age-adjusted	19
Table 5a. Poor physical or mental health 14 or more days within the past month among adults in Oswego County and New York State	20
Table 5b. Poor physical or mental health 14 or more days within the past month among adults in Oswego County and New York State - Age-adjusted	21
Quality Health Care	
Table 6a. Health insurance among adults in Oswego County and New York State	22
Table 6b. Health insurance among adults in Oswego County and New York State - Age-adjusted	23
Table 7a. Health insurance among adults aged 18-64 in Oswego County and New York State	24
Table 7b. Health insurance among adults aged 18-64 in Oswego County and New York State - Age-adjusted	25

Contents

Table 8a. No health insurance among adults in Oswego County and New York State	26
Table 8b. No health insurance among adults in Oswego County and New York State - Age-adjusted	27
Table 9a. No health insurance among adults aged 18-64 in Oswego County and New York State	28
Table 9b. No health insurance among adults aged 18-64 in Oswego County and New York State - Age-adjusted	29
Table 10a. Adults with regular health care provider in Oswego County and New York State	30
Table 10b. Adults with regular health care provider in Oswego County and New York State - Age-adjusted	31
Table 11a. Cost prevented visit to doctor within the past year among adults in Oswego County and New York State	32
Table 11b. Cost prevented visit to doctor within the past year among adults in Oswego County and New York State - Age-adjusted	33
Table 12a. Visited a doctor for routine checkup within the past year among adults in Oswego County and New York State	34
Table 12b. Visited a doctor for routine checkup within the past year among adults in Oswego County and New York State - Age-adjusted	35
Table 13a. Visited a doctor for routine checkup within the past 2 years among adults in Oswego County and New York State	36
Table 13b. Visited a doctor for routine checkup within the past 2 years among adults in Oswego County and New York State - Age-adjusted	37
Table 14a. Dentist visit within the past year among adults in Oswego County and New York State	38
Table 14b. Dentist visit within the past year among adults in Oswego County and New York State - Age-adjusted	39
Table 15a. Had teeth cleaned within the past year among adults in Oswego County and New York State	40
Table 15b. Had teeth cleaned within the past year among adults in Oswego County and New York State - Age-adjusted	41

Contents

Table 16a. Permanent teeth extracted due to decay or gum disease among adults in Oswego County and New York State	42
Table 16b. Permanent teeth extracted due to decay or gum disease among adults in Oswego County and New York State - Age-adjusted	43
Table 17a. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older in Oswego County and New York State	44
Table 17b. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older in Oswego County and New York State - Age-adjusted	45
Table 18a. Flu shot within the past 12 months in Oswego County and New York State	46
Table 18b. Flu shot within the past 12 months in Oswego County and New York State - Age-adjusted	47
Table 19a. Flu vaccine that was sprayed in nose within the past 12 months in Oswego County and New York State	48
Table 19b. Flu vaccine that was sprayed in nose within the past 12 months in Oswego County and New York State - Age-adjusted	49
Table 20a. Flu shot or flu vaccine sprayed in nose within the past 12 months in Oswego County and New York State	50
Table 20b. Flu shot or flu vaccine sprayed in nose within the past 12 months in Oswego County and New York State - Age-adjusted	51
Table 21a. Flu shot among adults aged 65 and older in Oswego County and New York State	52
Table 21b. Flu shot among adults aged 65 and older in Oswego County and New York State - Age-adjusted	53
Table 22a. Pneumonia shot or pneumococcal vaccine in Oswego County and New York State	54
Table 22b. Pneumonia shot or pneumococcal vaccine in Oswego County and New York State - Age-adjusted	55
Table 23a. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older in Oswego County and New York State	56
Table 23b. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older in Oswego County and New York State - Age-adjusted	57

Contents

Cancer Screening

Table 24a. Ever had a mammogram among women aged 40 and older in Oswego County and New York State	58
Table 24b. Ever had a mammogram among women aged 40 and older in Oswego County and New York State - Age-adjusted	59
Table 25a. Had mammogram within the past 2 years among women aged 40 and older in Oswego County and New York State	60
Table 25b. Had mammogram within the past 2 years among women aged 40 and older in Oswego County and New York State - Age-adjusted	61
Table 26a. Had mammogram within the past 2 years among women aged 50 and older in Oswego County and New York State	62
Table 26b. Had mammogram within the past 2 years among women aged 50 and older in Oswego County and New York State - Age-adjusted	63
Table 27a. Ever had a Pap test among women in Oswego County and New York State	64
Table 27b. Ever had a Pap test among women in Oswego County and New York State - Age-adjusted	65
Table 28a. Had a Pap test within the past 3 years among women in Oswego County and New York State	66
Table 28b. Had a Pap test within the past 3 years among women in Oswego County and New York State - Age-adjusted	67
Table 29a. Ever had a digital rectal exam among men aged 40 and older in Oswego County and New York State	68
Table 29b. Ever had a digital rectal exam among men aged 40 and older in Oswego County and New York State - Age-adjusted	69
Table 30a. Had digital rectal exam within the past 2 years among men aged 40 and older in Oswego County and New York State	70
Table 30b. Had digital rectal exam within the past 2 years among men aged 40 and older in Oswego County and New York State - Age-adjusted	71

Contents

Table 31a. Ever had a prostate specific antigen test among men aged 40 and older in Oswego County and New York State	72
Table 31b. Ever had a prostate specific antigen test among men aged 40 and older in Oswego County and New York State - Age-adjusted	73
Table 32a. Had prostate specific antigen test within the past 2 years among men aged 40 and older in Oswego County and New York State	74
Table 32b. Had prostate specific antigen test within the past 2 years among men aged 40 and older in Oswego County and New York State - Age-adjusted	75
Table 33a. Prostate cancer among men aged 40 and older in Oswego County and New York State	76
Table 33b. Prostate cancer among men aged 40 and older in Oswego County and New York State - Age-adjusted	77
Table 34a. Home blood stool test ever used among adults aged 50 and older in Oswego County and New York State	78
Table 34b. Home blood stool test ever used among adults aged 50 and older in Oswego County and New York State - Age-adjusted	79
Table 35a. Home blood stool test used within the past year among adults aged 50 and older in Oswego County and New York State	80
Table 35b. Home blood stool test used within the past year among adults aged 50 and older in Oswego County and New York State - Age-adjusted	81
Table 36a. Home blood stool test used within the past 2 years among adults aged 50 and older in Oswego County and New York State	82
Table 36b. Home blood stool test used within the past 2 years among adults aged 50 and older in Oswego County and New York State - Age-adjusted	83
Table 37a. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in Oswego County and New York State	84
Table 37b. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in Oswego County and New York State - Age-adjusted	85
Table 38a. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Oswego County and New York State	86

Contents

Table 38b. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Oswego County and New York State - Age-adjusted	87
Table 39a. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Oswego County and New York State	88
Table 39b. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Oswego County and New York State - Age-adjusted	89
Chronic Diseases	
Table 40a. High blood pressure among adults in Oswego County and New York State	90
Table 40b. High blood pressure among adults in Oswego County and New York State - Age-adjusted	91
Table 41a. Blood pressure medication among adults with high blood pressure in Oswego County and New York State	92
Table 41b. Blood pressure medication among adults with high blood pressure in Oswego County and New York State - Age-adjusted	93
Table 42a. Ever had blood cholesterol checked among adults in Oswego County and New York State	94
Table 42b. Ever had blood cholesterol checked among adults in Oswego County and New York State - Age-adjusted	95
Table 43a. Blood cholesterol checked within the past 5 years among adults in Oswego County and New York State	96
Table 43b. Blood cholesterol checked within the past 5 years among adults in Oswego County and New York State - Age-adjusted	97
Table 44a. Cardiovascular disease among adults in Oswego County and New York State	98
Table 44b. Cardiovascular disease among adults in Oswego County and New York State - Age-adjusted	99
Table 45a. Coronary heart disease among adults in Oswego County and New York State	100

Contents

Table 45b. Coronary heart disease among adults in Oswego County and New York State - Age-adjusted	101
Table 46a. Ever diagnosed with diabetes among adults in Oswego County and New York State	102
Table 46b. Ever diagnosed with diabetes among adults in Oswego County and New York State - Age-adjusted	103
Table 47a. Ever diagnosed with asthma (lifetime) among adults in Oswego County and New York State	104
Table 47b. Ever diagnosed with asthma (lifetime) among adults in Oswego County and New York State - Age-adjusted	105
Table 48a. Current asthma among adults in Oswego County and New York State	106
Table 48b. Current asthma among adults in Oswego County and New York State - Age-adjusted	107
Table 49a. Chronic joint symptoms among adults in Oswego County and New York State	108
Table 49b. Chronic joint symptoms among adults in Oswego County and New York State - Age-adjusted	109
Table 50a. Arthritis among adults in Oswego County and New York State	110
Table 50b. Arthritis among adults in Oswego County and New York State - Age-adjusted	111
Table 51a. Activity limitations because of physical, mental or emotional problems among adults in Oswego County and New York State	112
Table 51b. Activity limitations because of physical, mental or emotional problems among adults in Oswego County and New York State - Age-adjusted	113
Table 52a. Have health problem that requires the use of special equipment among adults in Oswego County and New York State	114
Table 52b. Have health problem that requires the use of special equipment among adults in Oswego County and New York State - Age-adjusted	115
Table 53a. Disability among adults in Oswego County and New York State	116
Table 53b. Disability among adults in Oswego County and New York State - Age-adjusted	117

Contents

Health Behaviors

Table 54a. Current smoking among adults in Oswego County and New York State	118
Table 54b. Current smoking among adults in Oswego County and New York State - Age-adjusted	119
Table 55a. Everyday smoking among adults in Oswego County and New York State	120
Table 55b. Everyday smoking among adults in Oswego County and New York State - Age-adjusted	121
Table 56a. Adults living in homes in which smoking is prohibited in Oswego County and New York State	122
Table 56b. Adults living in homes in which smoking is prohibited in Oswego County and New York State - Age-adjusted	123
Table 57a. Binge drinking within the past month among adults in Oswego County and New York State	124
Table 57b. Binge drinking within the past month among adults in Oswego County and New York State - Age-adjusted	125
Table 58a. Heavy drinking within the past month among adults in Oswego County and New York State	126
Table 58b. Heavy drinking within the past month among adults in Oswego County and New York State - Age-adjusted	127
Table 59a. Never or rarely been asked about sexual history during routine check-up among adults in Oswego County and New York State	128
Table 59b. Never or rarely been asked about sexual history during routine check-up among adults in Oswego County and New York State - Age-adjusted	129
Table 60a. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults in Oswego County and New York State	130
Table 60b. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults in Oswego County and New York State - Age-adjusted	131

Contents

Table 61a. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults in Oswego County and New York State	132
Table 61b. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults in Oswego County and New York State - Age-adjusted	133

Overweight / Obesity

Table 62a. Overweight among adults in Oswego County and New York State	134
Table 62b. Overweight among adults in Oswego County and New York State - Age-adjusted	135
Table 63a. Obesity among adults in Oswego County and New York State	136
Table 63b. Obesity among adults in Oswego County and New York State - Age-adjusted	137
Table 64a. Overweight or obesity among adults in Oswego County and New York State	138
Table 64b. Overweight or obesity among adults in Oswego County and New York State - Age-adjusted	139
Table 65a. Received advice about weight by a health professional among adults in Oswego County and New York State	140
Table 65b. Received advice about weight by a health professional among adults in Oswego County and New York State - Age-adjusted	141
Table 66a. Received advice to lose weight by a health professional among adults provided weight advice in Oswego County and New York State	142
Table 66b. Received advice to lose weight by a health professional among adults provided weight advice in Oswego County and New York State - Age-adjusted	143

Physical Activity / Nutrition

Table 67a. No leisure-time physical activity among adults in Oswego County and New York State	144
Table 67b. No leisure-time physical activity among adults in Oswego County and New York State - Age-adjusted	145

Contents

Table 68a. Consumption of 5 or more servings of fruits and vegetables per day among adults in Oswego County and New York State	146
Table 68b. Consumption of 5 or more servings of fruits and vegetables per day among adults in Oswego County and New York State - Age-adjusted	147

Injury

Table 69a. Had a fall within the past 3 months among adults in Oswego County and New York State	148
Table 69b. Had a fall within the past 3 months among adults in Oswego County and New York State - Age-adjusted	149
Table 70a. Had a fall that caused injury within the past 3 months among adults in Oswego County and New York State	150
Table 70b. Had a fall that caused injury within the past 3 months among adults in Oswego County and New York State - Age-adjusted	151

Introduction

The New York State Expanded Behavioral Risk Factor Surveillance System (Expanded BRFSS) is a random digit dialed telephone survey of adults (18 years of age and older) that is representative of the non-institutionalized civilian population with landline telephones living in New York State. The goal of the Expanded BRFSS is to collect county-specific data on preventive health practices, risk behaviors, injuries and preventable chronic and infectious diseases. Topics assessed by the Expanded BRFSS include tobacco use, physical inactivity, diet, use of cancer screening services and other factors linked to the leading causes of morbidity and mortality. For each topic, health indicators are based on answers to one or more questions in the survey. The survey results are weighted to more accurately represent the adult population living in each county. Weighting procedures adjust for the differential selection probabilities by race, age and gender. More detailed information on weighting procedures is available in the technical brief.

Survey data collection occurred in 58 geographic areas of New York State, including each of the 57 counties outside of New York City and New York City (5 counties) as a single area. A standard questionnaire was used in all areas. Data collection for the 2009 Expanded BRFSS took place during the period of July 2008 through June 2009. The goal was to complete about 650 interviews in each area. The final number of interviews varies because of the sampling design of the survey and differential response rates. More detailed information on the sampling design is available in the technical brief.

The New York State Department of Health contracted with RTI International, which is responsible for the data collection, sampling, weighting and creation of the statistical reports. RTI International subcontracted with Clearwater Research Inc. for data collection and additional expertise in sample design and data quality control.

This report presents both crude estimates and estimates that have been age-adjusted to the Year 2000 US Standard Population when they represent the entire adult population of an area. Estimates for specific age groups are not age-adjusted. Age adjustment is a statistical process applied to rates that allows communities with different age structures to be compared. Counties can compare their data with results from New York State as a whole and with other counties. Confidence intervals at the 95% level associated with each health indicator give an estimated range of values which is likely to include the actual prevalence of the indicator. Confidence intervals are similar to margins of error. When the confidence intervals of two estimates of the same indicator from different areas do not overlap, they may be said to be statistically significantly different, i.e., these differences are unlikely related to chance and are considered true differences. If there is any value that is included in both intervals, the two estimates are not statistically significantly different.

The survey results for each of the 69 health indicators included in the survey are typically provided by gender, age, race and ethnicity, education and income in the Expanded BRFSS county reports. The estimates for some of these groupings may be suppressed in geographic areas with low proportions of adult respondents in these groupings.

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 1. Oswego County and New York State adult US Census Bureau population estimates,¹ 2007

	Oswego County		New York State	
	Adult Population ¹	Percent	Adult Population ¹	Percent
Total	94,009	100	14,884,315	100
Gender				
Male	46,064	49.0	7,103,791	47.7
Female	47,945	51.0	7,780,524	52.3
Age (years)				
18-34	30,583	32.5	4,497,479	30.2
35-44	16,665	17.7	2,844,115	19.1
45-54	18,704	19.9	2,854,316	19.2
55-64	13,719	14.6	2,142,000	14.4
≥ 65	14,338	15.3	2,546,405	17.1
Race/Ethnicity				
White Non-Hispanic	90,920	96.7	9,343,884	62.8
Black Non-Hispanic	817	0.9	2,156,941	14.5
Hispanic	1,297	1.4	2,258,717	15.2
Other Non-Hispanic	975	1.0	1,124,773	7.6

¹ Source: National Center for Health Statistics. Postcensal estimates of the resident population of the United States for July 1, 2000-July 1, 2007, by year, county, age, bridged race, Hispanic origin, and sex (Vintage 2007). Prepared under a collaborative arrangement with the U.S. Census Bureau; released August 7, 2008. Available from: http://www.cdc.gov/nchs/nvss/bridged_race.htm as of November 25, 2009.

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 2a. Fair or poor general health among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	665	14,700	15.7	(12.6 - 18.8)	38,124	2,531,400	17.1	(15.1 - 19.0)
Gender								
Male	252	7,900	17.2	(12.1 - 22.3)	14,599	1,069,700	15.1	(12.4 - 17.8)
Female	413	6,800	14.2	(10.6 - 17.9)	23,525	1,461,700	18.8	(16.1 - 21.6)
Age (years)								
18-34	*	*	*	*	3,940	535,800	12.0	(7.2 - 16.8)
35-44	79	2,900	17.5	(8.3 - 26.8)	4,985	289,200	10.0	(6.8 - 13.3)
45-54	159	2,800	15.0	(9.0 - 21.0)	7,819	507,800	17.9	(14.0 - 21.8)
55-64	129	3,100	22.8	(14.8 - 30.8)	8,757	424,700	19.9	(16.0 - 23.9)
≥ 65	206	4,200	29.8	(22.5 - 37.0)	12,623	773,900	30.5	(26.8 - 34.2)
Race/Ethnicity								
White Non-Hispanic	646	14,500	16.1	(12.9 - 19.2)	35,811	1,226,600	13.2	(11.7 - 14.6)
Black Non-Hispanic	*	*	*	*	739	457,300	21.3	(14.8 - 27.9)
Hispanic	*	*	*	*	861	684,100	30.4	(22.2 - 38.6)
Other Non-Hispanic	*	*	*	*	713	163,400	14.5	(5.1 - 24.0)
Education								
≤ High School	334	9,400	20.8	(15.6 - 26.0)	15,682	1,466,500	26.3	(22.6 - 30.0)
Some College	155	3,800	16.3	(9.9 - 22.7)	9,904	508,500	14.3	(10.9 - 17.8)
≥ College Degree	173	1,500	6.0	(2.3 - 9.8)	12,457	493,500	8.8	(6.5 - 11.1)
Income								
≤ \$24,999	175	5,300	26.9	(18.9 - 34.9)	9,633	1,269,200	36.3	(30.1 - 42.4)
\$25,000-\$49,999	175	4,200	17.9	(11.6 - 24.1)	9,649	447,600	15.4	(11.9 - 18.9)
\$50,000-\$74,999	*	*	*	*	5,769	237,800	12.1	(8.0 - 16.2)
≥ \$75,000	*	*	*	*	8,562	218,300	4.9	(3.6 - 6.1)

¹ Defined as reporting fair or poor general health

² Sample size from survey

³ Estimated number of adults who have fair or poor health (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 2b. Fair or poor general health among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	665	16.1	(12.8 - 19.4)	38,124	16.7	(14.8 - 18.6)
Gender						
Male	252	18.3	(12.8 - 23.8)	14,599	15.3	(12.6 - 18.0)
Female	413	14.0	(10.3 - 17.7)	23,525	18.0	(15.2 - 20.7)
Race/Ethnicity						
White Non-Hispanic	646	16.5	(13.1 - 19.8)	35,811	12.9	(11.4 - 14.4)
Black Non-Hispanic	*	*	*	739	20.8	(14.2 - 27.3)
Hispanic	*	*	*	861	29.9	(21.8 - 37.9)
Other Non-Hispanic	*	*	*	713	13.8	(4.8 - 22.8)
Education						
≤ High School	334	21.4	(15.9 - 26.9)	15,682	25.8	(22.1 - 29.5)
Some College	155	16.4	(10.0 - 22.7)	9,904	14.1	(10.7 - 17.6)
≥ College Degree	173	6.2	(2.2 - 10.1)	12,457	8.5	(6.3 - 10.8)
Income						
≤ \$24,999	175	26.2	(18.3 - 34.1)	9,633	35.9	(29.7 - 42.0)
\$25,000-\$49,999	175	18.8	(12.2 - 25.5)	9,649	15.0	(11.6 - 18.5)
\$50,000-\$74,999	*	*	*	5,769	11.8	(7.7 - 16.0)
≥ \$75,000	*	*	*	8,562	4.7	(3.5 - 6.0)

¹ Defined as reporting fair or poor general health

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 3a. Poor physical health 14 or more days within the past month among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	653	10,900	11.8	(9.1 - 14.4)	37,327	1,472,900	10.1	(8.7 - 11.6)
Gender								
Male	250	5,200	11.5	(7.5 - 15.5)	14,325	604,200	8.7	(7.0 - 10.5)
Female	403	5,700	12.0	(8.5 - 15.6)	23,002	868,600	11.4	(9.2 - 13.6)
Age (years)								
18-34	*	*	*	*	3,891	268,100	6.1	(2.7 - 9.5)
35-44	79	2,300	13.7	(5.8 - 21.7)	4,948	232,100	8.1	(5.2 - 11.1)
45-54	159	1,900	10.1	(5.3 - 14.9)	7,764	350,900	12.5	(9.4 - 15.6)
55-64	125	2,400	17.7	(10.0 - 25.3)	8,621	271,200	13.0	(10.3 - 15.8)
≥ 65	198	2,600	19.6	(13.3 - 26.0)	12,103	350,600	14.5	(12.1 - 16.9)
Race/Ethnicity								
White Non-Hispanic	633	10,700	12.0	(9.3 - 14.7)	35,073	929,700	10.2	(8.8 - 11.6)
Black Non-Hispanic	*	*	*	*	724	208,200	9.9	(4.8 - 14.9)
Hispanic	*	*	*	*	835	250,300	11.5	(6.5 - 16.4)
Other Non-Hispanic	*	*	*	*	695	84,600	7.6	(1.3 - 13.9)
Education								
≤ High School	324	6,100	13.7	(9.5 - 17.9)	15,178	692,800	12.9	(10.6 - 15.3)
Some College	155	2,800	12.2	(7.0 - 17.4)	9,751	362,500	10.3	(7.4 - 13.2)
≥ College Degree	171	2,000	8.1	(3.4 - 12.8)	12,327	359,300	6.4	(4.8 - 8.1)
Income								
≤ \$24,999	172	4,900	25.3	(17.2 - 33.3)	9,294	625,600	18.4	(13.6 - 23.3)
\$25,000-\$49,999	169	2,500	10.6	(5.5 - 15.7)	9,495	279,100	9.8	(7.3 - 12.3)
\$50,000-\$74,999	*	*	*	*	5,733	186,400	9.5	(5.8 - 13.2)
≥ \$75,000	*	*	*	*	8,510	219,100	4.9	(3.6 - 6.2)

¹ Defined as 14 or more days within the past month when physical health was not good

² Sample size from survey

³ Estimated number of adults with poor physical health 14 or more days within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 3b. Poor physical health 14 or more days within the past month among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	653	12.0	(9.2 - 14.8)	37,327	9.9	(8.4 - 11.4)
Gender						
Male	250	11.9	(7.6 - 16.2)	14,325	8.7	(6.9 - 10.5)
Female	403	12.1	(8.4 - 15.7)	23,002	11.0	(8.7 - 13.2)
Race/Ethnicity						
White Non-Hispanic	633	12.2	(9.3 - 15.1)	35,073	9.9	(8.5 - 11.3)
Black Non-Hispanic	*	*	*	724	9.7	(4.6 - 14.9)
Hispanic	*	*	*	835	11.3	(6.3 - 16.2)
Other Non-Hispanic	*	*	*	695	7.6	(1.2 - 14.0)
Education						
≤ High School	324	14.1	(9.6 - 18.5)	15,178	12.7	(10.2 - 15.1)
Some College	155	12.3	(7.0 - 17.6)	9,751	10.1	(7.2 - 12.9)
≥ College Degree	171	8.1	(3.3 - 12.9)	12,327	6.3	(4.7 - 8.0)
Income						
≤ \$24,999	172	25.2	(17.1 - 33.3)	9,294	18.4	(13.5 - 23.4)
\$25,000-\$49,999	169	11.2	(5.7 - 16.7)	9,495	9.4	(7.0 - 11.9)
\$50,000-\$74,999	*	*	*	5,733	9.0	(5.3 - 12.6)
≥ \$75,000	*	*	*	8,510	4.9	(3.5 - 6.2)

¹ Defined as 14 or more days within the past month when physical health was not good

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 4a. Poor mental health 14 or more days within the past month among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	656	11,700	12.7	(9.4 - 16.0)	37,528	1,473,200	10.1	(8.6 - 11.5)
Gender								
Male	252	4,800	10.6	(5.7 - 15.4)	14,407	555,200	7.9	(6.3 - 9.6)
Female	404	6,900	14.7	(10.4 - 19.1)	23,121	918,000	12.0	(9.7 - 14.3)
Age (years)								
18-34	90	4,100	13.8	(6.3 - 21.2)	3,889	496,700	11.2	(7.4 - 15.1)
35-44	78	2,800	17.3	(8.8 - 25.8)	4,947	254,100	8.9	(6.7 - 11.1)
45-54	158	2,200	11.9	(6.2 - 17.6)	7,749	347,200	12.3	(9.0 - 15.6)
55-64	126	1,800	13.1	(6.4 - 19.7)	8,623	211,300	10.1	(7.9 - 12.3)
≥ 65	204	800	5.7	(2.4 - 9.0)	12,320	163,900	6.7	(4.7 - 8.7)
Race/Ethnicity								
White Non-Hispanic	636	11,600	13.0	(9.6 - 16.3)	35,266	899,200	9.8	(8.4 - 11.2)
Black Non-Hispanic	*	*	*	*	722	227,800	10.9	(5.4 - 16.4)
Hispanic	*	*	*	*	845	263,700	11.8	(6.8 - 16.8)
Other Non-Hispanic	*	*	*	*	695	82,500	7.4	(1.4 - 13.5)
Education								
≤ High School	325	6,200	14.0	(9.1 - 18.9)	15,335	665,000	12.2	(10.0 - 14.5)
Some College	154	3,200	14.1	(6.8 - 21.4)	9,771	404,500	11.5	(8.3 - 14.8)
≥ College Degree	174	2,300	9.3	(4.1 - 14.4)	12,342	346,600	6.2	(4.5 - 7.9)
Income								
≤ \$24,999	175	5,000	25.8	(16.6 - 34.9)	9,398	611,500	17.9	(13.0 - 22.8)
\$25,000-\$49,999	172	2,000	8.6	(3.3 - 13.8)	9,547	260,300	9.0	(6.7 - 11.4)
\$50,000-\$74,999	99	1,600	9.9	(2.8 - 16.9)	5,726	187,300	9.6	(6.0 - 13.1)
≥ \$75,000	*	*	*	*	8,520	276,500	6.2	(4.6 - 7.7)

¹ Defined as 14 or more days within the past month when mental health was not good

² Sample size from survey

³ Estimated number of adults with poor mental health 14 or more days within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 4b. Poor mental health 14 or more days within the past month among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	656	12.7	(9.4 - 16.0)	37,528	10.2	(8.7 - 11.7)
Gender						
Male	252	10.2	(5.5 - 14.9)	14,407	7.9	(6.3 - 9.6)
Female	404	15.1	(10.6 - 19.7)	23,121	12.2	(9.8 - 14.6)
Race/Ethnicity						
White Non-Hispanic	636	12.9	(9.6 - 16.3)	35,266	10.0	(8.5 - 11.4)
Black Non-Hispanic	*	*	*	722	11.1	(5.5 - 16.7)
Hispanic	*	*	*	845	11.5	(6.5 - 16.4)
Other Non-Hispanic	*	*	*	695	7.6	(1.5 - 13.7)
Education						
≤ High School	325	14.2	(9.2 - 19.2)	15,335	12.3	(10.0 - 14.6)
Some College	154	13.7	(6.6 - 20.8)	9,771	11.8	(8.5 - 15.1)
≥ College Degree	174	9.3	(4.1 - 14.4)	12,342	6.3	(4.6 - 8.0)
Income						
≤ \$24,999	175	26.1	(16.7 - 35.4)	9,398	18.2	(13.2 - 23.1)
\$25,000-\$49,999	172	8.8	(3.3 - 14.3)	9,547	9.4	(6.9 - 11.9)
\$50,000-\$74,999	99	9.3	(2.6 - 16.0)	5,726	9.5	(5.9 - 13.1)
≥ \$75,000	*	*	*	8,520	6.2	(4.6 - 7.7)

¹ Defined as 14 or more days within the past month when mental health was not good

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 5a. Poor physical or mental health 14 or more days within the past month among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	647	19,200	21.0	(17.2 - 24.9)	37,005	2,385,900	16.5	(14.8 - 18.2)
Gender								
Male	248	8,900	19.8	(13.9 - 25.8)	14,231	937,700	13.6	(11.4 - 15.8)
Female	399	10,300	22.2	(17.2 - 27.1)	22,774	1,448,200	19.1	(16.5 - 21.7)
Age (years)								
18-34	90	5,100	17.2	(9.2 - 25.2)	3,859	602,600	13.8	(9.7 - 17.9)
35-44	*	*	*	*	4,924	388,300	13.7	(10.4 - 17.0)
45-54	158	3,300	17.9	(11.5 - 24.3)	7,722	551,000	19.6	(15.8 - 23.4)
55-64	123	3,400	26.2	(17.3 - 35.2)	8,545	395,500	19.2	(16.1 - 22.3)
≥ 65	198	3,200	23.9	(17.1 - 30.6)	11,955	448,500	18.8	(15.9 - 21.7)
Race/Ethnicity								
White Non-Hispanic	627	18,900	21.4	(17.5 - 25.3)	34,778	1,485,700	16.3	(14.8 - 17.8)
Black Non-Hispanic	*	*	*	*	715	343,700	16.5	(10.3 - 22.7)
Hispanic	*	*	*	*	832	412,900	19.0	(12.6 - 25.3)
Other Non-Hispanic	*	*	*	*	680	143,500	13.0	(5.0 - 21.0)
Education								
≤ High School	320	10,300	23.4	(17.6 - 29.3)	15,018	1,077,200	20.3	(17.3 - 23.2)
Some College	153	5,300	23.7	(15.5 - 31.9)	9,672	615,500	17.6	(14.1 - 21.1)
≥ College Degree	171	3,600	14.6	(8.3 - 20.9)	12,244	634,200	11.4	(9.2 - 13.6)
Income								
≤ \$24,999	*	*	*	*	9,199	924,000	27.3	(21.9 - 32.8)
\$25,000-\$49,999	168	4,400	19.1	(11.9 - 26.2)	9,425	456,000	16.1	(13.0 - 19.2)
\$50,000-\$74,999	99	2,600	16.6	(8.1 - 25.2)	5,696	312,000	16.0	(11.4 - 20.5)
≥ \$75,000	*	*	*	*	8,475	447,600	10.0	(8.1 - 12.0)

¹ Defined as 14 or more days within the past month when physical or mental health prevented usual activities, such as self-care, work or recreation

² Sample size from survey

³ Estimated number of adults with poor physical or mental health 14 or more days within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 5b. Poor physical or mental health 14 or more days within the past month among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	647	21.2	(17.3 - 25.2)	37,005	16.4	(14.7 - 18.2)
Gender						
Male	248	19.8	(13.9 - 25.8)	14,231	13.6	(11.4 - 15.8)
Female	399	22.5	(17.4 - 27.6)	22,774	19.0	(16.3 - 21.6)
Race/Ethnicity						
White Non-Hispanic	627	21.6	(17.5 - 25.6)	34,778	16.3	(14.7 - 17.8)
Black Non-Hispanic	*	*	*	715	16.7	(10.3 - 23.0)
Hispanic	*	*	*	832	18.5	(12.2 - 24.8)
Other Non-Hispanic	*	*	*	680	13.1	(5.0 - 21.3)
Education						
≤ High School	320	24.0	(17.9 - 30.1)	15,018	20.1	(17.1 - 23.1)
Some College	153	23.5	(15.4 - 31.7)	9,672	17.6	(14.1 - 21.1)
≥ College Degree	171	14.4	(8.1 - 20.7)	12,244	11.4	(9.2 - 13.7)
Income						
≤ \$24,999	*	*	*	9,199	27.6	(22.1 - 33.1)
\$25,000-\$49,999	168	20.0	(12.5 - 27.6)	9,425	16.1	(12.9 - 19.3)
\$50,000-\$74,999	99	16.1	(7.7 - 24.5)	5,696	15.6	(11.0 - 20.1)
≥ \$75,000	*	*	*	8,475	10.0	(8.0 - 12.0)

¹ Defined as 14 or more days within the past month when physical or mental health prevented usual activities, such as self-care, work or recreation

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 6a. Health insurance among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	668	80,000	85.7	(81.1 - 90.4)	38,119	12,854,200	86.6	(84.2 - 89.0)
Gender								
Male	254	37,000	81.4	(72.8 - 90.0)	14,597	5,881,100	83.0	(78.9 - 87.1)
Female	414	43,000	89.9	(86.4 - 93.3)	23,522	6,973,100	89.9	(87.2 - 92.6)
Age (years)								
18-34	*	*	*	*	3,910	3,302,100	74.3	(67.8 - 80.7)
35-44	80	14,600	87.8	(78.9 - 96.7)	4,986	2,560,300	88.9	(84.7 - 93.1)
45-54	159	16,100	86.2	(80.8 - 91.6)	7,831	2,530,100	88.9	(84.6 - 93.2)
55-64	129	12,400	90.5	(85.6 - 95.3)	8,764	1,986,700	92.8	(90.2 - 95.5)
≥ 65	209	13,900	97.7	(95.0 - 100)	12,628	2,475,000	97.8	(96.7 - 98.8)
Race/Ethnicity								
White Non-Hispanic	648	78,400	86.9	(82.8 - 91.1)	35,811	8,591,200	92.2	(90.6 - 93.8)
Black Non-Hispanic	*	*	*	*	736	1,781,600	82.7	(75.3 - 90.1)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	334	34,400	77.0	(68.5 - 85.4)	15,673	4,448,900	79.7	(74.9 - 84.5)
Some College	156	21,200	90.6	(85.3 - 95.9)	9,906	3,004,600	85.0	(80.1 - 89.8)
≥ College Degree	175	24,000	96.7	(94.0 - 99.4)	12,459	5,361,700	95.3	(93.3 - 97.2)
Income								
≤ \$24,999	*	*	*	*	9,640	2,437,200	69.7	(62.5 - 76.8)
\$25,000-\$49,999	*	*	*	*	9,662	2,384,000	82.2	(76.1 - 88.3)
\$50,000-\$74,999	100	15,200	94.2	(89.0 - 99.3)	5,772	1,855,700	94.0	(91.2 - 96.7)
≥ \$75,000	116	20,700	97.4	(94.4 - 100)	8,567	4,388,400	97.5	(96.6 - 98.5)

¹ Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

² Sample size from survey

³ Estimated number of adults who have health insurance (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 6b. Health insurance among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	668	86.2	(81.6 - 90.7)	38,119	86.7	(84.4 - 89.1)
Gender						
Male	254	82.3	(74.0 - 90.6)	14,597	83.6	(79.7 - 87.5)
Female	414	89.9	(86.3 - 93.4)	23,522	89.6	(86.8 - 92.3)
Race/Ethnicity						
White Non-Hispanic	648	87.3	(83.2 - 91.4)	35,811	92.1	(90.5 - 93.7)
Black Non-Hispanic	*	*	*	736	82.5	(75.0 - 90.0)
Hispanic	*	*	*	860	68.7	(59.0 - 78.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	334	77.9	(69.6 - 86.2)	15,673	80.1	(75.4 - 84.7)
Some College	156	90.7	(85.5 - 96.0)	9,906	84.6	(79.7 - 89.4)
≥ College Degree	175	96.8	(94.2 - 99.5)	12,459	95.4	(93.6 - 97.2)
Income						
≤ \$24,999	*	*	*	9,640	70.2	(63.2 - 77.1)
\$25,000-\$49,999	176	82.9	(73.4 - 92.4)	9,662	81.9	(76.0 - 87.9)
\$50,000-\$74,999	100	94.1	(88.9 - 99.3)	5,772	94.0	(91.3 - 96.7)
≥ \$75,000	116	97.5	(94.5 - 100)	8,567	97.5	(96.6 - 98.5)

¹ Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 7a. Health insurance among adults aged 18-64¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	455	65,700	83.6	(78.1 - 89.0)	25,165	10,251,300	84.2	(81.3 - 87.1)
Gender								
Male	192	30,600	78.3	(68.6 - 88.1)	10,078	4,839,900	80.3	(75.5 - 85.0)
Female	263	35,100	88.8	(84.7 - 92.8)	15,087	5,411,400	88.0	(84.7 - 91.3)
Age (years)								
18-34	*	*	*	*	3,910	3,302,100	74.3	(67.8 - 80.7)
35-44	80	14,600	87.8	(78.9 - 96.7)	4,981	2,556,600	88.9	(84.7 - 93.1)
45-54	157	15,800	85.9	(80.4 - 91.4)	7,791	2,472,900	88.7	(84.3 - 93.0)
55-64	127	12,300	90.9	(86.1 - 95.7)	8,483	1,919,700	92.8	(90.1 - 95.5)
≥ 65	-	-	-	-	-	-	-	-
Race/Ethnicity								
White Non-Hispanic	440	64,400	85.0	(80.2 - 89.8)	23,354	6,582,200	90.3	(88.3 - 92.4)
Black Non-Hispanic	*	*	*	*	567	1,464,100	80.2	(71.6 - 88.7)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	9,037	3,191,500	74.6	(68.6 - 80.6)
Some College	112	18,000	89.3	(83.2 - 95.5)	7,031	2,540,400	82.9	(77.4 - 88.4)
≥ College Degree	136	21,200	96.3	(93.3 - 99.3)	9,068	4,502,400	94.5	(92.2 - 96.8)
Income								
≤ \$24,999	*	*	*	*	4,918	1,707,200	62.6	(54.1 - 71.2)
\$25,000-\$49,999	*	*	*	*	6,180	1,783,000	77.7	(70.3 - 85.1)
\$50,000-\$74,999	88	14,200	93.8	(88.3 - 99.2)	4,554	1,553,400	92.9	(89.7 - 96.1)
≥ \$75,000	108	20,100	98.2	(95.6 - 100)	7,324	3,987,600	97.3	(96.3 - 98.3)

¹ Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

² Sample size from survey

³ Estimated number of adults aged 18-64 who have health insurance (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 7b. Health insurance among adults aged 18-64¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	455	83.8	(78.3 - 89.2)	25,165	84.3	(81.5 - 87.2)
Gender						
Male	192	78.5	(68.6 - 88.3)	10,078	80.5	(75.9 - 85.1)
Female	263	88.8	(84.6 - 92.9)	15,087	87.9	(84.6 - 91.2)
Race/Ethnicity						
White Non-Hispanic	440	85.1	(80.2 - 90.0)	23,354	90.3	(88.3 - 92.3)
Black Non-Hispanic	*	*	*	567	80.0	(71.3 - 88.6)
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	9,037	75.1	(69.3 - 80.9)
Some College	112	89.2	(83.0 - 95.5)	7,031	82.6	(77.1 - 88.1)
≥ College Degree	136	96.4	(93.4 - 99.4)	9,068	94.6	(92.5 - 96.8)
Income						
≤ \$24,999	*	*	*	4,918	63.2	(54.9 - 71.6)
\$25,000-\$49,999	*	*	*	6,180	77.5	(70.2 - 84.7)
\$50,000-\$74,999	88	93.7	(88.1 - 99.3)	4,554	93.0	(89.8 - 96.2)
≥ \$75,000	108	98.3	(95.8 - 100)	7,324	97.3	(96.3 - 98.3)

¹ Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 8a. No health insurance among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	668	13,300	14.3	(9.6 - 18.9)	38,119	1,989,300	13.4	(11.0 - 15.8)
Gender								
Male	254	8,500	18.6	(10.0 - 27.2)	14,597	1,207,400	17.0	(12.9 - 21.1)
Female	414	4,800	10.1	(6.7 - 13.6)	23,522	781,900	10.1	(7.4 - 12.8)
Age (years)								
18-34	*	*	*	*	3,910	1,144,100	25.7	(19.3 - 32.2)
35-44	*	*	*	*	4,986	319,200	11.1	(6.9 - 15.3)
45-54	159	2,600	13.8	(8.4 - 19.2)	7,831	316,500	11.1	(6.8 - 15.4)
55-64	129	1,300	9.5	(4.7 - 14.4)	8,764	153,300	7.2	(4.5 - 9.8)
≥ 65	*	*	*	*	12,628	56,200	2.2	(1.2 - 3.3)
Race/Ethnicity								
White Non-Hispanic	648	11,800	13.1	(8.9 - 17.2)	35,811	727,100	7.8	(6.2 - 9.4)
Black Non-Hispanic	*	*	*	*	736	372,800	17.3	(9.9 - 24.7)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	334	10,300	23.0	(14.6 - 31.5)	15,673	1,134,400	20.3	(15.5 - 25.1)
Some College	156	2,200	9.4	(4.1 - 14.7)	9,906	531,400	15.0	(10.2 - 19.9)
≥ College Degree	*	*	*	*	12,459	266,300	4.7	(2.8 - 6.7)
Income								
≤ \$24,999	*	*	*	*	9,640	1,061,200	30.3	(23.2 - 37.5)
\$25,000-\$49,999	*	*	*	*	9,662	516,700	17.8	(11.7 - 23.9)
\$50,000-\$74,999	*	*	*	*	5,772	119,100	6.0	(3.3 - 8.8)
≥ \$75,000	*	*	*	*	8,567	111,100	2.5	(1.5 - 3.4)

¹ Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

² Sample size from survey

³ Estimated number of adults who have no health insurance (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 8b. No health insurance among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	668	13.8	(9.3 - 18.4)	38,119	13.3	(10.9 - 15.6)
Gender						
Male	254	17.7	(9.4 - 26.0)	14,597	16.4	(12.5 - 20.3)
Female	414	10.1	(6.6 - 13.7)	23,522	10.4	(7.7 - 13.2)
Race/Ethnicity						
White Non-Hispanic	648	12.7	(8.6 - 16.8)	35,811	7.9	(6.3 - 9.5)
Black Non-Hispanic	*	*	*	736	17.5	(10.0 - 25.0)
Hispanic	*	*	*	860	31.3	(21.6 - 41.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	334	22.1	(13.8 - 30.4)	15,673	19.9	(15.3 - 24.6)
Some College	156	9.3	(4.0 - 14.5)	9,906	15.4	(10.6 - 20.3)
≥ College Degree	*	*	*	12,459	4.6	(2.8 - 6.4)
Income						
≤ \$24,999	*	*	*	9,640	29.8	(22.9 - 36.8)
\$25,000-\$49,999	176	17.1	(7.6 - 26.6)	9,662	18.1	(12.1 - 24.0)
\$50,000-\$74,999	*	*	*	5,772	6.0	(3.3 - 8.7)
≥ \$75,000	*	*	*	8,567	2.5	(1.5 - 3.4)

¹ Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 9a. No health insurance among adults aged 18-64¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	455	12,900	16.4	(11.0 - 21.9)	25,165	1,928,400	15.8	(12.9 - 18.7)
Gender								
Male	192	8,500	21.7	(11.9 - 31.4)	10,078	1,190,200	19.7	(15.0 - 24.5)
Female	263	4,500	11.2	(7.2 - 15.3)	15,087	738,200	12.0	(8.7 - 15.3)
Age (years)								
18-34	*	*	*	*	3,910	1,144,100	25.7	(19.3 - 32.2)
35-44	*	*	*	*	4,981	319,200	11.1	(6.9 - 15.3)
45-54	157	2,600	14.1	(8.6 - 19.6)	7,791	316,100	11.3	(7.0 - 15.7)
55-64	127	1,200	9.1	(4.3 - 13.9)	8,483	148,900	7.2	(4.5 - 9.9)
≥ 65	-	-	-	-	-	-	-	-
Race/Ethnicity								
White Non-Hispanic	440	11,400	15.0	(10.2 - 19.8)	23,354	703,100	9.7	(7.6 - 11.7)
Black Non-Hispanic	*	*	*	*	567	361,800	19.8	(11.3 - 28.4)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	9,037	1,087,700	25.4	(19.4 - 31.4)
Some College	112	2,100	10.7	(4.5 - 16.8)	7,031	522,800	17.1	(11.6 - 22.6)
≥ College Degree	*	*	*	*	9,068	261,200	5.5	(3.2 - 7.8)
Income								
≤ \$24,999	*	*	*	*	4,918	1,018,000	37.4	(28.8 - 45.9)
\$25,000-\$49,999	*	*	*	*	6,180	511,100	22.3	(14.9 - 29.7)
\$50,000-\$74,999	*	*	*	*	4,554	118,800	7.1	(3.9 - 10.3)
≥ \$75,000	*	*	*	*	7,324	110,700	2.7	(1.7 - 3.7)

¹ Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

² Sample size from survey

³ Estimated number of adults aged 18-64 who have no health insurance (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 9b. No health insurance among adults aged 18-64¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	455	16.2	(10.8 - 21.7)	25,165	15.7	(12.8 - 18.5)
Gender						
Male	192	21.5	(11.7 - 31.4)	10,078	19.5	(14.9 - 24.1)
Female	263	11.2	(7.1 - 15.4)	15,087	12.1	(8.8 - 15.4)
Race/Ethnicity						
White Non-Hispanic	440	14.9	(10.0 - 19.8)	23,354	9.7	(7.7 - 11.7)
Black Non-Hispanic	*	*	*	567	20.0	(11.4 - 28.7)
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	9,037	24.9	(19.1 - 30.7)
Some College	112	10.8	(4.5 - 17.0)	7,031	17.4	(11.9 - 22.9)
≥ College Degree	*	*	*	9,068	5.4	(3.2 - 7.5)
Income						
≤ \$24,999	*	*	*	4,918	36.8	(28.4 - 45.1)
\$25,000-\$49,999	*	*	*	6,180	22.5	(15.3 - 29.8)
\$50,000-\$74,999	*	*	*	4,554	7.0	(3.8 - 10.2)
≥ \$75,000	*	*	*	7,324	2.7	(1.7 - 3.7)

¹ Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 10a. Adults with regular health care provider¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	670	76,700	81.6	(76.7 - 86.5)	38,135	12,324,300	83.0	(80.4 - 85.6)
Gender								
Male	255	34,300	74.5	(66.0 - 83.1)	14,593	5,494,300	77.6	(73.2 - 82.0)
Female	415	42,400	88.4	(84.2 - 92.6)	23,542	6,830,000	87.8	(85.1 - 90.5)
Age (years)								
18-34	*	*	*	*	3,929	2,931,600	65.8	(59.2 - 72.5)
35-44	80	14,300	85.8	(76.5 - 95.1)	4,988	2,471,000	86.1	(81.9 - 90.3)
45-54	159	16,100	86.0	(79.9 - 92.1)	7,823	2,529,300	88.6	(84.6 - 92.7)
55-64	129	12,700	92.7	(87.8 - 97.5)	8,770	2,002,000	93.6	(91.7 - 95.4)
≥ 65	210	13,500	94.2	(90.1 - 98.3)	12,625	2,390,400	94.1	(92.2 - 96.0)
Race/Ethnicity								
White Non-Hispanic	650	75,600	83.2	(78.7 - 87.6)	35,821	8,227,700	88.2	(86.4 - 89.9)
Black Non-Hispanic	*	*	*	*	740	1,753,200	81.3	(72.6 - 89.9)
Hispanic	*	*	*	*	860	1,453,100	64.7	(54.7 - 74.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	336	34,800	76.7	(68.6 - 84.8)	15,692	4,249,200	75.9	(70.9 - 80.8)
Some College	156	18,800	80.2	(71.1 - 89.3)	9,904	2,935,000	82.8	(77.2 - 88.4)
≥ College Degree	175	22,700	91.5	(85.7 - 97.2)	12,456	5,046,100	89.8	(87.1 - 92.6)
Income								
≤ \$24,999	*	*	*	*	9,644	2,554,200	72.7	(65.9 - 79.6)
\$25,000-\$49,999	*	*	*	*	9,655	2,243,100	77.3	(70.0 - 84.6)
\$50,000-\$74,999	100	14,300	88.7	(81.1 - 96.4)	5,771	1,767,500	89.5	(85.2 - 93.9)
≥ \$75,000	116	19,000	89.5	(82.7 - 96.2)	8,560	4,122,800	91.9	(90.2 - 93.7)

¹ Defined as having one or more persons respondent thinks of as a personal doctor or health care provider

² Sample size from survey

³ Estimated number of adults who have at least one person they think of as their personal doctor or health care provider (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 10b. Adults with regular health care provider¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	670	82.0	(77.2 - 86.8)	38,135	83.0	(80.4 - 85.5)
Gender						
Male	255	75.2	(66.9 - 83.6)	14,593	78.1	(73.9 - 82.3)
Female	415	88.5	(84.3 - 92.7)	23,542	87.4	(84.6 - 90.1)
Race/Ethnicity						
White Non-Hispanic	650	83.5	(79.1 - 87.9)	35,821	88.0	(86.2 - 89.7)
Black Non-Hispanic	*	*	*	740	81.4	(73.0 - 89.8)
Hispanic	*	*	*	860	65.2	(55.5 - 75.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	336	77.1	(69.2 - 85.1)	15,692	75.8	(71.0 - 80.6)
Some College	156	80.9	(72.1 - 89.7)	9,904	82.9	(77.5 - 88.3)
≥ College Degree	175	91.7	(86.0 - 97.4)	12,456	89.8	(87.1 - 92.4)
Income						
≤ \$24,999	*	*	*	9,644	72.9	(66.1 - 79.6)
\$25,000-\$49,999	177	81.0	(71.2 - 90.8)	9,655	77.4	(70.5 - 84.3)
\$50,000-\$74,999	100	89.5	(82.4 - 96.7)	5,771	89.6	(85.3 - 94.0)
≥ \$75,000	116	89.8	(83.2 - 96.4)	8,560	91.6	(89.8 - 93.4)

¹ Defined as having one or more persons respondent thinks of as a personal doctor or health care provider

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 11a. Cost prevented visit to doctor within the past year among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	667	10,200	10.9	(7.7 - 14.1)	38,143	2,034,800	13.7	(11.6 - 15.8)
Gender								
Male	253	4,700	10.2	(5.0 - 15.4)	14,607	903,300	12.7	(9.5 - 16.0)
Female	414	5,500	11.6	(7.7 - 15.4)	23,536	1,131,500	14.6	(11.8 - 17.4)
Age (years)								
18-34	92	3,700	12.0	(4.6 - 19.4)	3,939	947,300	21.2	(15.6 - 26.9)
35-44	80	2,300	13.9	(5.5 - 22.3)	4,987	347,900	12.1	(8.0 - 16.2)
45-54	159	2,800	14.9	(8.7 - 21.2)	7,825	382,200	13.5	(9.9 - 17.1)
55-64	128	900	6.9	(1.9 - 11.9)	8,766	236,800	11.1	(8.0 - 14.2)
≥ 65	*	*	*	*	12,626	120,600	4.8	(2.5 - 7.0)
Race/Ethnicity								
White Non-Hispanic	648	10,200	11.2	(7.9 - 14.6)	35,833	984,600	10.6	(8.9 - 12.2)
Black Non-Hispanic	*	*	*	*	738	280,000	13.1	(7.2 - 19.0)
Hispanic	*	*	*	*	859	597,900	26.5	(17.3 - 35.7)
Other Non-Hispanic	*	*	*	*	713	172,300	15.3	(5.7 - 24.9)
Education								
≤ High School	333	7,000	15.5	(9.8 - 21.2)	15,696	996,600	17.8	(13.8 - 21.9)
Some College	156	2,600	11.1	(5.2 - 16.9)	9,911	512,400	14.5	(10.4 - 18.6)
≥ College Degree	*	*	*	*	12,453	466,900	8.3	(6.0 - 10.6)
Income								
≤ \$24,999	176	3,600	18.4	(10.8 - 26.0)	9,639	854,500	24.3	(18.4 - 30.3)
\$25,000-\$49,999	175	3,000	12.6	(4.3 - 20.9)	9,656	532,900	18.4	(12.5 - 24.3)
\$50,000-\$74,999	*	*	*	*	5,776	202,700	10.3	(6.0 - 14.5)
≥ \$75,000	*	*	*	*	8,561	192,000	4.3	(2.9 - 5.6)

¹ Defined as needing to see a doctor within the past year but not being able to because of cost

² Sample size from survey

³ Estimated number of adults who did not see a doctor because of cost within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 11b. Cost prevented visit to doctor within the past year among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	667	10.9	(7.7 - 14.2)	38,143	13.8	(11.7 - 15.9)
Gender						
Male	253	10.1	(5.0 - 15.3)	14,607	12.5	(9.4 - 15.6)
Female	414	11.7	(7.7 - 15.7)	23,536	15.0	(12.1 - 17.9)
Race/Ethnicity						
White Non-Hispanic	648	11.3	(7.9 - 14.6)	35,833	10.8	(9.2 - 12.5)
Black Non-Hispanic	*	*	*	738	13.2	(7.2 - 19.3)
Hispanic	*	*	*	859	26.1	(17.2 - 35.1)
Other Non-Hispanic	*	*	*	713	14.8	(5.5 - 24.2)
Education						
≤ High School	333	15.7	(9.9 - 21.5)	15,696	17.8	(13.8 - 21.7)
Some College	156	10.6	(5.0 - 16.3)	9,911	15.3	(11.0 - 19.5)
≥ College Degree	*	*	*	12,453	8.2	(6.0 - 10.5)
Income						
≤ \$24,999	176	18.5	(10.7 - 26.2)	9,639	24.5	(18.6 - 30.4)
\$25,000-\$49,999	175	12.9	(4.7 - 21.1)	9,656	18.7	(12.9 - 24.6)
\$50,000-\$74,999	*	*	*	5,776	10.4	(6.1 - 14.7)
≥ \$75,000	*	*	*	8,561	4.4	(3.1 - 5.8)

¹ Defined as needing to see a doctor within the past year but not being able to because of cost

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 12a. Visited a doctor for routine checkup within the past year among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	667	63,500	67.6	(62.5 - 72.7)	38,026	10,809,300	72.9	(70.6 - 75.3)
Gender								
Male	255	30,200	65.6	(57.1 - 74.0)	14,592	4,738,400	66.9	(63.0 - 70.9)
Female	412	33,300	69.6	(63.9 - 75.3)	23,434	6,070,900	78.5	(75.9 - 81.0)
Age (years)								
18-34	*	*	*	*	3,921	2,747,400	61.7	(55.6 - 67.9)
35-44	*	*	*	*	4,973	1,883,100	65.5	(60.2 - 70.7)
45-54	159	12,500	67.0	(59.0 - 75.1)	7,808	2,174,100	76.6	(73.2 - 80.0)
55-64	128	11,000	80.7	(73.5 - 87.9)	8,751	1,728,600	81.4	(78.7 - 84.2)
≥ 65	208	12,700	89.3	(85.0 - 93.7)	12,573	2,276,000	90.0	(87.5 - 92.5)
Race/Ethnicity								
White Non-Hispanic	647	61,900	68.2	(63.4 - 73.1)	35,721	6,746,200	72.4	(70.6 - 74.2)
Black Non-Hispanic	*	*	*	*	736	1,688,900	78.8	(70.4 - 87.3)
Hispanic	*	*	*	*	857	1,550,900	69.3	(60.0 - 78.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	334	29,500	65.1	(57.2 - 73.1)	15,640	4,150,600	74.7	(70.7 - 78.7)
Some College	155	16,700	71.4	(62.1 - 80.8)	9,880	2,524,000	71.3	(66.1 - 76.5)
≥ College Degree	175	16,900	68.1	(59.3 - 76.8)	12,428	4,042,900	71.9	(68.5 - 75.3)
Income								
≤ \$24,999	*	*	*	*	9,616	2,469,200	70.4	(64.2 - 76.6)
\$25,000-\$49,999	176	16,700	69.9	(59.9 - 79.8)	9,631	2,068,400	71.3	(65.3 - 77.3)
\$50,000-\$74,999	*	*	*	*	5,765	1,466,000	74.8	(69.6 - 80.0)
≥ \$75,000	116	14,800	69.7	(59.8 - 79.6)	8,547	3,176,500	70.7	(67.3 - 74.2)

¹ Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past year

² Sample size from survey

³ Estimated number of adults who have visited a doctor for a routine checkup within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 12b. Visited a doctor for routine checkup within the past year among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	667	67.7	(62.6 - 72.8)	38,026	72.7	(70.3 - 75.0)
Gender						
Male	255	65.9	(57.6 - 74.3)	14,592	67.1	(63.3 - 70.9)
Female	412	69.4	(63.6 - 75.2)	23,434	77.7	(75.1 - 80.3)
Race/Ethnicity						
White Non-Hispanic	647	68.2	(63.3 - 73.1)	35,721	71.9	(70.1 - 73.7)
Black Non-Hispanic	*	*	*	736	79.3	(71.3 - 87.4)
Hispanic	*	*	*	857	69.2	(60.1 - 78.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	334	65.2	(57.3 - 73.1)	15,640	74.4	(70.4 - 78.4)
Some College	155	72.5	(63.4 - 81.7)	9,880	70.9	(65.9 - 76.0)
≥ College Degree	175	67.1	(58.1 - 76.1)	12,428	71.7	(68.3 - 75.0)
Income						
≤ \$24,999	*	*	*	9,616	70.2	(64.1 - 76.2)
\$25,000-\$49,999	176	70.1	(60.4 - 79.9)	9,631	71.1	(65.3 - 76.9)
\$50,000-\$74,999	*	*	*	5,765	74.5	(69.2 - 79.8)
≥ \$75,000	*	*	*	8,547	70.3	(66.9 - 73.8)

¹ Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past year

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 13a. Visited a doctor for routine checkup within the past 2 years among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	667	73,600	78.4	(73.6 - 83.3)	38,026	12,726,600	85.9	(83.9 - 87.9)
Gender								
Male	255	34,500	74.9	(66.6 - 83.3)	14,592	5,791,900	81.8	(78.2 - 85.4)
Female	412	39,100	81.8	(76.8 - 86.9)	23,434	6,934,700	89.6	(87.9 - 91.3)
Age (years)								
18-34	*	*	*	*	3,921	3,535,400	79.4	(73.9 - 85.0)
35-44	*	*	*	*	4,973	2,355,600	81.9	(78.0 - 85.7)
45-54	159	15,300	81.7	(74.9 - 88.5)	7,808	2,507,900	88.4	(85.6 - 91.2)
55-64	128	12,600	92.4	(87.7 - 97.1)	8,751	1,924,800	90.7	(88.6 - 92.7)
≥ 65	208	13,200	92.7	(88.9 - 96.5)	12,573	2,402,900	95.0	(93.5 - 96.5)
Race/Ethnicity								
White Non-Hispanic	647	72,000	79.4	(74.8 - 83.9)	35,721	8,070,000	86.6	(85.4 - 87.9)
Black Non-Hispanic	*	*	*	*	736	1,914,500	89.4	(82.3 - 96.5)
Hispanic	*	*	*	*	857	1,832,800	81.9	(73.7 - 90.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	334	34,200	75.6	(67.6 - 83.5)	15,640	4,713,700	84.8	(81.2 - 88.4)
Some College	155	19,500	83.5	(75.4 - 91.7)	9,880	2,990,500	84.5	(79.6 - 89.4)
≥ College Degree	175	19,500	78.5	(70.4 - 86.6)	12,428	4,928,300	87.6	(85.3 - 89.9)
Income								
≤ \$24,999	*	*	*	*	9,616	2,911,700	83.0	(77.4 - 88.6)
\$25,000-\$49,999	176	19,400	81.2	(72.1 - 90.3)	9,631	2,375,500	81.9	(76.1 - 87.7)
\$50,000-\$74,999	100	13,000	80.9	(71.7 - 90.0)	5,765	1,733,500	88.4	(84.9 - 91.9)
≥ \$75,000	116	16,600	78.2	(69.1 - 87.4)	8,547	3,907,000	87.0	(84.8 - 89.2)

¹ Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past 2 years

² Sample size from survey

³ Estimated number of adults who have visited a doctor for a routine checkup within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 13b. Visited a doctor for routine checkup within the past 2 years among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	667	78.4	(73.5 - 83.2)	38,026	85.7	(83.7 - 87.6)
Gender						
Male	255	75.2	(67.1 - 83.4)	14,592	81.9	(78.4 - 85.3)
Female	412	81.3	(76.1 - 86.6)	23,434	89.1	(87.4 - 90.9)
Race/Ethnicity						
White Non-Hispanic	647	79.2	(74.6 - 83.7)	35,721	86.2	(85.0 - 87.5)
Black Non-Hispanic	*	*	*	736	89.6	(83.0 - 96.3)
Hispanic	*	*	*	857	81.8	(73.8 - 89.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	334	75.7	(67.9 - 83.6)	15,640	84.6	(81.1 - 88.1)
Some College	155	84.1	(76.2 - 92.0)	9,880	84.4	(79.8 - 89.0)
≥ College Degree	175	77.3	(68.9 - 85.8)	12,428	87.3	(85.0 - 89.6)
Income						
≤ \$24,999	*	*	*	9,616	83.2	(77.8 - 88.5)
\$25,000-\$49,999	176	81.0	(71.9 - 90.0)	9,631	81.9	(76.3 - 87.4)
\$50,000-\$74,999	100	81.3	(72.3 - 90.4)	5,765	88.1	(84.4 - 91.8)
≥ \$75,000	116	77.4	(67.9 - 86.8)	8,547	86.4	(84.2 - 88.7)

¹ Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past 2 years

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 14a. Dentist visit within the past year among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	664	63,400	67.6	(62.7 - 72.5)	38,026	10,540,400	71.1	(68.6 - 73.6)
Gender								
Male	254	29,300	63.7	(55.5 - 71.9)	14,573	4,845,800	68.4	(64.5 - 72.2)
Female	410	34,000	71.4	(66.0 - 76.8)	23,453	5,694,600	73.6	(70.4 - 76.8)
Age (years)								
18-34	*	*	*	*	3,924	2,789,700	62.6	(56.3 - 69.0)
35-44	*	*	*	*	4,979	2,181,000	75.8	(70.9 - 80.7)
45-54	158	13,000	69.7	(61.8 - 77.5)	7,814	2,149,300	75.8	(71.9 - 79.7)
55-64	129	9,200	66.8	(57.7 - 75.8)	8,742	1,690,500	79.2	(75.9 - 82.6)
≥ 65	205	9,000	63.7	(56.1 - 71.4)	12,567	1,729,800	68.5	(64.7 - 72.3)
Race/Ethnicity								
White Non-Hispanic	645	62,100	68.5	(63.9 - 73.2)	35,721	6,959,900	74.7	(72.7 - 76.6)
Black Non-Hispanic	*	*	*	*	735	1,405,800	66.2	(57.9 - 74.5)
Hispanic	*	*	*	*	862	1,425,600	63.1	(53.7 - 72.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	330	26,000	57.7	(49.8 - 65.5)	15,613	3,620,700	65.1	(60.7 - 69.5)
Some College	156	16,500	70.7	(62.5 - 78.9)	9,880	2,504,700	70.8	(66.1 - 75.5)
≥ College Degree	175	20,500	82.6	(75.2 - 90.0)	12,453	4,402,400	78.2	(74.6 - 81.8)
Income								
≤ \$24,999	*	*	*	*	9,594	1,972,200	56.2	(49.8 - 62.6)
\$25,000-\$49,999	*	*	*	*	9,649	1,896,000	65.8	(59.5 - 72.1)
\$50,000-\$74,999	*	*	*	*	5,763	1,482,400	75.2	(69.4 - 81.0)
≥ \$75,000	116	17,400	81.9	(74.0 - 89.7)	8,564	3,707,900	82.4	(79.5 - 85.3)

¹ Defined as having visited a dentist or a dental clinic for any reason within the past year

² Sample size from survey

³ Estimated number of adults who have visited a dentist or a dental clinic for any reason within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 14b. Dentist visit within the past year among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	664	67.4	(62.5 - 72.3)	38,026	71.1	(68.6 - 73.5)
Gender						
Male	254	63.0	(54.8 - 71.2)	14,573	68.3	(64.6 - 72.1)
Female	410	71.6	(66.2 - 77.1)	23,453	73.6	(70.4 - 76.8)
Race/Ethnicity						
White Non-Hispanic	645	68.3	(63.5 - 73.0)	35,721	74.4	(72.4 - 76.3)
Black Non-Hispanic	*	*	*	735	66.6	(58.3 - 74.9)
Hispanic	*	*	*	862	63.6	(54.4 - 72.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	330	57.1	(49.3 - 65.0)	15,613	65.3	(60.9 - 69.6)
Some College	156	70.8	(62.6 - 79.1)	9,880	70.4	(65.7 - 75.1)
≥ College Degree	175	82.8	(75.3 - 90.3)	12,453	78.1	(74.6 - 81.7)
Income						
≤ \$24,999	*	*	*	9,594	56.4	(50.1 - 62.8)
\$25,000-\$49,999	*	*	*	9,649	65.8	(59.6 - 72.0)
\$50,000-\$74,999	*	*	*	5,763	75.0	(69.1 - 80.8)
≥ \$75,000	116	82.2	(74.2 - 90.2)	8,564	82.1	(79.1 - 85.0)

¹ Defined as having visited a dentist or a dental clinic for any reason within the past year

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 15a. Had teeth cleaned within the past year among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	589	60,600	69.9	(64.7 - 75.1)	34,070	9,985,900	71.7	(69.1 - 74.3)
Gender								
Male	225	27,700	65.0	(56.2 - 73.8)	13,173	4,616,400	69.0	(65.0 - 73.1)
Female	364	32,800	74.7	(69.1 - 80.3)	20,897	5,369,500	74.2	(70.8 - 77.5)
Age (years)								
18-34	*	*	*	*	3,860	2,746,400	63.0	(56.5 - 69.5)
35-44	*	*	*	*	4,875	2,112,100	75.5	(70.7 - 80.3)
45-54	148	12,000	68.5	(60.2 - 76.7)	7,438	2,022,600	73.3	(69.1 - 77.4)
55-64	115	9,200	73.3	(64.2 - 82.5)	7,906	1,560,900	77.7	(73.8 - 81.6)
≥ 65	161	8,400	77.8	(71.0 - 84.6)	9,991	1,543,900	77.1	(73.7 - 80.5)
Race/Ethnicity								
White Non-Hispanic	571	59,100	70.6	(65.7 - 75.6)	31,982	6,601,500	75.4	(73.4 - 77.4)
Black Non-Hispanic	*	*	*	*	680	1,227,200	61.6	(52.1 - 71.0)
Hispanic	*	*	*	*	789	1,421,300	66.6	(57.1 - 76.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	277	24,800	61.8	(53.1 - 70.4)	12,740	3,388,600	67.7	(63.1 - 72.3)
Some College	137	15,500	71.7	(62.7 - 80.7)	9,145	2,286,800	68.5	(62.9 - 74.1)
≥ College Degree	173	20,000	81.1	(73.5 - 88.8)	12,128	4,298,500	78.3	(74.8 - 81.7)
Income								
≤ \$24,999	*	*	*	*	7,525	1,861,800	60.3	(53.5 - 67.1)
\$25,000-\$49,999	*	*	*	*	8,709	1,681,700	62.6	(55.4 - 69.8)
\$50,000-\$74,999	*	*	*	*	5,534	1,447,800	76.3	(70.7 - 81.8)
≥ \$75,000	115	17,200	81.6	(73.5 - 89.6)	8,448	3,617,500	80.8	(77.8 - 83.8)

¹ Defined as having had teeth cleaned by a dentist or dental hygienist within the past year

² Sample size from survey

³ Estimated number of adults who have had their teeth cleaned by a dentist or dental hygienist within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 15b. Had teeth cleaned within the past year among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	589	70.0	(64.7 - 75.2)	34,070	71.7	(69.1 - 74.2)
Gender						
Male	225	64.6	(55.9 - 73.4)	13,173	69.2	(65.3 - 73.1)
Female	364	75.1	(69.4 - 80.7)	20,897	73.9	(70.5 - 77.3)
Race/Ethnicity						
White Non-Hispanic	571	70.6	(65.6 - 75.6)	31,982	75.1	(73.0 - 77.1)
Black Non-Hispanic	*	*	*	680	62.0	(52.7 - 71.4)
Hispanic	*	*	*	789	66.9	(57.5 - 76.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	277	61.6	(53.0 - 70.2)	12,740	67.7	(63.2 - 72.3)
Some College	137	72.3	(63.4 - 81.2)	9,145	68.1	(62.6 - 73.6)
≥ College Degree	173	81.2	(73.4 - 89.0)	12,128	78.3	(74.9 - 81.7)
Income						
≤ \$24,999	*	*	*	7,525	60.2	(53.4 - 67.0)
\$25,000-\$49,999	*	*	*	8,709	62.7	(55.7 - 69.7)
\$50,000-\$74,999	*	*	*	5,534	76.4	(70.9 - 81.9)
≥ \$75,000	115	81.6	(73.4 - 89.9)	8,448	80.4	(77.4 - 83.4)

¹ Defined as having had teeth cleaned by a dentist or dental hygienist within the past year

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 16a. Permanent teeth extracted due to decay or gum disease among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	663	44,000	47.0	(42.0 - 52.0)	37,652	7,426,500	50.4	(47.8 - 52.9)
Gender								
Male	254	20,800	45.2	(37.2 - 53.1)	14,457	3,367,000	47.8	(43.9 - 51.6)
Female	409	23,200	48.8	(42.8 - 54.7)	23,195	4,059,500	52.7	(49.5 - 56.0)
Age (years)								
18-34	92	7,100	23.2	(14.4 - 31.9)	3,921	1,121,600	25.4	(19.7 - 31.0)
35-44	*	*	*	*	4,964	1,250,000	43.6	(37.9 - 49.2)
45-54	159	9,800	52.5	(44.0 - 61.1)	7,780	1,595,300	56.1	(51.4 - 60.7)
55-64	126	9,500	70.5	(61.7 - 79.3)	8,648	1,382,800	65.5	(61.6 - 69.4)
≥ 65	206	12,500	88.9	(84.6 - 93.2)	12,339	2,076,900	83.0	(80.6 - 85.4)
Race/Ethnicity								
White Non-Hispanic	643	43,100	47.6	(42.7 - 52.6)	35,364	4,425,700	47.9	(46.0 - 49.9)
Black Non-Hispanic	*	*	*	*	727	1,244,700	58.2	(48.9 - 67.4)
Hispanic	*	*	*	*	853	1,220,500	54.2	(44.9 - 63.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	330	26,000	57.9	(49.6 - 66.2)	15,427	3,436,000	61.9	(57.2 - 66.6)
Some College	156	10,200	43.4	(34.0 - 52.8)	9,798	1,726,200	49.3	(44.3 - 54.2)
≥ College Degree	174	7,600	30.7	(23.0 - 38.3)	12,351	2,242,600	40.1	(36.5 - 43.6)
Income								
≤ \$24,999	*	*	*	*	9,483	2,171,900	62.3	(55.8 - 68.8)
\$25,000-\$49,999	*	*	*	*	9,552	1,577,400	54.7	(48.3 - 61.0)
\$50,000-\$74,999	*	*	*	*	5,744	1,000,700	51.5	(45.6 - 57.4)
≥ \$75,000	116	5,700	26.8	(18.2 - 35.3)	8,516	1,738,200	38.8	(35.3 - 42.2)

¹ Defined as having had at least one permanent tooth extracted because of tooth decay or gum disease

² Sample size from survey

³ Estimated number of adults who have had at least one permanent tooth extracted because of tooth decay or gum disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 16b. Permanent teeth extracted due to decay or gum disease among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	663	46.7	(41.7 - 51.7)	37,652	49.4	(46.9 - 51.9)
Gender						
Male	254	45.9	(37.9 - 53.9)	14,457	47.9	(44.1 - 51.7)
Female	409	47.5	(41.5 - 53.6)	23,195	50.7	(47.4 - 54.0)
Race/Ethnicity						
White Non-Hispanic	643	47.3	(42.3 - 52.3)	35,364	46.5	(44.5 - 48.5)
Black Non-Hispanic	*	*	*	727	57.7	(48.6 - 66.9)
Hispanic	*	*	*	853	53.8	(44.6 - 62.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	330	57.4	(49.1 - 65.6)	15,427	61.3	(56.7 - 66.0)
Some College	156	43.1	(33.8 - 52.5)	9,798	47.9	(43.0 - 52.9)
≥ College Degree	174	30.7	(23.0 - 38.4)	12,351	39.0	(35.5 - 42.5)
Income						
≤ \$24,999	*	*	*	9,483	61.6	(55.2 - 68.1)
\$25,000-\$49,999	*	*	*	9,552	54.2	(48.0 - 60.3)
\$50,000-\$74,999	*	*	*	5,744	49.9	(44.0 - 55.9)
≥ \$75,000	116	25.8	(17.2 - 34.4)	8,516	37.7	(34.2 - 41.1)

¹ Defined as having had at least one permanent tooth extracted because of tooth decay or gum disease

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 17a. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	206	3,300	23.5	(16.5 - 30.5)	12,339	491,200	19.6	(16.0 - 23.3)
Gender								
Male	*	*	*	*	4,352	167,200	16.4	(11.8 - 20.9)
Female	147	1,700	21.6	(13.9 - 29.3)	7,987	324,000	21.8	(16.6 - 27.0)
Age (years)								
≥ 65	206	3,300	23.5	(16.5 - 30.5)	12,339	491,200	19.6	(16.0 - 23.3)
Race/Ethnicity								
White Non-Hispanic	201	3,200	23.1	(16.1 - 30.2)	11,885	322,600	16.8	(14.3 - 19.2)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	124	2,400	29.4	(19.6 - 39.2)	6,358	314,100	25.3	(19.9 - 30.7)
Some College	*	*	*	*	2,736	90,600	20.2	(12.1 - 28.2)
≥ College Degree	*	*	*	*	3,214	80,500	10.1	(3.6 - 16.5)
Income								
≤ \$24,999	*	*	*	*	4,558	199,600	26.9	(19.3 - 34.4)
\$25,000-\$49,999	*	*	*	*	3,365	123,900	21.4	(13.1 - 29.6)
\$50,000-\$74,999	*	*	*	*	1,179	21,400	7.3	(3.8 - 10.8)
≥ \$75,000	*	*	*	*	1,187	9,600	2.5	(1.0 - 4.0)

¹ Defined as having all permanent teeth extracted because of tooth decay or gum disease among adults aged 65 and older

² Sample size from survey

³ Estimated number of adults aged 65 and older who have had all permanent teeth extracted because of tooth decay or gum disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 17b. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	206	23.8	(16.5 - 31.2)	12,339	19.4	(15.7 - 23.2)
Gender						
Male	*	*	*	4,352	16.5	(11.5 - 21.4)
Female	147	21.6	(13.9 - 29.3)	7,987	22.1	(16.6 - 27.7)
Race/Ethnicity						
White Non-Hispanic	201	23.4	(15.9 - 30.8)	11,885	16.6	(13.9 - 19.3)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	6,358	24.9	(19.3 - 30.5)
Some College	*	*	*	2,736	21.0	(12.1 - 29.9)
≥ College Degree	*	*	*	3,214	10.0	(3.6 - 16.3)
Income						
≤ \$24,999	*	*	*	4,558	27.3	(19.1 - 35.5)
\$25,000-\$49,999	*	*	*	3,365	21.0	(12.9 - 29.1)
\$50,000-\$74,999	*	*	*	1,179	7.6	(3.6 - 11.5)
≥ \$75,000	*	*	*	1,187	2.2	(0.9 - 3.5)

¹ Defined as having all permanent teeth extracted because of tooth decay or gum disease among adults aged 65 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 18a. Flu shot within the past 12 months¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	669	36,800	39.1	(34.4 - 43.8)	38,079	6,255,900	42.3	(39.9 - 44.8)
Gender								
Male	255	15,800	34.4	(27.3 - 41.5)	14,579	2,798,900	39.6	(35.8 - 43.5)
Female	414	20,900	43.7	(37.8 - 49.6)	23,500	3,456,900	44.8	(41.6 - 47.9)
Age (years)								
18-49	243	14,600	26.2	(19.9 - 32.6)	12,505	2,685,300	31.1	(27.3 - 34.9)
50-64	217	12,500	52.4	(45.1 - 59.7)	12,972	1,662,900	46.0	(42.6 - 49.5)
≥ 65	209	9,600	67.3	(60.0 - 74.7)	12,602	1,907,700	75.2	(72.0 - 78.4)
Race/Ethnicity								
White Non-Hispanic	649	36,200	39.9	(35.1 - 44.6)	35,770	4,029,000	43.3	(41.4 - 45.3)
Black Non-Hispanic	*	*	*	*	738	907,000	42.1	(33.1 - 51.1)
Hispanic	*	*	*	*	860	882,900	39.6	(30.4 - 48.9)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	335	15,300	33.7	(27.1 - 40.4)	15,663	2,457,800	44.3	(39.8 - 48.8)
Some College	156	9,100	39.1	(29.9 - 48.2)	9,898	1,377,700	38.9	(34.0 - 43.9)
≥ College Degree	175	12,200	49.1	(39.9 - 58.2)	12,436	2,404,400	43.0	(39.5 - 46.4)
Income								
≤ \$24,999	177	6,200	31.7	(22.9 - 40.4)	9,635	1,656,800	47.6	(41.2 - 54.0)
\$25,000-\$49,999	177	10,200	42.7	(33.0 - 52.5)	9,647	1,200,600	41.4	(35.6 - 47.3)
\$50,000-\$74,999	*	*	*	*	5,763	792,300	40.2	(34.3 - 46.2)
≥ \$75,000	*	*	*	*	8,548	1,776,100	39.6	(36.3 - 42.8)

¹ Defined as having had a flu shot within the past 12 months

² Sample size from survey

³ Estimated number of adults who have had a flu shot within the past 12 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 18b. Flu shot within the past 12 months¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	669	39.0	(34.2 - 43.7)	38,079	41.7	(39.2 - 44.1)
Gender						
Male	255	34.3	(27.3 - 41.3)	14,579	40.0	(36.2 - 43.8)
Female	414	43.4	(37.4 - 49.4)	23,500	43.1	(40.0 - 46.3)
Race/Ethnicity						
White Non-Hispanic	649	39.6	(34.9 - 44.4)	35,770	42.4	(40.4 - 44.3)
Black Non-Hispanic	*	*	*	738	42.3	(33.3 - 51.3)
Hispanic	*	*	*	860	39.0	(29.9 - 48.1)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	335	33.7	(27.0 - 40.3)	15,663	43.7	(39.1 - 48.2)
Some College	156	40.0	(30.7 - 49.3)	9,898	37.8	(33.0 - 42.7)
≥ College Degree	175	47.9	(38.7 - 57.1)	12,436	42.5	(39.1 - 46.0)
Income						
≤ \$24,999	177	31.6	(22.8 - 40.4)	9,635	47.3	(41.0 - 53.7)
\$25,000-\$49,999	177	43.4	(33.8 - 53.1)	9,647	40.3	(34.5 - 46.0)
\$50,000-\$74,999	*	*	*	5,763	39.4	(33.4 - 45.4)
≥ \$75,000	*	*	*	8,548	39.1	(35.8 - 42.4)

¹ Defined as having had a flu shot within the past 12 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 19a. Flu vaccine that was sprayed in nose within the past 12 months¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	*	*	*	*	38,106	248,300	1.7	(0.6 - 2.8)
Gender								
Male	*	*	*	*	14,586	148,000	2.1	(0.0 - 4.2)
Female	*	*	*	*	23,520	100,300	1.3	(0.5 - 2.1)
Age (years)								
18-49	*	*	*	*	12,535	184,900	2.1	(0.3 - 3.9)
50-64	*	*	*	*	12,968	36,000	1.0	(0.2 - 1.8)
≥ 65	*	*	*	*	12,603	27,400	1.1	(0.4 - 1.8)
Race/Ethnicity								
White Non-Hispanic	*	*	*	*	35,796	66,000	0.7	(0.4 - 1.0)
Black Non-Hispanic	*	*	*	*	740	38,500	1.8	(0.0 - 4.1)
Hispanic	*	*	*	*	860	133,700	5.9	(0.0 - 12.3)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,679	172,500	3.1	(0.3 - 5.9)
Some College	*	*	*	*	9,897	43,500	1.2	(0.4 - 2.0)
≥ College Degree	*	*	*	*	12,449	31,600	0.6	(0.3 - 0.8)
Income								
≤ \$24,999	*	*	*	*	9,636	138,400	3.9	(0.0 - 8.1)
\$25,000-\$49,999	*	*	*	*	9,648	41,500	1.4	(0.0 - 3.1)
\$50,000-\$74,999	*	*	*	*	5,766	13,300	0.7	(0.2 - 1.2)
≥ \$75,000	*	*	*	*	8,561	35,300	0.8	(0.3 - 1.3)

¹ Defined as having had flu vaccine sprayed in nose within the past 12 months

² Sample size from survey

³ Estimated number of adults who have had flu vaccine sprayed in nose within the past 12 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 19b. Flu vaccine that was sprayed in nose within the past 12 months¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	*	*	*	38,106	1.7	(0.6 - 2.7)
Gender						
Male	*	*	*	14,586	2.0	(0.1 - 3.9)
Female	*	*	*	23,520	1.4	(0.5 - 2.3)
Race/Ethnicity						
White Non-Hispanic	*	*	*	35,796	0.7	(0.4 - 1.0)
Black Non-Hispanic	*	*	*	740	1.9	(0.0 - 4.4)
Hispanic	*	*	*	860	5.9	(0.0 - 11.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	15,679	3.1	(0.4 - 5.8)
Some College	*	*	*	9,897	1.2	(0.4 - 2.0)
≥ College Degree	*	*	*	12,449	0.6	(0.3 - 0.8)
Income						
≤ \$24,999	*	*	*	9,636	3.8	(0.0 - 7.7)
\$25,000-\$49,999	*	*	*	9,648	1.5	(0.0 - 3.4)
\$50,000-\$74,999	*	*	*	5,766	0.7	(0.1 - 1.2)
≥ \$75,000	*	*	*	8,561	0.8	(0.3 - 1.3)

¹ Defined as having had flu vaccine sprayed in nose within the past 12 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 20a. Flu shot or flu vaccine sprayed in nose within the past 12 months¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	669	37,200	39.6	(34.8 - 44.3)	38,059	6,302,300	42.7	(40.2 - 45.1)
Gender								
Male	255	16,200	35.3	(28.1 - 42.5)	14,568	2,822,700	40.0	(36.1 - 43.9)
Female	414	20,900	43.7	(37.8 - 49.6)	23,491	3,479,500	45.1	(42.0 - 48.2)
Age (years)								
18-49	243	15,000	26.9	(20.5 - 33.4)	12,501	2,726,400	31.6	(27.8 - 35.4)
50-64	217	12,500	52.4	(45.1 - 59.7)	12,964	1,664,200	46.1	(42.7 - 49.5)
≥ 65	209	9,600	67.3	(60.0 - 74.7)	12,594	1,911,700	75.4	(72.2 - 78.6)
Race/Ethnicity								
White Non-Hispanic	649	36,600	40.3	(35.5 - 45.0)	35,750	4,048,700	43.6	(41.7 - 45.5)
Black Non-Hispanic	*	*	*	*	738	910,000	42.2	(33.2 - 51.2)
Hispanic	*	*	*	*	860	903,300	40.6	(31.3 - 49.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	335	15,300	33.7	(27.1 - 40.4)	15,651	2,480,100	44.7	(40.2 - 49.3)
Some College	156	9,500	40.8	(31.4 - 50.1)	9,895	1,387,300	39.2	(34.3 - 44.1)
≥ College Degree	175	12,200	49.1	(39.9 - 58.2)	12,431	2,418,300	43.2	(39.8 - 46.7)
Income								
≤ \$24,999	177	6,200	31.7	(22.9 - 40.4)	9,626	1,675,100	48.2	(41.8 - 54.6)
\$25,000-\$49,999	177	10,600	44.4	(34.5 - 54.3)	9,643	1,208,300	41.7	(35.8 - 47.6)
\$50,000-\$74,999	*	*	*	*	5,761	795,600	40.4	(34.5 - 46.3)
≥ \$75,000	*	*	*	*	8,548	1,791,200	39.9	(36.6 - 43.2)

¹ Defined as having had a flu shot or flu vaccine sprayed in nose within the past 12 months

² Sample size from survey

³ Estimated number of adults who have had a flu shot or flu vaccine sprayed in nose within the past 12 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 20b. Flu shot or flu vaccine sprayed in nose within the past 12 months¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	669	39.3	(34.6 - 44.1)	38,059	42.0	(39.6 - 44.5)
Gender						
Male	255	35.1	(28.0 - 42.2)	14,568	40.4	(36.6 - 44.2)
Female	414	43.4	(37.4 - 49.4)	23,491	43.5	(40.3 - 46.7)
Race/Ethnicity						
White Non-Hispanic	649	40.0	(35.3 - 44.8)	35,750	42.6	(40.7 - 44.6)
Black Non-Hispanic	*	*	*	738	42.4	(33.4 - 51.4)
Hispanic	*	*	*	860	40.0	(30.9 - 49.1)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	335	33.7	(27.0 - 40.3)	15,651	44.1	(39.6 - 48.6)
Some College	156	41.6	(32.1 - 51.1)	9,895	38.1	(33.2 - 43.0)
≥ College Degree	175	47.9	(38.7 - 57.1)	12,431	42.8	(39.4 - 46.3)
Income						
≤ \$24,999	177	31.6	(22.8 - 40.4)	9,626	48.0	(41.6 - 54.3)
\$25,000-\$49,999	177	45.0	(35.2 - 54.7)	9,643	40.5	(34.8 - 46.3)
\$50,000-\$74,999	*	*	*	5,761	39.6	(33.6 - 45.6)
≥ \$75,000	*	*	*	8,548	39.4	(36.1 - 42.7)

¹ Defined as having had a flu shot or flu vaccine sprayed in nose within the past 12 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 21a. Flu shot among adults aged 65 and older¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	209	9,600	67.3	(60.0 - 74.7)	12,602	1,907,700	75.2	(72.0 - 78.4)
Gender								
Male	*	*	*	*	4,417	772,900	74.6	(69.3 - 79.8)
Female	149	5,500	67.0	(58.6 - 75.5)	8,185	1,134,800	75.6	(71.6 - 79.6)
Age (years)								
≥ 65	209	9,600	67.3	(60.0 - 74.7)	12,602	1,907,700	75.2	(72.0 - 78.4)
Race/Ethnicity								
White Non-Hispanic	204	9,400	67.4	(59.9 - 74.9)	12,137	1,487,100	76.0	(73.5 - 78.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	127	5,600	66.3	(56.7 - 76.0)	6,518	930,800	74.0	(68.9 - 79.2)
Some College	*	*	*	*	2,790	317,800	69.7	(62.7 - 76.7)
≥ College Degree	*	*	*	*	3,261	654,000	80.5	(76.4 - 84.6)
Income								
≤ \$24,999	*	*	*	*	4,665	545,300	72.4	(65.6 - 79.3)
\$25,000-\$49,999	*	*	*	*	3,422	430,600	72.8	(67.0 - 78.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	1,198	301,000	77.5	(70.9 - 84.2)

¹ Defined as having had a flu shot during the past 12 months among adults aged 65 and older

² Sample size from survey

³ Estimated number of adults aged 65 and older who have had a flu shot within the past 12 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 21b. Flu shot among adults aged 65 and older¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	209	67.4	(59.7 - 75.1)	12,602	75.0	(71.5 - 78.5)
Gender						
Male	*	*	*	4,417	74.4	(68.7 - 80.1)
Female	149	67.0	(58.6 - 75.5)	8,185	75.5	(71.3 - 79.8)
Race/Ethnicity						
White Non-Hispanic	204	67.4	(59.6 - 75.3)	12,137	75.9	(73.0 - 78.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	6,518	73.3	(67.5 - 79.1)
Some College	*	*	*	2,790	70.0	(62.7 - 77.2)
≥ College Degree	*	*	*	3,261	80.7	(76.2 - 85.1)
Income						
≤ \$24,999	*	*	*	4,665	72.6	(65.1 - 80.1)
\$25,000-\$49,999	*	*	*	3,422	71.4	(65.0 - 77.9)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	1,198	78.0	(70.8 - 85.1)

¹ Defined as having had a flu shot during the past 12 months among adults aged 65 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 22a. Pneumonia shot or pneumococcal vaccine¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	613	23,400	28.9	(24.6 - 33.1)	35,303	3,456,800	26.1	(23.9 - 28.3)
Gender								
Male	223	10,100	26.7	(19.9 - 33.4)	13,052	1,542,200	25.2	(21.5 - 28.8)
Female	390	13,300	30.8	(25.5 - 36.1)	22,251	1,914,600	26.9	(24.2 - 29.6)
Age (years)								
18-49	211	6,900	14.9	(9.2 - 20.6)	10,943	1,097,000	14.6	(11.3 - 17.9)
50-64	195	6,600	31.4	(24.4 - 38.5)	12,179	758,000	23.0	(19.8 - 26.2)
≥ 65	207	10,000	70.1	(63.0 - 77.2)	12,181	1,601,700	65.7	(61.7 - 69.7)
Race/Ethnicity								
White Non-Hispanic	598	23,000	28.8	(24.5 - 33.2)	33,225	2,358,700	28.7	(26.9 - 30.5)
Black Non-Hispanic	*	*	*	*	685	584,400	28.4	(19.5 - 37.3)
Hispanic	*	*	*	*	769	408,000	19.5	(12.6 - 26.4)
Other Non-Hispanic	*	*	*	*	624	105,600	12.2	(2.3 - 22.2)
Education								
≤ High School	308	11,600	31.0	(24.7 - 37.3)	14,700	1,535,300	30.4	(26.4 - 34.3)
Some College	142	5,300	25.5	(16.8 - 34.2)	9,151	836,200	27.2	(21.9 - 32.5)
≥ College Degree	160	6,200	28.3	(20.4 - 36.1)	11,375	1,070,700	21.4	(18.6 - 24.1)
Income								
≤ \$24,999	165	6,200	36.6	(27.4 - 45.9)	9,082	1,050,900	33.1	(27.6 - 38.6)
\$25,000-\$49,999	165	6,800	32.5	(24.0 - 40.9)	9,033	797,500	30.3	(24.1 - 36.6)
\$50,000-\$74,999	82	2,400	18.8	(9.3 - 28.3)	5,320	378,800	21.8	(17.4 - 26.1)
≥ \$75,000	107	4,200	22.1	(14.0 - 30.3)	7,773	648,700	16.3	(14.1 - 18.5)

¹ Defined as having ever had a pneumonia shot or pneumococcal vaccine

² Sample size from survey

³ Estimated number of adults who have ever had a pneumonia shot or pneumococcal vaccine (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 22b. Pneumonia shot or pneumococcal vaccine¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	613	28.7	(24.4 - 32.9)	35,303	25.4	(23.2 - 27.6)
Gender						
Male	223	27.3	(20.6 - 34.1)	13,052	25.8	(22.3 - 29.4)
Female	390	29.8	(24.5 - 35.2)	22,251	25.1	(22.4 - 27.8)
Race/Ethnicity						
White Non-Hispanic	598	28.7	(24.3 - 33.0)	33,225	27.7	(25.9 - 29.5)
Black Non-Hispanic	*	*	*	685	27.9	(19.1 - 36.7)
Hispanic	*	*	*	769	19.6	(12.8 - 26.5)
Other Non-Hispanic	*	*	*	624	11.3	(2.0 - 20.7)
Education						
≤ High School	308	30.8	(24.5 - 37.2)	14,700	29.5	(25.6 - 33.5)
Some College	142	26.1	(17.3 - 34.9)	9,151	26.2	(21.0 - 31.3)
≥ College Degree	160	27.3	(19.5 - 35.0)	11,375	21.0	(18.3 - 23.8)
Income						
≤ \$24,999	165	35.5	(26.3 - 44.7)	9,082	32.3	(26.9 - 37.8)
\$25,000-\$49,999	165	33.7	(25.1 - 42.3)	9,033	29.0	(23.0 - 35.0)
\$50,000-\$74,999	82	18.5	(9.0 - 27.9)	5,320	21.2	(16.8 - 25.7)
≥ \$75,000	107	21.5	(13.2 - 29.7)	7,773	16.2	(13.9 - 18.5)

¹ Defined as having ever had a pneumonia shot or pneumococcal vaccine

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 23a. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	207	10,000	70.1	(63.0 - 77.2)	12,181	1,601,700	65.7	(61.7 - 69.7)
Gender								
Male	*	*	*	*	4,180	606,400	62.1	(55.5 - 68.7)
Female	148	5,800	72.0	(64.2 - 79.9)	8,001	995,300	68.0	(63.1 - 73.0)
Age (years)								
≥ 65	207	10,000	70.1	(63.0 - 77.2)	12,181	1,601,700	65.7	(61.7 - 69.7)
Race/Ethnicity								
White Non-Hispanic	202	9,800	70.2	(63.0 - 77.4)	11,735	1,281,300	68.2	(64.9 - 71.5)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	126	6,000	70.6	(61.6 - 79.6)	6,345	806,200	65.6	(59.5 - 71.7)
Some College	*	*	*	*	2,694	289,500	66.8	(60.1 - 73.4)
≥ College Degree	*	*	*	*	3,111	501,100	65.6	(58.5 - 72.7)
Income								
≤ \$24,999	*	*	*	*	4,533	497,100	67.7	(59.9 - 75.5)
\$25,000-\$49,999	*	*	*	*	3,324	344,300	61.5	(53.3 - 69.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	1,138	236,300	64.5	(55.2 - 73.8)

¹ Defined as having ever had a pneumonia shot or pneumococcal vaccine among adults aged 65 and older

² Sample size from survey

³ Estimated number of adults aged 65 and older who have ever had a pneumonia shot or pneumococcal vaccine (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 23b. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	207	69.8	(62.3 - 77.2)	12,181	64.7	(60.4 - 69.1)
Gender						
Male	*	*	*	4,180	61.5	(54.3 - 68.6)
Female	148	72.0	(64.2 - 79.9)	8,001	67.6	(62.4 - 72.9)
Race/Ethnicity						
White Non-Hispanic	202	69.9	(62.3 - 77.4)	11,735	67.1	(63.2 - 70.9)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	126	70.5	(61.1 - 79.9)	6,345	64.2	(57.4 - 71.0)
Some College	*	*	*	2,694	65.8	(58.4 - 73.3)
≥ College Degree	*	*	*	3,111	65.3	(57.7 - 72.9)
Income						
≤ \$24,999	*	*	*	4,533	66.0	(57.0 - 74.9)
\$25,000-\$49,999	*	*	*	3,324	60.3	(51.9 - 68.6)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	*	*

¹ Defined as having ever had a pneumonia shot or pneumococcal vaccine among adults aged 65 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 24a. Ever had a mammogram among women aged 40 and older¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	327	25,600	90.5	(86.2 - 94.9)	19,491	4,328,800	91.8	(89.7 - 94.0)
Gender								
Male	-	-	-	-	-	-	-	-
Female	327	25,600	90.5	(86.2 - 94.9)	19,491	4,328,800	91.8	(89.7 - 94.0)
Age (years)								
40-49	*	*	*	*	3,851	1,287,500	85.0	(79.7 - 90.3)
50-64	111	10,000	94.0	(88.5 - 99.6)	7,452	1,641,700	96.0	(94.1 - 97.9)
≥ 65	148	7,800	96.9	(94.0 - 99.7)	8,188	1,399,600	94.1	(90.5 - 97.6)
Race/Ethnicity								
White Non-Hispanic	322	25,200	90.4	(86.0 - 94.8)	18,548	2,976,100	93.9	(92.7 - 95.1)
Black Non-Hispanic	*	*	*	*	342	746,600	93.8	(87.3 - 100)
Hispanic	*	*	*	*	333	447,600	91.8	(84.6 - 99.0)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	168	12,100	90.0	(83.4 - 96.6)	8,266	1,698,700	91.0	(87.4 - 94.5)
Some College	76	6,900	89.0	(79.9 - 98.0)	5,322	1,055,500	93.6	(90.2 - 97.0)
≥ College Degree	82	6,500	93.1	(86.2 - 100)	5,872	1,560,700	91.6	(87.5 - 95.6)
Income								
≤ \$24,999	101	6,500	91.2	(84.7 - 97.6)	5,691	1,023,400	93.5	(89.7 - 97.4)
\$25,000-\$49,999	86	6,500	90.0	(81.8 - 98.1)	4,857	860,300	89.5	(83.6 - 95.5)
\$50,000-\$74,999	*	*	*	*	2,648	547,100	94.6	(92.2 - 96.9)
≥ \$75,000	*	*	*	*	3,582	1,185,100	92.2	(88.2 - 96.1)

¹ Defined as having ever had a mammogram among women aged 40 and older

² Sample size from survey

³ Estimated number of women aged 40 and older who have ever had a mammogram (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 24b. Ever had a mammogram among women aged 40 and older¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	327	89.8	(85.1 - 94.5)	19,491	91.1	(88.7 - 93.6)
Gender						
Male	-	-	-	-	-	-
Female	327	89.8	(85.1 - 94.5)	19,491	91.1	(88.7 - 93.6)
Race/Ethnicity						
White Non-Hispanic	322	89.7	(84.9 - 94.4)	18,548	93.3	(92.0 - 94.7)
Black Non-Hispanic	*	*	*	342	93.1	(85.7 - 100)
Hispanic	*	*	*	333	91.0	(82.8 - 99.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	168	90.0	(83.1 - 96.9)	8,266	90.2	(86.1 - 94.2)
Some College	76	87.9	(78.2 - 97.6)	5,322	92.9	(88.9 - 96.9)
≥ College Degree	82	91.5	(83.0 - 100)	5,872	90.9	(86.6 - 95.3)
Income						
≤ \$24,999	101	90.1	(82.8 - 97.5)	5,691	93.0	(88.5 - 97.4)
\$25,000-\$49,999	86	89.5	(80.8 - 98.2)	4,857	88.8	(82.5 - 95.1)
\$50,000-\$74,999	*	*	*	2,648	94.3	(91.7 - 96.8)
≥ \$75,000	*	*	*	3,582	91.2	(86.8 - 95.6)

¹ Defined as having ever had a mammogram among women aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 25a. Had mammogram within the past 2 years among women aged 40 and older¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	324	22,500	79.7	(74.4 - 85.0)	19,376	3,760,000	80.3	(77.8 - 82.9)
Gender								
Male	-	-	-	-	-	-	-	-
Female	324	22,500	79.7	(74.4 - 85.0)	19,376	3,760,000	80.3	(77.8 - 82.9)
Age (years)								
40-49	*	*	*	*	3,842	1,122,700	74.4	(68.5 - 80.2)
50-64	111	8,900	83.4	(75.8 - 90.9)	7,428	1,487,900	87.2	(84.5 - 89.8)
≥ 65	145	6,800	85.9	(79.4 - 92.4)	8,106	1,149,400	78.5	(74.2 - 82.9)
Race/Ethnicity								
White Non-Hispanic	319	22,200	79.9	(74.6 - 85.2)	18,438	2,531,500	80.2	(78.3 - 82.2)
Black Non-Hispanic	*	*	*	*	339	667,100	85.6	(77.5 - 93.8)
Hispanic	*	*	*	*	332	419,700	86.1	(77.5 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	165	10,600	79.4	(71.5 - 87.3)	8,204	1,486,900	80.5	(76.4 - 84.6)
Some College	*	*	*	*	5,297	929,400	83.0	(78.8 - 87.1)
≥ College Degree	82	6,000	84.9	(76.2 - 93.6)	5,845	1,334,800	78.6	(73.9 - 83.2)
Income								
≤ \$24,999	101	5,600	79.1	(69.8 - 88.4)	5,650	882,300	80.8	(75.9 - 85.7)
\$25,000-\$49,999	86	6,000	82.6	(73.3 - 92.0)	4,846	732,300	76.3	(69.7 - 82.9)
\$50,000-\$74,999	*	*	*	*	2,641	494,200	85.6	(81.9 - 89.4)
≥ \$75,000	*	*	*	*	3,572	1,025,900	80.0	(75.2 - 84.8)

¹ Defined as having had a mammogram within the past 2 years among women aged 40 and older

² Sample size from survey

³ Estimated number of women aged 40 and older who have had a mammogram within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 25b. Had mammogram within the past 2 years among women aged 40 and older¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	324	79.1	(73.5 - 84.7)	19,376	79.7	(76.9 - 82.5)
Gender						
Male	-	-	-	-	-	-
Female	324	79.1	(73.5 - 84.7)	19,376	79.7	(76.9 - 82.5)
Race/Ethnicity						
White Non-Hispanic	319	79.4	(73.8 - 85.0)	18,438	79.7	(77.6 - 81.8)
Black Non-Hispanic	*	*	*	339	85.3	(76.5 - 94.2)
Hispanic	*	*	*	332	85.2	(75.6 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	165	78.9	(70.5 - 87.2)	8,204	80.1	(75.6 - 84.7)
Some College	*	*	*	5,297	82.3	(77.7 - 86.9)
≥ College Degree	82	83.6	(73.8 - 93.4)	5,845	77.8	(72.9 - 82.8)
Income						
≤ \$24,999	*	*	*	5,650	80.8	(75.4 - 86.2)
\$25,000-\$49,999	86	82.5	(72.8 - 92.2)	4,846	75.7	(68.7 - 82.6)
\$50,000-\$74,999	*	*	*	2,641	85.5	(81.6 - 89.4)
≥ \$75,000	*	*	*	3,572	79.0	(73.8 - 84.2)

¹ Defined as having had a mammogram within the past 2 years among women aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 26a. Had mammogram within the past 2 years among women aged 50 and older¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	256	15,700	84.4	(79.3 - 89.6)	15,534	2,637,300	83.2	(80.7 - 85.7)
Gender								
Male	-	-	-	-	-	-	-	-
Female	256	15,700	84.4	(79.3 - 89.6)	15,534	2,637,300	83.2	(80.7 - 85.7)
Age (years)								
50-64	111	8,900	83.4	(75.8 - 90.9)	7,428	1,487,900	87.2	(84.5 - 89.8)
≥ 65	145	6,800	85.9	(79.4 - 92.4)	8,106	1,149,400	78.5	(74.2 - 82.9)
Race/Ethnicity								
White Non-Hispanic	252	15,400	84.2	(79.0 - 89.4)	14,887	1,907,000	82.3	(80.3 - 84.4)
Black Non-Hispanic	*	*	*	*	249	385,600	89.9	(83.2 - 96.7)
Hispanic	*	*	*	*	220	286,700	93.9	(89.1 - 98.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	137	7,800	82.9	(75.1 - 90.7)	7,026	1,094,500	83.3	(80.0 - 86.6)
Some College	*	*	*	*	4,124	646,700	87.3	(84.2 - 90.3)
≥ College Degree	61	4,100	92.0	(85.2 - 98.8)	4,356	890,500	80.6	(75.1 - 86.1)
Income								
≤ \$24,999	88	4,700	86.0	(78.4 - 93.7)	4,991	649,100	81.0	(76.8 - 85.2)
\$25,000-\$49,999	66	4,100	87.4	(77.9 - 96.8)	3,913	533,300	78.6	(70.9 - 86.2)
\$50,000-\$74,999	*	*	*	*	1,943	359,300	89.6	(86.3 - 93.0)
≥ \$75,000	*	*	*	*	2,303	601,300	84.6	(79.3 - 89.9)

¹ Defined as having had a mammogram within the past 2 years among women aged 50 and older

² Sample size from survey

³ Estimated number of women aged 50 and older who have had a mammogram within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 26b. Had mammogram within the past 2 years among women aged 50 and older¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	256	84.4	(79.4 - 89.5)	15,534	83.1	(80.3 - 85.8)
Gender						
Male	-	-	-	-	-	-
Female	256	84.4	(79.4 - 89.5)	15,534	83.1	(80.3 - 85.8)
Race/Ethnicity						
White Non-Hispanic	252	84.2	(79.1 - 89.4)	14,887	82.1	(79.9 - 84.3)
Black Non-Hispanic	*	*	*	249	90.4	(83.6 - 97.2)
Hispanic	*	*	*	220	94.1	(89.3 - 99.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	137	83.0	(75.4 - 90.7)	7,026	83.6	(80.1 - 87.0)
Some College	*	*	*	4,124	87.3	(84.1 - 90.5)
≥ College Degree	61	91.8	(84.7 - 98.8)	4,356	79.9	(73.9 - 86.0)
Income						
≤ \$24,999	88	86.0	(78.4 - 93.6)	4,991	81.3	(76.9 - 85.6)
\$25,000-\$49,999	66	87.4	(78.0 - 96.8)	3,913	78.1	(69.9 - 86.4)
\$50,000-\$74,999	*	*	*	1,943	90.2	(87.1 - 93.4)
≥ \$75,000	*	*	*	2,303	83.9	(77.9 - 89.9)

¹ Defined as having had a mammogram within the past 2 years among women aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 27a. Ever had a Pap test among women¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	411	45,400	95.4	(92.0 - 98.7)	23,381	7,045,300	92.1	(90.0 - 94.3)
Gender								
Male	-	-	-	-	-	-	-	-
Female	411	45,400	95.4	(92.0 - 98.7)	23,381	7,045,300	92.1	(90.0 - 94.3)
Age (years)								
18-34	61	13,700	90.6	(81.1 - 100)	2,377	1,896,600	85.2	(79.3 - 91.0)
35-44	*	*	*	*	3,023	1,301,500	92.6	(87.7 - 97.5)
45-54	78	8,800	100	(100 - 100)	4,657	1,420,300	98.3	(97.0 - 99.6)
55-64	78	6,900	100	(100 - 100)	5,187	1,071,200	97.2	(95.0 - 99.3)
≥ 65	147	7,700	95.7	(92.0 - 99.4)	8,137	1,355,700	92.3	(88.5 - 96.1)
Race/Ethnicity								
White Non-Hispanic	401	44,200	95.3	(91.8 - 98.7)	21,976	4,439,700	94.2	(92.1 - 96.3)
Black Non-Hispanic	*	*	*	*	481	1,231,600	91.2	(84.0 - 98.4)
Hispanic	*	*	*	*	533	984,500	90.1	(85.3 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	204	20,200	96.6	(93.5 - 99.8)	9,526	2,540,500	88.3	(83.8 - 92.7)
Some College	96	12,000	93.4	(85.3 - 100)	6,462	1,782,200	93.5	(90.4 - 96.6)
≥ College Degree	108	12,700	95.2	(87.7 - 100)	7,345	2,643,200	95.0	(92.2 - 97.9)
Income								
≤ \$24,999	120	10,200	97.2	(94.2 - 100)	6,526	1,717,200	93.7	(90.9 - 96.6)
\$25,000-\$49,999	100	9,900	98.3	(94.9 - 100)	5,787	1,378,400	91.0	(85.6 - 96.4)
\$50,000-\$74,999	64	10,100	98.3	(95.0 - 100)	3,310	933,900	96.0	(91.8 - 100)
≥ \$75,000	*	*	*	*	4,570	1,943,300	95.4	(92.6 - 98.2)

¹ Defined as having ever had a Pap test among women

² Sample size from survey

³ Estimated number of women who have ever had a Pap test (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 27b. Ever had a Pap test among women¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	411	95.2	(91.8 - 98.6)	23,381	91.8	(89.6 - 94.0)
Gender						
Male	-	-	-	-	-	-
Female	411	95.2	(91.8 - 98.6)	23,381	91.8	(89.6 - 94.0)
Race/Ethnicity						
White Non-Hispanic	401	95.1	(91.6 - 98.6)	21,976	94.0	(91.8 - 96.2)
Black Non-Hispanic	*	*	*	481	90.8	(83.5 - 98.1)
Hispanic	*	*	*	533	89.7	(84.6 - 94.7)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	204	96.4	(93.0 - 99.7)	9,526	87.8	(83.2 - 92.4)
Some College	96	93.2	(85.1 - 100)	6,462	93.1	(89.7 - 96.4)
≥ College Degree	108	95.3	(87.8 - 100)	7,345	94.9	(92.0 - 97.7)
Income						
≤ \$24,999	120	97.2	(94.3 - 100)	6,526	93.3	(90.1 - 96.5)
\$25,000-\$49,999	100	97.9	(93.7 - 100)	5,787	91.1	(85.7 - 96.5)
\$50,000-\$74,999	64	97.9	(93.9 - 100)	3,310	96.0	(91.9 - 100)
≥ \$75,000	58	94.7	(84.7 - 100)	4,570	95.0	(91.9 - 98.1)

¹ Defined as having ever had a Pap test among women

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 28a. Had a Pap test within the past 3 years among women¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	405	38,600	82.0	(77.3 - 86.6)	23,109	6,205,600	82.2	(79.8 - 84.6)
Gender								
Male	-	-	-	-	-	-	-	-
Female	405	38,600	82.0	(77.3 - 86.6)	23,109	6,205,600	82.2	(79.8 - 84.6)
Age (years)								
18-34	*	*	*	*	2,371	1,818,400	81.8	(75.5 - 88.1)
35-44	*	*	*	*	3,012	1,220,300	86.9	(81.8 - 92.0)
45-54	78	7,600	85.4	(76.0 - 94.8)	4,640	1,311,500	92.0	(89.8 - 94.3)
55-64	77	5,600	83.6	(75.0 - 92.3)	5,152	924,300	84.5	(81.0 - 88.0)
≥ 65	142	4,900	63.0	(54.3 - 71.7)	7,934	931,100	66.4	(61.8 - 71.0)
Race/Ethnicity								
White Non-Hispanic	395	37,500	81.6	(76.9 - 86.4)	21,720	3,807,100	81.5	(79.2 - 83.8)
Black Non-Hispanic	*	*	*	*	473	1,142,800	86.7	(79.1 - 94.3)
Hispanic	*	*	*	*	527	894,900	83.6	(76.3 - 91.0)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	199	15,600	76.3	(69.1 - 83.5)	9,384	2,132,600	76.0	(71.1 - 81.0)
Some College	95	10,800	84.5	(75.5 - 93.5)	6,401	1,594,300	84.0	(80.4 - 87.5)
≥ College Degree	108	11,700	87.6	(79.4 - 95.9)	7,281	2,406,000	86.9	(83.7 - 90.1)
Income								
≤ \$24,999	120	8,100	77.1	(68.8 - 85.4)	6,441	1,453,900	82.0	(77.9 - 86.0)
\$25,000-\$49,999	99	8,200	81.6	(73.2 - 90.1)	5,740	1,182,000	78.3	(72.6 - 83.9)
\$50,000-\$74,999	63	9,000	90.0	(81.2 - 98.9)	3,296	825,000	85.0	(78.0 - 91.9)
≥ \$75,000	*	*	*	*	4,555	1,810,200	89.0	(85.7 - 92.2)

¹ Defined as having had a Pap test within the past 3 years among women

² Sample size from survey

³ Estimated number of women who have had a Pap test within the past 3 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 28b. Had a Pap test within the past 3 years among women¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	405	82.0	(77.2 - 86.7)	23,109	82.7	(80.2 - 85.1)
Gender						
Male	-	-	-	-	-	-
Female	405	82.0	(77.2 - 86.7)	23,109	82.7	(80.2 - 85.1)
Race/Ethnicity						
White Non-Hispanic	395	81.6	(76.7 - 86.5)	21,720	82.3	(80.0 - 84.6)
Black Non-Hispanic	*	*	*	473	87.0	(79.3 - 94.6)
Hispanic	*	*	*	527	83.4	(75.9 - 90.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	199	75.7	(68.1 - 83.2)	9,384	76.4	(71.3 - 81.5)
Some College	95	85.3	(76.4 - 94.1)	6,401	84.2	(80.5 - 88.0)
≥ College Degree	108	87.8	(79.6 - 96.0)	7,281	87.5	(84.3 - 90.6)
Income						
≤ \$24,999	120	77.5	(68.9 - 86.1)	6,441	82.7	(78.5 - 86.9)
\$25,000-\$49,999	99	81.4	(72.3 - 90.4)	5,740	79.2	(73.5 - 84.8)
\$50,000-\$74,999	63	89.0	(79.4 - 98.6)	3,296	85.7	(78.7 - 92.8)
≥ \$75,000	*	*	*	4,555	88.9	(85.5 - 92.3)

¹ Defined as having had a Pap test within the past 3 years among women

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 29a. Ever had a digital rectal exam among men aged 40 and older¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	208	21,600	77.7	(71.2 - 84.2)	12,033	3,176,500	73.8	(69.7 - 77.8)
Gender								
Male	208	21,600	77.7	(71.2 - 84.2)	12,033	3,176,500	73.8	(69.7 - 77.8)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	3,098	922,800	66.8	(59.9 - 73.8)
55-64	51	6,200	91.5	(84.0 - 99.1)	3,504	865,600	87.1	(82.5 - 91.7)
≥ 65	58	5,100	84.9	(75.3 - 94.4)	4,348	953,900	92.9	(91.0 - 94.9)
Race/Ethnicity								
White Non-Hispanic	201	21,200	77.7	(71.1 - 84.3)	11,411	2,452,100	80.3	(77.7 - 82.9)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	105	10,200	74.0	(64.0 - 84.0)	5,013	931,600	64.1	(57.3 - 71.0)
Some College	*	*	*	*	2,695	674,200	75.5	(67.8 - 83.2)
≥ College Degree	*	*	*	*	4,302	1,568,000	80.5	(74.0 - 87.0)
Income								
≤ \$24,999	*	*	*	*	2,573	573,300	65.9	(56.0 - 75.9)
\$25,000-\$49,999	*	*	*	*	3,168	530,100	74.6	(67.2 - 82.1)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,309	1,351,100	78.0	(72.6 - 83.5)

¹ Defined as having ever had a digital rectal exam among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have ever had a digital rectal exam (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 29b. Ever had a digital rectal exam among men aged 40 and older¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	208	77.0	(69.9 - 84.1)	12,033	73.7	(69.6 - 77.8)
Gender						
Male	208	77.0	(69.9 - 84.1)	12,033	73.7	(69.6 - 77.8)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	201	76.9	(69.7 - 84.1)	11,411	79.9	(77.2 - 82.5)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	5,013	64.7	(57.8 - 71.7)
Some College	*	*	*	2,695	75.0	(67.1 - 82.9)
≥ College Degree	*	*	*	4,302	80.3	(73.8 - 86.8)
Income						
≤ \$24,999	*	*	*	2,573	66.9	(56.9 - 76.9)
\$25,000-\$49,999	*	*	*	3,168	74.8	(67.3 - 82.2)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,309	77.5	(72.0 - 83.0)

¹ Defined as having ever had a digital rectal exam among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 30a. Had digital rectal exam within the past 2 years among men aged 40 and older¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	206	15,600	56.9	(48.9 - 64.9)	11,923	2,380,300	55.8	(51.8 - 59.8)
Gender								
Male	206	15,600	56.9	(48.9 - 64.9)	11,923	2,380,300	55.8	(51.8 - 59.8)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	1,078	246,800	27.5	(18.5 - 36.5)
45-54	*	*	*	*	3,074	650,000	47.6	(40.8 - 54.4)
55-64	*	*	*	*	3,467	673,500	68.1	(62.4 - 73.8)
≥ 65	*	*	*	*	4,304	809,900	79.7	(75.0 - 84.3)
Race/Ethnicity								
White Non-Hispanic	199	15,400	57.0	(48.9 - 65.0)	11,305	1,875,600	62.2	(59.2 - 65.2)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	4,963	718,700	49.6	(43.2 - 56.1)
Some College	*	*	*	*	2,665	511,500	58.5	(50.8 - 66.3)
≥ College Degree	*	*	*	*	4,277	1,148,400	59.4	(52.8 - 66.0)
Income								
≤ \$24,999	*	*	*	*	2,553	388,000	45.0	(35.0 - 55.0)
\$25,000-\$49,999	*	*	*	*	3,139	422,400	60.5	(53.0 - 68.0)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,295	978,300	56.9	(51.1 - 62.7)

¹ Defined as having had a digital rectal exam within the past 2 years among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have had a digital rectal exam within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 30b. Had digital rectal exam within the past 2 years among men aged 40 and older¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	206	55.4	(46.9 - 63.9)	11,923	55.9	(51.7 - 60.0)
Gender						
Male	206	55.4	(46.9 - 63.9)	11,923	55.9	(51.7 - 60.0)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	199	55.3	(46.7 - 63.9)	11,305	62.0	(58.9 - 65.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	4,963	50.5	(43.7 - 57.3)
Some College	*	*	*	2,665	58.3	(50.4 - 66.2)
≥ College Degree	*	*	*	4,277	59.2	(52.4 - 65.9)
Income						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,139	60.8	(53.1 - 68.4)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,295	56.5	(50.5 - 62.4)

¹ Defined as having had a digital rectal exam within the past 2 years among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 31a. Ever had a prostate specific antigen test among men aged 40 and older¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	192	15,400	61.3	(52.8 - 69.7)	11,557	2,822,600	67.6	(63.9 - 71.2)
Gender								
Male	192	15,400	61.3	(52.8 - 69.7)	11,557	2,822,600	67.6	(63.9 - 71.2)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,938	760,600	57.4	(50.6 - 64.3)
55-64	*	*	*	*	3,373	811,400	83.4	(78.8 - 88.1)
≥ 65	56	5,100	89.3	(80.4 - 98.2)	4,215	905,400	90.4	(87.1 - 93.6)
Race/Ethnicity								
White Non-Hispanic	184	15,000	61.1	(52.5 - 69.6)	10,950	1,965,400	67.0	(63.9 - 70.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	4,775	829,200	59.0	(52.4 - 65.6)
Some College	*	*	*	*	2,584	535,000	61.9	(53.7 - 70.2)
≥ College Degree	*	*	*	*	4,176	1,455,700	76.8	(72.5 - 81.1)
Income								
≤ \$24,999	*	*	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	*	3,048	444,100	64.1	(56.0 - 72.3)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,188	1,211,100	72.3	(67.6 - 77.0)

¹ Defined as having ever had a prostate specific antigen test among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have ever had a prostate specific antigen test (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 31b. Ever had a prostate specific antigen test among men aged 40 and older¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	192	59.4	(50.3 - 68.5)	11,557	67.4	(63.7 - 71.1)
Gender						
Male	192	59.4	(50.3 - 68.5)	11,557	67.4	(63.7 - 71.1)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	184	59.0	(49.7 - 68.2)	10,950	66.6	(63.4 - 69.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	4,775	59.7	(52.9 - 66.4)
Some College	*	*	*	2,584	61.4	(53.0 - 69.9)
≥ College Degree	*	*	*	4,176	76.3	(71.9 - 80.7)
Income						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,048	64.2	(56.1 - 72.4)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,188	71.5	(66.6 - 76.3)

¹ Defined as having ever had a prostate specific antigen test among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 32a. Had prostate specific antigen test within the past 2 years among men aged 40 and older¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	189	13,400	54.3	(45.8 - 62.7)	11,437	2,493,800	59.9	(56.0 - 63.9)
Gender								
Male	189	13,400	54.3	(45.8 - 62.7)	11,437	2,493,800	59.9	(56.0 - 63.9)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,917	659,400	49.9	(42.8 - 57.0)
55-64	*	*	*	*	3,326	719,700	74.5	(69.4 - 79.7)
≥ 65	*	*	*	*	4,166	830,000	83.3	(78.4 - 88.3)
Race/Ethnicity								
White Non-Hispanic	181	13,200	54.3	(45.8 - 62.9)	10,836	1,762,200	60.4	(57.3 - 63.5)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	4,727	734,200	52.4	(45.7 - 59.1)
Some College	*	*	*	*	2,558	483,800	56.3	(48.2 - 64.4)
≥ College Degree	*	*	*	*	4,133	1,273,200	67.4	(61.6 - 73.3)
Income								
≤ \$24,999	*	*	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	*	3,021	410,800	59.6	(51.5 - 67.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,170	1,080,100	64.6	(59.1 - 70.0)

¹ Defined as having had a prostate specific antigen test within the past 2 years among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have had a prostate specific antigen test within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 32b. Had prostate specific antigen test within the past 2 years among men aged 40 and older¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	189	52.9	(44.0 - 61.9)	11,437	59.8	(55.8 - 63.9)
Gender						
Male	189	52.9	(44.0 - 61.9)	11,437	59.8	(55.8 - 63.9)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	181	52.9	(43.8 - 62.0)	10,836	60.1	(56.9 - 63.4)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	4,727	52.9	(45.9 - 59.9)
Some College	*	*	*	2,558	56.0	(47.7 - 64.2)
≥ College Degree	*	*	*	4,133	67.1	(61.2 - 73.1)
Income						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,021	59.8	(51.7 - 67.9)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,170	63.9	(58.3 - 69.6)

¹ Defined as having had a prostate specific antigen test within the past 2 years among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 33a. Prostate cancer among men aged 40 and older¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	*	*	*	*	12,133	178,200	4.1	(3.0 - 5.2)
Gender								
Male	*	*	*	*	12,133	178,200	4.1	(3.0 - 5.2)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	3,120	6,100	0.4	(0.0 - 0.9)
55-64	*	*	*	*	3,532	31,600	3.1	(1.5 - 4.8)
≥ 65	*	*	*	*	4,388	140,400	13.6	(9.7 - 17.4)
Race/Ethnicity								
White Non-Hispanic	*	*	*	*	11,502	143,700	4.7	(3.5 - 5.8)
Black Non-Hispanic	*	*	*	*	184	23,500	5.1	(0.0 - 11.0)
Hispanic	*	*	*	*	197	7,700	1.6	(0.0 - 3.7)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	5,065	46,300	3.1	(2.2 - 4.1)
Some College	*	*	*	*	2,709	39,200	4.4	(1.3 - 7.5)
≥ College Degree	*	*	*	*	4,336	92,500	4.7	(3.0 - 6.5)
Income								
≤ \$24,999	*	*	*	*	2,597	35,800	4.1	(2.2 - 6.0)
\$25,000-\$49,999	*	*	*	*	3,196	29,600	4.1	(2.9 - 5.4)
\$50,000-\$74,999	*	*	*	*	2,007	20,300	2.8	(1.2 - 4.4)
≥ \$75,000	*	*	*	*	3,322	56,500	3.2	(1.7 - 4.8)

¹ Defined as having ever been told by a health professional that respondent had prostate cancer among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have ever been told they had prostate cancer (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 33b. Prostate cancer among men aged 40 and older¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	*	*	*	12,133	4.5	(3.2 - 5.7)
Gender						
Male	*	*	*	12,133	4.5	(3.2 - 5.7)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	*	*	*	11,502	5.1	(3.8 - 6.4)
Black Non-Hispanic	*	*	*	184	5.7	(0.0 - 12.8)
Hispanic	*	*	*	197	1.8	(0.0 - 4.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	5,065	3.3	(2.3 - 4.3)
Some College	*	*	*	2,709	5.0	(1.1 - 9.0)
≥ College Degree	*	*	*	4,336	5.2	(3.1 - 7.2)
Income						
≤ \$24,999	*	*	*	2,597	4.5	(2.2 - 6.7)
\$25,000-\$49,999	*	*	*	3,196	4.3	(3.0 - 5.6)
\$50,000-\$74,999	*	*	*	2,007	3.0	(1.1 - 5.0)
≥ \$75,000	*	*	*	3,322	3.5	(1.6 - 5.4)

¹ Defined as having ever been told by a health professional that respondent had prostate cancer among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 34a. Home blood stool test ever used among adults aged 50 and older¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	422	18,800	49.6	(44.1 - 55.0)	25,322	2,288,500	38.0	(35.5 - 40.4)
Gender								
Male	162	9,400	49.4	(41.0 - 57.8)	9,571	1,081,900	38.9	(35.1 - 42.6)
Female	260	9,400	49.8	(42.9 - 56.7)	15,751	1,206,600	37.2	(34.0 - 40.4)
Age (years)								
50-64	216	10,500	44.2	(36.9 - 51.5)	12,834	1,192,000	34.0	(30.8 - 37.2)
≥ 65	206	8,300	58.6	(50.9 - 66.3)	12,472	1,093,400	43.6	(39.8 - 47.4)
Race/Ethnicity								
White Non-Hispanic	413	18,500	49.4	(43.9 - 54.9)	24,213	1,754,500	39.1	(37.1 - 41.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	218	8,100	44.5	(37.0 - 52.0)	11,146	823,500	34.9	(30.5 - 39.2)
Some College	*	*	*	*	6,261	480,300	37.0	(32.6 - 41.4)
≥ College Degree	*	*	*	*	7,851	973,700	41.6	(37.8 - 45.3)
Income								
≤ \$24,999	129	4,700	48.5	(38.8 - 58.2)	7,222	505,000	35.4	(29.7 - 41.2)
\$25,000-\$49,999	*	*	*	*	6,609	467,600	40.3	(35.2 - 45.4)
\$50,000-\$74,999	*	*	*	*	3,500	313,000	38.2	(32.6 - 43.9)
≥ \$75,000	*	*	*	*	4,647	657,800	37.3	(33.0 - 41.6)

¹ Defined as having ever used a home blood stool test among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have ever used a home blood stool test (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 34b. Home blood stool test ever used among adults aged 50 and older¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	422	50.2	(44.8 - 55.7)	25,322	38.1	(35.5 - 40.8)
Gender						
Male	162	50.2	(41.8 - 58.6)	9,571	39.4	(35.3 - 43.5)
Female	260	50.2	(43.4 - 57.0)	15,751	36.9	(33.5 - 40.3)
Race/Ethnicity						
White Non-Hispanic	413	50.0	(44.5 - 55.5)	24,213	38.9	(36.7 - 41.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	218	45.5	(37.9 - 53.1)	11,146	35.1	(30.3 - 39.9)
Some College	*	*	*	6,261	36.6	(32.0 - 41.3)
≥ College Degree	*	*	*	7,851	41.9	(37.9 - 46.0)
Income						
≤ \$24,999	129	49.5	(39.9 - 59.1)	7,222	35.9	(29.5 - 42.4)
\$25,000-\$49,999	*	*	*	6,609	40.2	(34.8 - 45.6)
\$50,000-\$74,999	*	*	*	3,500	38.1	(31.9 - 44.3)
≥ \$75,000	*	*	*	4,647	37.5	(32.9 - 42.2)

¹ Defined as having ever used a home blood stool test among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 35a. Home blood stool test used within the past year among adults aged 50 and older¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	414	7,200	19.4	(15.1 - 23.7)	25,074	863,300	14.4	(12.6 - 16.2)
Gender								
Male	161	3,900	20.4	(13.7 - 27.1)	9,508	444,400	16.0	(13.2 - 18.9)
Female	253	3,400	18.3	(13.0 - 23.7)	15,566	418,900	13.1	(10.8 - 15.4)
Age (years)								
50-64	213	3,900	16.8	(11.2 - 22.3)	12,759	454,400	13.0	(10.6 - 15.5)
≥ 65	201	3,300	23.9	(17.1 - 30.6)	12,299	406,400	16.4	(13.7 - 19.0)
Race/Ethnicity								
White Non-Hispanic	405	7,100	19.2	(14.9 - 23.6)	23,971	634,300	14.3	(12.8 - 15.7)
Black Non-Hispanic	*	*	*	*	396	136,000	18.6	(9.4 - 27.8)
Hispanic	*	*	*	*	349	53,400	9.9	(2.5 - 17.2)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	215	3,500	19.2	(13.3 - 25.1)	11,026	288,600	12.3	(9.4 - 15.3)
Some College	93	1,800	20.4	(10.8 - 30.1)	6,208	212,500	16.4	(12.5 - 20.4)
≥ College Degree	105	1,900	19.0	(10.7 - 27.2)	7,778	352,400	15.2	(12.5 - 17.9)
Income								
≤ \$24,999	128	1,900	19.9	(12.1 - 27.6)	7,157	196,300	13.9	(10.1 - 17.6)
\$25,000-\$49,999	109	2,400	24.9	(16.0 - 33.8)	6,562	221,500	19.2	(14.1 - 24.3)
\$50,000-\$74,999	*	*	*	*	3,469	111,900	13.7	(10.1 - 17.4)
≥ \$75,000	*	*	*	*	4,623	205,100	11.7	(8.8 - 14.6)

¹ Defined as having used a home blood stool test within the past year among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have used a home blood stool test within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 35b. Home blood stool test used within the past year among adults aged 50 and older¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	414	19.9	(15.5 - 24.3)	25,074	14.6	(12.6 - 16.7)
Gender						
Male	161	21.2	(14.3 - 28.1)	9,508	16.4	(13.2 - 19.6)
Female	253	18.5	(13.2 - 23.8)	15,566	12.9	(10.4 - 15.4)
Race/Ethnicity						
White Non-Hispanic	405	19.8	(15.3 - 24.2)	23,971	14.2	(12.7 - 15.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	349	10.2	(1.7 - 18.7)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	215	20.2	(14.0 - 26.4)	11,026	12.7	(9.2 - 16.1)
Some College	93	19.6	(10.3 - 28.9)	6,208	16.6	(12.3 - 20.9)
≥ College Degree	105	19.9	(11.4 - 28.4)	7,778	15.3	(12.4 - 18.3)
Income						
≤ \$24,999	128	20.7	(12.9 - 28.5)	7,157	14.3	(9.8 - 18.7)
\$25,000-\$49,999	109	25.9	(16.5 - 35.4)	6,562	19.3	(13.8 - 24.8)
\$50,000-\$74,999	*	*	*	3,469	13.8	(9.8 - 17.7)
≥ \$75,000	*	*	*	4,623	11.8	(8.6 - 15.0)

¹ Defined as having used a home blood stool test within the past year among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 36a. Home blood stool test used within the past 2 years among adults aged 50 and older¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	414	10,500	28.1	(23.3 - 33.0)	25,074	1,218,000	20.4	(18.3 - 22.5)
Gender								
Male	161	5,500	29.0	(21.4 - 36.6)	9,508	593,100	21.4	(18.3 - 24.5)
Female	253	5,000	27.3	(21.1 - 33.4)	15,566	624,900	19.5	(16.6 - 22.4)
Age (years)								
50-64	213	5,800	24.6	(18.2 - 30.9)	12,759	629,900	18.1	(15.4 - 20.7)
≥ 65	201	4,700	34.3	(26.9 - 41.7)	12,299	585,500	23.6	(20.2 - 27.0)
Race/Ethnicity								
White Non-Hispanic	405	10,300	28.0	(23.1 - 33.0)	23,971	877,500	19.7	(18.2 - 21.3)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	215	5,100	28.4	(21.6 - 35.1)	11,026	454,600	19.4	(15.6 - 23.3)
Some College	*	*	*	*	6,208	262,200	20.3	(16.3 - 24.2)
≥ College Degree	105	2,900	28.6	(18.9 - 38.3)	7,778	491,400	21.2	(18.1 - 24.2)
Income								
≤ \$24,999	128	2,500	26.0	(17.6 - 34.3)	7,157	263,900	18.6	(14.6 - 22.7)
\$25,000-\$49,999	109	3,200	33.6	(23.8 - 43.4)	6,562	285,900	24.8	(19.6 - 30.0)
\$50,000-\$74,999	*	*	*	*	3,469	167,500	20.6	(15.6 - 25.5)
≥ \$75,000	*	*	*	*	4,623	294,900	16.8	(13.7 - 20.0)

¹ Defined as having used a home blood stool test within the past 2 years among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have used a home blood stool test within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 36b. Home blood stool test used within the past 2 years among adults aged 50 and older¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	414	28.6	(23.7 - 33.6)	25,074	20.5	(18.2 - 22.8)
Gender						
Male	161	29.7	(22.1 - 37.3)	9,508	21.7	(18.3 - 25.1)
Female	253	27.5	(21.4 - 33.6)	15,566	19.4	(16.3 - 22.5)
Race/Ethnicity						
White Non-Hispanic	405	28.6	(23.6 - 33.5)	23,971	19.7	(18.0 - 21.3)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	215	29.5	(22.5 - 36.4)	11,026	19.5	(15.4 - 23.7)
Some College	*	*	*	6,208	20.4	(16.1 - 24.7)
≥ College Degree	105	29.2	(19.5 - 38.9)	7,778	21.4	(18.1 - 24.7)
Income						
≤ \$24,999	128	27.1	(18.7 - 35.5)	7,157	18.9	(14.2 - 23.6)
\$25,000-\$49,999	*	*	*	6,562	24.8	(19.3 - 30.3)
\$50,000-\$74,999	*	*	*	3,469	20.6	(15.3 - 25.9)
≥ \$75,000	*	*	*	4,623	16.9	(13.4 - 20.4)

¹ Defined as having used a home blood stool test within the past 2 years among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 37a. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	422	24,500	64.6	(59.4 - 69.8)	25,357	4,019,100	66.6	(64.1 - 69.1)
Gender								
Male	162	12,300	64.3	(56.3 - 72.4)	9,582	1,894,100	68.0	(64.2 - 71.7)
Female	260	12,200	64.9	(58.3 - 71.5)	15,775	2,125,000	65.5	(62.1 - 68.8)
Age (years)								
50-54	*	*	*	*	4,179	718,900	50.6	(44.6 - 56.5)
55-64	129	10,000	72.9	(64.6 - 81.1)	8,695	1,493,500	71.0	(67.3 - 74.8)
≥ 65	206	10,200	72.8	(66.2 - 79.4)	12,466	1,802,000	72.1	(68.6 - 75.7)
Race/Ethnicity								
White Non-Hispanic	412	24,100	64.6	(59.4 - 69.9)	24,248	3,093,900	68.8	(66.8 - 70.8)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	218	11,300	61.9	(54.5 - 69.3)	11,121	1,445,100	61.2	(56.7 - 65.8)
Some College	*	*	*	*	6,287	835,300	64.3	(59.7 - 69.0)
≥ College Degree	*	*	*	*	7,884	1,732,400	73.7	(70.0 - 77.4)
Income								
≤ \$24,999	129	5,500	57.2	(47.5 - 66.8)	7,220	892,600	62.7	(56.8 - 68.6)
\$25,000-\$49,999	113	7,000	70.2	(60.7 - 79.7)	6,619	697,300	60.1	(54.6 - 65.7)
\$50,000-\$74,999	*	*	*	*	3,514	613,700	74.2	(68.9 - 79.6)
≥ \$75,000	*	*	*	*	4,665	1,294,400	73.3	(69.2 - 77.5)

¹ Defined as having ever had a sigmoidoscopy or colonoscopy among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have ever had a sigmoidoscopy or colonoscopy (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 37b. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	422	65.3	(60.2 - 70.5)	25,357	66.6	(63.9 - 69.3)
Gender						
Male	162	65.8	(57.9 - 73.6)	9,582	68.5	(64.5 - 72.5)
Female	260	64.8	(58.2 - 71.4)	15,775	64.8	(61.2 - 68.4)
Race/Ethnicity						
White Non-Hispanic	412	65.3	(60.0 - 70.5)	24,248	68.7	(66.5 - 70.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	218	63.1	(55.8 - 70.4)	11,121	61.3	(56.4 - 66.2)
Some College	*	*	*	6,287	63.8	(58.8 - 68.8)
≥ College Degree	*	*	*	7,884	74.0	(70.2 - 77.9)
Income						
≤ \$24,999	129	58.2	(48.8 - 67.6)	7,220	63.0	(56.7 - 69.3)
\$25,000-\$49,999	113	71.3	(61.8 - 80.7)	6,619	59.7	(53.8 - 65.6)
\$50,000-\$74,999	*	*	*	3,514	74.4	(68.3 - 80.5)
≥ \$75,000	*	*	*	4,665	73.6	(69.3 - 78.0)

¹ Defined as having ever had a sigmoidoscopy or colonoscopy among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 38a. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	417	23,400	62.1	(56.8 - 67.4)	25,169	3,855,800	64.6	(62.1 - 67.1)
Gender								
Male	162	11,900	62.6	(54.5 - 70.8)	9,521	1,835,400	66.4	(62.6 - 70.1)
Female	255	11,400	61.5	(54.7 - 68.3)	15,648	2,020,400	63.1	(59.8 - 66.4)
Age (years)								
50-54	*	*	*	*	4,169	700,500	49.3	(43.3 - 55.3)
55-64	127	9,700	71.1	(62.7 - 79.6)	8,652	1,455,100	69.4	(65.7 - 73.2)
≥ 65	203	9,500	68.2	(61.0 - 75.3)	12,331	1,695,700	69.5	(65.9 - 73.1)
Race/Ethnicity								
White Non-Hispanic	407	23,000	62.0	(56.7 - 67.4)	24,069	2,981,500	66.6	(64.6 - 68.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	213	10,400	58.0	(50.4 - 65.6)	11,017	1,360,400	59.0	(54.5 - 63.5)
Some College	*	*	*	*	6,251	807,100	62.4	(57.8 - 67.1)
≥ College Degree	*	*	*	*	7,841	1,682,500	71.7	(68.0 - 75.4)
Income								
≤ \$24,999	129	5,400	56.0	(46.4 - 65.7)	7,163	858,900	61.0	(55.1 - 67.0)
\$25,000-\$49,999	111	6,900	70.0	(60.4 - 79.6)	6,584	664,900	57.7	(52.2 - 63.2)
\$50,000-\$74,999	*	*	*	*	3,501	602,600	73.1	(67.7 - 78.5)
≥ \$75,000	*	*	*	*	4,655	1,260,400	71.5	(67.3 - 75.7)

¹ Defined as having had a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have had a sigmoidoscopy or colonoscopy within the past 10 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 38b. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	417	62.6	(57.3 - 67.9)	25,169	64.6	(61.9 - 67.3)
Gender						
Male	162	63.9	(55.9 - 71.9)	9,521	66.9	(62.9 - 70.9)
Female	255	61.2	(54.5 - 68.0)	15,648	62.4	(58.9 - 66.0)
Race/Ethnicity						
White Non-Hispanic	407	62.5	(57.2 - 67.9)	24,069	66.5	(64.3 - 68.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	213	59.0	(51.4 - 66.6)	11,017	59.0	(54.1 - 63.9)
Some College	*	*	*	6,251	61.9	(56.9 - 66.9)
≥ College Degree	*	*	*	7,841	72.1	(68.2 - 75.9)
Income						
≤ \$24,999	129	57.0	(47.6 - 66.5)	7,163	61.3	(54.9 - 67.6)
\$25,000-\$49,999	111	71.1	(61.5 - 80.6)	6,584	57.2	(51.4 - 63.0)
\$50,000-\$74,999	*	*	*	3,501	73.3	(67.2 - 79.4)
≥ \$75,000	*	*	*	4,655	71.8	(67.4 - 76.2)

¹ Defined as having had a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 39a. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	412	25,500	68.5	(63.3 - 73.7)	24,792	4,049,200	69.1	(66.7 - 71.6)
Gender								
Male	161	13,200	69.7	(61.9 - 77.5)	9,418	1,945,200	70.7	(67.0 - 74.3)
Female	251	12,300	67.2	(60.5 - 73.9)	15,374	2,104,100	67.8	(64.6 - 71.1)
Age (years)								
50-54	*	*	*	*	4,132	751,200	54.1	(48.0 - 60.1)
55-64	125	9,900	73.8	(65.5 - 82.1)	8,366	1,487,600	73.7	(70.0 - 77.4)
≥ 65	202	10,800	77.2	(70.9 - 83.6)	12,294	1,810,500	74.0	(70.6 - 77.4)
Race/Ethnicity								
White Non-Hispanic	402	25,000	68.5	(63.3 - 73.7)	23,726	3,091,400	70.4	(68.4 - 72.4)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	212	11,900	66.0	(58.6 - 73.4)	10,872	1,461,600	64.2	(59.8 - 68.7)
Some College	*	*	*	*	6,170	873,300	68.4	(63.9 - 72.8)
≥ College Degree	*	*	*	*	7,708	1,709,000	74.6	(70.9 - 78.3)
Income								
≤ \$24,999	127	6,100	63.8	(54.3 - 73.3)	7,092	923,300	66.0	(60.1 - 71.9)
\$25,000-\$49,999	111	7,700	78.3	(69.6 - 87.1)	6,526	730,000	64.2	(58.9 - 69.6)
\$50,000-\$74,999	*	*	*	*	3,467	621,500	76.6	(71.4 - 81.9)
≥ \$75,000	*	*	*	*	4,615	1,292,500	73.8	(69.8 - 77.9)

¹ Defined as having used a home blood stool test within the past year or a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 years and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have used a home blood stool test in the past year or sigmoidoscopy or colonoscopy within the past 10 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 39b. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	412	69.3	(64.3 - 74.4)	24,792	69.3	(66.7 - 71.9)
Gender						
Male	161	71.2	(63.7 - 78.8)	9,418	71.5	(67.6 - 75.4)
Female	251	67.3	(60.6 - 74.0)	15,374	67.2	(63.7 - 70.7)
Race/Ethnicity						
White Non-Hispanic	402	69.3	(64.1 - 74.4)	23,726	70.3	(68.1 - 72.4)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	212	67.4	(60.1 - 74.6)	10,872	64.5	(59.8 - 69.3)
Some College	*	*	*	6,170	68.2	(63.4 - 72.9)
≥ College Degree	106	72.4	(62.5 - 82.3)	7,708	75.0	(71.2 - 78.8)
Income						
≤ \$24,999	127	65.1	(55.9 - 74.3)	7,092	66.4	(60.2 - 72.6)
\$25,000-\$49,999	111	79.2	(70.7 - 87.8)	6,526	63.8	(58.2 - 69.5)
\$50,000-\$74,999	*	*	*	3,467	76.7	(70.6 - 82.7)
≥ \$75,000	*	*	*	4,615	74.4	(70.2 - 78.6)

¹ Defined as having used a home blood stool test within the past year or a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 years and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 40a. High blood pressure among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	666	26,700	28.5	(24.2 - 32.9)	38,125	3,974,800	26.8	(24.9 - 28.7)
Gender								
Male	255	15,200	33.0	(25.4 - 40.5)	14,611	1,960,500	27.7	(24.7 - 30.7)
Female	411	11,500	24.2	(19.8 - 28.6)	23,514	2,014,300	26.0	(23.6 - 28.3)
Age (years)								
18-34	*	*	*	*	3,935	312,200	7.0	(4.2 - 9.8)
35-44	80	2,600	15.5	(6.4 - 24.6)	4,984	426,200	14.8	(11.6 - 18.1)
45-54	158	4,800	25.8	(18.5 - 33.0)	7,825	862,400	30.3	(25.9 - 34.8)
55-64	128	6,400	47.1	(37.6 - 56.7)	8,765	890,800	41.6	(37.6 - 45.7)
≥ 65	208	9,200	64.4	(57.0 - 71.8)	12,616	1,483,200	58.5	(54.7 - 62.3)
Race/Ethnicity								
White Non-Hispanic	646	24,900	27.5	(23.5 - 31.5)	35,814	2,650,900	28.4	(26.9 - 30.0)
Black Non-Hispanic	*	*	*	*	736	558,500	26.2	(19.6 - 32.8)
Hispanic	*	*	*	*	863	538,100	23.8	(17.0 - 30.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	333	14,600	32.4	(25.3 - 39.5)	15,688	1,851,200	33.1	(29.4 - 36.9)
Some College	155	6,200	26.5	(18.6 - 34.5)	9,906	851,100	24.1	(20.9 - 27.3)
≥ College Degree	175	5,800	23.4	(16.6 - 30.2)	12,452	1,253,400	22.3	(19.7 - 24.9)
Income								
≤ \$24,999	*	*	*	*	9,640	1,153,000	32.8	(27.6 - 38.1)
\$25,000-\$49,999	176	8,200	34.3	(25.8 - 42.9)	9,657	807,800	27.9	(23.7 - 32.1)
\$50,000-\$74,999	100	3,700	23.1	(13.3 - 32.8)	5,773	507,600	25.8	(21.4 - 30.1)
≥ \$75,000	116	3,700	17.6	(10.3 - 24.9)	8,560	942,500	21.0	(18.4 - 23.6)

¹ Defined as having ever been told by a health professional that respondent has high blood pressure

² Sample size from survey

³ Estimated number of adults who have ever been told by a health professional that they have high blood pressure (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 40b. High blood pressure among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	666	28.4	(24.0 - 32.7)	38,125	25.7	(23.8 - 27.5)
Gender						
Male	255	33.9	(26.3 - 41.5)	14,611	27.7	(24.7 - 30.7)
Female	411	23.0	(18.7 - 27.3)	23,514	23.8	(21.5 - 26.0)
Race/Ethnicity						
White Non-Hispanic	646	27.4	(23.3 - 31.4)	35,814	27.2	(25.6 - 28.7)
Black Non-Hispanic	*	*	*	736	24.9	(18.4 - 31.3)
Hispanic	*	*	*	863	23.3	(16.6 - 30.0)
Other Non-Hispanic	*	*	*	712	19.3	(9.5 - 29.1)
Education						
≤ High School	333	32.9	(25.8 - 40.1)	15,688	32.1	(28.4 - 35.8)
Some College	155	25.6	(17.9 - 33.4)	9,906	22.8	(19.7 - 25.9)
≥ College Degree	175	22.7	(15.9 - 29.5)	12,452	21.2	(18.6 - 23.8)
Income						
≤ \$24,999	*	*	*	9,640	31.8	(26.7 - 37.0)
\$25,000-\$49,999	176	35.0	(26.4 - 43.7)	9,657	26.9	(22.8 - 30.9)
\$50,000-\$74,999	100	22.2	(12.3 - 32.0)	5,773	24.8	(20.4 - 29.2)
≥ \$75,000	116	16.0	(9.0 - 23.1)	8,560	19.9	(17.3 - 22.5)

¹ Defined as having ever been told by a health professional that respondent has high blood pressure

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 41a. Blood pressure medication among adults with high blood pressure¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	258	21,000	78.6	(68.6 - 88.6)	15,288	3,145,800	79.2	(76.0 - 82.4)
Gender								
Male	*	*	*	*	6,080	1,432,100	73.1	(67.6 - 78.5)
Female	152	10,300	89.6	(84.1 - 95.0)	9,208	1,713,700	85.2	(82.1 - 88.2)
Age (years)								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,315	645,500	74.9	(66.8 - 82.9)
55-64	60	6,000	94.2	(88.3 - 100)	4,045	782,300	87.8	(84.5 - 91.1)
≥ 65	135	8,500	92.1	(87.7 - 96.6)	7,675	1,383,400	93.4	(91.1 - 95.7)
Race/Ethnicity								
White Non-Hispanic	250	20,500	82.4	(75.6 - 89.2)	14,442	2,117,300	79.9	(77.5 - 82.4)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	7,354	1,482,800	80.2	(74.7 - 85.6)
Some College	*	*	*	*	3,868	632,000	74.3	(67.8 - 80.7)
≥ College Degree	*	*	*	*	4,035	1,013,200	80.9	(76.8 - 84.9)
Income								
≤ \$24,999	*	*	*	*	4,898	948,200	82.3	(75.7 - 88.9)
\$25,000-\$49,999	*	*	*	*	4,088	668,300	82.8	(77.6 - 88.0)
\$50,000-\$74,999	*	*	*	*	1,971	391,500	77.2	(67.7 - 86.6)
≥ \$75,000	*	*	*	*	2,397	683,700	72.5	(66.1 - 78.9)

¹ Defined as currently taking medicine for high blood pressure among adults with high blood pressure

² Sample size from survey

³ Estimated number of adults with high blood pressure who are taking medicine for high blood pressure (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 41b. Blood pressure medication among adults with high blood pressure¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	258	79.3	(69.6 - 89.0)	15,288	78.5	(75.2 - 81.7)
Gender						
Male	*	*	*	6,080	73.3	(67.9 - 78.7)
Female	152	89.1	(83.3 - 94.8)	9,208	84.0	(80.7 - 87.2)
Race/Ethnicity						
White Non-Hispanic	250	82.9	(76.3 - 89.6)	14,442	79.0	(76.5 - 81.6)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	7,354	79.7	(74.3 - 85.2)
Some College	*	*	*	3,868	72.5	(65.7 - 79.3)
≥ College Degree	*	*	*	4,035	80.4	(76.1 - 84.6)
Income						
≤ \$24,999	*	*	*	4,898	81.6	(75.0 - 88.3)
\$25,000-\$49,999	*	*	*	4,088	82.1	(76.7 - 87.5)
\$50,000-\$74,999	*	*	*	1,971	76.4	(66.7 - 86.1)
≥ \$75,000	*	*	*	2,397	71.6	(64.8 - 78.3)

¹ Defined as currently taking medicine for high blood pressure among adults with high blood pressure

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 42a. Ever had blood cholesterol checked among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	655	71,900	79.1	(73.9 - 84.3)	37,450	11,496,600	80.3	(77.7 - 83.0)
Gender								
Male	250	33,800	76.0	(67.1 - 84.9)	14,389	5,288,400	77.7	(73.3 - 82.0)
Female	405	38,200	82.1	(76.8 - 87.4)	23,061	6,208,200	82.7	(79.6 - 85.8)
Age (years)								
18-34	*	*	*	*	3,684	2,296,400	55.3	(48.7 - 61.9)
35-44	78	14,700	90.4	(83.6 - 97.2)	4,883	2,264,200	81.6	(76.6 - 86.5)
45-54	156	16,200	88.2	(83.2 - 93.1)	7,731	2,539,400	91.0	(87.0 - 94.9)
55-64	129	12,900	94.2	(89.4 - 98.9)	8,718	2,014,900	95.4	(93.5 - 97.2)
≥ 65	206	13,500	95.5	(92.2 - 98.7)	12,434	2,381,700	95.9	(94.3 - 97.6)
Race/Ethnicity								
White Non-Hispanic	635	70,500	80.3	(75.5 - 85.1)	35,198	7,832,600	86.3	(84.8 - 87.9)
Black Non-Hispanic	*	*	*	*	720	1,588,900	77.8	(68.4 - 87.2)
Hispanic	*	*	*	*	842	1,224,600	56.8	(46.8 - 66.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	326	31,500	72.9	(64.3 - 81.5)	15,342	3,795,700	70.6	(65.6 - 75.6)
Some College	154	18,700	81.5	(72.7 - 90.2)	9,732	2,599,800	77.4	(71.7 - 83.2)
≥ College Degree	172	21,300	87.6	(80.3 - 94.9)	12,296	5,008,300	91.3	(88.9 - 93.6)
Income								
≤ \$24,999	*	*	*	*	9,437	2,277,500	68.4	(61.5 - 75.3)
\$25,000-\$49,999	*	*	*	*	9,492	2,053,000	71.9	(64.5 - 79.3)
\$50,000-\$74,999	*	*	*	*	5,721	1,736,100	90.0	(87.7 - 92.3)
≥ \$75,000	114	19,400	93.9	(88.6 - 99.2)	8,460	4,019,300	92.2	(90.8 - 93.6)

¹ Defined as ever having blood cholesterol checked

² Sample size from survey

³ Estimated number of adults who have ever had blood cholesterol checked (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 42b. Ever had blood cholesterol checked among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	655	79.7	(74.7 - 84.8)	37,450	79.9	(77.3 - 82.5)
Gender						
Male	250	77.0	(68.5 - 85.6)	14,389	78.0	(73.9 - 82.1)
Female	405	82.3	(77.1 - 87.6)	23,061	81.6	(78.4 - 84.8)
Race/Ethnicity						
White Non-Hispanic	635	80.9	(76.2 - 85.5)	35,198	85.6	(84.0 - 87.2)
Black Non-Hispanic	*	*	*	720	77.5	(68.2 - 86.7)
Hispanic	*	*	*	842	57.2	(47.4 - 67.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	326	74.1	(65.8 - 82.3)	15,342	70.2	(65.2 - 75.1)
Some College	154	82.0	(73.5 - 90.5)	9,732	76.7	(71.1 - 82.3)
≥ College Degree	172	87.5	(80.2 - 94.8)	12,296	90.9	(88.6 - 93.2)
Income						
≤ \$24,999	*	*	*	9,437	68.1	(61.3 - 74.9)
\$25,000-\$49,999	*	*	*	9,492	71.6	(64.4 - 78.7)
\$50,000-\$74,999	*	*	*	5,721	89.6	(87.2 - 92.0)
≥ \$75,000	114	93.8	(88.4 - 99.2)	8,460	91.7	(90.1 - 93.2)

¹ Defined as ever having blood cholesterol checked

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 43a. Blood cholesterol checked within the past 5 years among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	651	68,100	75.5	(70.3 - 80.8)	37,201	11,035,200	77.8	(75.1 - 80.4)
Gender								
Male	249	32,600	74.0	(65.1 - 82.9)	14,307	5,020,700	74.5	(70.2 - 78.9)
Female	402	35,500	77.0	(71.4 - 82.7)	22,894	6,014,600	80.7	(77.6 - 83.8)
Age (years)								
18-34	*	*	*	*	3,647	2,179,800	53.3	(46.7 - 59.9)
35-44	78	13,700	84.5	(76.3 - 92.8)	4,862	2,161,200	78.0	(72.8 - 83.2)
45-54	155	15,100	82.6	(76.2 - 88.9)	7,697	2,408,600	86.6	(82.4 - 90.8)
55-64	128	12,700	92.9	(87.9 - 97.9)	8,677	1,957,900	93.3	(91.4 - 95.3)
≥ 65	206	13,200	93.6	(89.7 - 97.4)	12,318	2,327,800	94.8	(93.1 - 96.6)
Race/Ethnicity								
White Non-Hispanic	631	66,700	76.7	(71.7 - 81.6)	34,960	7,486,600	83.4	(81.8 - 85.0)
Black Non-Hispanic	*	*	*	*	719	1,561,700	76.5	(67.1 - 85.9)
Hispanic	*	*	*	*	838	1,184,000	55.6	(45.6 - 65.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	324	28,800	67.6	(59.1 - 76.2)	15,215	3,612,500	68.3	(63.2 - 73.3)
Some College	153	18,300	80.1	(71.3 - 88.9)	9,673	2,524,200	75.4	(69.8 - 81.1)
≥ College Degree	171	20,500	84.7	(77.1 - 92.4)	12,236	4,806,500	88.1	(85.5 - 90.6)
Income								
≤ \$24,999	*	*	*	*	9,350	2,214,800	66.9	(60.1 - 73.8)
\$25,000-\$49,999	*	*	*	*	9,454	1,936,600	68.8	(61.5 - 76.1)
\$50,000-\$74,999	*	*	*	*	5,701	1,640,400	85.9	(82.4 - 89.5)
≥ \$75,000	114	18,600	89.8	(83.4 - 96.3)	8,435	3,872,400	89.2	(87.5 - 90.9)

¹ Defined as having had blood cholesterol checked within the past 5 years

² Sample size from survey

³ Estimated number of adults who have had blood cholesterol checked within the past 5 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 43b. Blood cholesterol checked within the past 5 years among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	651	76.1	(70.9 - 81.2)	37,201	77.3	(74.7 - 79.9)
Gender						
Male	249	75.2	(66.6 - 83.8)	14,307	74.8	(70.7 - 79.0)
Female	402	76.9	(71.2 - 82.6)	22,894	79.5	(76.3 - 82.7)
Race/Ethnicity						
White Non-Hispanic	631	77.1	(72.3 - 82.0)	34,960	82.6	(81.0 - 84.3)
Black Non-Hispanic	*	*	*	719	76.2	(66.9 - 85.5)
Hispanic	*	*	*	838	55.9	(46.1 - 65.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	324	68.6	(60.3 - 77.0)	15,215	67.8	(62.9 - 72.8)
Some College	153	80.7	(72.1 - 89.3)	9,673	74.7	(69.1 - 80.2)
≥ College Degree	171	84.5	(76.8 - 92.2)	12,236	87.5	(85.0 - 90.1)
Income						
≤ \$24,999	*	*	*	9,350	66.6	(59.8 - 73.4)
\$25,000-\$49,999	*	*	*	9,454	68.5	(61.4 - 75.6)
\$50,000-\$74,999	*	*	*	5,701	85.4	(81.7 - 89.0)
≥ \$75,000	114	89.8	(83.3 - 96.3)	8,435	88.5	(86.7 - 90.3)

¹ Defined as having had blood cholesterol checked within the past 5 years

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 44a. Cardiovascular disease among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	665	7,500	8.0	(6.0 - 10.1)	37,810	1,135,000	7.7	(6.5 - 8.8)
Gender								
Male	253	4,300	9.3	(5.8 - 12.8)	14,477	664,600	9.4	(7.5 - 11.4)
Female	412	3,200	6.7	(4.5 - 8.9)	23,333	470,400	6.1	(4.8 - 7.4)
Age (years)								
18-34	*	*	*	*	3,930	100,800	2.3	(0.0 - 4.8)
35-44	*	*	*	*	4,971	103,400	3.6	(1.5 - 5.7)
45-54	158	1,200	6.5	(2.7 - 10.3)	7,768	131,300	4.6	(2.9 - 6.4)
55-64	128	1,500	11.0	(5.0 - 17.0)	8,699	224,200	10.6	(8.3 - 12.9)
≥ 65	207	3,800	27.3	(20.2 - 34.4)	12,442	575,300	23.0	(19.9 - 26.1)
Race/Ethnicity								
White Non-Hispanic	645	7,400	8.1	(6.0 - 10.2)	35,522	772,200	8.3	(7.4 - 9.2)
Black Non-Hispanic	*	*	*	*	728	140,700	6.6	(3.3 - 9.9)
Hispanic	*	*	*	*	854	163,300	7.3	(2.0 - 12.5)
Other Non-Hispanic	*	*	*	*	706	58,800	5.2	(0.9 - 9.5)
Education								
≤ High School	334	4,900	10.8	(7.2 - 14.3)	15,486	590,800	10.7	(8.2 - 13.2)
Some College	155	1,900	8.3	(4.3 - 12.3)	9,849	227,100	6.4	(4.7 - 8.1)
≥ College Degree	173	700	2.7	(0.8 - 4.7)	12,393	314,600	5.6	(4.3 - 6.9)
Income								
≤ \$24,999	176	2,900	14.6	(9.0 - 20.3)	9,505	448,100	12.9	(9.1 - 16.7)
\$25,000-\$49,999	175	2,300	9.8	(5.4 - 14.1)	9,571	179,300	6.2	(4.8 - 7.6)
\$50,000-\$74,999	*	*	*	*	5,751	166,900	8.5	(5.5 - 11.4)
≥ \$75,000	*	*	*	*	8,538	183,500	4.1	(2.9 - 5.3)

¹ Defined as having ever been told by a health professional that respondent had a heart attack or angina or a stroke

² Sample size from survey

³ Estimated number of adults with cardiovascular disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 44b. Cardiovascular disease among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	665	8.3	(6.1 - 10.5)	37,810	7.6	(6.4 - 8.7)
Gender						
Male	253	10.2	(6.3 - 14.2)	14,477	9.7	(7.8 - 11.7)
Female	412	6.5	(4.4 - 8.7)	23,333	5.6	(4.3 - 6.9)
Race/Ethnicity						
White Non-Hispanic	645	8.4	(6.2 - 10.7)	35,522	8.2	(7.2 - 9.1)
Black Non-Hispanic	*	*	*	728	6.8	(3.2 - 10.4)
Hispanic	*	*	*	854	7.0	(2.0 - 12.0)
Other Non-Hispanic	*	*	*	706	5.5	(0.9 - 10.0)
Education						
≤ High School	334	11.1	(7.3 - 14.9)	15,486	10.5	(8.0 - 13.0)
Some College	155	8.8	(4.5 - 13.2)	9,849	6.4	(4.5 - 8.2)
≥ College Degree	173	2.9	(0.7 - 5.0)	12,393	5.5	(4.2 - 6.9)
Income						
≤ \$24,999	176	14.8	(9.2 - 20.3)	9,505	12.8	(9.0 - 16.6)
\$25,000-\$49,999	175	10.8	(5.8 - 15.7)	9,571	6.0	(4.7 - 7.4)
\$50,000-\$74,999	*	*	*	5,751	8.5	(5.4 - 11.6)
≥ \$75,000	*	*	*	8,538	4.0	(2.7 - 5.2)

¹ Defined as having ever been told by a health professional that respondent had a heart attack or angina or a stroke

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 45a. Coronary heart disease among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	666	6,600	7.1	(5.1 - 9.1)	37,838	929,400	6.3	(5.3 - 7.3)
Gender								
Male	253	4,200	9.2	(5.7 - 12.7)	14,479	598,600	8.5	(6.6 - 10.4)
Female	413	2,400	5.1	(3.1 - 7.0)	23,359	330,800	4.3	(3.4 - 5.2)
Age (years)								
18-34	*	*	*	*	3,932	63,400	1.4	(0.0 - 3.6)
35-44	*	*	*	*	4,973	95,300	3.3	(1.2 - 5.4)
45-54	158	1,200	6.5	(2.7 - 10.3)	7,778	108,300	3.8	(2.1 - 5.6)
55-64	128	1,400	10.6	(4.7 - 16.6)	8,703	188,600	8.9	(6.7 - 11.1)
≥ 65	208	3,200	22.4	(15.5 - 29.2)	12,452	473,800	18.9	(16.1 - 21.7)
Race/Ethnicity								
White Non-Hispanic	646	6,500	7.2	(5.2 - 9.2)	35,552	636,700	6.9	(6.0 - 7.7)
Black Non-Hispanic	*	*	*	*	728	113,100	5.3	(2.3 - 8.4)
Hispanic	*	*	*	*	853	128,000	5.7	(1.1 - 10.3)
Other Non-Hispanic	*	*	*	*	705	51,500	4.6	(0.5 - 8.7)
Education								
≤ High School	335	4,100	9.1	(5.8 - 12.5)	15,485	474,500	8.6	(6.3 - 10.8)
Some College	155	1,900	8.0	(4.1 - 12.0)	9,860	193,000	5.5	(3.9 - 7.1)
≥ College Degree	*	*	*	*	12,411	259,700	4.6	(3.4 - 5.8)
Income								
≤ \$24,999	176	2,500	12.6	(7.3 - 17.8)	9,507	372,300	10.7	(7.3 - 14.1)
\$25,000-\$49,999	176	2,100	8.8	(4.5 - 13.0)	9,585	136,300	4.7	(3.8 - 5.7)
\$50,000-\$74,999	*	*	*	*	5,755	154,600	7.8	(4.9 - 10.7)
≥ \$75,000	*	*	*	*	8,541	151,800	3.4	(2.3 - 4.5)

¹ Defined as having ever been told by a health professional that respondent had a heart attack or angina

² Sample size from survey

³ Estimated number of adults with coronary heart disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 45b. Coronary heart disease among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	666	7.4	(5.3 - 9.6)	37,838	6.3	(5.2 - 7.3)
Gender						
Male	253	10.1	(6.1 - 14.0)	14,479	8.8	(6.9 - 10.8)
Female	413	4.9	(3.0 - 6.8)	23,359	3.9	(3.0 - 4.8)
Race/Ethnicity						
White Non-Hispanic	646	7.5	(5.3 - 9.7)	35,552	6.8	(5.9 - 7.7)
Black Non-Hispanic	*	*	*	728	5.6	(2.2 - 9.0)
Hispanic	*	*	*	853	5.4	(1.1 - 9.6)
Other Non-Hispanic	*	*	*	705	4.8	(0.4 - 9.2)
Education						
≤ High School	335	9.4	(5.8 - 13.1)	15,485	8.5	(6.2 - 10.7)
Some College	155	8.5	(4.2 - 12.9)	9,860	5.5	(3.7 - 7.2)
≥ College Degree	*	*	*	12,411	4.7	(3.4 - 5.9)
Income						
≤ \$24,999	176	12.6	(7.5 - 17.7)	9,507	10.7	(7.2 - 14.1)
\$25,000-\$49,999	176	9.8	(5.0 - 14.7)	9,585	4.7	(3.6 - 5.7)
\$50,000-\$74,999	*	*	*	5,755	8.0	(4.8 - 11.1)
≥ \$75,000	*	*	*	8,541	3.3	(2.1 - 4.5)

¹ Defined as having ever been told by a health professional that respondent had a heart attack or angina

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 46a. Ever diagnosed with diabetes among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	669	9,300	9.9	(7.6 - 12.3)	38,174	1,382,700	9.3	(8.0 - 10.5)
Gender								
Male	254	4,700	10.3	(6.4 - 14.1)	14,626	530,400	7.5	(6.2 - 8.8)
Female	415	4,600	9.6	(6.9 - 12.4)	23,548	852,300	11.0	(8.9 - 13.0)
Age (years)								
18-34	*	*	*	*	3,941	150,200	3.4	(1.0 - 5.8)
35-44	*	*	*	*	4,987	165,100	5.7	(3.6 - 7.9)
45-54	159	2,300	12.5	(7.0 - 17.9)	7,830	279,000	9.8	(6.7 - 12.8)
55-64	129	2,800	20.2	(12.8 - 27.7)	8,771	332,900	15.5	(12.3 - 18.8)
≥ 65	209	3,100	21.7	(15.3 - 28.0)	12,645	455,600	17.9	(15.0 - 20.8)
Race/Ethnicity								
White Non-Hispanic	649	9,200	10.2	(7.8 - 12.6)	35,862	809,300	8.7	(7.8 - 9.5)
Black Non-Hispanic	*	*	*	*	739	261,000	12.1	(6.6 - 17.6)
Hispanic	*	*	*	*	862	227,300	10.1	(5.4 - 14.8)
Other Non-Hispanic	*	*	*	*	711	85,100	7.6	(2.3 - 12.8)
Education								
≤ High School	335	5,000	11.1	(7.3 - 14.9)	15,709	730,000	13.0	(10.5 - 15.6)
Some College	156	2,100	8.9	(4.8 - 13.0)	9,918	303,500	8.6	(5.9 - 11.2)
≥ College Degree	175	2,200	8.7	(4.8 - 12.6)	12,464	337,500	6.0	(4.8 - 7.2)
Income								
≤ \$24,999	177	2,800	14.1	(8.6 - 19.7)	9,652	543,400	15.5	(11.4 - 19.5)
\$25,000-\$49,999	177	2,600	11.0	(6.4 - 15.7)	9,667	291,000	10.0	(7.5 - 12.6)
\$50,000-\$74,999	100	1,200	7.6	(1.7 - 13.6)	5,777	151,000	7.6	(5.0 - 10.3)
≥ \$75,000	116	1,600	7.6	(3.3 - 11.9)	8,565	220,800	4.9	(3.8 - 6.1)

¹ Defined as having ever been told by a doctor that respondent had diabetes, excluding pre-diabetes and women with diabetes only when pregnant

² Sample size from survey

³ Estimated number of adults who have diabetes (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 46b. Ever diagnosed with diabetes among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	669	9.9	(7.4 - 12.4)	38,174	9.0	(7.8 - 10.3)
Gender						
Male	254	10.8	(6.5 - 15.0)	14,626	7.5	(6.1 - 8.9)
Female	415	9.0	(6.4 - 11.6)	23,548	10.4	(8.4 - 12.4)
Race/Ethnicity						
White Non-Hispanic	649	10.1	(7.5 - 12.6)	35,862	8.4	(7.5 - 9.2)
Black Non-Hispanic	*	*	*	739	11.4	(6.0 - 16.7)
Hispanic	*	*	*	862	10.1	(5.4 - 14.9)
Other Non-Hispanic	*	*	*	711	7.7	(2.3 - 13.0)
Education						
≤ High School	335	11.5	(7.3 - 15.8)	15,709	12.6	(10.0 - 15.2)
Some College	156	8.5	(4.6 - 12.5)	9,918	8.3	(5.6 - 10.9)
≥ College Degree	175	8.0	(4.4 - 11.5)	12,464	5.9	(4.7 - 7.2)
Income						
≤ \$24,999	177	13.5	(8.2 - 18.8)	9,652	15.2	(11.1 - 19.2)
\$25,000-\$49,999	177	11.1	(6.3 - 15.9)	9,667	9.7	(7.1 - 12.2)
\$50,000-\$74,999	100	8.1	(1.2 - 15.0)	5,777	7.6	(4.9 - 10.2)
≥ \$75,000	116	6.9	(2.9 - 10.8)	8,565	4.7	(3.6 - 5.9)

¹ Defined as having ever been told by a doctor that respondent had diabetes, excluding pre-diabetes and women with diabetes only when pregnant

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 47a. Ever diagnosed with asthma (lifetime) among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	669	14,500	15.5	(11.7 - 19.3)	38,096	2,225,100	15.0	(13.2 - 16.8)
Gender								
Male	255	6,400	13.8	(7.7 - 19.9)	14,601	875,800	12.4	(10.1 - 14.7)
Female	414	8,200	17.1	(12.5 - 21.6)	23,495	1,349,300	17.4	(14.6 - 20.1)
Age (years)								
18-34	92	5,500	18.0	(8.6 - 27.3)	3,931	825,100	18.5	(13.8 - 23.3)
35-44	80	2,600	15.4	(7.4 - 23.4)	4,972	431,700	15.1	(11.2 - 19.1)
45-54	159	3,500	18.6	(11.7 - 25.4)	7,819	365,900	12.8	(9.7 - 16.0)
55-64	128	1,900	14.1	(7.8 - 20.4)	8,756	316,700	14.8	(11.9 - 17.7)
≥ 65	210	1,100	7.6	(4.0 - 11.1)	12,618	285,700	11.2	(8.6 - 13.8)
Race/Ethnicity								
White Non-Hispanic	649	14,200	15.6	(11.7 - 19.5)	35,783	1,365,700	14.6	(13.0 - 16.3)
Black Non-Hispanic	*	*	*	*	737	367,000	17.2	(10.2 - 24.2)
Hispanic	*	*	*	*	863	317,100	14.1	(9.0 - 19.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	335	5,300	11.6	(6.9 - 16.3)	15,675	831,600	14.9	(12.0 - 17.8)
Some College	156	4,500	19.0	(11.0 - 27.0)	9,893	587,500	16.6	(13.1 - 20.1)
≥ College Degree	175	4,800	19.5	(11.2 - 27.8)	12,444	739,000	13.2	(10.4 - 15.9)
Income								
≤ \$24,999	177	3,000	15.2	(8.8 - 21.6)	9,643	708,900	20.2	(14.9 - 25.5)
\$25,000-\$49,999	177	3,500	14.7	(7.0 - 22.4)	9,641	389,800	13.4	(10.0 - 16.9)
\$50,000-\$74,999	100	2,400	14.9	(5.4 - 24.4)	5,762	257,800	13.2	(10.0 - 16.5)
≥ \$75,000	116	4,500	21.0	(11.2 - 30.7)	8,562	577,900	12.8	(10.4 - 15.3)

¹ Defined as adults having ever been told by a health professional that they had asthma

² Sample size from survey

³ Estimated number of adults who have lifetime asthma (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 47b. Ever diagnosed with asthma (lifetime) among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	669	15.2	(11.5 - 19.0)	38,096	15.2	(13.3 - 17.0)
Gender						
Male	255	13.3	(7.4 - 19.2)	14,601	12.6	(10.3 - 14.9)
Female	414	17.1	(12.4 - 21.7)	23,495	17.5	(14.8 - 20.3)
Race/Ethnicity						
White Non-Hispanic	649	15.3	(11.4 - 19.1)	35,783	14.9	(13.2 - 16.6)
Black Non-Hispanic	*	*	*	737	17.1	(10.1 - 24.1)
Hispanic	*	*	*	863	14.5	(9.3 - 19.7)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	335	11.1	(6.6 - 15.6)	15,675	15.1	(12.1 - 18.0)
Some College	156	18.9	(10.9 - 26.9)	9,893	16.9	(13.4 - 20.4)
≥ College Degree	175	19.7	(11.3 - 28.0)	12,444	13.3	(10.6 - 16.0)
Income						
≤ \$24,999	177	15.0	(8.5 - 21.6)	9,643	20.3	(15.0 - 25.6)
\$25,000-\$49,999	177	14.6	(7.0 - 22.1)	9,641	13.7	(10.2 - 17.2)
\$50,000-\$74,999	100	14.6	(5.3 - 23.9)	5,762	13.5	(10.0 - 17.0)
≥ \$75,000	116	20.3	(10.7 - 29.9)	8,562	13.0	(10.6 - 15.4)

¹ Defined as adults having ever been told by a health professional that they had asthma

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 48a. Current asthma among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	667	11,500	12.2	(8.6 - 15.8)	37,933	1,415,000	9.6	(8.1 - 11.0)
Gender								
Male	254	5,200	11.2	(5.3 - 17.2)	14,546	440,700	6.3	(4.8 - 7.7)
Female	413	6,300	13.2	(9.0 - 17.4)	23,387	974,300	12.6	(10.2 - 15.0)
Age (years)								
18-34	92	4,300	13.9	(4.9 - 22.9)	3,909	467,200	10.5	(6.8 - 14.3)
35-44	80	2,300	13.7	(6.0 - 21.4)	4,947	246,800	8.7	(5.8 - 11.6)
45-54	157	2,900	15.7	(9.1 - 22.3)	7,787	240,000	8.4	(6.1 - 10.8)
55-64	128	1,300	9.2	(4.2 - 14.1)	8,726	231,100	10.8	(8.1 - 13.5)
≥ 65	210	800	5.3	(2.2 - 8.4)	12,564	229,900	9.1	(6.5 - 11.6)
Race/Ethnicity								
White Non-Hispanic	647	11,300	12.4	(8.7 - 16.1)	35,629	888,400	9.6	(8.2 - 11.0)
Black Non-Hispanic	*	*	*	*	736	246,900	11.6	(5.7 - 17.5)
Hispanic	*	*	*	*	859	186,800	8.3	(4.4 - 12.2)
Other Non-Hispanic	*	*	*	*	709	93,000	8.3	(1.4 - 15.2)
Education								
≤ High School	334	3,900	8.6	(4.2 - 13.0)	15,594	595,100	10.7	(8.2 - 13.2)
Some College	156	3,700	16.0	(8.3 - 23.7)	9,861	322,000	9.1	(7.0 - 11.3)
≥ College Degree	174	3,800	15.4	(7.3 - 23.5)	12,394	430,900	7.7	(5.8 - 9.7)
Income								
≤ \$24,999	177	1,900	9.7	(4.7 - 14.8)	9,600	489,200	14.0	(9.7 - 18.3)
\$25,000-\$49,999	177	2,700	11.3	(4.0 - 18.6)	9,590	268,400	9.3	(6.2 - 12.4)
\$50,000-\$74,999	98	2,100	13.3	(3.9 - 22.8)	5,741	143,400	7.4	(5.1 - 9.6)
≥ \$75,000	116	4,000	19.0	(9.3 - 28.6)	8,541	339,200	7.6	(5.8 - 9.4)

¹ Defined as adults having ever been told by a health professional that they had asthma and still have asthma

² Sample size from survey

³ Estimated number of adults who currently have asthma (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 48b. Current asthma among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	667	12.1	(8.6 - 15.7)	37,933	9.7	(8.2 - 11.2)
Gender						
Male	254	11.0	(5.3 - 16.7)	14,546	6.4	(4.9 - 7.9)
Female	413	13.2	(8.9 - 17.5)	23,387	12.7	(10.2 - 15.1)
Race/Ethnicity						
White Non-Hispanic	647	12.3	(8.6 - 15.9)	35,629	9.7	(8.3 - 11.2)
Black Non-Hispanic	*	*	*	736	11.5	(5.6 - 17.3)
Hispanic	*	*	*	859	8.6	(4.5 - 12.6)
Other Non-Hispanic	*	*	*	709	8.3	(1.3 - 15.3)
Education						
≤ High School	334	8.4	(4.2 - 12.6)	15,594	10.8	(8.3 - 13.3)
Some College	156	16.0	(8.2 - 23.8)	9,861	9.4	(7.2 - 11.6)
≥ College Degree	174	15.6	(7.5 - 23.7)	12,394	7.8	(5.8 - 9.7)
Income						
≤ \$24,999	177	9.6	(4.4 - 14.9)	9,600	14.1	(9.7 - 18.4)
\$25,000-\$49,999	177	11.5	(4.3 - 18.7)	9,590	9.5	(6.3 - 12.7)
\$50,000-\$74,999	98	13.1	(3.8 - 22.4)	5,741	7.5	(5.0 - 10.1)
≥ \$75,000	116	18.6	(9.1 - 28.2)	8,541	7.7	(5.9 - 9.5)

¹ Defined as adults having ever been told by a health professional that they had asthma and still have asthma

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 49a. Chronic joint symptoms among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	668	46,800	50.2	(45.1 - 55.3)	38,099	6,377,100	42.9	(40.5 - 45.3)
Gender								
Male	253	24,800	54.7	(46.4 - 63.0)	14,595	3,033,600	42.8	(39.0 - 46.6)
Female	415	22,000	45.9	(40.1 - 51.8)	23,504	3,343,500	43.0	(40.0 - 46.0)
Age (years)								
18-34	*	*	*	*	3,933	1,261,300	28.3	(22.6 - 34.0)
35-44	*	*	*	*	4,984	1,110,900	38.6	(33.1 - 44.0)
45-54	158	9,200	49.6	(41.1 - 58.2)	7,817	1,456,500	51.2	(46.4 - 55.9)
55-64	129	9,100	66.3	(57.5 - 75.1)	8,751	1,167,400	54.6	(50.4 - 58.8)
≥ 65	210	8,500	59.6	(52.0 - 67.1)	12,614	1,381,000	54.3	(50.5 - 58.1)
Race/Ethnicity								
White Non-Hispanic	648	44,600	49.5	(44.5 - 54.5)	35,785	4,267,600	45.8	(43.8 - 47.7)
Black Non-Hispanic	*	*	*	*	740	890,500	41.3	(32.5 - 50.0)
Hispanic	*	*	*	*	860	827,500	36.7	(27.7 - 45.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	335	24,700	54.6	(46.9 - 62.4)	15,676	2,450,000	43.8	(39.4 - 48.1)
Some College	155	11,900	52.1	(42.4 - 61.7)	9,900	1,598,300	45.1	(40.1 - 50.1)
≥ College Degree	175	10,200	40.9	(32.1 - 49.8)	12,440	2,313,100	41.1	(37.8 - 44.5)
Income								
≤ \$24,999	*	*	*	*	9,641	1,483,000	42.2	(36.3 - 48.1)
\$25,000-\$49,999	*	*	*	*	9,647	1,332,100	45.9	(39.8 - 52.1)
\$50,000-\$74,999	*	*	*	*	5,768	894,200	45.3	(39.5 - 51.1)
≥ \$75,000	116	7,300	34.5	(24.8 - 44.2)	8,549	1,874,200	41.7	(38.4 - 45.1)

¹ Defined as adults having symptoms of pain, aching or stiffness in or around a joint during the past 30 days and began more than 3 months ago

² Sample size from survey

³ Estimated number of adults who have chronic joint symptoms (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 49b. Chronic joint symptoms among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	668	49.8	(44.7 - 54.9)	38,099	42.1	(39.7 - 44.5)
Gender						
Male	253	54.6	(46.3 - 62.9)	14,595	42.6	(38.8 - 46.3)
Female	415	45.2	(39.2 - 51.2)	23,504	41.6	(38.6 - 44.7)
Race/Ethnicity						
White Non-Hispanic	648	49.1	(44.1 - 54.2)	35,785	44.7	(42.7 - 46.6)
Black Non-Hispanic	*	*	*	740	40.2	(31.5 - 48.9)
Hispanic	*	*	*	860	36.6	(27.8 - 45.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	335	54.5	(46.7 - 62.2)	15,676	43.1	(38.8 - 47.5)
Some College	155	51.4	(41.7 - 61.1)	9,900	43.9	(39.0 - 48.8)
≥ College Degree	175	40.1	(31.2 - 49.0)	12,440	40.3	(37.0 - 43.7)
Income						
≤ \$24,999	*	*	*	9,641	41.6	(35.7 - 47.4)
\$25,000-\$49,999	*	*	*	9,647	45.0	(39.0 - 51.0)
\$50,000-\$74,999	*	*	*	5,768	44.3	(38.4 - 50.2)
≥ \$75,000	116	33.3	(23.5 - 43.1)	8,549	40.8	(37.4 - 44.2)

¹ Defined as adults having symptoms of pain, aching or stiffness in or around a joint during the past 30 days and began more than 3 months ago

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 50a. Arthritis among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	661	32,300	34.7	(30.2 - 39.3)	38,013	4,102,800	27.7	(25.9 - 29.6)
Gender								
Male	252	15,400	33.8	(26.6 - 41.1)	14,555	1,642,700	23.3	(20.7 - 25.8)
Female	409	16,900	35.7	(30.2 - 41.1)	23,458	2,460,100	31.8	(29.2 - 34.4)
Age (years)								
18-34	90	4,100	13.6	(6.2 - 21.1)	3,927	312,100	7.0	(4.9 - 9.1)
35-44	*	*	*	*	4,972	412,500	14.5	(11.3 - 17.6)
45-54	158	6,000	32.4	(24.5 - 40.3)	7,789	869,100	30.7	(26.6 - 34.8)
55-64	126	8,000	60.0	(50.7 - 69.3)	8,729	999,300	47.0	(42.9 - 51.1)
≥ 65	208	9,000	63.3	(56.0 - 70.7)	12,596	1,509,800	59.6	(55.8 - 63.3)
Race/Ethnicity								
White Non-Hispanic	641	31,800	35.5	(30.9 - 40.0)	35,710	2,967,700	31.9	(30.3 - 33.6)
Black Non-Hispanic	*	*	*	*	737	588,100	27.3	(20.2 - 34.4)
Hispanic	*	*	*	*	856	380,700	17.1	(11.8 - 22.5)
Other Non-Hispanic	*	*	*	*	710	166,300	14.8	(7.8 - 21.8)
Education								
≤ High School	333	17,400	38.6	(31.5 - 45.7)	15,652	1,742,700	31.3	(27.8 - 34.8)
Some College	153	7,100	30.9	(22.6 - 39.3)	9,874	902,000	25.5	(22.3 - 28.6)
≥ College Degree	172	7,700	31.5	(23.4 - 39.7)	12,407	1,437,800	25.6	(22.9 - 28.4)
Income								
≤ \$24,999	*	*	*	*	9,618	1,112,500	32.0	(27.1 - 36.9)
\$25,000-\$49,999	174	9,700	41.2	(31.6 - 50.7)	9,624	821,900	28.4	(24.2 - 32.6)
\$50,000-\$74,999	*	*	*	*	5,753	542,500	27.5	(23.2 - 31.9)
≥ \$75,000	115	4,600	22.0	(13.9 - 30.0)	8,536	1,012,800	22.6	(19.8 - 25.3)

¹ Defined as adults having ever been told by a health professional that they have some form of arthritis, rheumatoid arthritis, gout, lupus or fibromyalgia (excluding back or neck)

² Sample size from survey

³ Estimated number of adults who have ever been told by a health professional they have some form of arthritis (excluding back or neck) (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 50b. Arthritis among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	661	34.7	(30.1 - 39.3)	38,013	26.6	(24.8 - 28.5)
Gender						
Male	252	34.6	(27.2 - 42.0)	14,555	23.5	(20.9 - 26.1)
Female	409	34.8	(29.2 - 40.3)	23,458	29.5	(27.0 - 32.1)
Race/Ethnicity						
White Non-Hispanic	641	35.3	(30.7 - 40.0)	35,710	30.4	(28.7 - 32.0)
Black Non-Hispanic	*	*	*	737	26.5	(19.3 - 33.6)
Hispanic	*	*	*	856	17.2	(11.8 - 22.6)
Other Non-Hispanic	*	*	*	710	14.9	(7.7 - 22.0)
Education						
≤ High School	333	38.6	(31.5 - 45.8)	15,652	30.4	(26.9 - 33.8)
Some College	153	31.1	(22.6 - 39.5)	9,874	24.1	(21.0 - 27.1)
≥ College Degree	172	31.0	(22.8 - 39.3)	12,407	24.7	(21.9 - 27.5)
Income						
≤ \$24,999	*	*	*	9,618	31.2	(26.3 - 36.1)
\$25,000-\$49,999	174	41.9	(32.3 - 51.4)	9,624	27.3	(23.2 - 31.3)
\$50,000-\$74,999	*	*	*	5,753	26.3	(21.9 - 30.7)
≥ \$75,000	115	19.7	(12.1 - 27.4)	8,536	21.6	(18.8 - 24.3)

¹ Defined as adults having ever been told by a health professional that they have some form of arthritis, rheumatoid arthritis, gout, lupus or fibromyalgia (excluding back or neck)

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 51a. Activity limitations because of physical, mental or emotional problems among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	664	23,600	25.4	(21.6 - 29.3)	37,998	3,053,000	20.6	(18.7 - 22.5)
Gender								
Male	251	12,200	27.2	(20.9 - 33.4)	14,566	1,318,400	18.6	(15.7 - 21.5)
Female	413	11,300	23.8	(19.1 - 28.5)	23,432	1,734,600	22.4	(19.8 - 25.0)
Age (years)								
18-34	90	2,800	9.4	(4.0 - 14.7)	3,929	580,000	13.0	(8.2 - 17.9)
35-44	80	4,200	25.2	(15.2 - 35.1)	4,973	418,600	14.5	(11.0 - 18.1)
45-54	157	4,700	25.5	(18.1 - 32.9)	7,812	620,200	21.7	(18.3 - 25.2)
55-64	128	6,500	47.2	(37.6 - 56.9)	8,739	651,600	30.5	(26.5 - 34.6)
≥ 65	209	5,400	38.3	(30.7 - 45.8)	12,545	782,700	31.1	(27.7 - 34.6)
Race/Ethnicity								
White Non-Hispanic	644	23,200	25.9	(22.0 - 29.7)	35,688	2,037,200	21.9	(20.3 - 23.5)
Black Non-Hispanic	*	*	*	*	735	429,700	20.1	(13.5 - 26.7)
Hispanic	*	*	*	*	863	448,200	19.8	(11.9 - 27.8)
Other Non-Hispanic	*	*	*	*	712	137,900	12.3	(5.2 - 19.3)
Education								
≤ High School	334	11,600	25.7	(20.1 - 31.3)	15,626	1,359,600	24.4	(20.6 - 28.1)
Some College	154	5,500	24.3	(16.8 - 31.8)	9,875	700,800	19.8	(16.5 - 23.2)
≥ College Degree	173	6,200	25.3	(17.9 - 32.8)	12,415	924,800	16.5	(14.2 - 18.7)
Income								
≤ \$24,999	*	*	*	*	9,603	1,055,800	30.1	(24.7 - 35.5)
\$25,000-\$49,999	176	6,900	28.8	(20.7 - 37.0)	9,628	630,900	21.8	(16.6 - 27.0)
\$50,000-\$74,999	100	2,600	16.2	(8.4 - 23.9)	5,760	408,100	20.7	(16.1 - 25.2)
≥ \$75,000	115	2,800	13.4	(6.9 - 19.9)	8,544	583,300	13.0	(11.1 - 14.9)

¹ Defined as activity limitations because of physical, mental or emotional problems

² Sample size from survey

³ Estimated number of adults who are limited in any activities because of physical, mental or emotional problems (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 51b. Activity limitations because of physical, mental or emotional problems among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	664	25.5	(21.6 - 29.4)	37,998	20.1	(18.2 - 22.0)
Gender						
Male	251	27.3	(21.0 - 33.6)	14,566	18.6	(15.8 - 21.4)
Female	413	23.8	(18.9 - 28.6)	23,432	21.5	(18.9 - 24.1)
Race/Ethnicity						
White Non-Hispanic	644	25.8	(21.8 - 29.8)	35,688	21.3	(19.7 - 22.9)
Black Non-Hispanic	*	*	*	735	19.9	(13.2 - 26.5)
Hispanic	*	*	*	863	19.1	(11.5 - 26.8)
Other Non-Hispanic	*	*	*	712	12.4	(5.3 - 19.6)
Education						
≤ High School	334	25.6	(20.0 - 31.3)	15,626	23.8	(20.1 - 27.4)
Some College	154	24.6	(17.0 - 32.2)	9,875	19.3	(16.0 - 22.6)
≥ College Degree	173	25.2	(17.6 - 32.8)	12,415	16.1	(13.9 - 18.3)
Income						
≤ \$24,999	*	*	*	9,603	29.7	(24.3 - 35.2)
\$25,000-\$49,999	176	29.9	(21.6 - 38.3)	9,628	21.2	(16.1 - 26.3)
\$50,000-\$74,999	100	16.2	(8.2 - 24.1)	5,760	20.1	(15.6 - 24.6)
≥ \$75,000	115	12.1	(6.0 - 18.2)	8,544	12.7	(10.8 - 14.6)

¹ Defined as activity limitations because of physical, mental or emotional problems

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 52a. Have health problem that requires the use of special equipment among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	669	8,600	9.1	(6.8 - 11.5)	38,171	1,169,400	7.9	(6.9 - 8.9)
Gender								
Male	255	4,500	9.8	(5.9 - 13.7)	14,617	460,700	6.5	(5.3 - 7.7)
Female	414	4,100	8.5	(5.7 - 11.4)	23,554	708,700	9.1	(7.6 - 10.7)
Age (years)								
18-34	*	*	*	*	3,940	68,800	1.5	(0.7 - 2.4)
35-44	*	*	*	*	4,989	122,500	4.3	(2.4 - 6.1)
45-54	159	1,600	8.6	(3.5 - 13.7)	7,836	182,500	6.4	(4.7 - 8.1)
55-64	129	1,700	12.2	(6.0 - 18.4)	8,773	219,600	10.3	(7.8 - 12.8)
≥ 65	209	2,600	17.9	(12.6 - 23.3)	12,633	576,100	22.7	(19.0 - 26.3)
Race/Ethnicity								
White Non-Hispanic	649	8,400	9.3	(6.8 - 11.7)	35,853	802,000	8.6	(7.6 - 9.6)
Black Non-Hispanic	*	*	*	*	739	235,800	10.9	(6.4 - 15.5)
Hispanic	*	*	*	*	864	104,100	4.6	(2.1 - 7.1)
Other Non-Hispanic	*	*	*	*	715	27,600	2.5	(0.8 - 4.1)
Education								
≤ High School	335	4,500	9.9	(6.4 - 13.4)	15,711	604,200	10.8	(8.7 - 12.8)
Some College	156	2,000	8.4	(4.1 - 12.8)	9,914	260,000	7.3	(5.5 - 9.1)
≥ College Degree	175	2,100	8.6	(3.6 - 13.5)	12,463	298,900	5.3	(4.1 - 6.5)
Income								
≤ \$24,999	177	3,200	16.4	(10.1 - 22.6)	9,656	430,500	12.2	(9.6 - 14.8)
\$25,000-\$49,999	177	2,300	9.8	(5.4 - 14.2)	9,663	238,900	8.2	(5.9 - 10.5)
\$50,000-\$74,999	*	*	*	*	5,775	103,000	5.2	(3.6 - 6.8)
≥ \$75,000	*	*	*	*	8,566	136,000	3.0	(2.1 - 4.0)

¹ Defined as having any health problem that requires the use of special equipment such as a cane, a wheelchair, a special bed, or a special telephone

² Sample size from survey

³ Estimated number of adults who have any health problem that requires the use of special equipment (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 52b. Have health problem that requires the use of special equipment among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	669	9.4	(6.9 - 11.9)	38,171	7.6	(6.6 - 8.6)
Gender						
Male	255	10.3	(6.2 - 14.5)	14,617	6.8	(5.5 - 8.0)
Female	414	8.4	(5.5 - 11.3)	23,554	8.4	(6.9 - 9.9)
Race/Ethnicity						
White Non-Hispanic	649	9.4	(6.9 - 12.0)	35,853	8.3	(7.3 - 9.3)
Black Non-Hispanic	*	*	*	739	10.5	(6.1 - 15.0)
Hispanic	*	*	*	864	4.5	(2.1 - 6.9)
Other Non-Hispanic	*	*	*	715	2.2	(0.7 - 3.8)
Education						
≤ High School	335	10.1	(6.4 - 13.8)	15,711	10.3	(8.3 - 12.3)
Some College	156	8.4	(4.0 - 12.7)	9,914	7.1	(5.2 - 8.9)
≥ College Degree	175	9.0	(3.8 - 14.2)	12,463	5.3	(4.1 - 6.6)
Income						
≤ \$24,999	177	16.9	(10.4 - 23.4)	9,656	11.9	(9.4 - 14.5)
\$25,000-\$49,999	177	10.2	(5.4 - 14.9)	9,663	7.7	(5.5 - 10.0)
\$50,000-\$74,999	*	*	*	5,775	5.1	(3.4 - 6.8)
≥ \$75,000	*	*	*	8,566	2.9	(2.0 - 3.9)

¹ Defined as having any health problem that requires the use of special equipment such as a cane, a wheelchair, a special bed, or a special telephone

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 53a. Disability among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	664	25,600	27.5	(23.5 - 31.6)	38,016	3,394,200	22.9	(20.9 - 24.9)
Gender								
Male	251	13,400	29.7	(23.1 - 36.3)	14,568	1,469,700	20.7	(17.8 - 23.7)
Female	413	12,200	25.5	(20.7 - 30.3)	23,448	1,924,600	24.8	(22.1 - 27.5)
Age (years)								
18-34	91	3,400	11.5	(5.4 - 17.5)	3,931	590,300	13.3	(8.4 - 18.1)
35-44	*	*	*	*	4,974	447,900	15.6	(11.9 - 19.3)
45-54	157	4,800	25.9	(18.5 - 33.3)	7,811	643,900	22.6	(19.1 - 26.0)
55-64	128	6,800	49.5	(40.0 - 59.1)	8,740	702,100	33.0	(28.9 - 37.0)
≥ 65	208	6,000	42.3	(34.7 - 50.0)	12,560	1,010,000	39.9	(36.1 - 43.7)
Race/Ethnicity								
White Non-Hispanic	644	25,100	28.0	(23.9 - 32.0)	35,703	2,253,400	24.2	(22.6 - 25.9)
Black Non-Hispanic	*	*	*	*	736	511,700	23.8	(16.8 - 30.8)
Hispanic	*	*	*	*	864	484,300	21.4	(13.4 - 29.5)
Other Non-Hispanic	*	*	*	*	713	144,800	12.9	(5.8 - 19.9)
Education								
≤ High School	333	12,800	28.2	(22.3 - 34.2)	15,638	1,544,200	27.6	(23.7 - 31.4)
Some College	155	5,900	25.7	(18.0 - 33.3)	9,876	755,200	21.4	(17.9 - 24.8)
≥ College Degree	173	6,700	27.4	(19.6 - 35.2)	12,419	1,021,300	18.2	(15.9 - 20.5)
Income								
≤ \$24,999	*	*	*	*	9,613	1,179,100	33.6	(28.0 - 39.2)
\$25,000-\$49,999	176	7,100	30.0	(21.7 - 38.2)	9,629	699,000	24.2	(18.9 - 29.4)
\$50,000-\$74,999	100	3,000	18.7	(9.9 - 27.6)	5,757	430,100	21.8	(17.2 - 26.4)
≥ \$75,000	115	3,300	15.6	(8.4 - 22.9)	8,546	632,700	14.1	(12.1 - 16.1)

¹ Defined as having activity limitations due to physical, mental or emotional problems or have health problems that require the use of special equipment

² Sample size from survey

³ Estimated number of adults with a disability (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 53b. Disability among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	664	27.6	(23.5 - 31.7)	38,016	22.3	(20.4 - 24.3)
Gender						
Male	251	29.9	(23.2 - 36.6)	14,568	20.9	(18.0 - 23.8)
Female	413	25.4	(20.5 - 30.3)	23,448	23.6	(21.0 - 26.3)
Race/Ethnicity						
White Non-Hispanic	644	27.9	(23.8 - 32.1)	35,703	23.6	(21.9 - 25.3)
Black Non-Hispanic	*	*	*	736	23.5	(16.5 - 30.5)
Hispanic	*	*	*	864	20.6	(12.9 - 28.3)
Other Non-Hispanic	*	*	*	713	12.9	(5.8 - 20.1)
Education						
≤ High School	333	28.2	(22.2 - 34.3)	15,638	26.8	(23.0 - 30.6)
Some College	155	26.0	(18.3 - 33.7)	9,876	20.9	(17.4 - 24.3)
≥ College Degree	173	27.2	(19.3 - 35.1)	12,419	17.9	(15.5 - 20.3)
Income						
≤ \$24,999	*	*	*	9,613	33.2	(27.6 - 38.8)
\$25,000-\$49,999	176	30.9	(22.5 - 39.4)	9,629	23.5	(18.3 - 28.6)
\$50,000-\$74,999	100	19.3	(9.9 - 28.8)	5,757	21.2	(16.6 - 25.8)
≥ \$75,000	115	14.2	(7.3 - 21.1)	8,546	13.8	(11.7 - 15.8)

¹ Defined as having activity limitations due to physical, mental or emotional problems or have health problems that require the use of special equipment

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 54a. Current smoking among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	669	23,300	24.8	(20.1 - 29.5)	38,029	2,483,500	16.8	(15.1 - 18.6)
Gender								
Male	255	11,400	24.7	(17.0 - 32.4)	14,559	1,356,400	19.3	(16.3 - 22.2)
Female	414	11,900	24.8	(19.4 - 30.3)	23,470	1,127,100	14.6	(12.7 - 16.6)
Age (years)								
18-34	*	*	*	*	3,929	993,500	22.3	(17.5 - 27.1)
35-44	*	*	*	*	4,977	421,100	14.8	(11.6 - 17.9)
45-54	159	5,000	26.5	(19.0 - 34.1)	7,807	508,800	18.3	(15.4 - 21.3)
55-64	129	1,900	13.6	(7.4 - 19.7)	8,744	314,500	14.7	(12.1 - 17.3)
≥ 65	209	1,700	12.1	(6.7 - 17.5)	12,572	245,600	9.7	(7.2 - 12.3)
Race/Ethnicity								
White Non-Hispanic	649	21,400	23.6	(19.3 - 27.9)	35,729	1,615,500	17.3	(15.9 - 18.8)
Black Non-Hispanic	*	*	*	*	735	284,100	13.6	(8.9 - 18.2)
Hispanic	*	*	*	*	858	350,900	15.8	(9.6 - 21.9)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	336	13,400	29.6	(22.1 - 37.1)	15,654	1,237,300	22.4	(19.1 - 25.6)
Some College	155	6,500	27.7	(18.4 - 37.0)	9,875	670,600	19.1	(15.5 - 22.8)
≥ College Degree	175	3,100	12.6	(6.8 - 18.4)	12,421	563,300	10.0	(7.5 - 12.5)
Income								
≤ \$24,999	*	*	*	*	9,626	835,100	24.1	(19.1 - 29.1)
\$25,000-\$49,999	177	6,200	25.9	(17.6 - 34.3)	9,629	558,100	19.4	(15.6 - 23.2)
\$50,000-\$74,999	*	*	*	*	5,756	338,100	17.3	(13.0 - 21.7)
≥ \$75,000	116	2,100	9.8	(3.6 - 16.1)	8,549	468,700	10.4	(8.6 - 12.3)

¹ Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day or some days

² Sample size from survey

³ Estimated number of adult current smokers (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 54b. Current smoking among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	669	24.7	(20.0 - 29.4)	38,029	17.0	(15.3 - 18.8)
Gender						
Male	255	24.4	(16.8 - 32.1)	14,559	19.2	(16.3 - 22.1)
Female	414	25.0	(19.5 - 30.6)	23,470	15.0	(13.0 - 17.0)
Race/Ethnicity						
White Non-Hispanic	649	23.6	(19.3 - 28.0)	35,729	17.7	(16.2 - 19.2)
Black Non-Hispanic	*	*	*	735	13.5	(8.8 - 18.2)
Hispanic	*	*	*	858	15.9	(9.8 - 22.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	336	29.6	(22.1 - 37.1)	15,654	22.8	(19.5 - 26.0)
Some College	155	27.2	(18.0 - 36.3)	9,875	19.4	(15.8 - 22.9)
≥ College Degree	175	12.6	(6.6 - 18.5)	12,421	10.0	(7.6 - 12.5)
Income						
≤ \$24,999	*	*	*	9,626	24.1	(19.2 - 29.1)
\$25,000-\$49,999	177	25.3	(17.2 - 33.5)	9,629	19.9	(16.0 - 23.7)
\$50,000-\$74,999	*	*	*	5,756	17.8	(13.3 - 22.3)
≥ \$75,000	116	9.7	(3.4 - 16.1)	8,549	10.6	(8.6 - 12.5)

¹ Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day or some days

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 55a. Everyday smoking among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	669	18,900	20.1	(15.6 - 24.6)	38,029	1,626,800	11.0	(9.8 - 12.2)
Gender								
Male	255	10,400	22.7	(15.0 - 30.3)	14,559	848,400	12.0	(10.0 - 14.1)
Female	414	8,500	17.7	(13.0 - 22.4)	23,470	778,400	10.1	(8.7 - 11.5)
Age (years)								
18-34	*	*	*	*	3,929	559,800	12.6	(9.6 - 15.6)
35-44	*	*	*	*	4,977	315,200	11.0	(8.4 - 13.7)
45-54	159	3,800	20.4	(13.5 - 27.4)	7,807	377,000	13.6	(11.1 - 16.0)
55-64	129	1,400	10.4	(5.0 - 15.8)	8,744	203,000	9.5	(7.5 - 11.5)
≥ 65	209	1,600	11.0	(5.7 - 16.4)	12,572	171,700	6.8	(4.8 - 8.8)
Race/Ethnicity								
White Non-Hispanic	649	17,100	18.8	(14.8 - 22.9)	35,729	1,158,100	12.4	(11.2 - 13.7)
Black Non-Hispanic	*	*	*	*	735	197,700	9.4	(5.6 - 13.3)
Hispanic	*	*	*	*	858	176,200	7.9	(4.2 - 11.6)
Other Non-Hispanic	*	*	*	*	707	94,900	8.5	(3.3 - 13.7)
Education								
≤ High School	336	11,200	24.7	(17.3 - 32.1)	15,654	881,900	15.9	(13.4 - 18.5)
Some College	155	5,000	21.2	(12.5 - 29.9)	9,875	410,200	11.7	(9.6 - 13.8)
≥ College Degree	175	2,500	10.1	(5.0 - 15.2)	12,421	323,500	5.8	(4.3 - 7.2)
Income								
≤ \$24,999	*	*	*	*	9,626	452,800	13.1	(10.3 - 15.8)
\$25,000-\$49,999	177	5,800	24.4	(16.1 - 32.6)	9,629	425,900	14.8	(11.9 - 17.7)
\$50,000-\$74,999	*	*	*	*	5,756	222,700	11.4	(8.5 - 14.4)
≥ \$75,000	*	*	*	*	8,549	320,200	7.1	(5.5 - 8.8)

¹ Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day

² Sample size from survey

³ Estimated number of adults who have smoked at least 100 cigarettes in lifetime and currently smoking everyday (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 55b. Everyday smoking among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	669	20.3	(15.8 - 24.8)	38,029	11.3	(10.0 - 12.5)
Gender						
Male	255	22.6	(14.9 - 30.2)	14,559	12.1	(10.1 - 14.1)
Female	414	18.1	(13.2 - 23.0)	23,470	10.5	(9.0 - 12.0)
Race/Ethnicity						
White Non-Hispanic	649	19.1	(15.0 - 23.2)	35,729	12.8	(11.5 - 14.1)
Black Non-Hispanic	*	*	*	735	9.4	(5.5 - 13.4)
Hispanic	*	*	*	858	8.0	(4.3 - 11.7)
Other Non-Hispanic	*	*	*	707	8.5	(3.2 - 13.8)
Education						
≤ High School	336	24.9	(17.5 - 32.3)	15,654	16.4	(13.8 - 19.0)
Some College	155	20.9	(12.4 - 29.4)	9,875	12.1	(9.9 - 14.3)
≥ College Degree	175	10.2	(5.0 - 15.4)	12,421	5.8	(4.3 - 7.3)
Income						
≤ \$24,999	*	*	*	9,626	13.4	(10.6 - 16.2)
\$25,000-\$49,999	177	24.0	(15.9 - 32.1)	9,629	15.3	(12.3 - 18.3)
\$50,000-\$74,999	*	*	*	5,756	11.7	(8.7 - 14.8)
≥ \$75,000	*	*	*	8,549	7.2	(5.5 - 8.9)

¹ Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 56a. Adults living in homes in which smoking is prohibited¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	667	65,600	70.3	(65.4 - 75.2)	38,045	11,997,100	80.8	(79.0 - 82.6)
Gender								
Male	254	29,200	64.3	(56.0 - 72.6)	14,577	5,634,400	79.4	(76.6 - 82.2)
Female	413	36,300	76.1	(71.0 - 81.1)	23,468	6,362,800	82.1	(79.9 - 84.3)
Age (years)								
18-34	*	*	*	*	3,934	3,655,200	82.1	(77.8 - 86.4)
35-44	*	*	*	*	4,985	2,479,700	86.1	(82.5 - 89.7)
45-54	157	12,700	68.3	(60.3 - 76.3)	7,808	2,194,400	77.2	(73.5 - 80.8)
55-64	129	10,400	75.8	(68.1 - 83.5)	8,753	1,732,500	81.0	(78.3 - 83.7)
≥ 65	210	9,800	68.7	(61.2 - 76.1)	12,565	1,935,400	76.5	(73.4 - 79.5)
Race/Ethnicity								
White Non-Hispanic	647	64,200	71.2	(66.6 - 75.8)	35,737	7,426,900	79.8	(78.3 - 81.2)
Black Non-Hispanic	*	*	*	*	738	1,692,100	78.6	(71.9 - 85.3)
Hispanic	*	*	*	*	863	2,003,900	88.7	(84.1 - 93.3)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	334	27,300	61.2	(53.3 - 69.1)	15,647	4,307,200	77.1	(74.0 - 80.2)
Some College	156	17,200	73.7	(64.9 - 82.5)	9,883	2,795,400	78.9	(75.0 - 82.8)
≥ College Degree	174	20,600	83.3	(77.4 - 89.3)	12,437	4,817,300	85.7	(83.3 - 88.2)
Income								
≤ \$24,999	*	*	*	*	9,608	2,627,700	74.9	(70.0 - 79.7)
\$25,000-\$49,999	*	*	*	*	9,640	2,314,500	79.8	(76.0 - 83.5)
\$50,000-\$74,999	*	*	*	*	5,766	1,551,300	78.7	(74.0 - 83.4)
≥ \$75,000	116	19,300	90.9	(85.8 - 95.9)	8,556	3,883,200	86.5	(84.3 - 88.7)

¹ Defined as having a rule that prohibits smoking anywhere inside the home

² Sample size from survey

³ Estimated number of adults who live in a home with a rule that prohibits smoking anywhere inside the home (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 56b. Adults living in homes in which smoking is prohibited¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	667	70.4	(65.5 - 75.3)	38,045	80.9	(79.2 - 82.7)
Gender						
Male	254	64.2	(56.0 - 72.5)	14,577	79.4	(76.7 - 82.2)
Female	413	76.3	(71.2 - 81.4)	23,468	82.3	(80.0 - 84.5)
Race/Ethnicity						
White Non-Hispanic	647	71.2	(66.5 - 75.9)	35,737	79.9	(78.4 - 81.4)
Black Non-Hispanic	*	*	*	738	78.6	(71.8 - 85.4)
Hispanic	*	*	*	863	88.9	(84.4 - 93.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	334	61.1	(53.2 - 69.0)	15,647	76.9	(73.8 - 80.0)
Some College	156	73.9	(65.3 - 82.5)	9,883	79.1	(75.2 - 83.0)
≥ College Degree	174	84.1	(78.3 - 89.9)	12,437	86.0	(83.5 - 88.5)
Income						
≤ \$24,999	*	*	*	9,608	74.9	(70.1 - 79.7)
\$25,000-\$49,999	176	66.0	(56.3 - 75.8)	9,640	79.8	(76.1 - 83.6)
\$50,000-\$74,999	*	*	*	5,766	78.6	(73.8 - 83.3)
≥ \$75,000	116	92.0	(87.4 - 96.5)	8,556	86.6	(84.3 - 88.9)

¹ Defined as having a rule that prohibits smoking anywhere inside the home

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 57a. Binge drinking within the past month among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	663	21,300	22.8	(17.7 - 27.9)	37,745	2,602,000	17.7	(15.7 - 19.8)
Gender								
Male	251	14,700	32.1	(23.5 - 40.7)	14,420	1,712,400	24.5	(20.9 - 28.1)
Female	412	6,600	13.9	(9.1 - 18.6)	23,325	889,600	11.6	(9.5 - 13.7)
Age (years)								
18-34	*	*	*	*	3,895	1,254,300	28.5	(22.8 - 34.2)
35-44	80	3,400	20.2	(11.0 - 29.3)	4,934	584,800	20.6	(16.5 - 24.7)
45-54	157	3,400	18.4	(11.5 - 25.2)	7,738	467,600	16.6	(13.3 - 19.9)
55-64	126	1,000	7.6	(2.6 - 12.7)	8,663	207,200	9.8	(7.9 - 11.8)
≥ 65	208	900	6.1	(2.0 - 10.2)	12,515	88,100	3.5	(2.5 - 4.5)
Race/Ethnicity								
White Non-Hispanic	644	19,800	21.9	(17.1 - 26.6)	35,453	1,761,900	19.2	(17.5 - 20.8)
Black Non-Hispanic	*	*	*	*	728	247,000	11.6	(6.3 - 17.0)
Hispanic	*	*	*	*	854	460,600	20.5	(11.3 - 29.8)
Other Non-Hispanic	*	*	*	*	710	132,500	12.1	(3.4 - 20.7)
Education								
≤ High School	333	11,000	24.4	(16.3 - 32.4)	15,491	945,800	17.4	(13.2 - 21.6)
Some College	154	5,800	24.9	(15.7 - 34.0)	9,819	636,600	18.1	(14.6 - 21.5)
≥ College Degree	173	4,500	18.3	(9.7 - 27.0)	12,358	1,018,700	18.2	(15.3 - 21.1)
Income								
≤ \$24,999	*	*	*	*	9,557	469,100	13.5	(8.5 - 18.5)
\$25,000-\$49,999	174	5,400	23.0	(13.3 - 32.6)	9,575	534,300	18.5	(13.0 - 24.0)
\$50,000-\$74,999	*	*	*	*	5,726	378,200	19.2	(14.1 - 24.3)
≥ \$75,000	*	*	*	*	8,501	1,011,400	22.6	(19.7 - 25.6)

¹ Defined as men having 5 or more drinks or women having 4 or more drinks on 1 or more occasion within the past month

² Sample size from survey

³ Estimated number of adult binge drinkers within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 57b. Binge drinking within the past month among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	663	22.5	(17.6 - 27.5)	37,745	18.1	(16.1 - 20.2)
Gender						
Male	251	31.4	(23.0 - 39.8)	14,420	24.5	(21.0 - 27.9)
Female	412	14.1	(9.3 - 18.9)	23,325	12.4	(10.2 - 14.5)
Race/Ethnicity						
White Non-Hispanic	644	21.7	(17.1 - 26.4)	35,453	19.8	(18.1 - 21.6)
Black Non-Hispanic	*	*	*	728	11.8	(6.4 - 17.1)
Hispanic	*	*	*	854	20.2	(11.3 - 29.1)
Other Non-Hispanic	*	*	*	710	12.1	(3.4 - 20.8)
Education						
≤ High School	333	23.7	(15.8 - 31.5)	15,491	17.5	(13.5 - 21.5)
Some College	154	25.2	(16.1 - 34.3)	9,819	18.6	(15.1 - 22.0)
≥ College Degree	173	18.3	(9.7 - 26.9)	12,358	18.8	(15.9 - 21.7)
Income						
≤ \$24,999	*	*	*	9,557	13.4	(8.6 - 18.2)
\$25,000-\$49,999	174	22.5	(13.2 - 31.9)	9,575	18.9	(13.5 - 24.2)
\$50,000-\$74,999	*	*	*	5,726	19.9	(14.8 - 25.1)
≥ \$75,000	*	*	*	8,501	23.2	(20.2 - 26.1)

¹ Defined as men having 5 or more drinks or women having 4 or more drinks on 1 or more occasion within the past month

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 58a. Heavy drinking within the past month among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	652	7,000	7.6	(4.6 - 10.7)	37,553	718,700	4.9	(4.2 - 5.6)
Gender								
Male	250	4,300	9.5	(4.2 - 14.8)	14,330	333,300	4.8	(3.9 - 5.7)
Female	402	2,700	5.8	(2.8 - 8.8)	23,223	385,400	5.0	(4.0 - 6.1)
Age (years)								
18-34	*	*	*	*	3,866	224,900	5.1	(3.5 - 6.8)
35-44	*	*	*	*	4,924	137,200	4.8	(3.0 - 6.7)
45-54	156	2,100	11.5	(5.9 - 17.1)	7,715	144,600	5.2	(4.1 - 6.2)
55-64	*	*	*	*	8,624	117,400	5.6	(3.9 - 7.3)
≥ 65	*	*	*	*	12,424	94,600	3.8	(2.7 - 4.9)
Race/Ethnicity								
White Non-Hispanic	632	7,000	7.8	(4.7 - 11.0)	35,279	572,400	6.3	(5.4 - 7.1)
Black Non-Hispanic	*	*	*	*	723	58,100	2.8	(0.6 - 4.9)
Hispanic	*	*	*	*	848	60,900	2.7	(0.5 - 4.9)
Other Non-Hispanic	*	*	*	*	703	27,200	2.5	(0.8 - 4.2)
Education								
≤ High School	327	3,900	8.8	(3.6 - 14.0)	15,411	228,100	4.2	(3.2 - 5.2)
Some College	151	1,400	6.0	(2.1 - 9.8)	9,759	163,700	4.7	(3.6 - 5.8)
≥ College Degree	172	1,800	7.2	(2.2 - 12.3)	12,306	326,800	5.9	(4.5 - 7.3)
Income								
≤ \$24,999	173	1,300	6.7	(0.4 - 13.0)	9,512	104,200	3.0	(1.8 - 4.1)
\$25,000-\$49,999	172	1,300	5.7	(2.3 - 9.0)	9,537	158,700	5.5	(3.9 - 7.2)
\$50,000-\$74,999	*	*	*	*	5,708	95,400	4.9	(3.1 - 6.7)
≥ \$75,000	115	2,100	10.1	(3.5 - 16.7)	8,481	313,900	7.0	(5.4 - 8.6)

¹ Defined as adult men averaging more than 2 alcoholic drinks per day and adult women averaging more than 1 alcoholic drink per day within the past month

² Sample size from survey

³ Estimated number of adult heavy drinkers within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 58b. Heavy drinking within the past month among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	652	7.6	(4.7 - 10.6)	37,553	5.0	(4.3 - 5.8)
Gender						
Male	250	9.7	(4.5 - 14.9)	14,330	4.9	(3.9 - 5.9)
Female	402	5.6	(2.7 - 8.5)	23,223	5.1	(4.0 - 6.2)
Race/Ethnicity						
White Non-Hispanic	632	7.8	(4.8 - 10.9)	35,279	6.4	(5.5 - 7.2)
Black Non-Hispanic	*	*	*	723	2.9	(0.6 - 5.1)
Hispanic	*	*	*	848	2.8	(0.5 - 5.1)
Other Non-Hispanic	*	*	*	703	2.5	(0.8 - 4.2)
Education						
≤ High School	327	8.7	(3.6 - 13.8)	15,411	4.3	(3.2 - 5.3)
Some College	151	6.0	(2.2 - 9.8)	9,759	4.8	(3.6 - 5.9)
≥ College Degree	172	7.3	(2.2 - 12.4)	12,306	6.0	(4.5 - 7.4)
Income						
≤ \$24,999	173	6.4	(0.4 - 12.4)	9,512	3.0	(1.9 - 4.1)
\$25,000-\$49,999	172	5.9	(2.4 - 9.5)	9,537	5.8	(4.0 - 7.5)
\$50,000-\$74,999	*	*	*	5,708	5.0	(3.1 - 7.0)
≥ \$75,000	115	10.3	(3.5 - 17.1)	8,481	7.0	(5.4 - 8.7)

¹ Defined as adult men averaging more than 2 alcoholic drinks per day and adult women averaging more than 1 alcoholic drink per day within the past month

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 59a. Never or rarely been asked about sexual history during routine check-up among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	612	53,100	63.2	(58.1 - 68.4)	35,193	8,480,600	61.9	(59.2 - 64.5)
Gender								
Male	229	27,800	70.0	(61.6 - 78.5)	13,299	4,357,400	67.7	(63.6 - 71.7)
Female	383	25,300	57.1	(50.9 - 63.4)	21,894	4,123,300	56.8	(53.3 - 60.2)
Age (years)								
18-34	*	*	*	*	3,507	1,804,600	44.3	(37.8 - 50.8)
35-44	*	*	*	*	4,531	1,635,300	62.2	(56.6 - 67.8)
45-54	144	11,500	68.8	(60.7 - 76.9)	7,215	1,715,600	65.4	(60.5 - 70.3)
55-64	116	8,500	68.5	(59.1 - 77.9)	8,168	1,416,200	70.7	(66.6 - 74.8)
≥ 65	196	11,100	82.1	(76.1 - 88.1)	11,772	1,908,900	80.3	(76.9 - 83.8)
Race/Ethnicity								
White Non-Hispanic	597	52,300	63.4	(58.2 - 68.6)	33,090	5,940,300	68.5	(66.3 - 70.7)
Black Non-Hispanic	*	*	*	*	683	831,800	40.8	(31.8 - 49.9)
Hispanic	*	*	*	*	784	1,071,700	53.4	(43.6 - 63.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	305	25,900	65.9	(58.0 - 73.8)	14,264	3,001,200	60.1	(55.4 - 64.9)
Some College	*	*	*	*	9,156	1,972,800	60.3	(54.9 - 65.7)
≥ College Degree	160	15,100	66.8	(58.0 - 75.6)	11,706	3,476,000	65.0	(61.2 - 68.7)
Income								
≤ \$24,999	164	11,000	64.2	(54.6 - 73.8)	8,772	1,529,400	49.3	(42.6 - 56.0)
\$25,000-\$49,999	*	*	*	*	8,932	1,705,000	64.4	(58.0 - 70.7)
\$50,000-\$74,999	*	*	*	*	5,433	1,217,000	64.5	(58.5 - 70.5)
≥ \$75,000	*	*	*	*	8,108	2,934,300	68.2	(64.7 - 71.7)

¹ Defined as respondents who never or rarely (hardly ever) were asked about their sexual history, such as about sexual partners and sexual practices, during a routine check-up at a doctor's office or clinic

² Sample size from survey

³ Estimated number of adults who were never or rarely asked about their sexual history during a routine check-up at a doctor's office or clinic (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 59b. Never or rarely been asked about sexual history during routine check-up among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	612	63.3	(58.2 - 68.5)	35,193	61.4	(58.7 - 64.0)
Gender						
Male	229	71.2	(63.0 - 79.4)	13,299	67.9	(64.0 - 71.9)
Female	383	56.3	(49.9 - 62.7)	21,894	55.5	(52.0 - 59.0)
Race/Ethnicity						
White Non-Hispanic	597	63.5	(58.3 - 68.8)	33,090	67.8	(65.6 - 70.1)
Black Non-Hispanic	*	*	*	683	41.1	(32.0 - 50.2)
Hispanic	*	*	*	784	53.0	(43.3 - 62.6)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	305	66.3	(58.4 - 74.1)	14,264	59.5	(54.7 - 64.3)
Some College	*	*	*	9,156	59.7	(54.4 - 65.0)
≥ College Degree	160	67.1	(58.2 - 76.0)	11,706	64.5	(60.8 - 68.2)
Income						
≤ \$24,999	164	64.0	(54.1 - 73.8)	8,772	48.7	(42.0 - 55.4)
\$25,000-\$49,999	*	*	*	8,932	63.7	(57.5 - 70.0)
\$50,000-\$74,999	*	*	*	5,433	64.2	(58.2 - 70.2)
≥ \$75,000	*	*	*	8,108	67.5	(64.0 - 71.0)

¹ Defined as respondents who never or rarely (hardly ever) were asked about their sexual history, such as about sexual partners and sexual practices, during a routine check-up at a doctor's office or clinic

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 60a. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	386	43,000	71.7	(65.3 - 78.1)	22,788	7,011,100	77.9	(75.2 - 80.5)
Gender								
Male	149	25,700	84.3	(74.5 - 94.0)	9,034	3,618,500	81.0	(77.2 - 84.8)
Female	237	17,200	58.7	(50.8 - 66.6)	13,754	3,392,600	74.8	(71.2 - 78.4)
Age (years)								
18-34	*	*	*	*	3,213	2,362,500	70.8	(65.1 - 76.6)
35-44	*	*	*	*	3,513	1,313,400	71.7	(65.6 - 77.7)
45-54	87	7,800	77.9	(68.9 - 86.9)	4,975	1,290,600	80.3	(75.9 - 84.8)
55-64	72	7,100	88.7	(80.3 - 97.2)	4,955	932,200	87.2	(84.1 - 90.3)
≥ 65	99	6,300	96.1	(92.5 - 99.7)	6,132	1,112,400	95.9	(94.4 - 97.5)
Race/Ethnicity								
White Non-Hispanic	375	42,300	73.4	(67.7 - 79.2)	21,540	4,875,200	78.3	(75.6 - 80.9)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	491	1,002,400	84.8	(77.8 - 91.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	8,879	2,581,700	82.1	(77.9 - 86.3)
Some College	*	*	*	*	5,953	1,599,500	77.5	(72.6 - 82.3)
≥ College Degree	*	*	*	*	7,928	2,824,800	75.6	(71.8 - 79.5)
Income								
≤ \$24,999	*	*	*	*	5,491	1,602,900	78.7	(71.9 - 85.6)
\$25,000-\$49,999	101	12,200	78.9	(69.2 - 88.6)	5,912	1,335,900	78.8	(73.8 - 83.9)
\$50,000-\$74,999	*	*	*	*	3,717	987,500	76.6	(70.4 - 82.8)
≥ \$75,000	*	*	*	*	5,771	2,321,600	77.6	(74.0 - 81.2)

¹ Defined as respondent estimating there were hardly any (0-1 out of 10) or a few (2-3 out of 10) persons their age in their community who have had a sexually transmitted disease such as gonorrhea, chlamydia, herpes or syphilis

² Sample size from survey

³ Estimated number of adults who believed hardly any or a few people their age in their community have had a sexually transmitted disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 60b. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	386	70.9	(64.5 - 77.4)	22,788	77.3	(74.7 - 79.9)
Gender						
Male	149	83.8	(74.2 - 93.5)	9,034	81.1	(77.4 - 84.8)
Female	237	57.9	(49.8 - 65.9)	13,754	73.6	(70.0 - 77.3)
Race/Ethnicity						
White Non-Hispanic	375	72.5	(66.6 - 78.4)	21,540	77.5	(74.9 - 80.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	491	84.6	(77.6 - 91.5)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	8,879	81.6	(77.4 - 85.7)
Some College	*	*	*	5,953	76.8	(72.0 - 81.6)
≥ College Degree	*	*	*	7,928	75.1	(71.3 - 79.0)
Income						
≤ \$24,999	*	*	*	5,491	77.9	(71.0 - 84.7)
\$25,000-\$49,999	*	*	*	5,912	77.9	(72.8 - 83.0)
\$50,000-\$74,999	*	*	*	3,717	76.1	(70.0 - 82.3)
≥ \$75,000	*	*	*	5,771	77.0	(73.4 - 80.6)

¹ Defined as respondent estimating there were hardly any (0-1 out of 10) or a few (2-3 out of 10) persons their age in their community who have had a sexually transmitted disease such as gonorrhea, chlamydia, herpes or syphilis

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 61a. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	629	84,800	93.4	(91.3 - 95.5)	36,287	12,738,300	89.7	(88.2 - 91.3)
Gender								
Male	245	41,600	92.2	(88.4 - 96.0)	13,893	6,104,600	89.4	(87.2 - 91.5)
Female	384	43,300	94.5	(92.6 - 96.5)	22,394	6,633,700	90.1	(87.9 - 92.2)
Age (years)								
18-34	90	29,800	98.0	(94.2 - 100)	3,850	4,007,400	91.3	(87.5 - 95.1)
35-44	79	15,900	96.3	(92.5 - 100)	4,875	2,578,600	93.2	(90.4 - 96.0)
45-54	156	16,800	91.7	(87.3 - 96.1)	7,607	2,416,800	90.5	(87.9 - 93.2)
55-64	123	12,200	92.4	(86.8 - 98.0)	8,411	1,876,500	92.2	(90.0 - 94.5)
≥ 65	181	10,100	81.5	(75.2 - 87.9)	11,544	1,858,900	79.7	(76.6 - 82.8)
Race/Ethnicity								
White Non-Hispanic	611	82,100	93.4	(91.2 - 95.5)	34,102	8,050,700	89.4	(87.7 - 91.0)
Black Non-Hispanic	*	*	*	*	702	1,822,200	91.9	(88.2 - 95.6)
Hispanic	*	*	*	*	820	1,967,200	89.9	(85.2 - 94.7)
Other Non-Hispanic	*	*	*	*	663	898,200	88.6	(79.5 - 97.6)
Education								
≤ High School	305	38,100	88.5	(84.4 - 92.7)	14,507	4,385,300	84.6	(81.9 - 87.2)
Some College	151	22,600	98.9	(97.8 - 100)	9,571	3,094,300	90.0	(86.7 - 93.4)
≥ College Degree	170	23,700	96.6	(93.7 - 99.4)	12,142	5,228,800	95.4	(94.0 - 96.7)
Income								
≤ \$24,999	165	16,300	86.6	(79.3 - 94.0)	9,021	2,833,500	86.5	(82.6 - 90.3)
\$25,000-\$49,999	161	21,600	95.6	(92.8 - 98.3)	9,261	2,529,900	89.5	(86.5 - 92.5)
\$50,000-\$74,999	99	15,900	98.8	(97.1 - 100)	5,644	1,740,000	91.2	(87.4 - 95.1)
≥ \$75,000	115	20,200	95.9	(92.6 - 99.3)	8,397	4,184,300	94.4	(92.5 - 96.3)

¹ Defined as believing it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community

² Sample size from survey

³ Estimated number of adults who believed it is at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 61b. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	629	93.3	(91.2 - 95.4)	36,287	89.8	(88.3 - 91.4)
Gender						
Male	245	91.9	(88.1 - 95.7)	13,893	89.1	(86.9 - 91.3)
Female	384	94.7	(92.7 - 96.6)	22,394	90.5	(88.3 - 92.7)
Race/Ethnicity						
White Non-Hispanic	611	93.4	(91.2 - 95.5)	34,102	89.5	(87.8 - 91.2)
Black Non-Hispanic	*	*	*	702	91.7	(87.8 - 95.6)
Hispanic	*	*	*	820	89.6	(84.7 - 94.5)
Other Non-Hispanic	*	*	*	663	89.0	(80.5 - 97.6)
Education						
≤ High School	305	88.6	(84.6 - 92.7)	14,507	84.6	(81.8 - 87.4)
Some College	151	98.9	(97.9 - 99.9)	9,571	90.2	(86.9 - 93.5)
≥ College Degree	170	96.3	(93.2 - 99.4)	12,142	95.4	(94.0 - 96.7)
Income						
≤ \$24,999	165	86.8	(79.7 - 93.9)	9,021	86.5	(82.6 - 90.4)
\$25,000-\$49,999	161	95.6	(92.8 - 98.3)	9,261	89.6	(86.5 - 92.6)
\$50,000-\$74,999	99	99.0	(97.6 - 100)	5,644	91.1	(86.9 - 95.3)
≥ \$75,000	115	95.6	(91.9 - 99.3)	8,397	94.5	(92.6 - 96.4)

¹ Defined as believing it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 62a. Overweight among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	638	32,100	35.8	(30.8 - 40.9)	36,710	5,213,000	36.4	(34.0 - 38.8)
Gender								
Male	252	18,900	42.2	(34.0 - 50.4)	14,502	3,073,200	43.7	(39.9 - 47.6)
Female	386	13,200	29.5	(23.8 - 35.1)	22,208	2,139,800	29.3	(26.4 - 32.2)
Age (years)								
18-34	*	*	*	*	3,789	1,295,200	30.1	(24.7 - 35.6)
35-44	*	*	*	*	4,785	1,052,000	37.5	(31.7 - 43.3)
45-54	154	7,000	38.4	(30.0 - 46.9)	7,495	1,053,300	38.8	(34.1 - 43.4)
55-64	119	4,300	33.5	(24.0 - 43.0)	8,356	798,100	38.8	(34.8 - 42.7)
≥ 65	202	5,500	39.9	(32.1 - 47.7)	12,285	1,014,400	41.5	(37.6 - 45.4)
Race/Ethnicity								
White Non-Hispanic	619	31,800	36.7	(31.6 - 41.7)	34,501	3,282,200	36.4	(34.4 - 38.3)
Black Non-Hispanic	*	*	*	*	708	740,000	36.7	(28.3 - 45.1)
Hispanic	*	*	*	*	813	781,500	35.6	(27.1 - 44.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	320	16,300	37.5	(29.6 - 45.4)	15,079	1,889,700	35.4	(31.2 - 39.5)
Some College	148	6,500	30.0	(21.3 - 38.7)	9,515	1,314,500	38.7	(33.8 - 43.7)
≥ College Degree	167	9,100	37.7	(28.8 - 46.7)	12,062	1,996,900	36.3	(32.8 - 39.7)
Income								
≤ \$24,999	169	5,800	30.4	(20.6 - 40.1)	9,346	1,170,900	34.0	(28.2 - 39.8)
\$25,000-\$49,999	*	*	*	*	9,368	1,136,000	40.4	(34.6 - 46.2)
\$50,000-\$74,999	*	*	*	*	5,615	738,700	38.3	(32.3 - 44.3)
≥ \$75,000	*	*	*	*	8,337	1,659,100	37.5	(34.1 - 40.9)

¹ Defined as having a body mass index (BMI) more than 24.9 but less than 30.0. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Sample size from survey

³ Estimated number of adults who are overweight (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 62b. Overweight among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	638	35.9	(30.9 - 40.9)	36,710	36.2	(33.8 - 38.6)
Gender						
Male	252	42.3	(34.1 - 50.4)	14,502	43.9	(40.1 - 47.6)
Female	386	29.5	(23.7 - 35.3)	22,208	28.9	(25.9 - 31.8)
Race/Ethnicity						
White Non-Hispanic	619	36.7	(31.7 - 41.8)	34,501	36.0	(34.0 - 38.0)
Black Non-Hispanic	*	*	*	708	37.0	(28.5 - 45.5)
Hispanic	*	*	*	813	36.0	(27.5 - 44.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	320	37.2	(29.4 - 45.0)	15,079	35.2	(31.1 - 39.4)
Some College	148	30.4	(21.5 - 39.2)	9,515	38.3	(33.3 - 43.2)
≥ College Degree	167	37.9	(28.8 - 47.0)	12,062	36.2	(32.7 - 39.8)
Income						
≤ \$24,999	169	29.7	(20.2 - 39.3)	9,346	33.8	(28.0 - 39.5)
\$25,000-\$49,999	*	*	*	9,368	40.6	(34.8 - 46.4)
\$50,000-\$74,999	*	*	*	5,615	38.6	(32.5 - 44.7)
≥ \$75,000	*	*	*	8,337	37.2	(33.7 - 40.6)

¹ Defined as having a body mass index (BMI) more than 24.9 but less than 30.0. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 63a. Obesity among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	638	28,800	32.1	(27.3 - 36.9)	36,710	3,327,400	23.2	(21.2 - 25.3)
Gender								
Male	252	15,300	34.0	(26.1 - 41.9)	14,502	1,505,800	21.4	(19.0 - 23.9)
Female	386	13,500	30.2	(24.7 - 35.7)	22,208	1,821,600	25.0	(21.8 - 28.1)
Age (years)								
18-34	*	*	*	*	3,789	730,600	17.0	(12.4 - 21.6)
35-44	*	*	*	*	4,785	662,200	23.6	(19.4 - 27.9)
45-54	154	6,600	36.1	(27.8 - 44.5)	7,495	816,800	30.1	(25.3 - 34.8)
55-64	119	5,400	42.5	(32.6 - 52.4)	8,356	563,900	27.4	(23.7 - 31.1)
≥ 65	202	4,200	30.3	(23.2 - 37.5)	12,285	554,000	22.7	(19.2 - 26.1)
Race/Ethnicity								
White Non-Hispanic	619	26,700	30.8	(26.4 - 35.3)	34,501	2,062,700	22.9	(21.2 - 24.6)
Black Non-Hispanic	*	*	*	*	708	703,100	34.9	(26.1 - 43.7)
Hispanic	*	*	*	*	813	456,400	20.8	(14.3 - 27.2)
Other Non-Hispanic	*	*	*	*	688	105,200	9.6	(4.0 - 15.3)
Education								
≤ High School	320	13,900	32.1	(24.6 - 39.5)	15,079	1,561,600	29.2	(25.3 - 33.2)
Some College	148	7,900	36.4	(27.0 - 45.8)	9,515	792,800	23.4	(19.4 - 27.3)
≥ College Degree	167	6,900	28.6	(20.4 - 36.8)	12,062	915,000	16.6	(14.4 - 18.8)
Income								
≤ \$24,999	*	*	*	*	9,346	974,800	28.3	(22.6 - 34.0)
\$25,000-\$49,999	171	7,200	31.1	(22.6 - 39.7)	9,368	584,800	20.8	(17.1 - 24.5)
\$50,000-\$74,999	*	*	*	*	5,615	475,300	24.6	(20.0 - 29.3)
≥ \$75,000	113	6,000	28.8	(19.3 - 38.3)	8,337	867,700	19.6	(17.0 - 22.3)

¹ Defined as having a body mass index (BMI) of 30.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Sample size from survey

³ Estimated number of adults who are obese (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 63b. Obesity among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	638	32.0	(27.2 - 36.8)	36,710	23.1	(21.0 - 25.1)
Gender						
Male	252	34.2	(26.4 - 42.1)	14,502	21.5	(19.0 - 24.0)
Female	386	29.8	(24.2 - 35.4)	22,208	24.6	(21.5 - 27.8)
Race/Ethnicity						
White Non-Hispanic	619	30.8	(26.3 - 35.4)	34,501	22.8	(21.0 - 24.5)
Black Non-Hispanic	*	*	*	708	34.8	(25.9 - 43.6)
Hispanic	*	*	*	813	20.5	(14.1 - 26.9)
Other Non-Hispanic	*	*	*	688	9.5	(3.8 - 15.2)
Education						
≤ High School	320	32.2	(24.8 - 39.6)	15,079	29.4	(25.4 - 33.3)
Some College	148	36.3	(26.9 - 45.8)	9,515	23.1	(19.2 - 27.1)
≥ College Degree	167	28.1	(19.8 - 36.3)	12,062	16.4	(14.2 - 18.5)
Income						
≤ \$24,999	*	*	*	9,346	28.3	(22.6 - 34.1)
\$25,000-\$49,999	171	30.3	(21.9 - 38.7)	9,368	21.0	(17.2 - 24.8)
\$50,000-\$74,999	*	*	*	5,615	24.0	(19.4 - 28.7)
≥ \$75,000	113	28.6	(18.9 - 38.3)	8,337	19.3	(16.7 - 21.9)

¹ Defined as having a body mass index (BMI) of 30.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 64a. Overweight or obesity among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	638	60,900	67.9	(63.1 - 72.8)	36,710	8,540,500	59.6	(57.0 - 62.2)
Gender								
Male	252	34,200	76.2	(68.8 - 83.5)	14,502	4,579,100	65.2	(61.0 - 69.4)
Female	386	26,700	59.7	(53.5 - 65.9)	22,208	3,961,400	54.3	(51.0 - 57.6)
Age (years)								
18-34	*	*	*	*	3,789	2,025,800	47.1	(40.8 - 53.5)
35-44	*	*	*	*	4,785	1,714,200	61.1	(55.6 - 66.6)
45-54	154	13,500	74.6	(67.0 - 82.2)	7,495	1,870,100	68.8	(64.6 - 73.1)
55-64	119	9,600	76.1	(68.2 - 83.9)	8,356	1,362,000	66.2	(61.9 - 70.4)
≥ 65	202	9,700	70.2	(63.3 - 77.2)	12,285	1,568,400	64.1	(60.6 - 67.7)
Race/Ethnicity								
White Non-Hispanic	619	58,500	67.5	(62.6 - 72.4)	34,501	5,344,900	59.3	(57.2 - 61.3)
Black Non-Hispanic	*	*	*	*	708	1,443,100	71.6	(62.5 - 80.7)
Hispanic	*	*	*	*	813	1,237,900	56.4	(46.5 - 66.2)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	320	30,200	69.6	(62.4 - 76.7)	15,079	3,451,400	64.6	(59.8 - 69.4)
Some College	148	14,400	66.3	(56.6 - 76.1)	9,515	2,107,300	62.1	(57.0 - 67.2)
≥ College Degree	167	16,000	66.3	(57.1 - 75.6)	12,062	2,911,900	52.9	(49.2 - 56.6)
Income								
≤ \$24,999	*	*	*	*	9,346	2,145,700	62.3	(55.7 - 68.9)
\$25,000-\$49,999	171	16,900	73.4	(64.0 - 82.9)	9,368	1,720,700	61.2	(54.5 - 67.9)
\$50,000-\$74,999	*	*	*	*	5,615	1,214,000	62.9	(57.1 - 68.8)
≥ \$75,000	*	*	*	*	8,337	2,526,700	57.1	(53.6 - 60.6)

¹ Defined as having a body mass index (BMI) of 25.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Sample size from survey

³ Estimated number of adults who are overweight or obese (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 64b. Overweight or obesity among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	638	67.9	(63.0 - 72.8)	36,710	59.3	(56.8 - 61.9)
Gender						
Male	252	76.5	(69.2 - 83.8)	14,502	65.4	(61.3 - 69.4)
Female	386	59.3	(53.0 - 65.7)	22,208	53.5	(50.1 - 56.8)
Race/Ethnicity						
White Non-Hispanic	619	67.6	(62.6 - 72.5)	34,501	58.8	(56.7 - 60.8)
Black Non-Hispanic	*	*	*	708	71.8	(63.0 - 80.6)
Hispanic	*	*	*	813	56.5	(46.8 - 66.1)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	320	69.4	(62.2 - 76.6)	15,079	64.6	(59.9 - 69.3)
Some College	148	66.7	(57.0 - 76.4)	9,515	61.4	(56.4 - 66.5)
≥ College Degree	167	65.9	(56.7 - 75.2)	12,062	52.6	(48.9 - 56.3)
Income						
≤ \$24,999	*	*	*	9,346	62.1	(55.6 - 68.6)
\$25,000-\$49,999	171	73.6	(64.5 - 82.8)	9,368	61.6	(55.1 - 68.1)
\$50,000-\$74,999	*	*	*	5,615	62.6	(56.7 - 68.6)
≥ \$75,000	*	*	*	8,337	56.5	(52.9 - 60.0)

¹ Defined as having a body mass index (BMI) of 25.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 65a. Received advice about weight by a health professional among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	640	22,800	25.3	(21.2 - 29.4)	36,729	3,929,700	27.4	(25.1 - 29.7)
Gender								
Male	253	11,400	25.3	(18.9 - 31.7)	14,499	1,654,800	23.6	(20.6 - 26.6)
Female	387	11,400	25.4	(20.2 - 30.5)	22,230	2,274,900	31.0	(27.7 - 34.3)
Age (years)								
18-34	89	4,100	13.9	(6.6 - 21.2)	3,796	895,500	20.7	(15.3 - 26.2)
35-44	*	*	*	*	4,779	722,700	25.8	(20.9 - 30.6)
45-54	155	6,600	36.0	(27.7 - 44.3)	7,490	910,900	33.5	(29.0 - 38.1)
55-64	119	5,100	40.1	(30.2 - 50.0)	8,351	741,100	36.1	(32.0 - 40.3)
≥ 65	203	3,100	22.5	(16.0 - 29.0)	12,313	659,500	26.9	(23.0 - 30.7)
Race/Ethnicity								
White Non-Hispanic	621	22,700	26.1	(21.9 - 30.3)	34,502	2,386,200	26.5	(24.6 - 28.4)
Black Non-Hispanic	*	*	*	*	709	615,700	30.4	(22.0 - 38.8)
Hispanic	*	*	*	*	830	555,400	25.1	(17.8 - 32.4)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	321	8,900	20.5	(15.1 - 25.9)	15,116	1,469,700	27.5	(23.6 - 31.4)
Some College	149	6,700	30.9	(22.0 - 39.7)	9,515	946,800	27.8	(23.2 - 32.4)
≥ College Degree	167	7,100	29.5	(21.2 - 37.7)	12,041	1,445,500	26.2	(23.0 - 29.4)
Income								
≤ \$24,999	169	3,600	18.8	(11.9 - 25.7)	9,360	1,079,400	31.4	(25.3 - 37.5)
\$25,000-\$49,999	172	5,900	25.6	(18.0 - 33.3)	9,365	696,900	24.6	(19.8 - 29.4)
\$50,000-\$74,999	*	*	*	*	5,603	507,100	26.3	(21.6 - 31.0)
≥ \$75,000	113	6,100	29.4	(20.0 - 38.8)	8,334	1,207,100	27.3	(24.2 - 30.5)

¹ Defined as being advised by a doctor, nurse or other health professional about their weight (lose, gain or maintain)

² Sample size from survey

³ Estimated number of adults provided advice by a health professional about their weight (lose, gain or maintain) (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 65b. Received advice about weight by a health professional among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	640	24.9	(20.8 - 29.0)	36,729	27.1	(24.8 - 29.4)
Gender						
Male	253	25.0	(18.6 - 31.4)	14,499	23.4	(20.5 - 26.3)
Female	387	24.8	(19.6 - 30.0)	22,230	30.6	(27.3 - 34.0)
Race/Ethnicity						
White Non-Hispanic	621	25.7	(21.5 - 29.9)	34,502	26.1	(24.2 - 28.0)
Black Non-Hispanic	*	*	*	709	30.4	(21.9 - 38.9)
Hispanic	*	*	*	830	24.9	(17.7 - 32.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	321	20.6	(15.0 - 26.2)	15,116	27.3	(23.4 - 31.2)
Some College	149	30.8	(21.9 - 39.7)	9,515	27.4	(22.9 - 32.0)
≥ College Degree	167	27.8	(19.8 - 35.8)	12,041	25.9	(22.7 - 29.0)
Income						
≤ \$24,999	169	18.5	(11.6 - 25.4)	9,360	31.0	(24.9 - 37.0)
\$25,000-\$49,999	172	25.2	(17.6 - 32.9)	9,365	24.9	(20.0 - 29.8)
\$50,000-\$74,999	*	*	*	5,603	25.9	(21.2 - 30.6)
≥ \$75,000	113	28.2	(18.9 - 37.6)	8,334	26.9	(23.7 - 30.0)

¹ Defined as being advised by a doctor, nurse or other health professional about their weight (lose, gain or maintain)

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 66a. Received advice to lose weight by a health professional among adults provided weight advice¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	177	20,800	91.2	(86.5 - 95.8)	9,724	3,412,900	86.8	(83.9 - 89.8)
Gender								
Male	73	10,400	91.4	(85.4 - 97.5)	3,807	1,468,700	88.8	(85.7 - 91.8)
Female	104	10,300	90.9	(83.8 - 98.0)	5,917	1,944,200	85.5	(80.8 - 90.1)
Age (years)								
18-34	*	*	*	*	681	779,700	87.1	(80.6 - 93.6)
35-44	*	*	*	*	1,215	622,100	86.1	(76.9 - 95.2)
45-54	54	5,900	90.0	(81.6 - 98.5)	2,238	813,300	89.3	(84.4 - 94.2)
55-64	*	*	*	*	2,697	645,500	87.1	(81.2 - 93.0)
≥ 65	*	*	*	*	2,893	552,300	83.7	(76.9 - 90.6)
Race/Ethnicity								
White Non-Hispanic	176	20,700	91.2	(86.5 - 95.8)	9,082	2,076,700	87.0	(84.5 - 89.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	251	483,600	87.1	(78.5 - 95.7)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	79	8,000	90.1	(82.3 - 97.8)	3,962	1,239,700	84.3	(78.6 - 90.1)
Some College	*	*	*	*	2,633	814,800	86.1	(80.1 - 92.0)
≥ College Degree	52	6,400	90.5	(81.4 - 99.6)	3,115	1,290,900	89.3	(85.3 - 93.3)
Income								
≤ \$24,999	*	*	*	*	2,539	922,100	85.4	(78.7 - 92.1)
\$25,000-\$49,999	*	*	*	*	2,482	594,100	85.2	(78.4 - 92.1)
\$50,000-\$74,999	*	*	*	*	1,521	446,100	88.0	(83.2 - 92.7)
≥ \$75,000	*	*	*	*	2,270	1,073,200	88.9	(83.7 - 94.1)

¹ Defined as provided advice by a health professional about their weight and advised to lose weight

² Sample size from survey

³ Estimated number of adults provided advice by a health professional about their weight and advised to lose weight (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 66b. Received advice to lose weight by a health professional among adults provided weight advice¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	177	91.3	(86.6 - 96.0)	9,724	86.9	(83.8 - 89.9)
Gender						
Male	73	91.9	(86.2 - 97.7)	3,807	88.7	(85.7 - 91.8)
Female	104	90.6	(83.2 - 98.0)	5,917	85.5	(80.7 - 90.3)
Race/Ethnicity						
White Non-Hispanic	176	91.2	(86.5 - 95.9)	9,082	87.0	(84.4 - 89.6)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	251	87.6	(79.3 - 95.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	79	90.6	(83.1 - 98.2)	3,962	84.6	(78.7 - 90.5)
Some College	*	*	*	2,633	85.6	(79.5 - 91.7)
≥ College Degree	*	*	*	3,115	89.4	(85.4 - 93.3)
Income						
≤ \$24,999	*	*	*	2,539	85.7	(79.0 - 92.3)
\$25,000-\$49,999	*	*	*	2,482	85.3	(78.4 - 92.3)
\$50,000-\$74,999	*	*	*	1,521	88.0	(83.1 - 92.8)
≥ \$75,000	*	*	*	2,270	88.7	(83.1 - 94.2)

¹ Defined as provided advice by a health professional about their weight and advised to lose weight

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 67a. No leisure-time physical activity among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	670	21,400	22.7	(18.9 - 26.6)	38,128	3,541,700	23.9	(21.6 - 26.1)
Gender								
Male	255	9,600	20.8	(14.8 - 26.7)	14,602	1,482,500	20.9	(17.8 - 24.0)
Female	415	11,800	24.6	(19.8 - 29.4)	23,526	2,059,200	26.6	(23.4 - 29.7)
Age (years)								
18-34	92	4,400	14.3	(6.8 - 21.7)	3,938	982,900	22.0	(16.6 - 27.5)
35-44	*	*	*	*	4,988	556,000	19.3	(15.0 - 23.6)
45-54	159	4,300	23.0	(15.7 - 30.3)	7,827	759,000	26.8	(21.7 - 31.9)
55-64	129	3,500	25.2	(17.2 - 33.1)	8,765	464,900	21.7	(18.2 - 25.2)
≥ 65	210	5,000	34.8	(27.5 - 42.1)	12,610	778,800	30.7	(27.3 - 34.2)
Race/Ethnicity								
White Non-Hispanic	650	20,600	22.7	(18.8 - 26.5)	35,816	1,892,800	20.3	(18.6 - 22.0)
Black Non-Hispanic	*	*	*	*	739	690,000	32.0	(23.4 - 40.6)
Hispanic	*	*	*	*	860	600,800	26.7	(19.1 - 34.4)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	336	13,700	30.2	(23.7 - 36.7)	15,683	1,810,200	32.5	(28.4 - 36.5)
Some College	156	3,700	15.7	(9.8 - 21.5)	9,906	814,200	23.0	(18.6 - 27.3)
≥ College Degree	175	3,900	15.9	(9.9 - 21.9)	12,458	841,000	14.9	(11.9 - 18.0)
Income								
≤ \$24,999	*	*	*	*	9,640	1,332,200	38.0	(31.7 - 44.3)
\$25,000-\$49,999	177	6,500	27.3	(18.3 - 36.3)	9,658	648,000	22.3	(18.3 - 26.3)
\$50,000-\$74,999	100	1,200	7.7	(2.6 - 12.8)	5,773	430,700	21.8	(16.6 - 27.0)
≥ \$75,000	116	2,400	11.3	(5.1 - 17.5)	8,565	560,800	12.5	(10.3 - 14.6)

¹ Defined as having not participated in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise

² Sample size from survey

³ Estimated number of adults who have not participated in any physical activities or exercises (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 67b. No leisure-time physical activity among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	670	23.0	(19.1 - 26.9)	38,128	23.7	(21.5 - 25.9)
Gender						
Male	255	21.2	(15.2 - 27.2)	14,602	20.9	(17.8 - 23.9)
Female	415	24.8	(19.8 - 29.8)	23,526	26.3	(23.1 - 29.5)
Race/Ethnicity						
White Non-Hispanic	650	22.9	(19.0 - 26.8)	35,816	20.0	(18.3 - 21.7)
Black Non-Hispanic	*	*	*	739	32.0	(23.3 - 40.6)
Hispanic	*	*	*	860	26.9	(19.2 - 34.5)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	336	30.7	(24.1 - 37.3)	15,683	32.3	(28.3 - 36.4)
Some College	156	15.2	(9.4 - 20.9)	9,906	22.8	(18.5 - 27.2)
≥ College Degree	175	16.3	(9.9 - 22.6)	12,458	14.9	(11.8 - 17.9)
Income						
≤ \$24,999	*	*	*	9,640	37.9	(31.6 - 44.2)
\$25,000-\$49,999	177	27.4	(18.5 - 36.4)	9,658	22.0	(18.0 - 26.0)
\$50,000-\$74,999	100	8.3	(2.5 - 14.1)	5,773	21.6	(16.3 - 26.8)
≥ \$75,000	116	11.4	(5.0 - 17.8)	8,565	12.6	(10.3 - 14.9)

¹ Defined as having not participated in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 68a. Consumption of 5 or more servings of fruits and vegetables per day among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	669	25,300	27.1	(22.5 - 31.7)	38,203	4,023,400	27.0	(24.9 - 29.2)
Gender								
Male	254	8,800	19.4	(12.2 - 26.5)	14,634	1,603,900	22.6	(19.4 - 25.8)
Female	415	16,500	34.5	(28.7 - 40.2)	23,569	2,419,500	31.1	(28.2 - 34.0)
Age (years)								
18-34	*	*	*	*	3,940	1,261,700	28.3	(22.7 - 33.9)
35-44	79	3,900	24.3	(14.5 - 34.0)	4,989	688,000	23.9	(19.5 - 28.3)
45-54	159	3,500	18.9	(12.5 - 25.3)	7,837	746,800	26.2	(22.3 - 30.1)
55-64	129	3,600	26.3	(17.9 - 34.8)	8,777	579,100	27.0	(23.6 - 30.5)
≥ 65	210	3,500	24.7	(18.3 - 31.1)	12,660	747,800	29.4	(26.0 - 32.8)
Race/Ethnicity								
White Non-Hispanic	649	24,900	27.5	(22.9 - 32.2)	35,884	2,538,000	27.2	(25.4 - 28.9)
Black Non-Hispanic	*	*	*	*	740	572,400	26.5	(19.2 - 33.9)
Hispanic	*	*	*	*	864	672,200	29.8	(21.0 - 38.5)
Other Non-Hispanic	*	*	*	*	715	240,800	21.4	(12.0 - 30.8)
Education								
≤ High School	335	10,100	22.5	(15.7 - 29.4)	15,726	1,309,800	23.4	(19.3 - 27.4)
Some College	156	7,100	30.5	(21.3 - 39.8)	9,924	964,300	27.2	(23.0 - 31.3)
≥ College Degree	175	7,800	31.6	(23.4 - 39.9)	12,471	1,747,500	31.0	(27.9 - 34.1)
Income								
≤ \$24,999	176	4,100	21.3	(13.0 - 29.6)	9,667	831,500	23.7	(18.6 - 28.7)
\$25,000-\$49,999	177	5,900	24.8	(15.6 - 34.0)	9,669	917,200	31.6	(25.6 - 37.6)
\$50,000-\$74,999	*	*	*	*	5,777	501,000	25.3	(21.4 - 29.3)
≥ \$75,000	116	6,000	28.3	(19.0 - 37.6)	8,569	1,260,100	28.0	(25.0 - 31.0)

¹ Defined as having consumed 5 or more servings of fruits and vegetables in an average day

² Sample size from survey

³ Estimated number of adults who consume 5 or more servings of fruits and vegetables per day (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 68b. Consumption of 5 or more servings of fruits and vegetables per day among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	669	27.1	(22.5 - 31.6)	38,203	27.1	(24.9 - 29.2)
Gender						
Male	254	19.2	(12.3 - 26.1)	14,634	22.7	(19.6 - 25.9)
Female	415	34.5	(28.7 - 40.4)	23,569	31.0	(28.1 - 34.0)
Race/Ethnicity						
White Non-Hispanic	649	27.4	(22.8 - 32.1)	35,884	27.1	(25.4 - 28.9)
Black Non-Hispanic	*	*	*	740	26.8	(19.4 - 34.2)
Hispanic	*	*	*	864	29.5	(20.9 - 38.1)
Other Non-Hispanic	*	*	*	715	21.8	(12.2 - 31.5)
Education						
≤ High School	335	22.0	(15.4 - 28.6)	15,726	23.3	(19.3 - 27.3)
Some College	156	31.1	(21.7 - 40.4)	9,924	27.1	(22.9 - 31.2)
≥ College Degree	175	31.9	(23.5 - 40.2)	12,471	31.2	(28.0 - 34.3)
Income						
≤ \$24,999	176	21.9	(13.5 - 30.2)	9,667	23.9	(18.8 - 28.9)
\$25,000-\$49,999	177	24.4	(15.5 - 33.2)	9,669	32.0	(26.0 - 38.0)
\$50,000-\$74,999	*	*	*	5,777	25.0	(21.0 - 28.9)
≥ \$75,000	116	27.8	(18.4 - 37.2)	8,569	28.0	(24.9 - 31.1)

¹ Defined as having consumed 5 or more servings of fruits and vegetables in an average day

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Oswego County**

Table 69a. Had a fall within the past 3 months among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	491	6,400	13.9	(10.6 - 17.3)	28,783	1,244,800	17.0	(14.9 - 19.0)
Gender								
Male	188	3,000	13.4	(8.3 - 18.4)	10,966	489,400	14.4	(11.9 - 16.8)
Female	303	3,500	14.5	(10.0 - 18.9)	17,817	755,300	19.2	(16.1 - 22.3)
Age (years)								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	156	2,300	12.3	(6.8 - 17.8)	7,741	445,500	16.0	(12.0 - 20.0)
55-64	127	2,200	16.1	(9.2 - 23.0)	8,454	336,000	16.5	(13.4 - 19.6)
≥ 65	208	2,000	14.0	(8.9 - 19.1)	12,588	463,300	18.4	(15.2 - 21.6)
Race/Ethnicity								
White Non-Hispanic	479	6,300	13.9	(10.5 - 17.3)	27,456	869,000	16.6	(15.1 - 18.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	424	76,100	11.0	(4.5 - 17.4)
Other Non-Hispanic	*	*	*	*	445	44,300	10.0	(3.1 - 16.8)
Education								
≤ High School	254	3,600	16.4	(11.4 - 21.4)	12,417	493,000	17.3	(13.3 - 21.3)
Some College	111	1,500	13.4	(6.4 - 20.3)	7,232	299,400	18.7	(14.6 - 22.7)
≥ College Degree	125	1,300	10.2	(4.2 - 16.2)	9,086	441,500	15.4	(12.9 - 17.9)
Income								
≤ \$24,999	140	2,500	22.8	(14.6 - 31.0)	7,824	423,900	25.0	(18.5 - 31.6)
\$25,000-\$49,999	133	1,900	15.6	(8.8 - 22.5)	7,445	223,600	16.0	(12.8 - 19.2)
\$50,000-\$74,999	*	*	*	*	4,141	151,200	14.6	(10.8 - 18.4)
≥ \$75,000	*	*	*	*	5,874	287,100	12.5	(10.0 - 15.1)

¹ Defined as respondent unintentionally coming to rest on the ground or another lower level one or more times within the past 3 months

² Sample size from survey

³ Estimated number of adults who have had a fall within the past 3 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 69b. Had a fall within the past 3 months among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	491	14.0	(10.6 - 17.4)	28,783	17.3	(15.0 - 19.5)
Gender						
Male	188	13.6	(8.5 - 18.7)	10,966	15.0	(12.1 - 17.8)
Female	303	14.4	(10.0 - 18.8)	17,817	19.4	(16.0 - 22.8)
Race/Ethnicity						
White Non-Hispanic	479	13.9	(10.5 - 17.3)	27,456	16.9	(15.2 - 18.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	424	10.7	(4.2 - 17.1)
Other Non-Hispanic	*	*	*	445	10.0	(2.9 - 17.0)
Education						
≤ High School	254	15.9	(11.1 - 20.8)	12,417	17.4	(13.0 - 21.7)
Some College	111	13.8	(6.7 - 20.9)	7,232	19.0	(14.7 - 23.2)
≥ College Degree	125	10.7	(4.4 - 16.9)	9,086	16.0	(13.1 - 18.8)
Income						
≤ \$24,999	140	22.6	(14.6 - 30.6)	7,824	25.3	(18.3 - 32.4)
\$25,000-\$49,999	133	16.1	(8.9 - 23.2)	7,445	15.9	(12.6 - 19.2)
\$50,000-\$74,999	*	*	*	4,141	14.7	(10.6 - 18.8)
≥ \$75,000	*	*	*	5,874	12.9	(9.9 - 15.8)

¹ Defined as respondent unintentionally coming to rest on the ground or another lower level one or more times within the past 3 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 70a. Had a fall that caused injury within the past 3 months among adults¹ in Oswego County and New York State

	Oswego County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	491	2,300	5.0	(3.0 - 7.1)	28,760	335,700	4.6	(3.7 - 5.5)
Gender								
Male	*	*	*	*	10,957	120,000	3.5	(2.5 - 4.6)
Female	303	1,400	6.0	(3.1 - 8.9)	17,803	215,600	5.5	(4.1 - 6.9)
Age (years)								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	7,737	115,100	4.1	(2.9 - 5.4)
55-64	*	*	*	*	8,449	90,300	4.4	(2.7 - 6.1)
≥ 65	208	800	5.8	(2.5 - 9.1)	12,574	130,300	5.2	(3.4 - 6.9)
Race/Ethnicity								
White Non-Hispanic	479	2,300	5.1	(3.1 - 7.2)	27,433	270,800	5.2	(4.2 - 6.1)
Black Non-Hispanic	*	*	*	*	458	47,900	4.9	(0.9 - 9.0)
Hispanic	*	*	*	*	424	12,400	1.8	(0.2 - 3.3)
Other Non-Hispanic	*	*	*	*	445	4,600	1.0	(0.4 - 1.6)
Education								
≤ High School	254	1,300	5.8	(2.9 - 8.8)	12,404	148,900	5.2	(3.6 - 6.8)
Some College	*	*	*	*	7,228	85,400	5.3	(3.1 - 7.5)
≥ College Degree	*	*	*	*	9,081	101,300	3.5	(2.4 - 4.6)
Income								
≤ \$24,999	140	1,200	11.5	(5.6 - 17.5)	7,812	90,200	5.3	(3.6 - 7.1)
\$25,000-\$49,999	*	*	*	*	7,442	93,200	6.7	(3.9 - 9.4)
\$50,000-\$74,999	*	*	*	*	4,139	37,000	3.6	(1.8 - 5.3)
≥ \$75,000	*	*	*	*	5,872	54,700	2.4	(1.5 - 3.3)

¹ Defined as as respondent unintentionally coming to rest on the ground or another lower level and caused respondent to limit regular activities for at least a day or to go see a doctor one or more times within the past 3 months

² Sample size from survey

³ Estimated number of adults who have had a fall that caused an injury within the past 3 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Oswego County

Table 70b. Had a fall that caused injury within the past 3 months among adults¹ in Oswego County and New York State - Age-Adjusted²

	Oswego County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	491	5.1	(3.0 - 7.1)	28,760	4.6	(3.6 - 5.5)
Gender						
Male	*	*	*	10,957	3.6	(2.4 - 4.7)
Female	303	6.0	(3.1 - 8.9)	17,803	5.5	(4.1 - 6.9)
Race/Ethnicity						
White Non-Hispanic	479	5.2	(3.1 - 7.3)	27,433	5.2	(4.2 - 6.2)
Black Non-Hispanic	*	*	*	458	4.7	(0.7 - 8.7)
Hispanic	*	*	*	424	1.8	(0.1 - 3.4)
Other Non-Hispanic	*	*	*	445	1.0	(0.4 - 1.6)
Education						
≤ High School	254	5.7	(2.8 - 8.6)	12,404	5.2	(3.5 - 6.8)
Some College	*	*	*	7,228	5.3	(3.1 - 7.5)
≥ College Degree	*	*	*	9,081	3.6	(2.4 - 4.7)
Income						
≤ \$24,999	140	11.3	(5.5 - 17.1)	7,812	5.1	(3.3 - 7.0)
\$25,000-\$49,999	*	*	*	7,442	6.6	(3.9 - 9.2)
\$50,000-\$74,999	*	*	*	4,139	3.7	(1.7 - 5.8)
≥ \$75,000	*	*	*	5,872	2.4	(1.4 - 3.3)

¹ Defined as as respondent unintentionally coming to rest on the ground or another lower level and caused respondent to limit regular activities for at least a day or to go see a doctor one or more times within the past 3 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)