# New York State Expanded Behavioral Risk Factor Surveillance System

### **Rockland County**

Final Report July 2008 - June 2009

New York State Department of Health December 2009

Introduction	12
Table 1. Rockland County and New York State adult US Census Bureau population estimates, 2007	13
General Health	
Table 2a. Fair or poor general health among adults in Rockland County and New York State  Table 2b. Fair or poor general health among adults in Rockland County and New York  State - Age-adjusted	14 15
Table 3a. Poor physical health 14 or more days within the past month among adults in Rockland County and New York State	16
Table 3b. Poor physical health 14 or more days within the past month among adults in Rockland County and New York State - Age-adjusted	17
Table 4a. Poor mental health 14 or more days within the past month among adults in Rockland County and New York State	18
Table 4b. Poor mental health 14 or more days within the past month among adults in Rockland County and New York State - Age-adjusted	19
Table 5a. Poor physical or mental health 14 or more days within the past month among adults in Rockland County and New York State	20
Table 5b. Poor physical or mental health 14 or more days within the past month among adults in Rockland County and New York State - Age-adjusted	21
Quality Health Care	
Table 6a. Health insurance among adults in Rockland County and New York State	22
Table 6b. Health insurance among adults in Rockland County and New York State - Age-adjusted	23
Table 7a. Health insurance among adults aged 18-64 in Rockland County and New York State	24
Table 7b. Health insurance among adults aged 18-64 in Rockland County and New York State - Age-adjusted	25

Table 8a. l	No health insurance among adults in Rockland County and New York State	26
	No health insurance among adults in Rockland County and New York State - Age-adjusted	27
	No health insurance among adults aged 18-64 in Rockland County and New York State	28
	No health insurance among adults aged 18-64 in Rockland County and New York State - Age-adjusted	29
Table 10a.	Adults with regular health care provider in Rockland County and New York State	30
Table 10b.	Adults with regular health care provider in Rockland County and New York State - Age-adjusted	31
Table 11a.	Cost prevented visit to doctor within the past year among adults in Rockland County and New York State	32
Table 11b.	Cost prevented visit to doctor within the past year among adults in Rockland County and New York State - Age-adjusted	33
Table 12a.	Visited a doctor for routine checkup within the past year among adults in Rockland County and New York State	34
Table 12b.	Visited a doctor for routine checkup within the past year among adults in Rockland County and New York State - Age-adjusted	35
Table 13a.	Visited a doctor for routine checkup within the past 2 years among adults in Rockland County and New York State	36
Table 13b.	Visited a doctor for routine checkup within the past 2 years among adults in Rockland County and New York State - Age-adjusted	37
Table 14a.	Dentist visit within the past year among adults in Rockland County and New York State	38
Table 14b.	Dentist visit within the past year among adults in Rockland County and New York State - Age-adjusted	39
Table 15a.	Had teeth cleaned within the past year among adults in Rockland County and New York State	40
Table 15b.	Had teeth cleaned within the past year among adults in Rockland County and New York State - Age-adjusted	41

Table 16a	a. Permanent teeth extracted due to decay or gum disease among adults in Rockland County and New York State	42
Table 16b	b. Permanent teeth extracted due to decay or gum disease among adults in Rockland County and New York State - Age-adjusted	43
Table 17a	a. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older in Rockland County and New York State	44
Table 17t	b. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older in Rockland County and New York State - Age-adjusted	45
Table 18a	a. Flu shot within the past 12 months in Rockland County and New York State	46
Table 18b	b. Flu shot within the past 12 months in Rockland County and New York State - Age-adjusted	47
Table 19a	a. Flu vaccine that was sprayed in nose within the past 12 months in Rockland County and New York State	48
Table 19t	b. Flu vaccine that was sprayed in nose within the past 12 months in Rockland County and New York State - Age-adjusted	49
Table 20a	a. Flu shot or flu vaccine sprayed in nose within the past 12 months in Rockland County and New York State	50
Table 20t	b. Flu shot or flu vaccine sprayed in nose within the past 12 months in Rockland County and New York State - Age-adjusted	51
Table 21a	a. Flu shot among adults aged 65 and older in Rockland County and New York State	52
Table 21t	b. Flu shot among adults aged 65 and older in Rockland County and New York State - Age-adjusted	53
Table 22a	a. Pneumonia shot or pneumococcal vaccine in Rockland County and New York State	54
Table 22b	o. Pneumonia shot or pneumococcal vaccine in Rockland County and New York State - Age-adjusted	55
Table 23a	a. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older in Rockland County and New York State	56
Table 23b	o. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older in Rockland County and New York State - Age-adjusted	57

### **Cancer Screening**

Table 24a.	Ever had a mammogram among women aged 40 and older in Rockland County and New York State	58
Table 24b.	Ever had a mammogram among women aged 40 and older in Rockland County and New York State - Age-adjusted	59
Table 25a.	Had mammogram within the past 2 years among women aged 40 and older in Rockland County and New York State	60
Table 25b.	Had mammogram within the past 2 years among women aged 40 and older in Rockland County and New York State - Age-adjusted	61
Table 26a.	Had mammogram within the past 2 years among women aged 50 and older in Rockland County and New York State	62
Table 26b.	Had mammogram within the past 2 years among women aged 50 and older in Rockland County and New York State - Age-adjusted	63
Table 27a.	Ever had a Pap test among women in Rockland County and New York State	64
Table 27b.	Ever had a Pap test among women in Rockland County and New York State - Age-adjusted	65
Table 28a.	Had a Pap test within the past 3 years among women in Rockland County and New York State	66
Table 28b.	Had a Pap test within the past 3 years among women in Rockland County and New York State - Age-adjusted	67
Table 29a.	Ever had a digital rectal exam among men aged 40 and older in Rockland County and New York State	68
Table 29b.	Ever had a digital rectal exam among men aged 40 and older in Rockland County and New York State - Age-adjusted	69
Table 30a.	Had digital rectal exam within the past 2 years among men aged 40 and older in Rockland County and New York State	70
Table 30b.	Had digital rectal exam within the past 2 years among men aged 40 and older in Rockland County and New York State - Age-adjusted	71

Table 31a.	Ever had a prostate specific antigen test among men aged 40 and older in Rockland County and New York State	72
Table 31b.	Ever had a prostate specific antigen test among men aged 40 and older in Rockland County and New York State - Age-adjusted	73
Table 32a.	Had prostate specific antigen test within the past 2 years among men aged 40 and older in Rockland County and New York State	74
Table 32b.	Had prostate specific antigen test within the past 2 years among men aged 40 and older in Rockland County and New York State - Age-adjusted	75
Table 33a.	Prostate cancer among men aged 40 and older in Rockland County and New York State	76
Table 33b.	Prostate cancer among men aged 40 and older in Rockland County and New York State - Age-adjusted	77
Table 34a.	Home blood stool test ever used among adults aged 50 and older in Rockland County and New York State	78
Table 34b.	Home blood stool test ever used among adults aged 50 and older in Rockland County and New York State - Age-adjusted	79
Table 35a.	Home blood stool test used within the past year among adults aged 50 and older in Rockland County and New York State	80
Table 35b.	Home blood stool test used within the past year among adults aged 50 and older in Rockland County and New York State - Age-adjusted	81
Table 36a.	Home blood stool test used within the past 2 years among adults aged 50 and older in Rockland County and New York State	82
Table 36b.	Home blood stool test used within the past 2 years among adults aged 50 and older in Rockland County and New York State - Age-adjusted	83
Table 37a.	Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in Rockland County and New York State	84
Table 37b.	Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in Rockland County and New York State - Age-adjusted	85
Table 38a.	Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Rockland County and New York State	86

Table 38b.	Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Rockland County and New York State - Age-adjusted	87
Table 39a.	Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Rockland County and New York State	88
Table 39b.	Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Rockland County and New York State - Age-adjusted	89
Chronic	Diseases	
Table 40a.	High blood pressure among adults in Rockland County and New York State	90
Table 40b.	High blood pressure among adults in Rockland County and New York State - Age-adjusted	91
Table 41a.	Blood pressure medication among adults with high blood pressure in Rockland County and New York State	92
Table 41b.	Blood pressure medication among adults with high blood pressure in Rockland County and New York State - Age-adjusted	93
Table 42a.	Ever had blood cholesterol checked among adults in Rockland County and New York State	94
Table 42b.	Ever had blood cholesterol checked among adults in Rockland County and New York State - Age-adjusted	95
Table 43a.	Blood cholesterol checked within the past 5 years among adults in Rockland County and New York State	96
Table 43b.	Blood cholesterol checked within the past 5 years among adults in Rockland County and New York State - Age-adjusted	97
Table 44a.	Cardiovascular disease among adults in Rockland County and New York State	98
Table 44b.	Cardiovascular disease among adults in Rockland County and New York State - Age-adjusted	99
Table 45a.	Coronary heart disease among adults in Rockland County and New York State	100

Table 45b. Coronary heart disease among adults in Rockland County State - Age-adjusted	and New York 10
Table 46a. Ever diagnosed with diabetes among adults in Rockland C	ounty and New York State 10
Table 46b. Ever diagnosed with diabetes among adults in Rockland C State - Age-adjusted	ounty and New York 10
Table 47a. Ever diagnosed with asthma (lifetime) among adults in Ro York State	ckland County and New
Table 47b. Ever diagnosed with asthma (lifetime) among adults in Ro York State - Age-adjusted	ckland County and New 10
Table 48a. Current asthma among adults in Rockland County and New	w York State 10
Table 48b. Current asthma among adults in Rockland County and New State - Age-adjusted	w York 10
Table 49a. Chronic joint symptoms among adults in Rockland County	and New York State 10
Table 49b. Chronic joint symptoms among adults in Rockland County State - Age-adjusted	and New York 10
Table 50a. Arthritis among adults in Rockland County and New York	State 11
Table 50b. Arthritis among adults in Rockland County and New York	State - Age-adjusted 11
Table 51a. Activity limitations because of physical, mental or emotion in Rockland County and New York State	nal problems among adults
Table 51b. Activity limitations because of physical, mental or emotion in Rockland County and New York State - Age-adjusted	nal problems among adults 11
Table 52a. Have health problem that requires the use of special equiper Rockland County and New York State	ment among adults in
Table 52b. Have health problem that requires the use of special equip.  Rockland County and New York State - Age-adjusted	ment among adults in
Table 53a. Disability among adults in Rockland County and New Yor	k State 11
Table 53b. Disability among adults in Rockland County and New Yor	rk State - Age-adjusted 11

#### **Health Behaviors**

Table 54a.	Current smoking among adults in Rockland County and New York State	118
Table 54b.	Current smoking among adults in Rockland County and New York State - Age-adjusted	119
Table 55a.	Everyday smoking among adults in Rockland County and New York State	120
Table 55b.	Everyday smoking among adults in Rockland County and New York State - Age-adjusted	121
Table 56a.	Adults living in homes in which smoking is prohibited in Rockland County and New York State	122
Table 56b.	Adults living in homes in which smoking is prohibited in Rockland County and New York State - Age-adjusted	123
Table 57a.	Binge drinking within the past month among adults in Rockland County and New York State	124
Table 57b.	Binge drinking within the past month among adults in Rockland County and New York State - Age-adjusted	125
Table 58a.	Heavy drinking within the past month among adults in Rockland County and New York State	126
Table 58b.	Heavy drinking within the past month among adults in Rockland County and New York State - Age-adjusted	127
Table 59a.	Never or rarely been asked about sexual history during routine check-up among adults in Rockland County and New York State	128
Table 59b.	Never or rarely been asked about sexual history during routine check-up among adults in Rockland County and New York State - Age-adjusted	129
Table 60a.	Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults in Rockland County and New York State	130
Table 60b.	Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults in Rockland County and New York State - Age-adjusted	131

Table 61a.	Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults in Rockland County and New York State	132
Table 61b.	Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults in Rockland County and New York State - Age-adjusted	133
Overwei	ight / Obesity	
Table 62a.	Overweight among adults in Rockland County and New York State	134
Table 62b.	Overweight among adults in Rockland County and New York State - Age-adjusted	135
Table 63a.	Obesity among adults in Rockland County and New York State	136
Table 63b.	Obesity among adults in Rockland County and New York State - Age-adjusted	137
Table 64a.	Overweight or obesity among adults in Rockland County and New York State	138
Table 64b.	Overweight or obesity among adults in Rockland County and New York State - Age-adjusted	139
Table 65a.	Received advice about weight by a health professional among adults in Rockland County and New York State	140
Table 65b.	Received advice about weight by a health professional among adults in Rockland County and New York State - Age-adjusted	141
Table 66a.	Received advice to lose weight by a health professional among adults provided weight advice in Rockland County and New York State	142
Table 66b.	Received advice to lose weight by a health professional among adults provided weight advice in Rockland County and New York State - Age-adjusted	143
Physical	Activity / Nutrition	
Table 67a.	No leisure-time physical activity among adults in Rockland County and New York State	144
Table 67b.	No leisure-time physical activity among adults in Rockland County and New York State - Age-adjusted	145

Table 68a. Consumption of 5 or more servings of fruits and vegetables per day among adults in Rockland County and New York State	146
Table 68b. Consumption of 5 or more servings of fruits and vegetables per day among adults in Rockland County and New York State - Age-adjusted	147
Injury	
Table 69a. Had a fall within the past 3 months among adults in Rockland County and New York State	148
Table 69b. Had a fall within the past 3 months among adults in Rockland County and New York State - Age-adjusted	149
Table 70a. Had a fall that caused injury within the past 3 months among adults in Rockland County and New York State	150
Table 70b. Had a fall that caused injury within the past 3 months among adults in Rockland County and New York State - Age-adjusted	151

#### Introduction

The New York State Expanded Behavioral Risk Factor Surveillance System (Expanded BRFSS) is a random digit dialed telephone survey of adults (18 years of age and older) that is representative of the non-institutionalized civilian population with landline telephones living in New York State. The goal of the Expanded BRFSS is to collect county-specific data on preventive health practices, risk behaviors, injuries and preventable chronic and infectious diseases. Topics assessed by the Expanded BRFSS include tobacco use, physical inactivity, diet, use of cancer screening services and other factors linked to the leading causes of morbidity and mortality. For each topic, health indicators are based on answers to one or more questions in the survey. The survey results are weighted to more accurately represent the adult population living in each county. Weighting procedures adjust for the differential selection probabilities by race, age and gender. More detailed information on weighting procedures is available in the technical brief.

Survey data collection occurred in 58 geographic areas of New York State, including each of the 57 counties outside of New York City and New York City (5 counties) as a single area. A standard questionnaire was used in all areas. Data collection for the 2009 Expanded BRFSS took place during the period of July 2008 through June 2009. The goal was to complete about 650 interviews in each area. The final number of interviews varies because of the sampling design of the survey and differential response rates. More detailed information on the sampling design is available in the technical brief.

The New York State Department of Health contracted with RTI International, which is responsible for the data collection, sampling, weighting and creation of the statistical reports. RTI International subcontracted with Clearwater Research Inc. for data collection and additional expertise in sample design and data quality control.

This report presents both crude estimates and estimates that have been age-adjusted to the Year 2000 US Standard Population when they represent the entire adult population of an area. Estimates for specific age groups are not age-adjusted. Age adjustment is a statistical process applied to rates that allows communities with different age structures to be compared. Counties can compare their data with results from New York State as a whole and with other counties. Confidence intervals at the 95% level associated with each health indicator give an estimated range of values which is likely to include the actual prevalence of the indicator. Confidence intervals are similar to margins of error. When the confidence intervals of two estimates of the same indicator from different areas do not overlap, they may be said to be statistically significantly different, i.e., these differences are unlikely related to chance and are considered true differences. If there is any value that is included in both intervals, the two estimates are not statistically significantly different.

The survey results for each of the 69 health indicators included in the survey are typically provided by gender, age, race and ethnicity, education and income in the Expanded BRFSS county reports. The estimates for some of these groupings may be suppressed in geographic areas with low proportions of adult respondents in these groupings.

Table 1. Rockland County and New York State adult US Census Bureau population estimates, <sup>1</sup> 2007

	Rockland (	County	New York State		
	Adult Population <sup>1</sup>	Percent	Adult Population <sup>1</sup>	Percent	
Total	217,218	100	14,884,315	100	
Gender					
Male	105,337	48.5	7,103,791	47.7	
Female	111,881	51.5	7,780,524	52.3	
Age (years)					
18-34	60,157	27.7	4,497,479	30.2	
35-44	37,969	17.5	2,844,115	19.1	
45-54	43,528	20.0	2,854,316	19.2	
55-64	36,626	16.9	2,142,000	14.4	
≥ 65	38,938	17.9	2,546,405	17.1	
Race/Ethnicity					
White Non-Hispanic	150,607	69.3	9,343,884	62.8	
Black Non-Hispanic	23,668	10.9	2,156,941	14.5	
Hispanic	27,819	12.8	2,258,717	15.2	
Other Non-Hispanic	15,124	7.0	1,124,773	7.6	

<sup>&</sup>lt;sup>1</sup> Source: National Center for Health Statistics. Postcensal estimates of the resident population of the United States for July 1, 2000-July 1, 2007, by year, county, age, bridged race, Hispanic origin, and sex (Vintage 2007). Prepared under a collaborative arrangement with the U.S. Census Bureau; released August 7, 2008. Available from: http://www.cdc.gov/nchs/nvss/bridged\_race.htm as of November 25, 2009.

Table 2a. Fair or poor general health among adults in Rockland County and New York State

	Rockland County				New York State			
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	657	20,100	9.3	(6.7 - 11.9)	38,124	2,531,400	17.1	(15.1 - 19.0)
Gender								
Male	244	10,000	9.5	(5.7 - 13.3)	14,599	1,069,700	15.1	(12.4 - 17.8)
Female	413	10,100	9.0	(5.5 - 12.6)	23,525	1,461,700	18.8	(16.1 - 21.6)
Age (years)								
18-34	*	*	*	*	3,940	535,800	12.0	(7.2 - 16.8)
35-44	*	*	*	*	4,985	289,200	10.0	(6.8 - 13.3)
45-54	131	5,300	12.1	(5.4 - 18.9)	7,819	507,800	17.9	(14.0 - 21.8)
55-64	134	6,100	16.9	(7.8 - 26.0)	8,757	424,700	19.9	(16.0 - 23.9)
≥ 65	218	7,700	19.9	(13.9 - 25.8)	12,623	773,900	30.5	(26.8 - 34.2)
Race/Ethnicity								
White Non-Hispanic	550	13,600	9.0	(6.5 - 11.6)	35,811	1,226,600	13.2	(11.7 - 14.6)
Black Non-Hispanic	*	*	*	*	739	457,300	21.3	(14.8 - 27.9)
Hispanic	*	*	*	*	861	684,100	30.4	(22.2 - 38.6)
Other Non-Hispanic	*	*	*	*	713	163,400	14.5	(5.1 - 24.0)
Education								
≤ High School	198	8,600	13.0	(7.0 - 19.1)	15,682	1,466,500	26.3	(22.6 - 30.0)
Some College	116	3,500	8.5	(2.3 - 14.7)	9,904	508,500	14.3	(10.9 - 17.8)
≥ College Degree	339	8,100	7.4	(4.7 - 10.2)	12,457	493,500	8.8	(6.5 - 11.1)
Income								
≤ \$24,999	91	5,400	17.2	(8.3 - 26.0)	9,633	1,269,200	36.3	(30.1 - 42.4)
\$25,000-\$49,999	110	4,700	13.7	(4.4 - 22.9)	9,649	447,600	15.4	(11.9 - 18.9)
\$50,000-\$74,999	*	*	*	*	5,769	237,800	12.1	(8.0 - 16.2)
≥ \$75,000	270	5,300	5.3	(2.4 - 8.3)	8,562	218,300	4.9	(3.6 - 6.1)

 $<sup>^{\</sup>mathrm{1}}$  Defined as reporting fair or poor general health

 $<sup>^{\</sup>rm 2}$  Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of adults who have fair or poor health (rounded to the nearest hundred)

 <sup>4 95%</sup> confidence interval
 \* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 2b. Fair or poor general health among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland (	County	N	New York State				
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>			
Total	657	8.1	(5.8 - 10.4)	38,124	16.7	(14.8 - 18.6)			
Gender									
Male	244	8.9	(5.2 - 12.6)	14,599	15.3	(12.6 - 18.0)			
Female	413	7.3	(4.5 - 10.2)	23,525	18.0	(15.2 - 20.7)			
Race/Ethnicity									
White Non-Hispanic	550	8.1	(5.7 - 10.6)	35,811	12.9	(11.4 - 14.4)			
Black Non-Hispanic	*	*	*	739	20.8	(14.2 - 27.3)			
Hispanic	*	*	*	861	29.9	(21.8 - 37.9)			
Other Non-Hispanic	*	*	*	713	13.8	(4.8 - 22.8)			
Education									
≤ High School	198	11.3	(6.2 - 16.4)	15,682	25.8	(22.1 - 29.5)			
Some College	116	7.7	(1.6 - 13.9)	9,904	14.1	(10.7 - 17.6)			
≥ College Degree	339	6.4	(4.0 - 8.9)	12,457	8.5	(6.3 - 10.8)			
Income									
≤ \$24,999	91	15.4	(7.1 - 23.7)	9,633	35.9	(29.7 - 42.0)			
\$25,000-\$49,999	110	11.3	(3.9 - 18.7)	9,649	15.0	(11.6 - 18.5)			
\$50,000-\$74,999	*	*	*	5,769	11.8	(7.7 - 16.0)			
≥ \$75,000	270	4.7	(1.9 - 7.4)	8,562	4.7	(3.5 - 6.0)			

<sup>&</sup>lt;sup>1</sup> Defined as reporting fair or poor general health

 $<sup>^{2}</sup>$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 3a. Poor physical health 14 or more days within the past month among adults in Rockland County and New York State

		Rockla	nd Count	у		New Y	ork State	<b>e</b>
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	640	15,000	7.0	(4.6 - 9.4)	37,327	1,472,900	10.1	(8.7 - 11.6)
Gender								
Male	239	7,500	7.2	(3.8 - 10.6)	14,325	604,200	8.7	(7.0 - 10.5)
Female	401	7,500	6.8	(3.5 - 10.2)	23,002	868,600	11.4	(9.2 - 13.6)
Age (years)								
18-34	*	*	*	*	3,891	268,100	6.1	(2.7 - 9.5)
35-44	*	*	*	*	4,948	232,100	8.1	(5.2 - 11.1)
45-54	130	5,400	12.5	(4.6 - 20.3)	7,764	350,900	12.5	(9.4 - 15.6)
55-64	*	*	*	*	8,621	271,200	13.0	(10.3 - 15.8)
≥ 65	206	5,500	14.7	(8.9 - 20.4)	12,103	350,600	14.5	(12.1 - 16.9)
Race/Ethnicity								
White Non-Hispanic	534	10,700	7.2	(4.7 - 9.8)	35,073	929,700	10.2	(8.8 - 11.6)
Black Non-Hispanic	*	*	*	*	724	208,200	9.9	(4.8 - 14.9)
Hispanic	*	*	*	*	835	250,300	11.5	(6.5 - 16.4)
Other Non-Hispanic	*	*	*	*	695	84,600	7.6	(1.3 - 13.9)
Education								
≤ High School	192	7,400	11.4	(5.4 - 17.4)	15,178	692,800	12.9	(10.6 - 15.3)
Some College	*	*	*	*	9,751	362,500	10.3	(7.4 - 13.2)
≥ College Degree	333	5,200	4.9	(2.6 - 7.1)	12,327	359,300	6.4	(4.8 - 8.1)
Income								
≤ \$24,999	88	3,200	10.2	(3.4 - 17.1)	9,294	625,600	18.4	(13.6 - 23.3)
\$25,000-\$49,999	104	3,500	10.6	(3.7 - 17.6)	9,495	279,100	9.8	(7.3 - 12.3)
\$50,000-\$74,999	*	*	*	*	5,733	186,400	9.5	(5.8 - 13.2)
≥ \$75,000	267	5,500	5.6	(1.9 - 9.2)	8,510	219,100	4.9	(3.6 - 6.2)

 $<sup>^{\</sup>rm 1}$  Defined as 14 or more days within the past month when physical health was not good

 $<sup>^{2}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults with poor physical health 14 or more days within the past month (rounded to the nearest hundred)

<sup>4 95%</sup> confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 3b. Poor physical health 14 or more days within the past month among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland (	County	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	640	6.7	(4.4 - 9.1)	37,327	9.9	(8.4 - 11.4)		
Gender								
Male	239	6.9	(3.6 - 10.3)	14,325	8.7	(6.9 - 10.5)		
Female	401	6.5	(3.2 - 9.8)	23,002	11.0	(8.7 - 13.2)		
Race/Ethnicity								
White Non-Hispanic	534	7.1	(4.5 - 9.7)	35,073	9.9	(8.5 - 11.3)		
Black Non-Hispanic	*	*	*	724	9.7	(4.6 - 14.9)		
Hispanic	*	*	*	835	11.3	(6.3 - 16.2)		
Other Non-Hispanic	*	*	*	695	7.6	(1.2 - 14.0)		
Education								
≤ High School	192	10.7	(4.9 - 16.4)	15,178	12.7	(10.2 - 15.1)		
Some College	*	*	*	9,751	10.1	(7.2 - 12.9)		
≥ College Degree	333	4.5	(2.3 - 6.6)	12,327	6.3	(4.7 - 8.0)		
Income								
≤ \$24,999	88	9.6	(3.1 - 16.1)	9,294	18.4	(13.5 - 23.4)		
\$25,000-\$49,999	104	9.3	(3.1 - 15.4)	9,495	9.4	(7.0 - 11.9)		
\$50,000-\$74,999	*	*	*	5,733	9.0	(5.3 - 12.6)		
≥ \$75,000	267	5.3	(1.7 - 8.9)	8,510	4.9	(3.5 - 6.2)		

 $<sup>^{\</sup>rm l}$  Defined as 14 or more days within the past month when physical health was not good

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 4a. Poor mental health 14 or more days within the past month among adults<sup>1</sup> in Rockland County and New York State

		Rockla	ind Count	у		New Y	ork State	e
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	644	17,800	8.3	(5.8 - 10.8)	37,528	1,473,200	10.1	(8.6 - 11.5)
Gender								
Male	237	9,400	9.1	(5.0 - 13.2)	14,407	555,200	7.9	(6.3 - 9.6)
Female	407	8,400	7.6	(4.6 - 10.5)	23,121	918,000	12.0	(9.7 - 14.3)
Age (years)								
18-34	*	*	*	*	3,889	496,700	11.2	(7.4 - 15.1)
35-44	*	*	*	*	4,947	254,100	8.9	(6.7 - 11.1)
45-54	130	5,800	13.3	(6.2 - 20.4)	7,749	347,200	12.3	(9.0 - 15.6)
55-64	134	2,900	8.1	(2.8 - 13.3)	8,623	211,300	10.1	(7.9 - 12.3)
≥ 65	208	4,100	10.9	(5.9 - 15.9)	12,320	163,900	6.7	(4.7 - 8.7)
Race/Ethnicity								
White Non-Hispanic	538	13,600	9.2	(6.3 - 12.1)	35,266	899,200	9.8	(8.4 - 11.2)
Black Non-Hispanic	*	*	*	*	722	227,800	10.9	(5.4 - 16.4)
Hispanic	*	*	*	*	845	263,700	11.8	(6.8 - 16.8)
Other Non-Hispanic	*	*	*	*	695	82,500	7.4	(1.4 - 13.5)
Education								
≤ High School	193	5,300	8.2	(4.0 - 12.4)	15,335	665,000	12.2	(10.0 - 14.5)
Some College	111	3,500	8.7	(2.8 - 14.5)	9,771	404,500	11.5	(8.3 - 14.8)
≥ College Degree	336	8,400	7.8	(4.3 - 11.3)	12,342	346,600	6.2	(4.5 - 7.9)
Income								
≤ \$24,999	88	4,400	14.4	(5.7 - 23.0)	9,398	611,500	17.9	(13.0 - 22.8)
\$25,000-\$49,999	106	3,100	9.3	(3.3 - 15.2)	9,547	260,300	9.0	(6.7 - 11.4)
\$50,000-\$74,999	*	*	*	*	5,726	187,300	9.6	(6.0 - 13.1)
≥ \$75,000	268	6,800	6.9	(3.3 - 10.5)	8,520	276,500	6.2	(4.6 - 7.7)

 $<sup>^{\</sup>mathrm{1}}$  Defined as 14 or more days within the past month when mental health was not good

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults with poor mental health 14 or more days within the past month (rounded to the nearest hundred)

<sup>4 95%</sup> confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 4b. Poor mental health 14 or more days within the past month among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland (	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	644	8.1	(5.5 - 10.7)	37,528	10.2	(8.7 - 11.7)		
Gender								
Male	237	9.0	(4.7 - 13.3)	14,407	7.9	(6.3 - 9.6)		
Female	407	7.3	(4.3 - 10.3)	23,121	12.2	(9.8 - 14.6)		
Race/Ethnicity								
White Non-Hispanic	538	8.8	(5.9 - 11.6)	35,266	10.0	(8.5 - 11.4)		
Black Non-Hispanic	*	*	*	722	11.1	(5.5 - 16.7)		
Hispanic	*	*	*	845	11.5	(6.5 - 16.4)		
Other Non-Hispanic	*	*	*	695	7.6	(1.5 - 13.7)		
Education								
≤ High School	193	7.5	(3.6 - 11.4)	15,335	12.3	(10.0 - 14.6)		
Some College	111	8.9	(2.7 - 15.2)	9,771	11.8	(8.5 - 15.1)		
≥ College Degree	336	7.7	(3.9 - 11.5)	12,342	6.3	(4.6 - 8.0)		
Income								
≤ \$24,999	88	13.5	(5.0 - 21.9)	9,398	18.2	(13.2 - 23.1)		
\$25,000-\$49,999	106	8.0	(2.8 - 13.2)	9,547	9.4	(6.9 - 11.9)		
\$50,000-\$74,999	*	*	*	5,726	9.5	(5.9 - 13.1)		
≥ \$75,000	268	7.1	(3.1 - 11.0)	8,520	6.2	(4.6 - 7.7)		

 $<sup>^{\</sup>rm 1}$  Defined as 14 or more days within the past month when mental health was not good

 $<sup>^{2}</sup>$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 5a. Poor physical or mental health 14 or more days within the past month among adults<sup>1</sup> in Rockland County and New York State

		Rockla	ind Coun	ty		New Y	ork State	<b>e</b>
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	632	29,400	13.9	(10.7 - 17.2)	37,005	2,385,900	16.5	(14.8 - 18.2)
Gender								
Male	235	15,100	14.7	(9.7 - 19.7)	14,231	937,700	13.6	(11.4 - 15.8)
Female	397	14,300	13.2	(9.0 - 17.4)	22,774	1,448,200	19.1	(16.5 - 21.7)
Age (years)								
18-34	*	*	*	*	3,859	602,600	13.8	(9.7 - 17.9)
35-44	*	*	*	*	4,924	388,300	13.7	(10.4 - 17.0)
45-54	129	10,000	23.2	(13.6 - 32.7)	7,722	551,000	19.6	(15.8 - 23.4)
55-64	132	4,800	13.3	(6.3 - 20.3)	8,545	395,500	19.2	(16.1 - 22.3)
≥ 65	201	7,800	21.2	(14.6 - 27.9)	11,955	448,500	18.8	(15.9 - 21.7)
Race/Ethnicity								
White Non-Hispanic	527	21,000	14.3	(10.8 - 17.9)	34,778	1,485,700	16.3	(14.8 - 17.8)
Black Non-Hispanic	*	*	*	*	715	343,700	16.5	(10.3 - 22.7)
Hispanic	*	*	*	*	832	412,900	19.0	(12.6 - 25.3)
Other Non-Hispanic	*	*	*	*	680	143,500	13.0	(5.0 - 21.0)
Education								
≤ High School	187	11,400	18.2	(11.1 - 25.3)	15,018	1,077,200	20.3	(17.3 - 23.2)
Some College	109	5,500	14.0	(6.4 - 21.6)	9,672	615,500	17.6	(14.1 - 21.1)
≥ College Degree	332	11,900	11.1	(7.2 - 14.9)	12,244	634,200	11.4	(9.2 - 13.6)
Income								
≤ \$24,999	*	*	*	*	9,199	924,000	27.3	(21.9 - 32.8)
\$25,000-\$49,999	102	6,000	18.1	(9.3 - 26.8)	9,425	456,000	16.1	(13.0 - 19.2)
\$50,000-\$74,999	*	*	*	*	5,696	312,000	16.0	(11.4 - 20.5)
≥ \$75,000	266	11,400	11.6	(6.7 - 16.4)	8,475	447,600	10.0	(8.1 - 12.0)

 $<sup>^{1}</sup>$  Defined as 14 or more days within the past month when physical or mental health prevented usual activities, such as self-care, work or recreation <sup>2</sup> Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults with poor physical or mental health 14 or more days within the past month (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 5b. Poor physical or mental health 14 or more days within the past month among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	ew York	State
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	632	13.6	(10.2 - 16.9)	37,005	16.4	(14.7 - 18.2)
Gender						
Male	235	14.4	(9.3 - 19.6)	14,231	13.6	(11.4 - 15.8)
Female	397	12.8	(8.4 - 17.1)	22,774	19.0	(16.3 - 21.6)
Race/Ethnicity						
White Non-Hispanic	527	13.9	(10.3 - 17.6)	34,778	16.3	(14.7 - 17.8)
Black Non-Hispanic	*	*	*	715	16.7	(10.3 - 23.0)
Hispanic	*	*	*	832	18.5	(12.2 - 24.8)
Other Non-Hispanic	*	*	*	680	13.1	(5.0 - 21.3)
Education						
≤ High School	187	16.8	(10.1 - 23.6)	15,018	20.1	(17.1 - 23.1)
Some College	109	15.2	(6.7 - 23.7)	9,672	17.6	(14.1 - 21.1)
≥ College Degree	332	10.7	(6.6 - 14.7)	12,244	11.4	(9.2 - 13.7)
Income						
≤ \$24,999	*	*	*	9,199	27.6	(22.1 - 33.1)
\$25,000-\$49,999	102	15.9	(8.0 - 23.7)	9,425	16.1	(12.9 - 19.3)
\$50,000-\$74,999	*	*	*	5,696	15.6	(11.0 - 20.1)
≥ \$75,000	266	11.7	(6.6 - 16.7)	8,475	10.0	(8.0 - 12.0)

<sup>&</sup>lt;sup>1</sup> Defined as 14 or more days within the past month when physical or mental health prevented usual activities, such as self-care, work or recreation

<sup>&</sup>lt;sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

 $<sup>^4</sup>$  95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 6a. Health insurance among adults 1 in Rockland County and New York State

		Rockla	nd Coun	ty		New Y	ork State	•
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	656	197,700	91.4	(87.7 - 95.1)	38,119	12,854,200	86.6	(84.2 - 89.0)
Gender								
Male	244	93,800	89.1	(82.7 - 95.5)	14,597	5,881,100	83.0	(78.9 - 87.1)
Female	412	103,900	93.6	(90.0 - 97.3)	23,522	6,973,100	89.9	(87.2 - 92.6)
Age (years)								
18-34	80	51,600	87.1	(77.5 - 96.7)	3,910	3,302,100	74.3	(67.8 - 80.7)
35-44	*	*	*	*	4,986	2,560,300	88.9	(84.7 - 93.1)
45-54	131	39,500	90.7	(83.6 - 97.8)	7,831	2,530,100	88.9	(84.6 - 93.2)
55-64	135	33,400	91.1	(85.5 - 96.8)	8,764	1,986,700	92.8	(90.2 - 95.5)
≥ 65	218	38,400	98.7	(97.1 - 100)	12,628	2,475,000	97.8	(96.7 - 98.8)
Race/Ethnicity								
White Non-Hispanic	549	145,100	96.9	(95.4 - 98.5)	35,811	8,591,200	92.2	(90.6 - 93.8)
Black Non-Hispanic	*	*	*	*	736	1,781,600	82.7	(75.3 - 90.1)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	197	55,100	84.6	(76.2 - 92.9)	15,673	4,448,900	79.7	(74.9 - 84.5)
Some College	116	39,000	95.8	(90.6 - 100)	9,906	3,004,600	85.0	(80.1 - 89.8)
≥ College Degree	339	101,800	93.7	(89.0 - 98.5)	12,459	5,361,700	95.3	(93.3 - 97.2)
Income								
≤ \$24,999	*	*	*	*	9,640	2,437,200	69.7	(62.5 - 76.8)
\$25,000-\$49,999	*	*	*	*	9,662	2,384,000	82.2	(76.1 - 88.3)
\$50,000-\$74,999	78	19,300	91.4	(81.8 - 100)	5,772	1,855,700	94.0	(91.2 - 96.7)
≥ \$75,000	270	97,800	98.6	(95.9 - 100)	8,567	4,388,400	97.5	(96.6 - 98.5)

 $<sup>^{1}</sup>$  Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare  $^{2}$  Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of adults who have health insurance (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 6b. Health insurance among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	656	91.7	(88.0 - 95.5)	38,119	86.7	(84.4 - 89.1)		
Gender								
Male	244	89.6	(83.1 - 96.1)	14,597	83.6	(79.7 - 87.5)		
Female	412	93.7	(90.0 - 97.5)	23,522	89.6	(86.8 - 92.3)		
Race/Ethnicity								
White Non-Hispanic	549	97.2	(95.8 - 98.7)	35,811	92.1	(90.5 - 93.7)		
Black Non-Hispanic	*	*	*	736	82.5	(75.0 - 90.0)		
Hispanic	*	*	*	860	68.7	(59.0 - 78.4)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	197	85.3	(76.9 - 93.7)	15,673	80.1	(75.4 - 84.7)		
Some College	116	96.0	(91.0 - 100)	9,906	84.6	(79.7 - 89.4)		
≥ College Degree	339	93.8	(88.7 - 98.8)	12,459	95.4	(93.6 - 97.2)		
Income								
≤ \$24,999	*	*	*	9,640	70.2	(63.2 - 77.1)		
\$25,000-\$49,999	*	*	*	9,662	81.9	(76.0 - 87.9)		
\$50,000-\$74,999	78	91.7	(82.4 - 100)	5,772	94.0	(91.3 - 96.7)		
≥ \$75,000	270	98.7	(96.2 - 100)	8,567	97.5	(96.6 - 98.5)		

 $<sup>^{1}</sup>$  Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare <sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>&</sup>lt;sup>3</sup> Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval \* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 7a. Health insurance among adults aged 18-641 in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	e
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	427	157,000	89.8	(85.3 - 94.2)	25,165	10,251,300	84.2	(81.3 - 87.1)
Gender								
Male	155	77,000	87.4	(79.9 - 95.0)	10,078	4,839,900	80.3	(75.5 - 85.0)
Female	272	80,000	92.1	(87.5 - 96.7)	15,087	5,411,400	88.0	(84.7 - 91.3)
Age (years)								
18-34	80	51,600	87.1	(77.5 - 96.7)	3,910	3,302,100	74.3	(67.8 - 80.7)
35-44	*	*	*	*	4,981	2,556,600	88.9	(84.7 - 93.1)
45-54	129	39,200	90.6	(83.5 - 97.7)	7,791	2,472,900	88.7	(84.3 - 93.0)
55-64	127	31,900	91.1	(85.3 - 97.0)	8,483	1,919,700	92.8	(90.1 - 95.5)
≥ 65	-	-	-	-	-	-	-	-
Race/Ethnicity								
White Non-Hispanic	340	110,600	96.6	(94.6 - 98.5)	23,354	6,582,200	90.3	(88.3 - 92.4)
Black Non-Hispanic	*	*	*	*	567	1,464,100	80.2	(71.6 - 88.7)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	9,037	3,191,500	74.6	(68.6 - 80.6)
Some College	80	32,500	95.0	(88.8 - 100)	7,031	2,540,400	82.9	(77.4 - 88.4)
≥ College Degree	238	83,500	92.6	(86.9 - 98.3)	9,068	4,502,400	94.5	(92.2 - 96.8)
Income								
≤ \$24,999	*	*	*	*	4,918	1,707,200	62.6	(54.1 - 71.2)
\$25,000-\$49,999	*	*	*	*	6,180	1,783,000	77.7	(70.3 - 85.1)
\$50,000-\$74,999	*	*	*	*	4,554	1,553,400	92.9	(89.7 - 96.1)
≥ \$75,000	217	87,300	98.4	(95.4 - 100)	7,324	3,987,600	97.3	(96.3 - 98.3)

 $<sup>^{1}</sup>$  Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

<sup>&</sup>lt;sup>2</sup> Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of adults aged 18-64 who have health insurance (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 7b. Health insurance among adults aged 18-64<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	427	90.3	(85.8 - 94.8)	25,165	84.3	(81.5 - 87.2)		
Gender								
Male	155	88.0	(80.2 - 95.7)	10,078	80.5	(75.9 - 85.1)		
Female	272	92.5	(87.9 - 97.1)	15,087	87.9	(84.6 - 91.2)		
Race/Ethnicity								
White Non-Hispanic	340	97.0	(95.2 - 98.8)	23,354	90.3	(88.3 - 92.3)		
Black Non-Hispanic	*	*	*	567	80.0	(71.3 - 88.6)		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	9,037	75.1	(69.3 - 80.9)		
Some College	80	95.3	(89.5 - 100)	7,031	82.6	(77.1 - 88.1)		
≥ College Degree	238	92.7	(86.7 - 98.7)	9,068	94.6	(92.5 - 96.8)		
Income								
≤ \$24,999	*	*	*	4,918	63.2	(54.9 - 71.6)		
\$25,000-\$49,999	*	*	*	6,180	77.5	(70.2 - 84.7)		
\$50,000-\$74,999	*	*	*	4,554	93.0	(89.8 - 96.2)		
≥ \$75,000	217	98.5	(95.7 - 100)	7,324	97.3	(96.3 - 98.3)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64  $^2$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 8a. No health insurance among adults<sup>1</sup> in Rockland County and New York State

		Rockla	ınd Count	ty		New Y	ork State	e
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	656	18,600	8.6	(4.9 - 12.3)	38,119	1,989,300	13.4	(11.0 - 15.8)
Gender								
Male	244	11,500	10.9	(4.5 - 17.3)	14,597	1,207,400	17.0	(12.9 - 21.1)
Female	412	7,100	6.4	(2.7 - 10.0)	23,522	781,900	10.1	(7.4 - 12.8)
Age (years)								
18-34	*	*	*	*	3,910	1,144,100	25.7	(19.3 - 32.2)
35-44	*	*	*	*	4,986	319,200	11.1	(6.9 - 15.3)
45-54	131	4,100	9.3	(2.2 - 16.4)	7,831	316,500	11.1	(6.8 - 15.4)
55-64	135	3,200	8.9	(3.2 - 14.5)	8,764	153,300	7.2	(4.5 - 9.8)
≥ 65	*	*	*	*	12,628	56,200	2.2	(1.2 - 3.3)
Race/Ethnicity								
White Non-Hispanic	549	4,600	3.1	(1.5 - 4.6)	35,811	727,100	7.8	(6.2 - 9.4)
Black Non-Hispanic	*	*	*	*	736	372,800	17.3	(9.9 - 24.7)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	197	10,100	15.4	(7.1 - 23.8)	15,673	1,134,400	20.3	(15.5 - 25.1)
Some College	*	*	*	*	9,906	531,400	15.0	(10.2 - 19.9)
≥ College Degree	339	6,800	6.3	(1.5 - 11.0)	12,459	266,300	4.7	(2.8 - 6.7)
Income								
≤ \$24,999	*	*	*	*	9,640	1,061,200	30.3	(23.2 - 37.5)
\$25,000-\$49,999	*	*	*	*	9,662	516,700	17.8	(11.7 - 23.9)
\$50,000-\$74,999	*	*	*	*	5,772	119,100	6.0	(3.3 - 8.8)
≥ \$75,000	*	*	*	*	8,567	111,100	2.5	(1.5 - 3.4)

<sup>&</sup>lt;sup>1</sup> Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare <sup>2</sup> Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of adults who have no health insurance (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 8b. No health insurance among adults 1 in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland (	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	656	8.3	(4.5 - 12.0)	38,119	13.3	(10.9 - 15.6)		
Gender								
Male	244	10.4	(3.9 - 16.9)	14,597	16.4	(12.5 - 20.3)		
Female	412	6.3	(2.5 - 10.0)	23,522	10.4	(7.7 - 13.2)		
Race/Ethnicity								
White Non-Hispanic	549	2.8	(1.3 - 4.2)	35,811	7.9	(6.3 - 9.5)		
Black Non-Hispanic	*	*	*	736	17.5	(10.0 - 25.0)		
Hispanic	*	*	*	860	31.3	(21.6 - 41.0)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	197	14.7	(6.3 - 23.1)	15,673	19.9	(15.3 - 24.6)		
Some College	*	*	*	9,906	15.4	(10.6 - 20.3)		
≥ College Degree	339	6.2	(1.2 - 11.3)	12,459	4.6	(2.8 - 6.4)		
Income								
≤ \$24,999	*	*	*	9,640	29.8	(22.9 - 36.8)		
\$25,000-\$49,999	*	*	*	9,662	18.1	(12.1 - 24.0)		
\$50,000-\$74,999	*	*	*	5,772	6.0	(3.3 - 8.7)		
≥ \$75,000	*	*	*	8,567	2.5	(1.5 - 3.4)		

<sup>&</sup>lt;sup>1</sup> Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare <sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 9a. No health insurance among adults aged 18-641 in Rockland County and New York State

		Rockla	ind Count	ty		New Y	ork State	e
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	427	17,900	10.2	(5.8 - 14.7)	25,165	1,928,400	15.8	(12.9 - 18.7)
Gender								
Male	155	11,100	12.6	(5.0 - 20.1)	10,078	1,190,200	19.7	(15.0 - 24.5)
Female	272	6,900	7.9	(3.3 - 12.5)	15,087	738,200	12.0	(8.7 - 15.3)
Age (years)								
18-34	*	*	*	*	3,910	1,144,100	25.7	(19.3 - 32.2)
35-44	*	*	*	*	4,981	319,200	11.1	(6.9 - 15.3)
45-54	129	4,100	9.4	(2.3 - 16.5)	7,791	316,100	11.3	(7.0 - 15.7)
55-64	127	3,100	8.9	(3.0 - 14.7)	8,483	148,900	7.2	(4.5 - 9.9)
≥ 65	-	-	-	-	-	-	-	-
Race/Ethnicity								
White Non-Hispanic	340	3,900	3.4	(1.5 - 5.4)	23,354	703,100	9.7	(7.6 - 11.7)
Black Non-Hispanic	*	*	*	*	567	361,800	19.8	(11.3 - 28.4)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	9,037	1,087,700	25.4	(19.4 - 31.4)
Some College	*	*	*	*	7,031	522,800	17.1	(11.6 - 22.6)
≥ College Degree	238	6,700	7.4	(1.7 - 13.1)	9,068	261,200	5.5	(3.2 - 7.8)
Income								
≤ \$24,999	*	*	*	*	4,918	1,018,000	37.4	(28.8 - 45.9)
\$25,000-\$49,999	*	*	*	*	6,180	511,100	22.3	(14.9 - 29.7)
\$50,000-\$74,999	*	*	*	*	4,554	118,800	7.1	(3.9 - 10.3)
≥ \$75,000	*	*	*	*	7,324	110,700	2.7	(1.7 - 3.7)

Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

<sup>&</sup>lt;sup>2</sup> Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults aged 18-64 who have no health insurance (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 9b. No health insurance among adults aged 18-64<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland (	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	427	9.7	(5.2 - 14.2)	25,165	15.7	(12.8 - 18.5)		
Gender								
Male	155	12.0	(4.3 - 19.8)	10,078	19.5	(14.9 - 24.1)		
Female	272	7.5	(2.9 - 12.1)	15,087	12.1	(8.8 - 15.4)		
Race/Ethnicity								
White Non-Hispanic	340	3.0	(1.2 - 4.8)	23,354	9.7	(7.7 - 11.7)		
Black Non-Hispanic	*	*	*	567	20.0	(11.4 - 28.7)		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	9,037	24.9	(19.1 - 30.7)		
Some College	*	*	*	7,031	17.4	(11.9 - 22.9)		
≥ College Degree	238	7.3	(1.3 - 13.3)	9,068	5.4	(3.2 - 7.5)		
Income								
≤ \$24,999	*	*	*	4,918	36.8	(28.4 - 45.1)		
\$25,000-\$49,999	*	*	*	6,180	22.5	(15.3 - 29.8)		
\$50,000-\$74,999	*	*	*	4,554	7.0	(3.8 - 10.2)		
≥ \$75,000	*	*	*	7,324	2.7	(1.7 - 3.7)		

<sup>&</sup>lt;sup>1</sup> Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64 <sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 10a. Adults with regular health care provider in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	656	190,700	88.0	(84.3 - 91.8)	38,135	12,324,300	83.0	(80.4 - 85.6)
Gender								
Male	244	89,300	84.8	(78.5 - 91.1)	14,593	5,494,300	77.6	(73.2 - 82.0)
Female	412	101,400	91.1	(87.0 - 95.2)	23,542	6,830,000	87.8	(85.1 - 90.5)
Age (years)								
18-34	*	*	*	*	3,929	2,931,600	65.8	(59.2 - 72.5)
35-44	92	34,300	90.3	(84.0 - 96.7)	4,988	2,471,000	86.1	(81.9 - 90.3)
45-54	130	38,000	87.7	(80.1 - 95.2)	7,823	2,529,300	88.6	(84.6 - 92.7)
55-64	135	34,100	93.1	(88.7 - 97.5)	8,770	2,002,000	93.6	(91.7 - 95.4)
≥ 65	218	37,900	97.3	(94.8 - 99.8)	12,625	2,390,400	94.1	(92.2 - 96.0)
Race/Ethnicity								
White Non-Hispanic	549	136,900	91.3	(88.1 - 94.4)	35,821	8,227,700	88.2	(86.4 - 89.9)
Black Non-Hispanic	*	*	*	*	740	1,753,200	81.3	(72.6 - 89.9)
Hispanic	*	*	*	*	860	1,453,100	64.7	(54.7 - 74.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	197	50,600	77.3	(67.8 - 86.8)	15,692	4,249,200	75.9	(70.9 - 80.8)
Some College	116	36,600	89.8	(82.7 - 96.8)	9,904	2,935,000	82.8	(77.2 - 88.4)
≥ College Degree	339	101,700	93.7	(90.5 - 96.8)	12,456	5,046,100	89.8	(87.1 - 92.6)
Income								
≤ \$24,999	*	*	*	*	9,644	2,554,200	72.7	(65.9 - 79.6)
\$25,000-\$49,999	*	*	*	*	9,655	2,243,100	77.3	(70.0 - 84.6)
\$50,000-\$74,999	78	19,100	90.8	(81.4 - 100)	5,771	1,767,500	89.5	(85.2 - 93.9)
≥ \$75,000	270	92,400	93.1	(89.4 - 96.8)	8,560	4,122,800	91.9	(90.2 - 93.7)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having one or more persons respondent thinks of as a personal doctor or health care provider

 $<sup>^{2} \, \</sup>mathrm{Sample} \, \mathrm{size} \, \mathrm{from} \, \mathrm{survey}$ 

<sup>&</sup>lt;sup>3</sup> Estimated number of adults who have at least one person they think of as their personal doctor or health care provider (rounded to the nearest hundred)
<sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 10b. Adults with regular health care provider<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	656	87.8	(83.9 - 91.7)	38,135	83.0	(80.4 - 85.5)		
Gender								
Male	244	85.1	(78.8 - 91.3)	14,593	78.1	(73.9 - 82.3)		
Female	412	90.4	(85.8 - 95.0)	23,542	87.4	(84.6 - 90.1)		
Race/Ethnicity								
White Non-Hispanic	549	90.7	(87.2 - 94.3)	35,821	88.0	(86.2 - 89.7)		
Black Non-Hispanic	*	*	*	740	81.4	(73.0 - 89.8)		
Hispanic	*	*	*	860	65.2	(55.5 - 75.0)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	197	75.9	(66.1 - 85.8)	15,692	75.8	(71.0 - 80.6)		
Some College	116	90.3	(83.3 - 97.3)	9,904	82.9	(77.5 - 88.3)		
≥ College Degree	339	93.6	(90.3 - 96.9)	12,456	89.8	(87.1 - 92.4)		
Income								
≤ \$24,999	*	*	*	9,644	72.9	(66.1 - 79.6)		
\$25,000-\$49,999	*	*	*	9,655	77.4	(70.5 - 84.3)		
\$50,000-\$74,999	78	91.4	(82.6 - 100)	5,771	89.6	(85.3 - 94.0)		
≥ \$75,000	270	93.0	(89.1 - 96.9)	8,560	91.6	(89.8 - 93.4)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as having one or more persons respondent thinks of as a personal doctor or health care provider

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{\</sup>rm 3}$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 11a. Cost prevented visit to doctor within the past year among adults<sup>1</sup> in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	658	30,200	13.9	(9.7 - 18.1)	38,143	2,034,800	13.7	(11.6 - 15.8)
Gender								
Male	244	18,700	17.7	(10.2 - 25.2)	14,607	903,300	12.7	(9.5 - 16.0)
Female	414	11,500	10.3	(6.6 - 14.0)	23,536	1,131,500	14.6	(11.8 - 17.4)
Age (years)								
18-34	*	*	*	*	3,939	947,300	21.2	(15.6 - 26.9)
35-44	*	*	*	*	4,987	347,900	12.1	(8.0 - 16.2)
45-54	131	6,100	14.0	(6.6 - 21.4)	7,825	382,200	13.5	(9.9 - 17.1)
55-64	135	4,400	12.0	(6.2 - 17.9)	8,766	236,800	11.1	(8.0 - 14.2)
≥ 65	*	*	*	*	12,626	120,600	4.8	(2.5 - 7.0)
Race/Ethnicity								
White Non-Hispanic	551	13,400	8.9	(5.5 - 12.3)	35,833	984,600	10.6	(8.9 - 12.2)
Black Non-Hispanic	*	*	*	*	738	280,000	13.1	(7.2 - 19.0)
Hispanic	*	*	*	*	859	597,900	26.5	(17.3 - 35.7)
Other Non-Hispanic	*	*	*	*	713	172,300	15.3	(5.7 - 24.9)
Education								
≤ High School	199	13,400	20.2	(11.2 - 29.3)	15,696	996,600	17.8	(13.8 - 21.9)
Some College	*	*	*	*	9,911	512,400	14.5	(10.4 - 18.6)
≥ College Degree	339	9,300	8.6	(4.2 - 12.9)	12,453	466,900	8.3	(6.0 - 10.6)
Income								
≤ \$24,999	*	*	*	*	9,639	854,500	24.3	(18.4 - 30.3)
\$25,000-\$49,999	*	*	*	*	9,656	532,900	18.4	(12.5 - 24.3)
\$50,000-\$74,999	*	*	*	*	5,776	202,700	10.3	(6.0 - 14.5)
≥ \$75,000	270	5,800	5.9	(1.5 - 10.3)	8,561	192,000	4.3	(2.9 - 5.6)

 $<sup>^{\</sup>mathrm{1}}$  Defined as needing to see a doctor within the past year but not being able to because of cost

 $<sup>^{2} \, \</sup>mathrm{Sample} \, \mathrm{size} \, \mathrm{from} \, \mathrm{survey}$ 

 $<sup>^3</sup>$  Estimated number of adults who did not see a doctor because of cost within the past year (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 11b. Cost prevented visit to doctor within the past year among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	<b>Rockland County</b>			N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	658	13.6	(9.3 - 17.8)	38,143	13.8	(11.7 - 15.9)		
Gender								
Male	244	17.5	(9.9 - 25.1)	14,607	12.5	(9.4 - 15.6)		
Female	414	9.8	(6.0 - 13.6)	23,536	15.0	(12.1 - 17.9)		
Race/Ethnicity								
White Non-Hispanic	551	8.6	(5.1 - 12.2)	35,833	10.8	(9.2 - 12.5)		
Black Non-Hispanic	*	*	*	738	13.2	(7.2 - 19.3)		
Hispanic	*	*	*	859	26.1	(17.2 - 35.1)		
Other Non-Hispanic	*	*	*	713	14.8	(5.5 - 24.2)		
Education								
≤ High School	199	21.0	(11.6 - 30.4)	15,696	17.8	(13.8 - 21.7)		
Some College	*	*	*	9,911	15.3	(11.0 - 19.5)		
≥ College Degree	339	8.0	(3.6 - 12.4)	12,453	8.2	(6.0 - 10.5)		
Income								
≤ \$24,999	*	*	*	9,639	24.5	(18.6 - 30.4)		
\$25,000-\$49,999	*	*	*	9,656	18.7	(12.9 - 24.6)		
\$50,000-\$74,999	*	*	*	5,776	10.4	(6.1 - 14.7)		
≥ \$75,000	270	5.6	(1.1 - 10.2)	8,561	4.4	(3.1 - 5.8)		

 $<sup>^{\</sup>rm 1}$  Defined as needing to see a doctor within the past year but not being able to because of cost

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 12a. Visited a doctor for routine checkup within the past year among adults in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	•
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	655	147,800	68.5	(63.4 - 73.6)	38,026	10,809,300	72.9	(70.6 - 75.3)
Gender								
Male	243	67,600	64.2	(56.0 - 72.5)	14,592	4,738,400	66.9	(63.0 - 70.9)
Female	412	80,300	72.6	(66.6 - 78.6)	23,434	6,070,900	78.5	(75.9 - 81.0)
Age (years)								
18-34	*	*	*	*	3,921	2,747,400	61.7	(55.6 - 67.9)
35-44	*	*	*	*	4,973	1,883,100	65.5	(60.2 - 70.7)
45-54	131	31,600	72.7	(63.6 - 81.8)	7,808	2,174,100	76.6	(73.2 - 80.0)
55-64	135	27,400	74.9	(65.1 - 84.7)	8,751	1,728,600	81.4	(78.7 - 84.2)
≥ 65	217	35,500	91.3	(87.2 - 95.4)	12,573	2,276,000	90.0	(87.5 - 92.5)
Race/Ethnicity								
White Non-Hispanic	549	107,500	71.7	(66.9 - 76.6)	35,721	6,746,200	72.4	(70.6 - 74.2)
Black Non-Hispanic	*	*	*	*	736	1,688,900	78.8	(70.4 - 87.3)
Hispanic	*	*	*	*	857	1,550,900	69.3	(60.0 - 78.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	197	45,000	68.9	(59.3 - 78.6)	15,640	4,150,600	74.7	(70.7 - 78.7)
Some College	*	*	*	*	9,880	2,524,000	71.3	(66.1 - 76.5)
≥ College Degree	338	75,400	69.8	(62.9 - 76.8)	12,428	4,042,900	71.9	(68.5 - 75.3)
Income								
≤ \$24,999	*	*	*	*	9,616	2,469,200	70.4	(64.2 - 76.6)
\$25,000-\$49,999	*	*	*	*	9,631	2,068,400	71.3	(65.3 - 77.3)
\$50,000-\$74,999	*	*	*	*	5,765	1,466,000	74.8	(69.6 - 80.0)
≥ \$75,000	269	73,300	74.4	(67.5 - 81.3)	8,547	3,176,500	70.7	(67.3 - 74.2)

<sup>&</sup>lt;sup>1</sup> Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past year

<sup>2</sup> Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults who have visited a doctor for a routine checkup within the past year (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 12b. Visited a doctor for routine checkup within the past year among adults in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	<b>Rockland County</b>			N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	655	67.0	(61.6 - 72.5)	38,026	72.7	(70.3 - 75.0)		
Gender								
Male	243	63.9	(55.3 - 72.5)	14,592	67.1	(63.3 - 70.9)		
Female	412	70.0	(63.3 - 76.7)	23,434	77.7	(75.1 - 80.3)		
Race/Ethnicity								
White Non-Hispanic	549	70.7	(65.5 - 75.9)	35,721	71.9	(70.1 - 73.7)		
Black Non-Hispanic	*	*	*	736	79.3	(71.3 - 87.4)		
Hispanic	*	*	*	857	69.2	(60.1 - 78.2)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	197	67.9	(58.1 - 77.7)	15,640	74.4	(70.4 - 78.4)		
Some College	*	*	*	9,880	70.9	(65.9 - 76.0)		
≥ College Degree	338	67.7	(60.0 - 75.4)	12,428	71.7	(68.3 - 75.0)		
Income								
≤ \$24,999	*	*	*	9,616	70.2	(64.1 - 76.2)		
\$25,000-\$49,999	*	*	*	9,631	71.1	(65.3 - 76.9)		
\$50,000-\$74,999	*	*	*	5,765	74.5	(69.2 - 79.8)		
≥ \$75,000	269	72.8	(65.2 - 80.4)	8,547	70.3	(66.9 - 73.8)		

<sup>&</sup>lt;sup>1</sup> Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past year  $^2$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 13a. Visited a doctor for routine checkup within the past 2 years among adults in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	•
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	655	183,300	84.9	(80.5 - 89.4)	38,026	12,726,600	85.9	(83.9 - 87.9)
Gender								
Male	243	85,100	80.9	(73.2 - 88.6)	14,592	5,791,900	81.8	(78.2 - 85.4)
Female	412	98,200	88.8	(84.5 - 93.1)	23,434	6,934,700	89.6	(87.9 - 91.3)
Age (years)								
18-34	*	*	*	*	3,921	3,535,400	79.4	(73.9 - 85.0)
35-44	*	*	*	*	4,973	2,355,600	81.9	(78.0 - 85.7)
45-54	131	39,400	90.4	(84.8 - 96.1)	7,808	2,507,900	88.4	(85.6 - 91.2)
55-64	135	31,500	86.1	(77.2 - 95.1)	8,751	1,924,800	90.7	(88.6 - 92.7)
≥ 65	217	37,900	97.6	(95.1 - 100)	12,573	2,402,900	95.0	(93.5 - 96.5)
Race/Ethnicity								
White Non-Hispanic	549	133,500	89.1	(85.9 - 92.3)	35,721	8,070,000	86.6	(85.4 - 87.9)
Black Non-Hispanic	*	*	*	*	736	1,914,500	89.4	(82.3 - 96.5)
Hispanic	*	*	*	*	857	1,832,800	81.9	(73.7 - 90.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	197	54,900	84.0	(75.5 - 92.5)	15,640	4,713,700	84.8	(81.2 - 88.4)
Some College	*	*	*	*	9,880	2,990,500	84.5	(79.6 - 89.4)
≥ College Degree	338	91,700	84.9	(78.8 - 91.1)	12,428	4,928,300	87.6	(85.3 - 89.9)
Income								
≤ \$24,999	*	*	*	*	9,616	2,911,700	83.0	(77.4 - 88.6)
\$25,000-\$49,999	109	30,400	89.5	(82.4 - 96.6)	9,631	2,375,500	81.9	(76.1 - 87.7)
\$50,000-\$74,999	*	*	*	*	5,765	1,733,500	88.4	(84.9 - 91.9)
≥ \$75,000	269	85,100	86.3	(80.1 - 92.4)	8,547	3,907,000	87.0	(84.8 - 89.2)

<sup>&</sup>lt;sup>1</sup> Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past 2 years

<sup>2</sup> Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults who have visited a doctor for a routine checkup within the past 2 years (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 13b. Visited a doctor for routine checkup within the past 2 years among adults in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	655	84.0	(79.2 - 88.9)	38,026	85.7	(83.7 - 87.6)		
Gender								
Male	243	80.7	(72.6 - 88.8)	14,592	81.9	(78.4 - 85.3)		
Female	412	87.3	(82.0 - 92.5)	23,434	89.1	(87.4 - 90.9)		
Race/Ethnicity								
White Non-Hispanic	549	88.7	(85.2 - 92.2)	35,721	86.2	(85.0 - 87.5)		
Black Non-Hispanic	*	*	*	736	89.6	(83.0 - 96.3)		
Hispanic	*	*	*	857	81.8	(73.8 - 89.8)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	197	84.2	(75.9 - 92.5)	15,640	84.6	(81.1 - 88.1)		
Some College	*	*	*	9,880	84.4	(79.8 - 89.0)		
≥ College Degree	338	83.3	(76.1 - 90.5)	12,428	87.3	(85.0 - 89.6)		
Income								
≤ \$24,999	*	*	*	9,616	83.2	(77.8 - 88.5)		
\$25,000-\$49,999	109	88.5	(80.5 - 96.5)	9,631	81.9	(76.3 - 87.4)		
\$50,000-\$74,999	*	*	*	5,765	88.1	(84.4 - 91.8)		
≥ \$75,000	269	85.2	(78.3 - 92.1)	8,547	86.4	(84.2 - 88.7)		

<sup>&</sup>lt;sup>1</sup> Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past 2 years  $^2$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 14a. Dentist visit within the past year among adults<sup>1</sup> in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	•
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	657	164,200	75.6	(70.8 - 80.5)	38,026	10,540,400	71.1	(68.6 - 73.6)
Gender								
Male	243	74,300	70.6	(62.5 - 78.8)	14,573	4,845,800	68.4	(64.5 - 72.2)
Female	414	89,900	80.3	(75.1 - 85.5)	23,453	5,694,600	73.6	(70.4 - 76.8)
Age (years)								
18-34	*	*	*	*	3,924	2,789,700	62.6	(56.3 - 69.0)
35-44	*	*	*	*	4,979	2,181,000	75.8	(70.9 - 80.7)
45-54	131	33,000	75.9	(66.2 - 85.6)	7,814	2,149,300	75.8	(71.9 - 79.7)
55-64	*	*	*	*	8,742	1,690,500	79.2	(75.9 - 82.6)
≥ 65	217	30,900	79.5	(73.7 - 85.3)	12,567	1,729,800	68.5	(64.7 - 72.3)
Race/Ethnicity								
White Non-Hispanic	550	125,100	83.2	(79.5 - 86.8)	35,721	6,959,900	74.7	(72.7 - 76.6)
Black Non-Hispanic	*	*	*	*	735	1,405,800	66.2	(57.9 - 74.5)
Hispanic	*	*	*	*	862	1,425,600	63.1	(53.7 - 72.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	198	45,400	68.9	(59.6 - 78.1)	15,613	3,620,700	65.1	(60.7 - 69.5)
Some College	*	*	*	*	9,880	2,504,700	70.8	(66.1 - 75.5)
≥ College Degree	339	89,600	82.5	(76.1 - 88.9)	12,453	4,402,400	78.2	(74.6 - 81.8)
Income								
≤ \$24,999	*	*	*	*	9,594	1,972,200	56.2	(49.8 - 62.6)
\$25,000-\$49,999	*	*	*	*	9,649	1,896,000	65.8	(59.5 - 72.1)
\$50,000-\$74,999	*	*	*	*	5,763	1,482,400	75.2	(69.4 - 81.0)
≥ \$75,000	270	85,100	85.7	(79.9 - 91.6)	8,564	3,707,900	82.4	(79.5 - 85.3)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having visited a dentist or a dental clinic for any reason within the past year

 $<sup>^{2} \, \</sup>mathrm{Sample} \, \mathrm{size} \, \mathrm{from} \, \mathrm{survey}$ 

<sup>&</sup>lt;sup>3</sup> Estimated number of adults who have visited a dentist or a dental clinic for any reason within the past year (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a \_\_numerator less than 10)

Table 14b. Dentist visit within the past year among adults<sup>1</sup> in Rockland County and New York
State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	ew York	State
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	657	76.0	(71.1 - 81.0)	38,026	71.1	(68.6 - 73.5)
Gender						
Male	243	71.1	(62.8 - 79.4)	14,573	68.3	(64.6 - 72.1)
Female	414	80.7	(75.3 - 86.0)	23,453	73.6	(70.4 - 76.8)
Race/Ethnicity						
White Non-Hispanic	550	83.2	(79.3 - 87.1)	35,721	74.4	(72.4 - 76.3)
Black Non-Hispanic	*	*	*	735	66.6	(58.3 - 74.9)
Hispanic	*	*	*	862	63.6	(54.4 - 72.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	198	69.5	(60.3 - 78.7)	15,613	65.3	(60.9 - 69.6)
Some College	*	*	*	9,880	70.4	(65.7 - 75.1)
≥ College Degree	339	82.6	(75.9 - 89.2)	12,453	78.1	(74.6 - 81.7)
Income						
≤ \$24,999	*	*	*	9,594	56.4	(50.1 - 62.8)
\$25,000-\$49,999	*	*	*	9,649	65.8	(59.6 - 72.0)
\$50,000-\$74,999	*	*	*	5,763	75.0	(69.1 - 80.8)
≥ \$75,000	270	86.5	(80.9 - 92.1)	8,564	82.1	(79.1 - 85.0)

 $<sup>^{\</sup>rm 1}$  Defined as having visited a dentist or a dental clinic for any reason within the past year

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 15a. Had teeth cleaned within the past year among adults<sup>1</sup> in Rockland County and New York State

		Rockla	nd Coun	ty		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	626	157,700	75.4	(70.4 - 80.3)	34,070	9,985,900	71.7	(69.1 - 74.3)
Gender								
Male	231	72,800	72.3	(63.9 - 80.6)	13,173	4,616,400	69.0	(65.0 - 73.1)
Female	395	85,000	78.2	(72.7 - 83.7)	20,897	5,369,500	74.2	(70.8 - 77.5)
Age (years)								
18-34	*	*	*	*	3,860	2,746,400	63.0	(56.5 - 69.5)
35-44	*	*	*	*	4,875	2,112,100	75.5	(70.7 - 80.3)
45-54	129	31,200	73.0	(63.1 - 82.9)	7,438	2,022,600	73.3	(69.1 - 77.4)
55-64	*	*	*	*	7,906	1,560,900	77.7	(73.8 - 81.6)
≥ 65	195	28,800	81.5	(75.3 - 87.7)	9,991	1,543,900	77.1	(73.7 - 80.5)
Race/Ethnicity								
White Non-Hispanic	522	120,600	83.2	(79.4 - 87.1)	31,982	6,601,500	75.4	(73.4 - 77.4)
Black Non-Hispanic	*	*	*	*	680	1,227,200	61.6	(52.1 - 71.0)
Hispanic	*	*	*	*	789	1,421,300	66.6	(57.1 - 76.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	180	41,800	67.9	(58.1 - 77.7)	12,740	3,388,600	67.7	(63.1 - 72.3)
Some College	*	*	*	*	9,145	2,286,800	68.5	(62.9 - 74.1)
≥ College Degree	333	88,500	83.2	(77.0 - 89.4)	12,128	4,298,500	78.3	(74.8 - 81.7)
Income								
≤ \$24,999	*	*	*	*	7,525	1,861,800	60.3	(53.5 - 67.1)
\$25,000-\$49,999	*	*	*	*	8,709	1,681,700	62.6	(55.4 - 69.8)
\$50,000-\$74,999	*	*	*	*	5,534	1,447,800	76.3	(70.7 - 81.8)
≥ \$75,000	265	83,500	86.1	(80.4 - 91.8)	8,448	3,617,500	80.8	(77.8 - 83.8)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having had teeth cleaned by a dentist or dental hygienist within the past year

 $<sup>^{2} \, \</sup>mathrm{Sample} \, \mathrm{size} \, \mathrm{from} \, \mathrm{survey}$ 

 $<sup>^3</sup>$  Estimated number of adults who have had their teeth cleaned by a dentist or dental hygienist within the past year (rounded to the nearest hundred)
4 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 15b. Had teeth cleaned within the past year among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	<b>Rockland County</b>			N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	626	75.6	(70.5 - 80.7)	34,070	71.7	(69.1 - 74.2)		
Gender								
Male	231	72.5	(64.1 - 81.0)	13,173	69.2	(65.3 - 73.1)		
Female	395	78.5	(72.8 - 84.2)	20,897	73.9	(70.5 - 77.3)		
Race/Ethnicity								
White Non-Hispanic	522	83.0	(78.8 - 87.2)	31,982	75.1	(73.0 - 77.1)		
Black Non-Hispanic	*	*	*	680	62.0	(52.7 - 71.4)		
Hispanic	*	*	*	789	66.9	(57.5 - 76.2)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	180	68.7	(59.0 - 78.3)	12,740	67.7	(63.2 - 72.3)		
Some College	*	*	*	9,145	68.1	(62.6 - 73.6)		
≥ College Degree	333	83.2	(76.7 - 89.7)	12,128	78.3	(74.9 - 81.7)		
Income								
≤ \$24,999	*	*	*	7,525	60.2	(53.4 - 67.0)		
\$25,000-\$49,999	*	*	*	8,709	62.7	(55.7 - 69.7)		
\$50,000-\$74,999	*	*	*	5,534	76.4	(70.9 - 81.9)		
≥ \$75,000	265	86.5	(80.9 - 92.1)	8,448	80.4	(77.4 - 83.4)		

<sup>&</sup>lt;sup>1</sup> Defined as having had teeth cleaned by a dentist or dental hygienist within the past year

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 16a. Permanent teeth extracted due to decay or gum disease among adults in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	652	105,600	48.9	(43.8 - 54.1)	37,652	7,426,500	50.4	(47.8 - 52.9)
Gender								
Male	243	49,200	46.8	(38.7 - 54.8)	14,457	3,367,000	47.8	(43.9 - 51.6)
Female	409	56,500	51.0	(44.6 - 57.3)	23,195	4,059,500	52.7	(49.5 - 56.0)
Age (years)								
18-34	82	13,000	21.6	(12.1 - 31.1)	3,921	1,121,600	25.4	(19.7 - 31.0)
35-44	*	*	*	*	4,964	1,250,000	43.6	(37.9 - 49.2)
45-54	*	*	*	*	7,780	1,595,300	56.1	(51.4 - 60.7)
55-64	*	*	*	*	8,648	1,382,800	65.5	(61.6 - 69.4)
≥ 65	216	30,300	79.0	(73.3 - 84.8)	12,339	2,076,900	83.0	(80.6 - 85.4)
Race/Ethnicity								
White Non-Hispanic	546	75,700	50.7	(45.3 - 56.0)	35,364	4,425,700	47.9	(46.0 - 49.9)
Black Non-Hispanic	*	*	*	*	727	1,244,700	58.2	(48.9 - 67.4)
Hispanic	*	*	*	*	853	1,220,500	54.2	(44.9 - 63.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	197	33,300	50.9	(41.2 - 60.6)	15,427	3,436,000	61.9	(57.2 - 66.6)
Some College	*	*	*	*	9,798	1,726,200	49.3	(44.3 - 54.2)
≥ College Degree	336	47,200	43.7	(36.7 - 50.7)	12,351	2,242,600	40.1	(36.5 - 43.6)
Income								
≤ \$24,999	*	*	*	*	9,483	2,171,900	62.3	(55.8 - 68.8)
\$25,000-\$49,999	*	*	*	*	9,552	1,577,400	54.7	(48.3 - 61.0)
\$50,000-\$74,999	*	*	*	*	5,744	1,000,700	51.5	(45.6 - 57.4)
≥ \$75,000	268	42,000	42.5	(35.0 - 49.9)	8,516	1,738,200	38.8	(35.3 - 42.2)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having had at least one permanent tooth extracted because of tooth decay or gum disease

 $<sup>^{2}\,\</sup>mathrm{Sample}$  size from survey

 $<sup>^3</sup>$  Estimated number of adults who have had at least one permanent tooth extracted because of tooth decay or gum disease (rounded to the nearest hundred)
4 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 16b. Permanent teeth extracted due to decay or gum disease among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	ew York	State
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	652	46.1	(40.8 - 51.4)	37,652	49.4	(46.9 - 51.9)
Gender						
Male	243	44.6	(36.4 - 52.8)	14,457	47.9	(44.1 - 51.7)
Female	409	47.5	(40.7 - 54.3)	23,195	50.7	(47.4 - 54.0)
Race/Ethnicity						
White Non-Hispanic	546	47.1	(41.5 - 52.6)	35,364	46.5	(44.5 - 48.5)
Black Non-Hispanic	*	*	*	727	57.7	(48.6 - 66.9)
Hispanic	*	*	*	853	53.8	(44.6 - 62.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	197	47.5	(37.8 - 57.3)	15,427	61.3	(56.7 - 66.0)
Some College	*	*	*	9,798	47.9	(43.0 - 52.9)
≥ College Degree	336	41.6	(34.2 - 48.9)	12,351	39.0	(35.5 - 42.5)
Income						
≤ \$24,999	*	*	*	9,483	61.6	(55.2 - 68.1)
\$25,000-\$49,999	*	*	*	9,552	54.2	(48.0 - 60.3)
\$50,000-\$74,999	*	*	*	5,744	49.9	(44.0 - 55.9)
≥ \$75,000	268	39.9	(32.2 - 47.5)	8,516	37.7	(34.2 - 41.1)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having had at least one permanent tooth extracted because of tooth decay or gum disease

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>&</sup>lt;sup>3</sup> Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 17a. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older in Rockland County and New York State

		Rockla	nd Count	у		New Y	ork State	9
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	216	3,300	8.6	(4.6 - 12.6)	12,339	491,200	19.6	(16.0 - 23.3)
Gender								
Male	*	*	*	*	4,352	167,200	16.4	(11.8 - 20.9)
Female	130	1,900	8.5	(4.0 - 13.1)	7,987	324,000	21.8	(16.6 - 27.0)
Age (years)								
≥ 65	216	3,300	8.6	(4.6 - 12.6)	12,339	491,200	19.6	(16.0 - 23.3)
Race/Ethnicity								
White Non-Hispanic	200	3,300	9.9	(5.3 - 14.4)	11,885	322,600	16.8	(14.3 - 19.2)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	86	2,100	14.1	(5.6 - 22.7)	6,358	314,100	25.3	(19.9 - 30.7)
Some College	*	*	*	*	2,736	90,600	20.2	(12.1 - 28.2)
≥ College Degree	*	*	*	*	3,214	80,500	10.1	(3.6 - 16.5)
Income								
≤ \$24,999	*	*	*	*	4,558	199,600	26.9	(19.3 - 34.4)
\$25,000-\$49,999	*	*	*	*	3,365	123,900	21.4	(13.1 - 29.6)
\$50,000-\$74,999	*	*	*	*	1,179	21,400	7.3	(3.8 - 10.8)
≥ \$75,000	*	*	*	*	1,187	9,600	2.5	(1.0 - 4.0)

 $<sup>^{</sup>m 1}$  Defined as having all permanent teeth extracted because of tooth decay or gum disease among adults aged 65 and older

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults aged 65 and older who have had all permanent teeth extracted because of tooth decay or gum disease (rounded to the nearest hundred)

4 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 17b. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland (	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	216	8.6	(4.4 - 12.8)	12,339	19.4	(15.7 - 23.2)		
Gender								
Male	*	*	*	4,352	16.5	(11.5 - 21.4)		
Female	130	8.5	(4.0 - 13.1)	7,987	22.1	(16.6 - 27.7)		
Race/Ethnicity								
White Non-Hispanic	200	9.8	(5.1 - 14.5)	11,885	16.6	(13.9 - 19.3)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	86	14.7	(5.6 - 23.8)	6,358	24.9	(19.3 - 30.5)		
Some College	*	*	*	2,736	21.0	(12.1 - 29.9)		
≥ College Degree	*	*	*	3,214	10.0	(3.6 - 16.3)		
Income								
≤ \$24,999	*	*	*	4,558	27.3	(19.1 - 35.5)		
\$25,000-\$49,999	*	*	*	3,365	21.0	(12.9 - 29.1)		
\$50,000-\$74,999	*	*	*	1,179	7.6	(3.6 - 11.5)		
≥ \$75,000	*	*	*	1,187	2.2	(0.9 - 3.5)		

 $<sup>^{</sup>m 1}$  Defined as having all permanent teeth extracted because of tooth decay or gum disease among adults aged 65 and older

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 18a. Flu shot within the past 12 months<sup>1</sup> in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	9
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	656	88,000	40.6	(35.8 - 45.5)	38,079	6,255,900	42.3	(39.9 - 44.8)
Gender								
Male	242	41,400	39.6	(31.9 - 47.2)	14,579	2,798,900	39.6	(35.8 - 43.5)
Female	414	46,600	41.6	(35.5 - 47.8)	23,500	3,456,900	44.8	(41.6 - 47.9)
Age (years)								
18-49	230	28,500	25.0	(17.9 - 32.2)	12,505	2,685,300	31.1	(27.3 - 34.9)
50-64	209	29,200	45.7	(37.3 - 54.1)	12,972	1,662,900	46.0	(42.6 - 49.5)
≥ 65	217	30,300	78.0	(71.7 - 84.3)	12,602	1,907,700	75.2	(72.0 - 78.4)
Race/Ethnicity								
White Non-Hispanic	549	64,300	42.9	(37.8 - 48.0)	35,770	4,029,000	43.3	(41.4 - 45.3)
Black Non-Hispanic	*	*	*	*	738	907,000	42.1	(33.1 - 51.1)
Hispanic	*	*	*	*	860	882,900	39.6	(30.4 - 48.9)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	197	24,500	37.5	(28.8 - 46.1)	15,663	2,457,800	44.3	(39.8 - 48.8)
Some College	*	*	*	*	9,898	1,377,700	38.9	(34.0 - 43.9)
≥ College Degree	339	45,600	42.0	(35.2 - 48.8)	12,436	2,404,400	43.0	(39.5 - 46.4)
Income								
≤ \$24,999	*	*	*	*	9,635	1,656,800	47.6	(41.2 - 54.0)
\$25,000-\$49,999	*	*	*	*	9,647	1,200,600	41.4	(35.6 - 47.3)
\$50,000-\$74,999	*	*	*	*	5,763	792,300	40.2	(34.3 - 46.2)
≥ \$75,000	270	43,500	43.8	(36.4 - 51.3)	8,548	1,776,100	39.6	(36.3 - 42.8)

<sup>&</sup>lt;sup>1</sup> Defined as having had a flu shot within the past 12 months

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults who have had a flu shot within the past 12 months (rounded to the nearest hundred)

 <sup>4 95%</sup> confidence interval
 \* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 18b. Flu shot within the past 12 months<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	656	38.2	(33.2 - 43.2)	38,079	41.7	(39.2 - 44.1)		
Gender								
Male	242	38.3	(30.5 - 46.1)	14,579	40.0	(36.2 - 43.8)		
Female	414	38.1	(31.7 - 44.6)	23,500	43.1	(40.0 - 46.3)		
Race/Ethnicity								
White Non-Hispanic	549	39.8	(34.5 - 45.0)	35,770	42.4	(40.4 - 44.3)		
Black Non-Hispanic	*	*	*	738	42.3	(33.3 - 51.3)		
Hispanic	*	*	*	860	39.0	(29.9 - 48.1)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	197	34.3	(25.8 - 42.7)	15,663	43.7	(39.1 - 48.2)		
Some College	*	*	*	9,898	37.8	(33.0 - 42.7)		
≥ College Degree	339	39.9	(32.8 - 47.0)	12,436	42.5	(39.1 - 46.0)		
Income								
≤ \$24,999	*	*	*	9,635	47.3	(41.0 - 53.7)		
\$25,000-\$49,999	*	*	*	9,647	40.3	(34.5 - 46.0)		
\$50,000-\$74,999	*	*	*	5,763	39.4	(33.4 - 45.4)		
≥ \$75,000	270	41.6	(33.9 - 49.4)	8,548	39.1	(35.8 - 42.4)		

 $<sup>^{\</sup>rm l}$  Defined as having had a flu shot within the past 12 months

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 19a. Flu vaccine that was sprayed in nose within the past 12 months<sup>1</sup> in Rockland County and New York State

		Rockla	nd Count	у		New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	*	*	*	*	38,106	248,300	1.7	(0.6 - 2.8)		
Gender										
Male	*	*	*	*	14,586	148,000	2.1	(0.0 - 4.2)		
Female	*	*	*	*	23,520	100,300	1.3	(0.5 - 2.1)		
Age (years)										
18-49	*	*	*	*	12,535	184,900	2.1	(0.3 - 3.9)		
50-64	*	*	*	*	12,968	36,000	1.0	(0.2 - 1.8)		
≥ 65	*	*	*	*	12,603	27,400	1.1	(0.4 - 1.8)		
Race/Ethnicity										
White Non-Hispanic	*	*	*	*	35,796	66,000	0.7	(0.4 - 1.0)		
Black Non-Hispanic	*	*	*	*	740	38,500	1.8	(0.0 - 4.1)		
Hispanic	*	*	*	*	860	133,700	5.9	(0.0 - 12.3)		
Other Non-Hispanic	*	*	*	*	*	*	*	*		
Education										
≤ High School	*	*	*	*	15,679	172,500	3.1	(0.3 - 5.9)		
Some College	*	*	*	*	9,897	43,500	1.2	(0.4 - 2.0)		
≥ College Degree	*	*	*	*	12,449	31,600	0.6	(0.3 - 0.8)		
Income										
≤ \$24,999	*	*	*	*	9,636	138,400	3.9	(0.0 - 8.1)		
\$25,000-\$49,999	*	*	*	*	9,648	41,500	1.4	(0.0 - 3.1)		
\$50,000-\$74,999	*	*	*	*	5,766	13,300	0.7	(0.2 - 1.2)		
≥ \$75,000	*	*	*	*	8,561	35,300	0.8	(0.3 - 1.3)		

 $<sup>^{\</sup>rm l}$  Defined as having had flu vaccine sprayed in nose within the past 12 months

<sup>&</sup>lt;sup>2</sup> Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults who have had flu vaccine sprayed in nose within the past 12 months (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 19b. Flu vaccine that was sprayed in nose within the past 12 months<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland C	ounty	N	ew York	State
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>
Total	*	*	*	38,106	1.7	(0.6 - 2.7)
Gender						
Male	*	*	*	14,586	2.0	(0.1 - 3.9)
Female	*	*	*	23,520	1.4	(0.5 - 2.3)
Race/Ethnicity						
White Non-Hispanic	*	*	*	35,796	0.7	(0.4 - 1.0)
Black Non-Hispanic	*	*	*	740	1.9	(0.0 - 4.4)
Hispanic	*	*	*	860	5.9	(0.0 - 11.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	15,679	3.1	(0.4 - 5.8)
Some College	*	*	*	9,897	1.2	(0.4 - 2.0)
≥ College Degree	*	*	*	12,449	0.6	(0.3 - 0.8)
Income						
≤ \$24,999	*	*	*	9,636	3.8	(0.0 - 7.7)
\$25,000-\$49,999	*	*	*	9,648	1.5	(0.0 - 3.4)
\$50,000-\$74,999	*	*	*	5,766	0.7	(0.1 - 1.2)
≥ \$75,000	*	*	*	8,561	0.8	(0.3 - 1.3)

 $<sup>^{\</sup>rm I}$  Defined as having had flu vaccine sprayed in nose within the past 12 months

<sup>&</sup>lt;sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 20a. Flu shot or flu vaccine sprayed in nose within the past 12 months<sup>1</sup> in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	<b>e</b>
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	655	88,100	40.9	(36.0 - 45.8)	38,059	6,302,300	42.7	(40.2 - 45.1)
Gender								
Male	242	41,500	39.7	(32.0 - 47.4)	14,568	2,822,700	40.0	(36.1 - 43.9)
Female	413	46,600	42.1	(35.9 - 48.2)	23,491	3,479,500	45.1	(42.0 - 48.2)
Age (years)								
18-49	229	28,600	25.4	(18.2 - 32.6)	12,501	2,726,400	31.6	(27.8 - 35.4)
50-64	209	29,200	45.7	(37.3 - 54.1)	12,964	1,664,200	46.1	(42.7 - 49.5)
≥ 65	217	30,300	78.0	(71.7 - 84.3)	12,594	1,911,700	75.4	(72.2 - 78.6)
Race/Ethnicity								
White Non-Hispanic	548	64,500	43.3	(38.2 - 48.5)	35,750	4,048,700	43.6	(41.7 - 45.5)
Black Non-Hispanic	*	*	*	*	738	910,000	42.2	(33.2 - 51.2)
Hispanic	*	*	*	*	860	903,300	40.6	(31.3 - 49.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	196	24,500	38.1	(29.3 - 46.9)	15,651	2,480,100	44.7	(40.2 - 49.3)
Some College	*	*	*	*	9,895	1,387,300	39.2	(34.3 - 44.1)
≥ College Degree	339	45,700	42.1	(35.3 - 48.9)	12,431	2,418,300	43.2	(39.8 - 46.7)
Income								
≤ \$24,999	*	*	*	*	9,626	1,675,100	48.2	(41.8 - 54.6)
\$25,000-\$49,999	*	*	*	*	9,643	1,208,300	41.7	(35.8 - 47.6)
\$50,000-\$74,999	*	*	*	*	5,761	795,600	40.4	(34.5 - 46.3)
≥ \$75,000	270	43,600	44.0	(36.5 - 51.4)	8,548	1,791,200	39.9	(36.6 - 43.2)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having had a flu shot or flu vaccine sprayed in nose within the past 12 months

<sup>&</sup>lt;sup>2</sup> Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults who have had a flu shot or flu vaccine sprayed in nose within the past 12 months (rounded to the nearest hundred)

4 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a

numerator less than 10)

Table 20b. Flu shot or flu vaccine sprayed in nose within the past 12 months<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	ew York	State
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	655	38.5	(33.5 - 43.5)	38,059	42.0	(39.6 - 44.5)
Gender						
Male	242	38.4	(30.6 - 46.2)	14,568	40.4	(36.6 - 44.2)
Female	413	38.6	(32.1 - 45.1)	23,491	43.5	(40.3 - 46.7)
Race/Ethnicity						
White Non-Hispanic	548	40.3	(35.0 - 45.5)	35,750	42.6	(40.7 - 44.6)
Black Non-Hispanic	*	*	*	738	42.4	(33.4 - 51.4)
Hispanic	*	*	*	860	40.0	(30.9 - 49.1)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	196	35.0	(26.5 - 43.6)	15,651	44.1	(39.6 - 48.6)
Some College	*	*	*	9,895	38.1	(33.2 - 43.0)
≥ College Degree	339	40.0	(32.9 - 47.1)	12,431	42.8	(39.4 - 46.3)
Income						
≤ \$24,999	*	*	*	9,626	48.0	(41.6 - 54.3)
\$25,000-\$49,999	*	*	*	9,643	40.5	(34.8 - 46.3)
\$50,000-\$74,999	*	*	*	5,761	39.6	(33.6 - 45.6)
≥ \$75,000	270	41.8	(34.0 - 49.5)	8,548	39.4	(36.1 - 42.7)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having had a flu shot or flu vaccine sprayed in nose within the past 12 months

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 21a. Flu shot among adults aged 65 and older in Rockland County and New York State

		Rockla	nd Coun	ty		New Y	ork State	e
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	217	30,300	78.0	(71.7 - 84.3)	12,602	1,907,700	75.2	(72.0 - 78.4)
Gender								
Male	86	14,300	85.3	(76.9 - 93.6)	4,417	772,900	74.6	(69.3 - 79.8)
Female	131	16,000	72.6	(63.6 - 81.5)	8,185	1,134,800	75.6	(71.6 - 79.6)
Age (years)								
≥ 65	217	30,300	78.0	(71.7 - 84.3)	12,602	1,907,700	75.2	(72.0 - 78.4)
Race/Ethnicity								
White Non-Hispanic	201	27,100	80.1	(73.9 - 86.3)	12,137	1,487,100	76.0	(73.5 - 78.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	87	12,300	80.3	(71.6 - 89.1)	6,518	930,800	74.0	(68.9 - 79.2)
Some College	*	*	*	*	2,790	317,800	69.7	(62.7 - 76.7)
≥ College Degree	95	13,700	79.3	(69.4 - 89.2)	3,261	654,000	80.5	(76.4 - 84.6)
Income								
≤ \$24,999	*	*	*	*	4,665	545,300	72.4	(65.6 - 79.3)
\$25,000-\$49,999	*	*	*	*	3,422	430,600	72.8	(67.0 - 78.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	53	9,300	88.6	(79.1 - 98.1)	1,198	301,000	77.5	(70.9 - 84.2)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having had a flu shot during the past 12 months among adults aged 65 and older

<sup>&</sup>lt;sup>2</sup> Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults aged 65 and older who have had a flu shot within the past 12 months (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interva

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 21b. Flu shot among adults aged 65 and older<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	Ro	ockland (	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total 2	217	78.6	(72.3 - 84.8)	12,602	75.0	(71.5 - 78.5)		
Gender								
Male	86	85.1	(76.7 - 93.5)	4,417	74.4	(68.7 - 80.1)		
Female 1	l31	72.5	(63.5 - 81.5)	8,185	75.5	(71.3 - 79.8)		
Race/Ethnicity								
White Non-Hispanic	201	80.8	(74.7 - 86.9)	12,137	75.9	(73.0 - 78.8)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	87	80.9	(72.3 - 89.4)	6,518	73.3	(67.5 - 79.1)		
Some College	*	*	*	2,790	70.0	(62.7 - 77.2)		
≥ College Degree	95	79.6	(69.7 - 89.6)	3,261	80.7	(76.2 - 85.1)		
Income								
≤ \$24,999	*	*	*	4,665	72.6	(65.1 - 80.1)		
\$25,000-\$49,999	*	*	*	3,422	71.4	(65.0 - 77.9)		
\$50,000-\$74,999	*	*	*	*	*	*		
≥ \$75,000	53	89.4	(80.5 - 98.3)	1,198	78.0	(70.8 - 85.1)		

 $<sup>^{\</sup>rm 1}$  Defined as having had a flu shot during the past 12 months among adults aged 65 and older

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 22a. Pneumonia shot or pneumococcal vaccine<sup>1</sup> in Rockland County and New York State

		Rockla	and Coun	ty		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	594	39,400	20.8	(17.2 - 24.4)	35,303	3,456,800	26.1	(23.9 - 28.3)
Gender								
Male	210	18,400	20.5	(14.6 - 26.3)	13,052	1,542,200	25.2	(21.5 - 28.8)
Female	384	21,000	21.1	(16.8 - 25.4)	22,251	1,914,600	26.9	(24.2 - 29.6)
Age (years)								
18-49	199	6,300	6.6	(2.3 - 10.8)	10,943	1,097,000	14.6	(11.3 - 17.9)
50-64	191	7,400	12.8	(7.9 - 17.7)	12,179	758,000	23.0	(19.8 - 26.2)
≥ 65	204	25,700	70.4	(63.4 - 77.4)	12,181	1,601,700	65.7	(61.7 - 69.7)
Race/Ethnicity								
White Non-Hispanic	504	33,400	24.9	(20.6 - 29.2)	33,225	2,358,700	28.7	(26.9 - 30.5)
Black Non-Hispanic	*	*	*	*	685	584,400	28.4	(19.5 - 37.3)
Hispanic	*	*	*	*	769	408,000	19.5	(12.6 - 26.4)
Other Non-Hispanic	*	*	*	*	624	105,600	12.2	(2.3 - 22.2)
Education								
≤ High School	184	12,000	20.2	(14.2 - 26.2)	14,700	1,535,300	30.4	(26.4 - 34.3)
Some College	108	7,100	19.0	(11.1 - 26.8)	9,151	836,200	27.2	(21.9 - 32.5)
≥ College Degree	299	20,300	22.2	(16.8 - 27.6)	11,375	1,070,700	21.4	(18.6 - 24.1)
Income								
≤ \$24,999	87	7,000	24.4	(14.7 - 34.2)	9,082	1,050,900	33.1	(27.6 - 38.6)
\$25,000-\$49,999	100	7,300	24.9	(15.6 - 34.3)	9,033	797,500	30.3	(24.1 - 36.6)
\$50,000-\$74,999	*	*	*	*	5,320	378,800	21.8	(17.4 - 26.1)
≥ \$75,000	241	15,000	17.2	(11.7 - 22.6)	7,773	648,700	16.3	(14.1 - 18.5)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having ever had a pneumonia shot or pneumococcal vaccine

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults who have ever had a pneumonia shot or pneumococcal vaccine (rounded to the nearest hundred)

<sup>4 95%</sup> confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 22b. Pneumonia shot or pneumococcal vaccine<sup>1</sup> in Rockland County and New York
State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	594	19.6	(16.0 - 23.2)	35,303	25.4	(23.2 - 27.6)		
Gender								
Male	210	20.8	(14.7 - 26.8)	13,052	25.8	(22.3 - 29.4)		
Female	384	18.5	(14.4 - 22.6)	22,251	25.1	(22.4 - 27.8)		
Race/Ethnicity								
White Non-Hispanic	504	23.6	(19.2 - 28.0)	33,225	27.7	(25.9 - 29.5)		
Black Non-Hispanic	*	*	*	685	27.9	(19.1 - 36.7)		
Hispanic	*	*	*	769	19.6	(12.8 - 26.5)		
Other Non-Hispanic	*	*	*	624	11.3	(2.0 - 20.7)		
Education								
≤ High School	184	18.9	(13.0 - 24.8)	14,700	29.5	(25.6 - 33.5)		
Some College	108	17.2	(9.4 - 24.9)	9,151	26.2	(21.0 - 31.3)		
≥ College Degree	299	21.2	(15.7 - 26.7)	11,375	21.0	(18.3 - 23.8)		
Income								
≤ \$24,999	87	21.6	(12.6 - 30.5)	9,082	32.3	(26.9 - 37.8)		
\$25,000-\$49,999	100	23.4	(14.3 - 32.6)	9,033	29.0	(23.0 - 35.0)		
\$50,000-\$74,999	*	*	*	5,320	21.2	(16.8 - 25.7)		
≥ \$75,000	241	16.8	(11.1 - 22.4)	7,773	16.2	(13.9 - 18.5)		

 $<sup>^{\</sup>rm 1}$  Defined as having ever had a pneumonia shot or pneumococcal vaccine

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 23a. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older in Rockland County and New York State

		Rockla	nd Coun	ty		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	204	25,700	70.4	(63.4 - 77.4)	12,181	1,601,700	65.7	(61.7 - 69.7)
Gender								
Male	*	*	*	*	4,180	606,400	62.1	(55.5 - 68.7)
Female	127	15,200	70.3	(61.2 - 79.4)	8,001	995,300	68.0	(63.1 - 73.0)
Age (years)								
≥ 65	204	25,700	70.4	(63.4 - 77.4)	12,181	1,601,700	65.7	(61.7 - 69.7)
Race/Ethnicity								
White Non-Hispanic	190	22,800	71.5	(64.4 - 78.6)	11,735	1,281,300	68.2	(64.9 - 71.5)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	6,345	806,200	65.6	(59.5 - 71.7)
Some College	*	*	*	*	2,694	289,500	66.8	(60.1 - 73.4)
≥ College Degree	*	*	*	*	3,111	501,100	65.6	(58.5 - 72.7)
Income								
≤ \$24,999	*	*	*	*	4,533	497,100	67.7	(59.9 - 75.5)
\$25,000-\$49,999	*	*	*	*	3,324	344,300	61.5	(53.3 - 69.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	1,138	236,300	64.5	(55.2 - 73.8)

 $<sup>^{</sup>m 1}$  Defined as having ever had a pneumonia shot or pneumococcal vaccine among adults aged 65 and older

 $<sup>^{2}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults aged 65 and older who have ever had a pneumonia shot or pneumococcal vaccine (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 23b. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	Re	ockland	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	204	70.3	(63.2 - 77.4)	12,181	64.7	(60.4 - 69.1)		
Gender								
Male	*	*	*	4,180	61.5	(54.3 - 68.6)		
Female	127	70.3	(61.1 - 79.4)	8,001	67.6	(62.4 - 72.9)		
Race/Ethnicity								
White Non-Hispanic	190	71.4	(64.2 - 78.6)	11,735	67.1	(63.2 - 70.9)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	6,345	64.2	(57.4 - 71.0)		
Some College	*	*	*	2,694	65.8	(58.4 - 73.3)		
≥ College Degree	*	*	*	3,111	65.3	(57.7 - 72.9)		
Income								
≤ \$24,999	*	*	*	4,533	66.0	(57.0 - 74.9)		
\$25,000-\$49,999	*	*	*	3,324	60.3	(51.9 - 68.6)		
\$50,000-\$74,999	*	*	*	*	*	*		
≥ \$75,000	*	*	*	*	*	*		

 $<sup>^{\</sup>mathrm{1}}$  Defined as having ever had a pneumonia shot or pneumococcal vaccine among adults aged 65 and older

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 24a. Ever had a mammogram among women aged 40 and older<sup>1</sup> in Rockland County and New York State

		Rockla	nd Coun	ty		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	328	69,700	95.8	(93.4 - 98.1)	19,491	4,328,800	91.8	(89.7 - 94.0)
Gender								
Male	-	-	-	-	-	-	-	-
Female	328	69,700	95.8	(93.4 - 98.1)	19,491	4,328,800	91.8	(89.7 - 94.0)
Age (years)								
40-49	80	18,500	91.7	(84.9 - 98.4)	3,851	1,287,500	85.0	(79.7 - 90.3)
50-64	119	30,300	97.9	(95.2 - 100)	7,452	1,641,700	96.0	(94.1 - 97.9)
≥ 65	129	20,900	96.6	(93.8 - 99.4)	8,188	1,399,600	94.1	(90.5 - 97.6)
Race/Ethnicity								
White Non-Hispanic	279	48,200	94.8	(91.7 - 97.9)	18,548	2,976,100	93.9	(92.7 - 95.1)
Black Non-Hispanic	*	*	*	*	342	746,600	93.8	(87.3 - 100)
Hispanic	*	*	*	*	333	447,600	91.8	(84.6 - 99.0)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	105	23,100	95.9	(92.6 - 99.2)	8,266	1,698,700	91.0	(87.4 - 94.5)
Some College	67	16,600	94.8	(88.6 - 100)	5,322	1,055,500	93.6	(90.2 - 97.0)
≥ College Degree	155	29,500	96.2	(92.7 - 99.6)	5,872	1,560,700	91.6	(87.5 - 95.6)
Income								
≤ \$24,999	*	*	*	*	5,691	1,023,400	93.5	(89.7 - 97.4)
\$25,000-\$49,999	61	14,200	93.0	(85.6 - 100)	4,857	860,300	89.5	(83.6 - 95.5)
\$50,000-\$74,999	*	*	*	*	2,648	547,100	94.6	(92.2 - 96.9)
≥ \$75,000	119	28,200	97.5	(94.6 - 100)	3,582	1,185,100	92.2	(88.2 - 96.1)

 $<sup>^{\</sup>rm 1}$  Defined as having ever had a mammogram among women aged 40 and older

<sup>&</sup>lt;sup>2</sup> Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of women aged 40 and older who have ever had a mammogram (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 24b. Ever had a mammogram among women aged 40 and older<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	328	95.3	(92.4 - 98.1)	19,491	91.1	(88.7 - 93.6)		
Gender								
Male	-	-	-	-	-	-		
Female	328	95.3	(92.4 - 98.1)	19,491	91.1	(88.7 - 93.6)		
Race/Ethnicity								
White Non-Hispanic	279	94.1	(90.3 - 97.9)	18,548	93.3	(92.0 - 94.7)		
Black Non-Hispanic	*	*	*	342	93.1	(85.7 - 100)		
Hispanic	*	*	*	333	91.0	(82.8 - 99.2)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	105	95.4	(91.5 - 99.3)	8,266	90.2	(86.1 - 94.2)		
Some College	67	93.4	(85.1 - 100)	5,322	92.9	(88.9 - 96.9)		
≥ College Degree	155	96.2	(92.8 - 99.6)	5,872	90.9	(86.6 - 95.3)		
Income								
≤ \$24,999	*	*	*	5,691	93.0	(88.5 - 97.4)		
\$25,000-\$49,999	*	*	*	4,857	88.8	(82.5 - 95.1)		
\$50,000-\$74,999	*	*	*	2,648	94.3	(91.7 - 96.8)		
≥ \$75,000	119	97.5	(94.5 - 100)	3,582	91.2	(86.8 - 95.6)		

 $<sup>^{\</sup>rm I}$  Defined as having ever had a mammogram among women aged 40 and older

 $<sup>^{2}</sup>$  Prevalence estimates are age-adjusted to the 2000 US Standard population  $\,$ 

 $<sup>^{\</sup>rm 3}$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 25a. Had mammogram within the past 2 years among women aged 40 and older in Rockland County and New York State

		Rockla	nd Coun	ty		New Y	ork State	9
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	325	58,900	81.6	(76.0 - 87.2)	19,376	3,760,000	80.3	(77.8 - 82.9)
Gender								
Male	-	-	-	-	-	-	-	-
Female	325	58,900	81.6	(76.0 - 87.2)	19,376	3,760,000	80.3	(77.8 - 82.9)
Age (years)								
40-49	*	*	*	*	3,842	1,122,700	74.4	(68.5 - 80.2)
50-64	118	25,400	82.8	(74.4 - 91.2)	7,428	1,487,900	87.2	(84.5 - 89.8)
≥ 65	128	17,900	83.2	(76.1 - 90.3)	8,106	1,149,400	78.5	(74.2 - 82.9)
Race/Ethnicity								
White Non-Hispanic	276	41,100	82.0	(76.8 - 87.2)	18,438	2,531,500	80.2	(78.3 - 82.2)
Black Non-Hispanic	*	*	*	*	339	667,100	85.6	(77.5 - 93.8)
Hispanic	*	*	*	*	332	419,700	86.1	(77.5 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	8,204	1,486,900	80.5	(76.4 - 84.6)
Some College	*	*	*	*	5,297	929,400	83.0	(78.8 - 87.1)
≥ College Degree	154	27,100	89.0	(83.6 - 94.5)	5,845	1,334,800	78.6	(73.9 - 83.2)
Income								
≤ \$24,999	*	*	*	*	5,650	882,300	80.8	(75.9 - 85.7)
\$25,000-\$49,999	*	*	*	*	4,846	732,300	76.3	(69.7 - 82.9)
\$50,000-\$74,999	*	*	*	*	2,641	494,200	85.6	(81.9 - 89.4)
≥ \$75,000	*	*	*	*	3,572	1,025,900	80.0	(75.2 - 84.8)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having had a mammogram within the past 2 years among women aged 40 and older

<sup>&</sup>lt;sup>2</sup> Sample size from survey

 $<sup>^3</sup>$  Estimated number of women aged 40 and older who have had a mammogram within the past 2 years (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 25b. Had mammogram within the past 2 years among women aged 40 and older in Rockland County and New York State - Age-Adjusted 2

	Ro	ockland (	County	N	ew York	State
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	325	81.4	(75.5 - 87.3)	19,376	79.7	(76.9 - 82.5)
Gender						
Male	-	-	-	-	-	-
Female	325	81.4	(75.5 - 87.3)	19,376	79.7	(76.9 - 82.5)
Race/Ethnicity						
White Non-Hispanic	276	81.6	(76.0 - 87.1)	18,438	79.7	(77.6 - 81.8)
Black Non-Hispanic	*	*	*	339	85.3	(76.5 - 94.2)
Hispanic	*	*	*	332	85.2	(75.6 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	8,204	80.1	(75.6 - 84.7)
Some College	*	*	*	5,297	82.3	(77.7 - 86.9)
≥ College Degree	154	89.3	(83.7 - 94.8)	5,845	77.8	(72.9 - 82.8)
Income						
≤ \$24,999	*	*	*	5,650	80.8	(75.4 - 86.2)
\$25,000-\$49,999	*	*	*	4,846	75.7	(68.7 - 82.6)
\$50,000-\$74,999	*	*	*	2,641	85.5	(81.6 - 89.4)
≥ \$75,000	*	*	*	3,572	79.0	(73.8 - 84.2)

 $<sup>^{1}</sup>$  Defined as having had a mammogram within the past 2 years among women aged 40 and older

 $<sup>^{2}</sup>$  Prevalence estimates are age-adjusted to the 2000 US Standard population  $\,$ 

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 26a. Had mammogram within the past 2 years among women aged 50 and older 1 in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	9
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	246	43,300	83.0	(77.2 - 88.7)	15,534	2,637,300	83.2	(80.7 - 85.7)
Gender								
Male	-	-	-	-	-	-	-	-
Female	246	43,300	83.0	(77.2 - 88.7)	15,534	2,637,300	83.2	(80.7 - 85.7)
Age (years)								
50-64	118	25,400	82.8	(74.4 - 91.2)	7,428	1,487,900	87.2	(84.5 - 89.8)
≥ 65	128	17,900	83.2	(76.1 - 90.3)	8,106	1,149,400	78.5	(74.2 - 82.9)
Race/Ethnicity								
White Non-Hispanic	210	30,000	82.3	(76.5 - 88.1)	14,887	1,907,000	82.3	(80.3 - 84.4)
Black Non-Hispanic	*	*	*	*	249	385,600	89.9	(83.2 - 96.7)
Hispanic	*	*	*	*	220	286,700	93.9	(89.1 - 98.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	94	16,500	80.0	(70.0 - 90.0)	7,026	1,094,500	83.3	(80.0 - 86.6)
Some College	*	*	*	*	4,124	646,700	87.3	(84.2 - 90.3)
≥ College Degree	103	17,300	88.3	(81.5 - 95.2)	4,356	890,500	80.6	(75.1 - 86.1)
Income								
≤ \$24,999	*	*	*	*	4,991	649,100	81.0	(76.8 - 85.2)
\$25,000-\$49,999	*	*	*	*	3,913	533,300	78.6	(70.9 - 86.2)
\$50,000-\$74,999	*	*	*	*	1,943	359,300	89.6	(86.3 - 93.0)
≥ \$75,000	74	15,100	86.9	(77.0 - 96.8)	2,303	601,300	84.6	(79.3 - 89.9)

<sup>&</sup>lt;sup>1</sup> Defined as having had a mammogram within the past 2 years among women aged 50 and older

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of women aged 50 and older who have had a mammogram within the past 2 years (rounded to the nearest

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a

Table 26b. Had mammogram within the past 2 years among women aged 50 and older in Rockland County and New York State - Age-Adjusted 2

	R	ockland	County	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	246	82.9	(77.1 - 88.7)	15,534	83.1	(80.3 - 85.8)		
Gender								
Male	-	-	-	-	-	-		
Female	246	82.9	(77.1 - 88.7)	15,534	83.1	(80.3 - 85.8)		
Race/Ethnicity								
White Non-Hispanic	210	82.4	(76.7 - 88.2)	14,887	82.1	(79.9 - 84.3)		
Black Non-Hispanic	*	*	*	249	90.4	(83.6 - 97.2)		
Hispanic	*	*	*	220	94.1	(89.3 - 99.0)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	94	80.4	(70.6 - 90.1)	7,026	83.6	(80.1 - 87.0)		
Some College	*	*	*	4,124	87.3	(84.1 - 90.5)		
≥ College Degree	103	88.7	(81.9 - 95.4)	4,356	79.9	(73.9 - 86.0)		
Income								
≤ \$24,999	*	*	*	4,991	81.3	(76.9 - 85.6)		
\$25,000-\$49,999	*	*	*	3,913	78.1	(69.9 - 86.4)		
\$50,000-\$74,999	*	*	*	1,943	90.2	(87.1 - 93.4)		
≥ \$75,000	*	*	*	2,303	83.9	(77.9 - 89.9)		

 $<sup>^{\</sup>rm I}$  Defined as having had a mammogram within the past 2 years among women aged 50 and older

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{\</sup>rm 3}$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 27a. Ever had a Pap test among women in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	408	103,000	93.4	(89.9 - 96.9)	23,381	7,045,300	92.1	(90.0 - 94.3)
Gender								
Male	-	-	-	-	-	-	-	-
Female	408	103,000	93.4	(89.9 - 96.9)	23,381	7,045,300	92.1	(90.0 - 94.3)
Age (years)								
18-34	*	*	*	*	2,377	1,896,600	85.2	(79.3 - 91.0)
35-44	64	19,000	98.7	(96.3 - 100)	3,023	1,301,500	92.6	(87.7 - 97.5)
45-54	86	21,300	96.4	(92.2 - 100)	4,657	1,420,300	98.3	(97.0 - 99.6)
55-64	81	18,900	98.7	(96.0 - 100)	5,187	1,071,200	97.2	(95.0 - 99.3)
≥ 65	129	20,400	94.2	(90.2 - 98.3)	8,137	1,355,700	92.3	(88.5 - 96.1)
Race/Ethnicity								
White Non-Hispanic	335	67,600	93.6	(89.6 - 97.6)	21,976	4,439,700	94.2	(92.1 - 96.3)
Black Non-Hispanic	*	*	*	*	481	1,231,600	91.2	(84.0 - 98.4)
Hispanic	*	*	*	*	533	984,500	90.1	(85.3 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	136	35,000	90.7	(84.0 - 97.5)	9,526	2,540,500	88.3	(83.8 - 92.7)
Some College	79	23,100	93.8	(86.9 - 100)	6,462	1,782,200	93.5	(90.4 - 96.6)
≥ College Degree	191	44,300	95.2	(90.4 - 100)	7,345	2,643,200	95.0	(92.2 - 97.9)
Income								
≤ \$24,999	56	10,500	87.9	(78.1 - 97.7)	6,526	1,717,200	93.7	(90.9 - 96.6)
\$25,000-\$49,999	72	22,000	99.6	(98.7 - 100)	5,787	1,378,400	91.0	(85.6 - 96.4)
\$50,000-\$74,999	*	*	*	*	3,310	933,900	96.0	(91.8 - 100)
≥ \$75,000	146	41,500	98.6	(96.7 - 100)	4,570	1,943,300	95.4	(92.6 - 98.2)

 $<sup>^{1}</sup>$  Defined as having ever had a Pap test among women

 $<sup>^{2} \, \</sup>mathrm{Sample} \, \mathrm{size} \, \mathrm{from} \, \mathrm{survey}$ 

 $<sup>^{3}</sup>$  Estimated number of women who have ever had a Pap test (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 27b. Ever had a Pap test among women<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	408	92.8	(88.8 - 96.8)	23,381	91.8	(89.6 - 94.0)		
Gender								
Male	-	-	-	-	-	-		
Female	408	92.8	(88.8 - 96.8)	23,381	91.8	(89.6 - 94.0)		
Race/Ethnicity								
White Non-Hispanic	335	92.7	(87.8 - 97.5)	21,976	94.0	(91.8 - 96.2)		
Black Non-Hispanic	*	*	*	481	90.8	(83.5 - 98.1)		
Hispanic	*	*	*	533	89.7	(84.6 - 94.7)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	136	89.4	(81.2 - 97.6)	9,526	87.8	(83.2 - 92.4)		
Some College	79	93.2	(85.3 - 100)	6,462	93.1	(89.7 - 96.4)		
≥ College Degree	191	95.2	(90.1 - 100)	7,345	94.9	(92.0 - 97.7)		
Income								
≤ \$24,999	*	*	*	6,526	93.3	(90.1 - 96.5)		
\$25,000-\$49,999	72	99.6	(98.8 - 100)	5,787	91.1	(85.7 - 96.5)		
\$50,000-\$74,999	*	*	*	3,310	96.0	(91.9 - 100)		
≥ \$75,000	146	98.6	(96.7 - 100)	4,570	95.0	(91.9 - 98.1)		

<sup>&</sup>lt;sup>1</sup> Defined as having ever had a Pap test among women

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{\</sup>rm 3}$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 28a. Had a Pap test within the past 3 years among women<sup>1</sup> in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	403	89,700	82.4	(77.7 - 87.1)	23,109	6,205,600	82.2	(79.8 - 84.6)
Gender								
Male	-	-	-	-	-	-	-	-
Female	403	89,700	82.4	(77.7 - 87.1)	23,109	6,205,600	82.2	(79.8 - 84.6)
Age (years)								
18-34	*	*	*	*	2,371	1,818,400	81.8	(75.5 - 88.1)
35-44	64	17,800	92.8	(86.5 - 99.2)	3,012	1,220,300	86.9	(81.8 - 92.0)
45-54	85	19,800	91.4	(85.1 - 97.6)	4,640	1,311,500	92.0	(89.8 - 94.3)
55-64	*	*	*	*	5,152	924,300	84.5	(81.0 - 88.0)
≥ 65	127	13,600	64.1	(54.7 - 73.6)	7,934	931,100	66.4	(61.8 - 71.0)
Race/Ethnicity								
White Non-Hispanic	331	58,300	81.8	(76.7 - 86.9)	21,720	3,807,100	81.5	(79.2 - 83.8)
Black Non-Hispanic	*	*	*	*	473	1,142,800	86.7	(79.1 - 94.3)
Hispanic	*	*	*	*	527	894,900	83.6	(76.3 - 91.0)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	134	29,700	78.2	(69.7 - 86.7)	9,384	2,132,600	76.0	(71.1 - 81.0)
Some College	78	20,100	82.3	(72.8 - 91.8)	6,401	1,594,300	84.0	(80.4 - 87.5)
≥ College Degree	189	39,400	85.8	(78.7 - 92.9)	7,281	2,406,000	86.9	(83.7 - 90.1)
Income								
≤ \$24,999	*	*	*	*	6,441	1,453,900	82.0	(77.9 - 86.0)
\$25,000-\$49,999	71	18,000	82.2	(72.4 - 92.0)	5,740	1,182,000	78.3	(72.6 - 83.9)
\$50,000-\$74,999	*	*	*	*	3,296	825,000	85.0	(78.0 - 91.9)
≥ \$75,000	146	38,600	91.7	(85.6 - 97.8)	4,555	1,810,200	89.0	(85.7 - 92.2)

<sup>&</sup>lt;sup>1</sup> Defined as having had a Pap test within the past 3 years among women

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of women who have had a Pap test within the past 3 years (rounded to the nearest hundred)

<sup>4 95%</sup> confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 28b. Had a Pap test within the past 3 years among women<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	403	83.4	(78.6 - 88.2)	23,109	82.7	(80.2 - 85.1)		
Gender								
Male	-	-	-	-	-	-		
Female	403	83.4	(78.6 - 88.2)	23,109	82.7	(80.2 - 85.1)		
Race/Ethnicity								
White Non-Hispanic	331	82.3	(76.7 - 87.8)	21,720	82.3	(80.0 - 84.6)		
Black Non-Hispanic	*	*	*	473	87.0	(79.3 - 94.6)		
Hispanic	*	*	*	527	83.4	(75.9 - 90.9)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	134	78.2	(68.8 - 87.5)	9,384	76.4	(71.3 - 81.5)		
Some College	78	83.7	(74.1 - 93.4)	6,401	84.2	(80.5 - 88.0)		
≥ College Degree	189	87.3	(80.7 - 94.0)	7,281	87.5	(84.3 - 90.6)		
Income								
≤ \$24,999	*	*	*	6,441	82.7	(78.5 - 86.9)		
\$25,000-\$49,999	71	84.7	(76.0 - 93.4)	5,740	79.2	(73.5 - 84.8)		
\$50,000-\$74,999	*	*	*	3,296	85.7	(78.7 - 92.8)		
≥ \$75,000	146	92.8	(87.7 - 97.9)	4,555	88.9	(85.5 - 92.3)		

 $<sup>^{1}</sup>$  Defined as having had a Pap test within the past 3 years among women

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{\</sup>rm 3}$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 29a. Ever had a digital rectal exam among men aged 40 and older in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	•
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	198	53,100	80.6	(72.3 - 88.8)	12,033	3,176,500	73.8	(69.7 - 77.8)
Gender								
Male	198	53,100	80.6	(72.3 - 88.8)	12,033	3,176,500	73.8	(69.7 - 77.8)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	3,098	922,800	66.8	(59.9 - 73.8)
55-64	52	16,000	93.3	(87.1 - 99.4)	3,504	865,600	87.1	(82.5 - 91.7)
≥ 65	85	15,900	96.7	(92.9 - 100)	4,348	953,900	92.9	(91.0 - 94.9)
Race/Ethnicity								
White Non-Hispanic	179	46,700	87.1	(80.5 - 93.7)	11,411	2,452,100	80.3	(77.7 - 82.9)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	5,013	931,600	64.1	(57.3 - 71.0)
Some College	*	*	*	*	2,695	674,200	75.5	(67.8 - 83.2)
≥ College Degree	*	*	*	*	4,302	1,568,000	80.5	(74.0 - 87.0)
Income								
≤ \$24,999	*	*	*	*	2,573	573,300	65.9	(56.0 - 75.9)
\$25,000-\$49,999	*	*	*	*	3,168	530,100	74.6	(67.2 - 82.1)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	99	33,700	91.2	(84.8 - 97.6)	3,309	1,351,100	78.0	(72.6 - 83.5)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having ever had a digital rectal exam among men aged 40 and older

 $<sup>^2 \, {\</sup>rm Sample \; size \; from \; survey}$ 

 $<sup>^3</sup>$  Estimated number of men aged 40 and older who have ever had a digital rectal exam (rounded to the nearest hundred)

<sup>4 95%</sup> confidence interval \* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 29b. Ever had a digital rectal exam among men aged 40 and older in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	ew York	State
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	198	79.0	(69.8 - 88.2)	12,033	73.7	(69.6 - 77.8)
Gender						
Male	198	79.0	(69.8 - 88.2)	12,033	73.7	(69.6 - 77.8)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	179	85.7	(78.3 - 93.1)	11,411	79.9	(77.2 - 82.5)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	5,013	64.7	(57.8 - 71.7)
Some College	*	*	*	2,695	75.0	(67.1 - 82.9)
≥ College Degree	*	*	*	4,302	80.3	(73.8 - 86.8)
Income						
≤ \$24,999	*	*	*	2,573	66.9	(56.9 - 76.9)
\$25,000-\$49,999	*	*	*	3,168	74.8	(67.3 - 82.2)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	99	89.5	(81.6 - 97.4)	3,309	77.5	(72.0 - 83.0)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having ever had a digital rectal exam among men aged 40 and older

 $<sup>^{2}</sup>$  Prevalence estimates are age-adjusted to the 2000 US Standard population  $\,$ 

 $<sup>^{\</sup>rm 3}$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval \* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 30a. Had digital rectal exam within the past 2 years among men aged 40 and older in Rockland County and New York State

		Rockla	ınd Coun	ty		New York State			
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	
Total	196	37,500	57.4	(48.3 - 66.6)	11,923	2,380,300	55.8	(51.8 - 59.8)	
Gender									
Male	196	37,500	57.4	(48.3 - 66.6)	11,923	2,380,300	55.8	(51.8 - 59.8)	
Female	-	-	-	-	-	-	-	-	
Age (years)									
40-44	*	*	*	*	1,078	246,800	27.5	(18.5 - 36.5)	
45-54	*	*	*	*	3,074	650,000	47.6	(40.8 - 54.4)	
55-64	*	*	*	*	3,467	673,500	68.1	(62.4 - 73.8)	
≥ 65	83	12,600	79.8	(70.5 - 89.1)	4,304	809,900	79.7	(75.0 - 84.3)	
Race/Ethnicity									
White Non-Hispanic	177	34,300	64.7	(56.1 - 73.2)	11,305	1,875,600	62.2	(59.2 - 65.2)	
Black Non-Hispanic	*	*	*	*	*	*	*	*	
Hispanic	*	*	*	*	*	*	*	*	
Other Non-Hispanic	*	*	*	*	*	*	*	*	
Education									
≤ High School	*	*	*	*	4,963	718,700	49.6	(43.2 - 56.1)	
Some College	*	*	*	*	2,665	511,500	58.5	(50.8 - 66.3)	
≥ College Degree	*	*	*	*	4,277	1,148,400	59.4	(52.8 - 66.0)	
Income									
≤ \$24,999	*	*	*	*	2,553	388,000	45.0	(35.0 - 55.0)	
\$25,000-\$49,999	*	*	*	*	3,139	422,400	60.5	(53.0 - 68.0)	
\$50,000-\$74,999	*	*	*	*	*	*	*	*	
≥ \$75,000	*	*	*	*	3,295	978,300	56.9	(51.1 - 62.7)	

 $<sup>^{\</sup>mathrm{1}}$  Defined as having had a digital rectal exam within the past 2 years among men aged 40 and older

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of men aged 40 and older who have had a digital rectal exam within the past 2 years (rounded to the nearest hundred)
4 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 30b. Had digital rectal exam within the past 2 years among men aged 40 and older in Rockland County and New York State - Age-Adjusted 2

	R	ockland	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	196	56.8	(47.1 - 66.4)	11,923	55.9	(51.7 - 60.0)		
Gender								
Male	196	56.8	(47.1 - 66.4)	11,923	55.9	(51.7 - 60.0)		
Female	-	-	-	-	-	-		
Race/Ethnicity								
White Non-Hispanic	177	64.7	(55.8 - 73.7)	11,305	62.0	(58.9 - 65.1)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	4,963	50.5	(43.7 - 57.3)		
Some College	*	*	*	2,665	58.3	(50.4 - 66.2)		
≥ College Degree	*	*	*	4,277	59.2	(52.4 - 65.9)		
Income								
≤ \$24,999	*	*	*	*	*	*		
\$25,000-\$49,999	*	*	*	3,139	60.8	(53.1 - 68.4)		
\$50,000-\$74,999	*	*	*	*	*	*		
≥ \$75,000	*	*	*	3,295	56.5	(50.5 - 62.4)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as having had a digital rectal exam within the past 2 years among men aged 40 and older

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>&</sup>lt;sup>3</sup> Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 31a. Ever had a prostate specific antigen test among men aged 40 and older in Rockland County and New York State

		Rockla	nd Coun	ty		New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	195	46,000	71.1	(61.9 - 80.2)	11,557	2,822,600	67.6	(63.9 - 71.2)		
Gender										
Male	195	46,000	71.1	(61.9 - 80.2)	11,557	2,822,600	67.6	(63.9 - 71.2)		
Female	-	-	-	-	-	-	-	-		
Age (years)										
40-44	*	*	*	*	*	*	*	*		
45-54	*	*	*	*	2,938	760,600	57.4	(50.6 - 64.3)		
55-64	*	*	*	*	3,373	811,400	83.4	(78.8 - 88.1)		
≥ 65	84	15,600	95.8	(91.3 - 100)	4,215	905,400	90.4	(87.1 - 93.6)		
Race/Ethnicity										
White Non-Hispanic	176	40,700	77.6	(69.7 - 85.5)	10,950	1,965,400	67.0	(63.9 - 70.1)		
Black Non-Hispanic	*	*	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*	*	*		
Education										
≤ High School	*	*	*	*	4,775	829,200	59.0	(52.4 - 65.6)		
Some College	*	*	*	*	2,584	535,000	61.9	(53.7 - 70.2)		
≥ College Degree	*	*	*	*	4,176	1,455,700	76.8	(72.5 - 81.1)		
Income										
≤ \$24,999	*	*	*	*	*	*	*	*		
\$25,000-\$49,999	*	*	*	*	3,048	444,100	64.1	(56.0 - 72.3)		
\$50,000-\$74,999	*	*	*	*	*	*	*	*		
≥ \$75,000	*	*	*	*	3,188	1,211,100	72.3	(67.6 - 77.0)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as having ever had a prostate specific antigen test among men aged 40 and older

 $<sup>^{2}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of men aged 40 and older who have ever had a prostate specific antigen test (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 31b. Ever had a prostate specific antigen test among men aged 40 and older<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	195	70.1	(60.1 - 80.1)	11,557	67.4	(63.7 - 71.1)		
Gender								
Male	195	70.1	(60.1 - 80.1)	11,557	67.4	(63.7 - 71.1)		
Female	-	-	-	-	-	-		
Race/Ethnicity								
White Non-Hispanic	176	77.7	(69.6 - 85.9)	10,950	66.6	(63.4 - 69.8)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	4,775	59.7	(52.9 - 66.4)		
Some College	*	*	*	2,584	61.4	(53.0 - 69.9)		
≥ College Degree	*	*	*	4,176	76.3	(71.9 - 80.7)		
Income								
≤ \$24,999	*	*	*	*	*	*		
\$25,000-\$49,999	*	*	*	3,048	64.2	(56.1 - 72.4)		
\$50,000-\$74,999	*	*	*	*	*	*		
≥ \$75,000	*	*	*	3,188	71.5	(66.6 - 76.3)		

<sup>&</sup>lt;sup>1</sup> Defined as having ever had a prostate specific antigen test among men aged 40 and older

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>&</sup>lt;sup>3</sup> Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 32a. Had prostate specific antigen test within the past 2 years among men aged 40 and older<sup>1</sup> in Rockland County and New York State

		Rockla	nd Coun	ty		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	192	38,200	59.6	(50.1 - 69.0)	11,437	2,493,800	59.9	(56.0 - 63.9)
Gender								
Male	192	38,200	59.6	(50.1 - 69.0)	11,437	2,493,800	59.9	(56.0 - 63.9)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,917	659,400	49.9	(42.8 - 57.0)
55-64	*	*	*	*	3,326	719,700	74.5	(69.4 - 79.7)
≥ 65	82	14,300	89.9	(82.9 - 96.9)	4,166	830,000	83.3	(78.4 - 88.3)
Race/Ethnicity								
White Non-Hispanic	173	34,900	67.4	(58.6 - 76.2)	10,836	1,762,200	60.4	(57.3 - 63.5)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	4,727	734,200	52.4	(45.7 - 59.1)
Some College	*	*	*	*	2,558	483,800	56.3	(48.2 - 64.4)
≥ College Degree	*	*	*	*	4,133	1,273,200	67.4	(61.6 - 73.3)
Income								
≤ \$24,999	*	*	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	*	3,021	410,800	59.6	(51.5 - 67.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,170	1,080,100	64.6	(59.1 - 70.0)

 $<sup>^{</sup>m 1}$  Defined as having had a prostate specific antigen test within the past 2 years among men aged 40 and older

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of men aged 40 and older who have had a prostate specific antigen test within the past 2 years (rounded to the nearest hundred)
4 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 32b. Had prostate specific antigen test within the past 2 years among men aged 40 and older in Rockland County and New York State - Age-Adjusted 2

	R	ockland	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	192	59.2	(49.2 - 69.2)	11,437	59.8	(55.8 - 63.9)		
Gender								
Male	192	59.2	(49.2 - 69.2)	11,437	59.8	(55.8 - 63.9)		
Female	-	-	-	-	-	-		
Race/Ethnicity								
White Non-Hispanic	173	67.5	(58.3 - 76.7)	10,836	60.1	(56.9 - 63.4)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	4,727	52.9	(45.9 - 59.9)		
Some College	*	*	*	2,558	56.0	(47.7 - 64.2)		
≥ College Degree	*	*	*	4,133	67.1	(61.2 - 73.1)		
Income								
≤ \$24,999	*	*	*	*	*	*		
\$25,000-\$49,999	*	*	*	3,021	59.8	(51.7 - 67.9)		
\$50,000-\$74,999	*	*	*	*	*	*		
≥ \$75,000	*	*	*	3,170	63.9	(58.3 - 69.6)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as having had a prostate specific antigen test within the past 2 years among men aged 40 and older

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 33a. Prostate cancer among men aged 40 and older<sup>1</sup> in Rockland County and New York State

		Rockla	ind County	/		New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	*	*	*	*	12,133	178,200	4.1	(3.0 - 5.2)		
Gender										
Male	*	*	*	*	12,133	178,200	4.1	(3.0 - 5.2)		
Female	-	-	=	-	-	-	-	-		
Age (years)										
40-44	*	*	*	*	*	*	*	*		
45-54	*	*	*	*	3,120	6,100	0.4	(0.0 - 0.9)		
55-64	*	*	*	*	3,532	31,600	3.1	(1.5 - 4.8)		
≥ 65	*	*	*	*	4,388	140,400	13.6	(9.7 - 17.4)		
Race/Ethnicity										
White Non-Hispanic	*	*	*	*	11,502	143,700	4.7	(3.5 - 5.8)		
Black Non-Hispanic	*	*	*	*	184	23,500	5.1	(0.0 - 11.0)		
Hispanic	*	*	*	*	197	7,700	1.6	(0.0 - 3.7)		
Other Non-Hispanic	*	*	*	*	*	*	*	*		
Education										
≤ High School	*	*	*	*	5,065	46,300	3.1	(2.2 - 4.1)		
Some College	*	*	*	*	2,709	39,200	4.4	(1.3 - 7.5)		
≥ College Degree	*	*	*	*	4,336	92,500	4.7	(3.0 - 6.5)		
Income										
≤ \$24,999	*	*	*	*	2,597	35,800	4.1	(2.2 - 6.0)		
\$25,000-\$49,999	*	*	*	*	3,196	29,600	4.1	(2.9 - 5.4)		
\$50,000-\$74,999	*	*	*	*	2,007	20,300	2.8	(1.2 - 4.4)		
≥ \$75,000	*	*	*	*	3,322	56,500	3.2	(1.7 - 4.8)		

 $<sup>^{</sup>m 1}$  Defined as having ever been told by a health professional that respondent had prostate cancer among men aged 40 and older

 $<sup>^{2}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of men aged 40 and older who have ever been told they had prostate cancer (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 33b. Prostate cancer among men aged 40 and older<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland C	ounty	N	ew York	State
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	*	*	*	12,133	4.5	(3.2 - 5.7)
Gender						
Male	*	*	*	12,133	4.5	(3.2 - 5.7)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	*	*	*	11,502	5.1	(3.8 - 6.4)
Black Non-Hispanic	*	*	*	184	5.7	(0.0 - 12.8)
Hispanic	*	*	*	197	1.8	(0.0 - 4.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	5,065	3.3	(2.3 - 4.3)
Some College	*	*	*	2,709	5.0	(1.1 - 9.0)
≥ College Degree	*	*	*	4,336	5.2	(3.1 - 7.2)
Income						
≤ \$24,999	*	*	*	2,597	4.5	(2.2 - 6.7)
\$25,000-\$49,999	*	*	*	3,196	4.3	(3.0 - 5.6)
\$50,000-\$74,999	*	*	*	2,007	3.0	(1.1 - 5.0)
≥ \$75,000	*	*	*	3,322	3.5	(1.6 - 5.4)

 $<sup>^{</sup>m 1}$  Defined as having ever been told by a health professional that respondent had prostate cancer among men aged 40 and older

 $<sup>^{2}</sup>$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 34a. Home blood stool test ever used among adults aged 50 and older<sup>1</sup> in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	420	33,000	32.3	(26.7 - 37.9)	25,322	2,288,500	38.0	(35.5 - 40.4)
Gender								
Male	163	16,200	34.2	(25.2 - 43.2)	9,571	1,081,900	38.9	(35.1 - 42.6)
Female	257	16,800	30.6	(23.7 - 37.6)	15,751	1,206,600	37.2	(34.0 - 40.4)
Age (years)								
50-64	207	17,500	27.4	(19.6 - 35.2)	12,834	1,192,000	34.0	(30.8 - 37.2)
≥ 65	210	15,300	40.9	(33.5 - 48.3)	12,472	1,093,400	43.6	(39.8 - 47.4)
Race/Ethnicity								
White Non-Hispanic	368	26,400	33.3	(27.7 - 39.0)	24,213	1,754,500	39.1	(37.1 - 41.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	145	9,800	26.8	(17.0 - 36.6)	11,146	823,500	34.9	(30.5 - 39.2)
Some College	*	*	*	*	6,261	480,300	37.0	(32.6 - 41.4)
≥ College Degree	201	17,200	36.7	(28.7 - 44.8)	7,851	973,700	41.6	(37.8 - 45.3)
Income								
≤ \$24,999	*	*	*	*	7,222	505,000	35.4	(29.7 - 41.2)
\$25,000-\$49,999	78	3,000	16.9	(7.9 - 25.8)	6,609	467,600	40.3	(35.2 - 45.4)
\$50,000-\$74,999	*	*	*	*	3,500	313,000	38.2	(32.6 - 43.9)
≥ \$75,000	147	17,300	41.5	(31.5 - 51.5)	4,647	657,800	37.3	(33.0 - 41.6)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having ever used a home blood stool test among adults aged 50 and older

<sup>&</sup>lt;sup>2</sup> Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults aged 50 and older who have ever used a home blood stool test (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 34b. Home blood stool test ever used among adults aged 50 and older<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	420	32.3	(26.8 - 37.8)	25,322	38.1	(35.5 - 40.8)		
Gender								
Male	163	34.4	(25.7 - 43.1)	9,571	39.4	(35.3 - 43.5)		
Female	257	30.3	(23.4 - 37.2)	15,751	36.9	(33.5 - 40.3)		
Race/Ethnicity								
White Non-Hispanic	368	33.6	(27.9 - 39.3)	24,213	38.9	(36.7 - 41.1)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	145	26.9	(17.5 - 36.4)	11,146	35.1	(30.3 - 39.9)		
Some College	*	*	*	6,261	36.6	(32.0 - 41.3)		
≥ College Degree	201	36.7	(28.7 - 44.6)	7,851	41.9	(37.9 - 46.0)		
Income								
≤ \$24,999	*	*	*	7,222	35.9	(29.5 - 42.4)		
\$25,000-\$49,999	78	17.0	(8.1 - 25.9)	6,609	40.2	(34.8 - 45.6)		
\$50,000-\$74,999	*	*	*	3,500	38.1	(31.9 - 44.3)		
≥ \$75,000	147	40.3	(30.6 - 50.1)	4,647	37.5	(32.9 - 42.2)		

 $<sup>^{\</sup>rm I}$  Defined as having ever used a home blood stool test among adults aged 50 and older

 $<sup>^{2}</sup>$  Prevalence estimates are age-adjusted to the 2000 US Standard population  $\,$ 

 $<sup>^{\</sup>rm 3}$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 35a. Home blood stool test used within the past year among adults aged 50 and older in Rockland County and New York State

		Rockla	ind Count	ty		New Y	ork State	9
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	420	9,000	8.8	(5.7 - 12.0)	25,074	863,300	14.4	(12.6 - 16.2)
Gender								
Male	163	3,600	7.6	(3.3 - 12.0)	9,508	444,400	16.0	(13.2 - 18.9)
Female	257	5,400	9.8	(5.3 - 14.4)	15,566	418,900	13.1	(10.8 - 15.4)
Age (years)								
50-64	207	4,900	7.7	(3.5 - 11.9)	12,759	454,400	13.0	(10.6 - 15.5)
≥ 65	210	4,100	11.0	(6.2 - 15.7)	12,299	406,400	16.4	(13.7 - 19.0)
Race/Ethnicity								
White Non-Hispanic	368	7,700	9.7	(6.2 - 13.3)	23,971	634,300	14.3	(12.8 - 15.7)
Black Non-Hispanic	*	*	*	*	396	136,000	18.6	(9.4 - 27.8)
Hispanic	*	*	*	*	349	53,400	9.9	(2.5 - 17.2)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	145	2,700	7.4	(2.2 - 12.6)	11,026	288,600	12.3	(9.4 - 15.3)
Some College	*	*	*	*	6,208	212,500	16.4	(12.5 - 20.4)
≥ College Degree	201	4,400	9.4	(5.0 - 13.9)	7,778	352,400	15.2	(12.5 - 17.9)
Income								
≤ \$24,999	*	*	*	*	7,157	196,300	13.9	(10.1 - 17.6)
\$25,000-\$49,999	*	*	*	*	6,562	221,500	19.2	(14.1 - 24.3)
\$50,000-\$74,999	*	*	*	*	3,469	111,900	13.7	(10.1 - 17.4)
≥ \$75,000	147	4,200	10.1	(4.7 - 15.5)	4,623	205,100	11.7	(8.8 - 14.6)

 $<sup>^{</sup>m 1}$  Defined as having used a home blood stool test within the past year among adults aged 50 and older

 $<sup>^{2}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults aged 50 and older who have used a home blood stool test within the past year (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 35b. Home blood stool test used within the past year among adults aged 50 and older in Rockland County and New York State - Age-Adjusted 2

	R	ockland (	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	420	8.9	(5.7 - 12.2)	25,074	14.6	(12.6 - 16.7)		
Gender								
Male	163	7.9	(3.6 - 12.3)	9,508	16.4	(13.2 - 19.6)		
Female	257	9.9	(5.2 - 14.6)	15,566	12.9	(10.4 - 15.4)		
Race/Ethnicity								
White Non-Hispanic	368	9.9	(6.3 - 13.5)	23,971	14.2	(12.7 - 15.7)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	349	10.2	(1.7 - 18.7)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	145	7.5	(2.3 - 12.7)	11,026	12.7	(9.2 - 16.1)		
Some College	*	*	*	6,208	16.6	(12.3 - 20.9)		
≥ College Degree	201	9.6	(5.0 - 14.1)	7,778	15.3	(12.4 - 18.3)		
Income								
≤ \$24,999	*	*	*	7,157	14.3	(9.8 - 18.7)		
\$25,000-\$49,999	*	*	*	6,562	19.3	(13.8 - 24.8)		
\$50,000-\$74,999	*	*	*	3,469	13.8	(9.8 - 17.7)		
≥ \$75,000	147	9.9	(4.5 - 15.2)	4,623	11.8	(8.6 - 15.0)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as having used a home blood stool test within the past year among adults aged 50 and older

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 36a. Home blood stool test used within the past 2 years among adults aged 50 and older<sup>1</sup> in Rockland County and New York State

		Rockla	ind Count	ty .		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	420	12,900	12.7	(8.9 - 16.4)	25,074	1,218,000	20.4	(18.3 - 22.5)
Gender								
Male	163	5,800	12.3	(6.4 - 18.2)	9,508	593,100	21.4	(18.3 - 24.5)
Female	257	7,100	13.0	(8.1 - 17.8)	15,566	624,900	19.5	(16.6 - 22.4)
Age (years)								
50-64	207	7,300	11.4	(6.3 - 16.6)	12,759	629,900	18.1	(15.4 - 20.7)
≥ 65	210	5,500	14.6	(9.3 - 19.8)	12,299	585,500	23.6	(20.2 - 27.0)
Race/Ethnicity								
White Non-Hispanic	368	11,300	14.2	(9.9 - 18.6)	23,971	877,500	19.7	(18.2 - 21.3)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	145	3,800	10.3	(4.6 - 15.9)	11,026	454,600	19.4	(15.6 - 23.3)
Some College	*	*	*	*	6,208	262,200	20.3	(16.3 - 24.2)
≥ College Degree	201	7,300	15.6	(9.5 - 21.6)	7,778	491,400	21.2	(18.1 - 24.2)
Income								
≤ \$24,999	63	2,000	14.7	(5.1 - 24.3)	7,157	263,900	18.6	(14.6 - 22.7)
\$25,000-\$49,999	*	*	*	*	6,562	285,900	24.8	(19.6 - 30.0)
\$50,000-\$74,999	*	*	*	*	3,469	167,500	20.6	(15.6 - 25.5)
≥ \$75,000	147	5,800	14.0	(7.7 - 20.3)	4,623	294,900	16.8	(13.7 - 20.0)

 $<sup>^{</sup>m I}$  Defined as having used a home blood stool test within the past 2 years among adults aged 50 and older

 $<sup>^{2} \, \</sup>mathrm{Sample} \, \mathrm{size} \, \mathrm{from} \, \mathrm{survey}$ 

<sup>&</sup>lt;sup>3</sup> Estimated number of adults aged 50 and older who have used a home blood stool test within the past 2 years (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 36b. Home blood stool test used within the past 2 years among adults aged 50 and older<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	ew York	State
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>
Total	420	12.9	(9.0 - 16.8)	25,074	20.5	(18.2 - 22.8)
Gender						
Male	163	12.7	(6.8 - 18.7)	9,508	21.7	(18.3 - 25.1)
Female	257	13.1	(8.1 - 18.1)	15,566	19.4	(16.3 - 22.5)
Race/Ethnicity						
White Non-Hispanic	368	14.5	(10.1 - 19.0)	23,971	19.7	(18.0 - 21.3)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	145	10.6	(4.9 - 16.3)	11,026	19.5	(15.4 - 23.7)
Some College	*	*	*	6,208	20.4	(16.1 - 24.7)
≥ College Degree	201	15.7	(9.6 - 21.9)	7,778	21.4	(18.1 - 24.7)
Income						
≤ \$24,999	*	*	*	7,157	18.9	(14.2 - 23.6)
\$25,000-\$49,999	*	*	*	6,562	24.8	(19.3 - 30.3)
\$50,000-\$74,999	*	*	*	3,469	20.6	(15.3 - 25.9)
≥ \$75,000	147	13.9	(7.5 - 20.3)	4,623	16.9	(13.4 - 20.4)

 $<sup>^{</sup>m 1}$  Defined as having used a home blood stool test within the past 2 years among adults aged 50 and older

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 37a. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in Rockland County and New York State

		Rockla	and Coun	ty		New Y	ork State	e
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	421	67,300	66.2	(60.0 - 72.3)	25,357	4,019,100	66.6	(64.1 - 69.1)
Gender								
Male	164	30,900	65.6	(55.9 - 75.2)	9,582	1,894,100	68.0	(64.2 - 71.7)
Female	257	36,400	66.7	(58.7 - 74.6)	15,775	2,125,000	65.5	(62.1 - 68.8)
Age (years)								
50-54	*	*	*	*	4,179	718,900	50.6	(44.6 - 56.5)
55-64	*	*	*	*	8,695	1,493,500	71.0	(67.3 - 74.8)
≥ 65	211	29,900	80.6	(74.6 - 86.6)	12,466	1,802,000	72.1	(68.6 - 75.7)
Race/Ethnicity								
White Non-Hispanic	369	56,400	71.6	(65.8 - 77.4)	24,248	3,093,900	68.8	(66.8 - 70.8)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	11,121	1,445,100	61.2	(56.7 - 65.8)
Some College	*	*	*	*	6,287	835,300	64.3	(59.7 - 69.0)
≥ College Degree	202	33,200	70.8	(62.9 - 78.7)	7,884	1,732,400	73.7	(70.0 - 77.4)
Income								
≤ \$24,999	*	*	*	*	7,220	892,600	62.7	(56.8 - 68.6)
\$25,000-\$49,999	*	*	*	*	6,619	697,300	60.1	(54.6 - 65.7)
\$50,000-\$74,999	*	*	*	*	3,514	613,700	74.2	(68.9 - 79.6)
≥ \$75,000	148	29,400	70.4	(60.4 - 80.4)	4,665	1,294,400	73.3	(69.2 - 77.5)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having ever had a sigmoidoscopy or colonoscopy among adults aged 50 and older

 $<sup>^2\,\</sup>mathrm{Sample}$  size from survey

 $<sup>^3</sup>$  Estimated number of adults aged 50 and older who have ever had a sigmoidoscopy or colonoscopy (rounded to the nearest hundred)

 <sup>4 95%</sup> confidence interval
 \* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 37b. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	421	66.6	(60.6 - 72.7)	25,357	66.6	(63.9 - 69.3)		
Gender								
Male	164	66.9	(57.7 - 76.2)	9,582	68.5	(64.5 - 72.5)		
Female	257	66.3	(58.4 - 74.2)	15,775	64.8	(61.2 - 68.4)		
Race/Ethnicity								
White Non-Hispanic	369	71.7	(65.9 - 77.6)	24,248	68.7	(66.5 - 70.8)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	11,121	61.3	(56.4 - 66.2)		
Some College	*	*	*	6,287	63.8	(58.8 - 68.8)		
≥ College Degree	202	71.4	(63.6 - 79.1)	7,884	74.0	(70.2 - 77.9)		
Income								
≤ \$24,999	*	*	*	7,220	63.0	(56.7 - 69.3)		
\$25,000-\$49,999	*	*	*	6,619	59.7	(53.8 - 65.6)		
\$50,000-\$74,999	*	*	*	3,514	74.4	(68.3 - 80.5)		
≥ \$75,000	148	70.9	(61.2 - 80.5)	4,665	73.6	(69.3 - 78.0)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as having ever had a sigmoidoscopy or colonoscopy among adults aged 50 and older

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 38a. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older<sup>1</sup> in Rockland County and New York State

		Rockla	nd Coun	ty		New Y	ork State	•
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	415	65,400	64.9	(58.7 - 71.1)	25,169	3,855,800	64.6	(62.1 - 67.1)
Gender								
Male	163	30,400	64.7	(55.0 - 74.3)	9,521	1,835,400	66.4	(62.6 - 70.1)
Female	252	35,000	65.1	(57.1 - 73.1)	15,648	2,020,400	63.1	(59.8 - 66.4)
Age (years)								
50-54	*	*	*	*	4,169	700,500	49.3	(43.3 - 55.3)
55-64	*	*	*	*	8,652	1,455,100	69.4	(65.7 - 73.2)
≥ 65	206	28,700	78.8	(72.6 - 85.0)	12,331	1,695,700	69.5	(65.9 - 73.1)
Race/Ethnicity								
White Non-Hispanic	363	54,500	70.0	(64.1 - 75.9)	24,069	2,981,500	66.6	(64.6 - 68.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	11,017	1,360,400	59.0	(54.5 - 63.5)
Some College	*	*	*	*	6,251	807,100	62.4	(57.8 - 67.1)
≥ College Degree	200	32,300	69.2	(61.2 - 77.2)	7,841	1,682,500	71.7	(68.0 - 75.4)
Income								
≤ \$24,999	*	*	*	*	7,163	858,900	61.0	(55.1 - 67.0)
\$25,000-\$49,999	*	*	*	*	6,584	664,900	57.7	(52.2 - 63.2)
\$50,000-\$74,999	*	*	*	*	3,501	602,600	73.1	(67.7 - 78.5)
≥ \$75,000	147	28,900	69.5	(59.5 - 79.4)	4,655	1,260,400	71.5	(67.3 - 75.7)

<sup>&</sup>lt;sup>1</sup> Defined as having had a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults aged 50 and older who have had a sigmoidoscopy or colonoscopy within the past 10 years (rounded to the nearest hundred)
4 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 38b. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older  $^1$  in Rockland County and New York State - Age-Adjusted  $^2$ 

	R	ockland	County	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	415	65.4	(59.3 - 71.5)	25,169	64.6	(61.9 - 67.3)		
Gender								
Male	163	66.2	(56.9 - 75.4)	9,521	66.9	(62.9 - 70.9)		
Female	252	64.7	(56.7 - 72.7)	15,648	62.4	(58.9 - 66.0)		
Race/Ethnicity								
White Non-Hispanic	363	70.3	(64.4 - 76.2)	24,069	66.5	(64.3 - 68.7)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	11,017	59.0	(54.1 - 63.9)		
Some College	*	*	*	6,251	61.9	(56.9 - 66.9)		
≥ College Degree	200	69.8	(61.9 - 77.7)	7,841	72.1	(68.2 - 75.9)		
Income								
≤ \$24,999	*	*	*	7,163	61.3	(54.9 - 67.6)		
\$25,000-\$49,999	*	*	*	6,584	57.2	(51.4 - 63.0)		
\$50,000-\$74,999	*	*	*	3,501	73.3	(67.2 - 79.4)		
≥ \$75 <b>,</b> 000	147	70.0	(60.3 - 79.6)	4,655	71.8	(67.4 - 76.2)		

 $<sup>^{</sup>m I}$  Defined as having had a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 39a. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	e
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	403	66,100	67.3	(61.0 - 73.6)	24,792	4,049,200	69.1	(66.7 - 71.6)
Gender								
Male	160	30,400	65.4	(55.6 - 75.1)	9,418	1,945,200	70.7	(67.0 - 74.3)
Female	243	35,700	69.0	(61.0 - 77.1)	15,374	2,104,100	67.8	(64.6 - 71.1)
Age (years)								
50-54	*	*	*	*	4,132	751,200	54.1	(48.0 - 60.1)
55-64	*	*	*	*	8,366	1,487,600	73.7	(70.0 - 77.4)
≥ 65	205	28,800	79.3	(73.1 - 85.5)	12,294	1,810,500	74.0	(70.6 - 77.4)
Race/Ethnicity								
White Non-Hispanic	355	54,800	71.7	(65.7 - 77.6)	23,726	3,091,400	70.4	(68.4 - 72.4)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	10,872	1,461,600	64.2	(59.8 - 68.7)
Some College	*	*	*	*	6,170	873,300	68.4	(63.9 - 72.8)
≥ College Degree	193	32,300	71.2	(63.1 - 79.3)	7,708	1,709,000	74.6	(70.9 - 78.3)
Income								
≤ \$24,999	*	*	*	*	7,092	923,300	66.0	(60.1 - 71.9)
\$25,000-\$49,999	*	*	*	*	6,526	730,000	64.2	(58.9 - 69.6)
\$50,000-\$74,999	*	*	*	*	3,467	621,500	76.6	(71.4 - 81.9)
≥ \$75,000	*	*	*	*	4,615	1,292,500	73.8	(69.8 - 77.9)

<sup>&</sup>lt;sup>1</sup> Defined as having used a home blood stool test within the past year or a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 years and older  $^2$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults aged 50 and older who have used a home blood stool test in the past year or sigmoidoscopy or colonoscopy within the past 10 years (rounded to the nearest hundred)  $^4$  95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 39b. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	Ro	ockland (	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	403	67.9	(61.7 - 74.0)	24,792	69.3	(66.7 - 71.9)		
Gender								
Male	160	66.8	(57.5 - 76.2)	9,418	71.5	(67.6 - 75.4)		
Female	243	68.9	(60.9 - 76.8)	15,374	67.2	(63.7 - 70.7)		
Race/Ethnicity								
White Non-Hispanic	355	71.8	(65.9 - 77.7)	23,726	70.3	(68.1 - 72.4)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	10,872	64.5	(59.8 - 69.3)		
Some College	*	*	*	6,170	68.2	(63.4 - 72.9)		
≥ College Degree	193	71.7	(63.8 - 79.7)	7,708	75.0	(71.2 - 78.8)		
Income								
≤ \$24,999	*	*	*	7,092	66.4	(60.2 - 72.6)		
\$25,000-\$49,999	*	*	*	6,526	63.8	(58.2 - 69.5)		
\$50,000-\$74,999	*	*	*	3,467	76.7	(70.6 - 82.7)		
≥ \$75,000	146	71.2	(61.6 - 80.9)	4,615	74.4	(70.2 - 78.6)		

<sup>&</sup>lt;sup>1</sup> Defined as having used a home blood stool test within the past year or a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 years and older

Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 40a. High blood pressure among adults<sup>1</sup> in Rockland County and New York State

		Rockla	nd Coun	ty		New Y	ork State	<b>e</b>
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	657	62,400	28.8	(24.3 - 33.2)	38,125	3,974,800	26.8	(24.9 - 28.7)
Gender								
Male	243	31,300	29.7	(22.7 - 36.7)	14,611	1,960,500	27.7	(24.7 - 30.7)
Female	414	31,200	27.9	(22.3 - 33.4)	23,514	2,014,300	26.0	(23.6 - 28.3)
Age (years)								
18-34	*	*	*	*	3,935	312,200	7.0	(4.2 - 9.8)
35-44	*	*	*	*	4,984	426,200	14.8	(11.6 - 18.1)
45-54	*	*	*	*	7,825	862,400	30.3	(25.9 - 34.8)
55-64	*	*	*	*	8,765	890,800	41.6	(37.6 - 45.7)
≥ 65	217	23,200	59.7	(52.4 - 66.9)	12,616	1,483,200	58.5	(54.7 - 62.3)
Race/Ethnicity								
White Non-Hispanic	550	43,800	29.1	(24.6 - 33.6)	35,814	2,650,900	28.4	(26.9 - 30.0)
Black Non-Hispanic	*	*	*	*	736	558,500	26.2	(19.6 - 32.8)
Hispanic	*	*	*	*	863	538,100	23.8	(17.0 - 30.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	199	22,000	33.2	(24.6 - 41.9)	15,688	1,851,200	33.1	(29.4 - 36.9)
Some College	*	*	*	*	9,906	851,100	24.1	(20.9 - 27.3)
≥ College Degree	339	28,700	26.4	(20.4 - 32.4)	12,452	1,253,400	22.3	(19.7 - 24.9)
Income								
≤ \$24,999	*	*	*	*	9,640	1,153,000	32.8	(27.6 - 38.1)
\$25,000-\$49,999	*	*	*	*	9,657	807,800	27.9	(23.7 - 32.1)
\$50,000-\$74,999	*	*	*	*	5,773	507,600	25.8	(21.4 - 30.1)
≥ \$75,000	270	25,100	25.3	(18.8 - 31.8)	8,560	942,500	21.0	(18.4 - 23.6)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having ever been told by a health professional that respondent has high blood pressure

<sup>&</sup>lt;sup>2</sup> Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults who have ever been told by a health professional that they have high blood pressure (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 40b. High blood pressure among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	657	26.1	(21.7 - 30.4)	38,125	25.7	(23.8 - 27.5)		
Gender								
Male	243	28.0	(21.0 - 35.0)	14,611	27.7	(24.7 - 30.7)		
Female	414	24.2	(18.8 - 29.7)	23,514	23.8	(21.5 - 26.0)		
Race/Ethnicity								
White Non-Hispanic	550	26.2	(21.8 - 30.6)	35,814	27.2	(25.6 - 28.7)		
Black Non-Hispanic	*	*	*	736	24.9	(18.4 - 31.3)		
Hispanic	*	*	*	863	23.3	(16.6 - 30.0)		
Other Non-Hispanic	*	*	*	712	19.3	(9.5 - 29.1)		
Education								
≤ High School	199	30.5	(22.0 - 39.0)	15,688	32.1	(28.4 - 35.8)		
Some College	*	*	*	9,906	22.8	(19.7 - 25.9)		
≥ College Degree	339	23.9	(17.9 - 29.9)	12,452	21.2	(18.6 - 23.8)		
Income								
≤ \$24,999	*	*	*	9,640	31.8	(26.7 - 37.0)		
\$25,000-\$49,999	*	*	*	9,657	26.9	(22.8 - 30.9)		
\$50,000-\$74,999	*	*	*	5,773	24.8	(20.4 - 29.2)		
≥ \$75,000	270	22.2	(16.1 - 28.3)	8,560	19.9	(17.3 - 22.5)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as having ever been told by a health professional that respondent has high blood pressure

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 41a. Blood pressure medication among adults with high blood pressure in Rockland County and New York State

		Rockla	nd Coun	ty		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	233	45,800	73.4	(64.0 - 82.7)	15,288	3,145,800	79.2	(76.0 - 82.4)
Gender								
Male	*	*	*	*	6,080	1,432,100	73.1	(67.6 - 78.5)
Female	138	25,400	81.4	(71.4 - 91.3)	9,208	1,713,700	85.2	(82.1 - 88.2)
Age (years)								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,315	645,500	74.9	(66.8 - 82.9)
55-64	*	*	*	*	4,045	782,300	87.8	(84.5 - 91.1)
≥ 65	128	22,200	95.9	(92.5 - 99.3)	7,675	1,383,400	93.4	(91.1 - 95.7)
Race/Ethnicity								
White Non-Hispanic	198	35,200	80.2	(72.5 - 87.9)	14,442	2,117,300	79.9	(77.5 - 82.4)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	7,354	1,482,800	80.2	(74.7 - 85.6)
Some College	*	*	*	*	3,868	632,000	74.3	(67.8 - 80.7)
≥ College Degree	*	*	*	*	4,035	1,013,200	80.9	(76.8 - 84.9)
Income								
≤ \$24,999	*	*	*	*	4,898	948,200	82.3	(75.7 - 88.9)
\$25,000-\$49,999	*	*	*	*	4,088	668,300	82.8	(77.6 - 88.0)
\$50,000-\$74,999	*	*	*	*	1,971	391,500	77.2	(67.7 - 86.6)
≥ \$75,000	*	*	*	*	2,397	683,700	72.5	(66.1 - 78.9)

 $<sup>^{\</sup>mathrm{1}}$  Defined as currently taking medicine for high blood pressure among adults with high blood pressure

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults with high blood pressure who are taking medicine for high blood pressure (rounded to the nearest hundred)
<sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 41b. Blood pressure medication among adults with high blood pressure<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	*	*	*	15,288	78.5	(75.2 - 81.7)		
Gender								
Male	*	*	*	6,080	73.3	(67.9 - 78.7)		
Female	*	*	*	9,208	84.0	(80.7 - 87.2)		
Race/Ethnicity								
White Non-Hispanic	198	80.3	(72.2 - 88.4)	14,442	79.0	(76.5 - 81.6)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	*	*	*		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	7,354	79.7	(74.3 - 85.2)		
Some College	*	*	*	3,868	72.5	(65.7 - 79.3)		
≥ College Degree	*	*	*	4,035	80.4	(76.1 - 84.6)		
Income								
≤ \$24,999	*	*	*	4,898	81.6	(75.0 - 88.3)		
\$25,000-\$49,999	*	*	*	4,088	82.1	(76.7 - 87.5)		
\$50,000-\$74,999	*	*	*	1,971	76.4	(66.7 - 86.1)		
≥ \$75,000	*	*	*	2,397	71.6	(64.8 - 78.3)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as currently taking medicine for high blood pressure among adults with high blood pressure

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 42a. Ever had blood cholesterol checked among adults<sup>1</sup> in Rockland County and New York State

		Rockla	nd Coun	ty		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	635	172,200	83.3	(78.5 - 88.1)	37,450	11,496,600	80.3	(77.7 - 83.0)
Gender								
Male	238	84,900	83.9	(76.4 - 91.5)	14,389	5,288,400	77.7	(73.3 - 82.0)
Female	397	87,300	82.6	(76.6 - 88.6)	23,061	6,208,200	82.7	(79.6 - 85.8)
Age (years)								
18-34	*	*	*	*	3,684	2,296,400	55.3	(48.7 - 61.9)
35-44	*	*	*	*	4,883	2,264,200	81.6	(76.6 - 86.5)
45-54	128	40,400	94.4	(89.8 - 98.9)	7,731	2,539,400	91.0	(87.0 - 94.9)
55-64	133	34,300	98.5	(96.6 - 100)	8,718	2,014,900	95.4	(93.5 - 97.2)
≥ 65	214	38,000	99.7	(99.2 - 100)	12,434	2,381,700	95.9	(94.3 - 97.6)
Race/Ethnicity								
White Non-Hispanic	532	124,800	87.3	(83.0 - 91.7)	35,198	7,832,600	86.3	(84.8 - 87.9)
Black Non-Hispanic	*	*	*	*	720	1,588,900	77.8	(68.4 - 87.2)
Hispanic	*	*	*	*	842	1,224,600	56.8	(46.8 - 66.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	188	46,100	76.9	(67.1 - 86.7)	15,342	3,795,700	70.6	(65.6 - 75.6)
Some College	*	*	*	*	9,732	2,599,800	77.4	(71.7 - 83.2)
≥ College Degree	332	93,000	87.7	(81.3 - 94.1)	12,296	5,008,300	91.3	(88.9 - 93.6)
Income								
≤ \$24,999	*	*	*	*	9,437	2,277,500	68.4	(61.5 - 75.3)
\$25,000-\$49,999	*	*	*	*	9,492	2,053,000	71.9	(64.5 - 79.3)
\$50,000-\$74,999	*	*	*	*	5,721	1,736,100	90.0	(87.7 - 92.3)
≥ \$75,000	265	87,100	90.2	(84.6 - 95.8)	8,460	4,019,300	92.2	(90.8 - 93.6)

<sup>&</sup>lt;sup>1</sup> Defined as ever having blood cholesterol checked

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults who have ever had blood cholesterol checked (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 42b. Ever had blood cholesterol checked among adults<sup>1</sup> in Rockland County and New York
State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	635	81.1	(75.7 - 86.4)	37,450	79.9	(77.3 - 82.5)		
Gender								
Male	238	83.3	(75.3 - 91.3)	14,389	78.0	(73.9 - 82.1)		
Female	397	78.9	(71.8 - 86.0)	23,061	81.6	(78.4 - 84.8)		
Race/Ethnicity								
White Non-Hispanic	532	85.1	(80.1 - 90.1)	35,198	85.6	(84.0 - 87.2)		
Black Non-Hispanic	*	*	*	720	77.5	(68.2 - 86.7)		
Hispanic	*	*	*	842	57.2	(47.4 - 67.0)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	15,342	70.2	(65.2 - 75.1)		
Some College	*	*	*	9,732	76.7	(71.1 - 82.3)		
≥ College Degree	332	86.2	(78.9 - 93.5)	12,296	90.9	(88.6 - 93.2)		
Income								
≤ \$24,999	*	*	*	9,437	68.1	(61.3 - 74.9)		
\$25,000-\$49,999	*	*	*	9,492	71.6	(64.4 - 78.7)		
\$50,000-\$74,999	*	*	*	5,721	89.6	(87.2 - 92.0)		
≥ \$75,000	265	88.6	(82.0 - 95.2)	8,460	91.7	(90.1 - 93.2)		

<sup>&</sup>lt;sup>1</sup> Defined as ever having blood cholesterol checked

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 43a. Blood cholesterol checked within the past 5 years among adults<sup>1</sup> in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	•
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	632	165,600	80.3	(75.4 - 85.2)	37,201	11,035,200	77.8	(75.1 - 80.4)
Gender								
Male	238	81,800	80.8	(73.1 - 88.6)	14,307	5,020,700	74.5	(70.2 - 78.9)
Female	394	83,800	79.8	(73.7 - 85.9)	22,894	6,014,600	80.7	(77.6 - 83.8)
Age (years)								
18-34	*	*	*	*	3,647	2,179,800	53.3	(46.7 - 59.9)
35-44	*	*	*	*	4,862	2,161,200	78.0	(72.8 - 83.2)
45-54	128	39,400	92.1	(86.5 - 97.7)	7,697	2,408,600	86.6	(82.4 - 90.8)
55-64	131	32,800	95.4	(92.2 - 98.6)	8,677	1,957,900	93.3	(91.4 - 95.3)
≥ 65	213	37,400	98.3	(96.0 - 100)	12,318	2,327,800	94.8	(93.1 - 96.6)
Race/Ethnicity								
White Non-Hispanic	530	118,900	83.3	(78.6 - 88.1)	34,960	7,486,600	83.4	(81.8 - 85.0)
Black Non-Hispanic	*	*	*	*	719	1,561,700	76.5	(67.1 - 85.9)
Hispanic	*	*	*	*	838	1,184,000	55.6	(45.6 - 65.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	188	44,900	74.9	(65.1 - 84.8)	15,215	3,612,500	68.3	(63.2 - 73.3)
Some College	*	*	*	*	9,673	2,524,200	75.4	(69.8 - 81.1)
≥ College Degree	330	88,900	84.0	(77.3 - 90.6)	12,236	4,806,500	88.1	(85.5 - 90.6)
Income								
≤ \$24,999	*	*	*	*	9,350	2,214,800	66.9	(60.1 - 73.8)
\$25,000-\$49,999	*	*	*	*	9,454	1,936,600	68.8	(61.5 - 76.1)
\$50,000-\$74,999	*	*	*	*	5,701	1,640,400	85.9	(82.4 - 89.5)
≥ \$75,000	265	83,700	86.7	(80.7 - 92.8)	8,435	3,872,400	89.2	(87.5 - 90.9)

 $<sup>^{\</sup>rm 1}$  Defined as having had blood cholesterol checked within the past 5 years

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults who have had blood cholesterol checked within the past 5 years (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 43b. Blood cholesterol checked within the past 5 years among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	632	77.9	(72.5 - 83.4)	37,201	77.3	(74.7 - 79.9)		
Gender								
Male	238	80.0	(71.7 - 88.2)	14,307	74.8	(70.7 - 79.0)		
Female	394	76.0	(68.8 - 83.1)	22,894	79.5	(76.3 - 82.7)		
Race/Ethnicity								
White Non-Hispanic	530	80.9	(75.5 - 86.3)	34,960	82.6	(81.0 - 84.3)		
Black Non-Hispanic	*	*	*	719	76.2	(66.9 - 85.5)		
Hispanic	*	*	*	838	55.9	(46.1 - 65.8)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	15,215	67.8	(62.9 - 72.8)		
Some College	*	*	*	9,673	74.7	(69.1 - 80.2)		
≥ College Degree	330	82.0	(74.5 - 89.6)	12,236	87.5	(85.0 - 90.1)		
Income								
≤ \$24,999	*	*	*	9,350	66.6	(59.8 - 73.4)		
\$25,000-\$49,999	*	*	*	9,454	68.5	(61.4 - 75.6)		
\$50,000-\$74,999	*	*	*	5,701	85.4	(81.7 - 89.0)		
≥ \$75,000	265	84.6	(77.5 - 91.7)	8,435	88.5	(86.7 - 90.3)		

 $<sup>^{\</sup>rm 1}$  Defined as having had blood cholesterol checked within the past 5 years

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 44a. Cardiovascular disease among adults<sup>1</sup> in Rockland County and New York State

		Rockla	nd Coun	ty		New Y	ork State	•
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	655	13,200	6.1	(4.3 - 7.8)	37,810	1,135,000	7.7	(6.5 - 8.8)
Gender								
Male	244	7,900	7.5	(4.5 - 10.6)	14,477	664,600	9.4	(7.5 - 11.4)
Female	411	5,300	4.7	(2.9 - 6.6)	23,333	470,400	6.1	(4.8 - 7.4)
Age (years)								
18-34	*	*	*	*	3,930	100,800	2.3	(0.0 - 4.8)
35-44	*	*	*	*	4,971	103,400	3.6	(1.5 - 5.7)
45-54	*	*	*	*	7,768	131,300	4.6	(2.9 - 6.4)
55-64	*	*	*	*	8,699	224,200	10.6	(8.3 - 12.9)
≥ 65	215	9,100	23.8	(17.4 - 30.1)	12,442	575,300	23.0	(19.9 - 26.1)
Race/Ethnicity								
White Non-Hispanic	548	11,100	7.4	(5.2 - 9.6)	35,522	772,200	8.3	(7.4 - 9.2)
Black Non-Hispanic	*	*	*	*	728	140,700	6.6	(3.3 - 9.9)
Hispanic	*	*	*	*	854	163,300	7.3	(2.0 - 12.5)
Other Non-Hispanic	*	*	*	*	706	58,800	5.2	(0.9 - 9.5)
Education								
≤ High School	199	5,700	8.6	(4.9 - 12.4)	15,486	590,800	10.7	(8.2 - 13.2)
Some College	*	*	*	*	9,849	227,100	6.4	(4.7 - 8.1)
≥ College Degree	336	5,300	4.9	(2.8 - 7.0)	12,393	314,600	5.6	(4.3 - 6.9)
Income								
≤ \$24,999	92	4,300	13.4	(6.1 - 20.7)	9,505	448,100	12.9	(9.1 - 16.7)
\$25,000-\$49,999	*	*	*	*	9,571	179,300	6.2	(4.8 - 7.6)
\$50,000-\$74,999	*	*	*	*	5,751	166,900	8.5	(5.5 - 11.4)
≥ \$75,000	269	2,900	2.9	(1.2 - 4.6)	8,538	183,500	4.1	(2.9 - 5.3)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having ever been told by a health professional that respondent had a heart attack or angina or a stroke

<sup>&</sup>lt;sup>2</sup> Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults with cardiovascular disease (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 44b. Cardiovascular disease among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland (	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	655	5.5	(3.9 - 7.1)	37,810	7.6	(6.4 - 8.7)		
Gender								
Male	244	7.2	(4.3 - 10.0)	14,477	9.7	(7.8 - 11.7)		
Female	411	4.0	(2.4 - 5.5)	23,333	5.6	(4.3 - 6.9)		
Race/Ethnicity								
White Non-Hispanic	548	6.7	(4.7 - 8.7)	35,522	8.2	(7.2 - 9.1)		
Black Non-Hispanic	*	*	*	728	6.8	(3.2 - 10.4)		
Hispanic	*	*	*	854	7.0	(2.0 - 12.0)		
Other Non-Hispanic	*	*	*	706	5.5	(0.9 - 10.0)		
Education								
≤ High School	199	7.8	(4.4 - 11.2)	15,486	10.5	(8.0 - 13.0)		
Some College	*	*	*	9,849	6.4	(4.5 - 8.2)		
≥ College Degree	336	4.4	(2.5 - 6.4)	12,393	5.5	(4.2 - 6.9)		
Income								
≤ \$24,999	92	12.4	(5.4 - 19.3)	9,505	12.8	(9.0 - 16.6)		
\$25,000-\$49,999	*	*	*	9,571	6.0	(4.7 - 7.4)		
\$50,000-\$74,999	*	*	*	5,751	8.5	(5.4 - 11.6)		
≥ \$75,000	269	2.6	(1.1 - 4.1)	8,538	4.0	(2.7 - 5.2)		

 $<sup>^{1}</sup>$  Defined as having ever been told by a health professional that respondent had a heart attack or angina or a stroke

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{\</sup>rm 3}$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 45a. Coronary heart disease among adults<sup>1</sup> in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	e
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	655	11,000	5.1	(3.5 - 6.7)	37,838	929,400	6.3	(5.3 - 7.3)
Gender								
Male	244	7,000	6.6	(3.8 - 9.5)	14,479	598,600	8.5	(6.6 - 10.4)
Female	411	4,000	3.6	(2.0 - 5.2)	23,359	330,800	4.3	(3.4 - 5.2)
Age (years)								
18-34	*	*	*	*	3,932	63,400	1.4	(0.0 - 3.6)
35-44	*	*	*	*	4,973	95,300	3.3	(1.2 - 5.4)
45-54	*	*	*	*	7,778	108,300	3.8	(2.1 - 5.6)
55-64	*	*	*	*	8,703	188,600	8.9	(6.7 - 11.1)
≥ 65	215	7,600	19.8	(13.9 - 25.7)	12,452	473,800	18.9	(16.1 - 21.7)
Race/Ethnicity								
White Non-Hispanic	548	9,200	6.1	(4.1 - 8.1)	35,552	636,700	6.9	(6.0 - 7.7)
Black Non-Hispanic	*	*	*	*	728	113,100	5.3	(2.3 - 8.4)
Hispanic	*	*	*	*	853	128,000	5.7	(1.1 - 10.3)
Other Non-Hispanic	*	*	*	*	705	51,500	4.6	(0.5 - 8.7)
Education								
≤ High School	199	4,700	7.1	(3.6 - 10.5)	15,485	474,500	8.6	(6.3 - 10.8)
Some College	*	*	*	*	9,860	193,000	5.5	(3.9 - 7.1)
≥ College Degree	336	4,600	4.2	(2.3 - 6.2)	12,411	259,700	4.6	(3.4 - 5.8)
Income								
≤ \$24,999	92	3,300	10.3	(3.8 - 16.9)	9,507	372,300	10.7	(7.3 - 14.1)
\$25,000-\$49,999	*	*	*	*	9,585	136,300	4.7	(3.8 - 5.7)
\$50,000-\$74,999	*	*	*	*	5,755	154,600	7.8	(4.9 - 10.7)
≥ \$75,000	269	2,400	2.4	(1.0 - 3.9)	8,541	151,800	3.4	(2.3 - 4.5)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having ever been told by a health professional that respondent had a heart attack or angina

<sup>&</sup>lt;sup>2</sup> Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults with coronary heart disease (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 45b. Coronary heart disease among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	Ro	ockland C	County	N	ew York	State
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	655	4.6	(3.2 - 6.1)	37,838	6.3	(5.2 - 7.3)
Gender						
Male	244	6.3	(3.7 - 9.0)	14,479	8.8	(6.9 - 10.8)
Female	411	3.0	(1.7 - 4.4)	23,359	3.9	(3.0 - 4.8)
Race/Ethnicity						
White Non-Hispanic	548	5.6	(3.8 - 7.4)	35,552	6.8	(5.9 - 7.7)
Black Non-Hispanic	*	*	*	728	5.6	(2.2 - 9.0)
Hispanic	*	*	*	853	5.4	(1.1 - 9.6)
Other Non-Hispanic	*	*	*	705	4.8	(0.4 - 9.2)
Education						
≤ High School	199	6.4	(3.4 - 9.4)	15,485	8.5	(6.2 - 10.7)
Some College	*	*	*	9,860	5.5	(3.7 - 7.2)
≥ College Degree	336	3.9	(2.1 - 5.7)	12,411	4.7	(3.4 - 5.9)
Income						
≤ \$24,999	92	9.5	(3.3 - 15.7)	9,507	10.7	(7.2 - 14.1)
\$25,000-\$49,999	*	*	*	9,585	4.7	(3.6 - 5.7)
\$50,000-\$74,999	*	*	*	5,755	8.0	(4.8 - 11.1)
≥ \$75,000 Z	269	2.2	(0.9 - 3.5)	8,541	3.3	(2.1 - 4.5)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having ever been told by a health professional that respondent had a heart attack or angina

 $<sup>^{2}</sup>$  Prevalence estimates are age-adjusted to the 2000 US Standard population  $\,$ 

 $<sup>^{3}</sup>$  Sample size from survey

 $<sup>^4</sup>$  95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 46a. Ever diagnosed with diabetes among adults<sup>1</sup> in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	657	18,900	8.7	(6.4 - 11.0)	38,174	1,382,700	9.3	(8.0 - 10.5)
Gender								
Male	243	9,500	9.0	(5.6 - 12.4)	14,626	530,400	7.5	(6.2 - 8.8)
Female	414	9,400	8.4	(5.4 - 11.4)	23,548	852,300	11.0	(8.9 - 13.0)
Age (years)								
18-34	*	*	*	*	3,941	150,200	3.4	(1.0 - 5.8)
35-44	*	*	*	*	4,987	165,100	5.7	(3.6 - 7.9)
45-54	*	*	*	*	7,830	279,000	9.8	(6.7 - 12.8)
55-64	135	5,600	15.3	(7.9 - 22.6)	8,771	332,900	15.5	(12.3 - 18.8)
≥ 65	217	7,300	18.9	(13.2 - 24.6)	12,645	455,600	17.9	(15.0 - 20.8)
Race/Ethnicity								
White Non-Hispanic	550	14,300	9.5	(6.9 - 12.1)	35,862	809,300	8.7	(7.8 - 9.5)
Black Non-Hispanic	*	*	*	*	739	261,000	12.1	(6.6 - 17.6)
Hispanic	*	*	*	*	862	227,300	10.1	(5.4 - 14.8)
Other Non-Hispanic	*	*	*	*	711	85,100	7.6	(2.3 - 12.8)
Education								
≤ High School	199	8,800	13.3	(7.9 - 18.7)	15,709	730,000	13.0	(10.5 - 15.6)
Some College	115	2,600	6.4	(2.2 - 10.7)	9,918	303,500	8.6	(5.9 - 11.2)
≥ College Degree	339	7,500	6.9	(4.2 - 9.6)	12,464	337,500	6.0	(4.8 - 7.2)
Income								
≤ \$24,999	91	2,000	6.4	(2.1 - 10.7)	9,652	543,400	15.5	(11.4 - 19.5)
\$25,000-\$49,999	110	5,800	17.1	(8.2 - 25.9)	9,667	291,000	10.0	(7.5 - 12.6)
\$50,000-\$74,999	*	*	*	*	5,777	151,000	7.6	(5.0 - 10.3)
≥ \$75,000	270	8,100	8.1	(4.9 - 11.4)	8,565	220,800	4.9	(3.8 - 6.1)

 $<sup>^{1}</sup>$  Defined as having ever been told by a doctor that respondent had diabetes, excluding pre-diabetes and women with diabetes only when  $\begin{array}{c} \text{pregnant} \\ ^2 \text{ Sample size from survey} \end{array}$ 

 $<sup>^{3}</sup>$  Estimated number of adults who have diabetes (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 46b. Ever diagnosed with diabetes among adults in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland (	County	N	New York State				
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>			
Total	657	8.0	(5.8 - 10.1)	38,174	9.0	(7.8 - 10.3)			
Gender									
Male	243	8.2	(5.1 - 11.3)	14,626	7.5	(6.1 - 8.9)			
Female	414	7.7	(4.7 - 10.7)	23,548	10.4	(8.4 - 12.4)			
Race/Ethnicity									
White Non-Hispanic	550	8.6	(6.2 - 11.0)	35,862	8.4	(7.5 - 9.2)			
Black Non-Hispanic	*	*	*	739	11.4	(6.0 - 16.7)			
Hispanic	*	*	*	862	10.1	(5.4 - 14.9)			
Other Non-Hispanic	*	*	*	711	7.7	(2.3 - 13.0)			
Education									
≤ High School	199	12.2	(7.1 - 17.4)	15,709	12.6	(10.0 - 15.2)			
Some College	115	5.4	(1.6 - 9.3)	9,918	8.3	(5.6 - 10.9)			
≥ College Degree	339	6.5	(3.9 - 9.1)	12,464	5.9	(4.7 - 7.2)			
Income									
≤ \$24,999	91	5.2	(1.7 - 8.7)	9,652	15.2	(11.1 - 19.2)			
\$25,000-\$49,999	110	15.8	(7.1 - 24.5)	9,667	9.7	(7.1 - 12.2)			
\$50,000-\$74,999	*	*	*	5,777	7.6	(4.9 - 10.2)			
≥ \$75,000	270	7.5	(4.5 - 10.5)	8,565	4.7	(3.6 - 5.9)			

<sup>&</sup>lt;sup>1</sup> Defined as having ever been told by a doctor that respondent had diabetes, excluding pre-diabetes and women with diabetes only when  $\begin{array}{l} \text{pregnant} \\ \text{^{2}} \text{ Prevalence estimates are age-adjusted to the 2000 US Standard population} \end{array}$ 

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 47a. Ever diagnosed with asthma (lifetime) among adults<sup>1</sup> in Rockland County and New York State

		Rockla	nd Count	ty		New Y	ork State	9
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	657	21,800	10.1	(7.2 - 13.1)	38,096	2,225,100	15.0	(13.2 - 16.8)
Gender								
Male	244	7,700	7.3	(3.6 - 11.0)	14,601	875,800	12.4	(10.1 - 14.7)
Female	413	14,200	12.8	(8.3 - 17.3)	23,495	1,349,300	17.4	(14.6 - 20.1)
Age (years)								
18-34	*	*	*	*	3,931	825,100	18.5	(13.8 - 23.3)
35-44	92	3,500	9.1	(2.9 - 15.3)	4,972	431,700	15.1	(11.2 - 19.1)
45-54	131	3,900	8.9	(3.9 - 13.8)	7,819	365,900	12.8	(9.7 - 16.0)
55-64	135	3,100	8.4	(1.6 - 15.3)	8,756	316,700	14.8	(11.9 - 17.7)
≥ 65	218	5,000	12.8	(7.4 - 18.3)	12,618	285,700	11.2	(8.6 - 13.8)
Race/Ethnicity								
White Non-Hispanic	550	14,600	9.8	(6.7 - 12.8)	35,783	1,365,700	14.6	(13.0 - 16.3)
Black Non-Hispanic	*	*	*	*	737	367,000	17.2	(10.2 - 24.2)
Hispanic	*	*	*	*	863	317,100	14.1	(9.0 - 19.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	198	5,100	7.9	(3.4 - 12.4)	15,675	831,600	14.9	(12.0 - 17.8)
Some College	116	6,400	15.8	(6.6 - 25.0)	9,893	587,500	16.6	(13.1 - 20.1)
≥ College Degree	339	10,300	9.5	(5.7 - 13.3)	12,444	739,000	13.2	(10.4 - 15.9)
Income								
≤ \$24,999	92	2,200	6.8	(2.2 - 11.4)	9,643	708,900	20.2	(14.9 - 25.5)
\$25,000-\$49,999	*	*	*	*	9,641	389,800	13.4	(10.0 - 16.9)
\$50,000-\$74,999	*	*	*	*	5,762	257,800	13.2	(10.0 - 16.5)
≥ \$75,000	270	10,900	11.0	(6.4 - 15.6)	8,562	577,900	12.8	(10.4 - 15.3)

 $<sup>^{\</sup>mathrm{1}}$  Defined as adults having ever been told by a health professional that they had asthma

 $<sup>^{2}\,\</sup>mathrm{Sample}$  size from survey

 $<sup>^{3}</sup>$  Estimated number of adults who have lifetime asthma (rounded to the nearest hundred)

<sup>4 95%</sup> confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 47b. Ever diagnosed with asthma (lifetime) among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland (	County	N	ew York	State
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	657	10.1	(7.1 - 13.2)	38,096	15.2	(13.3 - 17.0)
Gender						
Male	244	7.4	(3.5 - 11.3)	14,601	12.6	(10.3 - 14.9)
Female	413	12.8	(8.2 - 17.4)	23,495	17.5	(14.8 - 20.3)
Race/Ethnicity						
White Non-Hispanic	550	10.1	(6.7 - 13.4)	35,783	14.9	(13.2 - 16.6)
Black Non-Hispanic	*	*	*	737	17.1	(10.1 - 24.1)
Hispanic	*	*	*	863	14.5	(9.3 - 19.7)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	198	8.0	(3.2 - 12.8)	15,675	15.1	(12.1 - 18.0)
Some College	116	16.3	(6.5 - 26.1)	9,893	16.9	(13.4 - 20.4)
≥ College Degree	339	9.2	(5.4 - 13.0)	12,444	13.3	(10.6 - 16.0)
Income						
≤ \$24,999	92	6.3	(1.8 - 10.7)	9,643	20.3	(15.0 - 25.6)
\$25,000-\$49,999	*	*	*	9,641	13.7	(10.2 - 17.2)
\$50,000-\$74,999	*	*	*	5,762	13.5	(10.0 - 17.0)
≥ \$75,000	270	10.8	(6.2 - 15.4)	8,562	13.0	(10.6 - 15.4)

 $<sup>^{\</sup>rm 1}$  Defined as adults having ever been told by a health professional that they had asthma

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{\</sup>rm 3}$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 48a. Current asthma among adults in Rockland County and New York State

		Rockla	ind Count	ty .		New Y	ork State	•
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	653	15,500	7.2	(4.6 - 9.8)	37,933	1,415,000	9.6	(8.1 - 11.0)
Gender								
Male	243	5,100	4.9	(1.8 - 7.9)	14,546	440,700	6.3	(4.8 - 7.7)
Female	410	10,400	9.4	(5.3 - 13.5)	23,387	974,300	12.6	(10.2 - 15.0)
Age (years)								
18-34	*	*	*	*	3,909	467,200	10.5	(6.8 - 14.3)
35-44	*	*	*	*	4,947	246,800	8.7	(5.8 - 11.6)
45-54	131	3,100	7.0	(2.5 - 11.6)	7,787	240,000	8.4	(6.1 - 10.8)
55-64	*	*	*	*	8,726	231,100	10.8	(8.1 - 13.5)
≥ 65	216	3,400	8.8	(3.9 - 13.7)	12,564	229,900	9.1	(6.5 - 11.6)
Race/Ethnicity								
White Non-Hispanic	546	8,800	5.9	(3.5 - 8.3)	35,629	888,400	9.6	(8.2 - 11.0)
Black Non-Hispanic	*	*	*	*	736	246,900	11.6	(5.7 - 17.5)
Hispanic	*	*	*	*	859	186,800	8.3	(4.4 - 12.2)
Other Non-Hispanic	*	*	*	*	709	93,000	8.3	(1.4 - 15.2)
Education								
≤ High School	197	3,500	5.5	(1.5 - 9.4)	15,594	595,100	10.7	(8.2 - 13.2)
Some College	116	5,100	12.5	(3.9 - 21.1)	9,861	322,000	9.1	(7.0 - 11.3)
≥ College Degree	336	6,900	6.3	(3.1 - 9.6)	12,394	430,900	7.7	(5.8 - 9.7)
Income								
≤ \$24,999	*	*	*	*	9,600	489,200	14.0	(9.7 - 18.3)
\$25,000-\$49,999	*	*	*	*	9,590	268,400	9.3	(6.2 - 12.4)
\$50,000-\$74,999	*	*	*	*	5,741	143,400	7.4	(5.1 - 9.6)
≥ \$75,000	270	8,500	8.5	(4.4 - 12.6)	8,541	339,200	7.6	(5.8 - 9.4)

 $<sup>^{\</sup>mathrm{1}}$  Defined as adults having ever been told by a health professional that they had asthma and still have asthma

<sup>&</sup>lt;sup>2</sup> Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults who currently have asthma (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 48b. Current asthma among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland (	County	N	ew York	State
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>
Total	653	7.1	(4.5 - 9.8)	37,933	9.7	(8.2 - 11.2)
Gender						
Male	243	5.0	(1.7 - 8.2)	14,546	6.4	(4.9 - 7.9)
Female	410	9.2	(5.0 - 13.3)	23,387	12.7	(10.2 - 15.1)
Race/Ethnicity						
White Non-Hispanic	546	6.1	(3.5 - 8.7)	35,629	9.7	(8.3 - 11.2)
Black Non-Hispanic	*	*	*	736	11.5	(5.6 - 17.3)
Hispanic	*	*	*	859	8.6	(4.5 - 12.6)
Other Non-Hispanic	*	*	*	709	8.3	(1.3 - 15.3)
Education						
≤ High School	197	5.4	(1.3 - 9.5)	15,594	10.8	(8.3 - 13.3)
Some College	116	12.9	(3.8 - 22.0)	9,861	9.4	(7.2 - 11.6)
≥ College Degree	336	6.1	(3.0 - 9.2)	12,394	7.8	(5.8 - 9.7)
Income						
≤ \$24,999	*	*	*	9,600	14.1	(9.7 - 18.4)
\$25,000-\$49,999	*	*	*	9,590	9.5	(6.3 - 12.7)
\$50,000-\$74,999	*	*	*	5,741	7.5	(5.0 - 10.1)
≥ \$75,000	270	8.3	(4.3 - 12.3)	8,541	7.7	(5.9 - 9.5)

 $<sup>^{\</sup>mathrm{1}}$  Defined as adults having ever been told by a health professional that they had asthma and still have asthma

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 49a. Chronic joint symptoms among adults in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	•
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	656	77,700	35.9	(31.2 - 40.6)	38,099	6,377,100	42.9	(40.5 - 45.3)
Gender								
Male	242	38,300	36.7	(29.1 - 44.3)	14,595	3,033,600	42.8	(39.0 - 46.6)
Female	414	39,400	35.2	(29.5 - 40.9)	23,504	3,343,500	43.0	(40.0 - 46.0)
Age (years)								
18-34	82	9,300	15.4	(7.0 - 23.7)	3,933	1,261,300	28.3	(22.6 - 34.0)
35-44	*	*	*	*	4,984	1,110,900	38.6	(33.1 - 44.0)
45-54	*	*	*	*	7,817	1,456,500	51.2	(46.4 - 55.9)
55-64	*	*	*	*	8,751	1,167,400	54.6	(50.4 - 58.8)
≥ 65	218	21,600	55.5	(48.2 - 62.8)	12,614	1,381,000	54.3	(50.5 - 58.1)
Race/Ethnicity								
White Non-Hispanic	550	58,800	39.2	(34.1 - 44.3)	35,785	4,267,600	45.8	(43.8 - 47.7)
Black Non-Hispanic	*	*	*	*	740	890,500	41.3	(32.5 - 50.0)
Hispanic	*	*	*	*	860	827,500	36.7	(27.7 - 45.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	198	20,200	30.8	(23.0 - 38.7)	15,676	2,450,000	43.8	(39.4 - 48.1)
Some College	*	*	*	*	9,900	1,598,300	45.1	(40.1 - 50.1)
≥ College Degree	338	38,600	35.8	(29.0 - 42.5)	12,440	2,313,100	41.1	(37.8 - 44.5)
Income								
≤ \$24,999	*	*	*	*	9,641	1,483,000	42.2	(36.3 - 48.1)
\$25,000-\$49,999	*	*	*	*	9,647	1,332,100	45.9	(39.8 - 52.1)
\$50,000-\$74,999	*	*	*	*	5,768	894,200	45.3	(39.5 - 51.1)
≥ \$75,000	269	33,400	33.9	(27.0 - 40.8)	8,549	1,874,200	41.7	(38.4 - 45.1)

 $<sup>^1</sup>$  Defined as adults having symptoms of pain, aching or stiffness in or around a joint during the past 30 days and began more than 3 months ago

<sup>&</sup>lt;sup>2</sup> Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of adults who have chronic joint symptoms (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 49b. Chronic joint symptoms among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	Re	ockland	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	656	34.5	(29.6 - 39.4)	38,099	42.1	(39.7 - 44.5)		
Gender								
Male	242	36.2	(28.3 - 44.0)	14,595	42.6	(38.8 - 46.3)		
Female	414	32.9	(27.0 - 38.8)	23,504	41.6	(38.6 - 44.7)		
Race/Ethnicity								
White Non-Hispanic	550	37.2	(31.9 - 42.6)	35,785	44.7	(42.7 - 46.6)		
Black Non-Hispanic	*	*	*	740	40.2	(31.5 - 48.9)		
Hispanic	*	*	*	860	36.6	(27.8 - 45.4)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	198	28.8	(21.1 - 36.5)	15,676	43.1	(38.8 - 47.5)		
Some College	*	*	*	9,900	43.9	(39.0 - 48.8)		
≥ College Degree	338	34.6	(27.5 - 41.6)	12,440	40.3	(37.0 - 43.7)		
Income								
≤ \$24,999	*	*	*	9,641	41.6	(35.7 - 47.4)		
\$25,000-\$49,999	*	*	*	9,647	45.0	(39.0 - 51.0)		
\$50,000-\$74,999	*	*	*	5,768	44.3	(38.4 - 50.2)		
≥ \$75,000	269	32.9	(25.8 - 40.1)	8,549	40.8	(37.4 - 44.2)		

 $<sup>^{1}</sup>$  Defined as adults having symptoms of pain, aching or stiffness in or around a joint during the past 30 days and began more than 3 months ago

Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 50a. Arthritis among adults<sup>1</sup> in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	e
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	652	53,100	24.6	(20.9 - 28.4)	38,013	4,102,800	27.7	(25.9 - 29.6)
Gender								
Male	241	22,700	21.8	(16.0 - 27.6)	14,555	1,642,700	23.3	(20.7 - 25.8)
Female	411	30,300	27.3	(22.5 - 32.1)	23,458	2,460,100	31.8	(29.2 - 34.4)
Age (years)								
18-34	*	*	*	*	3,927	312,100	7.0	(4.9 - 9.1)
35-44	*	*	*	*	4,972	412,500	14.5	(11.3 - 17.6)
45-54	130	8,300	19.2	(12.2 - 26.1)	7,789	869,100	30.7	(26.6 - 34.8)
55-64	*	*	*	*	8,729	999,300	47.0	(42.9 - 51.1)
≥ 65	216	22,900	59.7	(52.6 - 66.9)	12,596	1,509,800	59.6	(55.8 - 63.3)
Race/Ethnicity								
White Non-Hispanic	547	42,000	28.1	(23.8 - 32.3)	35,710	2,967,700	31.9	(30.3 - 33.6)
Black Non-Hispanic	*	*	*	*	737	588,100	27.3	(20.2 - 34.4)
Hispanic	*	*	*	*	856	380,700	17.1	(11.8 - 22.5)
Other Non-Hispanic	*	*	*	*	710	166,300	14.8	(7.8 - 21.8)
Education								
≤ High School	198	15,900	24.2	(17.1 - 31.3)	15,652	1,742,700	31.3	(27.8 - 34.8)
Some College	114	8,800	21.9	(14.0 - 29.8)	9,874	902,000	25.5	(22.3 - 28.6)
≥ College Degree	336	28,000	25.9	(20.5 - 31.3)	12,407	1,437,800	25.6	(22.9 - 28.4)
Income								
≤ \$24,999	91	7,800	24.5	(14.6 - 34.4)	9,618	1,112,500	32.0	(27.1 - 36.9)
\$25,000-\$49,999	110	8,600	25.1	(16.1 - 34.2)	9,624	821,900	28.4	(24.2 - 32.6)
\$50,000-\$74,999	*	*	*	*	5,753	542,500	27.5	(23.2 - 31.9)
≥ \$75,000	268	21,800	22.1	(16.4 - 27.9)	8,536	1,012,800	22.6	(19.8 - 25.3)

 $<sup>^{1}</sup>$  Defined as adults having ever been told by a health professional that they have some form of arthritis, rheumatoid arthritis, gout, lupus or fibromyalgia (excluding back or neck)  $^{2}$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults who have ever been told by a health professional they have some form of arthritis (excluding back or neck) (rounded to the nearest hundred)
4 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 50b. Arthritis among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	652	22.0	(18.4 - 25.6)	38,013	26.6	(24.8 - 28.5)		
Gender								
Male	241	20.2	(14.7 - 25.8)	14,555	23.5	(20.9 - 26.1)		
Female	411	23.6	(19.0 - 28.3)	23,458	29.5	(27.0 - 32.1)		
Race/Ethnicity								
White Non-Hispanic	547	25.0	(20.9 - 29.2)	35,710	30.4	(28.7 - 32.0)		
Black Non-Hispanic	*	*	*	737	26.5	(19.3 - 33.6)		
Hispanic	*	*	*	856	17.2	(11.8 - 22.6)		
Other Non-Hispanic	*	*	*	710	14.9	(7.7 - 22.0)		
Education								
≤ High School	198	21.6	(15.2 - 28.1)	15,652	30.4	(26.9 - 33.8)		
Some College	114	18.6	(11.4 - 25.8)	9,874	24.1	(21.0 - 27.1)		
≥ College Degree	336	23.6	(18.2 - 28.9)	12,407	24.7	(21.9 - 27.5)		
Income								
≤ \$24,999	91	22.0	(12.5 - 31.4)	9,618	31.2	(26.3 - 36.1)		
\$25,000-\$49,999	110	22.9	(14.3 - 31.5)	9,624	27.3	(23.2 - 31.3)		
\$50,000-\$74,999	*	*	*	5,753	26.3	(21.9 - 30.7)		
≥ \$75,000	268	19.3	(13.9 - 24.7)	8,536	21.6	(18.8 - 24.3)		

 $<sup>^{1}</sup>$  Defined as adults having ever been told by a health professional that they have some form of arthritis, rheumatoid arthritis, gout, lupus or fibromyalgia (excluding back or neck)  $^{2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 51a. Activity limitations because of physical, mental or emotional problems among adults<sup>1</sup> in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	•
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	653	33,700	15.6	(12.5 - 18.8)	37,998	3,053,000	20.6	(18.7 - 22.5)
Gender								
Male	242	15,800	15.1	(10.3 - 19.9)	14,566	1,318,400	18.6	(15.7 - 21.5)
Female	411	17,900	16.1	(12.1 - 20.1)	23,432	1,734,600	22.4	(19.8 - 25.0)
Age (years)								
18-34	*	*	*	*	3,929	580,000	13.0	(8.2 - 17.9)
35-44	*	*	*	*	4,973	418,600	14.5	(11.0 - 18.1)
45-54	130	10,300	23.8	(15.1 - 32.6)	7,812	620,200	21.7	(18.3 - 25.2)
55-64	134	7,500	20.7	(12.8 - 28.5)	8,739	651,600	30.5	(26.5 - 34.6)
≥ 65	216	10,900	28.5	(21.7 - 35.3)	12,545	782,700	31.1	(27.7 - 34.6)
Race/Ethnicity								
White Non-Hispanic	546	23,900	16.0	(12.7 - 19.4)	35,688	2,037,200	21.9	(20.3 - 23.5)
Black Non-Hispanic	*	*	*	*	735	429,700	20.1	(13.5 - 26.7)
Hispanic	*	*	*	*	863	448,200	19.8	(11.9 - 27.8)
Other Non-Hispanic	*	*	*	*	712	137,900	12.3	(5.2 - 19.3)
Education								
≤ High School	198	10,700	16.3	(10.3 - 22.2)	15,626	1,359,600	24.4	(20.6 - 28.1)
Some College	114	7,400	18.5	(10.4 - 26.5)	9,875	700,800	19.8	(16.5 - 23.2)
≥ College Degree	337	15,600	14.5	(10.3 - 18.6)	12,415	924,800	16.5	(14.2 - 18.7)
Income								
≤ \$24,999	*	*	*	*	9,603	1,055,800	30.1	(24.7 - 35.5)
\$25,000-\$49,999	109	6,400	18.8	(10.0 - 27.7)	9,628	630,900	21.8	(16.6 - 27.0)
\$50,000-\$74,999	77	3,100	15.2	(6.3 - 24.1)	5,760	408,100	20.7	(16.1 - 25.2)
≥ \$75,000	269	11,300	11.4	(7.4 - 15.4)	8,544	583,300	13.0	(11.1 - 14.9)

 $<sup>^{1}</sup>$  Defined as activity limitations because of physical, mental or emotional problems

 $<sup>^{2}% \</sup>left( -1\right) =0$  Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults who are limited in any activities because of physical, mental or emotional problems (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 51b. Activity limitations because of physical, mental or emotional problems among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	653	14.2	(11.2 - 17.2)	37,998	20.1	(18.2 - 22.0)		
Gender								
Male	242	14.0	(9.4 - 18.6)	14,566	18.6	(15.8 - 21.4)		
Female	411	14.4	(10.5 - 18.3)	23,432	21.5	(18.9 - 24.1)		
Race/Ethnicity								
White Non-Hispanic	546	14.5	(11.3 - 17.8)	35,688	21.3	(19.7 - 22.9)		
Black Non-Hispanic	*	*	*	735	19.9	(13.2 - 26.5)		
Hispanic	*	*	*	863	19.1	(11.5 - 26.8)		
Other Non-Hispanic	*	*	*	712	12.4	(5.3 - 19.6)		
Education								
≤ High School	198	15.0	(9.4 - 20.6)	15,626	23.8	(20.1 - 27.4)		
Some College	114	15.8	(8.3 - 23.2)	9,875	19.3	(16.0 - 22.6)		
≥ College Degree	337	13.4	(9.3 - 17.5)	12,415	16.1	(13.9 - 18.3)		
Income								
≤ \$24,999	92	19.1	(9.7 - 28.5)	9,603	29.7	(24.3 - 35.2)		
\$25,000-\$49,999	109	17.7	(9.0 - 26.3)	9,628	21.2	(16.1 - 26.3)		
\$50,000-\$74,999	*	*	*	5,760	20.1	(15.6 - 24.6)		
≥ \$75,000	269	10.1	(6.4 - 13.8)	8,544	12.7	(10.8 - 14.6)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as activity limitations because of physical, mental or emotional problems

 $<sup>^{2}</sup>$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 52a. Have health problem that requires the use of special equipment among adults in Rockland County and New York State

		Rockla	nd Coun	ty		New Y	ork State	9
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	658	13,000	6.0	(4.2 - 7.8)	38,171	1,169,400	7.9	(6.9 - 8.9)
Gender								
Male	244	5,700	5.4	(2.5 - 8.3)	14,617	460,700	6.5	(5.3 - 7.7)
Female	414	7,300	6.5	(4.3 - 8.8)	23,554	708,700	9.1	(7.6 - 10.7)
Age (years)								
18-34	*	*	*	*	3,940	68,800	1.5	(0.7 - 2.4)
35-44	*	*	*	*	4,989	122,500	4.3	(2.4 - 6.1)
45-54	131	4,900	11.3	(4.8 - 17.7)	7,836	182,500	6.4	(4.7 - 8.1)
55-64	*	*	*	*	8,773	219,600	10.3	(7.8 - 12.8)
≥ 65	218	7,100	18.1	(12.4 - 23.8)	12,633	576,100	22.7	(19.0 - 26.3)
Race/Ethnicity								
White Non-Hispanic	551	8,800	5.8	(4.0 - 7.6)	35,853	802,000	8.6	(7.6 - 9.6)
Black Non-Hispanic	*	*	*	*	739	235,800	10.9	(6.4 - 15.5)
Hispanic	*	*	*	*	864	104,100	4.6	(2.1 - 7.1)
Other Non-Hispanic	*	*	*	*	715	27,600	2.5	(0.8 - 4.1)
Education								
≤ High School	199	6,700	10.1	(5.6 - 14.6)	15,711	604,200	10.8	(8.7 - 12.8)
Some College	116	2,500	6.2	(1.6 - 10.9)	9,914	260,000	7.3	(5.5 - 9.1)
≥ College Degree	339	3,800	3.5	(1.9 - 5.2)	12,463	298,900	5.3	(4.1 - 6.5)
Income								
≤ \$24,999	92	4,400	13.8	(5.2 - 22.5)	9,656	430,500	12.2	(9.6 - 14.8)
\$25,000-\$49,999	110	2,600	7.7	(2.6 - 12.8)	9,663	238,900	8.2	(5.9 - 10.5)
\$50,000-\$74,999	*	*	*	*	5,775	103,000	5.2	(3.6 - 6.8)
≥ \$75,000	*	*	*	*	8,566	136,000	3.0	(2.1 - 4.0)

 $<sup>^{1}</sup>$  Defined as having any health problem that requires the use of special equipment such as a cane, a wheelchair, a special bed, or a special telephone

2 Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults who have any health problem that requires the use of special equipment (rounded to the nearest hundred)

 $<sup>^4</sup>$  95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 52b. Have health problem that requires the use of special equipment among adults 1 in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland (	County	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	658	5.4	(3.7 - 7.1)	38,171	7.6	(6.6 - 8.6)		
Gender								
Male	244	5.2	(2.4 - 8.0)	14,617	6.8	(5.5 - 8.0)		
Female	414	5.6	(3.6 - 7.5)	23,554	8.4	(6.9 - 9.9)		
Race/Ethnicity								
White Non-Hispanic	551	5.3	(3.6 - 6.9)	35,853	8.3	(7.3 - 9.3)		
Black Non-Hispanic	*	*	*	739	10.5	(6.1 - 15.0)		
Hispanic	*	*	*	864	4.5	(2.1 - 6.9)		
Other Non-Hispanic	*	*	*	715	2.2	(0.7 - 3.8)		
Education								
≤ High School	199	9.4	(5.1 - 13.7)	15,711	10.3	(8.3 - 12.3)		
Some College	116	5.5	(1.3 - 9.7)	9,914	7.1	(5.2 - 8.9)		
≥ College Degree	339	3.0	(1.6 - 4.5)	12,463	5.3	(4.1 - 6.6)		
Income								
≤ \$24,999	92	12.7	(4.6 - 20.8)	9,656	11.9	(9.4 - 14.5)		
\$25,000-\$49,999	110	6.9	(2.3 - 11.6)	9,663	7.7	(5.5 - 10.0)		
\$50,000-\$74,999	*	*	*	5,775	5.1	(3.4 - 6.8)		
≥ \$75,000	*	*	*	8,566	2.9	(2.0 - 3.9)		

 $<sup>^{1}</sup>$  Defined as having any health problem that requires the use of special equipment such as a cane, a wheelchair, a special bed, or a special telephone  $^2$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 53a. Disability among adults<sup>1</sup> in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	e
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	653	36,400	16.9	(13.7 - 20.1)	38,016	3,394,200	22.9	(20.9 - 24.9)
Gender								
Male	242	16,700	15.9	(11.0 - 20.8)	14,568	1,469,700	20.7	(17.8 - 23.7)
Female	411	19,800	17.8	(13.7 - 22.0)	23,448	1,924,600	24.8	(22.1 - 27.5)
Age (years)								
18-34	*	*	*	*	3,931	590,300	13.3	(8.4 - 18.1)
35-44	*	*	*	*	4,974	447,900	15.6	(11.9 - 19.3)
45-54	130	10,300	23.8	(15.1 - 32.6)	7,811	643,900	22.6	(19.1 - 26.0)
55-64	134	7,600	20.9	(13.1 - 28.8)	8,740	702,100	33.0	(28.9 - 37.0)
≥ 65	216	13,500	35.4	(28.3 - 42.6)	12,560	1,010,000	39.9	(36.1 - 43.7)
Race/Ethnicity								
White Non-Hispanic	546	26,300	17.6	(14.2 - 21.1)	35,703	2,253,400	24.2	(22.6 - 25.9)
Black Non-Hispanic	*	*	*	*	736	511,700	23.8	(16.8 - 30.8)
Hispanic	*	*	*	*	864	484,300	21.4	(13.4 - 29.5)
Other Non-Hispanic	*	*	*	*	713	144,800	12.9	(5.8 - 19.9)
Education								
≤ High School	198	12,300	18.7	(12.5 - 25.0)	15,638	1,544,200	27.6	(23.7 - 31.4)
Some College	114	7,500	18.7	(10.7 - 26.8)	9,876	755,200	21.4	(17.9 - 24.8)
≥ College Degree	337	16,600	15.4	(11.2 - 19.6)	12,419	1,021,300	18.2	(15.9 - 20.5)
Income								
≤ \$24,999	*	*	*	*	9,613	1,179,100	33.6	(28.0 - 39.2)
\$25,000-\$49,999	109	7,100	21.0	(11.8 - 30.1)	9,629	699,000	24.2	(18.9 - 29.4)
\$50,000-\$74,999	77	3,600	17.3	(8.0 - 26.7)	5,757	430,100	21.8	(17.2 - 26.4)
≥ \$75,000	269	11,300	11.4	(7.4 - 15.4)	8,546	632,700	14.1	(12.1 - 16.1)

 $<sup>^{1}</sup>$  Defined as having activity limitations due to physical, mental or emotional problems or have health problems that require the use of special equipment

2 Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of adults with a disability (rounded to the nearest hundred)

 $<sup>^4</sup>$  95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 53b. Disability among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	653	15.4	(12.3 - 18.5)	38,016	22.3	(20.4 - 24.3)		
Gender								
Male	242	14.9	(10.2 - 19.6)	14,568	20.9	(18.0 - 23.8)		
Female	411	15.8	(11.8 - 19.8)	23,448	23.6	(21.0 - 26.3)		
Race/Ethnicity								
White Non-Hispanic	546	16.0	(12.7 - 19.4)	35,703	23.6	(21.9 - 25.3)		
Black Non-Hispanic	*	*	*	736	23.5	(16.5 - 30.5)		
Hispanic	*	*	*	864	20.6	(12.9 - 28.3)		
Other Non-Hispanic	*	*	*	713	12.9	(5.8 - 20.1)		
Education								
≤ High School	198	17.4	(11.5 - 23.3)	15,638	26.8	(23.0 - 30.6)		
Some College	114	16.1	(8.6 - 23.5)	9,876	20.9	(17.4 - 24.3)		
≥ College Degree	337	14.2	(10.0 - 18.3)	12,419	17.9	(15.5 - 20.3)		
Income								
≤ \$24,999	92	21.0	(11.2 - 30.8)	9,613	33.2	(27.6 - 38.8)		
\$25,000-\$49,999	109	19.5	(10.6 - 28.4)	9,629	23.5	(18.3 - 28.6)		
\$50,000-\$74,999	*	*	*	5,757	21.2	(16.6 - 25.8)		
≥ \$75,000	269	10.1	(6.4 - 13.8)	8,546	13.8	(11.7 - 15.8)		

<sup>1</sup> Defined as having activity limitations due to physical, mental or emotional problems or have health problems that require the use of special equipment <sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 54a. Current smoking among adults<sup>1</sup> in Rockland County and New York State

		Rockla	ind Count	<b>Y</b>		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	657	20,700	9.5	(6.6 - 12.5)	38,029	2,483,500	16.8	(15.1 - 18.6)
Gender								
Male	244	11,300	10.7	(5.6 - 15.8)	14,559	1,356,400	19.3	(16.3 - 22.2)
Female	413	9,400	8.4	(5.2 - 11.5)	23,470	1,127,100	14.6	(12.7 - 16.6)
Age (years)								
18-34	*	*	*	*	3,929	993,500	22.3	(17.5 - 27.1)
35-44	92	3,400	9.0	(3.2 - 14.8)	4,977	421,100	14.8	(11.6 - 17.9)
45-54	130	4,400	10.1	(4.6 - 15.5)	7,807	508,800	18.3	(15.4 - 21.3)
55-64	135	1,600	4.4	(1.5 - 7.4)	8,744	314,500	14.7	(12.1 - 17.3)
≥ 65	218	3,800	9.7	(4.8 - 14.6)	12,572	245,600	9.7	(7.2 - 12.3)
Race/Ethnicity								
White Non-Hispanic	550	14,800	9.8	(6.2 - 13.4)	35,729	1,615,500	17.3	(15.9 - 18.8)
Black Non-Hispanic	*	*	*	*	735	284,100	13.6	(8.9 - 18.2)
Hispanic	*	*	*	*	858	350,900	15.8	(9.6 - 21.9)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	199	7,500	11.4	(4.9 - 17.9)	15,654	1,237,300	22.4	(19.1 - 25.6)
Some College	116	7,100	17.4	(8.8 - 26.0)	9,875	670,600	19.1	(15.5 - 22.8)
≥ College Degree	338	6,100	5.6	(2.7 - 8.5)	12,421	563,300	10.0	(7.5 - 12.5)
Income								
≤ \$24,999	92	4,600	14.5	(6.4 - 22.6)	9,626	835,100	24.1	(19.1 - 29.1)
\$25,000-\$49,999	110	3,600	10.7	(4.5 - 16.9)	9,629	558,100	19.4	(15.6 - 23.2)
\$50,000-\$74,999	*	*	*	*	5,756	338,100	17.3	(13.0 - 21.7)
≥ \$75,000	270	9,000	9.1	(3.9 - 14.3)	8,549	468,700	10.4	(8.6 - 12.3)

 $<sup>^{</sup>m 1}$  Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day or some days

 $<sup>^{2} \, \</sup>mathrm{Sample} \, \mathrm{size} \, \mathrm{from} \, \mathrm{survey}$ 

 $<sup>^{3}</sup>$  Estimated number of adult current smokers (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 54b. Current smoking among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	<b>Rockland County</b>			N	New York State		
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	
Total	657	9.7	(6.5 - 12.8)	38,029	17.0	(15.3 - 18.8)	
Gender							
Male	244	11.2	(5.8 - 16.6)	14,559	19.2	(16.3 - 22.1)	
Female	413	8.2	(4.8 - 11.6)	23,470	15.0	(13.0 - 17.0)	
Race/Ethnicity							
White Non-Hispanic	550	10.0	(6.1 - 14.0)	35,729	17.7	(16.2 - 19.2)	
Black Non-Hispanic	*	*	*	735	13.5	(8.8 - 18.2)	
Hispanic	*	*	*	858	15.9	(9.8 - 22.0)	
Other Non-Hispanic	*	*	*	*	*	*	
Education							
≤ High School	199	11.4	(4.5 - 18.3)	15,654	22.8	(19.5 - 26.0)	
Some College	116	17.7	(8.5 - 26.9)	9,875	19.4	(15.8 - 22.9)	
≥ College Degree	338	5.7	(2.6 - 8.8)	12,421	10.0	(7.6 - 12.5)	
Income							
≤ \$24,999	92	13.3	(5.5 - 21.0)	9,626	24.1	(19.2 - 29.1)	
\$25,000-\$49,999	110	10.9	(4.3 - 17.5)	9,629	19.9	(16.0 - 23.7)	
\$50,000-\$74,999	*	*	*	5,756	17.8	(13.3 - 22.3)	
≥ \$75,000	270	9.6	(4.0 - 15.2)	8,549	10.6	(8.6 - 12.5)	

 $<sup>^{\</sup>mathrm{1}}$  Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day or some days

 $<sup>^{2}</sup>$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 55a. Everyday smoking among adults<sup>1</sup> in Rockland County and New York State

		Rockla	ind Count	ty		New Y	ork State	•
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	657	10,900	5.0	(3.2 - 6.8)	38,029	1,626,800	11.0	(9.8 - 12.2)
Gender								
Male	244	4,100	3.9	(1.3 - 6.5)	14,559	848,400	12.0	(10.0 - 14.1)
Female	413	6,700	6.0	(3.6 - 8.4)	23,470	778,400	10.1	(8.7 - 11.5)
Age (years)								
18-34	*	*	*	*	3,929	559,800	12.6	(9.6 - 15.6)
35-44	*	*	*	*	4,977	315,200	11.0	(8.4 - 13.7)
45-54	130	3,100	7.1	(3.3 - 10.8)	7,807	377,000	13.6	(11.1 - 16.0)
55-64	*	*	*	*	8,744	203,000	9.5	(7.5 - 11.5)
≥ 65	218	2,300	5.9	(1.9 - 9.9)	12,572	171,700	6.8	(4.8 - 8.8)
Race/Ethnicity								
White Non-Hispanic	550	8,400	5.6	(3.5 - 7.7)	35,729	1,158,100	12.4	(11.2 - 13.7)
Black Non-Hispanic	*	*	*	*	735	197,700	9.4	(5.6 - 13.3)
Hispanic	*	*	*	*	858	176,200	7.9	(4.2 - 11.6)
Other Non-Hispanic	*	*	*	*	707	94,900	8.5	(3.3 - 13.7)
Education								
≤ High School	199	2,900	4.4	(2.0 - 6.9)	15,654	881,900	15.9	(13.4 - 18.5)
Some College	116	4,400	10.7	(4.7 - 16.7)	9,875	410,200	11.7	(9.6 - 13.8)
≥ College Degree	338	3,600	3.3	(1.0 - 5.6)	12,421	323,500	5.8	(4.3 - 7.2)
Income								
≤ \$24,999	92	3,100	9.6	(3.6 - 15.6)	9,626	452,800	13.1	(10.3 - 15.8)
\$25,000-\$49,999	110	2,300	6.6	(2.3 - 11.0)	9,629	425,900	14.8	(11.9 - 17.7)
\$50,000-\$74,999	*	*	*	*	5,756	222,700	11.4	(8.5 - 14.4)
≥ \$75,000	*	*	*	*	8,549	320,200	7.1	(5.5 - 8.8)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults who have smoked at least 100 cigarettes in lifetime and currently smoking everyday (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 55b. Everyday smoking among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland (	County	N	ew York	State
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>
Total	657	4.8	(3.0 - 6.6)	38,029	11.3	(10.0 - 12.5)
Gender						
Male	244	4.0	(1.2 - 6.7)	14,559	12.1	(10.1 - 14.1)
Female	413	5.6	(3.2 - 8.0)	23,470	10.5	(9.0 - 12.0)
Race/Ethnicity						
White Non-Hispanic	550	5.4	(3.2 - 7.5)	35,729	12.8	(11.5 - 14.1)
Black Non-Hispanic	*	*	*	735	9.4	(5.5 - 13.4)
Hispanic	*	*	*	858	8.0	(4.3 - 11.7)
Other Non-Hispanic	*	*	*	707	8.5	(3.2 - 13.8)
Education						
≤ High School	199	4.1	(1.8 - 6.4)	15,654	16.4	(13.8 - 19.0)
Some College	116	10.6	(4.3 - 16.8)	9,875	12.1	(9.9 - 14.3)
≥ College Degree	338	3.1	(0.8 - 5.5)	12,421	5.8	(4.3 - 7.3)
Income						
≤ \$24,999	92	8.9	(2.9 - 14.9)	9,626	13.4	(10.6 - 16.2)
\$25,000-\$49,999	110	6.3	(2.1 - 10.4)	9,629	15.3	(12.3 - 18.3)
\$50,000-\$74,999	*	*	*	5,756	11.7	(8.7 - 14.8)
≥ \$75,000	*	*	*	8,549	7.2	(5.5 - 8.9)

 $<sup>^{\</sup>mathrm{1}}$  Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 56a. Adults living in homes in which smoking is prohibited in Rockland County and New York State

		Rockla	nd Coun	ty		New York State					
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>			
Total	655	187,200	86.3	(82.9 - 89.7)	38,045	11,997,100	80.8	(79.0 - 82.6)			
Gender											
Male	243	92,300	87.7	(82.6 - 92.8)	14,577	5,634,400	79.4	(76.6 - 82.2)			
Female	412	94,900	85.0	(80.5 - 89.5)	23,468	6,362,800	82.1	(79.9 - 84.3)			
Age (years)											
18-34	82	52,900	87.9	(79.4 - 96.4)	3,934	3,655,200	82.1	(77.8 - 86.4)			
35-44	92	32,900	86.7	(77.9 - 95.6)	4,985	2,479,700	86.1	(82.5 - 89.7)			
45-54	130	38,300	88.3	(82.4 - 94.1)	7,808	2,194,400	77.2	(73.5 - 80.8)			
55-64	135	32,000	87.4	(81.4 - 93.4)	8,753	1,732,500	81.0	(78.3 - 83.7)			
≥ 65	216	31,000	80.2	(74.3 - 86.1)	12,565	1,935,400	76.5	(73.4 - 79.5)			
Race/Ethnicity											
White Non-Hispanic	549	129,000	85.8	(82.0 - 89.6)	35,737	7,426,900	79.8	(78.3 - 81.2)			
Black Non-Hispanic	*	*	*	*	738	1,692,100	78.6	(71.9 - 85.3)			
Hispanic	*	*	*	*	863	2,003,900	88.7	(84.1 - 93.3)			
Other Non-Hispanic	*	*	*	*	*	*	*	*			
Education											
≤ High School	198	54,200	82.3	(74.9 - 89.6)	15,647	4,307,200	77.1	(74.0 - 80.2)			
Some College	115	35,100	86.4	(78.8 - 94.1)	9,883	2,795,400	78.9	(75.0 - 82.8)			
≥ College Degree	338	96,000	88.5	(84.4 - 92.7)	12,437	4,817,300	85.7	(83.3 - 88.2)			
Income											
≤ \$24,999	*	*	*	*	9,608	2,627,700	74.9	(70.0 - 79.7)			
\$25,000-\$49,999	110	27,400	80.4	(70.8 - 90.0)	9,640	2,314,500	79.8	(76.0 - 83.5)			
\$50,000-\$74,999	*	*	*	*	5,766	1,551,300	78.7	(74.0 - 83.4)			
≥ \$75,000	269	91,200	92.1	(88.2 - 96.0)	8,556	3,883,200	86.5	(84.3 - 88.7)			

 $<sup>^{\</sup>rm l}$  Defined as having a rule that prohibits smoking anywhere inside the home

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults who live in a home with a rule that prohibits smoking anywhere inside the home (rounded to the nearest hundred)

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a

Table 56b. Adults living in homes in which smoking is prohibited<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	n³	D				
		Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>
Total	655	86.3	(82.7 - 90.0)	38,045	80.9	(79.2 - 82.7)
Gender						
Male	243	87.4	(82.0 - 92.9)	14,577	79.4	(76.7 - 82.2)
Female	412	85.3	(80.3 - 90.2)	23,468	82.3	(80.0 - 84.5)
Race/Ethnicity						
White Non-Hispanic	549	85.6	(81.5 - 89.8)	35,737	79.9	(78.4 - 81.4)
Black Non-Hispanic	*	*	*	738	78.6	(71.8 - 85.4)
Hispanic	*	*	*	863	88.9	(84.4 - 93.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	198	81.4	(73.3 - 89.5)	15,647	76.9	(73.8 - 80.0)
Some College	115	86.6	(78.0 - 95.2)	9,883	79.1	(75.2 - 83.0)
≥ College Degree	338	89.0	(84.7 - 93.3)	12,437	86.0	(83.5 - 88.5)
Income						
≤ \$24,999	*	*	*	9,608	74.9	(70.1 - 79.7)
\$25,000-\$49,999	*	*	*	9,640	79.8	(76.1 - 83.6)
\$50,000-\$74,999	*	*	*	5,766	78.6	(73.8 - 83.3)
≥ \$75,000	269	91.8	(87.5 - 96.1)	8,556	86.6	(84.3 - 88.9)

 $<sup>^{\</sup>rm 1}$  Defined as having a rule that prohibits smoking anywhere inside the home

 $<sup>^{2}% \,\,\</sup>mathrm{Prevalence}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{\</sup>rm 3}$  Sample size from survey

 $<sup>^4</sup>$  95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 57a. Binge drinking within the past month among adults<sup>1</sup> in Rockland County and New York State

		Rockla	nd Coun	ty		New York State					
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>			
Total	643	26,600	12.5	(8.5 - 16.5)	37,745	2,602,000	17.7	(15.7 - 19.8)			
Gender											
Male	240	18,400	17.7	(10.4 - 25.0)	14,420	1,712,400	24.5	(20.9 - 28.1)			
Female	403	8,300	7.5	(4.3 - 10.8)	23,325	889,600	11.6	(9.5 - 13.7)			
Age (years)											
18-34	*	*	*	*	3,895	1,254,300	28.5	(22.8 - 34.2)			
35-44	91	4,300	11.7	(5.0 - 18.5)	4,934	584,800	20.6	(16.5 - 24.7)			
45-54	127	3,800	8.8	(3.2 - 14.5)	7,738	467,600	16.6	(13.3 - 19.9)			
55-64	134	1,900	5.1	(2.0 - 8.2)	8,663	207,200	9.8	(7.9 - 11.8)			
≥ 65	*	*	*	*	12,515	88,100	3.5	(2.5 - 4.5)			
Race/Ethnicity											
White Non-Hispanic	538	17,700	12.0	(8.0 - 16.0)	35,453	1,761,900	19.2	(17.5 - 20.8)			
Black Non-Hispanic	*	*	*	*	728	247,000	11.6	(6.3 - 17.0)			
Hispanic	*	*	*	*	854	460,600	20.5	(11.3 - 29.8)			
Other Non-Hispanic	*	*	*	*	710	132,500	12.1	(3.4 - 20.7)			
Education											
≤ High School	193	6,200	9.6	(3.1 - 16.0)	15,491	945,800	17.4	(13.2 - 21.6)			
Some College	*	*	*	*	9,819	636,600	18.1	(14.6 - 21.5)			
≥ College Degree	330	13,200	12.5	(7.0 - 17.9)	12,358	1,018,700	18.2	(15.3 - 21.1)			
Income											
≤ \$24,999	*	*	*	*	9,557	469,100	13.5	(8.5 - 18.5)			
\$25,000-\$49,999	*	*	*	*	9,575	534,300	18.5	(13.0 - 24.0)			
\$50,000-\$74,999	*	*	*	*	5,726	378,200	19.2	(14.1 - 24.3)			
≥ \$75,000	264	14,500	15.0	(9.4 - 20.6)	8,501	1,011,400	22.6	(19.7 - 25.6)			

 $<sup>^{1}</sup>$  Defined as men having 5 or more drinks or women having 4 or more drinks on 1 or more occasion within the past month

 $<sup>^{2}</sup>$  Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of adult binge drinkers within the past month (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 57b. Binge drinking within the past month among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	ew York	State
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	643	13.1	(8.9 - 17.3)	37,745	18.1	(16.1 - 20.2)
Gender						
Male	240	18.3	(10.8 - 25.9)	14,420	24.5	(21.0 - 27.9)
Female	403	8.1	(4.4 - 11.8)	23,325	12.4	(10.2 - 14.5)
Race/Ethnicity						
White Non-Hispanic	538	13.0	(8.6 - 17.3)	35,453	19.8	(18.1 - 21.6)
Black Non-Hispanic	*	*	*	728	11.8	(6.4 - 17.1)
Hispanic	*	*	*	854	20.2	(11.3 - 29.1)
Other Non-Hispanic	*	*	*	710	12.1	(3.4 - 20.8)
Education						
≤ High School	193	9.5	(2.9 - 16.1)	15,491	17.5	(13.5 - 21.5)
Some College	*	*	*	9,819	18.6	(15.1 - 22.0)
≥ College Degree	330	13.2	(7.5 - 19.0)	12,358	18.8	(15.9 - 21.7)
Income						
≤ \$24,999	*	*	*	9,557	13.4	(8.6 - 18.2)
\$25,000-\$49,999	*	*	*	9,575	18.9	(13.5 - 24.2)
\$50,000-\$74,999	*	*	*	5,726	19.9	(14.8 - 25.1)
≥ \$75,000	264	16.3	(10.2 - 22.5)	8,501	23.2	(20.2 - 26.1)

 $<sup>^{1}</sup>$  Defined as men having 5 or more drinks or women having 4 or more drinks on 1 or more occasion within the past month

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 58a. Heavy drinking within the past month among adults in Rockland County and New York State

		Rockla	nd Count	У		New Y	ork State	
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	645	5,100	2.4	(1.2 - 3.6)	37,553	718,700	4.9	(4.2 - 5.6)
Gender								
Male	*	*	*	*	14,330	333,300	4.8	(3.9 - 5.7)
Female	405	3,000	2.7	(1.3 - 4.1)	23,223	385,400	5.0	(4.0 - 6.1)
Age (years)								
18-34	*	*	*	*	3,866	224,900	5.1	(3.5 - 6.8)
35-44	*	*	*	*	4,924	137,200	4.8	(3.0 - 6.7)
45-54	*	*	*	*	7,715	144,600	5.2	(4.1 - 6.2)
55-64	*	*	*	*	8,624	117,400	5.6	(3.9 - 7.3)
≥ 65	213	1,700	4.4	(1.4 - 7.3)	12,424	94,600	3.8	(2.7 - 4.9)
Race/Ethnicity								
White Non-Hispanic	541	4,000	2.7	(1.5 - 3.9)	35,279	572,400	6.3	(5.4 - 7.1)
Black Non-Hispanic	*	*	*	*	723	58,100	2.8	(0.6 - 4.9)
Hispanic	*	*	*	*	848	60,900	2.7	(0.5 - 4.9)
Other Non-Hispanic	*	*	*	*	703	27,200	2.5	(0.8 - 4.2)
Education								
≤ High School	196	2,800	4.4	(1.0 - 7.8)	15,411	228,100	4.2	(3.2 - 5.2)
Some College	*	*	*	*	9,759	163,700	4.7	(3.6 - 5.8)
≥ College Degree	329	1,500	1.4	(0.4 - 2.4)	12,306	326,800	5.9	(4.5 - 7.3)
Income								
≤ \$24,999	*	*	*	*	9,512	104,200	3.0	(1.8 - 4.1)
\$25,000-\$49,999	*	*	*	*	9,537	158,700	5.5	(3.9 - 7.2)
\$50,000-\$74,999	*	*	*	*	5,708	95,400	4.9	(3.1 - 6.7)
≥ \$75,000	265	2,000	2.0	(0.7 - 3.4)	8,481	313,900	7.0	(5.4 - 8.6)

 $<sup>^{1}</sup>$  Defined as adult men averaging more than 2 alcoholic drinks per day and adult women averaging more than 1 alcoholic drink per day within the past month  $^{2}$  Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of adult heavy drinkers within the past month (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 58b. Heavy drinking within the past month among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland C	County	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	645	2.1	(1.0 - 3.1)	37,553	5.0	(4.3 - 5.8)		
Gender								
Male	*	*	*	14,330	4.9	(3.9 - 5.9)		
Female	405	2.4	(1.1 - 3.7)	23,223	5.1	(4.0 - 6.2)		
Race/Ethnicity								
White Non-Hispanic	541	2.4	(1.3 - 3.6)	35,279	6.4	(5.5 - 7.2)		
Black Non-Hispanic	*	*	*	723	2.9	(0.6 - 5.1)		
Hispanic	*	*	*	848	2.8	(0.5 - 5.1)		
Other Non-Hispanic	*	*	*	703	2.5	(0.8 - 4.2)		
Education								
≤ High School	196	3.6	(0.8 - 6.4)	15,411	4.3	(3.2 - 5.3)		
Some College	*	*	*	9,759	4.8	(3.6 - 5.9)		
≥ College Degree	329	1.5	(0.3 - 2.6)	12,306	6.0	(4.5 - 7.4)		
Income								
≤ \$24,999	*	*	*	9,512	3.0	(1.9 - 4.1)		
\$25,000-\$49,999	*	*	*	9,537	5.8	(4.0 - 7.5)		
\$50,000-\$74,999	*	*	*	5,708	5.0	(3.1 - 7.0)		
≥ \$75,000	265	1.8	(0.5 - 3.1)	8,481	7.0	(5.4 - 8.7)		

 $<sup>^{1}</sup>$  Defined as adult men averaging more than 2 alcoholic drinks per day and adult women averaging more than 1 alcoholic drink per day within the past month <sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 59a. Never or rarely been asked about sexual history during routine check-up among adults<sup>1</sup> in Rockland County and New York State

		Rockla	nd Coun	ty		New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	596	144,400	74.5	(69.3 - 79.6)	35,193	8,480,600	61.9	(59.2 - 64.5)		
Gender										
Male	216	69,400	76.2	(68.3 - 84.1)	13,299	4,357,400	67.7	(63.6 - 71.7)		
Female	380	75,000	72.9	(66.2 - 79.6)	21,894	4,123,300	56.8	(53.3 - 60.2)		
Age (years)										
18-34	*	*	*	*	3,507	1,804,600	44.3	(37.8 - 50.8)		
35-44	*	*	*	*	4,531	1,635,300	62.2	(56.6 - 67.8)		
45-54	121	32,400	82.9	(74.7 - 91.0)	7,215	1,715,600	65.4	(60.5 - 70.3)		
55-64	*	*	*	*	8,168	1,416,200	70.7	(66.6 - 74.8)		
≥ 65	200	30,500	86.1	(80.7 - 91.4)	11,772	1,908,900	80.3	(76.9 - 83.8)		
Race/Ethnicity										
White Non-Hispanic	496	105,800	79.2	(74.4 - 84.0)	33,090	5,940,300	68.5	(66.3 - 70.7)		
Black Non-Hispanic	*	*	*	*	683	831,800	40.8	(31.8 - 49.9)		
Hispanic	*	*	*	*	784	1,071,700	53.4	(43.6 - 63.1)		
Other Non-Hispanic	*	*	*	*	*	*	*	*		
Education										
≤ High School	176	44,800	77.2	(67.7 - 86.7)	14,264	3,001,200	60.1	(55.4 - 64.9)		
Some College	*	*	*	*	9,156	1,972,800	60.3	(54.9 - 65.7)		
≥ College Degree	309	71,800	74.1	(66.8 - 81.4)	11,706	3,476,000	65.0	(61.2 - 68.7)		
Income										
≤ \$24,999	*	*	*	*	8,772	1,529,400	49.3	(42.6 - 56.0)		
\$25,000-\$49,999	*	*	*	*	8,932	1,705,000	64.4	(58.0 - 70.7)		
\$50,000-\$74,999	76	16,400	82.7	(72.8 - 92.5)	5,433	1,217,000	64.5	(58.5 - 70.5)		
≥ \$75,000	249	68,600	75.1	(67.8 - 82.5)	8,108	2,934,300	68.2	(64.7 - 71.7)		

<sup>&</sup>lt;sup>1</sup> Defined as respondents who never or rarely (hardly ever) were asked about their sexual history, such as about sexual partners and sexual practices, during a routine check-up at a doctor's office or clinic

 $<sup>^{2}\ \</sup>mathrm{Sample}\ \mathrm{size}\ \mathrm{from}\ \mathrm{survey}$ 

<sup>&</sup>lt;sup>3</sup> Estimated number of adults who were never or rarely asked about their sexual history during a routine check-up at a doctor's office or \_clinic (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 59b. Never or rarely been asked about sexual history during routine check-up among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	ew York	State
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	596	73.8	(68.4 - 79.3)	35,193	61.4	(58.7 - 64.0)
Gender						
Male	216	75.5	(67.3 - 83.7)	13,299	67.9	(64.0 - 71.9)
Female	380	72.4	(65.1 - 79.6)	21,894	55.5	(52.0 - 59.0)
Race/Ethnicity						
White Non-Hispanic	496	78.4	(73.2 - 83.6)	33,090	67.8	(65.6 - 70.1)
Black Non-Hispanic	*	*	*	683	41.1	(32.0 - 50.2)
Hispanic	*	*	*	784	53.0	(43.3 - 62.6)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	176	77.2	(67.6 - 86.8)	14,264	59.5	(54.7 - 64.3)
Some College	*	*	*	9,156	59.7	(54.4 - 65.0)
≥ College Degree	309	73.2	(65.4 - 81.0)	11,706	64.5	(60.8 - 68.2)
Income						
≤ \$24,999	*	*	*	8,772	48.7	(42.0 - 55.4)
\$25,000-\$49,999	*	*	*	8,932	63.7	(57.5 - 70.0)
\$50,000-\$74,999	*	*	*	5,433	64.2	(58.2 - 70.2)
≥ \$75,000	249	73.9	(65.9 - 81.8)	8,108	67.5	(64.0 - 71.0)

Defined as respondents who never or rarely (hardly ever) were asked about their sexual history, such as about sexual partners and sexual practices, during a routine check-up at a doctor's office or clinic

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>&</sup>lt;sup>3</sup> Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 60a. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults<sup>1</sup> in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	e
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	372	112,800	85.5	(80.6 - 90.4)	22,788	7,011,100	77.9	(75.2 - 80.5)
Gender								
Male	148	62,500	88.5	(82.3 - 94.6)	9,034	3,618,500	81.0	(77.2 - 84.8)
Female	224	50,300	82.0	(74.4 - 89.6)	13,754	3,392,600	74.8	(71.2 - 78.4)
Age (years)								
18-34	*	*	*	*	3,213	2,362,500	70.8	(65.1 - 76.6)
35-44	*	*	*	*	3,513	1,313,400	71.7	(65.6 - 77.7)
45-54	76	22,400	87.2	(78.5 - 95.9)	4,975	1,290,600	80.3	(75.9 - 84.8)
55-64	69	16,700	91.6	(85.1 - 98.1)	4,955	932,200	87.2	(84.1 - 90.3)
≥ 65	104	17,000	93.3	(88.5 - 98.1)	6,132	1,112,400	95.9	(94.4 - 97.5)
Race/Ethnicity								
White Non-Hispanic	321	84,000	86.4	(81.5 - 91.2)	21,540	4,875,200	78.3	(75.6 - 80.9)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	491	1,002,400	84.8	(77.8 - 91.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	103	34,300	95.8	(89.7 - 100)	8,879	2,581,700	82.1	(77.9 - 86.3)
Some College	*	*	*	*	5,953	1,599,500	77.5	(72.6 - 82.3)
≥ College Degree	202	56,600	81.2	(74.0 - 88.3)	7,928	2,824,800	75.6	(71.8 - 79.5)
Income								
≤ \$24,999	*	*	*	*	5,491	1,602,900	78.7	(71.9 - 85.6)
\$25,000-\$49,999	*	*	*	*	5,912	1,335,900	78.8	(73.8 - 83.9)
\$50,000-\$74,999	*	*	*	*	3,717	987,500	76.6	(70.4 - 82.8)
≥ \$75,000	175	57,000	83.2	(76.1 - 90.4)	5,771	2,321,600	77.6	(74.0 - 81.2)

<sup>&</sup>lt;sup>1</sup> Defined as respondent estimating there were hardly any (0-1 out of 10) or a few (2-3 out of 10) persons their age in their community who have had a sexually transmitted disease such as gonorrhea, chlamydia, herpes or syphilis

<sup>&</sup>lt;sup>2</sup> Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults who believed hardly any or a few people their age in their community have had a sexually transmitted disease (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 60b. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults<sup>1</sup> in Rockland County and New York

State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	ew York	State
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	372	84.3	(78.8 - 89.9)	22,788	77.3	(74.7 - 79.9)
Gender						
Male	148	88.3	(81.7 - 94.9)	9,034	81.1	(77.4 - 84.8)
Female	224	79.9	(71.1 - 88.7)	13,754	73.6	(70.0 - 77.3)
Race/Ethnicity						
White Non-Hispanic	321	85.3	(79.7 - 90.8)	21,540	77.5	(74.9 - 80.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	491	84.6	(77.6 - 91.5)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	103	94.9	(87.4 - 100)	8,879	81.6	(77.4 - 85.7)
Some College	*	*	*	5,953	76.8	(72.0 - 81.6)
≥ College Degree	202	79.7	(71.6 - 87.9)	7,928	75.1	(71.3 - 79.0)
Income						
≤ \$24,999	*	*	*	5,491	77.9	(71.0 - 84.7)
\$25,000-\$49,999	*	*	*	5,912	77.9	(72.8 - 83.0)
\$50,000-\$74,999	*	*	*	3,717	76.1	(70.0 - 82.3)
≥ \$75,000	175	81.8	(73.6 - 89.9)	5,771	77.0	(73.4 - 80.6)

<sup>&</sup>lt;sup>1</sup> Defined as respondent estimating there were hardly any (0-1 out of 10) or a few (2-3 out of 10) persons their age in their community who have had a sexually transmitted disease such as gonorrhea, chlamydia, herpes or syphilis

<sup>&</sup>lt;sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 61a. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults<sup>1</sup> in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	2
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	608	173,900	85.6	(82.0 - 89.2)	36,287	12,738,300	89.7	(88.2 - 91.3)
Gender								
Male	231	87,300	86.1	(80.6 - 91.6)	13,893	6,104,600	89.4	(87.2 - 91.5)
Female	377	86,600	85.1	(80.6 - 89.6)	22,394	6,633,700	90.1	(87.9 - 92.2)
Age (years)								
18-34	74	44,500	79.1	(70.0 - 88.3)	3,850	4,007,400	91.3	(87.5 - 95.1)
35-44	88	31,700	89.0	(80.8 - 97.1)	4,875	2,578,600	93.2	(90.4 - 96.0)
45-54	124	35,900	87.6	(79.6 - 95.5)	7,607	2,416,800	90.5	(87.9 - 93.2)
55-64	125	31,400	90.1	(84.1 - 96.1)	8,411	1,876,500	92.2	(90.0 - 94.5)
≥ 65	197	30,400	85.8	(80.6 - 91.0)	11,544	1,858,900	79.7	(76.6 - 82.8)
Race/Ethnicity								
White Non-Hispanic	509	117,000	83.3	(78.9 - 87.6)	34,102	8,050,700	89.4	(87.7 - 91.0)
Black Non-Hispanic	*	*	*	*	702	1,822,200	91.9	(88.2 - 95.6)
Hispanic	*	*	*	*	820	1,967,200	89.9	(85.2 - 94.7)
Other Non-Hispanic	*	*	*	*	663	898,200	88.6	(79.5 - 97.6)
Education								
≤ High School	174	41,800	69.3	(60.5 - 78.2)	14,507	4,385,300	84.6	(81.9 - 87.2)
Some College	111	35,400	90.0	(83.3 - 96.7)	9,571	3,094,300	90.0	(86.7 - 93.4)
≥ College Degree	320	95,600	93.5	(89.8 - 97.2)	12,142	5,228,800	95.4	(94.0 - 96.7)
Income								
≤ \$24,999	*	*	*	*	9,021	2,833,500	86.5	(82.6 - 90.3)
\$25,000-\$49,999	102	26,300	83.1	(74.4 - 91.8)	9,261	2,529,900	89.5	(86.5 - 92.5)
\$50,000-\$74,999	*	*	*	*	5,644	1,740,000	91.2	(87.4 - 95.1)
≥ \$75,000	262	90,600	93.9	(90.3 - 97.6)	8,397	4,184,300	94.4	(92.5 - 96.3)

<sup>&</sup>lt;sup>1</sup> Defined as believing it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community

<sup>&</sup>lt;sup>2</sup> Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults who believed it is at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 61b. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	608	84.8	(80.9 - 88.7)	36,287	89.8	(88.3 - 91.4)		
Gender								
Male	231	85.9	(80.1 - 91.6)	13,893	89.1	(86.9 - 91.3)		
Female	377	83.8	(78.5 - 89.0)	22,394	90.5	(88.3 - 92.7)		
Race/Ethnicity								
White Non-Hispanic	509	82.2	(77.4 - 87.0)	34,102	89.5	(87.8 - 91.2)		
Black Non-Hispanic	*	*	*	702	91.7	(87.8 - 95.6)		
Hispanic	*	*	*	820	89.6	(84.7 - 94.5)		
Other Non-Hispanic	*	*	*	663	89.0	(80.5 - 97.6)		
Education								
≤ High School	174	66.8	(57.2 - 76.4)	14,507	84.6	(81.8 - 87.4)		
Some College	111	89.6	(82.7 - 96.5)	9,571	90.2	(86.9 - 93.5)		
≥ College Degree	320	93.3	(89.2 - 97.4)	12,142	95.4	(94.0 - 96.7)		
Income								
≤ \$24,999	*	*	*	9,021	86.5	(82.6 - 90.4)		
\$25,000-\$49,999	102	81.6	(72.1 - 91.1)	9,261	89.6	(86.5 - 92.6)		
\$50,000-\$74,999	*	*	*	5,644	91.1	(86.9 - 95.3)		
≥ \$75,000	262	94.0	(90.2 - 97.9)	8,397	94.5	(92.6 - 96.4)		

 $<sup>^{1}</sup>$  Defined as believing it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community <sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 62a. Overweight among adults<sup>1</sup> in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	•
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	621	93,400	45.2	(39.8 - 50.5)	36,710	5,213,000	36.4	(34.0 - 38.8)
Gender								
Male	240	54,100	52.3	(44.1 - 60.5)	14,502	3,073,200	43.7	(39.9 - 47.6)
Female	381	39,400	38.0	(31.2 - 44.8)	22,208	2,139,800	29.3	(26.4 - 32.2)
Age (years)								
18-34	*	*	*	*	3,789	1,295,200	30.1	(24.7 - 35.6)
35-44	*	*	*	*	4,785	1,052,000	37.5	(31.7 - 43.3)
45-54	*	*	*	*	7,495	1,053,300	38.8	(34.1 - 43.4)
55-64	*	*	*	*	8,356	798,100	38.8	(34.8 - 42.7)
≥ 65	211	16,900	44.8	(37.3 - 52.3)	12,285	1,014,400	41.5	(37.6 - 45.4)
Race/Ethnicity								
White Non-Hispanic	522	60,600	42.1	(36.6 - 47.7)	34,501	3,282,200	36.4	(34.4 - 38.3)
Black Non-Hispanic	*	*	*	*	708	740,000	36.7	(28.3 - 45.1)
Hispanic	*	*	*	*	813	781,500	35.6	(27.1 - 44.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,079	1,889,700	35.4	(31.2 - 39.5)
Some College	*	*	*	*	9,515	1,314,500	38.7	(33.8 - 43.7)
≥ College Degree	321	47,500	45.7	(38.4 - 53.0)	12,062	1,996,900	36.3	(32.8 - 39.7)
Income								
≤ \$24,999	*	*	*	*	9,346	1,170,900	34.0	(28.2 - 39.8)
\$25,000-\$49,999	*	*	*	*	9,368	1,136,000	40.4	(34.6 - 46.2)
\$50,000-\$74,999	*	*	*	*	5,615	738,700	38.3	(32.3 - 44.3)
≥ \$75,000	264	48,100	49.5	(41.9 - 57.2)	8,337	1,659,100	37.5	(34.1 - 40.9)

<sup>&</sup>lt;sup>1</sup> Defined as having a body mass index (BMI) more than 24.9 but less than 30.0. BMI is calculated as weight in kilograms divided by the square of height in meters. <sup>2</sup> Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of adults who are overweight (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 62b. Overweight among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	621	44.8	(39.2 - 50.4)	36,710	36.2	(33.8 - 38.6)		
Gender								
Male	240	52.6	(44.1 - 61.1)	14,502	43.9	(40.1 - 47.6)		
Female	381	37.0	(29.7 - 44.2)	22,208	28.9	(25.9 - 31.8)		
Race/Ethnicity								
White Non-Hispanic	522	41.5	(35.7 - 47.4)	34,501	36.0	(34.0 - 38.0)		
Black Non-Hispanic	*	*	*	708	37.0	(28.5 - 45.5)		
Hispanic	*	*	*	813	36.0	(27.5 - 44.4)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	15,079	35.2	(31.1 - 39.4)		
Some College	*	*	*	9,515	38.3	(33.3 - 43.2)		
≥ College Degree	321	45.9	(38.1 - 53.7)	12,062	36.2	(32.7 - 39.8)		
Income								
≤ \$24,999	*	*	*	9,346	33.8	(28.0 - 39.5)		
\$25,000-\$49,999	*	*	*	9,368	40.6	(34.8 - 46.4)		
\$50,000-\$74,999	*	*	*	5,615	38.6	(32.5 - 44.7)		
≥ \$75,000	264	49.5	(41.4 - 57.6)	8,337	37.2	(33.7 - 40.6)		

 $<sup>^{1}</sup>$  Defined as having a body mass index (BMI) more than 24.9 but less than 30.0. BMI is calculated as weight in kilograms divided by the square of height in meters.  $^{2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 63a. Obesity among adults<sup>1</sup> in Rockland County and New York State

		Rockla	ınd Coun	ty		New Y	ork State	•
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	621	32,600	15.8	(12.2 - 19.4)	36,710	3,327,400	23.2	(21.2 - 25.3)
Gender								
Male	240	19,800	19.1	(13.0 - 25.3)	14,502	1,505,800	21.4	(19.0 - 23.9)
Female	381	12,900	12.4	(8.9 - 16.0)	22,208	1,821,600	25.0	(21.8 - 28.1)
Age (years)								
18-34	*	*	*	*	3,789	730,600	17.0	(12.4 - 21.6)
35-44	88	5,300	14.4	(6.0 - 22.8)	4,785	662,200	23.6	(19.4 - 27.9)
45-54	123	8,200	19.7	(11.3 - 28.1)	7,495	816,800	30.1	(25.3 - 34.8)
55-64	125	6,300	18.2	(10.6 - 25.9)	8,356	563,900	27.4	(23.7 - 31.1)
≥ 65	211	7,700	20.5	(14.2 - 26.9)	12,285	554,000	22.7	(19.2 - 26.1)
Race/Ethnicity								
White Non-Hispanic	522	25,200	17.5	(13.5 - 21.6)	34,501	2,062,700	22.9	(21.2 - 24.6)
Black Non-Hispanic	*	*	*	*	708	703,100	34.9	(26.1 - 43.7)
Hispanic	*	*	*	*	813	456,400	20.8	(14.3 - 27.2)
Other Non-Hispanic	*	*	*	*	688	105,200	9.6	(4.0 - 15.3)
Education								
≤ High School	186	13,000	21.1	(13.1 - 29.0)	15,079	1,561,600	29.2	(25.3 - 33.2)
Some College	111	6,400	16.3	(8.3 - 24.3)	9,515	792,800	23.4	(19.4 - 27.3)
≥ College Degree	321	13,200	12.7	(8.4 - 17.0)	12,062	915,000	16.6	(14.4 - 18.8)
Income								
≤ \$24,999	*	*	*	*	9,346	974,800	28.3	(22.6 - 34.0)
\$25,000-\$49,999	102	6,600	20.5	(10.6 - 30.5)	9,368	584,800	20.8	(17.1 - 24.5)
\$50,000-\$74,999	78	2,700	12.8	(4.6 - 20.9)	5,615	475,300	24.6	(20.0 - 29.3)
≥ \$75,000	264	15,400	15.8	(10.8 - 20.9)	8,337	867,700	19.6	(17.0 - 22.3)

 $<sup>^{1}</sup>$  Defined as having a body mass index (BMI) of 30.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

<sup>2</sup> Sample size from survey

 $<sup>^{\</sup>rm 3}$  Estimated number of adults who are obese (rounded to the nearest hundred)

 $<sup>^4</sup>$  95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 63b. Obesity among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	621	15.0	(11.4 - 18.6)	36,710	23.1	(21.0 - 25.1)		
Gender								
Male	240	18.6	(12.4 - 24.8)	14,502	21.5	(19.0 - 24.0)		
Female	381	11.4	(7.9 - 14.9)	22,208	24.6	(21.5 - 27.8)		
Race/Ethnicity								
White Non-Hispanic	522	16.7	(12.6 - 20.8)	34,501	22.8	(21.0 - 24.5)		
Black Non-Hispanic	*	*	*	708	34.8	(25.9 - 43.6)		
Hispanic	*	*	*	813	20.5	(14.1 - 26.9)		
Other Non-Hispanic	*	*	*	688	9.5	(3.8 - 15.2)		
Education								
≤ High School	186	20.2	(12.2 - 28.1)	15,079	29.4	(25.4 - 33.3)		
Some College	111	16.6	(7.9 - 25.3)	9,515	23.1	(19.2 - 27.1)		
≥ College Degree	321	11.7	(7.5 - 15.9)	12,062	16.4	(14.2 - 18.5)		
Income								
≤ \$24,999	*	*	*	9,346	28.3	(22.6 - 34.1)		
\$25,000-\$49,999	*	*	*	9,368	21.0	(17.2 - 24.8)		
\$50,000-\$74,999	78	12.8	(4.4 - 21.3)	5,615	24.0	(19.4 - 28.7)		
≥ \$75,000	264	15.1	(9.9 - 20.2)	8,337	19.3	(16.7 - 21.9)		

<sup>&</sup>lt;sup>1</sup> Defined as having a body mass index (BMI) of 30.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 64a. Overweight or obesity among adults<sup>1</sup> in Rockland County and New York State

		Rockla	ind Coun	ty		New Y	ork State	e
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	621	126,100	61.0	(55.9 - 66.0)	36,710	8,540,500	59.6	(57.0 - 62.2)
Gender								
Male	240	73,800	71.5	(63.9 - 79.0)	14,502	4,579,100	65.2	(61.0 - 69.4)
Female	381	52,200	50.5	(43.8 - 57.1)	22,208	3,961,400	54.3	(51.0 - 57.6)
Age (years)								
18-34	*	*	*	*	3,789	2,025,800	47.1	(40.8 - 53.5)
35-44	*	*	*	*	4,785	1,714,200	61.1	(55.6 - 66.6)
45-54	123	27,700	66.4	(56.5 - 76.4)	7,495	1,870,100	68.8	(64.6 - 73.1)
55-64	125	24,200	69.5	(60.3 - 78.8)	8,356	1,362,000	66.2	(61.9 - 70.4)
≥ 65	211	24,600	65.3	(58.2 - 72.5)	12,285	1,568,400	64.1	(60.6 - 67.7)
Race/Ethnicity								
White Non-Hispanic	522	85,800	59.7	(54.4 - 64.9)	34,501	5,344,900	59.3	(57.2 - 61.3)
Black Non-Hispanic	*	*	*	*	708	1,443,100	71.6	(62.5 - 80.7)
Hispanic	*	*	*	*	813	1,237,900	56.4	(46.5 - 66.2)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	186	40,400	65.2	(56.1 - 74.3)	15,079	3,451,400	64.6	(59.8 - 69.4)
Some College	*	*	*	*	9,515	2,107,300	62.1	(57.0 - 67.2)
≥ College Degree	321	60,700	58.4	(51.2 - 65.6)	12,062	2,911,900	52.9	(49.2 - 56.6)
Income								
≤ \$24,999	*	*	*	*	9,346	2,145,700	62.3	(55.7 - 68.9)
\$25,000-\$49,999	*	*	*	*	9,368	1,720,700	61.2	(54.5 - 67.9)
\$50,000-\$74,999	*	*	*	*	5,615	1,214,000	62.9	(57.1 - 68.8)
≥ \$75,000	264	63,500	65.4	(58.5 - 72.2)	8,337	2,526,700	57.1	(53.6 - 60.6)

<sup>&</sup>lt;sup>1</sup> Defined as having a body mass index (BMI) of 25.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters. <sup>2</sup> Sample size from survey

 $<sup>^{3}</sup>$  Estimated number of adults who are overweight or obese (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 64b. Overweight or obesity among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	621	59.8	(54.4 - 65.2)	36,710	59.3	(56.8 - 61.9)		
Gender								
Male	240	71.2	(63.4 - 79.0)	14,502	65.4	(61.3 - 69.4)		
Female	381	48.4	(41.2 - 55.5)	22,208	53.5	(50.1 - 56.8)		
Race/Ethnicity								
White Non-Hispanic	522	58.3	(52.7 - 63.9)	34,501	58.8	(56.7 - 60.8)		
Black Non-Hispanic	*	*	*	708	71.8	(63.0 - 80.6)		
Hispanic	*	*	*	813	56.5	(46.8 - 66.1)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	186	63.3	(53.7 - 72.9)	15,079	64.6	(59.9 - 69.3)		
Some College	*	*	*	9,515	61.4	(56.4 - 66.5)		
≥ College Degree	321	57.6	(50.0 - 65.2)	12,062	52.6	(48.9 - 56.3)		
Income								
≤ \$24,999	*	*	*	9,346	62.1	(55.6 - 68.6)		
\$25,000-\$49,999	*	*	*	9,368	61.6	(55.1 - 68.1)		
\$50,000-\$74,999	*	*	*	5,615	62.6	(56.7 - 68.6)		
≥ \$75,000	264	64.6	(57.3 - 71.9)	8,337	56.5	(52.9 - 60.0)		

 $<sup>^{1}</sup>$  Defined as having a body mass index (BMI) of 25.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.  $^2$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 65a. Received advice about weight by a health professional among adults<sup>1</sup> in Rockland County and New York State

		Rockla	ind Coun	ty		New Y	ork State	•
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	627	48,800	23.3	(19.2 - 27.5)	36,729	3,929,700	27.4	(25.1 - 29.7)
Gender								
Male	240	23,400	22.6	(16.5 - 28.7)	14,499	1,654,800	23.6	(20.6 - 26.6)
Female	387	25,400	24.0	(18.4 - 29.6)	22,230	2,274,900	31.0	(27.7 - 34.3)
Age (years)								
18-34	*	*	*	*	3,796	895,500	20.7	(15.3 - 26.2)
35-44	89	6,400	17.1	(8.3 - 25.9)	4,779	722,700	25.8	(20.9 - 30.6)
45-54	*	*	*	*	7,490	910,900	33.5	(29.0 - 38.1)
55-64	127	12,200	34.7	(24.7 - 44.6)	8,351	741,100	36.1	(32.0 - 40.3)
≥ 65	211	9,800	25.9	(19.4 - 32.5)	12,313	659,500	26.9	(23.0 - 30.7)
Race/Ethnicity								
White Non-Hispanic	523	37,200	25.8	(21.1 - 30.4)	34,502	2,386,200	26.5	(24.6 - 28.4)
Black Non-Hispanic	*	*	*	*	709	615,700	30.4	(22.0 - 38.8)
Hispanic	*	*	*	*	830	555,400	25.1	(17.8 - 32.4)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	189	11,100	17.5	(11.6 - 23.4)	15,116	1,469,700	27.5	(23.6 - 31.4)
Some College	*	*	*	*	9,515	946,800	27.8	(23.2 - 32.4)
≥ College Degree	323	25,800	24.7	(18.5 - 30.8)	12,041	1,445,500	26.2	(23.0 - 29.4)
Income								
≤ \$24,999	89	4,200	13.5	(4.8 - 22.2)	9,360	1,079,400	31.4	(25.3 - 37.5)
\$25,000-\$49,999	103	7,500	22.7	(13.0 - 32.5)	9,365	696,900	24.6	(19.8 - 29.4)
\$50,000-\$74,999	*	*	*	*	5,603	507,100	26.3	(21.6 - 31.0)
≥ \$75,000	265	26,300	27.1	(20.5 - 33.7)	8,334	1,207,100	27.3	(24.2 - 30.5)

 $<sup>^{1}</sup>$  Defined as being advised by a doctor, nurse or other health professional about their weight (lose, gain or maintain)

 $<sup>^2\,\</sup>mathrm{Sample}$  size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults provided advice by a health professional about their weight (lose, gain or maintain) (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 65b. Received advice about weight by a health professional among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	Ro	ockland	County	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	627	22.2	(17.9 - 26.6)	36,729	27.1	(24.8 - 29.4)		
Gender								
Male	240	21.0	(15.1 - 27.0)	14,499	23.4	(20.5 - 26.3)		
Female	387	23.4	(17.2 - 29.6)	22,230	30.6	(27.3 - 34.0)		
Race/Ethnicity								
White Non-Hispanic	523	24.0	(19.3 - 28.6)	34,502	26.1	(24.2 - 28.0)		
Black Non-Hispanic	*	*	*	709	30.4	(21.9 - 38.9)		
Hispanic	*	*	*	830	24.9	(17.7 - 32.2)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	189	16.0	(10.4 - 21.6)	15,116	27.3	(23.4 - 31.2)		
Some College	*	*	*	9,515	27.4	(22.9 - 32.0)		
≥ College Degree	323	23.6	(17.1 - 30.0)	12,041	25.9	(22.7 - 29.0)		
Income								
≤ \$24,999	89	12.9	(4.5 - 21.3)	9,360	31.0	(24.9 - 37.0)		
\$25,000-\$49,999	103	20.7	(11.3 - 30.0)	9,365	24.9	(20.0 - 29.8)		
\$50,000-\$74,999	*	*	*	5,603	25.9	(21.2 - 30.6)		
≥ \$75,000	265	26.3	(19.2 - 33.3)	8,334	26.9	(23.7 - 30.0)		

 $<sup>^{</sup>m 1}$  Defined as being advised by a doctor, nurse or other health professional about their weight (lose, gain or maintain)

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 66a. Received advice to lose weight by a health professional among adults provided weight advice<sup>1</sup> in Rockland County and New York State

		Rockla	nd Coun	ty		New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	163	40,100	82.2	(74.8 - 89.7)	9,724	3,412,900	86.8	(83.9 - 89.8)		
Gender										
Male	*	*	*	*	3,807	1,468,700	88.8	(85.7 - 91.8)		
Female	100	20,300	79.8	(69.8 - 89.7)	5,917	1,944,200	85.5	(80.8 - 90.1)		
Age (years)										
18-34	*	*	*	*	681	779,700	87.1	(80.6 - 93.6)		
35-44	*	*	*	*	1,215	622,100	86.1	(76.9 - 95.2)		
45-54	*	*	*	*	2,238	813,300	89.3	(84.4 - 94.2)		
55-64	*	*	*	*	2,697	645,500	87.1	(81.2 - 93.0)		
≥ 65	*	*	*	*	2,893	552,300	83.7	(76.9 - 90.6)		
Race/Ethnicity										
White Non-Hispanic	140	29,900	80.5	(72.1 - 89.0)	9,082	2,076,700	87.0	(84.5 - 89.6)		
Black Non-Hispanic	*	*	*	*	*	*	*	*		
Hispanic	*	*	*	*	251	483,600	87.1	(78.5 - 95.7)		
Other Non-Hispanic	*	*	*	*	*	*	*	*		
Education										
≤ High School	*	*	*	*	3,962	1,239,700	84.3	(78.6 - 90.1)		
Some College	*	*	*	*	2,633	814,800	86.1	(80.1 - 92.0)		
≥ College Degree	*	*	*	*	3,115	1,290,900	89.3	(85.3 - 93.3)		
Income										
≤ \$24,999	*	*	*	*	2,539	922,100	85.4	(78.7 - 92.1)		
\$25,000-\$49,999	*	*	*	*	2,482	594,100	85.2	(78.4 - 92.1)		
\$50,000-\$74,999	*	*	*	*	1,521	446,100	88.0	(83.2 - 92.7)		
≥ \$75,000	*	*	*	*	2,270	1,073,200	88.9	(83.7 - 94.1)		

 $<sup>^{\</sup>rm I}$  Defined as provided advice by a health professional about their weight and advised to lose weight

 $<sup>^{2}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>3</sup> Estimated number of adults provided advice by a health professional about their weight and advised to lose weight (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 66b. Received advice to lose weight by a health professional among adults provided weight advice<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	<b>Rockland County</b>			N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	163	81.8	(73.8 - 89.9)	9,724	86.9	(83.8 - 89.9)		
Gender								
Male	*	*	*	3,807	88.7	(85.7 - 91.8)		
Female	*	*	*	5,917	85.5	(80.7 - 90.3)		
Race/Ethnicity								
White Non-Hispanic	140	79.7	(70.5 - 89.0)	9,082	87.0	(84.4 - 89.6)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	251	87.6	(79.3 - 95.8)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	*	*	*	3,962	84.6	(78.7 - 90.5)		
Some College	*	*	*	2,633	85.6	(79.5 - 91.7)		
≥ College Degree	*	*	*	3,115	89.4	(85.4 - 93.3)		
Income								
≤ \$24,999	*	*	*	2,539	85.7	(79.0 - 92.3)		
\$25,000-\$49,999	*	*	*	2,482	85.3	(78.4 - 92.3)		
\$50,000-\$74,999	*	*	*	1,521	88.0	(83.1 - 92.8)		
≥ \$75,000	*	*	*	2,270	88.7	(83.1 - 94.2)		

 $<sup>^{1}</sup>$  Defined as provided advice by a health professional about their weight and advised to lose weight

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>&</sup>lt;sup>3</sup> Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 67a. No leisure-time physical activity among adults<sup>1</sup> in Rockland County and New York State

		Rockla	nd Coun	ty		New Y	ork State	<b>e</b>
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	654	60,000	27.7	(23.0 - 32.4)	38,128	3,541,700	23.9	(21.6 - 26.1)
Gender								
Male	242	25,200	24.0	(17.1 - 30.9)	14,602	1,482,500	20.9	(17.8 - 24.0)
Female	412	34,800	31.2	(24.9 - 37.5)	23,526	2,059,200	26.6	(23.4 - 29.7)
Age (years)								
18-34	*	*	*	*	3,938	982,900	22.0	(16.6 - 27.5)
35-44	92	5,000	13.2	(5.4 - 21.1)	4,988	556,000	19.3	(15.0 - 23.6)
45-54	131	10,300	23.7	(13.8 - 33.6)	7,827	759,000	26.8	(21.7 - 31.9)
55-64	*	*	*	*	8,765	464,900	21.7	(18.2 - 25.2)
≥ 65	215	12,600	32.8	(25.8 - 39.7)	12,610	778,800	30.7	(27.3 - 34.2)
Race/Ethnicity								
White Non-Hispanic	547	36,800	24.5	(20.0 - 29.1)	35,816	1,892,800	20.3	(18.6 - 22.0)
Black Non-Hispanic	*	*	*	*	739	690,000	32.0	(23.4 - 40.6)
Hispanic	*	*	*	*	860	600,800	26.7	(19.1 - 34.4)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	198	27,600	41.9	(32.3 - 51.4)	15,683	1,810,200	32.5	(28.4 - 36.5)
Some College	*	*	*	*	9,906	814,200	23.0	(18.6 - 27.3)
≥ College Degree	337	18,600	17.2	(11.5 - 22.9)	12,458	841,000	14.9	(11.9 - 18.0)
Income								
≤ \$24,999	*	*	*	*	9,640	1,332,200	38.0	(31.7 - 44.3)
\$25,000-\$49,999	*	*	*	*	9,658	648,000	22.3	(18.3 - 26.3)
\$50,000-\$74,999	*	*	*	*	5,773	430,700	21.8	(16.6 - 27.0)
≥ \$75,000	270	16,100	16.2	(9.7 - 22.7)	8,565	560,800	12.5	(10.3 - 14.6)

 $<sup>^{1}</sup>$  Defined as having not participated in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise

2 Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults who have not participated in any physical activities or exercises (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 67b. No leisure-time physical activity among adults in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	654	27.6	(22.7 - 32.5)	38,128	23.7	(21.5 - 25.9)		
Gender								
Male	242	23.2	(16.3 - 30.0)	14,602	20.9	(17.8 - 23.9)		
Female	412	31.7	(24.9 - 38.5)	23,526	26.3	(23.1 - 29.5)		
Race/Ethnicity								
White Non-Hispanic	547	24.7	(19.8 - 29.6)	35,816	20.0	(18.3 - 21.7)		
Black Non-Hispanic	*	*	*	739	32.0	(23.3 - 40.6)		
Hispanic	*	*	*	860	26.9	(19.2 - 34.5)		
Other Non-Hispanic	*	*	*	*	*	*		
Education								
≤ High School	198	41.7	(32.0 - 51.4)	15,683	32.3	(28.3 - 36.4)		
Some College	*	*	*	9,906	22.8	(18.5 - 27.2)		
≥ College Degree	337	17.0	(10.8 - 23.1)	12,458	14.9	(11.8 - 17.9)		
Income								
≤ \$24,999	*	*	*	9,640	37.9	(31.6 - 44.2)		
\$25,000-\$49,999	*	*	*	9,658	22.0	(18.0 - 26.0)		
\$50,000-\$74,999	*	*	*	5,773	21.6	(16.3 - 26.8)		
≥ \$75,000	270	16.2	(9.4 - 23.0)	8,565	12.6	(10.3 - 14.9)		

 $<sup>^{1}</sup>$  Defined as having not participated in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise

2 Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 68a. Consumption of 5 or more servings of fruits and vegetables per day among adults 1 in Rockland County and New York State

		Rockla	nd Coun	ty		New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	658	57,100	26.3	(21.8 - 30.8)	38,203	4,023,400	27.0	(24.9 - 29.2)		
Gender										
Male	244	18,500	17.5	(11.1 - 24.0)	14,634	1,603,900	22.6	(19.4 - 25.8)		
Female	414	38,600	34.5	(28.5 - 40.6)	23,569	2,419,500	31.1	(28.2 - 34.0)		
Age (years)										
18-34	*	*	*	*	3,940	1,261,700	28.3	(22.7 - 33.9)		
35-44	92	8,400	22.2	(12.6 - 31.8)	4,989	688,000	23.9	(19.5 - 28.3)		
45-54	131	11,000	25.2	(16.4 - 34.1)	7,837	746,800	26.2	(22.3 - 30.1)		
55-64	135	10,300	28.1	(18.3 - 37.9)	8,777	579,100	27.0	(23.6 - 30.5)		
≥ 65	218	9,500	24.3	(18.2 - 30.5)	12,660	747,800	29.4	(26.0 - 32.8)		
Race/Ethnicity										
White Non-Hispanic	551	41,700	27.7	(23.0 - 32.4)	35,884	2,538,000	27.2	(25.4 - 28.9)		
Black Non-Hispanic	*	*	*	*	740	572,400	26.5	(19.2 - 33.9)		
Hispanic	*	*	*	*	864	672,200	29.8	(21.0 - 38.5)		
Other Non-Hispanic	*	*	*	*	715	240,800	21.4	(12.0 - 30.8)		
Education										
≤ High School	199	15,000	22.8	(15.4 - 30.2)	15,726	1,309,800	23.4	(19.3 - 27.4)		
Some College	*	*	*	*	9,924	964,300	27.2	(23.0 - 31.3)		
≥ College Degree	339	31,900	29.4	(22.9 - 35.9)	12,471	1,747,500	31.0	(27.9 - 34.1)		
Income										
≤ \$24,999	*	*	*	*	9,667	831,500	23.7	(18.6 - 28.7)		
\$25,000-\$49,999	*	*	*	*	9,669	917,200	31.6	(25.6 - 37.6)		
\$50,000-\$74,999	*	*	*	*	5,777	501,000	25.3	(21.4 - 29.3)		
≥ \$75,000	270	21,400	21.5	(15.4 - 27.6)	8,569	1,260,100	28.0	(25.0 - 31.0)		

 $<sup>^{\</sup>mathrm{1}}$  Defined as having consumed 5 or more servings of fruits and vegetables in an average day

 $<sup>^{2} \, \</sup>mathrm{Sample} \, \mathrm{size} \, \mathrm{from} \, \mathrm{survey}$ 

 $<sup>^3</sup>$  Estimated number of adults who consume 5 or more servings of fruits and vegetables per day (rounded to the nearest hundred)

<sup>4 95%</sup> confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 68b. Consumption of 5 or more servings of fruits and vegetables per day among adults<sup>1</sup> in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland	County	N	New York State			
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	658	26.4	(21.7 - 31.1)	38,203	27.1	(24.9 - 29.2)		
Gender								
Male	244	17.5	(10.9 - 24.1)	14,634	22.7	(19.6 - 25.9)		
Female	414	34.8	(28.3 - 41.3)	23,569	31.0	(28.1 - 34.0)		
Race/Ethnicity								
White Non-Hispanic	551	27.9	(22.9 - 32.9)	35,884	27.1	(25.4 - 28.9)		
Black Non-Hispanic	*	*	*	740	26.8	(19.4 - 34.2)		
Hispanic	*	*	*	864	29.5	(20.9 - 38.1)		
Other Non-Hispanic	*	*	*	715	21.8	(12.2 - 31.5)		
Education								
≤ High School	199	23.2	(15.6 - 30.7)	15,726	23.3	(19.3 - 27.3)		
Some College	*	*	*	9,924	27.1	(22.9 - 31.2)		
≥ College Degree	339	29.3	(22.3 - 36.2)	12,471	31.2	(28.0 - 34.3)		
Income								
≤ \$24,999	*	*	*	9,667	23.9	(18.8 - 28.9)		
\$25,000-\$49,999	*	*	*	9,669	32.0	(26.0 - 38.0)		
\$50,000-\$74,999	*	*	*	5,777	25.0	(21.0 - 28.9)		
≥ \$75,000	270	21.4	(14.8 - 27.9)	8,569	28.0	(24.9 - 31.1)		

 $<sup>^{\</sup>rm 1}$  Defined as having consumed 5 or more servings of fruits and vegetables in an average day

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 69a. Had a fall within the past 3 months among adults<sup>1</sup> in Rockland County and New York State

		Rockla	and Count	ty		New Y	ork State	e
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
Total	470	14,600	12.6	(8.7 - 16.5)	28,783	1,244,800	17.0	(14.9 - 19.0)
Gender								
Male	181	5,900	10.8	(5.1 - 16.5)	10,966	489,400	14.4	(11.9 - 16.8)
Female	289	8,700	14.2	(8.9 - 19.4)	17,817	755,300	19.2	(16.1 - 22.3)
Age (years)								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	129	6,700	15.5	(8.2 - 22.9)	7,741	445,500	16.0	(12.0 - 20.0)
55-64	126	3,500	9.9	(2.5 - 17.3)	8,454	336,000	16.5	(13.4 - 19.6)
≥ 65	215	4,500	11.7	(6.7 - 16.6)	12,588	463,300	18.4	(15.2 - 21.6)
Race/Ethnicity								
White Non-Hispanic	414	10,500	11.6	(8.0 - 15.2)	27,456	869,000	16.6	(15.1 - 18.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	424	76,100	11.0	(4.5 - 17.4)
Other Non-Hispanic	*	*	*	*	445	44,300	10.0	(3.1 - 16.8)
Education								
≤ High School	149	3,200	8.4	(3.4 - 13.3)	12,417	493,000	17.3	(13.3 - 21.3)
Some College	*	*	*	*	7,232	299,400	18.7	(14.6 - 22.7)
≥ College Degree	238	8,100	14.1	(8.1 - 20.1)	9,086	441,500	15.4	(12.9 - 17.9)
Income								
≤ \$24,999	*	*	*	*	7,824	423,900	25.0	(18.5 - 31.6)
\$25,000-\$49,999	85	1,900	9.9	(3.1 - 16.7)	7,445	223,600	16.0	(12.8 - 19.2)
\$50,000-\$74,999	*	*	*	*	4,141	151,200	14.6	(10.8 - 18.4)
≥ \$75,000	179	7,100	13.6	(7.1 - 20.1)	5,874	287,100	12.5	(10.0 - 15.1)

 $<sup>^{1}</sup>$  Defined as respondent unintentionally coming to rest on the ground or another lower level one or more times within the past 3 months

 $<sup>^{2} \, \</sup>mathrm{Sample} \, \mathrm{size} \, \mathrm{from} \, \mathrm{survey}$ 

 $<sup>^3</sup>$  Estimated number of adults who have had a fall within the past 3 months (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 69b. Had a fall within the past 3 months among adults<sup>1</sup> in Rockland County and New York
State - Age-Adjusted<sup>2</sup>

	<b>Rockland County</b>			N	New York State			
n	3	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total 47	'0	12.7	(8.8 - 16.5)	28,783	17.3	(15.0 - 19.5)		
Gender								
Male 18	1	11.1	(5.2 - 17.0)	10,966	15.0	(12.1 - 17.8)		
Female 28	9	14.2	(9.1 - 19.3)	17,817	19.4	(16.0 - 22.8)		
Race/Ethnicity								
White Non-Hispanic 41	4	11.8	(8.1 - 15.4)	27,456	16.9	(15.2 - 18.7)		
Black Non-Hispanic	*	*	*	*	*	*		
Hispanic	*	*	*	424	10.7	(4.2 - 17.1)		
Other Non-Hispanic	*	*	*	445	10.0	(2.9 - 17.0)		
Education								
≤ High School 14	9	8.4	(3.3 - 13.5)	12,417	17.4	(13.0 - 21.7)		
Some College	*	*	*	7,232	19.0	(14.7 - 23.2)		
≥ College Degree 23	8	14.2	(8.3 - 20.2)	9,086	16.0	(13.1 - 18.8)		
Income								
≤ \$24,999	*	*	*	7,824	25.3	(18.3 - 32.4)		
\$25,000-\$49,999	35	10.3	(3.2 - 17.3)	7,445	15.9	(12.6 - 19.2)		
\$50,000-\$74,999	*	*	*	4,141	14.7	(10.6 - 18.8)		
≥ \$75,000 17	9	13.2	(7.0 - 19.5)	5,874	12.9	(9.9 - 15.8)		

<sup>&</sup>lt;sup>1</sup> Defined as respondent unintentionally coming to rest on the ground or another lower level one or more times within the past 3 months

 $<sup>^{\</sup>rm 2}$  Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{\</sup>rm 3}$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 70a. Had a fall that caused injury within the past 3 months among adults in Rockland County and New York State

		Rockla	nd Count	у		New York State				
	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n²	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>		
Total	470	3,800	3.2	(1.4 - 5.1)	28,760	335,700	4.6	(3.7 - 5.5)		
Gender										
Male	*	*	*	*	10,957	120,000	3.5	(2.5 - 4.6)		
Female	289	2,300	3.7	(1.3 - 6.0)	17,803	215,600	5.5	(4.1 - 6.9)		
Age (years)										
18-34	*	*	*	*	*	*	*	*		
35-44	*	*	*	*	*	*	*	*		
45-54	*	*	*	*	7,737	115,100	4.1	(2.9 - 5.4)		
55-64	*	*	*	*	8,449	90,300	4.4	(2.7 - 6.1)		
≥ 65	*	*	*	*	12,574	130,300	5.2	(3.4 - 6.9)		
Race/Ethnicity										
White Non-Hispanic	414	2,700	3.0	(1.3 - 4.8)	27,433	270,800	5.2	(4.2 - 6.1)		
Black Non-Hispanic	*	*	*	*	458	47,900	4.9	(0.9 - 9.0)		
Hispanic	*	*	*	*	424	12,400	1.8	(0.2 - 3.3)		
Other Non-Hispanic	*	*	*	*	445	4,600	1.0	(0.4 - 1.6)		
Education										
≤ High School	*	*	*	*	12,404	148,900	5.2	(3.6 - 6.8)		
Some College	*	*	*	*	7,228	85,400	5.3	(3.1 - 7.5)		
≥ College Degree	*	*	*	*	9,081	101,300	3.5	(2.4 - 4.6)		
Income										
≤ \$24,999	*	*	*	*	7,812	90,200	5.3	(3.6 - 7.1)		
\$25,000-\$49,999	*	*	*	*	7,442	93,200	6.7	(3.9 - 9.4)		
\$50,000-\$74,999	*	*	*	*	4,139	37,000	3.6	(1.8 - 5.3)		
≥ \$75,000	*	*	*	*	5,872	54,700	2.4	(1.5 - 3.3)		

<sup>&</sup>lt;sup>1</sup> Defined as as respondent unintentionally coming to rest on the ground or another lower level and caused respondent to limit regular activities for at least a day or to go see a doctor one or more times within the past 3 months  $^2$  Sample size from survey

 $<sup>^3</sup>$  Estimated number of adults who have had a fall that caused an injury within the past 3 months (rounded to the nearest hundred)

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Table 70b. Had a fall that caused injury within the past 3 months among adults in Rockland County and New York State - Age-Adjusted<sup>2</sup>

	R	ockland C	County	N	New York State			
	n³	Percent	(C.I.) <sup>4</sup>	n³	Percent	(C.I.) <sup>4</sup>		
Total	470	3.3	(1.3 - 5.2)	28,760	4.6	(3.6 - 5.5)		
Gender								
Male	*	*	*	10,957	3.6	(2.4 - 4.7)		
Female	289	3.7	(1.4 - 6.1)	17,803	5.5	(4.1 - 6.9)		
Race/Ethnicity								
White Non-Hispanic	414	2.9	(1.2 - 4.6)	27,433	5.2	(4.2 - 6.2)		
Black Non-Hispanic	*	*	*	458	4.7	(0.7 - 8.7)		
Hispanic	*	*	*	424	1.8	(0.1 - 3.4)		
Other Non-Hispanic	*	*	*	445	1.0	(0.4 - 1.6)		
Education								
≤ High School	*	*	*	12,404	5.2	(3.5 - 6.8)		
Some College	*	*	*	7,228	5.3	(3.1 - 7.5)		
≥ College Degree	*	*	*	9,081	3.6	(2.4 - 4.7)		
Income								
≤ \$24,999	*	*	*	7,812	5.1	(3.3 - 7.0)		
\$25,000-\$49,999	*	*	*	7,442	6.6	(3.9 - 9.2)		
\$50,000-\$74,999	*	*	*	4,139	3.7	(1.7 - 5.8)		
≥ \$75,000	*	*	*	5,872	2.4	(1.4 - 3.3)		

Defined as as respondent unintentionally coming to rest on the ground or another lower level and caused respondent to limit regular activities for at least a day or to go see a doctor one or more times within the past 3 months
Prevalence estimates are age-adjusted to the 2000 US Standard population

 $<sup>^{3}</sup>$  Sample size from survey

<sup>&</sup>lt;sup>4</sup> 95% confidence interval

<sup>\*</sup> Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)