

New York State Expanded Behavioral Risk Factor Surveillance System

Schuyler County

**Final Report
July 2008 - June 2009**

**New York State Department of Health
December 2009**

Contents

Introduction	12
Table 1. Schuyler County and New York State adult US Census Bureau population estimates, 2007	13
General Health	
Table 2a. Fair or poor general health among adults in Schuyler County and New York State	14
Table 2b. Fair or poor general health among adults in Schuyler County and New York State - Age-adjusted	15
Table 3a. Poor physical health 14 or more days within the past month among adults in Schuyler County and New York State	16
Table 3b. Poor physical health 14 or more days within the past month among adults in Schuyler County and New York State - Age-adjusted	17
Table 4a. Poor mental health 14 or more days within the past month among adults in Schuyler County and New York State	18
Table 4b. Poor mental health 14 or more days within the past month among adults in Schuyler County and New York State - Age-adjusted	19
Table 5a. Poor physical or mental health 14 or more days within the past month among adults in Schuyler County and New York State	20
Table 5b. Poor physical or mental health 14 or more days within the past month among adults in Schuyler County and New York State - Age-adjusted	21
Quality Health Care	
Table 6a. Health insurance among adults in Schuyler County and New York State	22
Table 6b. Health insurance among adults in Schuyler County and New York State - Age-adjusted	23
Table 7a. Health insurance among adults aged 18-64 in Schuyler County and New York State	24
Table 7b. Health insurance among adults aged 18-64 in Schuyler County and New York State - Age-adjusted	25

Contents

Table 8a. No health insurance among adults in Schuyler County and New York State	26
Table 8b. No health insurance among adults in Schuyler County and New York State - Age-adjusted	27
Table 9a. No health insurance among adults aged 18-64 in Schuyler County and New York State	28
Table 9b. No health insurance among adults aged 18-64 in Schuyler County and New York State - Age-adjusted	29
Table 10a. Adults with regular health care provider in Schuyler County and New York State	30
Table 10b. Adults with regular health care provider in Schuyler County and New York State - Age-adjusted	31
Table 11a. Cost prevented visit to doctor within the past year among adults in Schuyler County and New York State	32
Table 11b. Cost prevented visit to doctor within the past year among adults in Schuyler County and New York State - Age-adjusted	33
Table 12a. Visited a doctor for routine checkup within the past year among adults in Schuyler County and New York State	34
Table 12b. Visited a doctor for routine checkup within the past year among adults in Schuyler County and New York State - Age-adjusted	35
Table 13a. Visited a doctor for routine checkup within the past 2 years among adults in Schuyler County and New York State	36
Table 13b. Visited a doctor for routine checkup within the past 2 years among adults in Schuyler County and New York State - Age-adjusted	37
Table 14a. Dentist visit within the past year among adults in Schuyler County and New York State	38
Table 14b. Dentist visit within the past year among adults in Schuyler County and New York State - Age-adjusted	39
Table 15a. Had teeth cleaned within the past year among adults in Schuyler County and New York State	40
Table 15b. Had teeth cleaned within the past year among adults in Schuyler County and New York State - Age-adjusted	41

Contents

Table 16a. Permanent teeth extracted due to decay or gum disease among adults in Schuyler County and New York State	42
Table 16b. Permanent teeth extracted due to decay or gum disease among adults in Schuyler County and New York State - Age-adjusted	43
Table 17a. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older in Schuyler County and New York State	44
Table 17b. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older in Schuyler County and New York State - Age-adjusted	45
Table 18a. Flu shot within the past 12 months in Schuyler County and New York State	46
Table 18b. Flu shot within the past 12 months in Schuyler County and New York State - Age-adjusted	47
Table 19a. Flu vaccine that was sprayed in nose within the past 12 months in Schuyler County and New York State	48
Table 19b. Flu vaccine that was sprayed in nose within the past 12 months in Schuyler County and New York State - Age-adjusted	49
Table 20a. Flu shot or flu vaccine sprayed in nose within the past 12 months in Schuyler County and New York State	50
Table 20b. Flu shot or flu vaccine sprayed in nose within the past 12 months in Schuyler County and New York State - Age-adjusted	51
Table 21a. Flu shot among adults aged 65 and older in Schuyler County and New York State	52
Table 21b. Flu shot among adults aged 65 and older in Schuyler County and New York State - Age-adjusted	53
Table 22a. Pneumonia shot or pneumococcal vaccine in Schuyler County and New York State	54
Table 22b. Pneumonia shot or pneumococcal vaccine in Schuyler County and New York State - Age-adjusted	55
Table 23a. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older in Schuyler County and New York State	56
Table 23b. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older in Schuyler County and New York State - Age-adjusted	57

Contents

Cancer Screening

Table 24a. Ever had a mammogram among women aged 40 and older in Schuyler County and New York State	58
Table 24b. Ever had a mammogram among women aged 40 and older in Schuyler County and New York State - Age-adjusted	59
Table 25a. Had mammogram within the past 2 years among women aged 40 and older in Schuyler County and New York State	60
Table 25b. Had mammogram within the past 2 years among women aged 40 and older in Schuyler County and New York State - Age-adjusted	61
Table 26a. Had mammogram within the past 2 years among women aged 50 and older in Schuyler County and New York State	62
Table 26b. Had mammogram within the past 2 years among women aged 50 and older in Schuyler County and New York State - Age-adjusted	63
Table 27a. Ever had a Pap test among women in Schuyler County and New York State	64
Table 27b. Ever had a Pap test among women in Schuyler County and New York State - Age-adjusted	65
Table 28a. Had a Pap test within the past 3 years among women in Schuyler County and New York State	66
Table 28b. Had a Pap test within the past 3 years among women in Schuyler County and New York State - Age-adjusted	67
Table 29a. Ever had a digital rectal exam among men aged 40 and older in Schuyler County and New York State	68
Table 29b. Ever had a digital rectal exam among men aged 40 and older in Schuyler County and New York State - Age-adjusted	69
Table 30a. Had digital rectal exam within the past 2 years among men aged 40 and older in Schuyler County and New York State	70
Table 30b. Had digital rectal exam within the past 2 years among men aged 40 and older in Schuyler County and New York State - Age-adjusted	71

Contents

Table 31a. Ever had a prostate specific antigen test among men aged 40 and older in Schuyler County and New York State	72
Table 31b. Ever had a prostate specific antigen test among men aged 40 and older in Schuyler County and New York State - Age-adjusted	73
Table 32a. Had prostate specific antigen test within the past 2 years among men aged 40 and older in Schuyler County and New York State	74
Table 32b. Had prostate specific antigen test within the past 2 years among men aged 40 and older in Schuyler County and New York State - Age-adjusted	75
Table 33a. Prostate cancer among men aged 40 and older in Schuyler County and New York State	76
Table 33b. Prostate cancer among men aged 40 and older in Schuyler County and New York State - Age-adjusted	77
Table 34a. Home blood stool test ever used among adults aged 50 and older in Schuyler County and New York State	78
Table 34b. Home blood stool test ever used among adults aged 50 and older in Schuyler County and New York State - Age-adjusted	79
Table 35a. Home blood stool test used within the past year among adults aged 50 and older in Schuyler County and New York State	80
Table 35b. Home blood stool test used within the past year among adults aged 50 and older in Schuyler County and New York State - Age-adjusted	81
Table 36a. Home blood stool test used within the past 2 years among adults aged 50 and older in Schuyler County and New York State	82
Table 36b. Home blood stool test used within the past 2 years among adults aged 50 and older in Schuyler County and New York State - Age-adjusted	83
Table 37a. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in Schuyler County and New York State	84
Table 37b. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in Schuyler County and New York State - Age-adjusted	85
Table 38a. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Schuyler County and New York State	86

Contents

Table 38b. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Schuyler County and New York State - Age-adjusted	87
Table 39a. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Schuyler County and New York State	88
Table 39b. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Schuyler County and New York State - Age-adjusted	89
Chronic Diseases	
Table 40a. High blood pressure among adults in Schuyler County and New York State	90
Table 40b. High blood pressure among adults in Schuyler County and New York State - Age-adjusted	91
Table 41a. Blood pressure medication among adults with high blood pressure in Schuyler County and New York State	92
Table 41b. Blood pressure medication among adults with high blood pressure in Schuyler County and New York State - Age-adjusted	93
Table 42a. Ever had blood cholesterol checked among adults in Schuyler County and New York State	94
Table 42b. Ever had blood cholesterol checked among adults in Schuyler County and New York State - Age-adjusted	95
Table 43a. Blood cholesterol checked within the past 5 years among adults in Schuyler County and New York State	96
Table 43b. Blood cholesterol checked within the past 5 years among adults in Schuyler County and New York State - Age-adjusted	97
Table 44a. Cardiovascular disease among adults in Schuyler County and New York State	98
Table 44b. Cardiovascular disease among adults in Schuyler County and New York State - Age-adjusted	99
Table 45a. Coronary heart disease among adults in Schuyler County and New York State	100

Contents

Table 45b. Coronary heart disease among adults in Schuyler County and New York State - Age-adjusted	101
Table 46a. Ever diagnosed with diabetes among adults in Schuyler County and New York State	102
Table 46b. Ever diagnosed with diabetes among adults in Schuyler County and New York State - Age-adjusted	103
Table 47a. Ever diagnosed with asthma (lifetime) among adults in Schuyler County and New York State	104
Table 47b. Ever diagnosed with asthma (lifetime) among adults in Schuyler County and New York State - Age-adjusted	105
Table 48a. Current asthma among adults in Schuyler County and New York State	106
Table 48b. Current asthma among adults in Schuyler County and New York State - Age-adjusted	107
Table 49a. Chronic joint symptoms among adults in Schuyler County and New York State	108
Table 49b. Chronic joint symptoms among adults in Schuyler County and New York State - Age-adjusted	109
Table 50a. Arthritis among adults in Schuyler County and New York State	110
Table 50b. Arthritis among adults in Schuyler County and New York State - Age-adjusted	111
Table 51a. Activity limitations because of physical, mental or emotional problems among adults in Schuyler County and New York State	112
Table 51b. Activity limitations because of physical, mental or emotional problems among adults in Schuyler County and New York State - Age-adjusted	113
Table 52a. Have health problem that requires the use of special equipment among adults in Schuyler County and New York State	114
Table 52b. Have health problem that requires the use of special equipment among adults in Schuyler County and New York State - Age-adjusted	115
Table 53a. Disability among adults in Schuyler County and New York State	116
Table 53b. Disability among adults in Schuyler County and New York State - Age-adjusted	117

Contents

Health Behaviors

Table 54a. Current smoking among adults in Schuyler County and New York State	118
Table 54b. Current smoking among adults in Schuyler County and New York State - Age-adjusted	119
Table 55a. Everyday smoking among adults in Schuyler County and New York State	120
Table 55b. Everyday smoking among adults in Schuyler County and New York State - Age-adjusted	121
Table 56a. Adults living in homes in which smoking is prohibited in Schuyler County and New York State	122
Table 56b. Adults living in homes in which smoking is prohibited in Schuyler County and New York State - Age-adjusted	123
Table 57a. Binge drinking within the past month among adults in Schuyler County and New York State	124
Table 57b. Binge drinking within the past month among adults in Schuyler County and New York State - Age-adjusted	125
Table 58a. Heavy drinking within the past month among adults in Schuyler County and New York State	126
Table 58b. Heavy drinking within the past month among adults in Schuyler County and New York State - Age-adjusted	127
Table 59a. Never or rarely been asked about sexual history during routine check-up among adults in Schuyler County and New York State	128
Table 59b. Never or rarely been asked about sexual history during routine check-up among adults in Schuyler County and New York State - Age-adjusted	129
Table 60a. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults in Schuyler County and New York State	130
Table 60b. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults in Schuyler County and New York State - Age-adjusted	131

Contents

Table 61a. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults in Schuyler County and New York State	132
Table 61b. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults in Schuyler County and New York State - Age-adjusted	133

Overweight / Obesity

Table 62a. Overweight among adults in Schuyler County and New York State	134
Table 62b. Overweight among adults in Schuyler County and New York State - Age-adjusted	135
Table 63a. Obesity among adults in Schuyler County and New York State	136
Table 63b. Obesity among adults in Schuyler County and New York State - Age-adjusted	137
Table 64a. Overweight or obesity among adults in Schuyler County and New York State	138
Table 64b. Overweight or obesity among adults in Schuyler County and New York State - Age-adjusted	139
Table 65a. Received advice about weight by a health professional among adults in Schuyler County and New York State	140
Table 65b. Received advice about weight by a health professional among adults in Schuyler County and New York State - Age-adjusted	141
Table 66a. Received advice to lose weight by a health professional among adults provided weight advice in Schuyler County and New York State	142
Table 66b. Received advice to lose weight by a health professional among adults provided weight advice in Schuyler County and New York State - Age-adjusted	143

Physical Activity / Nutrition

Table 67a. No leisure-time physical activity among adults in Schuyler County and New York State	144
Table 67b. No leisure-time physical activity among adults in Schuyler County and New York State - Age-adjusted	145

Contents

Table 68a. Consumption of 5 or more servings of fruits and vegetables per day among adults in Schuyler County and New York State	146
Table 68b. Consumption of 5 or more servings of fruits and vegetables per day among adults in Schuyler County and New York State - Age-adjusted	147

Injury

Table 69a. Had a fall within the past 3 months among adults in Schuyler County and New York State	148
Table 69b. Had a fall within the past 3 months among adults in Schuyler County and New York State - Age-adjusted	149
Table 70a. Had a fall that caused injury within the past 3 months among adults in Schuyler County and New York State	150
Table 70b. Had a fall that caused injury within the past 3 months among adults in Schuyler County and New York State - Age-adjusted	151

Introduction

The New York State Expanded Behavioral Risk Factor Surveillance System (Expanded BRFSS) is a random digit dialed telephone survey of adults (18 years of age and older) that is representative of the non-institutionalized civilian population with landline telephones living in New York State. The goal of the Expanded BRFSS is to collect county-specific data on preventive health practices, risk behaviors, injuries and preventable chronic and infectious diseases. Topics assessed by the Expanded BRFSS include tobacco use, physical inactivity, diet, use of cancer screening services and other factors linked to the leading causes of morbidity and mortality. For each topic, health indicators are based on answers to one or more questions in the survey. The survey results are weighted to more accurately represent the adult population living in each county. Weighting procedures adjust for the differential selection probabilities by race, age and gender. More detailed information on weighting procedures is available in the technical brief.

Survey data collection occurred in 58 geographic areas of New York State, including each of the 57 counties outside of New York City and New York City (5 counties) as a single area. A standard questionnaire was used in all areas. Data collection for the 2009 Expanded BRFSS took place during the period of July 2008 through June 2009. The goal was to complete about 650 interviews in each area. The final number of interviews varies because of the sampling design of the survey and differential response rates. More detailed information on the sampling design is available in the technical brief.

The New York State Department of Health contracted with RTI International, which is responsible for the data collection, sampling, weighting and creation of the statistical reports. RTI International subcontracted with Clearwater Research Inc. for data collection and additional expertise in sample design and data quality control.

This report presents both crude estimates and estimates that have been age-adjusted to the Year 2000 US Standard Population when they represent the entire adult population of an area. Estimates for specific age groups are not age-adjusted. Age adjustment is a statistical process applied to rates that allows communities with different age structures to be compared. Counties can compare their data with results from New York State as a whole and with other counties. Confidence intervals at the 95% level associated with each health indicator give an estimated range of values which is likely to include the actual prevalence of the indicator. Confidence intervals are similar to margins of error. When the confidence intervals of two estimates of the same indicator from different areas do not overlap, they may be said to be statistically significantly different, i.e., these differences are unlikely related to chance and are considered true differences. If there is any value that is included in both intervals, the two estimates are not statistically significantly different.

The survey results for each of the 69 health indicators included in the survey are typically provided by gender, age, race and ethnicity, education and income in the Expanded BRFSS county reports. The estimates for some of these groupings may be suppressed in geographic areas with low proportions of adult respondents in these groupings.

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 1. Schuyler County and New York State adult US Census Bureau population estimates,¹ 2007

	Schuyler County		New York State	
	Adult Population ¹	Percent	Adult Population ¹	Percent
Total	15,009	100	14,884,315	100
Gender				
Male	7,412	49.4	7,103,791	47.7
Female	7,597	50.6	7,780,524	52.3
Age (years)				
18-34	4,128	27.5	4,497,479	30.2
35-44	2,452	16.3	2,844,115	19.1
45-54	2,938	19.6	2,854,316	19.2
55-64	2,520	16.8	2,142,000	14.4
≥ 65	2,971	19.8	2,546,405	17.1
Race/Ethnicity				
White Non-Hispanic	14,383	95.8	9,343,884	62.8
Black Non-Hispanic	277	1.8	2,156,941	14.5
Hispanic	205	1.4	2,258,717	15.2
Other Non-Hispanic	144	1.0	1,124,773	7.6

¹ Source: National Center for Health Statistics. Postcensal estimates of the resident population of the United States for July 1, 2000-July 1, 2007, by year, county, age, bridged race, Hispanic origin, and sex (Vintage 2007). Prepared under a collaborative arrangement with the U.S. Census Bureau; released August 7, 2008. Available from: http://www.cdc.gov/nchs/nvss/bridged_race.htm as of November 25, 2009.

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 2a. Fair or poor general health among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	668	2,400	16.3	(12.9 - 19.7)	38,124	2,531,400	17.1	(15.1 - 19.0)
Gender								
Male	279	1,300	17.0	(11.6 - 22.3)	14,599	1,069,700	15.1	(12.4 - 17.8)
Female	389	1,200	15.6	(11.4 - 19.8)	23,525	1,461,700	18.8	(16.1 - 21.6)
Age (years)								
18-34	*	*	*	*	3,940	535,800	12.0	(7.2 - 16.8)
35-44	77	400	17.0	(7.0 - 26.9)	4,985	289,200	10.0	(6.8 - 13.3)
45-54	127	400	13.7	(7.5 - 19.8)	7,819	507,800	17.9	(14.0 - 21.8)
55-64	174	500	18.7	(12.7 - 24.7)	8,757	424,700	19.9	(16.0 - 23.9)
≥ 65	233	800	26.6	(20.5 - 32.8)	12,623	773,900	30.5	(26.8 - 34.2)
Race/Ethnicity								
White Non-Hispanic	655	2,400	16.6	(13.1 - 20.1)	35,811	1,226,600	13.2	(11.7 - 14.6)
Black Non-Hispanic	*	*	*	*	739	457,300	21.3	(14.8 - 27.9)
Hispanic	*	*	*	*	861	684,100	30.4	(22.2 - 38.6)
Other Non-Hispanic	*	*	*	*	713	163,400	14.5	(5.1 - 24.0)
Education								
≤ High School	313	1,500	22.1	(16.5 - 27.8)	15,682	1,466,500	26.3	(22.6 - 30.0)
Some College	182	400	9.4	(5.3 - 13.5)	9,904	508,500	14.3	(10.9 - 17.8)
≥ College Degree	173	600	13.9	(7.2 - 20.6)	12,457	493,500	8.8	(6.5 - 11.1)
Income								
≤ \$24,999	192	900	25.4	(17.8 - 32.9)	9,633	1,269,200	36.3	(30.1 - 42.4)
\$25,000-\$49,999	196	700	15.6	(9.1 - 22.0)	9,649	447,600	15.4	(11.9 - 18.9)
\$50,000-\$74,999	*	*	*	*	5,769	237,800	12.1	(8.0 - 16.2)
≥ \$75,000	*	*	*	*	8,562	218,300	4.9	(3.6 - 6.1)

¹ Defined as reporting fair or poor general health

² Sample size from survey

³ Estimated number of adults who have fair or poor health (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 2b. Fair or poor general health among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	668	15.7	(12.0 - 19.3)	38,124	16.7	(14.8 - 18.6)
Gender						
Male	279	17.2	(11.3 - 23.1)	14,599	15.3	(12.6 - 18.0)
Female	389	14.2	(9.7 - 18.7)	23,525	18.0	(15.2 - 20.7)
Race/Ethnicity						
White Non-Hispanic	655	16.0	(12.2 - 19.8)	35,811	12.9	(11.4 - 14.4)
Black Non-Hispanic	*	*	*	739	20.8	(14.2 - 27.3)
Hispanic	*	*	*	861	29.9	(21.8 - 37.9)
Other Non-Hispanic	*	*	*	713	13.8	(4.8 - 22.8)
Education						
≤ High School	313	21.0	(14.9 - 27.1)	15,682	25.8	(22.1 - 29.5)
Some College	182	8.6	(4.4 - 12.8)	9,904	14.1	(10.7 - 17.6)
≥ College Degree	173	14.4	(6.7 - 22.1)	12,457	8.5	(6.3 - 10.8)
Income						
≤ \$24,999	192	24.5	(16.1 - 33.0)	9,633	35.9	(29.7 - 42.0)
\$25,000-\$49,999	196	14.3	(7.7 - 20.9)	9,649	15.0	(11.6 - 18.5)
\$50,000-\$74,999	*	*	*	5,769	11.8	(7.7 - 16.0)
≥ \$75,000	*	*	*	8,562	4.7	(3.5 - 6.0)

¹ Defined as reporting fair or poor general health

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 3a. Poor physical health 14 or more days within the past month among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	657	1,900	13.1	(10.0 - 16.2)	37,327	1,472,900	10.1	(8.7 - 11.6)
Gender								
Male	275	600	8.5	(5.3 - 11.8)	14,325	604,200	8.7	(7.0 - 10.5)
Female	382	1,300	17.5	(12.4 - 22.7)	23,002	868,600	11.4	(9.2 - 13.6)
Age (years)								
18-34	*	*	*	*	3,891	268,100	6.1	(2.7 - 9.5)
35-44	77	300	13.5	(5.4 - 21.7)	4,948	232,100	8.1	(5.2 - 11.1)
45-54	126	300	11.8	(6.0 - 17.5)	7,764	350,900	12.5	(9.4 - 15.6)
55-64	173	300	13.5	(8.2 - 18.8)	8,621	271,200	13.0	(10.3 - 15.8)
≥ 65	224	500	18.3	(12.6 - 23.9)	12,103	350,600	14.5	(12.1 - 16.9)
Race/Ethnicity								
White Non-Hispanic	644	1,800	12.5	(9.5 - 15.6)	35,073	929,700	10.2	(8.8 - 11.6)
Black Non-Hispanic	*	*	*	*	724	208,200	9.9	(4.8 - 14.9)
Hispanic	*	*	*	*	835	250,300	11.5	(6.5 - 16.4)
Other Non-Hispanic	*	*	*	*	695	84,600	7.6	(1.3 - 13.9)
Education								
≤ High School	306	1,100	16.3	(11.2 - 21.4)	15,178	692,800	12.9	(10.6 - 15.3)
Some College	181	400	10.1	(5.1 - 15.0)	9,751	362,500	10.3	(7.4 - 13.2)
≥ College Degree	170	400	11.0	(5.1 - 17.0)	12,327	359,300	6.4	(4.8 - 8.1)
Income								
≤ \$24,999	189	700	18.8	(11.9 - 25.8)	9,294	625,600	18.4	(13.6 - 23.3)
\$25,000-\$49,999	193	600	13.1	(7.1 - 19.1)	9,495	279,100	9.8	(7.3 - 12.3)
\$50,000-\$74,999	107	200	9.6	(2.0 - 17.1)	5,733	186,400	9.5	(5.8 - 13.2)
≥ \$75,000	*	*	*	*	8,510	219,100	4.9	(3.6 - 6.2)

¹ Defined as 14 or more days within the past month when physical health was not good

² Sample size from survey

³ Estimated number of adults with poor physical health 14 or more days within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 3b. Poor physical health 14 or more days within the past month among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	657	13.0	(9.5 - 16.5)	37,327	9.9	(8.4 - 11.4)
Gender						
Male	275	8.3	(4.8 - 11.7)	14,325	8.7	(6.9 - 10.5)
Female	382	17.7	(11.8 - 23.5)	23,002	11.0	(8.7 - 13.2)
Race/Ethnicity						
White Non-Hispanic	644	12.4	(9.0 - 15.8)	35,073	9.9	(8.5 - 11.3)
Black Non-Hispanic	*	*	*	724	9.7	(4.6 - 14.9)
Hispanic	*	*	*	835	11.3	(6.3 - 16.2)
Other Non-Hispanic	*	*	*	695	7.6	(1.2 - 14.0)
Education						
≤ High School	306	16.7	(10.7 - 22.6)	15,178	12.7	(10.2 - 15.1)
Some College	181	9.8	(4.5 - 15.1)	9,751	10.1	(7.2 - 12.9)
≥ College Degree	170	10.5	(4.2 - 16.9)	12,327	6.3	(4.7 - 8.0)
Income						
≤ \$24,999	189	18.6	(10.7 - 26.5)	9,294	18.4	(13.5 - 23.4)
\$25,000-\$49,999	193	13.1	(6.4 - 19.8)	9,495	9.4	(7.0 - 11.9)
\$50,000-\$74,999	107	9.9	(1.3 - 18.5)	5,733	9.0	(5.3 - 12.6)
≥ \$75,000	*	*	*	8,510	4.9	(3.5 - 6.2)

¹ Defined as 14 or more days within the past month when physical health was not good

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 4a. Poor mental health 14 or more days within the past month among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	662	1,400	9.4	(6.4 - 12.3)	37,528	1,473,200	10.1	(8.6 - 11.5)
Gender								
Male	279	500	6.5	(3.1 - 9.9)	14,407	555,200	7.9	(6.3 - 9.6)
Female	383	900	12.2	(7.5 - 16.9)	23,121	918,000	12.0	(9.7 - 14.3)
Age (years)								
18-34	*	*	*	*	3,889	496,700	11.2	(7.4 - 15.1)
35-44	*	*	*	*	4,947	254,100	8.9	(6.7 - 11.1)
45-54	127	200	8.4	(3.4 - 13.4)	7,749	347,200	12.3	(9.0 - 15.6)
55-64	171	200	9.4	(4.7 - 14.0)	8,623	211,300	10.1	(7.9 - 12.3)
≥ 65	231	200	8.4	(4.2 - 12.6)	12,320	163,900	6.7	(4.7 - 8.7)
Race/Ethnicity								
White Non-Hispanic	649	1,400	9.6	(6.6 - 12.6)	35,266	899,200	9.8	(8.4 - 11.2)
Black Non-Hispanic	*	*	*	*	722	227,800	10.9	(5.4 - 16.4)
Hispanic	*	*	*	*	845	263,700	11.8	(6.8 - 16.8)
Other Non-Hispanic	*	*	*	*	695	82,500	7.4	(1.4 - 13.5)
Education								
≤ High School	309	700	10.6	(6.2 - 15.0)	15,335	665,000	12.2	(10.0 - 14.5)
Some College	179	300	7.5	(2.3 - 12.7)	9,771	404,500	11.5	(8.3 - 14.8)
≥ College Degree	174	400	9.4	(3.3 - 15.5)	12,342	346,600	6.2	(4.5 - 7.9)
Income								
≤ \$24,999	191	600	17.2	(9.5 - 24.9)	9,398	611,500	17.9	(13.0 - 22.8)
\$25,000-\$49,999	193	200	4.3	(1.5 - 7.0)	9,547	260,300	9.0	(6.7 - 11.4)
\$50,000-\$74,999	*	*	*	*	5,726	187,300	9.6	(6.0 - 13.1)
≥ \$75,000	*	*	*	*	8,520	276,500	6.2	(4.6 - 7.7)

¹ Defined as 14 or more days within the past month when mental health was not good

² Sample size from survey

³ Estimated number of adults with poor mental health 14 or more days within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 4b. Poor mental health 14 or more days within the past month among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	662	9.8	(6.4 - 13.3)	37,528	10.2	(8.7 - 11.7)
Gender						
Male	279	6.8	(2.7 - 10.8)	14,407	7.9	(6.3 - 9.6)
Female	383	12.8	(7.3 - 18.3)	23,121	12.2	(9.8 - 14.6)
Race/Ethnicity						
White Non-Hispanic	649	10.1	(6.5 - 13.6)	35,266	10.0	(8.5 - 11.4)
Black Non-Hispanic	*	*	*	722	11.1	(5.5 - 16.7)
Hispanic	*	*	*	845	11.5	(6.5 - 16.4)
Other Non-Hispanic	*	*	*	695	7.6	(1.5 - 13.7)
Education						
≤ High School	309	11.0	(6.0 - 16.1)	15,335	12.3	(10.0 - 14.6)
Some College	179	7.9	(1.9 - 14.0)	9,771	11.8	(8.5 - 15.1)
≥ College Degree	174	9.8	(2.7 - 16.8)	12,342	6.3	(4.6 - 8.0)
Income						
≤ \$24,999	191	18.7	(9.4 - 27.9)	9,398	18.2	(13.2 - 23.1)
\$25,000-\$49,999	193	4.4	(1.2 - 7.6)	9,547	9.4	(6.9 - 11.9)
\$50,000-\$74,999	*	*	*	5,726	9.5	(5.9 - 13.1)
≥ \$75,000	*	*	*	8,520	6.2	(4.6 - 7.7)

¹ Defined as 14 or more days within the past month when mental health was not good

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 5a. Poor physical or mental health 14 or more days within the past month among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	650	2,600	18.0	(14.3 - 21.7)	37,005	2,385,900	16.5	(14.8 - 18.2)
Gender								
Male	274	900	12.4	(8.2 - 16.6)	14,231	937,700	13.6	(11.4 - 15.8)
Female	376	1,700	23.5	(17.7 - 29.3)	22,774	1,448,200	19.1	(16.5 - 21.7)
Age (years)								
18-34	*	*	*	*	3,859	602,600	13.8	(9.7 - 17.9)
35-44	*	*	*	*	4,924	388,300	13.7	(10.4 - 17.0)
45-54	126	500	17.7	(10.7 - 24.7)	7,722	551,000	19.6	(15.8 - 23.4)
55-64	170	500	19.5	(13.2 - 25.8)	8,545	395,500	19.2	(16.1 - 22.3)
≥ 65	221	600	21.5	(15.5 - 27.5)	11,955	448,500	18.8	(15.9 - 21.7)
Race/Ethnicity								
White Non-Hispanic	637	2,500	17.6	(13.9 - 21.3)	34,778	1,485,700	16.3	(14.8 - 17.8)
Black Non-Hispanic	*	*	*	*	715	343,700	16.5	(10.3 - 22.7)
Hispanic	*	*	*	*	832	412,900	19.0	(12.6 - 25.3)
Other Non-Hispanic	*	*	*	*	680	143,500	13.0	(5.0 - 21.0)
Education								
≤ High School	302	1,300	20.2	(14.7 - 25.8)	15,018	1,077,200	20.3	(17.3 - 23.2)
Some College	178	600	15.2	(8.5 - 21.9)	9,672	615,500	17.6	(14.1 - 21.1)
≥ College Degree	170	700	17.2	(10.0 - 24.5)	12,244	634,200	11.4	(9.2 - 13.6)
Income								
≤ \$24,999	187	1,000	27.4	(19.0 - 35.8)	9,199	924,000	27.3	(21.9 - 32.8)
\$25,000-\$49,999	190	700	16.0	(9.5 - 22.5)	9,425	456,000	16.1	(13.0 - 19.2)
\$50,000-\$74,999	107	300	10.9	(3.1 - 18.7)	5,696	312,000	16.0	(11.4 - 20.5)
≥ \$75,000	101	400	13.8	(5.3 - 22.2)	8,475	447,600	10.0	(8.1 - 12.0)

¹ Defined as 14 or more days within the past month when physical or mental health prevented usual activities, such as self-care, work or recreation

² Sample size from survey

³ Estimated number of adults with poor physical or mental health 14 or more days within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 5b. Poor physical or mental health 14 or more days within the past month among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	650	18.0	(13.9 - 22.2)	37,005	16.4	(14.7 - 18.2)
Gender						
Male	274	12.1	(7.5 - 16.7)	14,231	13.6	(11.4 - 15.8)
Female	376	23.9	(17.3 - 30.5)	22,774	19.0	(16.3 - 21.6)
Race/Ethnicity						
White Non-Hispanic	637	17.6	(13.5 - 21.7)	34,778	16.3	(14.7 - 17.8)
Black Non-Hispanic	*	*	*	715	16.7	(10.3 - 23.0)
Hispanic	*	*	*	832	18.5	(12.2 - 24.8)
Other Non-Hispanic	*	*	*	680	13.1	(5.0 - 21.3)
Education						
≤ High School	302	20.3	(14.1 - 26.6)	15,018	20.1	(17.1 - 23.1)
Some College	178	15.4	(7.9 - 22.8)	9,672	17.6	(14.1 - 21.1)
≥ College Degree	170	17.1	(9.0 - 25.1)	12,244	11.4	(9.2 - 13.7)
Income						
≤ \$24,999	187	27.9	(18.2 - 37.7)	9,199	27.6	(22.1 - 33.1)
\$25,000-\$49,999	190	16.2	(9.0 - 23.5)	9,425	16.1	(12.9 - 19.3)
\$50,000-\$74,999	107	11.1	(2.3 - 19.8)	5,696	15.6	(11.0 - 20.1)
≥ \$75,000	101	13.1	(4.1 - 22.1)	8,475	10.0	(8.0 - 12.0)

¹ Defined as 14 or more days within the past month when physical or mental health prevented usual activities, such as self-care, work or recreation

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 6a. Health insurance among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	669	11,900	79.5	(74.4 - 84.6)	38,119	12,854,200	86.6	(84.2 - 89.0)
Gender								
Male	279	5,300	72.3	(63.9 - 80.8)	14,597	5,881,100	83.0	(78.9 - 87.1)
Female	390	6,600	86.4	(81.2 - 91.6)	23,522	6,973,100	89.9	(87.2 - 92.6)
Age (years)								
18-34	*	*	*	*	3,910	3,302,100	74.3	(67.8 - 80.7)
35-44	76	2,000	84.1	(74.9 - 93.4)	4,986	2,560,300	88.9	(84.7 - 93.1)
45-54	127	2,400	80.8	(72.7 - 88.9)	7,831	2,530,100	88.9	(84.6 - 93.2)
55-64	174	2,200	88.2	(83.1 - 93.4)	8,764	1,986,700	92.8	(90.2 - 95.5)
≥ 65	235	2,900	98.2	(96.6 - 99.8)	12,628	2,475,000	97.8	(96.7 - 98.8)
Race/Ethnicity								
White Non-Hispanic	656	11,600	80.7	(75.8 - 85.6)	35,811	8,591,200	92.2	(90.6 - 93.8)
Black Non-Hispanic	*	*	*	*	736	1,781,600	82.7	(75.3 - 90.1)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	313	5,200	77.9	(70.8 - 85.0)	15,673	4,448,900	79.7	(74.9 - 84.5)
Some College	*	*	*	*	9,906	3,004,600	85.0	(80.1 - 89.8)
≥ College Degree	174	3,500	87.6	(78.9 - 96.3)	12,459	5,361,700	95.3	(93.3 - 97.2)
Income								
≤ \$24,999	193	2,800	74.0	(64.2 - 83.7)	9,640	2,437,200	69.7	(62.5 - 76.8)
\$25,000-\$49,999	*	*	*	*	9,662	2,384,000	82.2	(76.1 - 88.3)
\$50,000-\$74,999	109	2,300	89.4	(80.1 - 98.7)	5,772	1,855,700	94.0	(91.2 - 96.7)
≥ \$75,000	102	2,300	88.5	(79.6 - 97.3)	8,567	4,388,400	97.5	(96.6 - 98.5)

¹ Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

² Sample size from survey

³ Estimated number of adults who have health insurance (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 6b. Health insurance among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	669	78.3	(72.8 - 83.9)	38,119	86.7	(84.4 - 89.1)
Gender						
Male	279	71.4	(62.5 - 80.3)	14,597	83.6	(79.7 - 87.5)
Female	390	85.0	(78.9 - 91.1)	23,522	89.6	(86.8 - 92.3)
Race/Ethnicity						
White Non-Hispanic	656	79.5	(74.2 - 84.9)	35,811	92.1	(90.5 - 93.7)
Black Non-Hispanic	*	*	*	736	82.5	(75.0 - 90.0)
Hispanic	*	*	*	860	68.7	(59.0 - 78.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	313	76.2	(68.4 - 84.0)	15,673	80.1	(75.4 - 84.7)
Some College	*	*	*	9,906	84.6	(79.7 - 89.4)
≥ College Degree	174	87.0	(77.6 - 96.4)	12,459	95.4	(93.6 - 97.2)
Income						
≤ \$24,999	*	*	*	9,640	70.2	(63.2 - 77.1)
\$25,000-\$49,999	*	*	*	9,662	81.9	(76.0 - 87.9)
\$50,000-\$74,999	109	89.4	(79.8 - 99.0)	5,772	94.0	(91.3 - 96.7)
≥ \$75,000	102	88.4	(79.0 - 97.9)	8,567	97.5	(96.6 - 98.5)

¹ Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 7a. Health insurance among adults aged 18-64¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	427	8,900	74.8	(68.7 - 81.0)	25,165	10,251,300	84.2	(81.3 - 87.1)
Gender								
Male	185	4,000	66.8	(57.1 - 76.6)	10,078	4,839,900	80.3	(75.5 - 85.0)
Female	242	4,900	83.1	(76.5 - 89.7)	15,087	5,411,400	88.0	(84.7 - 91.3)
Age (years)								
18-34	*	*	*	*	3,910	3,302,100	74.3	(67.8 - 80.7)
35-44	76	2,000	84.1	(74.9 - 93.4)	4,981	2,556,600	88.9	(84.7 - 93.1)
45-54	127	2,400	80.8	(72.7 - 88.9)	7,791	2,472,900	88.7	(84.3 - 93.0)
55-64	167	2,200	88.9	(83.9 - 93.8)	8,483	1,919,700	92.8	(90.1 - 95.5)
≥ 65	-	-	-	-	-	-	-	-
Race/Ethnicity								
White Non-Hispanic	419	8,700	76.4	(70.4 - 82.3)	23,354	6,582,200	90.3	(88.3 - 92.4)
Black Non-Hispanic	*	*	*	*	567	1,464,100	80.2	(71.6 - 88.7)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	178	3,500	71.1	(62.0 - 80.2)	9,037	3,191,500	74.6	(68.6 - 80.6)
Some College	*	*	*	*	7,031	2,540,400	82.9	(77.4 - 88.4)
≥ College Degree	*	*	*	*	9,068	4,502,400	94.5	(92.2 - 96.8)
Income								
≤ \$24,999	*	*	*	*	4,918	1,707,200	62.6	(54.1 - 71.2)
\$25,000-\$49,999	*	*	*	*	6,180	1,783,000	77.7	(70.3 - 85.1)
\$50,000-\$74,999	*	*	*	*	4,554	1,553,400	92.9	(89.7 - 96.1)
≥ \$75,000	89	2,100	87.4	(77.8 - 97.0)	7,324	3,987,600	97.3	(96.3 - 98.3)

¹ Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

² Sample size from survey

³ Estimated number of adults aged 18-64 who have health insurance (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 7b. Health insurance among adults aged 18-64¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	427	74.2	(67.7 - 80.7)	25,165	84.3	(81.5 - 87.2)
Gender						
Male	*	*	*	10,078	80.5	(75.9 - 85.1)
Female	242	82.4	(75.0 - 89.7)	15,087	87.9	(84.6 - 91.2)
Race/Ethnicity						
White Non-Hispanic	419	75.6	(69.3 - 82.0)	23,354	90.3	(88.3 - 92.3)
Black Non-Hispanic	*	*	*	567	80.0	(71.3 - 88.6)
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	178	70.1	(60.4 - 79.8)	9,037	75.1	(69.3 - 80.9)
Some College	*	*	*	7,031	82.6	(77.1 - 88.1)
≥ College Degree	*	*	*	9,068	94.6	(92.5 - 96.8)
Income						
≤ \$24,999	*	*	*	4,918	63.2	(54.9 - 71.6)
\$25,000-\$49,999	*	*	*	6,180	77.5	(70.2 - 84.7)
\$50,000-\$74,999	*	*	*	4,554	93.0	(89.8 - 96.2)
≥ \$75,000	*	*	*	7,324	97.3	(96.3 - 98.3)

¹ Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 8a. No health insurance among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	669	3,100	20.5	(15.4 - 25.6)	38,119	1,989,300	13.4	(11.0 - 15.8)
Gender								
Male	279	2,000	27.7	(19.2 - 36.1)	14,597	1,207,400	17.0	(12.9 - 21.1)
Female	390	1,000	13.6	(8.4 - 18.8)	23,522	781,900	10.1	(7.4 - 12.8)
Age (years)								
18-34	*	*	*	*	3,910	1,144,100	25.7	(19.3 - 32.2)
35-44	76	400	15.9	(6.6 - 25.1)	4,986	319,200	11.1	(6.9 - 15.3)
45-54	127	600	19.2	(11.1 - 27.3)	7,831	316,500	11.1	(6.8 - 15.4)
55-64	174	300	11.8	(6.6 - 16.9)	8,764	153,300	7.2	(4.5 - 9.8)
≥ 65	*	*	*	*	12,628	56,200	2.2	(1.2 - 3.3)
Race/Ethnicity								
White Non-Hispanic	656	2,800	19.3	(14.4 - 24.2)	35,811	727,100	7.8	(6.2 - 9.4)
Black Non-Hispanic	*	*	*	*	736	372,800	17.3	(9.9 - 24.7)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	313	1,500	22.1	(15.0 - 29.2)	15,673	1,134,400	20.3	(15.5 - 25.1)
Some College	*	*	*	*	9,906	531,400	15.0	(10.2 - 19.9)
≥ College Degree	174	500	12.4	(3.7 - 21.1)	12,459	266,300	4.7	(2.8 - 6.7)
Income								
≤ \$24,999	193	1,000	26.0	(16.3 - 35.8)	9,640	1,061,200	30.3	(23.2 - 37.5)
\$25,000-\$49,999	*	*	*	*	9,662	516,700	17.8	(11.7 - 23.9)
\$50,000-\$74,999	*	*	*	*	5,772	119,100	6.0	(3.3 - 8.8)
≥ \$75,000	*	*	*	*	8,567	111,100	2.5	(1.5 - 3.4)

¹ Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

² Sample size from survey

³ Estimated number of adults who have no health insurance (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 8b. No health insurance among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	669	21.7	(16.1 - 27.2)	38,119	13.3	(10.9 - 15.6)
Gender						
Male	279	28.6	(19.7 - 37.5)	14,597	16.4	(12.5 - 20.3)
Female	390	15.0	(8.9 - 21.1)	23,522	10.4	(7.7 - 13.2)
Race/Ethnicity						
White Non-Hispanic	656	20.5	(15.1 - 25.8)	35,811	7.9	(6.3 - 9.5)
Black Non-Hispanic	*	*	*	736	17.5	(10.0 - 25.0)
Hispanic	*	*	*	860	31.3	(21.6 - 41.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	313	23.8	(16.0 - 31.6)	15,673	19.9	(15.3 - 24.6)
Some College	*	*	*	9,906	15.4	(10.6 - 20.3)
≥ College Degree	174	13.0	(3.6 - 22.4)	12,459	4.6	(2.8 - 6.4)
Income						
≤ \$24,999	*	*	*	9,640	29.8	(22.9 - 36.8)
\$25,000-\$49,999	*	*	*	9,662	18.1	(12.1 - 24.0)
\$50,000-\$74,999	*	*	*	5,772	6.0	(3.3 - 8.7)
≥ \$75,000	*	*	*	8,567	2.5	(1.5 - 3.4)

¹ Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 9a. No health insurance among adults aged 18-64¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	427	3,000	25.2	(19.0 - 31.3)	25,165	1,928,400	15.8	(12.9 - 18.7)
Gender								
Male	185	2,000	33.2	(23.4 - 42.9)	10,078	1,190,200	19.7	(15.0 - 24.5)
Female	242	1,000	16.9	(10.3 - 23.5)	15,087	738,200	12.0	(8.7 - 15.3)
Age (years)								
18-34	*	*	*	*	3,910	1,144,100	25.7	(19.3 - 32.2)
35-44	76	400	15.9	(6.6 - 25.1)	4,981	319,200	11.1	(6.9 - 15.3)
45-54	127	600	19.2	(11.1 - 27.3)	7,791	316,100	11.3	(7.0 - 15.7)
55-64	167	300	11.1	(6.2 - 16.1)	8,483	148,900	7.2	(4.5 - 9.9)
≥ 65	-	-	-	-	-	-	-	-
Race/Ethnicity								
White Non-Hispanic	419	2,700	23.6	(17.7 - 29.6)	23,354	703,100	9.7	(7.6 - 11.7)
Black Non-Hispanic	*	*	*	*	567	361,800	19.8	(11.3 - 28.4)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	178	1,400	28.9	(19.8 - 38.0)	9,037	1,087,700	25.4	(19.4 - 31.4)
Some College	*	*	*	*	7,031	522,800	17.1	(11.6 - 22.6)
≥ College Degree	*	*	*	*	9,068	261,200	5.5	(3.2 - 7.8)
Income								
≤ \$24,999	*	*	*	*	4,918	1,018,000	37.4	(28.8 - 45.9)
\$25,000-\$49,999	*	*	*	*	6,180	511,100	22.3	(14.9 - 29.7)
\$50,000-\$74,999	*	*	*	*	4,554	118,800	7.1	(3.9 - 10.3)
≥ \$75,000	*	*	*	*	7,324	110,700	2.7	(1.7 - 3.7)

¹ Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

² Sample size from survey

³ Estimated number of adults aged 18-64 who have no health insurance (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 9b. No health insurance among adults aged 18-64¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	427	25.8	(19.3 - 32.3)	25,165	15.7	(12.8 - 18.5)
Gender						
Male	*	*	*	10,078	19.5	(14.9 - 24.1)
Female	242	17.6	(10.3 - 25.0)	15,087	12.1	(8.8 - 15.4)
Race/Ethnicity						
White Non-Hispanic	419	24.4	(18.0 - 30.7)	23,354	9.7	(7.7 - 11.7)
Black Non-Hispanic	*	*	*	567	20.0	(11.4 - 28.7)
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	178	29.9	(20.2 - 39.6)	9,037	24.9	(19.1 - 30.7)
Some College	*	*	*	7,031	17.4	(11.9 - 22.9)
≥ College Degree	*	*	*	9,068	5.4	(3.2 - 7.5)
Income						
≤ \$24,999	*	*	*	4,918	36.8	(28.4 - 45.1)
\$25,000-\$49,999	*	*	*	6,180	22.5	(15.3 - 29.8)
\$50,000-\$74,999	*	*	*	4,554	7.0	(3.8 - 10.2)
≥ \$75,000	*	*	*	7,324	2.7	(1.7 - 3.7)

¹ Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 10a. Adults with regular health care provider¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	667	12,800	85.7	(81.9 - 89.5)	38,135	12,324,300	83.0	(80.4 - 85.6)
Gender								
Male	278	6,100	82.4	(76.2 - 88.6)	14,593	5,494,300	77.6	(73.2 - 82.0)
Female	389	6,700	88.9	(84.5 - 93.2)	23,542	6,830,000	87.8	(85.1 - 90.5)
Age (years)								
18-34	*	*	*	*	3,929	2,931,600	65.8	(59.2 - 72.5)
35-44	77	2,000	80.8	(71.4 - 90.1)	4,988	2,471,000	86.1	(81.9 - 90.3)
45-54	127	2,500	86.1	(79.5 - 92.6)	7,823	2,529,300	88.6	(84.6 - 92.7)
55-64	173	2,300	90.1	(85.1 - 95.1)	8,770	2,002,000	93.6	(91.7 - 95.4)
≥ 65	234	2,800	96.0	(93.6 - 98.3)	12,625	2,390,400	94.1	(92.2 - 96.0)
Race/Ethnicity								
White Non-Hispanic	654	12,300	85.7	(81.8 - 89.5)	35,821	8,227,700	88.2	(86.4 - 89.9)
Black Non-Hispanic	*	*	*	*	740	1,753,200	81.3	(72.6 - 89.9)
Hispanic	*	*	*	*	860	1,453,100	64.7	(54.7 - 74.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	312	5,600	84.7	(78.5 - 90.9)	15,692	4,249,200	75.9	(70.9 - 80.8)
Some College	181	3,700	86.6	(80.2 - 92.9)	9,904	2,935,000	82.8	(77.2 - 88.4)
≥ College Degree	174	3,500	86.4	(79.5 - 93.3)	12,456	5,046,100	89.8	(87.1 - 92.6)
Income								
≤ \$24,999	191	3,100	83.0	(74.8 - 91.1)	9,644	2,554,200	72.7	(65.9 - 79.6)
\$25,000-\$49,999	196	3,800	84.7	(77.2 - 92.1)	9,655	2,243,100	77.3	(70.0 - 84.6)
\$50,000-\$74,999	109	2,100	81.7	(72.1 - 91.3)	5,771	1,767,500	89.5	(85.2 - 93.9)
≥ \$75,000	102	2,300	88.2	(79.9 - 96.4)	8,560	4,122,800	91.9	(90.2 - 93.7)

¹ Defined as having one or more persons respondent thinks of as a personal doctor or health care provider

² Sample size from survey

³ Estimated number of adults who have at least one person they think of as their personal doctor or health care provider (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 10b. Adults with regular health care provider¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	667	84.7	(80.4 - 88.9)	38,135	83.0	(80.4 - 85.5)
Gender						
Male	278	81.5	(74.8 - 88.2)	14,593	78.1	(73.9 - 82.3)
Female	389	87.8	(82.7 - 92.9)	23,542	87.4	(84.6 - 90.1)
Race/Ethnicity						
White Non-Hispanic	654	84.5	(80.2 - 88.8)	35,821	88.0	(86.2 - 89.7)
Black Non-Hispanic	*	*	*	740	81.4	(73.0 - 89.8)
Hispanic	*	*	*	860	65.2	(55.5 - 75.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	312	84.1	(77.3 - 90.8)	15,692	75.8	(71.0 - 80.6)
Some College	181	85.3	(78.1 - 92.5)	9,904	82.9	(77.5 - 88.3)
≥ College Degree	174	84.9	(77.1 - 92.8)	12,456	89.8	(87.1 - 92.4)
Income						
≤ \$24,999	191	82.1	(72.9 - 91.2)	9,644	72.9	(66.1 - 79.6)
\$25,000-\$49,999	196	83.4	(75.2 - 91.6)	9,655	77.4	(70.5 - 84.3)
\$50,000-\$74,999	*	*	*	5,771	89.6	(85.3 - 94.0)
≥ \$75,000	102	87.6	(78.8 - 96.4)	8,560	91.6	(89.8 - 93.4)

¹ Defined as having one or more persons respondent thinks of as a personal doctor or health care provider

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 11a. Cost prevented visit to doctor within the past year among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	669	1,400	9.5	(6.3 - 12.6)	38,143	2,034,800	13.7	(11.6 - 15.8)
Gender								
Male	280	600	8.3	(3.3 - 13.4)	14,607	903,300	12.7	(9.5 - 16.0)
Female	389	800	10.6	(6.7 - 14.4)	23,536	1,131,500	14.6	(11.8 - 17.4)
Age (years)								
18-34	*	*	*	*	3,939	947,300	21.2	(15.6 - 26.9)
35-44	77	400	15.4	(6.2 - 24.6)	4,987	347,900	12.1	(8.0 - 16.2)
45-54	127	300	9.6	(4.3 - 14.8)	7,825	382,200	13.5	(9.9 - 17.1)
55-64	174	200	8.7	(4.3 - 13.2)	8,766	236,800	11.1	(8.0 - 14.2)
≥ 65	*	*	*	*	12,626	120,600	4.8	(2.5 - 7.0)
Race/Ethnicity								
White Non-Hispanic	656	1,400	9.7	(6.4 - 13.0)	35,833	984,600	10.6	(8.9 - 12.2)
Black Non-Hispanic	*	*	*	*	738	280,000	13.1	(7.2 - 19.0)
Hispanic	*	*	*	*	859	597,900	26.5	(17.3 - 35.7)
Other Non-Hispanic	*	*	*	*	713	172,300	15.3	(5.7 - 24.9)
Education								
≤ High School	314	700	9.9	(5.1 - 14.7)	15,696	996,600	17.8	(13.8 - 21.9)
Some College	181	400	9.8	(4.3 - 15.3)	9,911	512,400	14.5	(10.4 - 18.6)
≥ College Degree	174	300	8.3	(1.8 - 14.8)	12,453	466,900	8.3	(6.0 - 10.6)
Income								
≤ \$24,999	192	400	10.2	(4.9 - 15.5)	9,639	854,500	24.3	(18.4 - 30.3)
\$25,000-\$49,999	196	500	11.2	(4.5 - 17.8)	9,656	532,900	18.4	(12.5 - 24.3)
\$50,000-\$74,999	*	*	*	*	5,776	202,700	10.3	(6.0 - 14.5)
≥ \$75,000	*	*	*	*	8,561	192,000	4.3	(2.9 - 5.6)

¹ Defined as needing to see a doctor within the past year but not being able to because of cost

² Sample size from survey

³ Estimated number of adults who did not see a doctor because of cost within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 11b. Cost prevented visit to doctor within the past year among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	669	10.1	(6.5 - 13.7)	38,143	13.8	(11.7 - 15.9)
Gender						
Male	280	8.8	(3.2 - 14.4)	14,607	12.5	(9.4 - 15.6)
Female	389	11.4	(6.9 - 16.0)	23,536	15.0	(12.1 - 17.9)
Race/Ethnicity						
White Non-Hispanic	656	10.4	(6.7 - 14.1)	35,833	10.8	(9.2 - 12.5)
Black Non-Hispanic	*	*	*	738	13.2	(7.2 - 19.3)
Hispanic	*	*	*	859	26.1	(17.2 - 35.1)
Other Non-Hispanic	*	*	*	713	14.8	(5.5 - 24.2)
Education						
≤ High School	314	11.2	(5.5 - 16.9)	15,696	17.8	(13.8 - 21.7)
Some College	181	10.4	(4.3 - 16.5)	9,911	15.3	(11.0 - 19.5)
≥ College Degree	174	8.1	(1.4 - 14.9)	12,453	8.2	(6.0 - 10.5)
Income						
≤ \$24,999	192	11.2	(4.9 - 17.5)	9,639	24.5	(18.6 - 30.4)
\$25,000-\$49,999	196	12.0	(4.5 - 19.5)	9,656	18.7	(12.9 - 24.6)
\$50,000-\$74,999	*	*	*	5,776	10.4	(6.1 - 14.7)
≥ \$75,000	*	*	*	8,561	4.4	(3.1 - 5.8)

¹ Defined as needing to see a doctor within the past year but not being able to because of cost

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 12a. Visited a doctor for routine checkup within the past year among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	670	10,800	72.1	(67.3 - 76.9)	38,026	10,809,300	72.9	(70.6 - 75.3)
Gender								
Male	280	5,100	68.4	(60.8 - 75.9)	14,592	4,738,400	66.9	(63.0 - 70.9)
Female	390	5,800	75.8	(69.9 - 81.7)	23,434	6,070,900	78.5	(75.9 - 81.0)
Age (years)								
18-34	*	*	*	*	3,921	2,747,400	61.7	(55.6 - 67.9)
35-44	*	*	*	*	4,973	1,883,100	65.5	(60.2 - 70.7)
45-54	127	2,100	71.6	(63.3 - 79.9)	7,808	2,174,100	76.6	(73.2 - 80.0)
55-64	174	2,000	79.9	(73.5 - 86.4)	8,751	1,728,600	81.4	(78.7 - 84.2)
≥ 65	235	2,500	84.3	(79.1 - 89.5)	12,573	2,276,000	90.0	(87.5 - 92.5)
Race/Ethnicity								
White Non-Hispanic	657	10,500	72.7	(68.1 - 77.3)	35,721	6,746,200	72.4	(70.6 - 74.2)
Black Non-Hispanic	*	*	*	*	736	1,688,900	78.8	(70.4 - 87.3)
Hispanic	*	*	*	*	857	1,550,900	69.3	(60.0 - 78.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	314	5,100	75.8	(69.4 - 82.1)	15,640	4,150,600	74.7	(70.7 - 78.7)
Some College	*	*	*	*	9,880	2,524,000	71.3	(66.1 - 76.5)
≥ College Degree	174	2,800	69.4	(60.2 - 78.6)	12,428	4,042,900	71.9	(68.5 - 75.3)
Income								
≤ \$24,999	193	2,400	65.4	(55.7 - 75.1)	9,616	2,469,200	70.4	(64.2 - 76.6)
\$25,000-\$49,999	196	3,200	71.3	(61.7 - 81.0)	9,631	2,068,400	71.3	(65.3 - 77.3)
\$50,000-\$74,999	109	2,100	82.3	(73.6 - 91.0)	5,765	1,466,000	74.8	(69.6 - 80.0)
≥ \$75,000	102	1,900	75.1	(66.0 - 84.2)	8,547	3,176,500	70.7	(67.3 - 74.2)

¹ Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past year

² Sample size from survey

³ Estimated number of adults who have visited a doctor for a routine checkup within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 12b. Visited a doctor for routine checkup within the past year among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	670	70.8	(65.5 - 76.1)	38,026	72.7	(70.3 - 75.0)
Gender						
Male	280	67.7	(59.7 - 75.8)	14,592	67.1	(63.3 - 70.9)
Female	390	73.9	(67.0 - 80.7)	23,434	77.7	(75.1 - 80.3)
Race/Ethnicity						
White Non-Hispanic	657	71.4	(66.2 - 76.5)	35,721	71.9	(70.1 - 73.7)
Black Non-Hispanic	*	*	*	736	79.3	(71.3 - 87.4)
Hispanic	*	*	*	857	69.2	(60.1 - 78.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	314	75.0	(68.0 - 82.1)	15,640	74.4	(70.4 - 78.4)
Some College	*	*	*	9,880	70.9	(65.9 - 76.0)
≥ College Degree	*	*	*	12,428	71.7	(68.3 - 75.0)
Income						
≤ \$24,999	*	*	*	9,616	70.2	(64.1 - 76.2)
\$25,000-\$49,999	*	*	*	9,631	71.1	(65.3 - 76.9)
\$50,000-\$74,999	109	81.8	(72.2 - 91.4)	5,765	74.5	(69.2 - 79.8)
≥ \$75,000	*	*	*	8,547	70.3	(66.9 - 73.8)

¹ Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past year

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 13a. Visited a doctor for routine checkup within the past 2 years among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	670	12,300	82.1	(77.8 - 86.5)	38,026	12,726,600	85.9	(83.9 - 87.9)
Gender								
Male	280	5,700	76.8	(69.6 - 84.0)	14,592	5,791,900	81.8	(78.2 - 85.4)
Female	390	6,600	87.3	(82.6 - 92.1)	23,434	6,934,700	89.6	(87.9 - 91.3)
Age (years)								
18-34	*	*	*	*	3,921	3,535,400	79.4	(73.9 - 85.0)
35-44	77	2,100	85.6	(77.5 - 93.6)	4,973	2,355,600	81.9	(78.0 - 85.7)
45-54	127	2,400	80.7	(73.3 - 88.1)	7,808	2,507,900	88.4	(85.6 - 91.2)
55-64	174	2,200	88.0	(82.7 - 93.2)	8,751	1,924,800	90.7	(88.6 - 92.7)
≥ 65	235	2,700	90.7	(86.7 - 94.7)	12,573	2,402,900	95.0	(93.5 - 96.5)
Race/Ethnicity								
White Non-Hispanic	657	12,000	83.0	(79.0 - 87.0)	35,721	8,070,000	86.6	(85.4 - 87.9)
Black Non-Hispanic	*	*	*	*	736	1,914,500	89.4	(82.3 - 96.5)
Hispanic	*	*	*	*	857	1,832,800	81.9	(73.7 - 90.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	314	5,500	82.7	(76.8 - 88.6)	15,640	4,713,700	84.8	(81.2 - 88.4)
Some College	182	3,600	84.0	(74.7 - 93.2)	9,880	2,990,500	84.5	(79.6 - 89.4)
≥ College Degree	174	3,200	79.3	(71.1 - 87.5)	12,428	4,928,300	87.6	(85.3 - 89.9)
Income								
≤ \$24,999	193	3,000	79.5	(71.3 - 87.6)	9,616	2,911,700	83.0	(77.4 - 88.6)
\$25,000-\$49,999	196	3,600	79.8	(70.3 - 89.2)	9,631	2,375,500	81.9	(76.1 - 87.7)
\$50,000-\$74,999	109	2,300	89.4	(82.5 - 96.2)	5,765	1,733,500	88.4	(84.9 - 91.9)
≥ \$75,000	102	2,200	84.0	(76.4 - 91.6)	8,547	3,907,000	87.0	(84.8 - 89.2)

¹ Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past 2 years

² Sample size from survey

³ Estimated number of adults who have visited a doctor for a routine checkup within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 13b. Visited a doctor for routine checkup within the past 2 years among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	670	81.5	(76.8 - 86.2)	38,026	85.7	(83.7 - 87.6)
Gender						
Male	280	76.4	(68.8 - 84.0)	14,592	81.9	(78.4 - 85.3)
Female	390	86.5	(81.0 - 92.0)	23,434	89.1	(87.4 - 90.9)
Race/Ethnicity						
White Non-Hispanic	657	82.4	(78.0 - 86.8)	35,721	86.2	(85.0 - 87.5)
Black Non-Hispanic	*	*	*	736	89.6	(83.0 - 96.3)
Hispanic	*	*	*	857	81.8	(73.8 - 89.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	314	82.1	(75.6 - 88.7)	15,640	84.6	(81.1 - 88.1)
Some College	182	83.3	(73.3 - 93.3)	9,880	84.4	(79.8 - 89.0)
≥ College Degree	174	78.8	(69.9 - 87.6)	12,428	87.3	(85.0 - 89.6)
Income						
≤ \$24,999	193	78.4	(69.2 - 87.6)	9,616	83.2	(77.8 - 88.5)
\$25,000-\$49,999	*	*	*	9,631	81.9	(76.3 - 87.4)
\$50,000-\$74,999	109	89.5	(82.2 - 96.9)	5,765	88.1	(84.4 - 91.8)
≥ \$75,000	102	83.6	(75.4 - 91.8)	8,547	86.4	(84.2 - 88.7)

¹ Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past 2 years

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 14a. Dentist visit within the past year among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	664	10,300	69.0	(64.6 - 73.5)	38,026	10,540,400	71.1	(68.6 - 73.6)
Gender								
Male	277	4,900	66.3	(59.2 - 73.3)	14,573	4,845,800	68.4	(64.5 - 72.2)
Female	387	5,400	71.8	(66.4 - 77.2)	23,453	5,694,600	73.6	(70.4 - 76.8)
Age (years)								
18-34	57	3,400	81.3	(71.4 - 91.3)	3,924	2,789,700	62.6	(56.3 - 69.0)
35-44	*	*	*	*	4,979	2,181,000	75.8	(70.9 - 80.7)
45-54	126	2,000	70.5	(61.9 - 79.2)	7,814	2,149,300	75.8	(71.9 - 79.7)
55-64	171	1,700	69.8	(62.2 - 77.3)	8,742	1,690,500	79.2	(75.9 - 82.6)
≥ 65	233	1,700	59.0	(52.1 - 65.8)	12,567	1,729,800	68.5	(64.7 - 72.3)
Race/Ethnicity								
White Non-Hispanic	651	9,800	68.4	(63.9 - 72.9)	35,721	6,959,900	74.7	(72.7 - 76.6)
Black Non-Hispanic	*	*	*	*	735	1,405,800	66.2	(57.9 - 74.5)
Hispanic	*	*	*	*	862	1,425,600	63.1	(53.7 - 72.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	310	4,100	61.6	(54.7 - 68.5)	15,613	3,620,700	65.1	(60.7 - 69.5)
Some College	181	3,200	74.9	(67.5 - 82.3)	9,880	2,504,700	70.8	(66.1 - 75.5)
≥ College Degree	173	3,000	75.1	(66.2 - 83.9)	12,453	4,402,400	78.2	(74.6 - 81.8)
Income								
≤ \$24,999	192	1,900	51.9	(42.0 - 61.8)	9,594	1,972,200	56.2	(49.8 - 62.6)
\$25,000-\$49,999	193	3,200	72.1	(63.9 - 80.2)	9,649	1,896,000	65.8	(59.5 - 72.1)
\$50,000-\$74,999	*	*	*	*	5,763	1,482,400	75.2	(69.4 - 81.0)
≥ \$75,000	102	2,200	86.7	(79.5 - 94.0)	8,564	3,707,900	82.4	(79.5 - 85.3)

¹ Defined as having visited a dentist or a dental clinic for any reason within the past year

² Sample size from survey

³ Estimated number of adults who have visited a dentist or a dental clinic for any reason within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 14b. Dentist visit within the past year among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	664	69.1	(64.2 - 73.9)	38,026	71.1	(68.6 - 73.5)
Gender						
Male	277	65.0	(57.4 - 72.7)	14,573	68.3	(64.6 - 72.1)
Female	387	73.0	(67.0 - 78.9)	23,453	73.6	(70.4 - 76.8)
Race/Ethnicity						
White Non-Hispanic	651	68.4	(63.5 - 73.4)	35,721	74.4	(72.4 - 76.3)
Black Non-Hispanic	*	*	*	735	66.6	(58.3 - 74.9)
Hispanic	*	*	*	862	63.6	(54.4 - 72.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	310	62.3	(54.7 - 69.8)	15,613	65.3	(60.9 - 69.6)
Some College	181	74.6	(66.5 - 82.7)	9,880	70.4	(65.7 - 75.1)
≥ College Degree	173	73.9	(64.2 - 83.7)	12,453	78.1	(74.6 - 81.7)
Income						
≤ \$24,999	*	*	*	9,594	56.4	(50.1 - 62.8)
\$25,000-\$49,999	193	71.1	(62.1 - 80.1)	9,649	65.8	(59.6 - 72.0)
\$50,000-\$74,999	*	*	*	5,763	75.0	(69.1 - 80.8)
≥ \$75,000	102	86.4	(78.3 - 94.5)	8,564	82.1	(79.1 - 85.0)

¹ Defined as having visited a dentist or a dental clinic for any reason within the past year

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 15a. Had teeth cleaned within the past year among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	582	9,400	69.2	(64.3 - 74.1)	34,070	9,985,900	71.7	(69.1 - 74.3)
Gender								
Male	246	4,400	64.2	(56.4 - 72.0)	13,173	4,616,400	69.0	(65.0 - 73.1)
Female	336	5,100	74.2	(68.5 - 79.9)	20,897	5,369,500	74.2	(70.8 - 77.5)
Age (years)								
18-34	*	*	*	*	3,860	2,746,400	63.0	(56.5 - 69.5)
35-44	*	*	*	*	4,875	2,112,100	75.5	(70.7 - 80.3)
45-54	118	1,900	69.1	(60.0 - 78.1)	7,438	2,022,600	73.3	(69.1 - 77.4)
55-64	155	1,700	75.0	(67.7 - 82.2)	7,906	1,560,900	77.7	(73.8 - 81.6)
≥ 65	178	1,500	66.4	(58.9 - 74.0)	9,991	1,543,900	77.1	(73.7 - 80.5)
Race/Ethnicity								
White Non-Hispanic	571	9,000	68.4	(63.5 - 73.4)	31,982	6,601,500	75.4	(73.4 - 77.4)
Black Non-Hispanic	*	*	*	*	680	1,227,200	61.6	(52.1 - 71.0)
Hispanic	*	*	*	*	789	1,421,300	66.6	(57.1 - 76.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	244	3,600	64.4	(56.5 - 72.2)	12,740	3,388,600	67.7	(63.1 - 72.3)
Some College	169	2,900	71.6	(63.1 - 80.0)	9,145	2,286,800	68.5	(62.9 - 74.1)
≥ College Degree	169	2,900	73.5	(64.5 - 82.5)	12,128	4,298,500	78.3	(74.8 - 81.7)
Income								
≤ \$24,999	*	*	*	*	7,525	1,861,800	60.3	(53.5 - 67.1)
\$25,000-\$49,999	182	2,800	65.8	(56.5 - 75.2)	8,709	1,681,700	62.6	(55.4 - 69.8)
\$50,000-\$74,999	*	*	*	*	5,534	1,447,800	76.3	(70.7 - 81.8)
≥ \$75,000	100	2,100	82.8	(74.7 - 90.9)	8,448	3,617,500	80.8	(77.8 - 83.8)

¹ Defined as having had teeth cleaned by a dentist or dental hygienist within the past year

² Sample size from survey

³ Estimated number of adults who have had their teeth cleaned by a dentist or dental hygienist within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 15b. Had teeth cleaned within the past year among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	582	68.5	(63.2 - 73.9)	34,070	71.7	(69.1 - 74.2)
Gender						
Male	246	62.6	(54.2 - 71.1)	13,173	69.2	(65.3 - 73.1)
Female	336	74.4	(68.1 - 80.7)	20,897	73.9	(70.5 - 77.3)
Race/Ethnicity						
White Non-Hispanic	571	67.8	(62.4 - 73.2)	31,982	75.1	(73.0 - 77.1)
Black Non-Hispanic	*	*	*	680	62.0	(52.7 - 71.4)
Hispanic	*	*	*	789	66.9	(57.5 - 76.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	244	63.8	(55.2 - 72.4)	12,740	67.7	(63.2 - 72.3)
Some College	169	70.9	(61.6 - 80.1)	9,145	68.1	(62.6 - 73.6)
≥ College Degree	169	72.6	(62.8 - 82.5)	12,128	78.3	(74.9 - 81.7)
Income						
≤ \$24,999	*	*	*	7,525	60.2	(53.4 - 67.0)
\$25,000-\$49,999	*	*	*	8,709	62.7	(55.7 - 69.7)
\$50,000-\$74,999	*	*	*	5,534	76.4	(70.9 - 81.9)
≥ \$75,000	100	83.2	(74.6 - 91.8)	8,448	80.4	(77.4 - 83.4)

¹ Defined as having had teeth cleaned by a dentist or dental hygienist within the past year

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 16a. Permanent teeth extracted due to decay or gum disease among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	657	7,700	51.8	(46.7 - 57.0)	37,652	7,426,500	50.4	(47.8 - 52.9)
Gender								
Male	276	3,500	48.3	(40.5 - 56.0)	14,457	3,367,000	47.8	(43.9 - 51.6)
Female	381	4,100	55.3	(48.6 - 62.0)	23,195	4,059,500	52.7	(49.5 - 56.0)
Age (years)								
18-34	*	*	*	*	3,921	1,121,600	25.4	(19.7 - 31.0)
35-44	*	*	*	*	4,964	1,250,000	43.6	(37.9 - 49.2)
45-54	125	1,700	58.4	(48.9 - 67.9)	7,780	1,595,300	56.1	(51.4 - 60.7)
55-64	171	1,600	63.5	(55.3 - 71.7)	8,648	1,382,800	65.5	(61.6 - 69.4)
≥ 65	228	2,400	84.5	(79.6 - 89.4)	12,339	2,076,900	83.0	(80.6 - 85.4)
Race/Ethnicity								
White Non-Hispanic	644	7,400	52.0	(46.9 - 57.2)	35,364	4,425,700	47.9	(46.0 - 49.9)
Black Non-Hispanic	*	*	*	*	727	1,244,700	58.2	(48.9 - 67.4)
Hispanic	*	*	*	*	853	1,220,500	54.2	(44.9 - 63.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	307	4,100	62.9	(55.1 - 70.7)	15,427	3,436,000	61.9	(57.2 - 66.6)
Some College	*	*	*	*	9,798	1,726,200	49.3	(44.3 - 54.2)
≥ College Degree	172	1,500	37.5	(28.8 - 46.3)	12,351	2,242,600	40.1	(36.5 - 43.6)
Income								
≤ \$24,999	*	*	*	*	9,483	2,171,900	62.3	(55.8 - 68.8)
\$25,000-\$49,999	193	2,200	49.9	(40.2 - 59.7)	9,552	1,577,400	54.7	(48.3 - 61.0)
\$50,000-\$74,999	*	*	*	*	5,744	1,000,700	51.5	(45.6 - 57.4)
≥ \$75,000	*	*	*	*	8,516	1,738,200	38.8	(35.3 - 42.2)

¹ Defined as having had at least one permanent tooth extracted because of tooth decay or gum disease

² Sample size from survey

³ Estimated number of adults who have had at least one permanent tooth extracted because of tooth decay or gum disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 16b. Permanent teeth extracted due to decay or gum disease among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	657	48.9	(43.4 - 54.5)	37,652	49.4	(46.9 - 51.9)
Gender						
Male	276	46.1	(38.0 - 54.2)	14,457	47.9	(44.1 - 51.7)
Female	381	51.7	(44.2 - 59.2)	23,195	50.7	(47.4 - 54.0)
Race/Ethnicity						
White Non-Hispanic	644	49.2	(43.7 - 54.8)	35,364	46.5	(44.5 - 48.5)
Black Non-Hispanic	*	*	*	727	57.7	(48.6 - 66.9)
Hispanic	*	*	*	853	53.8	(44.6 - 62.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	307	60.7	(52.2 - 69.2)	15,427	61.3	(56.7 - 66.0)
Some College	*	*	*	9,798	47.9	(43.0 - 52.9)
≥ College Degree	172	35.1	(25.8 - 44.5)	12,351	39.0	(35.5 - 42.5)
Income						
≤ \$24,999	*	*	*	9,483	61.6	(55.2 - 68.1)
\$25,000-\$49,999	*	*	*	9,552	54.2	(48.0 - 60.3)
\$50,000-\$74,999	*	*	*	5,744	49.9	(44.0 - 55.9)
≥ \$75,000	*	*	*	8,516	37.7	(34.2 - 41.1)

¹ Defined as having had at least one permanent tooth extracted because of tooth decay or gum disease

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 17a. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	228	700	24.8	(18.7 - 30.8)	12,339	491,200	19.6	(16.0 - 23.3)
Gender								
Male	91	300	23.4	(14.3 - 32.6)	4,352	167,200	16.4	(11.8 - 20.9)
Female	137	400	25.8	(17.7 - 33.9)	7,987	324,000	21.8	(16.6 - 27.0)
Age (years)								
≥ 65	228	700	24.8	(18.7 - 30.8)	12,339	491,200	19.6	(16.0 - 23.3)
Race/Ethnicity								
White Non-Hispanic	223	700	24.2	(18.1 - 30.3)	11,885	322,600	16.8	(14.3 - 19.2)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	128	500	33.9	(25.1 - 42.8)	6,358	314,100	25.3	(19.9 - 30.7)
Some College	*	*	*	*	2,736	90,600	20.2	(12.1 - 28.2)
≥ College Degree	*	*	*	*	3,214	80,500	10.1	(3.6 - 16.5)
Income								
≤ \$24,999	*	*	*	*	4,558	199,600	26.9	(19.3 - 34.4)
\$25,000-\$49,999	*	*	*	*	3,365	123,900	21.4	(13.1 - 29.6)
\$50,000-\$74,999	*	*	*	*	1,179	21,400	7.3	(3.8 - 10.8)
≥ \$75,000	*	*	*	*	1,187	9,600	2.5	(1.0 - 4.0)

¹ Defined as having all permanent teeth extracted because of tooth decay or gum disease among adults aged 65 and older

² Sample size from survey

³ Estimated number of adults aged 65 and older who have had all permanent teeth extracted because of tooth decay or gum disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 17b. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	228	24.6	(18.5 - 30.7)	12,339	19.4	(15.7 - 23.2)
Gender						
Male	91	23.5	(14.3 - 32.6)	4,352	16.5	(11.5 - 21.4)
Female	137	25.8	(17.7 - 33.9)	7,987	22.1	(16.6 - 27.7)
Race/Ethnicity						
White Non-Hispanic	223	24.3	(18.1 - 30.4)	11,885	16.6	(13.9 - 19.3)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	128	33.9	(24.9 - 42.8)	6,358	24.9	(19.3 - 30.5)
Some College	*	*	*	2,736	21.0	(12.1 - 29.9)
≥ College Degree	*	*	*	3,214	10.0	(3.6 - 16.3)
Income						
≤ \$24,999	*	*	*	4,558	27.3	(19.1 - 35.5)
\$25,000-\$49,999	*	*	*	3,365	21.0	(12.9 - 29.1)
\$50,000-\$74,999	*	*	*	1,179	7.6	(3.6 - 11.5)
≥ \$75,000	*	*	*	1,187	2.2	(0.9 - 3.5)

¹ Defined as having all permanent teeth extracted because of tooth decay or gum disease among adults aged 65 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 18a. Flu shot within the past 12 months¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	668	5,600	37.4	(32.8 - 42.0)	38,079	6,255,900	42.3	(39.9 - 44.8)
Gender								
Male	279	2,300	31.1	(24.6 - 37.6)	14,579	2,798,900	39.6	(35.8 - 43.5)
Female	389	3,300	43.6	(37.2 - 49.9)	23,500	3,456,900	44.8	(41.6 - 47.9)
Age (years)								
18-49	189	1,900	24.3	(17.4 - 31.1)	12,505	2,685,300	31.1	(27.3 - 34.9)
50-64	245	1,800	43.8	(36.8 - 50.8)	12,972	1,662,900	46.0	(42.6 - 49.5)
≥ 65	234	1,900	63.6	(57.0 - 70.3)	12,602	1,907,700	75.2	(72.0 - 78.4)
Race/Ethnicity								
White Non-Hispanic	655	5,400	37.4	(32.8 - 42.1)	35,770	4,029,000	43.3	(41.4 - 45.3)
Black Non-Hispanic	*	*	*	*	738	907,000	42.1	(33.1 - 51.1)
Hispanic	*	*	*	*	860	882,900	39.6	(30.4 - 48.9)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	313	2,500	37.0	(30.3 - 43.7)	15,663	2,457,800	44.3	(39.8 - 48.8)
Some College	182	1,700	38.7	(29.6 - 47.9)	9,898	1,377,700	38.9	(34.0 - 43.9)
≥ College Degree	173	1,500	36.7	(27.8 - 45.6)	12,436	2,404,400	43.0	(39.5 - 46.4)
Income								
≤ \$24,999	193	1,400	37.1	(28.0 - 46.1)	9,635	1,656,800	47.6	(41.2 - 54.0)
\$25,000-\$49,999	195	1,500	33.9	(25.7 - 42.2)	9,647	1,200,600	41.4	(35.6 - 47.3)
\$50,000-\$74,999	*	*	*	*	5,763	792,300	40.2	(34.3 - 46.2)
≥ \$75,000	*	*	*	*	8,548	1,776,100	39.6	(36.3 - 42.8)

¹ Defined as having had a flu shot within the past 12 months

² Sample size from survey

³ Estimated number of adults who have had a flu shot within the past 12 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 18b. Flu shot within the past 12 months¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	668	35.0	(30.1 - 39.9)	38,079	41.7	(39.2 - 44.1)
Gender						
Male	279	29.7	(23.0 - 36.4)	14,579	40.0	(36.2 - 43.8)
Female	389	40.3	(33.3 - 47.3)	23,500	43.1	(40.0 - 46.3)
Race/Ethnicity						
White Non-Hispanic	655	35.1	(30.2 - 40.1)	35,770	42.4	(40.4 - 44.3)
Black Non-Hispanic	*	*	*	738	42.3	(33.3 - 51.3)
Hispanic	*	*	*	860	39.0	(29.9 - 48.1)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	313	34.8	(27.6 - 41.9)	15,663	43.7	(39.1 - 48.2)
Some College	182	36.0	(26.4 - 45.6)	9,898	37.8	(33.0 - 42.7)
≥ College Degree	173	34.5	(25.0 - 44.0)	12,436	42.5	(39.1 - 46.0)
Income						
≤ \$24,999	193	34.4	(24.6 - 44.2)	9,635	47.3	(41.0 - 53.7)
\$25,000-\$49,999	195	31.7	(23.1 - 40.3)	9,647	40.3	(34.5 - 46.0)
\$50,000-\$74,999	*	*	*	5,763	39.4	(33.4 - 45.4)
≥ \$75,000	*	*	*	8,548	39.1	(35.8 - 42.4)

¹ Defined as having had a flu shot within the past 12 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 19a. Flu vaccine that was sprayed in nose within the past 12 months¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	*	*	*	*	38,106	248,300	1.7	(0.6 - 2.8)
Gender								
Male	*	*	*	*	14,586	148,000	2.1	(0.0 - 4.2)
Female	*	*	*	*	23,520	100,300	1.3	(0.5 - 2.1)
Age (years)								
18-49	*	*	*	*	12,535	184,900	2.1	(0.3 - 3.9)
50-64	*	*	*	*	12,968	36,000	1.0	(0.2 - 1.8)
≥ 65	*	*	*	*	12,603	27,400	1.1	(0.4 - 1.8)
Race/Ethnicity								
White Non-Hispanic	*	*	*	*	35,796	66,000	0.7	(0.4 - 1.0)
Black Non-Hispanic	*	*	*	*	740	38,500	1.8	(0.0 - 4.1)
Hispanic	*	*	*	*	860	133,700	5.9	(0.0 - 12.3)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,679	172,500	3.1	(0.3 - 5.9)
Some College	*	*	*	*	9,897	43,500	1.2	(0.4 - 2.0)
≥ College Degree	*	*	*	*	12,449	31,600	0.6	(0.3 - 0.8)
Income								
≤ \$24,999	*	*	*	*	9,636	138,400	3.9	(0.0 - 8.1)
\$25,000-\$49,999	*	*	*	*	9,648	41,500	1.4	(0.0 - 3.1)
\$50,000-\$74,999	*	*	*	*	5,766	13,300	0.7	(0.2 - 1.2)
≥ \$75,000	*	*	*	*	8,561	35,300	0.8	(0.3 - 1.3)

¹ Defined as having had flu vaccine sprayed in nose within the past 12 months

² Sample size from survey

³ Estimated number of adults who have had flu vaccine sprayed in nose within the past 12 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 19b. Flu vaccine that was sprayed in nose within the past 12 months¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	*	*	*	38,106	1.7	(0.6 - 2.7)
Gender						
Male	*	*	*	14,586	2.0	(0.1 - 3.9)
Female	*	*	*	23,520	1.4	(0.5 - 2.3)
Race/Ethnicity						
White Non-Hispanic	*	*	*	35,796	0.7	(0.4 - 1.0)
Black Non-Hispanic	*	*	*	740	1.9	(0.0 - 4.4)
Hispanic	*	*	*	860	5.9	(0.0 - 11.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	15,679	3.1	(0.4 - 5.8)
Some College	*	*	*	9,897	1.2	(0.4 - 2.0)
≥ College Degree	*	*	*	12,449	0.6	(0.3 - 0.8)
Income						
≤ \$24,999	*	*	*	9,636	3.8	(0.0 - 7.7)
\$25,000-\$49,999	*	*	*	9,648	1.5	(0.0 - 3.4)
\$50,000-\$74,999	*	*	*	5,766	0.7	(0.1 - 1.2)
≥ \$75,000	*	*	*	8,561	0.8	(0.3 - 1.3)

¹ Defined as having had flu vaccine sprayed in nose within the past 12 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 20a. Flu shot or flu vaccine sprayed in nose within the past 12 months¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	666	5,700	38.0	(33.3 - 42.6)	38,059	6,302,300	42.7	(40.2 - 45.1)
Gender								
Male	278	2,400	32.1	(25.5 - 38.7)	14,568	2,822,700	40.0	(36.1 - 43.9)
Female	388	3,300	43.7	(37.3 - 50.0)	23,491	3,479,500	45.1	(42.0 - 48.2)
Age (years)								
18-49	189	2,000	24.7	(17.8 - 31.6)	12,501	2,726,400	31.6	(27.8 - 35.4)
50-64	245	1,800	44.8	(37.8 - 51.8)	12,964	1,664,200	46.1	(42.7 - 49.5)
≥ 65	232	1,900	64.1	(57.5 - 70.8)	12,594	1,911,700	75.4	(72.2 - 78.6)
Race/Ethnicity								
White Non-Hispanic	653	5,500	38.0	(33.4 - 42.7)	35,750	4,048,700	43.6	(41.7 - 45.5)
Black Non-Hispanic	*	*	*	*	738	910,000	42.2	(33.2 - 51.2)
Hispanic	*	*	*	*	860	903,300	40.6	(31.3 - 49.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	312	2,500	37.2	(30.5 - 43.9)	15,651	2,480,100	44.7	(40.2 - 49.3)
Some College	181	1,700	38.9	(29.7 - 48.1)	9,895	1,387,300	39.2	(34.3 - 44.1)
≥ College Degree	173	1,500	38.2	(29.2 - 47.2)	12,431	2,418,300	43.2	(39.8 - 46.7)
Income								
≤ \$24,999	192	1,400	37.1	(28.1 - 46.2)	9,626	1,675,100	48.2	(41.8 - 54.6)
\$25,000-\$49,999	195	1,500	33.9	(25.7 - 42.2)	9,643	1,208,300	41.7	(35.8 - 47.6)
\$50,000-\$74,999	*	*	*	*	5,761	795,600	40.4	(34.5 - 46.3)
≥ \$75,000	*	*	*	*	8,548	1,791,200	39.9	(36.6 - 43.2)

¹ Defined as having had a flu shot or flu vaccine sprayed in nose within the past 12 months

² Sample size from survey

³ Estimated number of adults who have had a flu shot or flu vaccine sprayed in nose within the past 12 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 20b. Flu shot or flu vaccine sprayed in nose within the past 12 months¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	666	35.6	(30.7 - 40.6)	38,059	42.0	(39.6 - 44.5)
Gender						
Male	278	30.8	(24.0 - 37.7)	14,568	40.4	(36.6 - 44.2)
Female	388	40.3	(33.3 - 47.3)	23,491	43.5	(40.3 - 46.7)
Race/Ethnicity						
White Non-Hispanic	653	35.7	(30.7 - 40.7)	35,750	42.6	(40.7 - 44.6)
Black Non-Hispanic	*	*	*	738	42.4	(33.4 - 51.4)
Hispanic	*	*	*	860	40.0	(30.9 - 49.1)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	312	35.0	(27.8 - 42.2)	15,651	44.1	(39.6 - 48.6)
Some College	181	36.1	(26.5 - 45.7)	9,895	38.1	(33.2 - 43.0)
≥ College Degree	173	36.2	(26.6 - 45.8)	12,431	42.8	(39.4 - 46.3)
Income						
≤ \$24,999	192	34.5	(24.7 - 44.3)	9,626	48.0	(41.6 - 54.3)
\$25,000-\$49,999	195	31.7	(23.1 - 40.3)	9,643	40.5	(34.8 - 46.3)
\$50,000-\$74,999	*	*	*	5,761	39.6	(33.6 - 45.6)
≥ \$75,000	*	*	*	8,548	39.4	(36.1 - 42.7)

¹ Defined as having had a flu shot or flu vaccine sprayed in nose within the past 12 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 21a. Flu shot among adults aged 65 and older¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	234	1,900	63.6	(57.0 - 70.3)	12,602	1,907,700	75.2	(72.0 - 78.4)
Gender								
Male	*	*	*	*	4,417	772,900	74.6	(69.3 - 79.8)
Female	142	1,100	65.6	(57.1 - 74.1)	8,185	1,134,800	75.6	(71.6 - 79.6)
Age (years)								
≥ 65	234	1,900	63.6	(57.0 - 70.3)	12,602	1,907,700	75.2	(72.0 - 78.4)
Race/Ethnicity								
White Non-Hispanic	229	1,800	63.8	(57.0 - 70.5)	12,137	1,487,100	76.0	(73.5 - 78.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	130	1,000	59.8	(50.7 - 69.0)	6,518	930,800	74.0	(68.9 - 79.2)
Some College	*	*	*	*	2,790	317,800	69.7	(62.7 - 76.7)
≥ College Degree	*	*	*	*	3,261	654,000	80.5	(76.4 - 84.6)
Income								
≤ \$24,999	*	*	*	*	4,665	545,300	72.4	(65.6 - 79.3)
\$25,000-\$49,999	*	*	*	*	3,422	430,600	72.8	(67.0 - 78.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	1,198	301,000	77.5	(70.9 - 84.2)

¹ Defined as having had a flu shot during the past 12 months among adults aged 65 and older

² Sample size from survey

³ Estimated number of adults aged 65 and older who have had a flu shot within the past 12 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 21b. Flu shot among adults aged 65 and older¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	234	63.4	(56.6 - 70.2)	12,602	75.0	(71.5 - 78.5)
Gender						
Male	*	*	*	4,417	74.4	(68.7 - 80.1)
Female	142	65.6	(57.1 - 74.1)	8,185	75.5	(71.3 - 79.8)
Race/Ethnicity						
White Non-Hispanic	229	63.5	(56.6 - 70.3)	12,137	75.9	(73.0 - 78.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	130	59.2	(49.8 - 68.6)	6,518	73.3	(67.5 - 79.1)
Some College	*	*	*	2,790	70.0	(62.7 - 77.2)
≥ College Degree	*	*	*	3,261	80.7	(76.2 - 85.1)
Income						
≤ \$24,999	*	*	*	4,665	72.6	(65.1 - 80.1)
\$25,000-\$49,999	*	*	*	3,422	71.4	(65.0 - 77.9)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	1,198	78.0	(70.8 - 85.1)

¹ Defined as having had a flu shot during the past 12 months among adults aged 65 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 22a. Pneumonia shot or pneumococcal vaccine¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	625	3,900	28.3	(23.9 - 32.6)	35,303	3,456,800	26.1	(23.9 - 28.3)
Gender								
Male	256	1,800	26.0	(19.1 - 32.8)	13,052	1,542,200	25.2	(21.5 - 28.8)
Female	369	2,100	30.5	(25.1 - 35.9)	22,251	1,914,600	26.9	(24.2 - 29.6)
Age (years)								
18-49	172	1,000	14.7	(7.9 - 21.5)	10,943	1,097,000	14.6	(11.3 - 17.9)
50-64	235	900	23.6	(17.7 - 29.6)	12,179	758,000	23.0	(19.8 - 26.2)
≥ 65	218	1,900	68.9	(62.1 - 75.7)	12,181	1,601,700	65.7	(61.7 - 69.7)
Race/Ethnicity								
White Non-Hispanic	613	3,600	27.6	(23.6 - 31.7)	33,225	2,358,700	28.7	(26.9 - 30.5)
Black Non-Hispanic	*	*	*	*	685	584,400	28.4	(19.5 - 37.3)
Hispanic	*	*	*	*	769	408,000	19.5	(12.6 - 26.4)
Other Non-Hispanic	*	*	*	*	624	105,600	12.2	(2.3 - 22.2)
Education								
≤ High School	298	1,800	28.2	(22.5 - 33.9)	14,700	1,535,300	30.4	(26.4 - 34.3)
Some College	*	*	*	*	9,151	836,200	27.2	(21.9 - 32.5)
≥ College Degree	162	900	24.4	(16.9 - 31.8)	11,375	1,070,700	21.4	(18.6 - 24.1)
Income								
≤ \$24,999	179	1,200	35.2	(26.2 - 44.1)	9,082	1,050,900	33.1	(27.6 - 38.6)
\$25,000-\$49,999	183	1,300	31.0	(21.8 - 40.3)	9,033	797,500	30.3	(24.1 - 36.6)
\$50,000-\$74,999	104	600	27.2	(17.2 - 37.1)	5,320	378,800	21.8	(17.4 - 26.1)
≥ \$75,000	96	300	14.1	(7.3 - 21.0)	7,773	648,700	16.3	(14.1 - 18.5)

¹ Defined as having ever had a pneumonia shot or pneumococcal vaccine

² Sample size from survey

³ Estimated number of adults who have ever had a pneumonia shot or pneumococcal vaccine (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 22b. Pneumonia shot or pneumococcal vaccine¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	625	26.0	(21.4 - 30.5)	35,303	25.4	(23.2 - 27.6)
Gender						
Male	256	25.0	(17.9 - 32.1)	13,052	25.8	(22.3 - 29.4)
Female	369	26.9	(21.2 - 32.6)	22,251	25.1	(22.4 - 27.8)
Race/Ethnicity						
White Non-Hispanic	613	25.2	(21.0 - 29.4)	33,225	27.7	(25.9 - 29.5)
Black Non-Hispanic	*	*	*	685	27.9	(19.1 - 36.7)
Hispanic	*	*	*	769	19.6	(12.8 - 26.5)
Other Non-Hispanic	*	*	*	624	11.3	(2.0 - 20.7)
Education						
≤ High School	298	25.5	(19.7 - 31.4)	14,700	29.5	(25.6 - 33.5)
Some College	*	*	*	9,151	26.2	(21.0 - 31.3)
≥ College Degree	162	22.5	(14.8 - 30.2)	11,375	21.0	(18.3 - 23.8)
Income						
≤ \$24,999	179	32.3	(22.7 - 41.8)	9,082	32.3	(26.9 - 37.8)
\$25,000-\$49,999	183	28.3	(18.7 - 37.8)	9,033	29.0	(23.0 - 35.0)
\$50,000-\$74,999	*	*	*	5,320	21.2	(16.8 - 25.7)
≥ \$75,000	96	13.4	(6.6 - 20.2)	7,773	16.2	(13.9 - 18.5)

¹ Defined as having ever had a pneumonia shot or pneumococcal vaccine

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 23a. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	218	1,900	68.9	(62.1 - 75.7)	12,181	1,601,700	65.7	(61.7 - 69.7)
Gender								
Male	*	*	*	*	4,180	606,400	62.1	(55.5 - 68.7)
Female	135	1,200	72.5	(63.9 - 81.0)	8,001	995,300	68.0	(63.1 - 73.0)
Age (years)								
≥ 65	218	1,900	68.9	(62.1 - 75.7)	12,181	1,601,700	65.7	(61.7 - 69.7)
Race/Ethnicity								
White Non-Hispanic	213	1,900	70.8	(64.1 - 77.6)	11,735	1,281,300	68.2	(64.9 - 71.5)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	124	1,100	67.1	(57.8 - 76.3)	6,345	806,200	65.6	(59.5 - 71.7)
Some College	*	*	*	*	2,694	289,500	66.8	(60.1 - 73.4)
≥ College Degree	*	*	*	*	3,111	501,100	65.6	(58.5 - 72.7)
Income								
≤ \$24,999	*	*	*	*	4,533	497,100	67.7	(59.9 - 75.5)
\$25,000-\$49,999	*	*	*	*	3,324	344,300	61.5	(53.3 - 69.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	1,138	236,300	64.5	(55.2 - 73.8)

¹ Defined as having ever had a pneumonia shot or pneumococcal vaccine among adults aged 65 and older

² Sample size from survey

³ Estimated number of adults aged 65 and older who have ever had a pneumonia shot or pneumococcal vaccine (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 23b. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	218	68.4	(61.5 - 75.4)	12,181	64.7	(60.4 - 69.1)
Gender						
Male	*	*	*	4,180	61.5	(54.3 - 68.6)
Female	135	72.5	(63.9 - 81.1)	8,001	67.6	(62.4 - 72.9)
Race/Ethnicity						
White Non-Hispanic	213	70.3	(63.4 - 77.2)	11,735	67.1	(63.2 - 70.9)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	124	66.4	(56.9 - 75.8)	6,345	64.2	(57.4 - 71.0)
Some College	*	*	*	2,694	65.8	(58.4 - 73.3)
≥ College Degree	*	*	*	3,111	65.3	(57.7 - 72.9)
Income						
≤ \$24,999	*	*	*	4,533	66.0	(57.0 - 74.9)
\$25,000-\$49,999	*	*	*	3,324	60.3	(51.9 - 68.6)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	*	*

¹ Defined as having ever had a pneumonia shot or pneumococcal vaccine among adults aged 65 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 24a. Ever had a mammogram among women aged 40 and older¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	340	4,900	94.3	(91.4 - 97.1)	19,491	4,328,800	91.8	(89.7 - 94.0)
Gender								
Male	-	-	-	-	-	-	-	-
Female	340	4,900	94.3	(91.4 - 97.1)	19,491	4,328,800	91.8	(89.7 - 94.0)
Age (years)								
40-49	60	1,300	89.9	(81.9 - 98.0)	3,851	1,287,500	85.0	(79.7 - 90.3)
50-64	137	2,000	97.7	(95.3 - 100)	7,452	1,641,700	96.0	(94.1 - 97.9)
≥ 65	143	1,600	93.8	(89.9 - 97.8)	8,188	1,399,600	94.1	(90.5 - 97.6)
Race/Ethnicity								
White Non-Hispanic	335	4,700	94.1	(91.2 - 97.0)	18,548	2,976,100	93.9	(92.7 - 95.1)
Black Non-Hispanic	*	*	*	*	342	746,600	93.8	(87.3 - 100)
Hispanic	*	*	*	*	333	447,600	91.8	(84.6 - 99.0)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	160	2,100	94.8	(91.1 - 98.5)	8,266	1,698,700	91.0	(87.4 - 94.5)
Some College	100	1,500	94.9	(90.5 - 99.4)	5,322	1,055,500	93.6	(90.2 - 97.0)
≥ College Degree	80	1,200	92.6	(85.5 - 99.8)	5,872	1,560,700	91.6	(87.5 - 95.6)
Income								
≤ \$24,999	111	1,300	90.1	(83.8 - 96.5)	5,691	1,023,400	93.5	(89.7 - 97.4)
\$25,000-\$49,999	88	1,300	97.0	(92.8 - 100)	4,857	860,300	89.5	(83.6 - 95.5)
\$50,000-\$74,999	52	800	93.0	(84.4 - 100)	2,648	547,100	94.6	(92.2 - 96.9)
≥ \$75,000	*	*	*	*	3,582	1,185,100	92.2	(88.2 - 96.1)

¹ Defined as having ever had a mammogram among women aged 40 and older

² Sample size from survey

³ Estimated number of women aged 40 and older who have ever had a mammogram (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 24b. Ever had a mammogram among women aged 40 and older¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	340	93.6	(90.1 - 97.1)	19,491	91.1	(88.7 - 93.6)
Gender						
Male	-	-	-	-	-	-
Female	340	93.6	(90.1 - 97.1)	19,491	91.1	(88.7 - 93.6)
Race/Ethnicity						
White Non-Hispanic	335	93.5	(89.9 - 97.1)	18,548	93.3	(92.0 - 94.7)
Black Non-Hispanic	*	*	*	342	93.1	(85.7 - 100)
Hispanic	*	*	*	333	91.0	(82.8 - 99.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	160	95.1	(91.4 - 98.7)	8,266	90.2	(86.1 - 94.2)
Some College	100	94.7	(89.4 - 100)	5,322	92.9	(88.9 - 96.9)
≥ College Degree	80	90.2	(80.4 - 100)	5,872	90.9	(86.6 - 95.3)
Income						
≤ \$24,999	111	89.3	(81.8 - 96.8)	5,691	93.0	(88.5 - 97.4)
\$25,000-\$49,999	88	96.3	(90.8 - 100)	4,857	88.8	(82.5 - 95.1)
\$50,000-\$74,999	*	*	*	2,648	94.3	(91.7 - 96.8)
≥ \$75,000	*	*	*	3,582	91.2	(86.8 - 95.6)

¹ Defined as having ever had a mammogram among women aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 25a. Had mammogram within the past 2 years among women aged 40 and older¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	340	4,100	79.7	(74.9 - 84.4)	19,376	3,760,000	80.3	(77.8 - 82.9)
Gender								
Male	-	-	-	-	-	-	-	-
Female	340	4,100	79.7	(74.9 - 84.4)	19,376	3,760,000	80.3	(77.8 - 82.9)
Age (years)								
40-49	*	*	*	*	3,842	1,122,700	74.4	(68.5 - 80.2)
50-64	137	1,700	82.9	(76.3 - 89.5)	7,428	1,487,900	87.2	(84.5 - 89.8)
≥ 65	143	1,300	78.1	(71.1 - 85.2)	8,106	1,149,400	78.5	(74.2 - 82.9)
Race/Ethnicity								
White Non-Hispanic	335	4,100	80.4	(75.8 - 85.0)	18,438	2,531,500	80.2	(78.3 - 82.2)
Black Non-Hispanic	*	*	*	*	339	667,100	85.6	(77.5 - 93.8)
Hispanic	*	*	*	*	332	419,700	86.1	(77.5 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	160	1,800	80.6	(74.1 - 87.1)	8,204	1,486,900	80.5	(76.4 - 84.6)
Some College	100	1,200	77.6	(68.5 - 86.6)	5,297	929,400	83.0	(78.8 - 87.1)
≥ College Degree	*	*	*	*	5,845	1,334,800	78.6	(73.9 - 83.2)
Income								
≤ \$24,999	*	*	*	*	5,650	882,300	80.8	(75.9 - 85.7)
\$25,000-\$49,999	88	1,100	84.4	(76.4 - 92.3)	4,846	732,300	76.3	(69.7 - 82.9)
\$50,000-\$74,999	*	*	*	*	2,641	494,200	85.6	(81.9 - 89.4)
≥ \$75,000	*	*	*	*	3,572	1,025,900	80.0	(75.2 - 84.8)

¹ Defined as having had a mammogram within the past 2 years among women aged 40 and older

² Sample size from survey

³ Estimated number of women aged 40 and older who have had a mammogram within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 25b. Had mammogram within the past 2 years among women aged 40 and older¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	340	79.7	(74.3 - 85.0)	19,376	79.7	(76.9 - 82.5)
Gender						
Male	-	-	-	-	-	-
Female	340	79.7	(74.3 - 85.0)	19,376	79.7	(76.9 - 82.5)
Race/Ethnicity						
White Non-Hispanic	335	80.4	(75.2 - 85.6)	18,438	79.7	(77.6 - 81.8)
Black Non-Hispanic	*	*	*	339	85.3	(76.5 - 94.2)
Hispanic	*	*	*	332	85.2	(75.6 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	160	81.6	(74.9 - 88.4)	8,204	80.1	(75.6 - 84.7)
Some College	100	78.6	(69.1 - 88.1)	5,297	82.3	(77.7 - 86.9)
≥ College Degree	*	*	*	5,845	77.8	(72.9 - 82.8)
Income						
≤ \$24,999	*	*	*	5,650	80.8	(75.4 - 86.2)
\$25,000-\$49,999	88	84.3	(75.4 - 93.3)	4,846	75.7	(68.7 - 82.6)
\$50,000-\$74,999	*	*	*	2,641	85.5	(81.6 - 89.4)
≥ \$75,000	*	*	*	3,572	79.0	(73.8 - 84.2)

¹ Defined as having had a mammogram within the past 2 years among women aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 26a. Had mammogram within the past 2 years among women aged 50 and older¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	280	3,000	80.7	(75.9 - 85.5)	15,534	2,637,300	83.2	(80.7 - 85.7)
Gender								
Male	-	-	-	-	-	-	-	-
Female	280	3,000	80.7	(75.9 - 85.5)	15,534	2,637,300	83.2	(80.7 - 85.7)
Age (years)								
50-64	137	1,700	82.9	(76.3 - 89.5)	7,428	1,487,900	87.2	(84.5 - 89.8)
≥ 65	143	1,300	78.1	(71.1 - 85.2)	8,106	1,149,400	78.5	(74.2 - 82.9)
Race/Ethnicity								
White Non-Hispanic	276	2,900	80.9	(76.1 - 85.7)	14,887	1,907,000	82.3	(80.3 - 84.4)
Black Non-Hispanic	*	*	*	*	249	385,600	89.9	(83.2 - 96.7)
Hispanic	*	*	*	*	220	286,700	93.9	(89.1 - 98.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	140	1,400	79.2	(72.1 - 86.2)	7,026	1,094,500	83.3	(80.0 - 86.6)
Some College	81	900	77.1	(67.4 - 86.8)	4,124	646,700	87.3	(84.2 - 90.3)
≥ College Degree	59	800	88.6	(80.6 - 96.5)	4,356	890,500	80.6	(75.1 - 86.1)
Income								
≤ \$24,999	98	800	71.0	(61.0 - 80.9)	4,991	649,100	81.0	(76.8 - 85.2)
\$25,000-\$49,999	71	800	84.0	(75.1 - 92.9)	3,913	533,300	78.6	(70.9 - 86.2)
\$50,000-\$74,999	*	*	*	*	1,943	359,300	89.6	(86.3 - 93.0)
≥ \$75,000	*	*	*	*	2,303	601,300	84.6	(79.3 - 89.9)

¹ Defined as having had a mammogram within the past 2 years among women aged 50 and older

² Sample size from survey

³ Estimated number of women aged 50 and older who have had a mammogram within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 26b. Had mammogram within the past 2 years among women aged 50 and older¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	280	80.9	(76.0 - 85.8)	15,534	83.1	(80.3 - 85.8)
Gender						
Male	-	-	-	-	-	-
Female	280	80.9	(76.0 - 85.8)	15,534	83.1	(80.3 - 85.8)
Race/Ethnicity						
White Non-Hispanic	276	81.1	(76.2 - 86.0)	14,887	82.1	(79.9 - 84.3)
Black Non-Hispanic	*	*	*	249	90.4	(83.6 - 97.2)
Hispanic	*	*	*	220	94.1	(89.3 - 99.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	140	79.2	(72.1 - 86.3)	7,026	83.6	(80.1 - 87.0)
Some College	81	77.5	(67.6 - 87.3)	4,124	87.3	(84.1 - 90.5)
≥ College Degree	59	88.6	(80.4 - 96.9)	4,356	79.9	(73.9 - 86.0)
Income						
≤ \$24,999	*	*	*	4,991	81.3	(76.9 - 85.6)
\$25,000-\$49,999	71	84.1	(75.1 - 93.2)	3,913	78.1	(69.9 - 86.4)
\$50,000-\$74,999	*	*	*	1,943	90.2	(87.1 - 93.4)
≥ \$75,000	*	*	*	2,303	83.9	(77.9 - 89.9)

¹ Defined as having had a mammogram within the past 2 years among women aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 27a. Ever had a Pap test among women¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	389	7,300	95.9	(93.1 - 98.8)	23,381	7,045,300	92.1	(90.0 - 94.3)
Gender								
Male	-	-	-	-	-	-	-	-
Female	389	7,300	95.9	(93.1 - 98.8)	23,381	7,045,300	92.1	(90.0 - 94.3)
Age (years)								
18-34	*	*	*	*	2,377	1,896,600	85.2	(79.3 - 91.0)
35-44	*	*	*	*	3,023	1,301,500	92.6	(87.7 - 97.5)
45-54	72	1,500	99.4	(98.1 - 100)	4,657	1,420,300	98.3	(97.0 - 99.6)
55-64	101	1,200	99.5	(98.4 - 100)	5,187	1,071,200	97.2	(95.0 - 99.3)
≥ 65	142	1,600	94.1	(90.3 - 97.8)	8,137	1,355,700	92.3	(88.5 - 96.1)
Race/Ethnicity								
White Non-Hispanic	383	7,100	95.8	(92.9 - 98.8)	21,976	4,439,700	94.2	(92.1 - 96.3)
Black Non-Hispanic	*	*	*	*	481	1,231,600	91.2	(84.0 - 98.4)
Hispanic	*	*	*	*	533	984,500	90.1	(85.3 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	181	3,100	97.3	(95.4 - 99.1)	9,526	2,540,500	88.3	(83.8 - 92.7)
Some College	113	2,100	91.3	(82.6 - 100)	6,462	1,782,200	93.5	(90.4 - 96.6)
≥ College Degree	95	2,100	98.9	(96.6 - 100)	7,345	2,643,200	95.0	(92.2 - 97.9)
Income								
≤ \$24,999	123	1,900	94.7	(91.0 - 98.4)	6,526	1,717,200	93.7	(90.9 - 96.6)
\$25,000-\$49,999	100	1,800	94.9	(88.4 - 100)	5,787	1,378,400	91.0	(85.6 - 96.4)
\$50,000-\$74,999	63	1,400	99.5	(98.4 - 100)	3,310	933,900	96.0	(91.8 - 100)
≥ \$75,000	54	1,300	100	(100 - 100)	4,570	1,943,300	95.4	(92.6 - 98.2)

¹ Defined as having ever had a Pap test among women

² Sample size from survey

³ Estimated number of women who have ever had a Pap test (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 27b. Ever had a Pap test among women¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	389	95.6	(92.2 - 99.1)	23,381	91.8	(89.6 - 94.0)
Gender						
Male	-	-	-	-	-	-
Female	389	95.6	(92.2 - 99.1)	23,381	91.8	(89.6 - 94.0)
Race/Ethnicity						
White Non-Hispanic	383	95.5	(92.0 - 99.1)	21,976	94.0	(91.8 - 96.2)
Black Non-Hispanic	*	*	*	481	90.8	(83.5 - 98.1)
Hispanic	*	*	*	533	89.7	(84.6 - 94.7)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	181	97.8	(96.3 - 99.3)	9,526	87.8	(83.2 - 92.4)
Some College	*	*	*	6,462	93.1	(89.7 - 96.4)
≥ College Degree	95	98.5	(95.7 - 100)	7,345	94.9	(92.0 - 97.7)
Income						
≤ \$24,999	123	94.9	(90.8 - 99.0)	6,526	93.3	(90.1 - 96.5)
\$25,000-\$49,999	100	94.0	(86.4 - 100)	5,787	91.1	(85.7 - 96.5)
\$50,000-\$74,999	63	99.6	(98.8 - 100)	3,310	96.0	(91.9 - 100)
≥ \$75,000	54	100	(100 - 100)	4,570	95.0	(91.9 - 98.1)

¹ Defined as having ever had a Pap test among women

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 28a. Had a Pap test within the past 3 years among women¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	386	5,600	74.7	(69.7 - 79.8)	23,109	6,205,600	82.2	(79.8 - 84.6)
Gender								
Male	-	-	-	-	-	-	-	-
Female	386	5,600	74.7	(69.7 - 79.8)	23,109	6,205,600	82.2	(79.8 - 84.6)
Age (years)								
18-34	*	*	*	*	2,371	1,818,400	81.8	(75.5 - 88.1)
35-44	*	*	*	*	3,012	1,220,300	86.9	(81.8 - 92.0)
45-54	72	1,200	79.5	(70.0 - 89.1)	4,640	1,311,500	92.0	(89.8 - 94.3)
55-64	101	900	72.5	(63.1 - 81.9)	5,152	924,300	84.5	(81.0 - 88.0)
≥ 65	139	800	46.2	(37.1 - 55.4)	7,934	931,100	66.4	(61.8 - 71.0)
Race/Ethnicity								
White Non-Hispanic	380	5,500	74.7	(69.6 - 79.8)	21,720	3,807,100	81.5	(79.2 - 83.8)
Black Non-Hispanic	*	*	*	*	473	1,142,800	86.7	(79.1 - 94.3)
Hispanic	*	*	*	*	527	894,900	83.6	(76.3 - 91.0)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	178	2,300	73.5	(66.2 - 80.7)	9,384	2,132,600	76.0	(71.1 - 81.0)
Some College	*	*	*	*	6,401	1,594,300	84.0	(80.4 - 87.5)
≥ College Degree	95	1,800	83.5	(75.2 - 91.7)	7,281	2,406,000	86.9	(83.7 - 90.1)
Income								
≤ \$24,999	*	*	*	*	6,441	1,453,900	82.0	(77.9 - 86.0)
\$25,000-\$49,999	*	*	*	*	5,740	1,182,000	78.3	(72.6 - 83.9)
\$50,000-\$74,999	*	*	*	*	3,296	825,000	85.0	(78.0 - 91.9)
≥ \$75,000	54	1,200	93.0	(86.1 - 99.9)	4,555	1,810,200	89.0	(85.7 - 92.2)

¹ Defined as having had a Pap test within the past 3 years among women

² Sample size from survey

³ Estimated number of women who have had a Pap test within the past 3 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 28b. Had a Pap test within the past 3 years among women¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	386	77.7	(72.4 - 83.0)	23,109	82.7	(80.2 - 85.1)
Gender						
Male	-	-	-	-	-	-
Female	386	77.7	(72.4 - 83.0)	23,109	82.7	(80.2 - 85.1)
Race/Ethnicity						
White Non-Hispanic	380	77.6	(72.3 - 83.0)	21,720	82.3	(80.0 - 84.6)
Black Non-Hispanic	*	*	*	473	87.0	(79.3 - 94.6)
Hispanic	*	*	*	527	83.4	(75.9 - 90.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	178	77.1	(69.7 - 84.6)	9,384	76.4	(71.3 - 81.5)
Some College	*	*	*	6,401	84.2	(80.5 - 88.0)
≥ College Degree	95	85.2	(76.9 - 93.6)	7,281	87.5	(84.3 - 90.6)
Income						
≤ \$24,999	*	*	*	6,441	82.7	(78.5 - 86.9)
\$25,000-\$49,999	*	*	*	5,740	79.2	(73.5 - 84.8)
\$50,000-\$74,999	63	81.1	(71.3 - 90.8)	3,296	85.7	(78.7 - 92.8)
≥ \$75,000	54	93.7	(87.1 - 100)	4,555	88.9	(85.5 - 92.3)

¹ Defined as having had a Pap test within the past 3 years among women

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 29a. Ever had a digital rectal exam among men aged 40 and older¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	236	3,500	74.2	(67.4 - 81.0)	12,033	3,176,500	73.8	(69.7 - 77.8)
Gender								
Male	236	3,500	74.2	(67.4 - 81.0)	12,033	3,176,500	73.8	(69.7 - 77.8)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	3,098	922,800	66.8	(59.9 - 73.8)
55-64	72	1,100	85.1	(76.7 - 93.6)	3,504	865,600	87.1	(82.5 - 91.7)
≥ 65	92	1,100	88.2	(81.4 - 94.9)	4,348	953,900	92.9	(91.0 - 94.9)
Race/Ethnicity								
White Non-Hispanic	230	3,300	74.8	(68.0 - 81.6)	11,411	2,452,100	80.3	(77.7 - 82.9)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	5,013	931,600	64.1	(57.3 - 71.0)
Some College	*	*	*	*	2,695	674,200	75.5	(67.8 - 83.2)
≥ College Degree	*	*	*	*	4,302	1,568,000	80.5	(74.0 - 87.0)
Income								
≤ \$24,999	*	*	*	*	2,573	573,300	65.9	(56.0 - 75.9)
\$25,000-\$49,999	*	*	*	*	3,168	530,100	74.6	(67.2 - 82.1)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,309	1,351,100	78.0	(72.6 - 83.5)

¹ Defined as having ever had a digital rectal exam among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have ever had a digital rectal exam (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 29b. Ever had a digital rectal exam among men aged 40 and older¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	236	72.1	(64.4 - 79.8)	12,033	73.7	(69.6 - 77.8)
Gender						
Male	236	72.1	(64.4 - 79.8)	12,033	73.7	(69.6 - 77.8)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	230	72.4	(64.5 - 80.2)	11,411	79.9	(77.2 - 82.5)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	5,013	64.7	(57.8 - 71.7)
Some College	*	*	*	2,695	75.0	(67.1 - 82.9)
≥ College Degree	*	*	*	4,302	80.3	(73.8 - 86.8)
Income						
≤ \$24,999	*	*	*	2,573	66.9	(56.9 - 76.9)
\$25,000-\$49,999	*	*	*	3,168	74.8	(67.3 - 82.2)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,309	77.5	(72.0 - 83.0)

¹ Defined as having ever had a digital rectal exam among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 30a. Had digital rectal exam within the past 2 years among men aged 40 and older¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	234	2,400	51.1	(43.7 - 58.5)	11,923	2,380,300	55.8	(51.8 - 59.8)
Gender								
Male	234	2,400	51.1	(43.7 - 58.5)	11,923	2,380,300	55.8	(51.8 - 59.8)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	1,078	246,800	27.5	(18.5 - 36.5)
45-54	*	*	*	*	3,074	650,000	47.6	(40.8 - 54.4)
55-64	*	*	*	*	3,467	673,500	68.1	(62.4 - 73.8)
≥ 65	*	*	*	*	4,304	809,900	79.7	(75.0 - 84.3)
Race/Ethnicity								
White Non-Hispanic	228	2,200	51.0	(43.6 - 58.5)	11,305	1,875,600	62.2	(59.2 - 65.2)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	4,963	718,700	49.6	(43.2 - 56.1)
Some College	*	*	*	*	2,665	511,500	58.5	(50.8 - 66.3)
≥ College Degree	*	*	*	*	4,277	1,148,400	59.4	(52.8 - 66.0)
Income								
≤ \$24,999	*	*	*	*	2,553	388,000	45.0	(35.0 - 55.0)
\$25,000-\$49,999	*	*	*	*	3,139	422,400	60.5	(53.0 - 68.0)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,295	978,300	56.9	(51.1 - 62.7)

¹ Defined as having had a digital rectal exam within the past 2 years among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have had a digital rectal exam within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 30b. Had digital rectal exam within the past 2 years among men aged 40 and older¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	234	48.9	(41.0 - 56.7)	11,923	55.9	(51.7 - 60.0)
Gender						
Male	234	48.9	(41.0 - 56.7)	11,923	55.9	(51.7 - 60.0)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	228	48.6	(40.6 - 56.6)	11,305	62.0	(58.9 - 65.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	4,963	50.5	(43.7 - 57.3)
Some College	*	*	*	2,665	58.3	(50.4 - 66.2)
≥ College Degree	*	*	*	4,277	59.2	(52.4 - 65.9)
Income						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,139	60.8	(53.1 - 68.4)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,295	56.5	(50.5 - 62.4)

¹ Defined as having had a digital rectal exam within the past 2 years among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 31a. Ever had a prostate specific antigen test among men aged 40 and older¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	226	2,800	62.1	(54.6 - 69.6)	11,557	2,822,600	67.6	(63.9 - 71.2)
Gender								
Male	226	2,800	62.1	(54.6 - 69.6)	11,557	2,822,600	67.6	(63.9 - 71.2)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,938	760,600	57.4	(50.6 - 64.3)
55-64	*	*	*	*	3,373	811,400	83.4	(78.8 - 88.1)
≥ 65	88	1,000	83.1	(74.4 - 91.8)	4,215	905,400	90.4	(87.1 - 93.6)
Race/Ethnicity								
White Non-Hispanic	220	2,600	61.9	(54.3 - 69.5)	10,950	1,965,400	67.0	(63.9 - 70.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	4,775	829,200	59.0	(52.4 - 65.6)
Some College	*	*	*	*	2,584	535,000	61.9	(53.7 - 70.2)
≥ College Degree	*	*	*	*	4,176	1,455,700	76.8	(72.5 - 81.1)
Income								
≤ \$24,999	*	*	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	*	3,048	444,100	64.1	(56.0 - 72.3)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,188	1,211,100	72.3	(67.6 - 77.0)

¹ Defined as having ever had a prostate specific antigen test among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have ever had a prostate specific antigen test (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 31b. Ever had a prostate specific antigen test among men aged 40 and older¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	226	59.3	(51.1 - 67.6)	11,557	67.4	(63.7 - 71.1)
Gender						
Male	226	59.3	(51.1 - 67.6)	11,557	67.4	(63.7 - 71.1)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	220	59.1	(50.7 - 67.5)	10,950	66.6	(63.4 - 69.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	4,775	59.7	(52.9 - 66.4)
Some College	*	*	*	2,584	61.4	(53.0 - 69.9)
≥ College Degree	*	*	*	4,176	76.3	(71.9 - 80.7)
Income						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,048	64.2	(56.1 - 72.4)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,188	71.5	(66.6 - 76.3)

¹ Defined as having ever had a prostate specific antigen test among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 32a. Had prostate specific antigen test within the past 2 years among men aged 40 and older¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	225	2,300	50.8	(43.3 - 58.4)	11,437	2,493,800	59.9	(56.0 - 63.9)
Gender								
Male	225	2,300	50.8	(43.3 - 58.4)	11,437	2,493,800	59.9	(56.0 - 63.9)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,917	659,400	49.9	(42.8 - 57.0)
55-64	*	*	*	*	3,326	719,700	74.5	(69.4 - 79.7)
≥ 65	88	1,000	77.5	(68.1 - 86.8)	4,166	830,000	83.3	(78.4 - 88.3)
Race/Ethnicity								
White Non-Hispanic	219	2,100	50.1	(42.5 - 57.6)	10,836	1,762,200	60.4	(57.3 - 63.5)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	4,727	734,200	52.4	(45.7 - 59.1)
Some College	*	*	*	*	2,558	483,800	56.3	(48.2 - 64.4)
≥ College Degree	*	*	*	*	4,133	1,273,200	67.4	(61.6 - 73.3)
Income								
≤ \$24,999	*	*	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	*	3,021	410,800	59.6	(51.5 - 67.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,170	1,080,100	64.6	(59.1 - 70.0)

¹ Defined as having had a prostate specific antigen test within the past 2 years among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have had a prostate specific antigen test within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 32b. Had prostate specific antigen test within the past 2 years among men aged 40 and older¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	225	48.3	(40.3 - 56.3)	11,437	59.8	(55.8 - 63.9)
Gender						
Male	225	48.3	(40.3 - 56.3)	11,437	59.8	(55.8 - 63.9)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	219	47.5	(39.4 - 55.6)	10,836	60.1	(56.9 - 63.4)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	4,727	52.9	(45.9 - 59.9)
Some College	*	*	*	2,558	56.0	(47.7 - 64.2)
≥ College Degree	*	*	*	4,133	67.1	(61.2 - 73.1)
Income						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,021	59.8	(51.7 - 67.9)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,170	63.9	(58.3 - 69.6)

¹ Defined as having had a prostate specific antigen test within the past 2 years among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 33a. Prostate cancer among men aged 40 and older¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	235	200	4.1	(1.7 - 6.5)	12,133	178,200	4.1	(3.0 - 5.2)
Gender								
Male	235	200	4.1	(1.7 - 6.5)	12,133	178,200	4.1	(3.0 - 5.2)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	3,120	6,100	0.4	(0.0 - 0.9)
55-64	*	*	*	*	3,532	31,600	3.1	(1.5 - 4.8)
≥ 65	91	100	11.6	(4.8 - 18.5)	4,388	140,400	13.6	(9.7 - 17.4)
Race/Ethnicity								
White Non-Hispanic	229	200	3.8	(1.5 - 6.1)	11,502	143,700	4.7	(3.5 - 5.8)
Black Non-Hispanic	*	*	*	*	184	23,500	5.1	(0.0 - 11.0)
Hispanic	*	*	*	*	197	7,700	1.6	(0.0 - 3.7)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	5,065	46,300	3.1	(2.2 - 4.1)
Some College	*	*	*	*	2,709	39,200	4.4	(1.3 - 7.5)
≥ College Degree	*	*	*	*	4,336	92,500	4.7	(3.0 - 6.5)
Income								
≤ \$24,999	*	*	*	*	2,597	35,800	4.1	(2.2 - 6.0)
\$25,000-\$49,999	*	*	*	*	3,196	29,600	4.1	(2.9 - 5.4)
\$50,000-\$74,999	*	*	*	*	2,007	20,300	2.8	(1.2 - 4.4)
≥ \$75,000	*	*	*	*	3,322	56,500	3.2	(1.7 - 4.8)

¹ Defined as having ever been told by a health professional that respondent had prostate cancer among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have ever been told they had prostate cancer (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 33b. Prostate cancer among men aged 40 and older¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	235	4.2	(1.8 - 6.6)	12,133	4.5	(3.2 - 5.7)
Gender						
Male	235	4.2	(1.8 - 6.6)	12,133	4.5	(3.2 - 5.7)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	229	3.8	(1.5 - 6.2)	11,502	5.1	(3.8 - 6.4)
Black Non-Hispanic	*	*	*	184	5.7	(0.0 - 12.8)
Hispanic	*	*	*	197	1.8	(0.0 - 4.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	5,065	3.3	(2.3 - 4.3)
Some College	*	*	*	2,709	5.0	(1.1 - 9.0)
≥ College Degree	*	*	*	4,336	5.2	(3.1 - 7.2)
Income						
≤ \$24,999	*	*	*	2,597	4.5	(2.2 - 6.7)
\$25,000-\$49,999	*	*	*	3,196	4.3	(3.0 - 5.6)
\$50,000-\$74,999	*	*	*	2,007	3.0	(1.1 - 5.0)
≥ \$75,000	*	*	*	3,322	3.5	(1.6 - 5.4)

¹ Defined as having ever been told by a health professional that respondent had prostate cancer among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 34a. Home blood stool test ever used among adults aged 50 and older¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	477	3,500	49.2	(44.2 - 54.3)	25,322	2,288,500	38.0	(35.5 - 40.4)
Gender								
Male	193	1,600	47.7	(39.8 - 55.5)	9,571	1,081,900	38.9	(35.1 - 42.6)
Female	284	1,900	50.6	(44.1 - 57.1)	15,751	1,206,600	37.2	(34.0 - 40.4)
Age (years)								
50-64	243	1,800	44.6	(37.6 - 51.6)	12,834	1,192,000	34.0	(30.8 - 37.2)
≥ 65	234	1,600	55.7	(48.8 - 62.6)	12,472	1,093,400	43.6	(39.8 - 47.4)
Race/Ethnicity								
White Non-Hispanic	468	3,300	49.0	(44.0 - 54.1)	24,213	1,754,500	39.1	(37.1 - 41.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	233	1,500	44.5	(37.3 - 51.7)	11,146	823,500	34.9	(30.5 - 39.2)
Some College	129	1,100	55.2	(45.6 - 64.7)	6,261	480,300	37.0	(32.6 - 41.4)
≥ College Degree	*	*	*	*	7,851	973,700	41.6	(37.8 - 45.3)
Income								
≤ \$24,999	149	1,000	50.4	(41.2 - 59.6)	7,222	505,000	35.4	(29.7 - 41.2)
\$25,000-\$49,999	137	900	46.7	(37.5 - 55.8)	6,609	467,600	40.3	(35.2 - 45.4)
\$50,000-\$74,999	*	*	*	*	3,500	313,000	38.2	(32.6 - 43.9)
≥ \$75,000	*	*	*	*	4,647	657,800	37.3	(33.0 - 41.6)

¹ Defined as having ever used a home blood stool test among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have ever used a home blood stool test (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 34b. Home blood stool test ever used among adults aged 50 and older¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	477	48.8	(43.6 - 54.0)	25,322	38.1	(35.5 - 40.8)
Gender						
Male	193	48.1	(40.1 - 56.0)	9,571	39.4	(35.3 - 43.5)
Female	284	49.5	(42.8 - 56.2)	15,751	36.9	(33.5 - 40.3)
Race/Ethnicity						
White Non-Hispanic	468	48.5	(43.4 - 53.7)	24,213	38.9	(36.7 - 41.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	233	43.6	(36.3 - 51.0)	11,146	35.1	(30.3 - 39.9)
Some College	129	54.5	(44.6 - 64.4)	6,261	36.6	(32.0 - 41.3)
≥ College Degree	*	*	*	7,851	41.9	(37.9 - 46.0)
Income						
≤ \$24,999	149	48.5	(39.2 - 57.8)	7,222	35.9	(29.5 - 42.4)
\$25,000-\$49,999	137	46.9	(37.6 - 56.1)	6,609	40.2	(34.8 - 45.6)
\$50,000-\$74,999	*	*	*	3,500	38.1	(31.9 - 44.3)
≥ \$75,000	*	*	*	4,647	37.5	(32.9 - 42.2)

¹ Defined as having ever used a home blood stool test among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 35a. Home blood stool test used within the past year among adults aged 50 and older¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	473	1,300	19.2	(15.2 - 23.2)	25,074	863,300	14.4	(12.6 - 16.2)
Gender								
Male	192	600	19.2	(12.8 - 25.7)	9,508	444,400	16.0	(13.2 - 18.9)
Female	281	700	19.1	(14.2 - 24.1)	15,566	418,900	13.1	(10.8 - 15.4)
Age (years)								
50-64	241	700	17.4	(11.9 - 23.0)	12,759	454,400	13.0	(10.6 - 15.5)
≥ 65	232	600	21.6	(15.9 - 27.2)	12,299	406,400	16.4	(13.7 - 19.0)
Race/Ethnicity								
White Non-Hispanic	464	1,200	18.5	(14.8 - 22.3)	23,971	634,300	14.3	(12.8 - 15.7)
Black Non-Hispanic	*	*	*	*	396	136,000	18.6	(9.4 - 27.8)
Hispanic	*	*	*	*	349	53,400	9.9	(2.5 - 17.2)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	230	700	20.6	(14.3 - 26.8)	11,026	288,600	12.3	(9.4 - 15.3)
Some College	129	400	20.8	(13.4 - 28.3)	6,208	212,500	16.4	(12.5 - 20.4)
≥ College Degree	114	300	14.8	(7.9 - 21.7)	7,778	352,400	15.2	(12.5 - 17.9)
Income								
≤ \$24,999	149	500	25.6	(17.7 - 33.4)	7,157	196,300	13.9	(10.1 - 17.6)
\$25,000-\$49,999	137	400	18.9	(11.7 - 26.0)	6,562	221,500	19.2	(14.1 - 24.3)
\$50,000-\$74,999	*	*	*	*	3,469	111,900	13.7	(10.1 - 17.4)
≥ \$75,000	*	*	*	*	4,623	205,100	11.7	(8.8 - 14.6)

¹ Defined as having used a home blood stool test within the past year among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have used a home blood stool test within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 35b. Home blood stool test used within the past year among adults aged 50 and older¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	473	19.5	(15.3 - 23.6)	25,074	14.6	(12.6 - 16.7)
Gender						
Male	192	20.1	(13.4 - 26.8)	9,508	16.4	(13.2 - 19.6)
Female	281	18.8	(13.8 - 23.9)	15,566	12.9	(10.4 - 15.4)
Race/Ethnicity						
White Non-Hispanic	464	18.7	(14.8 - 22.6)	23,971	14.2	(12.7 - 15.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	349	10.2	(1.7 - 18.7)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	230	20.5	(14.1 - 27.0)	11,026	12.7	(9.2 - 16.1)
Some College	129	21.1	(13.5 - 28.8)	6,208	16.6	(12.3 - 20.9)
≥ College Degree	114	15.7	(8.3 - 23.2)	7,778	15.3	(12.4 - 18.3)
Income						
≤ \$24,999	149	24.6	(16.8 - 32.4)	7,157	14.3	(9.8 - 18.7)
\$25,000-\$49,999	137	19.2	(11.8 - 26.6)	6,562	19.3	(13.8 - 24.8)
\$50,000-\$74,999	*	*	*	3,469	13.8	(9.8 - 17.7)
≥ \$75,000	*	*	*	4,623	11.8	(8.6 - 15.0)

¹ Defined as having used a home blood stool test within the past year among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 36a. Home blood stool test used within the past 2 years among adults aged 50 and older¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	473	2,000	28.8	(24.3 - 33.4)	25,074	1,218,000	20.4	(18.3 - 22.5)
Gender								
Male	192	1,000	29.1	(21.8 - 36.4)	9,508	593,100	21.4	(18.3 - 24.5)
Female	281	1,100	28.6	(22.9 - 34.3)	15,566	624,900	19.5	(16.6 - 22.4)
Age (years)								
50-64	241	1,100	26.1	(19.7 - 32.4)	12,759	629,900	18.1	(15.4 - 20.7)
≥ 65	232	1,000	32.7	(26.3 - 39.2)	12,299	585,500	23.6	(20.2 - 27.0)
Race/Ethnicity								
White Non-Hispanic	464	1,900	28.3	(23.8 - 32.7)	23,971	877,500	19.7	(18.2 - 21.3)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	230	900	27.9	(21.2 - 34.6)	11,026	454,600	19.4	(15.6 - 23.3)
Some College	129	700	33.2	(24.3 - 42.1)	6,208	262,200	20.3	(16.3 - 24.2)
≥ College Degree	114	500	25.8	(16.9 - 34.7)	7,778	491,400	21.2	(18.1 - 24.2)
Income								
≤ \$24,999	149	700	35.3	(26.7 - 43.9)	7,157	263,900	18.6	(14.6 - 22.7)
\$25,000-\$49,999	137	600	29.9	(21.5 - 38.2)	6,562	285,900	24.8	(19.6 - 30.0)
\$50,000-\$74,999	*	*	*	*	3,469	167,500	20.6	(15.6 - 25.5)
≥ \$75,000	*	*	*	*	4,623	294,900	16.8	(13.7 - 20.0)

¹ Defined as having used a home blood stool test within the past 2 years among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have used a home blood stool test within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

**Table 36b. Home blood stool test used within the past 2 years among adults aged 50 and older¹
in Schuyler County and New York State - Age-Adjusted²**

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	473	29.1	(24.4 - 33.8)	25,074	20.5	(18.2 - 22.8)
Gender						
Male	192	30.0	(22.5 - 37.5)	9,508	21.7	(18.3 - 25.1)
Female	281	28.3	(22.5 - 34.1)	15,566	19.4	(16.3 - 22.5)
Race/Ethnicity						
White Non-Hispanic	464	28.4	(23.8 - 33.0)	23,971	19.7	(18.0 - 21.3)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	230	27.6	(20.8 - 34.5)	11,026	19.5	(15.4 - 23.7)
Some College	129	33.3	(24.1 - 42.5)	6,208	20.4	(16.1 - 24.7)
≥ College Degree	114	27.3	(17.9 - 36.6)	7,778	21.4	(18.1 - 24.7)
Income						
≤ \$24,999	149	33.8	(25.2 - 42.3)	7,157	18.9	(14.2 - 23.6)
\$25,000-\$49,999	137	30.5	(21.9 - 39.0)	6,562	24.8	(19.3 - 30.3)
\$50,000-\$74,999	*	*	*	3,469	20.6	(15.3 - 25.9)
≥ \$75,000	*	*	*	4,623	16.9	(13.4 - 20.4)

¹ Defined as having used a home blood stool test within the past 2 years among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 37a. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	476	4,500	63.5	(58.6 - 68.5)	25,357	4,019,100	66.6	(64.1 - 69.1)
Gender								
Male	192	2,100	62.9	(55.0 - 70.8)	9,582	1,894,100	68.0	(64.2 - 71.7)
Female	284	2,400	64.1	(57.8 - 70.3)	15,775	2,125,000	65.5	(62.1 - 68.8)
Age (years)								
50-54	*	*	*	*	4,179	718,900	50.6	(44.6 - 56.5)
55-64	173	1,700	68.7	(61.3 - 76.2)	8,695	1,493,500	71.0	(67.3 - 74.8)
≥ 65	232	1,900	65.0	(58.2 - 71.7)	12,466	1,802,000	72.1	(68.6 - 75.7)
Race/Ethnicity								
White Non-Hispanic	467	4,400	64.9	(60.0 - 69.8)	24,248	3,093,900	68.8	(66.8 - 70.8)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	232	1,900	58.8	(51.6 - 66.1)	11,121	1,445,100	61.2	(56.7 - 65.8)
Some College	130	1,200	61.1	(51.7 - 70.5)	6,287	835,300	64.3	(59.7 - 69.0)
≥ College Degree	114	1,300	74.7	(65.4 - 84.1)	7,884	1,732,400	73.7	(70.0 - 77.4)
Income								
≤ \$24,999	149	1,100	56.8	(47.8 - 65.9)	7,220	892,600	62.7	(56.8 - 68.6)
\$25,000-\$49,999	137	1,300	65.6	(56.9 - 74.2)	6,619	697,300	60.1	(54.6 - 65.7)
\$50,000-\$74,999	*	*	*	*	3,514	613,700	74.2	(68.9 - 79.6)
≥ \$75,000	*	*	*	*	4,665	1,294,400	73.3	(69.2 - 77.5)

¹ Defined as having ever had a sigmoidoscopy or colonoscopy among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have ever had a sigmoidoscopy or colonoscopy (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 37b. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	476	63.2	(58.0 - 68.3)	25,357	66.6	(63.9 - 69.3)
Gender						
Male	192	62.6	(54.6 - 70.6)	9,582	68.5	(64.5 - 72.5)
Female	284	63.7	(57.3 - 70.2)	15,775	64.8	(61.2 - 68.4)
Race/Ethnicity						
White Non-Hispanic	467	64.6	(59.6 - 69.6)	24,248	68.7	(66.5 - 70.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	232	58.5	(51.1 - 66.0)	11,121	61.3	(56.4 - 66.2)
Some College	130	60.5	(50.7 - 70.3)	6,287	63.8	(58.8 - 68.8)
≥ College Degree	114	74.2	(64.6 - 83.9)	7,884	74.0	(70.2 - 77.9)
Income						
≤ \$24,999	149	55.9	(46.6 - 65.3)	7,220	63.0	(56.7 - 69.3)
\$25,000-\$49,999	137	65.6	(56.9 - 74.3)	6,619	59.7	(53.8 - 65.6)
\$50,000-\$74,999	*	*	*	3,514	74.4	(68.3 - 80.5)
≥ \$75,000	*	*	*	4,665	73.6	(69.3 - 78.0)

¹ Defined as having ever had a sigmoidoscopy or colonoscopy among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 38a. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	473	4,300	61.3	(56.2 - 66.3)	25,169	3,855,800	64.6	(62.1 - 67.1)
Gender								
Male	191	2,000	61.3	(53.3 - 69.2)	9,521	1,835,400	66.4	(62.6 - 70.1)
Female	282	2,300	61.2	(54.9 - 67.6)	15,648	2,020,400	63.1	(59.8 - 66.4)
Age (years)								
50-54	*	*	*	*	4,169	700,500	49.3	(43.3 - 55.3)
55-64	173	1,700	67.2	(59.7 - 74.8)	8,652	1,455,100	69.4	(65.7 - 73.2)
≥ 65	229	1,800	62.1	(55.2 - 68.9)	12,331	1,695,700	69.5	(65.9 - 73.1)
Race/Ethnicity								
White Non-Hispanic	464	4,200	62.6	(57.6 - 67.5)	24,069	2,981,500	66.6	(64.6 - 68.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	232	1,900	57.1	(49.8 - 64.3)	11,017	1,360,400	59.0	(54.5 - 63.5)
Some College	128	1,200	59.5	(50.0 - 69.0)	6,251	807,100	62.4	(57.8 - 67.1)
≥ College Degree	113	1,300	70.8	(60.9 - 80.7)	7,841	1,682,500	71.7	(68.0 - 75.4)
Income								
≤ \$24,999	149	1,100	54.5	(45.4 - 63.6)	7,163	858,900	61.0	(55.1 - 67.0)
\$25,000-\$49,999	136	1,200	62.3	(53.3 - 71.2)	6,584	664,900	57.7	(52.2 - 63.2)
\$50,000-\$74,999	*	*	*	*	3,501	602,600	73.1	(67.7 - 78.5)
≥ \$75,000	*	*	*	*	4,655	1,260,400	71.5	(67.3 - 75.7)

¹ Defined as having had a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have had a sigmoidoscopy or colonoscopy within the past 10 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 38b. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	473	60.8	(55.6 - 66.0)	25,169	64.6	(61.9 - 67.3)
Gender						
Male	191	60.8	(52.8 - 68.9)	9,521	66.9	(62.9 - 70.9)
Female	282	60.8	(54.2 - 67.4)	15,648	62.4	(58.9 - 66.0)
Race/Ethnicity						
White Non-Hispanic	464	62.2	(57.1 - 67.3)	24,069	66.5	(64.3 - 68.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	232	56.7	(49.2 - 64.2)	11,017	59.0	(54.1 - 63.9)
Some College	128	59.0	(49.1 - 68.9)	6,251	61.9	(56.9 - 66.9)
≥ College Degree	*	*	*	7,841	72.1	(68.2 - 75.9)
Income						
≤ \$24,999	149	53.5	(44.2 - 62.9)	7,163	61.3	(54.9 - 67.6)
\$25,000-\$49,999	136	61.9	(52.8 - 71.0)	6,584	57.2	(51.4 - 63.0)
\$50,000-\$74,999	*	*	*	3,501	73.3	(67.2 - 79.4)
≥ \$75,000	*	*	*	4,655	71.8	(67.4 - 76.2)

¹ Defined as having had a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 39a. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	469	4,900	70.5	(65.8 - 75.2)	24,792	4,049,200	69.1	(66.7 - 71.6)
Gender								
Male	192	2,400	71.7	(64.4 - 79.0)	9,418	1,945,200	70.7	(67.0 - 74.3)
Female	277	2,600	69.5	(63.4 - 75.5)	15,374	2,104,100	67.8	(64.6 - 71.1)
Age (years)								
50-54	*	*	*	*	4,132	751,200	54.1	(48.0 - 60.1)
55-64	167	1,800	75.6	(68.6 - 82.6)	8,366	1,487,600	73.7	(70.0 - 77.4)
≥ 65	231	2,100	71.3	(64.9 - 77.7)	12,294	1,810,500	74.0	(70.6 - 77.4)
Race/Ethnicity								
White Non-Hispanic	460	4,800	70.8	(66.1 - 75.6)	23,726	3,091,400	70.4	(68.4 - 72.4)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	229	2,100	65.6	(58.7 - 72.5)	10,872	1,461,600	64.2	(59.8 - 68.7)
Some College	126	1,400	72.0	(63.1 - 81.0)	6,170	873,300	68.4	(63.9 - 72.8)
≥ College Degree	114	1,400	77.8	(68.6 - 86.9)	7,708	1,709,000	74.6	(70.9 - 78.3)
Income								
≤ \$24,999	147	1,300	65.7	(56.9 - 74.4)	7,092	923,300	66.0	(60.1 - 71.9)
\$25,000-\$49,999	136	1,400	70.7	(62.5 - 78.9)	6,526	730,000	64.2	(58.9 - 69.6)
\$50,000-\$74,999	*	*	*	*	3,467	621,500	76.6	(71.4 - 81.9)
≥ \$75,000	*	*	*	*	4,615	1,292,500	73.8	(69.8 - 77.9)

¹ Defined as having used a home blood stool test within the past year or a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 years and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have used a home blood stool test in the past year or sigmoidoscopy or colonoscopy within the past 10 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 39b. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	469	70.5	(65.6 - 75.3)	24,792	69.3	(66.7 - 71.9)
Gender						
Male	192	71.9	(64.5 - 79.3)	9,418	71.5	(67.6 - 75.4)
Female	277	69.1	(62.8 - 75.4)	15,374	67.2	(63.7 - 70.7)
Race/Ethnicity						
White Non-Hispanic	460	70.7	(65.8 - 75.6)	23,726	70.3	(68.1 - 72.4)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	229	65.5	(58.4 - 72.6)	10,872	64.5	(59.8 - 69.3)
Some College	126	71.7	(62.2 - 81.1)	6,170	68.2	(63.4 - 72.9)
≥ College Degree	114	78.0	(68.7 - 87.3)	7,708	75.0	(71.2 - 78.8)
Income						
≤ \$24,999	147	64.6	(55.5 - 73.8)	7,092	66.4	(60.2 - 72.6)
\$25,000-\$49,999	136	70.9	(62.6 - 79.2)	6,526	63.8	(58.2 - 69.5)
\$50,000-\$74,999	*	*	*	3,467	76.7	(70.6 - 82.7)
≥ \$75,000	*	*	*	4,615	74.4	(70.2 - 78.6)

¹ Defined as having used a home blood stool test within the past year or a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 years and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 40a. High blood pressure among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	669	4,600	30.9	(26.7 - 35.0)	38,125	3,974,800	26.8	(24.9 - 28.7)
Gender								
Male	279	2,200	29.4	(23.1 - 35.7)	14,611	1,960,500	27.7	(24.7 - 30.7)
Female	390	2,500	32.3	(26.8 - 37.8)	23,514	2,014,300	26.0	(23.6 - 28.3)
Age (years)								
18-34	*	*	*	*	3,935	312,200	7.0	(4.2 - 9.8)
35-44	77	500	18.7	(9.3 - 28.1)	4,984	426,200	14.8	(11.6 - 18.1)
45-54	127	800	25.5	(17.2 - 33.8)	7,825	862,400	30.3	(25.9 - 34.8)
55-64	174	1,200	46.1	(38.0 - 54.2)	8,765	890,800	41.6	(37.6 - 45.7)
≥ 65	234	1,900	63.4	(56.6 - 70.2)	12,616	1,483,200	58.5	(54.7 - 62.3)
Race/Ethnicity								
White Non-Hispanic	656	4,500	31.2	(27.0 - 35.4)	35,814	2,650,900	28.4	(26.9 - 30.0)
Black Non-Hispanic	*	*	*	*	736	558,500	26.2	(19.6 - 32.8)
Hispanic	*	*	*	*	863	538,100	23.8	(17.0 - 30.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	314	2,600	38.3	(31.6 - 45.0)	15,688	1,851,200	33.1	(29.4 - 36.9)
Some College	182	1,100	25.0	(18.2 - 31.7)	9,906	851,100	24.1	(20.9 - 27.3)
≥ College Degree	173	1,000	24.8	(17.1 - 32.5)	12,452	1,253,400	22.3	(19.7 - 24.9)
Income								
≤ \$24,999	193	1,400	36.2	(27.2 - 45.2)	9,640	1,153,000	32.8	(27.6 - 38.1)
\$25,000-\$49,999	196	1,200	27.2	(20.0 - 34.3)	9,657	807,800	27.9	(23.7 - 32.1)
\$50,000-\$74,999	109	800	32.0	(22.7 - 41.4)	5,773	507,600	25.8	(21.4 - 30.1)
≥ \$75,000	*	*	*	*	8,560	942,500	21.0	(18.4 - 23.6)

¹ Defined as having ever been told by a health professional that respondent has high blood pressure

² Sample size from survey

³ Estimated number of adults who have ever been told by a health professional that they have high blood pressure (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 40b. High blood pressure among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	669	27.7	(23.4 - 32.0)	38,125	25.7	(23.8 - 27.5)
Gender						
Male	279	27.8	(21.4 - 34.3)	14,611	27.7	(24.7 - 30.7)
Female	390	27.5	(21.8 - 33.3)	23,514	23.8	(21.5 - 26.0)
Race/Ethnicity						
White Non-Hispanic	656	28.0	(23.6 - 32.3)	35,814	27.2	(25.6 - 28.7)
Black Non-Hispanic	*	*	*	736	24.9	(18.4 - 31.3)
Hispanic	*	*	*	863	23.3	(16.6 - 30.0)
Other Non-Hispanic	*	*	*	712	19.3	(9.5 - 29.1)
Education						
≤ High School	314	35.0	(28.0 - 42.1)	15,688	32.1	(28.4 - 35.8)
Some College	182	21.2	(14.7 - 27.7)	9,906	22.8	(19.7 - 25.9)
≥ College Degree	173	22.8	(14.6 - 31.1)	12,452	21.2	(18.6 - 23.8)
Income						
≤ \$24,999	193	33.1	(23.2 - 42.9)	9,640	31.8	(26.7 - 37.0)
\$25,000-\$49,999	196	24.3	(17.2 - 31.4)	9,657	26.9	(22.8 - 30.9)
\$50,000-\$74,999	109	26.7	(17.8 - 35.5)	5,773	24.8	(20.4 - 29.2)
≥ \$75,000	*	*	*	8,560	19.9	(17.3 - 22.5)

¹ Defined as having ever been told by a health professional that respondent has high blood pressure

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 41a. Blood pressure medication among adults with high blood pressure¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	287	3,700	79.2	(71.7 - 86.7)	15,288	3,145,800	79.2	(76.0 - 82.4)
Gender								
Male	*	*	*	*	6,080	1,432,100	73.1	(67.6 - 78.5)
Female	*	*	*	*	9,208	1,713,700	85.2	(82.1 - 88.2)
Age (years)								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,315	645,500	74.9	(66.8 - 82.9)
55-64	82	1,000	84.8	(77.0 - 92.7)	4,045	782,300	87.8	(84.5 - 91.1)
≥ 65	151	1,700	92.5	(87.7 - 97.3)	7,675	1,383,400	93.4	(91.1 - 95.7)
Race/Ethnicity								
White Non-Hispanic	283	3,500	78.5	(70.9 - 86.2)	14,442	2,117,300	79.9	(77.5 - 82.4)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	158	2,100	81.5	(71.8 - 91.2)	7,354	1,482,800	80.2	(74.7 - 85.6)
Some College	*	*	*	*	3,868	632,000	74.3	(67.8 - 80.7)
≥ College Degree	*	*	*	*	4,035	1,013,200	80.9	(76.8 - 84.9)
Income								
≤ \$24,999	*	*	*	*	4,898	948,200	82.3	(75.7 - 88.9)
\$25,000-\$49,999	*	*	*	*	4,088	668,300	82.8	(77.6 - 88.0)
\$50,000-\$74,999	*	*	*	*	1,971	391,500	77.2	(67.7 - 86.6)
≥ \$75,000	*	*	*	*	2,397	683,700	72.5	(66.1 - 78.9)

¹ Defined as currently taking medicine for high blood pressure among adults with high blood pressure

² Sample size from survey

³ Estimated number of adults with high blood pressure who are taking medicine for high blood pressure (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 41b. Blood pressure medication among adults with high blood pressure¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	287	76.1	(67.1 - 85.2)	15,288	78.5	(75.2 - 81.7)
Gender						
Male	*	*	*	6,080	73.3	(67.9 - 78.7)
Female	*	*	*	9,208	84.0	(80.7 - 87.2)
Race/Ethnicity						
White Non-Hispanic	283	75.5	(66.3 - 84.7)	14,442	79.0	(76.5 - 81.6)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	7,354	79.7	(74.3 - 85.2)
Some College	*	*	*	3,868	72.5	(65.7 - 79.3)
≥ College Degree	*	*	*	4,035	80.4	(76.1 - 84.6)
Income						
≤ \$24,999	*	*	*	4,898	81.6	(75.0 - 88.3)
\$25,000-\$49,999	*	*	*	4,088	82.1	(76.7 - 87.5)
\$50,000-\$74,999	*	*	*	1,971	76.4	(66.7 - 86.1)
≥ \$75,000	*	*	*	2,397	71.6	(64.8 - 78.3)

¹ Defined as currently taking medicine for high blood pressure among adults with high blood pressure

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 42a. Ever had blood cholesterol checked among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	657	10,900	75.4	(69.9 - 80.9)	37,450	11,496,600	80.3	(77.7 - 83.0)
Gender								
Male	275	5,100	71.4	(62.7 - 80.1)	14,389	5,288,400	77.7	(73.3 - 82.0)
Female	382	5,800	79.3	(72.6 - 86.0)	23,061	6,208,200	82.7	(79.6 - 85.8)
Age (years)								
18-34	*	*	*	*	3,684	2,296,400	55.3	(48.7 - 61.9)
35-44	76	1,900	80.7	(71.0 - 90.3)	4,883	2,264,200	81.6	(76.6 - 86.5)
45-54	124	2,600	90.2	(84.0 - 96.3)	7,731	2,539,400	91.0	(87.0 - 94.9)
55-64	174	2,400	95.4	(91.9 - 98.8)	8,718	2,014,900	95.4	(93.5 - 97.2)
≥ 65	231	2,800	97.1	(95.0 - 99.2)	12,434	2,381,700	95.9	(94.3 - 97.6)
Race/Ethnicity								
White Non-Hispanic	645	10,600	76.0	(70.6 - 81.4)	35,198	7,832,600	86.3	(84.8 - 87.9)
Black Non-Hispanic	*	*	*	*	720	1,588,900	77.8	(68.4 - 87.2)
Hispanic	*	*	*	*	842	1,224,600	56.8	(46.8 - 66.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	307	4,800	75.1	(67.2 - 83.1)	15,342	3,795,700	70.6	(65.6 - 75.6)
Some College	*	*	*	*	9,732	2,599,800	77.4	(71.7 - 83.2)
≥ College Degree	170	3,200	80.9	(71.5 - 90.2)	12,296	5,008,300	91.3	(88.9 - 93.6)
Income								
≤ \$24,999	*	*	*	*	9,437	2,277,500	68.4	(61.5 - 75.3)
\$25,000-\$49,999	*	*	*	*	9,492	2,053,000	71.9	(64.5 - 79.3)
\$50,000-\$74,999	*	*	*	*	5,721	1,736,100	90.0	(87.7 - 92.3)
≥ \$75,000	*	*	*	*	8,460	4,019,300	92.2	(90.8 - 93.6)

¹ Defined as ever having blood cholesterol checked

² Sample size from survey

³ Estimated number of adults who have ever had blood cholesterol checked (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 42b. Ever had blood cholesterol checked among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	657	72.4	(66.3 - 78.4)	37,450	79.9	(77.3 - 82.5)
Gender						
Male	275	69.3	(60.1 - 78.5)	14,389	78.0	(73.9 - 82.1)
Female	382	75.3	(67.5 - 83.1)	23,061	81.6	(78.4 - 84.8)
Race/Ethnicity						
White Non-Hispanic	645	72.9	(67.0 - 78.9)	35,198	85.6	(84.0 - 87.2)
Black Non-Hispanic	*	*	*	720	77.5	(68.2 - 86.7)
Hispanic	*	*	*	842	57.2	(47.4 - 67.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	307	72.1	(63.3 - 80.9)	15,342	70.2	(65.2 - 75.1)
Some College	*	*	*	9,732	76.7	(71.1 - 82.3)
≥ College Degree	*	*	*	12,296	90.9	(88.6 - 93.2)
Income						
≤ \$24,999	*	*	*	9,437	68.1	(61.3 - 74.9)
\$25,000-\$49,999	*	*	*	9,492	71.6	(64.4 - 78.7)
\$50,000-\$74,999	*	*	*	5,721	89.6	(87.2 - 92.0)
≥ \$75,000	*	*	*	8,460	91.7	(90.1 - 93.2)

¹ Defined as ever having blood cholesterol checked

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 43a. Blood cholesterol checked within the past 5 years among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	653	10,400	71.8	(66.3 - 77.3)	37,201	11,035,200	77.8	(75.1 - 80.4)
Gender								
Male	272	4,800	68.6	(59.9 - 77.2)	14,307	5,020,700	74.5	(70.2 - 78.9)
Female	381	5,500	74.9	(68.1 - 81.7)	22,894	6,014,600	80.7	(77.6 - 83.8)
Age (years)								
18-34	*	*	*	*	3,647	2,179,800	53.3	(46.7 - 59.9)
35-44	*	*	*	*	4,862	2,161,200	78.0	(72.8 - 83.2)
45-54	123	2,500	89.2	(82.9 - 95.6)	7,697	2,408,600	86.6	(82.4 - 90.8)
55-64	173	2,300	90.0	(85.4 - 94.6)	8,677	1,957,900	93.3	(91.4 - 95.3)
≥ 65	229	2,800	95.3	(92.8 - 97.8)	12,318	2,327,800	94.8	(93.1 - 96.6)
Race/Ethnicity								
White Non-Hispanic	641	10,000	72.3	(66.8 - 77.7)	34,960	7,486,600	83.4	(81.8 - 85.0)
Black Non-Hispanic	*	*	*	*	719	1,561,700	76.5	(67.1 - 85.9)
Hispanic	*	*	*	*	838	1,184,000	55.6	(45.6 - 65.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	303	4,600	72.0	(64.0 - 80.0)	15,215	3,612,500	68.3	(63.2 - 73.3)
Some College	*	*	*	*	9,673	2,524,200	75.4	(69.8 - 81.1)
≥ College Degree	170	2,900	74.4	(64.7 - 84.1)	12,236	4,806,500	88.1	(85.5 - 90.6)
Income								
≤ \$24,999	*	*	*	*	9,350	2,214,800	66.9	(60.1 - 73.8)
\$25,000-\$49,999	*	*	*	*	9,454	1,936,600	68.8	(61.5 - 76.1)
\$50,000-\$74,999	*	*	*	*	5,701	1,640,400	85.9	(82.4 - 89.5)
≥ \$75,000	*	*	*	*	8,435	3,872,400	89.2	(87.5 - 90.9)

¹ Defined as having had blood cholesterol checked within the past 5 years

² Sample size from survey

³ Estimated number of adults who have had blood cholesterol checked within the past 5 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 43b. Blood cholesterol checked within the past 5 years among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	653	68.6	(62.5 - 74.6)	37,201	77.3	(74.7 - 79.9)
Gender						
Male	272	66.6	(57.5 - 75.7)	14,307	74.8	(70.7 - 79.0)
Female	381	70.5	(62.6 - 78.3)	22,894	79.5	(76.3 - 82.7)
Race/Ethnicity						
White Non-Hispanic	641	69.0	(63.0 - 75.0)	34,960	82.6	(81.0 - 84.3)
Black Non-Hispanic	*	*	*	719	76.2	(66.9 - 85.5)
Hispanic	*	*	*	838	55.9	(46.1 - 65.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	303	68.8	(60.0 - 77.6)	15,215	67.8	(62.9 - 72.8)
Some College	*	*	*	9,673	74.7	(69.1 - 80.2)
≥ College Degree	*	*	*	12,236	87.5	(85.0 - 90.1)
Income						
≤ \$24,999	*	*	*	9,350	66.6	(59.8 - 73.4)
\$25,000-\$49,999	*	*	*	9,454	68.5	(61.4 - 75.6)
\$50,000-\$74,999	*	*	*	5,701	85.4	(81.7 - 89.0)
≥ \$75,000	*	*	*	8,435	88.5	(86.7 - 90.3)

¹ Defined as having had blood cholesterol checked within the past 5 years

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 44a. Cardiovascular disease among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	661	1,200	8.4	(6.4 - 10.5)	37,810	1,135,000	7.7	(6.5 - 8.8)
Gender								
Male	274	600	8.0	(5.0 - 11.0)	14,477	664,600	9.4	(7.5 - 11.4)
Female	387	700	8.8	(6.0 - 11.6)	23,333	470,400	6.1	(4.8 - 7.4)
Age (years)								
18-34	*	*	*	*	3,930	100,800	2.3	(0.0 - 4.8)
35-44	*	*	*	*	4,971	103,400	3.6	(1.5 - 5.7)
45-54	125	300	9.1	(3.7 - 14.4)	7,768	131,300	4.6	(2.9 - 6.4)
55-64	172	200	6.2	(2.4 - 9.9)	8,699	224,200	10.6	(8.3 - 12.9)
≥ 65	231	700	25.2	(19.0 - 31.3)	12,442	575,300	23.0	(19.9 - 26.1)
Race/Ethnicity								
White Non-Hispanic	648	1,200	8.6	(6.5 - 10.7)	35,522	772,200	8.3	(7.4 - 9.2)
Black Non-Hispanic	*	*	*	*	728	140,700	6.6	(3.3 - 9.9)
Hispanic	*	*	*	*	854	163,300	7.3	(2.0 - 12.5)
Other Non-Hispanic	*	*	*	*	706	58,800	5.2	(0.9 - 9.5)
Education								
≤ High School	307	700	10.6	(7.3 - 14.0)	15,486	590,800	10.7	(8.2 - 13.2)
Some College	182	300	6.5	(3.0 - 10.0)	9,849	227,100	6.4	(4.7 - 8.1)
≥ College Degree	172	300	6.9	(3.3 - 10.4)	12,393	314,600	5.6	(4.3 - 6.9)
Income								
≤ \$24,999	191	500	12.9	(8.1 - 17.7)	9,505	448,100	12.9	(9.1 - 16.7)
\$25,000-\$49,999	192	300	6.8	(3.2 - 10.4)	9,571	179,300	6.2	(4.8 - 7.6)
\$50,000-\$74,999	109	200	8.2	(3.3 - 13.0)	5,751	166,900	8.5	(5.5 - 11.4)
≥ \$75,000	*	*	*	*	8,538	183,500	4.1	(2.9 - 5.3)

¹ Defined as having ever been told by a health professional that respondent had a heart attack or angina or a stroke

² Sample size from survey

³ Estimated number of adults with cardiovascular disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 44b. Cardiovascular disease among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	661	7.5	(5.6 - 9.5)	37,810	7.6	(6.4 - 8.7)
Gender						
Male	274	7.9	(4.8 - 11.0)	14,477	9.7	(7.8 - 11.7)
Female	387	7.2	(4.7 - 9.7)	23,333	5.6	(4.3 - 6.9)
Race/Ethnicity						
White Non-Hispanic	648	7.7	(5.7 - 9.8)	35,522	8.2	(7.2 - 9.1)
Black Non-Hispanic	*	*	*	728	6.8	(3.2 - 10.4)
Hispanic	*	*	*	854	7.0	(2.0 - 12.0)
Other Non-Hispanic	*	*	*	706	5.5	(0.9 - 10.0)
Education						
≤ High School	307	9.8	(6.4 - 13.2)	15,486	10.5	(8.0 - 13.0)
Some College	182	5.6	(2.4 - 8.7)	9,849	6.4	(4.5 - 8.2)
≥ College Degree	172	6.1	(2.7 - 9.6)	12,393	5.5	(4.2 - 6.9)
Income						
≤ \$24,999	191	11.3	(6.8 - 15.8)	9,505	12.8	(9.0 - 16.6)
\$25,000-\$49,999	192	7.0	(3.0 - 11.0)	9,571	6.0	(4.7 - 7.4)
\$50,000-\$74,999	109	6.9	(2.6 - 11.3)	5,751	8.5	(5.4 - 11.6)
≥ \$75,000	*	*	*	8,538	4.0	(2.7 - 5.2)

¹ Defined as having ever been told by a health professional that respondent had a heart attack or angina or a stroke

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 45a. Coronary heart disease among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	662	1,100	7.4	(5.5 - 9.4)	37,838	929,400	6.3	(5.3 - 7.3)
Gender								
Male	275	500	6.8	(4.0 - 9.6)	14,479	598,600	8.5	(6.6 - 10.4)
Female	387	600	8.0	(5.3 - 10.7)	23,359	330,800	4.3	(3.4 - 5.2)
Age (years)								
18-34	*	*	*	*	3,932	63,400	1.4	(0.0 - 3.6)
35-44	*	*	*	*	4,973	95,300	3.3	(1.2 - 5.4)
45-54	126	300	9.0	(3.7 - 14.3)	7,778	108,300	3.8	(2.1 - 5.6)
55-64	172	200	6.2	(2.4 - 9.9)	8,703	188,600	8.9	(6.7 - 11.1)
≥ 65	231	600	20.1	(14.4 - 25.8)	12,452	473,800	18.9	(16.1 - 21.7)
Race/Ethnicity								
White Non-Hispanic	649	1,100	7.6	(5.6 - 9.6)	35,552	636,700	6.9	(6.0 - 7.7)
Black Non-Hispanic	*	*	*	*	728	113,100	5.3	(2.3 - 8.4)
Hispanic	*	*	*	*	853	128,000	5.7	(1.1 - 10.3)
Other Non-Hispanic	*	*	*	*	705	51,500	4.6	(0.5 - 8.7)
Education								
≤ High School	307	600	8.7	(5.6 - 11.8)	15,485	474,500	8.6	(6.3 - 10.8)
Some College	182	300	6.3	(2.8 - 9.8)	9,860	193,000	5.5	(3.9 - 7.1)
≥ College Degree	173	300	6.5	(3.0 - 9.9)	12,411	259,700	4.6	(3.4 - 5.8)
Income								
≤ \$24,999	192	400	10.4	(6.0 - 14.7)	9,507	372,300	10.7	(7.3 - 14.1)
\$25,000-\$49,999	192	300	6.4	(2.9 - 9.9)	9,585	136,300	4.7	(3.8 - 5.7)
\$50,000-\$74,999	109	200	7.5	(2.9 - 12.2)	5,755	154,600	7.8	(4.9 - 10.7)
≥ \$75,000	*	*	*	*	8,541	151,800	3.4	(2.3 - 4.5)

¹ Defined as having ever been told by a health professional that respondent had a heart attack or angina

² Sample size from survey

³ Estimated number of adults with coronary heart disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 45b. Coronary heart disease among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	662	6.6	(4.8 - 8.5)	37,838	6.3	(5.2 - 7.3)
Gender						
Male	275	6.7	(3.9 - 9.6)	14,479	8.8	(6.9 - 10.8)
Female	387	6.6	(4.1 - 9.0)	23,359	3.9	(3.0 - 4.8)
Race/Ethnicity						
White Non-Hispanic	649	6.8	(4.9 - 8.7)	35,552	6.8	(5.9 - 7.7)
Black Non-Hispanic	*	*	*	728	5.6	(2.2 - 9.0)
Hispanic	*	*	*	853	5.4	(1.1 - 9.6)
Other Non-Hispanic	*	*	*	705	4.8	(0.4 - 9.2)
Education						
≤ High School	307	8.0	(4.9 - 11.1)	15,485	8.5	(6.2 - 10.7)
Some College	182	5.4	(2.3 - 8.5)	9,860	5.5	(3.7 - 7.2)
≥ College Degree	173	5.8	(2.4 - 9.1)	12,411	4.7	(3.4 - 5.9)
Income						
≤ \$24,999	192	9.1	(5.1 - 13.0)	9,507	10.7	(7.2 - 14.1)
\$25,000-\$49,999	192	6.6	(2.7 - 10.6)	9,585	4.7	(3.6 - 5.7)
\$50,000-\$74,999	109	6.4	(2.2 - 10.5)	5,755	8.0	(4.8 - 11.1)
≥ \$75,000	*	*	*	8,541	3.3	(2.1 - 4.5)

¹ Defined as having ever been told by a health professional that respondent had a heart attack or angina

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 46a. Ever diagnosed with diabetes among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	670	1,700	11.5	(8.7 - 14.2)	38,174	1,382,700	9.3	(8.0 - 10.5)
Gender								
Male	280	800	10.8	(6.8 - 14.7)	14,626	530,400	7.5	(6.2 - 8.8)
Female	390	900	12.1	(8.4 - 15.9)	23,548	852,300	11.0	(8.9 - 13.0)
Age (years)								
18-34	*	*	*	*	3,941	150,200	3.4	(1.0 - 5.8)
35-44	*	*	*	*	4,987	165,100	5.7	(3.6 - 7.9)
45-54	127	200	6.8	(2.3 - 11.3)	7,830	279,000	9.8	(6.7 - 12.8)
55-64	174	500	18.7	(12.3 - 25.1)	8,771	332,900	15.5	(12.3 - 18.8)
≥ 65	235	700	22.7	(16.9 - 28.6)	12,645	455,600	17.9	(15.0 - 20.8)
Race/Ethnicity								
White Non-Hispanic	657	1,700	11.8	(9.0 - 14.6)	35,862	809,300	8.7	(7.8 - 9.5)
Black Non-Hispanic	*	*	*	*	739	261,000	12.1	(6.6 - 17.6)
Hispanic	*	*	*	*	862	227,300	10.1	(5.4 - 14.8)
Other Non-Hispanic	*	*	*	*	711	85,100	7.6	(2.3 - 12.8)
Education								
≤ High School	314	900	13.8	(9.5 - 18.1)	15,709	730,000	13.0	(10.5 - 15.6)
Some College	182	400	9.1	(4.0 - 14.1)	9,918	303,500	8.6	(5.9 - 11.2)
≥ College Degree	174	400	10.2	(5.4 - 15.0)	12,464	337,500	6.0	(4.8 - 7.2)
Income								
≤ \$24,999	193	600	14.9	(9.1 - 20.7)	9,652	543,400	15.5	(11.4 - 19.5)
\$25,000-\$49,999	196	600	12.3	(6.9 - 17.7)	9,667	291,000	10.0	(7.5 - 12.6)
\$50,000-\$74,999	109	200	6.7	(2.6 - 10.7)	5,777	151,000	7.6	(5.0 - 10.3)
≥ \$75,000	102	200	9.1	(3.6 - 14.6)	8,565	220,800	4.9	(3.8 - 6.1)

¹ Defined as having ever been told by a doctor that respondent had diabetes, excluding pre-diabetes and women with diabetes only when pregnant

² Sample size from survey

³ Estimated number of adults who have diabetes (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 46b. Ever diagnosed with diabetes among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	670	10.3	(7.5 - 13.1)	38,174	9.0	(7.8 - 10.3)
Gender						
Male	280	10.0	(6.0 - 14.0)	14,626	7.5	(6.1 - 8.9)
Female	390	10.6	(6.7 - 14.6)	23,548	10.4	(8.4 - 12.4)
Race/Ethnicity						
White Non-Hispanic	657	10.6	(7.7 - 13.5)	35,862	8.4	(7.5 - 9.2)
Black Non-Hispanic	*	*	*	739	11.4	(6.0 - 16.7)
Hispanic	*	*	*	862	10.1	(5.4 - 14.9)
Other Non-Hispanic	*	*	*	711	7.7	(2.3 - 13.0)
Education						
≤ High School	314	12.2	(7.9 - 16.6)	15,709	12.6	(10.0 - 15.2)
Some College	182	8.5	(2.9 - 14.1)	9,918	8.3	(5.6 - 10.9)
≥ College Degree	174	9.2	(4.3 - 14.1)	12,464	5.9	(4.7 - 7.2)
Income						
≤ \$24,999	193	14.1	(7.8 - 20.5)	9,652	15.2	(11.1 - 19.2)
\$25,000-\$49,999	196	10.8	(5.4 - 16.3)	9,667	9.7	(7.1 - 12.2)
\$50,000-\$74,999	109	5.4	(2.0 - 8.8)	5,777	7.6	(4.9 - 10.2)
≥ \$75,000	102	8.0	(2.9 - 13.1)	8,565	4.7	(3.6 - 5.9)

¹ Defined as having ever been told by a doctor that respondent had diabetes, excluding pre-diabetes and women with diabetes only when pregnant

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 47a. Ever diagnosed with asthma (lifetime) among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	662	2,500	16.5	(12.9 - 20.1)	38,096	2,225,100	15.0	(13.2 - 16.8)
Gender								
Male	278	1,000	13.6	(8.8 - 18.5)	14,601	875,800	12.4	(10.1 - 14.7)
Female	384	1,500	19.2	(14.0 - 24.4)	23,495	1,349,300	17.4	(14.6 - 20.1)
Age (years)								
18-34	57	600	14.3	(5.5 - 23.1)	3,931	825,100	18.5	(13.8 - 23.3)
35-44	*	*	*	*	4,972	431,700	15.1	(11.2 - 19.1)
45-54	126	500	15.9	(8.7 - 23.1)	7,819	365,900	12.8	(9.7 - 16.0)
55-64	173	400	15.9	(9.9 - 21.9)	8,756	316,700	14.8	(11.9 - 17.7)
≥ 65	230	400	13.8	(8.8 - 18.8)	12,618	285,700	11.2	(8.6 - 13.8)
Race/Ethnicity								
White Non-Hispanic	649	2,400	16.6	(12.9 - 20.3)	35,783	1,365,700	14.6	(13.0 - 16.3)
Black Non-Hispanic	*	*	*	*	737	367,000	17.2	(10.2 - 24.2)
Hispanic	*	*	*	*	863	317,100	14.1	(9.0 - 19.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	311	1,100	17.2	(11.7 - 22.6)	15,675	831,600	14.9	(12.0 - 17.8)
Some College	178	600	14.8	(8.5 - 21.0)	9,893	587,500	16.6	(13.1 - 20.1)
≥ College Degree	173	700	17.1	(9.8 - 24.4)	12,444	739,000	13.2	(10.4 - 15.9)
Income								
≤ \$24,999	190	600	17.2	(10.1 - 24.3)	9,643	708,900	20.2	(14.9 - 25.5)
\$25,000-\$49,999	195	900	19.4	(12.2 - 26.6)	9,641	389,800	13.4	(10.0 - 16.9)
\$50,000-\$74,999	108	400	16.1	(6.7 - 25.6)	5,762	257,800	13.2	(10.0 - 16.5)
≥ \$75,000	102	300	13.4	(6.2 - 20.7)	8,562	577,900	12.8	(10.4 - 15.3)

¹ Defined as adults having ever been told by a health professional that they had asthma

² Sample size from survey

³ Estimated number of adults who have lifetime asthma (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 47b. Ever diagnosed with asthma (lifetime) among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	662	17.0	(13.0 - 21.0)	38,096	15.2	(13.3 - 17.0)
Gender						
Male	278	13.8	(8.6 - 19.1)	14,601	12.6	(10.3 - 14.9)
Female	384	20.1	(14.1 - 26.1)	23,495	17.5	(14.8 - 20.3)
Race/Ethnicity						
White Non-Hispanic	649	17.2	(13.0 - 21.3)	35,783	14.9	(13.2 - 16.6)
Black Non-Hispanic	*	*	*	737	17.1	(10.1 - 24.1)
Hispanic	*	*	*	863	14.5	(9.3 - 19.7)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	311	18.2	(12.0 - 24.3)	15,675	15.1	(12.1 - 18.0)
Some College	178	14.6	(7.8 - 21.5)	9,893	16.9	(13.4 - 20.4)
≥ College Degree	173	17.6	(9.6 - 25.6)	12,444	13.3	(10.6 - 16.0)
Income						
≤ \$24,999	190	17.7	(9.5 - 25.9)	9,643	20.3	(15.0 - 25.6)
\$25,000-\$49,999	195	20.0	(12.2 - 27.9)	9,641	13.7	(10.2 - 17.2)
\$50,000-\$74,999	*	*	*	5,762	13.5	(10.0 - 17.0)
≥ \$75,000	102	14.1	(6.0 - 22.3)	8,562	13.0	(10.6 - 15.4)

¹ Defined as adults having ever been told by a health professional that they had asthma

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 48a. Current asthma among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	661	1,600	10.9	(8.1 - 13.7)	37,933	1,415,000	9.6	(8.1 - 11.0)
Gender								
Male	278	500	7.1	(3.8 - 10.4)	14,546	440,700	6.3	(4.8 - 7.7)
Female	383	1,100	14.6	(10.2 - 19.0)	23,387	974,300	12.6	(10.2 - 15.0)
Age (years)								
18-34	*	*	*	*	3,909	467,200	10.5	(6.8 - 14.3)
35-44	76	500	20.4	(10.7 - 30.2)	4,947	246,800	8.7	(5.8 - 11.6)
45-54	126	300	10.9	(5.1 - 16.7)	7,787	240,000	8.4	(6.1 - 10.8)
55-64	172	300	11.3	(6.1 - 16.5)	8,726	231,100	10.8	(8.1 - 13.5)
≥ 65	230	300	10.9	(6.3 - 15.5)	12,564	229,900	9.1	(6.5 - 11.6)
Race/Ethnicity								
White Non-Hispanic	648	1,500	10.8	(7.9 - 13.6)	35,629	888,400	9.6	(8.2 - 11.0)
Black Non-Hispanic	*	*	*	*	736	246,900	11.6	(5.7 - 17.5)
Hispanic	*	*	*	*	859	186,800	8.3	(4.4 - 12.2)
Other Non-Hispanic	*	*	*	*	709	93,000	8.3	(1.4 - 15.2)
Education								
≤ High School	311	800	11.5	(7.4 - 15.7)	15,594	595,100	10.7	(8.2 - 13.2)
Some College	177	500	12.1	(6.2 - 18.1)	9,861	322,000	9.1	(7.0 - 11.3)
≥ College Degree	173	300	8.6	(3.9 - 13.3)	12,394	430,900	7.7	(5.8 - 9.7)
Income								
≤ \$24,999	190	500	14.1	(7.5 - 20.6)	9,600	489,200	14.0	(9.7 - 18.3)
\$25,000-\$49,999	195	500	10.9	(5.8 - 16.0)	9,590	268,400	9.3	(6.2 - 12.4)
\$50,000-\$74,999	*	*	*	*	5,741	143,400	7.4	(5.1 - 9.6)
≥ \$75,000	102	300	10.4	(3.7 - 17.1)	8,541	339,200	7.6	(5.8 - 9.4)

¹ Defined as adults having ever been told by a health professional that they had asthma and still have asthma

² Sample size from survey

³ Estimated number of adults who currently have asthma (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 48b. Current asthma among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	661	11.2	(8.0 - 14.3)	37,933	9.7	(8.2 - 11.2)
Gender						
Male	278	7.1	(3.5 - 10.8)	14,546	6.4	(4.9 - 7.9)
Female	383	15.1	(10.0 - 20.2)	23,387	12.7	(10.2 - 15.1)
Race/Ethnicity						
White Non-Hispanic	648	11.1	(7.9 - 14.3)	35,629	9.7	(8.3 - 11.2)
Black Non-Hispanic	*	*	*	736	11.5	(5.6 - 17.3)
Hispanic	*	*	*	859	8.6	(4.5 - 12.6)
Other Non-Hispanic	*	*	*	709	8.3	(1.3 - 15.3)
Education						
≤ High School	311	12.0	(7.2 - 16.9)	15,594	10.8	(8.3 - 13.3)
Some College	177	12.6	(5.9 - 19.3)	9,861	9.4	(7.2 - 11.6)
≥ College Degree	173	8.4	(3.4 - 13.4)	12,394	7.8	(5.8 - 9.7)
Income						
≤ \$24,999	190	14.5	(7.0 - 22.0)	9,600	14.1	(9.7 - 18.4)
\$25,000-\$49,999	195	11.1	(5.4 - 16.9)	9,590	9.5	(6.3 - 12.7)
\$50,000-\$74,999	*	*	*	5,741	7.5	(5.0 - 10.1)
≥ \$75,000	102	11.5	(3.7 - 19.2)	8,541	7.7	(5.9 - 9.5)

¹ Defined as adults having ever been told by a health professional that they had asthma and still have asthma

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 49a. Chronic joint symptoms among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	668	7,400	49.7	(44.6 - 54.8)	38,099	6,377,100	42.9	(40.5 - 45.3)
Gender								
Male	278	3,600	49.0	(41.2 - 56.8)	14,595	3,033,600	42.8	(39.0 - 46.6)
Female	390	3,800	50.4	(43.9 - 56.9)	23,504	3,343,500	43.0	(40.0 - 46.0)
Age (years)								
18-34	*	*	*	*	3,933	1,261,300	28.3	(22.6 - 34.0)
35-44	*	*	*	*	4,984	1,110,900	38.6	(33.1 - 44.0)
45-54	127	1,600	54.4	(44.9 - 63.9)	7,817	1,456,500	51.2	(46.4 - 55.9)
55-64	173	1,300	52.6	(44.4 - 60.8)	8,751	1,167,400	54.6	(50.4 - 58.8)
≥ 65	234	1,800	61.1	(54.3 - 67.8)	12,614	1,381,000	54.3	(50.5 - 58.1)
Race/Ethnicity								
White Non-Hispanic	655	6,900	47.9	(42.9 - 53.0)	35,785	4,267,600	45.8	(43.8 - 47.7)
Black Non-Hispanic	*	*	*	*	740	890,500	41.3	(32.5 - 50.0)
Hispanic	*	*	*	*	860	827,500	36.7	(27.7 - 45.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	314	3,700	55.8	(48.4 - 63.2)	15,676	2,450,000	43.8	(39.4 - 48.1)
Some College	*	*	*	*	9,900	1,598,300	45.1	(40.1 - 50.1)
≥ College Degree	173	1,800	44.2	(34.9 - 53.6)	12,440	2,313,100	41.1	(37.8 - 44.5)
Income								
≤ \$24,999	*	*	*	*	9,641	1,483,000	42.2	(36.3 - 48.1)
\$25,000-\$49,999	196	2,400	52.2	(42.5 - 61.8)	9,647	1,332,100	45.9	(39.8 - 52.1)
\$50,000-\$74,999	*	*	*	*	5,768	894,200	45.3	(39.5 - 51.1)
≥ \$75,000	*	*	*	*	8,549	1,874,200	41.7	(38.4 - 45.1)

¹ Defined as adults having symptoms of pain, aching or stiffness in or around a joint during the past 30 days and began more than 3 months ago

² Sample size from survey

³ Estimated number of adults who have chronic joint symptoms (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 49b. Chronic joint symptoms among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	668	48.4	(42.8 - 53.9)	38,099	42.1	(39.7 - 44.5)
Gender						
Male	278	48.5	(40.2 - 56.8)	14,595	42.6	(38.8 - 46.3)
Female	390	48.3	(40.9 - 55.6)	23,504	41.6	(38.6 - 44.7)
Race/Ethnicity						
White Non-Hispanic	655	46.6	(41.2 - 52.1)	35,785	44.7	(42.7 - 46.6)
Black Non-Hispanic	*	*	*	740	40.2	(31.5 - 48.9)
Hispanic	*	*	*	860	36.6	(27.8 - 45.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	314	54.7	(46.5 - 62.8)	15,676	43.1	(38.8 - 47.5)
Some College	*	*	*	9,900	43.9	(39.0 - 48.8)
≥ College Degree	*	*	*	12,440	40.3	(37.0 - 43.7)
Income						
≤ \$24,999	*	*	*	9,641	41.6	(35.7 - 47.4)
\$25,000-\$49,999	*	*	*	9,647	45.0	(39.0 - 51.0)
\$50,000-\$74,999	*	*	*	5,768	44.3	(38.4 - 50.2)
≥ \$75,000	*	*	*	8,549	40.8	(37.4 - 44.2)

¹ Defined as adults having symptoms of pain, aching or stiffness in or around a joint during the past 30 days and began more than 3 months ago

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 50a. Arthritis among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	667	4,800	31.8	(27.6 - 36.1)	38,013	4,102,800	27.7	(25.9 - 29.6)
Gender								
Male	278	1,900	26.0	(20.1 - 32.0)	14,555	1,642,700	23.3	(20.7 - 25.8)
Female	389	2,800	37.5	(31.7 - 43.4)	23,458	2,460,100	31.8	(29.2 - 34.4)
Age (years)								
18-34	*	*	*	*	3,927	312,100	7.0	(4.9 - 9.1)
35-44	*	*	*	*	4,972	412,500	14.5	(11.3 - 17.6)
45-54	126	900	30.2	(21.7 - 38.6)	7,789	869,100	30.7	(26.6 - 34.8)
55-64	173	1,000	38.6	(30.7 - 46.4)	8,729	999,300	47.0	(42.9 - 51.1)
≥ 65	235	1,800	59.9	(53.1 - 66.6)	12,596	1,509,800	59.6	(55.8 - 63.3)
Race/Ethnicity								
White Non-Hispanic	654	4,600	32.2	(27.9 - 36.4)	35,710	2,967,700	31.9	(30.3 - 33.6)
Black Non-Hispanic	*	*	*	*	737	588,100	27.3	(20.2 - 34.4)
Hispanic	*	*	*	*	856	380,700	17.1	(11.8 - 22.5)
Other Non-Hispanic	*	*	*	*	710	166,300	14.8	(7.8 - 21.8)
Education								
≤ High School	314	2,500	38.2	(31.6 - 44.8)	15,652	1,742,700	31.3	(27.8 - 34.8)
Some College	179	1,300	30.7	(22.3 - 39.2)	9,874	902,000	25.5	(22.3 - 28.6)
≥ College Degree	174	900	22.5	(16.0 - 29.1)	12,407	1,437,800	25.6	(22.9 - 28.4)
Income								
≤ \$24,999	192	1,500	39.4	(30.2 - 48.6)	9,618	1,112,500	32.0	(27.1 - 36.9)
\$25,000-\$49,999	195	1,400	31.7	(23.9 - 39.4)	9,624	821,900	28.4	(24.2 - 32.6)
\$50,000-\$74,999	109	700	27.7	(18.7 - 36.7)	5,753	542,500	27.5	(23.2 - 31.9)
≥ \$75,000	102	500	20.9	(12.9 - 29.0)	8,536	1,012,800	22.6	(19.8 - 25.3)

¹ Defined as adults having ever been told by a health professional that they have some form of arthritis, rheumatoid arthritis, gout, lupus or fibromyalgia (excluding back or neck)

² Sample size from survey

³ Estimated number of adults who have ever been told by a health professional they have some form of arthritis (excluding back or neck) (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 50b. Arthritis among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	667	29.6	(25.1 - 34.1)	38,013	26.6	(24.8 - 28.5)
Gender						
Male	278	25.6	(19.3 - 32.0)	14,555	23.5	(20.9 - 26.1)
Female	389	33.5	(27.3 - 39.7)	23,458	29.5	(27.0 - 32.1)
Race/Ethnicity						
White Non-Hispanic	654	30.0	(25.5 - 34.5)	35,710	30.4	(28.7 - 32.0)
Black Non-Hispanic	*	*	*	737	26.5	(19.3 - 33.6)
Hispanic	*	*	*	856	17.2	(11.8 - 22.6)
Other Non-Hispanic	*	*	*	710	14.9	(7.7 - 22.0)
Education						
≤ High School	314	35.6	(28.5 - 42.6)	15,652	30.4	(26.9 - 33.8)
Some College	179	29.4	(20.3 - 38.6)	9,874	24.1	(21.0 - 27.1)
≥ College Degree	174	20.5	(13.9 - 27.1)	12,407	24.7	(21.9 - 27.5)
Income						
≤ \$24,999	192	36.0	(26.2 - 45.9)	9,618	31.2	(26.3 - 36.1)
\$25,000-\$49,999	195	29.8	(21.7 - 37.9)	9,624	27.3	(23.2 - 31.3)
\$50,000-\$74,999	109	23.9	(15.2 - 32.7)	5,753	26.3	(21.9 - 30.7)
≥ \$75,000	102	20.5	(11.9 - 29.0)	8,536	21.6	(18.8 - 24.3)

¹ Defined as adults having ever been told by a health professional that they have some form of arthritis, rheumatoid arthritis, gout, lupus or fibromyalgia (excluding back or neck)

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 51a. Activity limitations because of physical, mental or emotional problems among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	666	3,300	22.0	(18.4 - 25.5)	37,998	3,053,000	20.6	(18.7 - 22.5)
Gender								
Male	280	1,500	20.3	(15.2 - 25.3)	14,566	1,318,400	18.6	(15.7 - 21.5)
Female	386	1,800	23.6	(18.6 - 28.6)	23,432	1,734,600	22.4	(19.8 - 25.0)
Age (years)								
18-34	*	*	*	*	3,929	580,000	13.0	(8.2 - 17.9)
35-44	*	*	*	*	4,973	418,600	14.5	(11.0 - 18.1)
45-54	126	600	20.3	(12.8 - 27.8)	7,812	620,200	21.7	(18.3 - 25.2)
55-64	174	700	29.6	(22.2 - 37.0)	8,739	651,600	30.5	(26.5 - 34.6)
≥ 65	232	1,100	37.6	(30.8 - 44.5)	12,545	782,700	31.1	(27.7 - 34.6)
Race/Ethnicity								
White Non-Hispanic	653	3,200	22.4	(18.7 - 26.0)	35,688	2,037,200	21.9	(20.3 - 23.5)
Black Non-Hispanic	*	*	*	*	735	429,700	20.1	(13.5 - 26.7)
Hispanic	*	*	*	*	863	448,200	19.8	(11.9 - 27.8)
Other Non-Hispanic	*	*	*	*	712	137,900	12.3	(5.2 - 19.3)
Education								
≤ High School	313	1,600	23.9	(18.4 - 29.4)	15,626	1,359,600	24.4	(20.6 - 28.1)
Some College	181	700	16.7	(11.1 - 22.4)	9,875	700,800	19.8	(16.5 - 23.2)
≥ College Degree	172	1,000	24.3	(17.0 - 31.6)	12,415	924,800	16.5	(14.2 - 18.7)
Income								
≤ \$24,999	193	1,200	32.7	(24.3 - 41.1)	9,603	1,055,800	30.1	(24.7 - 35.5)
\$25,000-\$49,999	195	800	17.5	(11.6 - 23.5)	9,628	630,900	21.8	(16.6 - 27.0)
\$50,000-\$74,999	108	400	14.0	(7.7 - 20.3)	5,760	408,100	20.7	(16.1 - 25.2)
≥ \$75,000	102	400	15.7	(8.5 - 22.9)	8,544	583,300	13.0	(11.1 - 14.9)

¹ Defined as activity limitations because of physical, mental or emotional problems

² Sample size from survey

³ Estimated number of adults who are limited in any activities because of physical, mental or emotional problems (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 51b. Activity limitations because of physical, mental or emotional problems among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	666	20.7	(16.9 - 24.5)	37,998	20.1	(18.2 - 22.0)
Gender						
Male	280	19.3	(14.0 - 24.6)	14,566	18.6	(15.8 - 21.4)
Female	386	22.1	(16.7 - 27.5)	23,432	21.5	(18.9 - 24.1)
Race/Ethnicity						
White Non-Hispanic	653	21.1	(17.2 - 25.0)	35,688	21.3	(19.7 - 22.9)
Black Non-Hispanic	*	*	*	735	19.9	(13.2 - 26.5)
Hispanic	*	*	*	863	19.1	(11.5 - 26.8)
Other Non-Hispanic	*	*	*	712	12.4	(5.3 - 19.6)
Education						
≤ High School	313	22.6	(16.7 - 28.6)	15,626	23.8	(20.1 - 27.4)
Some College	181	15.5	(9.7 - 21.3)	9,875	19.3	(16.0 - 22.6)
≥ College Degree	172	23.0	(15.2 - 30.7)	12,415	16.1	(13.9 - 18.3)
Income						
≤ \$24,999	193	31.8	(22.6 - 41.0)	9,603	29.7	(24.3 - 35.2)
\$25,000-\$49,999	195	16.7	(10.2 - 23.1)	9,628	21.2	(16.1 - 26.3)
\$50,000-\$74,999	108	12.1	(6.0 - 18.1)	5,760	20.1	(15.6 - 24.6)
≥ \$75,000	102	13.8	(7.0 - 20.7)	8,544	12.7	(10.8 - 14.6)

¹ Defined as activity limitations because of physical, mental or emotional problems

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 52a. Have health problem that requires the use of special equipment among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	669	1,300	8.9	(6.6 - 11.2)	38,171	1,169,400	7.9	(6.9 - 8.9)
Gender								
Male	280	500	7.3	(4.4 - 10.2)	14,617	460,700	6.5	(5.3 - 7.7)
Female	389	800	10.5	(7.0 - 14.0)	23,554	708,700	9.1	(7.6 - 10.7)
Age (years)								
18-34	*	*	*	*	3,940	68,800	1.5	(0.7 - 2.4)
35-44	76	300	12.6	(4.8 - 20.3)	4,989	122,500	4.3	(2.4 - 6.1)
45-54	*	*	*	*	7,836	182,500	6.4	(4.7 - 8.1)
55-64	174	200	8.2	(4.0 - 12.4)	8,773	219,600	10.3	(7.8 - 12.8)
≥ 65	235	600	21.3	(15.6 - 27.0)	12,633	576,100	22.7	(19.0 - 26.3)
Race/Ethnicity								
White Non-Hispanic	656	1,200	8.6	(6.4 - 10.8)	35,853	802,000	8.6	(7.6 - 9.6)
Black Non-Hispanic	*	*	*	*	739	235,800	10.9	(6.4 - 15.5)
Hispanic	*	*	*	*	864	104,100	4.6	(2.1 - 7.1)
Other Non-Hispanic	*	*	*	*	715	27,600	2.5	(0.8 - 4.1)
Education								
≤ High School	314	900	13.6	(9.2 - 18.0)	15,711	604,200	10.8	(8.7 - 12.8)
Some College	181	200	4.2	(1.7 - 6.7)	9,914	260,000	7.3	(5.5 - 9.1)
≥ College Degree	174	200	6.0	(2.7 - 9.4)	12,463	298,900	5.3	(4.1 - 6.5)
Income								
≤ \$24,999	192	500	14.1	(8.6 - 19.7)	9,656	430,500	12.2	(9.6 - 14.8)
\$25,000-\$49,999	196	400	8.7	(4.0 - 13.3)	9,663	238,900	8.2	(5.9 - 10.5)
\$50,000-\$74,999	*	*	*	*	5,775	103,000	5.2	(3.6 - 6.8)
≥ \$75,000	*	*	*	*	8,566	136,000	3.0	(2.1 - 4.0)

¹ Defined as having any health problem that requires the use of special equipment such as a cane, a wheelchair, a special bed, or a special telephone

² Sample size from survey

³ Estimated number of adults who have any health problem that requires the use of special equipment (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 52b. Have health problem that requires the use of special equipment among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	669	8.6	(6.1 - 11.1)	38,171	7.6	(6.6 - 8.6)
Gender						
Male	280	7.5	(4.2 - 10.7)	14,617	6.8	(5.5 - 8.0)
Female	389	9.6	(5.9 - 13.4)	23,554	8.4	(6.9 - 9.9)
Race/Ethnicity						
White Non-Hispanic	656	8.2	(5.9 - 10.5)	35,853	8.3	(7.3 - 9.3)
Black Non-Hispanic	*	*	*	739	10.5	(6.1 - 15.0)
Hispanic	*	*	*	864	4.5	(2.1 - 6.9)
Other Non-Hispanic	*	*	*	715	2.2	(0.7 - 3.8)
Education						
≤ High School	314	13.4	(8.5 - 18.3)	15,711	10.3	(8.3 - 12.3)
Some College	181	3.8	(1.3 - 6.2)	9,914	7.1	(5.2 - 8.9)
≥ College Degree	174	5.9	(2.4 - 9.3)	12,463	5.3	(4.1 - 6.6)
Income						
≤ \$24,999	192	13.3	(7.4 - 19.2)	9,656	11.9	(9.4 - 14.5)
\$25,000-\$49,999	196	8.9	(3.6 - 14.2)	9,663	7.7	(5.5 - 10.0)
\$50,000-\$74,999	*	*	*	5,775	5.1	(3.4 - 6.8)
≥ \$75,000	*	*	*	8,566	2.9	(2.0 - 3.9)

¹ Defined as having any health problem that requires the use of special equipment such as a cane, a wheelchair, a special bed, or a special telephone

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 53a. Disability among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	667	3,600	23.9	(20.1 - 27.6)	38,016	3,394,200	22.9	(20.9 - 24.9)
Gender								
Male	280	1,600	21.6	(16.4 - 26.9)	14,568	1,469,700	20.7	(17.8 - 23.7)
Female	387	2,000	26.1	(20.8 - 31.4)	23,448	1,924,600	24.8	(22.1 - 27.5)
Age (years)								
18-34	*	*	*	*	3,931	590,300	13.3	(8.4 - 18.1)
35-44	*	*	*	*	4,974	447,900	15.6	(11.9 - 19.3)
45-54	126	600	20.3	(12.8 - 27.8)	7,811	643,900	22.6	(19.1 - 26.0)
55-64	174	700	29.6	(22.2 - 37.0)	8,740	702,100	33.0	(28.9 - 37.0)
≥ 65	233	1,300	42.5	(35.6 - 49.4)	12,560	1,010,000	39.9	(36.1 - 43.7)
Race/Ethnicity								
White Non-Hispanic	654	3,400	23.9	(20.1 - 27.6)	35,703	2,253,400	24.2	(22.6 - 25.9)
Black Non-Hispanic	*	*	*	*	736	511,700	23.8	(16.8 - 30.8)
Hispanic	*	*	*	*	864	484,300	21.4	(13.4 - 29.5)
Other Non-Hispanic	*	*	*	*	713	144,800	12.9	(5.8 - 19.9)
Education								
≤ High School	313	1,800	27.3	(21.4 - 33.3)	15,638	1,544,200	27.6	(23.7 - 31.4)
Some College	182	700	17.3	(11.6 - 23.0)	9,876	755,200	21.4	(17.9 - 24.8)
≥ College Degree	172	1,000	25.2	(17.8 - 32.6)	12,419	1,021,300	18.2	(15.9 - 20.5)
Income								
≤ \$24,999	193	1,300	33.6	(25.1 - 42.0)	9,613	1,179,100	33.6	(28.0 - 39.2)
\$25,000-\$49,999	195	1,000	21.1	(14.2 - 27.9)	9,629	699,000	24.2	(18.9 - 29.4)
\$50,000-\$74,999	108	400	15.2	(8.7 - 21.8)	5,757	430,100	21.8	(17.2 - 26.4)
≥ \$75,000	102	400	16.1	(8.9 - 23.4)	8,546	632,700	14.1	(12.1 - 16.1)

¹ Defined as having activity limitations due to physical, mental or emotional problems or have health problems that require the use of special equipment

² Sample size from survey

³ Estimated number of adults with a disability (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 53b. Disability among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	667	22.7	(18.7 - 26.8)	38,016	22.3	(20.4 - 24.3)
Gender						
Male	280	20.8	(15.3 - 26.3)	14,568	20.9	(18.0 - 23.8)
Female	387	24.6	(18.8 - 30.5)	23,448	23.6	(21.0 - 26.3)
Race/Ethnicity						
White Non-Hispanic	654	22.6	(18.7 - 26.6)	35,703	23.6	(21.9 - 25.3)
Black Non-Hispanic	*	*	*	736	23.5	(16.5 - 30.5)
Hispanic	*	*	*	864	20.6	(12.9 - 28.3)
Other Non-Hispanic	*	*	*	713	12.9	(5.8 - 20.1)
Education						
≤ High School	313	26.5	(19.9 - 33.0)	15,638	26.8	(23.0 - 30.6)
Some College	182	16.0	(10.1 - 21.8)	9,876	20.9	(17.4 - 24.3)
≥ College Degree	172	23.8	(16.0 - 31.6)	12,419	17.9	(15.5 - 20.3)
Income						
≤ \$24,999	193	32.6	(23.3 - 41.8)	9,613	33.2	(27.6 - 38.8)
\$25,000-\$49,999	195	21.0	(13.4 - 28.6)	9,629	23.5	(18.3 - 28.6)
\$50,000-\$74,999	108	13.1	(6.8 - 19.4)	5,757	21.2	(16.6 - 25.8)
≥ \$75,000	102	14.3	(7.4 - 21.2)	8,546	13.8	(11.7 - 15.8)

¹ Defined as having activity limitations due to physical, mental or emotional problems or have health problems that require the use of special equipment

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 54a. Current smoking among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	667	3,400	22.6	(18.3 - 27.0)	38,029	2,483,500	16.8	(15.1 - 18.6)
Gender								
Male	279	1,800	24.5	(17.8 - 31.2)	14,559	1,356,400	19.3	(16.3 - 22.2)
Female	388	1,600	20.8	(15.2 - 26.4)	23,470	1,127,100	14.6	(12.7 - 16.6)
Age (years)								
18-34	*	*	*	*	3,929	993,500	22.3	(17.5 - 27.1)
35-44	77	500	19.2	(9.9 - 28.6)	4,977	421,100	14.8	(11.6 - 17.9)
45-54	126	700	22.4	(14.4 - 30.4)	7,807	508,800	18.3	(15.4 - 21.3)
55-64	173	600	23.6	(16.8 - 30.5)	8,744	314,500	14.7	(12.1 - 17.3)
≥ 65	234	300	9.2	(4.9 - 13.4)	12,572	245,600	9.7	(7.2 - 12.3)
Race/Ethnicity								
White Non-Hispanic	654	3,200	22.5	(18.1 - 27.0)	35,729	1,615,500	17.3	(15.9 - 18.8)
Black Non-Hispanic	*	*	*	*	735	284,100	13.6	(8.9 - 18.2)
Hispanic	*	*	*	*	858	350,900	15.8	(9.6 - 21.9)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	313	2,300	34.8	(27.5 - 42.1)	15,654	1,237,300	22.4	(19.1 - 25.6)
Some College	180	500	12.3	(6.6 - 18.1)	9,875	670,600	19.1	(15.5 - 22.8)
≥ College Degree	174	500	13.5	(6.2 - 20.9)	12,421	563,300	10.0	(7.5 - 12.5)
Income								
≤ \$24,999	192	1,200	33.1	(23.5 - 42.7)	9,626	835,100	24.1	(19.1 - 29.1)
\$25,000-\$49,999	195	1,200	27.7	(19.1 - 36.2)	9,629	558,100	19.4	(15.6 - 23.2)
\$50,000-\$74,999	*	*	*	*	5,756	338,100	17.3	(13.0 - 21.7)
≥ \$75,000	102	300	11.0	(3.3 - 18.7)	8,549	468,700	10.4	(8.6 - 12.3)

¹ Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day or some days

² Sample size from survey

³ Estimated number of adult current smokers (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 54b. Current smoking among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	667	23.2	(18.4 - 28.0)	38,029	17.0	(15.3 - 18.8)
Gender						
Male	279	24.6	(17.6 - 31.7)	14,559	19.2	(16.3 - 22.1)
Female	388	21.9	(15.4 - 28.3)	23,470	15.0	(13.0 - 17.0)
Race/Ethnicity						
White Non-Hispanic	654	23.3	(18.4 - 28.1)	35,729	17.7	(16.2 - 19.2)
Black Non-Hispanic	*	*	*	735	13.5	(8.8 - 18.2)
Hispanic	*	*	*	858	15.9	(9.8 - 22.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	313	36.3	(28.3 - 44.3)	15,654	22.8	(19.5 - 26.0)
Some College	180	11.8	(5.7 - 17.9)	9,875	19.4	(15.8 - 22.9)
≥ College Degree	174	14.3	(6.2 - 22.4)	12,421	10.0	(7.6 - 12.5)
Income						
≤ \$24,999	*	*	*	9,626	24.1	(19.2 - 29.1)
\$25,000-\$49,999	195	28.2	(19.0 - 37.3)	9,629	19.9	(16.0 - 23.7)
\$50,000-\$74,999	*	*	*	5,756	17.8	(13.3 - 22.3)
≥ \$75,000	102	10.3	(2.3 - 18.3)	8,549	10.6	(8.6 - 12.5)

¹ Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day or some days

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 55a. Everyday smoking among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	667	2,700	17.8	(13.8 - 21.7)	38,029	1,626,800	11.0	(9.8 - 12.2)
Gender								
Male	279	1,400	19.4	(13.3 - 25.5)	14,559	848,400	12.0	(10.0 - 14.1)
Female	388	1,200	16.2	(11.1 - 21.2)	23,470	778,400	10.1	(8.7 - 11.5)
Age (years)								
18-34	*	*	*	*	3,929	559,800	12.6	(9.6 - 15.6)
35-44	77	400	16.3	(7.6 - 25.1)	4,977	315,200	11.0	(8.4 - 13.7)
45-54	126	600	18.9	(11.4 - 26.5)	7,807	377,000	13.6	(11.1 - 16.0)
55-64	173	500	18.0	(11.7 - 24.4)	8,744	203,000	9.5	(7.5 - 11.5)
≥ 65	234	200	6.8	(3.1 - 10.6)	12,572	171,700	6.8	(4.8 - 8.8)
Race/Ethnicity								
White Non-Hispanic	654	2,500	17.6	(13.6 - 21.6)	35,729	1,158,100	12.4	(11.2 - 13.7)
Black Non-Hispanic	*	*	*	*	735	197,700	9.4	(5.6 - 13.3)
Hispanic	*	*	*	*	858	176,200	7.9	(4.2 - 11.6)
Other Non-Hispanic	*	*	*	*	707	94,900	8.5	(3.3 - 13.7)
Education								
≤ High School	313	2,000	30.2	(23.1 - 37.3)	15,654	881,900	15.9	(13.4 - 18.5)
Some College	180	400	9.3	(4.1 - 14.5)	9,875	410,200	11.7	(9.6 - 13.8)
≥ College Degree	174	200	6.2	(1.9 - 10.4)	12,421	323,500	5.8	(4.3 - 7.2)
Income								
≤ \$24,999	192	1,000	28.0	(18.6 - 37.4)	9,626	452,800	13.1	(10.3 - 15.8)
\$25,000-\$49,999	195	1,200	26.4	(18.0 - 34.9)	9,629	425,900	14.8	(11.9 - 17.7)
\$50,000-\$74,999	*	*	*	*	5,756	222,700	11.4	(8.5 - 14.4)
≥ \$75,000	*	*	*	*	8,549	320,200	7.1	(5.5 - 8.8)

¹ Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day

² Sample size from survey

³ Estimated number of adults who have smoked at least 100 cigarettes in lifetime and currently smoking everyday (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 55b. Everyday smoking among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	667	18.2	(13.9 - 22.5)	38,029	11.3	(10.0 - 12.5)
Gender						
Male	279	19.4	(13.0 - 25.8)	14,559	12.1	(10.1 - 14.1)
Female	388	17.0	(11.2 - 22.8)	23,470	10.5	(9.0 - 12.0)
Race/Ethnicity						
White Non-Hispanic	654	18.2	(13.8 - 22.6)	35,729	12.8	(11.5 - 14.1)
Black Non-Hispanic	*	*	*	735	9.4	(5.5 - 13.4)
Hispanic	*	*	*	858	8.0	(4.3 - 11.7)
Other Non-Hispanic	*	*	*	707	8.5	(3.2 - 13.8)
Education						
≤ High School	313	31.8	(23.9 - 39.6)	15,654	16.4	(13.8 - 19.0)
Some College	180	9.1	(3.5 - 14.6)	9,875	12.1	(9.9 - 14.3)
≥ College Degree	174	6.3	(1.7 - 10.9)	12,421	5.8	(4.3 - 7.3)
Income						
≤ \$24,999	*	*	*	9,626	13.4	(10.6 - 16.2)
\$25,000-\$49,999	195	27.2	(18.1 - 36.3)	9,629	15.3	(12.3 - 18.3)
\$50,000-\$74,999	*	*	*	5,756	11.7	(8.7 - 14.8)
≥ \$75,000	*	*	*	8,549	7.2	(5.5 - 8.9)

¹ Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 56a. Adults living in homes in which smoking is prohibited¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	663	10,900	73.0	(68.5 - 77.4)	38,045	11,997,100	80.8	(79.0 - 82.6)
Gender								
Male	275	5,500	74.9	(68.4 - 81.4)	14,577	5,634,400	79.4	(76.6 - 82.2)
Female	388	5,400	71.0	(65.0 - 77.1)	23,468	6,362,800	82.1	(79.9 - 84.3)
Age (years)								
18-34	*	*	*	*	3,934	3,655,200	82.1	(77.8 - 86.4)
35-44	*	*	*	*	4,985	2,479,700	86.1	(82.5 - 89.7)
45-54	126	2,000	70.0	(61.4 - 78.5)	7,808	2,194,400	77.2	(73.5 - 80.8)
55-64	173	1,800	69.6	(62.1 - 77.2)	8,753	1,732,500	81.0	(78.3 - 83.7)
≥ 65	230	2,200	75.4	(69.5 - 81.3)	12,565	1,935,400	76.5	(73.4 - 79.5)
Race/Ethnicity								
White Non-Hispanic	650	10,500	73.0	(68.5 - 77.5)	35,737	7,426,900	79.8	(78.3 - 81.2)
Black Non-Hispanic	*	*	*	*	738	1,692,100	78.6	(71.9 - 85.3)
Hispanic	*	*	*	*	863	2,003,900	88.7	(84.1 - 93.3)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	312	4,200	63.4	(56.2 - 70.5)	15,647	4,307,200	77.1	(74.0 - 80.2)
Some College	179	3,200	75.4	(66.8 - 84.0)	9,883	2,795,400	78.9	(75.0 - 82.8)
≥ College Degree	172	3,500	86.2	(80.3 - 92.0)	12,437	4,817,300	85.7	(83.3 - 88.2)
Income								
≤ \$24,999	191	2,200	59.6	(49.8 - 69.4)	9,608	2,627,700	74.9	(70.0 - 79.7)
\$25,000-\$49,999	193	3,000	67.2	(58.1 - 76.2)	9,640	2,314,500	79.8	(76.0 - 83.5)
\$50,000-\$74,999	109	2,300	90.4	(84.7 - 96.2)	5,766	1,551,300	78.7	(74.0 - 83.4)
≥ \$75,000	102	2,300	88.4	(82.4 - 94.4)	8,556	3,883,200	86.5	(84.3 - 88.7)

¹ Defined as having a rule that prohibits smoking anywhere inside the home

² Sample size from survey

³ Estimated number of adults who live in a home with a rule that prohibits smoking anywhere inside the home (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 56b. Adults living in homes in which smoking is prohibited¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	663	73.0	(68.1 - 77.9)	38,045	80.9	(79.2 - 82.7)
Gender						
Male	275	75.6	(68.8 - 82.4)	14,577	79.4	(76.7 - 82.2)
Female	388	70.4	(63.4 - 77.4)	23,468	82.3	(80.0 - 84.5)
Race/Ethnicity						
White Non-Hispanic	650	73.1	(68.2 - 78.0)	35,737	79.9	(78.4 - 81.4)
Black Non-Hispanic	*	*	*	738	78.6	(71.8 - 85.4)
Hispanic	*	*	*	863	88.9	(84.4 - 93.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	312	62.7	(54.7 - 70.6)	15,647	76.9	(73.8 - 80.0)
Some College	179	74.9	(65.3 - 84.5)	9,883	79.1	(75.2 - 83.0)
≥ College Degree	172	87.1	(81.2 - 93.0)	12,437	86.0	(83.5 - 88.5)
Income						
≤ \$24,999	*	*	*	9,608	74.9	(70.1 - 79.7)
\$25,000-\$49,999	193	66.6	(56.8 - 76.5)	9,640	79.8	(76.1 - 83.6)
\$50,000-\$74,999	109	89.8	(82.8 - 96.7)	5,766	78.6	(73.8 - 83.3)
≥ \$75,000	102	90.2	(84.8 - 95.5)	8,556	86.6	(84.3 - 88.9)

¹ Defined as having a rule that prohibits smoking anywhere inside the home

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 57a. Binge drinking within the past month among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	662	3,100	20.7	(16.1 - 25.3)	37,745	2,602,000	17.7	(15.7 - 19.8)
Gender								
Male	274	1,900	25.6	(18.5 - 32.7)	14,420	1,712,400	24.5	(20.9 - 28.1)
Female	388	1,200	16.0	(10.0 - 22.0)	23,325	889,600	11.6	(9.5 - 13.7)
Age (years)								
18-34	*	*	*	*	3,895	1,254,300	28.5	(22.8 - 34.2)
35-44	*	*	*	*	4,934	584,800	20.6	(16.5 - 24.7)
45-54	124	600	20.1	(11.7 - 28.6)	7,738	467,600	16.6	(13.3 - 19.9)
55-64	172	300	11.4	(6.4 - 16.4)	8,663	207,200	9.8	(7.9 - 11.8)
≥ 65	233	100	4.4	(1.6 - 7.2)	12,515	88,100	3.5	(2.5 - 4.5)
Race/Ethnicity								
White Non-Hispanic	649	2,900	20.1	(15.5 - 24.7)	35,453	1,761,900	19.2	(17.5 - 20.8)
Black Non-Hispanic	*	*	*	*	728	247,000	11.6	(6.3 - 17.0)
Hispanic	*	*	*	*	854	460,600	20.5	(11.3 - 29.8)
Other Non-Hispanic	*	*	*	*	710	132,500	12.1	(3.4 - 20.7)
Education								
≤ High School	313	1,400	21.6	(14.8 - 28.4)	15,491	945,800	17.4	(13.2 - 21.6)
Some College	178	500	11.2	(5.2 - 17.2)	9,819	636,600	18.1	(14.6 - 21.5)
≥ College Degree	*	*	*	*	12,358	1,018,700	18.2	(15.3 - 21.1)
Income								
≤ \$24,999	189	400	10.7	(3.5 - 18.0)	9,557	469,100	13.5	(8.5 - 18.5)
\$25,000-\$49,999	195	1,000	21.1	(12.6 - 29.5)	9,575	534,300	18.5	(13.0 - 24.0)
\$50,000-\$74,999	*	*	*	*	5,726	378,200	19.2	(14.1 - 24.3)
≥ \$75,000	*	*	*	*	8,501	1,011,400	22.6	(19.7 - 25.6)

¹ Defined as men having 5 or more drinks or women having 4 or more drinks on 1 or more occasion within the past month

² Sample size from survey

³ Estimated number of adult binge drinkers within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 57b. Binge drinking within the past month among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	662	22.5	(17.3 - 27.7)	37,745	18.1	(16.1 - 20.2)
Gender						
Male	274	26.8	(19.2 - 34.5)	14,420	24.5	(21.0 - 27.9)
Female	388	18.4	(11.3 - 25.4)	23,325	12.4	(10.2 - 14.5)
Race/Ethnicity						
White Non-Hispanic	649	21.9	(16.7 - 27.2)	35,453	19.8	(18.1 - 21.6)
Black Non-Hispanic	*	*	*	728	11.8	(6.4 - 17.1)
Hispanic	*	*	*	854	20.2	(11.3 - 29.1)
Other Non-Hispanic	*	*	*	710	12.1	(3.4 - 20.8)
Education						
≤ High School	313	23.6	(16.0 - 31.2)	15,491	17.5	(13.5 - 21.5)
Some College	178	11.4	(4.6 - 18.2)	9,819	18.6	(15.1 - 22.0)
≥ College Degree	*	*	*	12,358	18.8	(15.9 - 21.7)
Income						
≤ \$24,999	189	11.8	(3.2 - 20.5)	9,557	13.4	(8.6 - 18.2)
\$25,000-\$49,999	195	23.2	(13.8 - 32.6)	9,575	18.9	(13.5 - 24.2)
\$50,000-\$74,999	*	*	*	5,726	19.9	(14.8 - 25.1)
≥ \$75,000	*	*	*	8,501	23.2	(20.2 - 26.1)

¹ Defined as men having 5 or more drinks or women having 4 or more drinks on 1 or more occasion within the past month

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 58a. Heavy drinking within the past month among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	660	1,300	8.5	(5.3 - 11.6)	37,553	718,700	4.9	(4.2 - 5.6)
Gender								
Male	273	700	9.0	(4.5 - 13.5)	14,330	333,300	4.8	(3.9 - 5.7)
Female	387	600	8.0	(3.5 - 12.4)	23,223	385,400	5.0	(4.0 - 6.1)
Age (years)								
18-34	*	*	*	*	3,866	224,900	5.1	(3.5 - 6.8)
35-44	*	*	*	*	4,924	137,200	4.8	(3.0 - 6.7)
45-54	125	200	8.1	(2.7 - 13.4)	7,715	144,600	5.2	(4.1 - 6.2)
55-64	*	*	*	*	8,624	117,400	5.6	(3.9 - 7.3)
≥ 65	*	*	*	*	12,424	94,600	3.8	(2.7 - 4.9)
Race/Ethnicity								
White Non-Hispanic	647	1,100	7.9	(4.8 - 10.9)	35,279	572,400	6.3	(5.4 - 7.1)
Black Non-Hispanic	*	*	*	*	723	58,100	2.8	(0.6 - 4.9)
Hispanic	*	*	*	*	848	60,900	2.7	(0.5 - 4.9)
Other Non-Hispanic	*	*	*	*	703	27,200	2.5	(0.8 - 4.2)
Education								
≤ High School	312	700	10.8	(5.5 - 16.1)	15,411	228,100	4.2	(3.2 - 5.2)
Some College	*	*	*	*	9,759	163,700	4.7	(3.6 - 5.8)
≥ College Degree	171	300	7.7	(2.0 - 13.4)	12,306	326,800	5.9	(4.5 - 7.3)
Income								
≤ \$24,999	*	*	*	*	9,512	104,200	3.0	(1.8 - 4.1)
\$25,000-\$49,999	193	400	9.1	(3.0 - 15.2)	9,537	158,700	5.5	(3.9 - 7.2)
\$50,000-\$74,999	*	*	*	*	5,708	95,400	4.9	(3.1 - 6.7)
≥ \$75,000	*	*	*	*	8,481	313,900	7.0	(5.4 - 8.6)

¹ Defined as adult men averaging more than 2 alcoholic drinks per day and adult women averaging more than 1 alcoholic drink per day within the past month

² Sample size from survey

³ Estimated number of adult heavy drinkers within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 58b. Heavy drinking within the past month among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	660	9.0	(5.5 - 12.5)	37,553	5.0	(4.3 - 5.8)
Gender						
Male	273	9.0	(4.3 - 13.7)	14,330	4.9	(3.9 - 5.9)
Female	387	9.0	(3.8 - 14.3)	23,223	5.1	(4.0 - 6.2)
Race/Ethnicity						
White Non-Hispanic	647	8.3	(4.9 - 11.8)	35,279	6.4	(5.5 - 7.2)
Black Non-Hispanic	*	*	*	723	2.9	(0.6 - 5.1)
Hispanic	*	*	*	848	2.8	(0.5 - 5.1)
Other Non-Hispanic	*	*	*	703	2.5	(0.8 - 4.2)
Education						
≤ High School	312	11.6	(5.7 - 17.5)	15,411	4.3	(3.2 - 5.3)
Some College	*	*	*	9,759	4.8	(3.6 - 5.9)
≥ College Degree	171	8.4	(1.9 - 14.9)	12,306	6.0	(4.5 - 7.4)
Income						
≤ \$24,999	*	*	*	9,512	3.0	(1.9 - 4.1)
\$25,000-\$49,999	193	9.7	(3.0 - 16.3)	9,537	5.8	(4.0 - 7.5)
\$50,000-\$74,999	*	*	*	5,708	5.0	(3.1 - 7.0)
≥ \$75,000	*	*	*	8,481	7.0	(5.4 - 8.7)

¹ Defined as adult men averaging more than 2 alcoholic drinks per day and adult women averaging more than 1 alcoholic drink per day within the past month

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 59a. Never or rarely been asked about sexual history during routine check-up among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	607	8,500	62.7	(57.3 - 68.2)	35,193	8,480,600	61.9	(59.2 - 64.5)
Gender								
Male	246	4,400	67.5	(58.9 - 76.1)	13,299	4,357,400	67.7	(63.6 - 71.7)
Female	361	4,100	58.3	(51.5 - 65.2)	21,894	4,123,300	56.8	(53.3 - 60.2)
Age (years)								
18-34	*	*	*	*	3,507	1,804,600	44.3	(37.8 - 50.8)
35-44	*	*	*	*	4,531	1,635,300	62.2	(56.6 - 67.8)
45-54	114	1,700	63.4	(53.8 - 73.1)	7,215	1,715,600	65.4	(60.5 - 70.3)
55-64	156	1,800	80.0	(73.2 - 86.7)	8,168	1,416,200	70.7	(66.6 - 74.8)
≥ 65	216	2,100	75.7	(69.1 - 82.2)	11,772	1,908,900	80.3	(76.9 - 83.8)
Race/Ethnicity								
White Non-Hispanic	597	8,300	63.6	(58.2 - 68.9)	33,090	5,940,300	68.5	(66.3 - 70.7)
Black Non-Hispanic	*	*	*	*	683	831,800	40.8	(31.8 - 49.9)
Hispanic	*	*	*	*	784	1,071,700	53.4	(43.6 - 63.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	286	3,900	64.6	(56.7 - 72.4)	14,264	3,001,200	60.1	(55.4 - 64.9)
Some College	*	*	*	*	9,156	1,972,800	60.3	(54.9 - 65.7)
≥ College Degree	162	2,400	64.4	(54.6 - 74.1)	11,706	3,476,000	65.0	(61.2 - 68.7)
Income								
≤ \$24,999	177	2,300	67.5	(57.5 - 77.5)	8,772	1,529,400	49.3	(42.6 - 56.0)
\$25,000-\$49,999	*	*	*	*	8,932	1,705,000	64.4	(58.0 - 70.7)
\$50,000-\$74,999	*	*	*	*	5,433	1,217,000	64.5	(58.5 - 70.5)
≥ \$75,000	*	*	*	*	8,108	2,934,300	68.2	(64.7 - 71.7)

¹ Defined as respondents who never or rarely (hardly ever) were asked about their sexual history, such as about sexual partners and sexual practices, during a routine check-up at a doctor's office or clinic

² Sample size from survey

³ Estimated number of adults who were never or rarely asked about their sexual history during a routine check-up at a doctor's office or clinic (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 59b. Never or rarely been asked about sexual history during routine check-up among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	607	60.9	(54.9 - 66.8)	35,193	61.4	(58.7 - 64.0)
Gender						
Male	246	67.5	(58.5 - 76.6)	13,299	67.9	(64.0 - 71.9)
Female	361	54.8	(47.1 - 62.6)	21,894	55.5	(52.0 - 59.0)
Race/Ethnicity						
White Non-Hispanic	597	61.8	(56.0 - 67.7)	33,090	67.8	(65.6 - 70.1)
Black Non-Hispanic	*	*	*	683	41.1	(32.0 - 50.2)
Hispanic	*	*	*	784	53.0	(43.3 - 62.6)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	286	62.5	(53.9 - 71.2)	14,264	59.5	(54.7 - 64.3)
Some College	*	*	*	9,156	59.7	(54.4 - 65.0)
≥ College Degree	*	*	*	11,706	64.5	(60.8 - 68.2)
Income						
≤ \$24,999	*	*	*	8,772	48.7	(42.0 - 55.4)
\$25,000-\$49,999	*	*	*	8,932	63.7	(57.5 - 70.0)
\$50,000-\$74,999	*	*	*	5,433	64.2	(58.2 - 70.2)
≥ \$75,000	*	*	*	8,108	67.5	(64.0 - 71.0)

¹ Defined as respondents who never or rarely (hardly ever) were asked about their sexual history, such as about sexual partners and sexual practices, during a routine check-up at a doctor's office or clinic

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 60a. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	404	8,000	76.7	(71.0 - 82.5)	22,788	7,011,100	77.9	(75.2 - 80.5)
Gender								
Male	175	4,400	82.8	(75.6 - 89.9)	9,034	3,618,500	81.0	(77.2 - 84.8)
Female	229	3,600	70.4	(61.7 - 79.0)	13,754	3,392,600	74.8	(71.2 - 78.4)
Age (years)								
18-34	*	*	*	*	3,213	2,362,500	70.8	(65.1 - 76.6)
35-44	*	*	*	*	3,513	1,313,400	71.7	(65.6 - 77.7)
45-54	80	1,400	78.4	(68.7 - 88.1)	4,975	1,290,600	80.3	(75.9 - 84.8)
55-64	100	1,300	89.2	(82.0 - 96.5)	4,955	932,200	87.2	(84.1 - 90.3)
≥ 65	114	1,400	95.5	(91.5 - 99.5)	6,132	1,112,400	95.9	(94.4 - 97.5)
Race/Ethnicity								
White Non-Hispanic	397	7,700	76.5	(70.7 - 82.3)	21,540	4,875,200	78.3	(75.6 - 80.9)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	491	1,002,400	84.8	(77.8 - 91.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	181	3,900	88.8	(82.7 - 95.0)	8,879	2,581,700	82.1	(77.9 - 86.3)
Some College	*	*	*	*	5,953	1,599,500	77.5	(72.6 - 82.3)
≥ College Degree	*	*	*	*	7,928	2,824,800	75.6	(71.8 - 79.5)
Income								
≤ \$24,999	*	*	*	*	5,491	1,602,900	78.7	(71.9 - 85.6)
\$25,000-\$49,999	122	2,700	80.5	(70.7 - 90.3)	5,912	1,335,900	78.8	(73.8 - 83.9)
\$50,000-\$74,999	*	*	*	*	3,717	987,500	76.6	(70.4 - 82.8)
≥ \$75,000	*	*	*	*	5,771	2,321,600	77.6	(74.0 - 81.2)

¹ Defined as respondent estimating there were hardly any (0-1 out of 10) or a few (2-3 out of 10) persons their age in their community who have had a sexually transmitted disease such as gonorrhea, chlamydia, herpes or syphilis

² Sample size from survey

³ Estimated number of adults who believed hardly any or a few people their age in their community have had a sexually transmitted disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 60b. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	404	74.3	(67.9 - 80.7)	22,788	77.3	(74.7 - 79.9)
Gender						
Male	175	81.0	(73.1 - 88.9)	9,034	81.1	(77.4 - 84.8)
Female	229	67.3	(57.6 - 77.0)	13,754	73.6	(70.0 - 77.3)
Race/Ethnicity						
White Non-Hispanic	397	74.1	(67.6 - 80.6)	21,540	77.5	(74.9 - 80.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	491	84.6	(77.6 - 91.5)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	181	87.4	(80.2 - 94.5)	8,879	81.6	(77.4 - 85.7)
Some College	*	*	*	5,953	76.8	(72.0 - 81.6)
≥ College Degree	*	*	*	7,928	75.1	(71.3 - 79.0)
Income						
≤ \$24,999	*	*	*	5,491	77.9	(71.0 - 84.7)
\$25,000-\$49,999	*	*	*	5,912	77.9	(72.8 - 83.0)
\$50,000-\$74,999	*	*	*	3,717	76.1	(70.0 - 82.3)
≥ \$75,000	*	*	*	5,771	77.0	(73.4 - 80.6)

¹ Defined as respondent estimating there were hardly any (0-1 out of 10) or a few (2-3 out of 10) persons their age in their community who have had a sexually transmitted disease such as gonorrhea, chlamydia, herpes or syphilis

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 61a. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	637	13,200	91.7	(89.5 - 93.9)	36,287	12,738,300	89.7	(88.2 - 91.3)
Gender								
Male	263	6,500	91.3	(87.7 - 94.8)	13,893	6,104,600	89.4	(87.2 - 91.5)
Female	374	6,800	92.1	(89.4 - 94.8)	22,394	6,633,700	90.1	(87.9 - 92.2)
Age (years)								
18-34	56	4,000	98.6	(95.8 - 100)	3,850	4,007,400	91.3	(87.5 - 95.1)
35-44	76	2,300	96.8	(92.3 - 100)	4,875	2,578,600	93.2	(90.4 - 96.0)
45-54	122	2,500	89.5	(83.7 - 95.4)	7,607	2,416,800	90.5	(87.9 - 93.2)
55-64	164	2,200	91.3	(86.4 - 96.2)	8,411	1,876,500	92.2	(90.0 - 94.5)
≥ 65	219	2,200	79.7	(73.9 - 85.5)	11,544	1,858,900	79.7	(76.6 - 82.8)
Race/Ethnicity								
White Non-Hispanic	625	12,800	91.9	(89.6 - 94.1)	34,102	8,050,700	89.4	(87.7 - 91.0)
Black Non-Hispanic	*	*	*	*	702	1,822,200	91.9	(88.2 - 95.6)
Hispanic	*	*	*	*	820	1,967,200	89.9	(85.2 - 94.7)
Other Non-Hispanic	*	*	*	*	663	898,200	88.6	(79.5 - 97.6)
Education								
≤ High School	292	5,400	86.7	(82.5 - 90.8)	14,507	4,385,300	84.6	(81.9 - 87.2)
Some College	173	4,000	95.7	(92.9 - 98.6)	9,571	3,094,300	90.0	(86.7 - 93.4)
≥ College Degree	172	3,800	95.3	(92.0 - 98.7)	12,142	5,228,800	95.4	(94.0 - 96.7)
Income								
≤ \$24,999	186	3,100	84.2	(78.3 - 90.1)	9,021	2,833,500	86.5	(82.6 - 90.3)
\$25,000-\$49,999	182	4,000	93.3	(89.6 - 96.9)	9,261	2,529,900	89.5	(86.5 - 92.5)
\$50,000-\$74,999	105	2,300	93.5	(88.7 - 98.4)	5,644	1,740,000	91.2	(87.4 - 95.1)
≥ \$75,000	101	2,400	95.7	(91.1 - 100)	8,397	4,184,300	94.4	(92.5 - 96.3)

¹ Defined as believing it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community

² Sample size from survey

³ Estimated number of adults who believed it is at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 61b. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	637	92.6	(90.4 - 94.7)	36,287	89.8	(88.3 - 91.4)
Gender						
Male	263	91.6	(88.0 - 95.2)	13,893	89.1	(86.9 - 91.3)
Female	374	93.5	(91.1 - 95.9)	22,394	90.5	(88.3 - 92.7)
Race/Ethnicity						
White Non-Hispanic	625	92.7	(90.5 - 94.9)	34,102	89.5	(87.8 - 91.2)
Black Non-Hispanic	*	*	*	702	91.7	(87.8 - 95.6)
Hispanic	*	*	*	820	89.6	(84.7 - 94.5)
Other Non-Hispanic	*	*	*	663	89.0	(80.5 - 97.6)
Education						
≤ High School	292	88.1	(84.2 - 92.1)	14,507	84.6	(81.8 - 87.4)
Some College	173	96.3	(93.8 - 98.9)	9,571	90.2	(86.9 - 93.5)
≥ College Degree	172	95.4	(91.5 - 99.2)	12,142	95.4	(94.0 - 96.7)
Income						
≤ \$24,999	186	86.0	(80.4 - 91.5)	9,021	86.5	(82.6 - 90.4)
\$25,000-\$49,999	182	94.1	(90.6 - 97.6)	9,261	89.6	(86.5 - 92.6)
\$50,000-\$74,999	105	94.5	(90.2 - 98.8)	5,644	91.1	(86.9 - 95.3)
≥ \$75,000	101	95.0	(89.2 - 100)	8,397	94.5	(92.6 - 96.4)

¹ Defined as believing it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 62a. Overweight among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	647	4,700	32.4	(27.6 - 37.2)	36,710	5,213,000	36.4	(34.0 - 38.8)
Gender								
Male	277	2,800	37.7	(30.1 - 45.3)	14,502	3,073,200	43.7	(39.9 - 47.6)
Female	370	1,900	26.9	(21.4 - 32.3)	22,208	2,139,800	29.3	(26.4 - 32.2)
Age (years)								
18-34	*	*	*	*	3,789	1,295,200	30.1	(24.7 - 35.6)
35-44	*	*	*	*	4,785	1,052,000	37.5	(31.7 - 43.3)
45-54	121	900	31.6	(22.4 - 40.8)	7,495	1,053,300	38.8	(34.1 - 43.4)
55-64	170	800	32.0	(24.4 - 39.6)	8,356	798,100	38.8	(34.8 - 42.7)
≥ 65	226	1,200	42.8	(35.8 - 49.9)	12,285	1,014,400	41.5	(37.6 - 45.4)
Race/Ethnicity								
White Non-Hispanic	634	4,600	33.1	(28.3 - 38.0)	34,501	3,282,200	36.4	(34.4 - 38.3)
Black Non-Hispanic	*	*	*	*	708	740,000	36.7	(28.3 - 45.1)
Hispanic	*	*	*	*	813	781,500	35.6	(27.1 - 44.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	302	2,400	36.4	(28.9 - 43.9)	15,079	1,889,700	35.4	(31.2 - 39.5)
Some College	175	1,100	27.3	(19.1 - 35.5)	9,515	1,314,500	38.7	(33.8 - 43.7)
≥ College Degree	170	1,200	31.2	(22.7 - 39.8)	12,062	1,996,900	36.3	(32.8 - 39.7)
Income								
≤ \$24,999	191	1,200	32.2	(22.6 - 41.8)	9,346	1,170,900	34.0	(28.2 - 39.8)
\$25,000-\$49,999	190	1,700	37.4	(28.1 - 46.6)	9,368	1,136,000	40.4	(34.6 - 46.2)
\$50,000-\$74,999	103	600	26.2	(16.9 - 35.5)	5,615	738,700	38.3	(32.3 - 44.3)
≥ \$75,000	*	*	*	*	8,337	1,659,100	37.5	(34.1 - 40.9)

¹ Defined as having a body mass index (BMI) more than 24.9 but less than 30.0. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Sample size from survey

³ Estimated number of adults who are overweight (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 62b. Overweight among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	647	31.4	(26.3 - 36.5)	36,710	36.2	(33.8 - 38.6)
Gender						
Male	277	37.7	(29.6 - 45.7)	14,502	43.9	(40.1 - 47.6)
Female	370	25.0	(19.1 - 30.9)	22,208	28.9	(25.9 - 31.8)
Race/Ethnicity						
White Non-Hispanic	634	32.1	(26.9 - 37.3)	34,501	36.0	(34.0 - 38.0)
Black Non-Hispanic	*	*	*	708	37.0	(28.5 - 45.5)
Hispanic	*	*	*	813	36.0	(27.5 - 44.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	302	35.3	(27.2 - 43.5)	15,079	35.2	(31.1 - 39.4)
Some College	175	25.5	(16.9 - 34.1)	9,515	38.3	(33.3 - 43.2)
≥ College Degree	170	31.3	(21.9 - 40.6)	12,062	36.2	(32.7 - 39.8)
Income						
≤ \$24,999	*	*	*	9,346	33.8	(28.0 - 39.5)
\$25,000-\$49,999	190	36.8	(27.0 - 46.6)	9,368	40.6	(34.8 - 46.4)
\$50,000-\$74,999	103	25.0	(15.2 - 34.7)	5,615	38.6	(32.5 - 44.7)
≥ \$75,000	*	*	*	8,337	37.2	(33.7 - 40.6)

¹ Defined as having a body mass index (BMI) more than 24.9 but less than 30.0. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 63a. Obesity among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	647	4,800	33.1	(28.3 - 37.9)	36,710	3,327,400	23.2	(21.2 - 25.3)
Gender								
Male	277	2,400	32.6	(25.4 - 39.7)	14,502	1,505,800	21.4	(19.0 - 23.9)
Female	370	2,400	33.7	(27.4 - 40.1)	22,208	1,821,600	25.0	(21.8 - 28.1)
Age (years)								
18-34	*	*	*	*	3,789	730,600	17.0	(12.4 - 21.6)
35-44	*	*	*	*	4,785	662,200	23.6	(19.4 - 27.9)
45-54	121	1,000	34.4	(25.2 - 43.6)	7,495	816,800	30.1	(25.3 - 34.8)
55-64	170	900	34.4	(26.6 - 42.2)	8,356	563,900	27.4	(23.7 - 31.1)
≥ 65	226	800	28.3	(21.9 - 34.8)	12,285	554,000	22.7	(19.2 - 26.1)
Race/Ethnicity								
White Non-Hispanic	634	4,800	34.2	(29.3 - 39.1)	34,501	2,062,700	22.9	(21.2 - 24.6)
Black Non-Hispanic	*	*	*	*	708	703,100	34.9	(26.1 - 43.7)
Hispanic	*	*	*	*	813	456,400	20.8	(14.3 - 27.2)
Other Non-Hispanic	*	*	*	*	688	105,200	9.6	(4.0 - 15.3)
Education								
≤ High School	302	2,100	33.1	(26.4 - 39.8)	15,079	1,561,600	29.2	(25.3 - 33.2)
Some College	*	*	*	*	9,515	792,800	23.4	(19.4 - 27.3)
≥ College Degree	170	1,000	26.9	(18.1 - 35.8)	12,062	915,000	16.6	(14.4 - 18.8)
Income								
≤ \$24,999	191	1,500	41.6	(31.7 - 51.6)	9,346	974,800	28.3	(22.6 - 34.0)
\$25,000-\$49,999	190	1,400	32.5	(23.4 - 41.7)	9,368	584,800	20.8	(17.1 - 24.5)
\$50,000-\$74,999	*	*	*	*	5,615	475,300	24.6	(20.0 - 29.3)
≥ \$75,000	100	700	27.7	(18.2 - 37.2)	8,337	867,700	19.6	(17.0 - 22.3)

¹ Defined as having a body mass index (BMI) of 30.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Sample size from survey

³ Estimated number of adults who are obese (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 63b. Obesity among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	647	33.5	(28.2 - 38.8)	36,710	23.1	(21.0 - 25.1)
Gender						
Male	277	33.2	(25.5 - 40.9)	14,502	21.5	(19.0 - 24.0)
Female	370	33.8	(26.6 - 41.0)	22,208	24.6	(21.5 - 27.8)
Race/Ethnicity						
White Non-Hispanic	634	34.6	(29.2 - 40.0)	34,501	22.8	(21.0 - 24.5)
Black Non-Hispanic	*	*	*	708	34.8	(25.9 - 43.6)
Hispanic	*	*	*	813	20.5	(14.1 - 26.9)
Other Non-Hispanic	*	*	*	688	9.5	(3.8 - 15.2)
Education						
≤ High School	302	32.9	(25.5 - 40.3)	15,079	29.4	(25.4 - 33.3)
Some College	*	*	*	9,515	23.1	(19.2 - 27.1)
≥ College Degree	170	28.1	(18.2 - 38.0)	12,062	16.4	(14.2 - 18.5)
Income						
≤ \$24,999	*	*	*	9,346	28.3	(22.6 - 34.1)
\$25,000-\$49,999	190	33.4	(23.5 - 43.2)	9,368	21.0	(17.2 - 24.8)
\$50,000-\$74,999	*	*	*	5,615	24.0	(19.4 - 28.7)
≥ \$75,000	*	*	*	8,337	19.3	(16.7 - 21.9)

¹ Defined as having a body mass index (BMI) of 30.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 64a. Overweight or obesity among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	647	9,500	65.5	(60.5 - 70.5)	36,710	8,540,500	59.6	(57.0 - 62.2)
Gender								
Male	277	5,200	70.3	(62.9 - 77.7)	14,502	4,579,100	65.2	(61.0 - 69.4)
Female	370	4,300	60.6	(54.0 - 67.2)	22,208	3,961,400	54.3	(51.0 - 57.6)
Age (years)								
18-34	*	*	*	*	3,789	2,025,800	47.1	(40.8 - 53.5)
35-44	*	*	*	*	4,785	1,714,200	61.1	(55.6 - 66.6)
45-54	121	1,900	66.0	(56.7 - 75.4)	7,495	1,870,100	68.8	(64.6 - 73.1)
55-64	170	1,600	66.5	(58.5 - 74.4)	8,356	1,362,000	66.2	(61.9 - 70.4)
≥ 65	226	2,000	71.1	(64.8 - 77.5)	12,285	1,568,400	64.1	(60.6 - 67.7)
Race/Ethnicity								
White Non-Hispanic	634	9,400	67.4	(62.5 - 72.2)	34,501	5,344,900	59.3	(57.2 - 61.3)
Black Non-Hispanic	*	*	*	*	708	1,443,100	71.6	(62.5 - 80.7)
Hispanic	*	*	*	*	813	1,237,900	56.4	(46.5 - 66.2)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	302	4,500	69.5	(62.6 - 76.3)	15,079	3,451,400	64.6	(59.8 - 69.4)
Some College	*	*	*	*	9,515	2,107,300	62.1	(57.0 - 67.2)
≥ College Degree	170	2,200	58.2	(48.6 - 67.7)	12,062	2,911,900	52.9	(49.2 - 56.6)
Income								
≤ \$24,999	191	2,700	73.9	(65.5 - 82.2)	9,346	2,145,700	62.3	(55.7 - 68.9)
\$25,000-\$49,999	190	3,100	69.9	(60.3 - 79.5)	9,368	1,720,700	61.2	(54.5 - 67.9)
\$50,000-\$74,999	*	*	*	*	5,615	1,214,000	62.9	(57.1 - 68.8)
≥ \$75,000	*	*	*	*	8,337	2,526,700	57.1	(53.6 - 60.6)

¹ Defined as having a body mass index (BMI) of 25.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Sample size from survey

³ Estimated number of adults who are overweight or obese (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 64b. Overweight or obesity among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	647	64.9	(59.4 - 70.4)	36,710	59.3	(56.8 - 61.9)
Gender						
Male	277	70.8	(63.1 - 78.5)	14,502	65.4	(61.3 - 69.4)
Female	370	58.8	(51.2 - 66.3)	22,208	53.5	(50.1 - 56.8)
Race/Ethnicity						
White Non-Hispanic	634	66.7	(61.4 - 72.0)	34,501	58.8	(56.7 - 60.8)
Black Non-Hispanic	*	*	*	708	71.8	(63.0 - 80.6)
Hispanic	*	*	*	813	56.5	(46.8 - 66.1)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	302	68.2	(60.5 - 76.0)	15,079	64.6	(59.9 - 69.3)
Some College	*	*	*	9,515	61.4	(56.4 - 66.5)
≥ College Degree	*	*	*	12,062	52.6	(48.9 - 56.3)
Income						
≤ \$24,999	191	73.7	(64.5 - 83.0)	9,346	62.1	(55.6 - 68.6)
\$25,000-\$49,999	*	*	*	9,368	61.6	(55.1 - 68.1)
\$50,000-\$74,999	*	*	*	5,615	62.6	(56.7 - 68.6)
≥ \$75,000	*	*	*	8,337	56.5	(52.9 - 60.0)

¹ Defined as having a body mass index (BMI) of 25.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 65a. Received advice about weight by a health professional among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	649	4,000	27.5	(23.1 - 31.9)	36,729	3,929,700	27.4	(25.1 - 29.7)
Gender								
Male	278	1,800	24.6	(18.5 - 30.7)	14,499	1,654,800	23.6	(20.6 - 26.6)
Female	371	2,200	30.5	(24.3 - 36.7)	22,230	2,274,900	31.0	(27.7 - 34.3)
Age (years)								
18-34	*	*	*	*	3,796	895,500	20.7	(15.3 - 26.2)
35-44	*	*	*	*	4,779	722,700	25.8	(20.9 - 30.6)
45-54	121	800	29.3	(20.8 - 37.7)	7,490	910,900	33.5	(29.0 - 38.1)
55-64	171	700	28.6	(21.4 - 35.9)	8,351	741,100	36.1	(32.0 - 40.3)
≥ 65	227	900	32.7	(25.9 - 39.6)	12,313	659,500	26.9	(23.0 - 30.7)
Race/Ethnicity								
White Non-Hispanic	636	3,900	27.7	(23.2 - 32.1)	34,502	2,386,200	26.5	(24.6 - 28.4)
Black Non-Hispanic	*	*	*	*	709	615,700	30.4	(22.0 - 38.8)
Hispanic	*	*	*	*	830	555,400	25.1	(17.8 - 32.4)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	303	2,000	31.4	(24.6 - 38.2)	15,116	1,469,700	27.5	(23.6 - 31.4)
Some College	176	1,000	23.3	(15.9 - 30.6)	9,515	946,800	27.8	(23.2 - 32.4)
≥ College Degree	170	1,000	25.7	(16.9 - 34.4)	12,041	1,445,500	26.2	(23.0 - 29.4)
Income								
≤ \$24,999	191	1,300	34.6	(24.6 - 44.6)	9,360	1,079,400	31.4	(25.3 - 37.5)
\$25,000-\$49,999	191	1,300	28.2	(20.2 - 36.3)	9,365	696,900	24.6	(19.8 - 29.4)
\$50,000-\$74,999	103	600	27.7	(17.8 - 37.6)	5,603	507,100	26.3	(21.6 - 31.0)
≥ \$75,000	100	600	22.2	(13.7 - 30.7)	8,334	1,207,100	27.3	(24.2 - 30.5)

¹ Defined as being advised by a doctor, nurse or other health professional about their weight (lose, gain or maintain)

² Sample size from survey

³ Estimated number of adults provided advice by a health professional about their weight (lose, gain or maintain) (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 65b. Received advice about weight by a health professional among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	649	27.1	(22.3 - 31.9)	36,729	27.1	(24.8 - 29.4)
Gender						
Male	278	24.6	(18.0 - 31.1)	14,499	23.4	(20.5 - 26.3)
Female	371	29.7	(22.8 - 36.7)	22,230	30.6	(27.3 - 34.0)
Race/Ethnicity						
White Non-Hispanic	636	27.2	(22.3 - 32.0)	34,502	26.1	(24.2 - 28.0)
Black Non-Hispanic	*	*	*	709	30.4	(21.9 - 38.9)
Hispanic	*	*	*	830	24.9	(17.7 - 32.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	303	31.5	(24.0 - 39.0)	15,116	27.3	(23.4 - 31.2)
Some College	176	21.0	(13.5 - 28.5)	9,515	27.4	(22.9 - 32.0)
≥ College Degree	170	26.4	(16.7 - 36.2)	12,041	25.9	(22.7 - 29.0)
Income						
≤ \$24,999	*	*	*	9,360	31.0	(24.9 - 37.0)
\$25,000-\$49,999	191	28.9	(20.0 - 37.8)	9,365	24.9	(20.0 - 29.8)
\$50,000-\$74,999	*	*	*	5,603	25.9	(21.2 - 30.6)
≥ \$75,000	100	21.3	(12.3 - 30.3)	8,334	26.9	(23.7 - 30.0)

¹ Defined as being advised by a doctor, nurse or other health professional about their weight (lose, gain or maintain)

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 66a. Received advice to lose weight by a health professional among adults provided weight advice¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	189	3,300	82.6	(75.2 - 90.0)	9,724	3,412,900	86.8	(83.9 - 89.8)
Gender								
Male	*	*	*	*	3,807	1,468,700	88.8	(85.7 - 91.8)
Female	109	1,800	84.1	(76.3 - 91.9)	5,917	1,944,200	85.5	(80.8 - 90.1)
Age (years)								
18-34	*	*	*	*	681	779,700	87.1	(80.6 - 93.6)
35-44	*	*	*	*	1,215	622,100	86.1	(76.9 - 95.2)
45-54	*	*	*	*	2,238	813,300	89.3	(84.4 - 94.2)
55-64	*	*	*	*	2,697	645,500	87.1	(81.2 - 93.0)
≥ 65	*	*	*	*	2,893	552,300	83.7	(76.9 - 90.6)
Race/Ethnicity								
White Non-Hispanic	185	3,200	83.8	(76.3 - 91.3)	9,082	2,076,700	87.0	(84.5 - 89.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	251	483,600	87.1	(78.5 - 95.7)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	3,962	1,239,700	84.3	(78.6 - 90.1)
Some College	*	*	*	*	2,633	814,800	86.1	(80.1 - 92.0)
≥ College Degree	*	*	*	*	3,115	1,290,900	89.3	(85.3 - 93.3)
Income								
≤ \$24,999	*	*	*	*	2,539	922,100	85.4	(78.7 - 92.1)
\$25,000-\$49,999	59	1,200	91.5	(84.1 - 98.9)	2,482	594,100	85.2	(78.4 - 92.1)
\$50,000-\$74,999	*	*	*	*	1,521	446,100	88.0	(83.2 - 92.7)
≥ \$75,000	*	*	*	*	2,270	1,073,200	88.9	(83.7 - 94.1)

¹ Defined as provided advice by a health professional about their weight and advised to lose weight

² Sample size from survey

³ Estimated number of adults provided advice by a health professional about their weight and advised to lose weight (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 66b. Received advice to lose weight by a health professional among adults provided weight advice¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	189	83.2	(75.3 - 91.2)	9,724	86.9	(83.8 - 89.9)
Gender						
Male	*	*	*	3,807	88.7	(85.7 - 91.8)
Female	109	86.0	(77.9 - 94.0)	5,917	85.5	(80.7 - 90.3)
Race/Ethnicity						
White Non-Hispanic	185	84.1	(76.0 - 92.2)	9,082	87.0	(84.4 - 89.6)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	251	87.6	(79.3 - 95.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	3,962	84.6	(78.7 - 90.5)
Some College	*	*	*	2,633	85.6	(79.5 - 91.7)
≥ College Degree	*	*	*	3,115	89.4	(85.4 - 93.3)
Income						
≤ \$24,999	*	*	*	2,539	85.7	(79.0 - 92.3)
\$25,000-\$49,999	59	91.6	(83.6 - 99.7)	2,482	85.3	(78.4 - 92.3)
\$50,000-\$74,999	*	*	*	1,521	88.0	(83.1 - 92.8)
≥ \$75,000	*	*	*	2,270	88.7	(83.1 - 94.2)

¹ Defined as provided advice by a health professional about their weight and advised to lose weight

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 67a. No leisure-time physical activity among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	668	3,200	21.1	(17.2 - 25.0)	38,128	3,541,700	23.9	(21.6 - 26.1)
Gender								
Male	279	1,500	19.7	(14.1 - 25.4)	14,602	1,482,500	20.9	(17.8 - 24.0)
Female	389	1,700	22.4	(17.2 - 27.7)	23,526	2,059,200	26.6	(23.4 - 29.7)
Age (years)								
18-34	*	*	*	*	3,938	982,900	22.0	(16.6 - 27.5)
35-44	77	500	20.2	(10.3 - 30.0)	4,988	556,000	19.3	(15.0 - 23.6)
45-54	127	600	19.2	(11.2 - 27.2)	7,827	759,000	26.8	(21.7 - 31.9)
55-64	173	600	24.3	(17.5 - 31.0)	8,765	464,900	21.7	(18.2 - 25.2)
≥ 65	234	900	31.8	(25.2 - 38.4)	12,610	778,800	30.7	(27.3 - 34.2)
Race/Ethnicity								
White Non-Hispanic	655	3,000	20.7	(16.9 - 24.6)	35,816	1,892,800	20.3	(18.6 - 22.0)
Black Non-Hispanic	*	*	*	*	739	690,000	32.0	(23.4 - 40.6)
Hispanic	*	*	*	*	860	600,800	26.7	(19.1 - 34.4)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	312	2,100	31.4	(24.7 - 38.2)	15,683	1,810,200	32.5	(28.4 - 36.5)
Some College	182	500	12.7	(7.1 - 18.3)	9,906	814,200	23.0	(18.6 - 27.3)
≥ College Degree	174	500	13.1	(7.4 - 18.7)	12,458	841,000	14.9	(11.9 - 18.0)
Income								
≤ \$24,999	193	1,200	31.9	(22.8 - 41.0)	9,640	1,332,200	38.0	(31.7 - 44.3)
\$25,000-\$49,999	195	900	20.6	(13.7 - 27.6)	9,658	648,000	22.3	(18.3 - 26.3)
\$50,000-\$74,999	109	300	13.2	(6.6 - 19.8)	5,773	430,700	21.8	(16.6 - 27.0)
≥ \$75,000	102	300	10.7	(3.9 - 17.4)	8,565	560,800	12.5	(10.3 - 14.6)

¹ Defined as having not participated in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise

² Sample size from survey

³ Estimated number of adults who have not participated in any physical activities or exercises (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 67b. No leisure-time physical activity among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	668	20.4	(16.3 - 24.6)	38,128	23.7	(21.5 - 25.9)
Gender						
Male	279	19.2	(13.3 - 25.0)	14,602	20.9	(17.8 - 23.9)
Female	389	21.7	(15.8 - 27.6)	23,526	26.3	(23.1 - 29.5)
Race/Ethnicity						
White Non-Hispanic	655	20.1	(15.9 - 24.4)	35,816	20.0	(18.3 - 21.7)
Black Non-Hispanic	*	*	*	739	32.0	(23.3 - 40.6)
Hispanic	*	*	*	860	26.9	(19.2 - 34.5)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	312	31.0	(23.6 - 38.4)	15,683	32.3	(28.3 - 36.4)
Some College	182	12.0	(6.0 - 18.0)	9,906	22.8	(18.5 - 27.2)
≥ College Degree	174	12.6	(6.6 - 18.6)	12,458	14.9	(11.8 - 17.9)
Income						
≤ \$24,999	193	30.2	(20.2 - 40.1)	9,640	37.9	(31.6 - 44.2)
\$25,000-\$49,999	195	21.0	(13.3 - 28.7)	9,658	22.0	(18.0 - 26.0)
\$50,000-\$74,999	109	12.4	(5.6 - 19.3)	5,773	21.6	(16.3 - 26.8)
≥ \$75,000	102	9.3	(3.1 - 15.5)	8,565	12.6	(10.3 - 14.9)

¹ Defined as having not participated in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schuyler County**

Table 68a. Consumption of 5 or more servings of fruits and vegetables per day among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	670	3,800	25.1	(20.8 - 29.4)	38,203	4,023,400	27.0	(24.9 - 29.2)
Gender								
Male	280	1,600	21.0	(14.4 - 27.6)	14,634	1,603,900	22.6	(19.4 - 25.8)
Female	390	2,200	29.1	(23.5 - 34.7)	23,569	2,419,500	31.1	(28.2 - 34.0)
Age (years)								
18-34	*	*	*	*	3,940	1,261,700	28.3	(22.7 - 33.9)
35-44	*	*	*	*	4,989	688,000	23.9	(19.5 - 28.3)
45-54	127	900	29.2	(20.5 - 37.9)	7,837	746,800	26.2	(22.3 - 30.1)
55-64	174	700	27.4	(20.0 - 34.9)	8,777	579,100	27.0	(23.6 - 30.5)
≥ 65	235	700	23.5	(17.6 - 29.4)	12,660	747,800	29.4	(26.0 - 32.8)
Race/Ethnicity								
White Non-Hispanic	657	3,600	25.1	(20.7 - 29.4)	35,884	2,538,000	27.2	(25.4 - 28.9)
Black Non-Hispanic	*	*	*	*	740	572,400	26.5	(19.2 - 33.9)
Hispanic	*	*	*	*	864	672,200	29.8	(21.0 - 38.5)
Other Non-Hispanic	*	*	*	*	715	240,800	21.4	(12.0 - 30.8)
Education								
≤ High School	314	1,300	19.8	(13.3 - 26.2)	15,726	1,309,800	23.4	(19.3 - 27.4)
Some College	182	1,200	28.1	(19.4 - 36.7)	9,924	964,300	27.2	(23.0 - 31.3)
≥ College Degree	174	1,200	30.7	(22.6 - 38.8)	12,471	1,747,500	31.0	(27.9 - 34.1)
Income								
≤ \$24,999	193	600	15.3	(8.9 - 21.7)	9,667	831,500	23.7	(18.6 - 28.7)
\$25,000-\$49,999	196	1,100	23.3	(14.7 - 32.0)	9,669	917,200	31.6	(25.6 - 37.6)
\$50,000-\$74,999	*	*	*	*	5,777	501,000	25.3	(21.4 - 29.3)
≥ \$75,000	*	*	*	*	8,569	1,260,100	28.0	(25.0 - 31.0)

¹ Defined as having consumed 5 or more servings of fruits and vegetables in an average day

² Sample size from survey

³ Estimated number of adults who consume 5 or more servings of fruits and vegetables per day (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 68b. Consumption of 5 or more servings of fruits and vegetables per day among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	670	24.6	(20.0 - 29.3)	38,203	27.1	(24.9 - 29.2)
Gender						
Male	280	21.0	(14.0 - 27.9)	14,634	22.7	(19.6 - 25.9)
Female	390	28.2	(21.9 - 34.4)	23,569	31.0	(28.1 - 34.0)
Race/Ethnicity						
White Non-Hispanic	657	24.5	(19.9 - 29.2)	35,884	27.1	(25.4 - 28.9)
Black Non-Hispanic	*	*	*	740	26.8	(19.4 - 34.2)
Hispanic	*	*	*	864	29.5	(20.9 - 38.1)
Other Non-Hispanic	*	*	*	715	21.8	(12.2 - 31.5)
Education						
≤ High School	314	19.7	(12.7 - 26.8)	15,726	23.3	(19.3 - 27.3)
Some College	182	27.5	(18.2 - 36.8)	9,924	27.1	(22.9 - 31.2)
≥ College Degree	174	29.3	(20.9 - 37.7)	12,471	31.2	(28.0 - 34.3)
Income						
≤ \$24,999	193	14.6	(7.5 - 21.8)	9,667	23.9	(18.8 - 28.9)
\$25,000-\$49,999	196	23.6	(14.3 - 32.8)	9,669	32.0	(26.0 - 38.0)
\$50,000-\$74,999	*	*	*	5,777	25.0	(21.0 - 28.9)
≥ \$75,000	*	*	*	8,569	28.0	(24.9 - 31.1)

¹ Defined as having consumed 5 or more servings of fruits and vegetables in an average day

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 69a. Had a fall within the past 3 months among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	526	1,700	19.9	(16.2 - 23.7)	28,783	1,244,800	17.0	(14.9 - 19.0)
Gender								
Male	216	700	18.0	(12.5 - 23.6)	10,966	489,400	14.4	(11.9 - 16.8)
Female	310	900	21.7	(16.5 - 26.8)	17,817	755,300	19.2	(16.1 - 22.3)
Age (years)								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	126	400	15.3	(8.7 - 22.0)	7,741	445,500	16.0	(12.0 - 20.0)
55-64	166	600	23.4	(16.3 - 30.5)	8,454	336,000	16.5	(13.4 - 19.6)
≥ 65	234	600	21.6	(15.8 - 27.4)	12,588	463,300	18.4	(15.2 - 21.6)
Race/Ethnicity								
White Non-Hispanic	515	1,600	19.9	(16.1 - 23.8)	27,456	869,000	16.6	(15.1 - 18.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	424	76,100	11.0	(4.5 - 17.4)
Other Non-Hispanic	*	*	*	*	445	44,300	10.0	(3.1 - 16.8)
Education								
≤ High School	255	800	21.3	(15.6 - 26.9)	12,417	493,000	17.3	(13.3 - 21.3)
Some College	143	500	20.1	(12.9 - 27.2)	7,232	299,400	18.7	(14.6 - 22.7)
≥ College Degree	128	400	17.3	(10.2 - 24.4)	9,086	441,500	15.4	(12.9 - 17.9)
Income								
≤ \$24,999	162	500	20.4	(13.6 - 27.3)	7,824	423,900	25.0	(18.5 - 31.6)
\$25,000-\$49,999	149	400	20.3	(13.4 - 27.1)	7,445	223,600	16.0	(12.8 - 19.2)
\$50,000-\$74,999	84	300	21.6	(11.9 - 31.3)	4,141	151,200	14.6	(10.8 - 18.4)
≥ \$75,000	76	300	15.7	(6.8 - 24.6)	5,874	287,100	12.5	(10.0 - 15.1)

¹ Defined as respondent unintentionally coming to rest on the ground or another lower level one or more times within the past 3 months

² Sample size from survey

³ Estimated number of adults who have had a fall within the past 3 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 69b. Had a fall within the past 3 months among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	526	19.4	(15.7 - 23.2)	28,783	17.3	(15.0 - 19.5)
Gender						
Male	216	17.7	(12.2 - 23.3)	10,966	15.0	(12.1 - 17.8)
Female	310	21.1	(15.9 - 26.2)	17,817	19.4	(16.0 - 22.8)
Race/Ethnicity						
White Non-Hispanic	515	19.5	(15.6 - 23.3)	27,456	16.9	(15.2 - 18.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	424	10.7	(4.2 - 17.1)
Other Non-Hispanic	*	*	*	445	10.0	(2.9 - 17.0)
Education						
≤ High School	255	20.8	(15.1 - 26.5)	12,417	17.4	(13.0 - 21.7)
Some College	143	19.7	(12.5 - 26.9)	7,232	19.0	(14.7 - 23.2)
≥ College Degree	128	16.6	(9.7 - 23.5)	9,086	16.0	(13.1 - 18.8)
Income						
≤ \$24,999	162	19.5	(12.9 - 26.2)	7,824	25.3	(18.3 - 32.4)
\$25,000-\$49,999	149	20.2	(13.3 - 27.1)	7,445	15.9	(12.6 - 19.2)
\$50,000-\$74,999	84	20.3	(10.7 - 29.9)	4,141	14.7	(10.6 - 18.8)
≥ \$75,000	76	15.4	(6.4 - 24.4)	5,874	12.9	(9.9 - 15.8)

¹ Defined as respondent unintentionally coming to rest on the ground or another lower level one or more times within the past 3 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 70a. Had a fall that caused injury within the past 3 months among adults¹ in Schuyler County and New York State

	Schuyler County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	525	500	6.3	(3.8 - 8.7)	28,760	335,700	4.6	(3.7 - 5.5)
Gender								
Male	216	300	7.1	(2.9 - 11.3)	10,957	120,000	3.5	(2.5 - 4.6)
Female	309	200	5.5	(2.8 - 8.2)	17,803	215,600	5.5	(4.1 - 6.9)
Age (years)								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	126	200	7.6	(2.7 - 12.6)	7,737	115,100	4.1	(2.9 - 5.4)
55-64	166	200	6.6	(2.1 - 11.2)	8,449	90,300	4.4	(2.7 - 6.1)
≥ 65	233	100	4.6	(1.6 - 7.5)	12,574	130,300	5.2	(3.4 - 6.9)
Race/Ethnicity								
White Non-Hispanic	514	500	6.2	(3.7 - 8.7)	27,433	270,800	5.2	(4.2 - 6.1)
Black Non-Hispanic	*	*	*	*	458	47,900	4.9	(0.9 - 9.0)
Hispanic	*	*	*	*	424	12,400	1.8	(0.2 - 3.3)
Other Non-Hispanic	*	*	*	*	445	4,600	1.0	(0.4 - 1.6)
Education								
≤ High School	254	300	7.8	(3.7 - 11.9)	12,404	148,900	5.2	(3.6 - 6.8)
Some College	*	*	*	*	7,228	85,400	5.3	(3.1 - 7.5)
≥ College Degree	*	*	*	*	9,081	101,300	3.5	(2.4 - 4.6)
Income								
≤ \$24,999	162	200	7.1	(2.9 - 11.4)	7,812	90,200	5.3	(3.6 - 7.1)
\$25,000-\$49,999	*	*	*	*	7,442	93,200	6.7	(3.9 - 9.4)
\$50,000-\$74,999	*	*	*	*	4,139	37,000	3.6	(1.8 - 5.3)
≥ \$75,000	*	*	*	*	5,872	54,700	2.4	(1.5 - 3.3)

¹ Defined as as respondent unintentionally coming to rest on the ground or another lower level and caused respondent to limit regular activities for at least a day or to go see a doctor one or more times within the past 3 months

² Sample size from survey

³ Estimated number of adults who have had a fall that caused an injury within the past 3 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schuyler County

Table 70b. Had a fall that caused injury within the past 3 months among adults¹ in Schuyler County and New York State - Age-Adjusted²

	Schuyler County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	525	6.2	(3.8 - 8.7)	28,760	4.6	(3.6 - 5.5)
Gender						
Male	216	7.1	(2.9 - 11.3)	10,957	3.6	(2.4 - 4.7)
Female	309	5.4	(2.7 - 8.1)	17,803	5.5	(4.1 - 6.9)
Race/Ethnicity						
White Non-Hispanic	514	6.3	(3.7 - 8.8)	27,433	5.2	(4.2 - 6.2)
Black Non-Hispanic	*	*	*	458	4.7	(0.7 - 8.7)
Hispanic	*	*	*	424	1.8	(0.1 - 3.4)
Other Non-Hispanic	*	*	*	445	1.0	(0.4 - 1.6)
Education						
≤ High School	254	8.1	(3.8 - 12.4)	12,404	5.2	(3.5 - 6.8)
Some College	*	*	*	7,228	5.3	(3.1 - 7.5)
≥ College Degree	*	*	*	9,081	3.6	(2.4 - 4.7)
Income						
≤ \$24,999	162	7.0	(2.8 - 11.2)	7,812	5.1	(3.3 - 7.0)
\$25,000-\$49,999	*	*	*	7,442	6.6	(3.9 - 9.2)
\$50,000-\$74,999	*	*	*	4,139	3.7	(1.7 - 5.8)
≥ \$75,000	*	*	*	5,872	2.4	(1.4 - 3.3)

¹ Defined as as respondent unintentionally coming to rest on the ground or another lower level and caused respondent to limit regular activities for at least a day or to go see a doctor one or more times within the past 3 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)