

# **New York State Expanded Behavioral Risk Factor Surveillance System**

**Washington County**

**Final Report  
July 2008 - June 2009**

**New York State Department of Health  
December 2009**

## Contents

<b>Introduction</b>	12
Table 1. Washington County and New York State adult US Census Bureau population estimates, 2007	13
<b>General Health</b>	
Table 2a. Fair or poor general health among adults in Washington County and New York State	14
Table 2b. Fair or poor general health among adults in Washington County and New York State - Age-adjusted	15
Table 3a. Poor physical health 14 or more days within the past month among adults in Washington County and New York State	16
Table 3b. Poor physical health 14 or more days within the past month among adults in Washington County and New York State - Age-adjusted	17
Table 4a. Poor mental health 14 or more days within the past month among adults in Washington County and New York State	18
Table 4b. Poor mental health 14 or more days within the past month among adults in Washington County and New York State - Age-adjusted	19
Table 5a. Poor physical or mental health 14 or more days within the past month among adults in Washington County and New York State	20
Table 5b. Poor physical or mental health 14 or more days within the past month among adults in Washington County and New York State - Age-adjusted	21
<b>Quality Health Care</b>	
Table 6a. Health insurance among adults in Washington County and New York State	22
Table 6b. Health insurance among adults in Washington County and New York State - Age-adjusted	23
Table 7a. Health insurance among adults aged 18-64 in Washington County and New York State	24
Table 7b. Health insurance among adults aged 18-64 in Washington County and New York State - Age-adjusted	25

## Contents

Table 8a. No health insurance among adults in Washington County and New York State	26
Table 8b. No health insurance among adults in Washington County and New York State - Age-adjusted	27
Table 9a. No health insurance among adults aged 18-64 in Washington County and New York State	28
Table 9b. No health insurance among adults aged 18-64 in Washington County and New York State - Age-adjusted	29
Table 10a. Adults with regular health care provider in Washington County and New York State	30
Table 10b. Adults with regular health care provider in Washington County and New York State - Age-adjusted	31
Table 11a. Cost prevented visit to doctor within the past year among adults in Washington County and New York State	32
Table 11b. Cost prevented visit to doctor within the past year among adults in Washington County and New York State - Age-adjusted	33
Table 12a. Visited a doctor for routine checkup within the past year among adults in Washington County and New York State	34
Table 12b. Visited a doctor for routine checkup within the past year among adults in Washington County and New York State - Age-adjusted	35
Table 13a. Visited a doctor for routine checkup within the past 2 years among adults in Washington County and New York State	36
Table 13b. Visited a doctor for routine checkup within the past 2 years among adults in Washington County and New York State - Age-adjusted	37
Table 14a. Dentist visit within the past year among adults in Washington County and New York State	38
Table 14b. Dentist visit within the past year among adults in Washington County and New York State - Age-adjusted	39
Table 15a. Had teeth cleaned within the past year among adults in Washington County and New York State	40
Table 15b. Had teeth cleaned within the past year among adults in Washington County and New York State - Age-adjusted	41

## Contents

Table 16a. Permanent teeth extracted due to decay or gum disease among adults in Washington County and New York State	42
Table 16b. Permanent teeth extracted due to decay or gum disease among adults in Washington County and New York State - Age-adjusted	43
Table 17a. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older in Washington County and New York State	44
Table 17b. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older in Washington County and New York State - Age-adjusted	45
Table 18a. Flu shot within the past 12 months in Washington County and New York State	46
Table 18b. Flu shot within the past 12 months in Washington County and New York State - Age-adjusted	47
Table 19a. Flu vaccine that was sprayed in nose within the past 12 months in Washington County and New York State	48
Table 19b. Flu vaccine that was sprayed in nose within the past 12 months in Washington County and New York State - Age-adjusted	49
Table 20a. Flu shot or flu vaccine sprayed in nose within the past 12 months in Washington County and New York State	50
Table 20b. Flu shot or flu vaccine sprayed in nose within the past 12 months in Washington County and New York State - Age-adjusted	51
Table 21a. Flu shot among adults aged 65 and older in Washington County and New York State	52
Table 21b. Flu shot among adults aged 65 and older in Washington County and New York State - Age-adjusted	53
Table 22a. Pneumonia shot or pneumococcal vaccine in Washington County and New York State	54
Table 22b. Pneumonia shot or pneumococcal vaccine in Washington County and New York State - Age-adjusted	55
Table 23a. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older in Washington County and New York State	56

## Contents

Table 23b. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older in Washington County and New York State - Age-adjusted	57
---	----

### Cancer Screening

Table 24a. Ever had a mammogram among women aged 40 and older in Washington County and New York State	58
---	----

Table 24b. Ever had a mammogram among women aged 40 and older in Washington County and New York State - Age-adjusted	59
--	----

Table 25a. Had mammogram within the past 2 years among women aged 40 and older in Washington County and New York State	60
--	----

Table 25b. Had mammogram within the past 2 years among women aged 40 and older in Washington County and New York State - Age-adjusted	61
---	----

Table 26a. Had mammogram within the past 2 years among women aged 50 and older in Washington County and New York State	62
--	----

Table 26b. Had mammogram within the past 2 years among women aged 50 and older in Washington County and New York State - Age-adjusted	63
---	----

Table 27a. Ever had a Pap test among women in Washington County and New York State	64
--	----

Table 27b. Ever had a Pap test among women in Washington County and New York State - Age-adjusted	65
---	----

Table 28a. Had a Pap test within the past 3 years among women in Washington County and New York State	66
---	----

Table 28b. Had a Pap test within the past 3 years among women in Washington County and New York State - Age-adjusted	67
--	----

Table 29a. Ever had a digital rectal exam among men aged 40 and older in Washington County and New York State	68
---	----

Table 29b. Ever had a digital rectal exam among men aged 40 and older in Washington County and New York State - Age-adjusted	69
--	----

Table 30a. Had digital rectal exam within the past 2 years among men aged 40 and older in Washington County and New York State	70
--	----

## Contents

Table 30b. Had digital rectal exam within the past 2 years among men aged 40 and older in Washington County and New York State - Age-adjusted	71
Table 31a. Ever had a prostate specific antigen test among men aged 40 and older in Washington County and New York State	72
Table 31b. Ever had a prostate specific antigen test among men aged 40 and older in Washington County and New York State - Age-adjusted	73
Table 32a. Had prostate specific antigen test within the past 2 years among men aged 40 and older in Washington County and New York State	74
Table 32b. Had prostate specific antigen test within the past 2 years among men aged 40 and older in Washington County and New York State - Age-adjusted	75
Table 33a. Prostate cancer among men aged 40 and older in Washington County and New York State	76
Table 33b. Prostate cancer among men aged 40 and older in Washington County and New York State - Age-adjusted	77
Table 34a. Home blood stool test ever used among adults aged 50 and older in Washington County and New York State	78
Table 34b. Home blood stool test ever used among adults aged 50 and older in Washington County and New York State - Age-adjusted	79
Table 35a. Home blood stool test used within the past year among adults aged 50 and older in Washington County and New York State	80
Table 35b. Home blood stool test used within the past year among adults aged 50 and older in Washington County and New York State - Age-adjusted	81
Table 36a. Home blood stool test used within the past 2 years among adults aged 50 and older in Washington County and New York State	82
Table 36b. Home blood stool test used within the past 2 years among adults aged 50 and older in Washington County and New York State - Age-adjusted	83
Table 37a. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in Washington County and New York State	84
Table 37b. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in Washington County and New York State - Age-adjusted	85

## Contents

Table 38a. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Washington County and New York State	86
Table 38b. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Washington County and New York State - Age-adjusted	87
Table 39a. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Washington County and New York State	88
Table 39b. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Washington County and New York State - Age-adjusted	89

### Chronic Diseases

Table 40a. High blood pressure among adults in Washington County and New York State	90
Table 40b. High blood pressure among adults in Washington County and New York State - Age-adjusted	91
Table 41a. Blood pressure medication among adults with high blood pressure in Washington County and New York State	92
Table 41b. Blood pressure medication among adults with high blood pressure in Washington County and New York State - Age-adjusted	93
Table 42a. Ever had blood cholesterol checked among adults in Washington County and New York State	94
Table 42b. Ever had blood cholesterol checked among adults in Washington County and New York State - Age-adjusted	95
Table 43a. Blood cholesterol checked within the past 5 years among adults in Washington County and New York State	96
Table 43b. Blood cholesterol checked within the past 5 years among adults in Washington County and New York State - Age-adjusted	97
Table 44a. Cardiovascular disease among adults in Washington County and New York State	98
Table 44b. Cardiovascular disease among adults in Washington County and New York State - Age-adjusted	99

## Contents

Table 45a. Coronary heart disease among adults in Washington County and New York State	100
Table 45b. Coronary heart disease among adults in Washington County and New York State - Age-adjusted	101
Table 46a. Ever diagnosed with diabetes among adults in Washington County and New York State	102
Table 46b. Ever diagnosed with diabetes among adults in Washington County and New York State - Age-adjusted	103
Table 47a. Ever diagnosed with asthma (lifetime) among adults in Washington County and New York State	104
Table 47b. Ever diagnosed with asthma (lifetime) among adults in Washington County and New York State - Age-adjusted	105
Table 48a. Current asthma among adults in Washington County and New York State	106
Table 48b. Current asthma among adults in Washington County and New York State - Age-adjusted	107
Table 49a. Chronic joint symptoms among adults in Washington County and New York State	108
Table 49b. Chronic joint symptoms among adults in Washington County and New York State - Age-adjusted	109
Table 50a. Arthritis among adults in Washington County and New York State	110
Table 50b. Arthritis among adults in Washington County and New York State - Age-adjusted	111
Table 51a. Activity limitations because of physical, mental or emotional problems among adults in Washington County and New York State	112
Table 51b. Activity limitations because of physical, mental or emotional problems among adults in Washington County and New York State - Age-adjusted	113
Table 52a. Have health problem that requires the use of special equipment among adults in Washington County and New York State	114
Table 52b. Have health problem that requires the use of special equipment among adults in Washington County and New York State - Age-adjusted	115



## Contents

Table 53a. Disability among adults in Washington County and New York State	116
Table 53b. Disability among adults in Washington County and New York State - Age-adjusted	117

### Health Behaviors

Table 54a. Current smoking among adults in Washington County and New York State	118
Table 54b. Current smoking among adults in Washington County and New York State - Age-adjusted	119
Table 55a. Everyday smoking among adults in Washington County and New York State	120
Table 55b. Everyday smoking among adults in Washington County and New York State - Age-adjusted	121
Table 56a. Adults living in homes in which smoking is prohibited in Washington County and New York State	122
Table 56b. Adults living in homes in which smoking is prohibited in Washington County and New York State - Age-adjusted	123
Table 57a. Binge drinking within the past month among adults in Washington County and New York State	124
Table 57b. Binge drinking within the past month among adults in Washington County and New York State - Age-adjusted	125
Table 58a. Heavy drinking within the past month among adults in Washington County and New York State	126
Table 58b. Heavy drinking within the past month among adults in Washington County and New York State - Age-adjusted	127
Table 59a. Never or rarely been asked about sexual history during routine check-up among adults in Washington County and New York State	128
Table 59b. Never or rarely been asked about sexual history during routine check-up among adults in Washington County and New York State - Age-adjusted	129
Table 60a. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults in Washington County and New York State	130

## Contents

Table 60b. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults in Washington County and New York State - Age-adjusted	131
Table 61a. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults in Washington County and New York State	132
Table 61b. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults in Washington County and New York State - Age-adjusted	133

### **Overweight / Obesity**

Table 62a. Overweight among adults in Washington County and New York State	134
Table 62b. Overweight among adults in Washington County and New York State - Age-adjusted	135
Table 63a. Obesity among adults in Washington County and New York State	136
Table 63b. Obesity among adults in Washington County and New York State - Age-adjusted	137
Table 64a. Overweight or obesity among adults in Washington County and New York State	138
Table 64b. Overweight or obesity among adults in Washington County and New York State - Age-adjusted	139
Table 65a. Received advice about weight by a health professional among adults in Washington County and New York State	140
Table 65b. Received advice about weight by a health professional among adults in Washington County and New York State - Age-adjusted	141
Table 66a. Received advice to lose weight by a health professional among adults provided weight advice in Washington County and New York State	142
Table 66b. Received advice to lose weight by a health professional among adults provided weight advice in Washington County and New York State - Age-adjusted	143

## Contents

### Physical Activity / Nutrition

Table 67a. No leisure-time physical activity among adults in Washington County and New York State	144
Table 67b. No leisure-time physical activity among adults in Washington County and New York State - Age-adjusted	145
Table 68a. Consumption of 5 or more servings of fruits and vegetables per day among adults in Washington County and New York State	146
Table 68b. Consumption of 5 or more servings of fruits and vegetables per day among adults in Washington County and New York State - Age-adjusted	147

### Injury

Table 69a. Had a fall within the past 3 months among adults in Washington County and New York State	148
Table 69b. Had a fall within the past 3 months among adults in Washington County and New York State - Age-adjusted	149
Table 70a. Had a fall that caused injury within the past 3 months among adults in Washington County and New York State	150
Table 70b. Had a fall that caused injury within the past 3 months among adults in Washington County and New York State - Age-adjusted	151

## Introduction

The New York State Expanded Behavioral Risk Factor Surveillance System (Expanded BRFSS) is a random digit dialed telephone survey of adults (18 years of age and older) that is representative of the non-institutionalized civilian population with landline telephones living in New York State. The goal of the Expanded BRFSS is to collect county-specific data on preventive health practices, risk behaviors, injuries and preventable chronic and infectious diseases. Topics assessed by the Expanded BRFSS include tobacco use, physical inactivity, diet, use of cancer screening services and other factors linked to the leading causes of morbidity and mortality. For each topic, health indicators are based on answers to one or more questions in the survey. The survey results are weighted to more accurately represent the adult population living in each county. Weighting procedures adjust for the differential selection probabilities by race, age and gender. More detailed information on weighting procedures is available in the technical brief.

Survey data collection occurred in 58 geographic areas of New York State, including each of the 57 counties outside of New York City and New York City (5 counties) as a single area. A standard questionnaire was used in all areas. Data collection for the 2009 Expanded BRFSS took place during the period of July 2008 through June 2009. The goal was to complete about 650 interviews in each area. The final number of interviews varies because of the sampling design of the survey and differential response rates. More detailed information on the sampling design is available in the technical brief.

The New York State Department of Health contracted with RTI International, which is responsible for the data collection, sampling, weighting and creation of the statistical reports. RTI International subcontracted with Clearwater Research Inc. for data collection and additional expertise in sample design and data quality control.

This report presents both crude estimates and estimates that have been age-adjusted to the Year 2000 US Standard Population when they represent the entire adult population of an area. Estimates for specific age groups are not age-adjusted. Age adjustment is a statistical process applied to rates that allows communities with different age structures to be compared. Counties can compare their data with results from New York State as a whole and with other counties. Confidence intervals at the 95% level associated with each health indicator give an estimated range of values which is likely to include the actual prevalence of the indicator. Confidence intervals are similar to margins of error. When the confidence intervals of two estimates of the same indicator from different areas do not overlap, they may be said to be statistically significantly different, i.e., these differences are unlikely related to chance and are considered true differences. If there is any value that is included in both intervals, the two estimates are not statistically significantly different.

The survey results for each of the 69 health indicators included in the survey are typically provided by gender, age, race and ethnicity, education and income in the Expanded BRFSS county reports. The estimates for some of these groupings may be suppressed in geographic areas with low proportions of adult respondents in these groupings.

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 1. Washington County and New York State adult US Census Bureau population estimates,<sup>1</sup> 2007**

	Washington County		New York State	
	Adult Population <sup>1</sup>	Percent	Adult Population <sup>1</sup>	Percent
<b>Total</b>	49,771	100	14,884,315	100
<b>Gender</b>				
Male	25,565	51.4	7,103,791	47.7
Female	24,206	48.6	7,780,524	52.3
<b>Age (years)</b>				
18-34	14,877	29.9	4,497,479	30.2
35-44	8,998	18.1	2,844,115	19.1
45-54	9,473	19.0	2,854,316	19.2
55-64	7,230	14.5	2,142,000	14.4
≥ 65	9,193	18.5	2,546,405	17.1
<b>Race/Ethnicity</b>				
White Non-Hispanic	46,377	93.2	9,343,884	62.8
Black Non-Hispanic	1,811	3.6	2,156,941	14.5
Hispanic	1,226	2.5	2,258,717	15.2
Other Non-Hispanic	357	0.7	1,124,773	7.6

<sup>1</sup> Source: National Center for Health Statistics. Postcensal estimates of the resident population of the United States for July 1, 2000-July 1, 2007, by year, county, age, bridged race, Hispanic origin, and sex (Vintage 2007). Prepared under a collaborative arrangement with the U.S. Census Bureau; released August 7, 2008. Available from: [http://www.cdc.gov/nchs/nvss/bridged\\_race.htm](http://www.cdc.gov/nchs/nvss/bridged_race.htm) as of November 25, 2009.

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 2a. Fair or poor general health among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	658	6,500	13.0	(9.9 - 16.1)	38,124	2,531,400	17.1	(15.1 - 19.0)
<b>Gender</b>								
Male	240	2,600	10.4	(5.9 - 14.9)	14,599	1,069,700	15.1	(12.4 - 17.8)
Female	418	3,800	15.8	(11.7 - 19.8)	23,525	1,461,700	18.8	(16.1 - 21.6)
<b>Age (years)</b>								
18-34	*	*	*	*	3,940	535,800	12.0	(7.2 - 16.8)
35-44	83	1,000	11.6	(4.6 - 18.6)	4,985	289,200	10.0	(6.8 - 13.3)
45-54	135	900	9.1	(4.6 - 13.6)	7,819	507,800	17.9	(14.0 - 21.8)
55-64	153	1,300	17.4	(10.9 - 23.9)	8,757	424,700	19.9	(16.0 - 23.9)
≥ 65	229	2,100	23.3	(17.5 - 29.0)	12,623	773,900	30.5	(26.8 - 34.2)
<b>Race/Ethnicity</b>								
White Non-Hispanic	636	5,700	12.2	(9.4 - 15.1)	35,811	1,226,600	13.2	(11.7 - 14.6)
Black Non-Hispanic	*	*	*	*	739	457,300	21.3	(14.8 - 27.9)
Hispanic	*	*	*	*	861	684,100	30.4	(22.2 - 38.6)
Other Non-Hispanic	*	*	*	*	713	163,400	14.5	(5.1 - 24.0)
<b>Education</b>								
≤ High School	312	4,500	19.8	(14.0 - 25.7)	15,682	1,466,500	26.3	(22.6 - 30.0)
Some College	161	1,500	11.6	(6.0 - 17.2)	9,904	508,500	14.3	(10.9 - 17.8)
≥ College Degree	183	500	3.5	(1.6 - 5.4)	12,457	493,500	8.8	(6.5 - 11.1)
<b>Income</b>								
≤ \$24,999	190	3,100	26.3	(17.3 - 35.3)	9,633	1,269,200	36.3	(30.1 - 42.4)
\$25,000-\$49,999	184	1,700	12.4	(7.3 - 17.6)	9,649	447,600	15.4	(11.9 - 18.9)
\$50,000-\$74,999	95	900	10.5	(1.6 - 19.3)	5,769	237,800	12.1	(8.0 - 16.2)
≥ \$75,000	*	*	*	*	8,562	218,300	4.9	(3.6 - 6.1)

<sup>1</sup> Defined as reporting fair or poor general health

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have fair or poor health (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 2b. Fair or poor general health among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	658	12.6	(9.4 - 15.7)	38,124	16.7	(14.8 - 18.6)
<b>Gender</b>						
Male	240	10.3	(5.9 - 14.7)	14,599	15.3	(12.6 - 18.0)
Female	418	14.9	(10.7 - 19.2)	23,525	18.0	(15.2 - 20.7)
<b>Race/Ethnicity</b>						
White Non-Hispanic	636	11.7	(8.8 - 14.6)	35,811	12.9	(11.4 - 14.4)
Black Non-Hispanic	*	*	*	739	20.8	(14.2 - 27.3)
Hispanic	*	*	*	861	29.9	(21.8 - 37.9)
Other Non-Hispanic	*	*	*	713	13.8	(4.8 - 22.8)
<b>Education</b>						
≤ High School	312	19.4	(13.5 - 25.4)	15,682	25.8	(22.1 - 29.5)
Some College	161	11.6	(6.0 - 17.3)	9,904	14.1	(10.7 - 17.6)
≥ College Degree	183	3.0	(1.3 - 4.7)	12,457	8.5	(6.3 - 10.8)
<b>Income</b>						
≤ \$24,999	190	26.9	(17.4 - 36.4)	9,633	35.9	(29.7 - 42.0)
\$25,000-\$49,999	184	12.4	(7.0 - 17.8)	9,649	15.0	(11.6 - 18.5)
\$50,000-\$74,999	95	9.5	(1.3 - 17.7)	5,769	11.8	(7.7 - 16.0)
≥ \$75,000	*	*	*	8,562	4.7	(3.5 - 6.0)

<sup>1</sup> Defined as reporting fair or poor general health

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 3a. Poor physical health 14 or more days within the past month among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	642	5,100	10.5	(7.7 - 13.2)	37,327	1,472,900	10.1	(8.7 - 11.6)
<b>Gender</b>								
Male	236	1,700	6.8	(3.4 - 10.1)	14,325	604,200	8.7	(7.0 - 10.5)
Female	406	3,400	14.4	(10.2 - 18.5)	23,002	868,600	11.4	(9.2 - 13.6)
<b>Age (years)</b>								
18-34	*	*	*	*	3,891	268,100	6.1	(2.7 - 9.5)
35-44	81	800	9.6	(3.4 - 15.8)	4,948	232,100	8.1	(5.2 - 11.1)
45-54	134	1,300	14.3	(7.1 - 21.5)	7,764	350,900	12.5	(9.4 - 15.6)
55-64	151	900	12.9	(7.4 - 18.4)	8,621	271,200	13.0	(10.3 - 15.8)
≥ 65	218	1,400	16.2	(11.0 - 21.4)	12,103	350,600	14.5	(12.1 - 16.9)
<b>Race/Ethnicity</b>								
White Non-Hispanic	620	4,400	9.7	(7.2 - 12.3)	35,073	929,700	10.2	(8.8 - 11.6)
Black Non-Hispanic	*	*	*	*	724	208,200	9.9	(4.8 - 14.9)
Hispanic	*	*	*	*	835	250,300	11.5	(6.5 - 16.4)
Other Non-Hispanic	*	*	*	*	695	84,600	7.6	(1.3 - 13.9)
<b>Education</b>								
≤ High School	303	3,300	15.0	(9.9 - 20.0)	15,178	692,800	12.9	(10.6 - 15.3)
Some College	156	1,000	7.9	(4.0 - 11.8)	9,751	362,500	10.3	(7.4 - 13.2)
≥ College Degree	181	800	5.7	(1.9 - 9.6)	12,327	359,300	6.4	(4.8 - 8.1)
<b>Income</b>								
≤ \$24,999	180	2,300	20.1	(12.1 - 28.0)	9,294	625,600	18.4	(13.6 - 23.3)
\$25,000-\$49,999	182	1,300	10.1	(4.8 - 15.4)	9,495	279,100	9.8	(7.3 - 12.3)
\$50,000-\$74,999	*	*	*	*	5,733	186,400	9.5	(5.8 - 13.2)
≥ \$75,000	*	*	*	*	8,510	219,100	4.9	(3.6 - 6.2)

<sup>1</sup> Defined as 14 or more days within the past month when physical health was not good

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults with poor physical health 14 or more days within the past month (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 3b. Poor physical health 14 or more days within the past month among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	642	10.2	(7.4 - 13.1)	37,327	9.9	(8.4 - 11.4)
<b>Gender</b>						
Male	236	6.7	(3.4 - 10.1)	14,325	8.7	(6.9 - 10.5)
Female	406	13.9	(9.5 - 18.3)	23,002	11.0	(8.7 - 13.2)
<b>Race/Ethnicity</b>						
White Non-Hispanic	620	9.4	(6.8 - 12.1)	35,073	9.9	(8.5 - 11.3)
Black Non-Hispanic	*	*	*	724	9.7	(4.6 - 14.9)
Hispanic	*	*	*	835	11.3	(6.3 - 16.2)
Other Non-Hispanic	*	*	*	695	7.6	(1.2 - 14.0)
<b>Education</b>						
≤ High School	303	14.9	(9.6 - 20.2)	15,178	12.7	(10.2 - 15.1)
Some College	156	7.7	(3.9 - 11.5)	9,751	10.1	(7.2 - 12.9)
≥ College Degree	181	5.6	(1.5 - 9.7)	12,327	6.3	(4.7 - 8.0)
<b>Income</b>						
≤ \$24,999	180	20.9	(12.3 - 29.6)	9,294	18.4	(13.5 - 23.4)
\$25,000-\$49,999	182	9.6	(4.4 - 14.8)	9,495	9.4	(7.0 - 11.9)
\$50,000-\$74,999	*	*	*	5,733	9.0	(5.3 - 12.6)
≥ \$75,000	*	*	*	8,510	4.9	(3.5 - 6.2)

<sup>1</sup> Defined as 14 or more days within the past month when physical health was not good

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 4a. Poor mental health 14 or more days within the past month among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	649	5,000	10.1	(7.0 - 13.2)	37,528	1,473,200	10.1	(8.6 - 11.5)
<b>Gender</b>								
Male	237	2,100	8.1	(4.0 - 12.1)	14,407	555,200	7.9	(6.3 - 9.6)
Female	412	2,900	12.2	(7.7 - 16.7)	23,121	918,000	12.0	(9.7 - 14.3)
<b>Age (years)</b>								
18-34	*	*	*	*	3,889	496,700	11.2	(7.4 - 15.1)
35-44	83	1,600	17.4	(7.9 - 27.0)	4,947	254,100	8.9	(6.7 - 11.1)
45-54	135	1,000	10.7	(5.5 - 15.9)	7,749	347,200	12.3	(9.0 - 15.6)
55-64	*	*	*	*	8,623	211,300	10.1	(7.9 - 12.3)
≥ 65	222	600	6.5	(2.9 - 10.1)	12,320	163,900	6.7	(4.7 - 8.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	627	4,600	10.0	(6.9 - 13.1)	35,266	899,200	9.8	(8.4 - 11.2)
Black Non-Hispanic	*	*	*	*	722	227,800	10.9	(5.4 - 16.4)
Hispanic	*	*	*	*	845	263,700	11.8	(6.8 - 16.8)
Other Non-Hispanic	*	*	*	*	695	82,500	7.4	(1.4 - 13.5)
<b>Education</b>								
≤ High School	306	2,600	11.5	(6.9 - 16.2)	15,335	665,000	12.2	(10.0 - 14.5)
Some College	158	1,500	11.6	(4.4 - 18.8)	9,771	404,500	11.5	(8.3 - 14.8)
≥ College Degree	183	900	6.1	(2.0 - 10.3)	12,342	346,600	6.2	(4.5 - 7.9)
<b>Income</b>								
≤ \$24,999	183	1,500	13.0	(6.3 - 19.7)	9,398	611,500	17.9	(13.0 - 22.8)
\$25,000-\$49,999	184	1,400	10.8	(5.7 - 15.9)	9,547	260,300	9.0	(6.7 - 11.4)
\$50,000-\$74,999	*	*	*	*	5,726	187,300	9.6	(6.0 - 13.1)
≥ \$75,000	*	*	*	*	8,520	276,500	6.2	(4.6 - 7.7)

<sup>1</sup> Defined as 14 or more days within the past month when mental health was not good

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults with poor mental health 14 or more days within the past month (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 4b. Poor mental health 14 or more days within the past month among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	649	10.9	(7.5 - 14.2)	37,528	10.2	(8.7 - 11.7)
<b>Gender</b>						
Male	237	8.5	(4.1 - 12.8)	14,407	7.9	(6.3 - 9.6)
Female	412	13.4	(8.2 - 18.5)	23,121	12.2	(9.8 - 14.6)
<b>Race/Ethnicity</b>						
White Non-Hispanic	627	10.8	(7.4 - 14.3)	35,266	10.0	(8.5 - 11.4)
Black Non-Hispanic	*	*	*	722	11.1	(5.5 - 16.7)
Hispanic	*	*	*	845	11.5	(6.5 - 16.4)
Other Non-Hispanic	*	*	*	695	7.6	(1.5 - 13.7)
<b>Education</b>						
≤ High School	306	12.7	(7.5 - 17.9)	15,335	12.3	(10.0 - 14.6)
Some College	158	12.7	(4.8 - 20.6)	9,771	11.8	(8.5 - 15.1)
≥ College Degree	183	6.1	(1.7 - 10.5)	12,342	6.3	(4.6 - 8.0)
<b>Income</b>						
≤ \$24,999	183	13.8	(6.7 - 21.0)	9,398	18.2	(13.2 - 23.1)
\$25,000-\$49,999	184	11.4	(5.8 - 17.0)	9,547	9.4	(6.9 - 11.9)
\$50,000-\$74,999	*	*	*	5,726	9.5	(5.9 - 13.1)
≥ \$75,000	*	*	*	8,520	6.2	(4.6 - 7.7)

<sup>1</sup> Defined as 14 or more days within the past month when mental health was not good

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 5a. Poor physical or mental health 14 or more days within the past month among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	637	8,500	17.5	(13.7 - 21.2)	37,005	2,385,900	16.5	(14.8 - 18.2)
<b>Gender</b>								
Male	234	3,100	12.5	(7.6 - 17.4)	14,231	937,700	13.6	(11.4 - 15.8)
Female	403	5,400	22.7	(17.4 - 28.1)	22,774	1,448,200	19.1	(16.5 - 21.7)
<b>Age (years)</b>								
18-34	*	*	*	*	3,859	602,600	13.8	(9.7 - 17.9)
35-44	*	*	*	*	4,924	388,300	13.7	(10.4 - 17.0)
45-54	134	1,800	19.3	(11.5 - 27.0)	7,722	551,000	19.6	(15.8 - 23.4)
55-64	149	1,200	16.7	(10.4 - 22.9)	8,545	395,500	19.2	(16.1 - 22.3)
≥ 65	215	1,800	20.3	(14.5 - 26.1)	11,955	448,500	18.8	(15.9 - 21.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	615	7,600	16.7	(13.1 - 20.3)	34,778	1,485,700	16.3	(14.8 - 17.8)
Black Non-Hispanic	*	*	*	*	715	343,700	16.5	(10.3 - 22.7)
Hispanic	*	*	*	*	832	412,900	19.0	(12.6 - 25.3)
Other Non-Hispanic	*	*	*	*	680	143,500	13.0	(5.0 - 21.0)
<b>Education</b>								
≤ High School	298	4,900	22.2	(15.9 - 28.5)	15,018	1,077,200	20.3	(17.3 - 23.2)
Some College	156	2,400	18.7	(10.8 - 26.7)	9,672	615,500	17.6	(14.1 - 21.1)
≥ College Degree	181	1,200	8.7	(4.2 - 13.2)	12,244	634,200	11.4	(9.2 - 13.6)
<b>Income</b>								
≤ \$24,999	177	3,000	26.8	(17.3 - 36.3)	9,199	924,000	27.3	(21.9 - 32.8)
\$25,000-\$49,999	182	2,400	17.9	(11.1 - 24.7)	9,425	456,000	16.1	(13.0 - 19.2)
\$50,000-\$74,999	95	1,000	11.5	(4.0 - 19.1)	5,696	312,000	16.0	(11.4 - 20.5)
≥ \$75,000	110	1,300	11.7	(3.9 - 19.5)	8,475	447,600	10.0	(8.1 - 12.0)

<sup>1</sup> Defined as 14 or more days within the past month when physical or mental health prevented usual activities, such as self-care, work or recreation

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults with poor physical or mental health 14 or more days within the past month (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 5b. Poor physical or mental health 14 or more days within the past month among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	637	17.9	(13.9 - 21.9)	37,005	16.4	(14.7 - 18.2)
<b>Gender</b>						
Male	234	12.8	(7.7 - 17.8)	14,231	13.6	(11.4 - 15.8)
Female	403	23.2	(17.3 - 29.1)	22,774	19.0	(16.3 - 21.6)
<b>Race/Ethnicity</b>						
White Non-Hispanic	615	17.2	(13.2 - 21.1)	34,778	16.3	(14.7 - 17.8)
Black Non-Hispanic	*	*	*	715	16.7	(10.3 - 23.0)
Hispanic	*	*	*	832	18.5	(12.2 - 24.8)
Other Non-Hispanic	*	*	*	680	13.1	(5.0 - 21.3)
<b>Education</b>						
≤ High School	298	23.0	(16.2 - 29.7)	15,018	20.1	(17.1 - 23.1)
Some College	156	19.6	(11.1 - 28.1)	9,672	17.6	(14.1 - 21.1)
≥ College Degree	181	8.4	(3.7 - 13.1)	12,244	11.4	(9.2 - 13.7)
<b>Income</b>						
≤ \$24,999	*	*	*	9,199	27.6	(22.1 - 33.1)
\$25,000-\$49,999	182	18.3	(11.2 - 25.4)	9,425	16.1	(12.9 - 19.3)
\$50,000-\$74,999	95	11.2	(3.3 - 19.2)	5,696	15.6	(11.0 - 20.1)
≥ \$75,000	110	12.3	(3.7 - 20.9)	8,475	10.0	(8.0 - 12.0)

<sup>1</sup> Defined as 14 or more days within the past month when physical or mental health prevented usual activities, such as self-care, work or recreation

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 6a. Health insurance among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	658	43,900	88.2	(83.8 - 92.6)	38,119	12,854,200	86.6	(84.2 - 89.0)
<b>Gender</b>								
Male	241	22,200	87.0	(79.8 - 94.3)	14,597	5,881,100	83.0	(78.9 - 87.1)
Female	417	21,600	89.5	(84.5 - 94.4)	23,522	6,973,100	89.9	(87.2 - 92.6)
<b>Age (years)</b>								
18-34	*	*	*	*	3,910	3,302,100	74.3	(67.8 - 80.7)
35-44	83	8,400	93.2	(87.5 - 98.9)	4,986	2,560,300	88.9	(84.7 - 93.1)
45-54	135	8,600	90.8	(84.6 - 96.9)	7,831	2,530,100	88.9	(84.6 - 93.2)
55-64	153	6,400	89.2	(84.2 - 94.2)	8,764	1,986,700	92.8	(90.2 - 95.5)
≥ 65	229	9,100	99.1	(97.9 - 100)	12,628	2,475,000	97.8	(96.7 - 98.8)
<b>Race/Ethnicity</b>								
White Non-Hispanic	636	41,600	89.9	(86.2 - 93.6)	35,811	8,591,200	92.2	(90.6 - 93.8)
Black Non-Hispanic	*	*	*	*	736	1,781,600	82.7	(75.3 - 90.1)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	312	19,800	88.1	(80.8 - 95.3)	15,673	4,448,900	79.7	(74.9 - 84.5)
Some College	161	10,900	84.1	(75.1 - 93.2)	9,906	3,004,600	85.0	(80.1 - 89.8)
≥ College Degree	183	13,000	92.5	(86.3 - 98.8)	12,459	5,361,700	95.3	(93.3 - 97.2)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,640	2,437,200	69.7	(62.5 - 76.8)
\$25,000-\$49,999	184	12,400	92.0	(85.8 - 98.3)	9,662	2,384,000	82.2	(76.1 - 88.3)
\$50,000-\$74,999	95	7,900	90.7	(81.6 - 99.9)	5,772	1,855,700	94.0	(91.2 - 96.7)
≥ \$75,000	112	10,300	94.3	(88.1 - 100)	8,567	4,388,400	97.5	(96.6 - 98.5)

<sup>1</sup> Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have health insurance (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 6b. Health insurance among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	658	88.1	(83.6 - 92.5)	38,119	86.7	(84.4 - 89.1)
<b>Gender</b>						
Male	241	87.7	(80.9 - 94.5)	14,597	83.6	(79.7 - 87.5)
Female	417	88.5	(82.8 - 94.2)	23,522	89.6	(86.8 - 92.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	636	89.5	(85.5 - 93.4)	35,811	92.1	(90.5 - 93.7)
Black Non-Hispanic	*	*	*	736	82.5	(75.0 - 90.0)
Hispanic	*	*	*	860	68.7	(59.0 - 78.4)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	312	87.8	(80.6 - 95.0)	15,673	80.1	(75.4 - 84.7)
Some College	161	83.4	(74.0 - 92.9)	9,906	84.6	(79.7 - 89.4)
≥ College Degree	183	92.9	(86.8 - 99.0)	12,459	95.4	(93.6 - 97.2)
<b>Income</b>						
≤ \$24,999	*	*	*	9,640	70.2	(63.2 - 77.1)
\$25,000-\$49,999	184	92.1	(85.9 - 98.3)	9,662	81.9	(76.0 - 87.9)
\$50,000-\$74,999	95	90.3	(80.4 - 100)	5,772	94.0	(91.3 - 96.7)
≥ \$75,000	112	93.6	(86.6 - 100)	8,567	97.5	(96.6 - 98.5)

<sup>1</sup> Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 7a. Health insurance among adults aged 18-64<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	426	34,700	85.8	(80.5 - 91.2)	25,165	10,251,300	84.2	(81.3 - 87.1)
<b>Gender</b>								
Male	158	18,200	84.8	(76.3 - 93.4)	10,078	4,839,900	80.3	(75.5 - 85.0)
Female	268	16,500	87.0	(80.8 - 93.2)	15,087	5,411,400	88.0	(84.7 - 91.3)
<b>Age (years)</b>								
18-34	*	*	*	*	3,910	3,302,100	74.3	(67.8 - 80.7)
35-44	83	8,400	93.2	(87.5 - 98.9)	4,981	2,556,600	88.9	(84.7 - 93.1)
45-54	135	8,600	90.8	(84.6 - 96.9)	7,791	2,472,900	88.7	(84.3 - 93.0)
55-64	150	6,400	89.8	(84.9 - 94.7)	8,483	1,919,700	92.8	(90.1 - 95.5)
≥ 65	-	-	-	-	-	-	-	-
<b>Race/Ethnicity</b>								
White Non-Hispanic	407	32,700	87.8	(83.2 - 92.3)	23,354	6,582,200	90.3	(88.3 - 92.4)
Black Non-Hispanic	*	*	*	*	567	1,464,100	80.2	(71.6 - 88.7)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	191	15,100	85.3	(76.3 - 94.3)	9,037	3,191,500	74.6	(68.6 - 80.6)
Some College	*	*	*	*	7,031	2,540,400	82.9	(77.4 - 88.4)
≥ College Degree	125	10,600	90.9	(83.4 - 98.5)	9,068	4,502,400	94.5	(92.2 - 96.8)
<b>Income</b>								
≤ \$24,999	*	*	*	*	4,918	1,707,200	62.6	(54.1 - 71.2)
\$25,000-\$49,999	121	9,600	90.0	(82.2 - 97.8)	6,180	1,783,000	77.7	(70.3 - 85.1)
\$50,000-\$74,999	80	7,300	90.1	(80.3 - 99.8)	4,554	1,553,400	92.9	(89.7 - 96.1)
≥ \$75,000	95	9,500	93.9	(87.1 - 100)	7,324	3,987,600	97.3	(96.3 - 98.3)

<sup>1</sup> Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 18-64 who have health insurance (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 7b. Health insurance among adults aged 18-64<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	426	85.9	(80.5 - 91.2)	25,165	84.3	(81.5 - 87.2)
<b>Gender</b>						
Male	158	85.3	(77.1 - 93.5)	10,078	80.5	(75.9 - 85.1)
Female	268	86.5	(79.7 - 93.2)	15,087	87.9	(84.6 - 91.2)
<b>Race/Ethnicity</b>						
White Non-Hispanic	407	87.4	(82.7 - 92.2)	23,354	90.3	(88.3 - 92.3)
Black Non-Hispanic	*	*	*	567	80.0	(71.3 - 88.6)
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	191	85.2	(76.3 - 94.0)	9,037	75.1	(69.3 - 80.9)
Some College	*	*	*	7,031	82.6	(77.1 - 88.1)
≥ College Degree	125	91.5	(84.2 - 98.8)	9,068	94.6	(92.5 - 96.8)
<b>Income</b>						
≤ \$24,999	*	*	*	4,918	63.2	(54.9 - 71.6)
\$25,000-\$49,999	121	90.1	(82.5 - 97.8)	6,180	77.5	(70.2 - 84.7)
\$50,000-\$74,999	*	*	*	4,554	93.0	(89.8 - 96.2)
≥ \$75,000	95	93.1	(85.5 - 100)	7,324	97.3	(96.3 - 98.3)

<sup>1</sup> Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 8a. No health insurance among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	658	5,900	11.8	(7.4 - 16.2)	38,119	1,989,300	13.4	(11.0 - 15.8)
<b>Gender</b>								
Male	241	3,300	13.0	(5.7 - 20.2)	14,597	1,207,400	17.0	(12.9 - 21.1)
Female	417	2,500	10.5	(5.6 - 15.5)	23,522	781,900	10.1	(7.4 - 12.8)
<b>Age (years)</b>								
18-34	*	*	*	*	3,910	1,144,100	25.7	(19.3 - 32.2)
35-44	*	*	*	*	4,986	319,200	11.1	(6.9 - 15.3)
45-54	135	900	9.2	(3.1 - 15.4)	7,831	316,500	11.1	(6.8 - 15.4)
55-64	153	800	10.8	(5.8 - 15.8)	8,764	153,300	7.2	(4.5 - 9.8)
≥ 65	*	*	*	*	12,628	56,200	2.2	(1.2 - 3.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	636	4,700	10.1	(6.4 - 13.8)	35,811	727,100	7.8	(6.2 - 9.4)
Black Non-Hispanic	*	*	*	*	736	372,800	17.3	(9.9 - 24.7)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	312	2,700	11.9	(4.7 - 19.2)	15,673	1,134,400	20.3	(15.5 - 25.1)
Some College	161	2,100	15.9	(6.8 - 24.9)	9,906	531,400	15.0	(10.2 - 19.9)
≥ College Degree	183	1,100	7.5	(1.2 - 13.7)	12,459	266,300	4.7	(2.8 - 6.7)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,640	1,061,200	30.3	(23.2 - 37.5)
\$25,000-\$49,999	184	1,100	8.0	(1.7 - 14.2)	9,662	516,700	17.8	(11.7 - 23.9)
\$50,000-\$74,999	*	*	*	*	5,772	119,100	6.0	(3.3 - 8.8)
≥ \$75,000	*	*	*	*	8,567	111,100	2.5	(1.5 - 3.4)

<sup>1</sup> Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have no health insurance (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 8b. No health insurance among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	658	11.9	(7.5 - 16.4)	38,119	13.3	(10.9 - 15.6)
<b>Gender</b>						
Male	241	12.3	(5.5 - 19.1)	14,597	16.4	(12.5 - 20.3)
Female	417	11.5	(5.8 - 17.2)	23,522	10.4	(7.7 - 13.2)
<b>Race/Ethnicity</b>						
White Non-Hispanic	636	10.5	(6.6 - 14.5)	35,811	7.9	(6.3 - 9.5)
Black Non-Hispanic	*	*	*	736	17.5	(10.0 - 25.0)
Hispanic	*	*	*	860	31.3	(21.6 - 41.0)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	312	12.2	(5.0 - 19.4)	15,673	19.9	(15.3 - 24.6)
Some College	161	16.6	(7.1 - 26.0)	9,906	15.4	(10.6 - 20.3)
≥ College Degree	183	7.1	(1.0 - 13.2)	12,459	4.6	(2.8 - 6.4)
<b>Income</b>						
≤ \$24,999	*	*	*	9,640	29.8	(22.9 - 36.8)
\$25,000-\$49,999	184	7.9	(1.7 - 14.1)	9,662	18.1	(12.1 - 24.0)
\$50,000-\$74,999	*	*	*	5,772	6.0	(3.3 - 8.7)
≥ \$75,000	*	*	*	8,567	2.5	(1.5 - 3.4)

<sup>1</sup> Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 9a. No health insurance among adults aged 18-64<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	426	5,700	14.2	(8.8 - 19.5)	25,165	1,928,400	15.8	(12.9 - 18.7)
<b>Gender</b>								
Male	158	3,300	15.2	(6.6 - 23.7)	10,078	1,190,200	19.7	(15.0 - 24.5)
Female	268	2,500	13.0	(6.8 - 19.2)	15,087	738,200	12.0	(8.7 - 15.3)
<b>Age (years)</b>								
18-34	*	*	*	*	3,910	1,144,100	25.7	(19.3 - 32.2)
35-44	*	*	*	*	4,981	319,200	11.1	(6.9 - 15.3)
45-54	135	900	9.2	(3.1 - 15.4)	7,791	316,100	11.3	(7.0 - 15.7)
55-64	150	700	10.2	(5.3 - 15.1)	8,483	148,900	7.2	(4.5 - 9.9)
≥ 65	-	-	-	-	-	-	-	-
<b>Race/Ethnicity</b>								
White Non-Hispanic	407	4,500	12.2	(7.7 - 16.8)	23,354	703,100	9.7	(7.6 - 11.7)
Black Non-Hispanic	*	*	*	*	567	361,800	19.8	(11.3 - 28.4)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	191	2,600	14.7	(5.7 - 23.7)	9,037	1,087,700	25.4	(19.4 - 31.4)
Some College	*	*	*	*	7,031	522,800	17.1	(11.6 - 22.6)
≥ College Degree	125	1,100	9.1	(1.5 - 16.6)	9,068	261,200	5.5	(3.2 - 7.8)
<b>Income</b>								
≤ \$24,999	*	*	*	*	4,918	1,018,000	37.4	(28.8 - 45.9)
\$25,000-\$49,999	121	1,100	10.0	(2.2 - 17.8)	6,180	511,100	22.3	(14.9 - 29.7)
\$50,000-\$74,999	*	*	*	*	4,554	118,800	7.1	(3.9 - 10.3)
≥ \$75,000	*	*	*	*	7,324	110,700	2.7	(1.7 - 3.7)

<sup>1</sup> Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 18-64 who have no health insurance (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 9b. No health insurance among adults aged 18-64<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	426	14.1	(8.8 - 19.5)	25,165	15.7	(12.8 - 18.5)
<b>Gender</b>						
Male	158	14.7	(6.5 - 22.9)	10,078	19.5	(14.9 - 24.1)
Female	268	13.5	(6.8 - 20.3)	15,087	12.1	(8.8 - 15.4)
<b>Race/Ethnicity</b>						
White Non-Hispanic	407	12.6	(7.8 - 17.3)	23,354	9.7	(7.7 - 11.7)
Black Non-Hispanic	*	*	*	567	20.0	(11.4 - 28.7)
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	191	14.8	(6.0 - 23.7)	9,037	24.9	(19.1 - 30.7)
Some College	*	*	*	7,031	17.4	(11.9 - 22.9)
≥ College Degree	125	8.5	(1.2 - 15.8)	9,068	5.4	(3.2 - 7.5)
<b>Income</b>						
≤ \$24,999	*	*	*	4,918	36.8	(28.4 - 45.1)
\$25,000-\$49,999	121	9.9	(2.2 - 17.5)	6,180	22.5	(15.3 - 29.8)
\$50,000-\$74,999	*	*	*	4,554	7.0	(3.8 - 10.2)
≥ \$75,000	*	*	*	7,324	2.7	(1.7 - 3.7)

<sup>1</sup> Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 10a. Adults with regular health care provider<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	657	40,900	82.3	(77.2 - 87.3)	38,135	12,324,300	83.0	(80.4 - 85.6)
<b>Gender</b>								
Male	239	20,100	78.9	(70.4 - 87.5)	14,593	5,494,300	77.6	(73.2 - 82.0)
Female	418	20,800	85.8	(80.7 - 90.8)	23,542	6,830,000	87.8	(85.1 - 90.5)
<b>Age (years)</b>								
18-34	*	*	*	*	3,929	2,931,600	65.8	(59.2 - 72.5)
35-44	83	7,300	81.7	(71.8 - 91.6)	4,988	2,471,000	86.1	(81.9 - 90.3)
45-54	134	8,300	88.3	(81.7 - 94.9)	7,823	2,529,300	88.6	(84.6 - 92.7)
55-64	153	6,300	87.6	(81.8 - 93.3)	8,770	2,002,000	93.6	(91.7 - 95.4)
≥ 65	229	8,400	92.3	(88.7 - 95.8)	12,625	2,390,400	94.1	(92.2 - 96.0)
<b>Race/Ethnicity</b>								
White Non-Hispanic	635	38,400	83.0	(78.3 - 87.7)	35,821	8,227,700	88.2	(86.4 - 89.9)
Black Non-Hispanic	*	*	*	*	740	1,753,200	81.3	(72.6 - 89.9)
Hispanic	*	*	*	*	860	1,453,100	64.7	(54.7 - 74.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	312	19,200	85.1	(77.8 - 92.4)	15,692	4,249,200	75.9	(70.9 - 80.8)
Some College	*	*	*	*	9,904	2,935,000	82.8	(77.2 - 88.4)
≥ College Degree	183	12,100	86.2	(78.7 - 93.6)	12,456	5,046,100	89.8	(87.1 - 92.6)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,644	2,554,200	72.7	(65.9 - 79.6)
\$25,000-\$49,999	*	*	*	*	9,655	2,243,100	77.3	(70.0 - 84.6)
\$50,000-\$74,999	*	*	*	*	5,771	1,767,500	89.5	(85.2 - 93.9)
≥ \$75,000	112	9,400	85.9	(77.4 - 94.3)	8,560	4,122,800	91.9	(90.2 - 93.7)

<sup>1</sup> Defined as having one or more persons respondent thinks of as a personal doctor or health care provider

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have at least one person they think of as their personal doctor or health care provider (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 10b. Adults with regular health care provider<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	657	81.8	(76.7 - 86.9)	38,135	83.0	(80.4 - 85.5)
<b>Gender</b>						
Male	239	79.1	(70.7 - 87.4)	14,593	78.1	(73.9 - 82.3)
Female	418	84.7	(79.0 - 90.5)	23,542	87.4	(84.6 - 90.1)
<b>Race/Ethnicity</b>						
White Non-Hispanic	635	82.3	(77.4 - 87.3)	35,821	88.0	(86.2 - 89.7)
Black Non-Hispanic	*	*	*	740	81.4	(73.0 - 89.8)
Hispanic	*	*	*	860	65.2	(55.5 - 75.0)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	312	84.3	(77.0 - 91.7)	15,692	75.8	(71.0 - 80.6)
Some College	*	*	*	9,904	82.9	(77.5 - 88.3)
≥ College Degree	183	86.1	(78.5 - 93.7)	12,456	89.8	(87.1 - 92.4)
<b>Income</b>						
≤ \$24,999	*	*	*	9,644	72.9	(66.1 - 79.6)
\$25,000-\$49,999	*	*	*	9,655	77.4	(70.5 - 84.3)
\$50,000-\$74,999	*	*	*	5,771	89.6	(85.3 - 94.0)
≥ \$75,000	112	84.9	(75.5 - 94.2)	8,560	91.6	(89.8 - 93.4)

<sup>1</sup> Defined as having one or more persons respondent thinks of as a personal doctor or health care provider

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 11a. Cost prevented visit to doctor within the past year among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	659	6,300	12.6	(8.9 - 16.3)	38,143	2,034,800	13.7	(11.6 - 15.8)
<b>Gender</b>								
Male	241	2,100	8.2	(3.6 - 12.8)	14,607	903,300	12.7	(9.5 - 16.0)
Female	418	4,200	17.3	(11.7 - 22.8)	23,536	1,131,500	14.6	(11.8 - 17.4)
<b>Age (years)</b>								
18-34	*	*	*	*	3,939	947,300	21.2	(15.6 - 26.9)
35-44	83	1,200	13.0	(4.7 - 21.3)	4,987	347,900	12.1	(8.0 - 16.2)
45-54	135	800	8.2	(2.7 - 13.8)	7,825	382,200	13.5	(9.9 - 17.1)
55-64	153	900	12.5	(7.5 - 17.5)	8,766	236,800	11.1	(8.0 - 14.2)
≥ 65	*	*	*	*	12,626	120,600	4.8	(2.5 - 7.0)
<b>Race/Ethnicity</b>								
White Non-Hispanic	637	5,300	11.5	(7.9 - 15.1)	35,833	984,600	10.6	(8.9 - 12.2)
Black Non-Hispanic	*	*	*	*	738	280,000	13.1	(7.2 - 19.0)
Hispanic	*	*	*	*	859	597,900	26.5	(17.3 - 35.7)
Other Non-Hispanic	*	*	*	*	713	172,300	15.3	(5.7 - 24.9)
<b>Education</b>								
≤ High School	313	3,800	16.9	(10.6 - 23.3)	15,696	996,600	17.8	(13.8 - 21.9)
Some College	161	1,600	12.6	(5.3 - 19.8)	9,911	512,400	14.5	(10.4 - 18.6)
≥ College Degree	183	800	5.8	(1.4 - 10.3)	12,453	466,900	8.3	(6.0 - 10.6)
<b>Income</b>								
≤ \$24,999	191	2,100	17.3	(9.4 - 25.3)	9,639	854,500	24.3	(18.4 - 30.3)
\$25,000-\$49,999	184	1,100	8.0	(3.4 - 12.7)	9,656	532,900	18.4	(12.5 - 24.3)
\$50,000-\$74,999	*	*	*	*	5,776	202,700	10.3	(6.0 - 14.5)
≥ \$75,000	*	*	*	*	8,561	192,000	4.3	(2.9 - 5.6)

<sup>1</sup> Defined as needing to see a doctor within the past year but not being able to because of cost

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who did not see a doctor because of cost within the past year (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 11b. Cost prevented visit to doctor within the past year among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	659	13.4	(9.4 - 17.4)	38,143	13.8	(11.7 - 15.9)
<b>Gender</b>						
Male	241	8.1	(3.5 - 12.7)	14,607	12.5	(9.4 - 15.6)
Female	418	19.0	(12.7 - 25.3)	23,536	15.0	(12.1 - 17.9)
<b>Race/Ethnicity</b>						
White Non-Hispanic	637	12.2	(8.3 - 16.2)	35,833	10.8	(9.2 - 12.5)
Black Non-Hispanic	*	*	*	738	13.2	(7.2 - 19.3)
Hispanic	*	*	*	859	26.1	(17.2 - 35.1)
Other Non-Hispanic	*	*	*	713	14.8	(5.5 - 24.2)
<b>Education</b>						
≤ High School	313	18.1	(11.2 - 25.1)	15,696	17.8	(13.8 - 21.7)
Some College	161	13.5	(5.8 - 21.3)	9,911	15.3	(11.0 - 19.5)
≥ College Degree	183	6.2	(1.2 - 11.1)	12,453	8.2	(6.0 - 10.5)
<b>Income</b>						
≤ \$24,999	191	18.0	(9.5 - 26.5)	9,639	24.5	(18.6 - 30.4)
\$25,000-\$49,999	184	8.7	(3.4 - 14.0)	9,656	18.7	(12.9 - 24.6)
\$50,000-\$74,999	*	*	*	5,776	10.4	(6.1 - 14.7)
≥ \$75,000	*	*	*	8,561	4.4	(3.1 - 5.8)

<sup>1</sup> Defined as needing to see a doctor within the past year but not being able to because of cost

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 12a. Visited a doctor for routine checkup within the past year among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	657	32,100	64.7	(58.8 - 70.5)	38,026	10,809,300	72.9	(70.6 - 75.3)
<b>Gender</b>								
Male	241	15,100	59.1	(49.4 - 68.9)	14,592	4,738,400	66.9	(63.0 - 70.9)
Female	416	17,000	70.6	(64.6 - 76.5)	23,434	6,070,900	78.5	(75.9 - 81.0)
<b>Age (years)</b>								
18-34	*	*	*	*	3,921	2,747,400	61.7	(55.6 - 67.9)
35-44	*	*	*	*	4,973	1,883,100	65.5	(60.2 - 70.7)
45-54	135	5,300	56.0	(46.1 - 65.9)	7,808	2,174,100	76.6	(73.2 - 80.0)
55-64	153	5,300	73.5	(65.6 - 81.4)	8,751	1,728,600	81.4	(78.7 - 84.2)
≥ 65	228	8,000	88.1	(83.9 - 92.4)	12,573	2,276,000	90.0	(87.5 - 92.5)
<b>Race/Ethnicity</b>								
White Non-Hispanic	636	30,200	65.2	(59.4 - 70.9)	35,721	6,746,200	72.4	(70.6 - 74.2)
Black Non-Hispanic	*	*	*	*	736	1,688,900	78.8	(70.4 - 87.3)
Hispanic	*	*	*	*	857	1,550,900	69.3	(60.0 - 78.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	313	14,100	62.3	(53.2 - 71.3)	15,640	4,150,600	74.7	(70.7 - 78.7)
Some College	*	*	*	*	9,880	2,524,000	71.3	(66.1 - 76.5)
≥ College Degree	182	9,900	70.6	(61.3 - 79.8)	12,428	4,042,900	71.9	(68.5 - 75.3)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,616	2,469,200	70.4	(64.2 - 76.6)
\$25,000-\$49,999	*	*	*	*	9,631	2,068,400	71.3	(65.3 - 77.3)
\$50,000-\$74,999	*	*	*	*	5,765	1,466,000	74.8	(69.6 - 80.0)
≥ \$75,000	*	*	*	*	8,547	3,176,500	70.7	(67.3 - 74.2)

<sup>1</sup> Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past year

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have visited a doctor for a routine checkup within the past year (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 12b. Visited a doctor for routine checkup within the past year among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	657	64.2	(58.3 - 70.1)	38,026	72.7	(70.3 - 75.0)
<b>Gender</b>						
Male	241	60.2	(50.6 - 69.8)	14,592	67.1	(63.3 - 70.9)
Female	416	68.4	(61.8 - 75.1)	23,434	77.7	(75.1 - 80.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	636	64.3	(58.5 - 70.2)	35,721	71.9	(70.1 - 73.7)
Black Non-Hispanic	*	*	*	736	79.3	(71.3 - 87.4)
Hispanic	*	*	*	857	69.2	(60.1 - 78.2)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	313	61.3	(52.2 - 70.5)	15,640	74.4	(70.4 - 78.4)
Some College	*	*	*	9,880	70.9	(65.9 - 76.0)
≥ College Degree	182	70.6	(61.2 - 80.0)	12,428	71.7	(68.3 - 75.0)
<b>Income</b>						
≤ \$24,999	*	*	*	9,616	70.2	(64.1 - 76.2)
\$25,000-\$49,999	*	*	*	9,631	71.1	(65.3 - 76.9)
\$50,000-\$74,999	*	*	*	5,765	74.5	(69.2 - 79.8)
≥ \$75,000	*	*	*	8,547	70.3	(66.9 - 73.8)

<sup>1</sup> Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past year

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 13a. Visited a doctor for routine checkup within the past 2 years among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	657	39,800	80.1	(75.1 - 85.1)	38,026	12,726,600	85.9	(83.9 - 87.9)
<b>Gender</b>								
Male	241	19,200	75.2	(66.6 - 83.8)	14,592	5,791,900	81.8	(78.2 - 85.4)
Female	416	20,600	85.3	(80.6 - 89.9)	23,434	6,934,700	89.6	(87.9 - 91.3)
<b>Age (years)</b>								
18-34	*	*	*	*	3,921	3,535,400	79.4	(73.9 - 85.0)
35-44	*	*	*	*	4,973	2,355,600	81.9	(78.0 - 85.7)
45-54	135	7,400	78.3	(69.7 - 86.9)	7,808	2,507,900	88.4	(85.6 - 91.2)
55-64	153	6,000	83.6	(77.1 - 90.1)	8,751	1,924,800	90.7	(88.6 - 92.7)
≥ 65	228	8,600	93.8	(90.8 - 96.9)	12,573	2,402,900	95.0	(93.5 - 96.5)
<b>Race/Ethnicity</b>								
White Non-Hispanic	636	37,600	81.1	(76.5 - 85.6)	35,721	8,070,000	86.6	(85.4 - 87.9)
Black Non-Hispanic	*	*	*	*	736	1,914,500	89.4	(82.3 - 96.5)
Hispanic	*	*	*	*	857	1,832,800	81.9	(73.7 - 90.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	313	17,800	78.7	(70.8 - 86.7)	15,640	4,713,700	84.8	(81.2 - 88.4)
Some College	*	*	*	*	9,880	2,990,500	84.5	(79.6 - 89.4)
≥ College Degree	182	11,700	82.9	(75.2 - 90.6)	12,428	4,928,300	87.6	(85.3 - 89.9)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,616	2,911,700	83.0	(77.4 - 88.6)
\$25,000-\$49,999	184	10,800	80.2	(70.7 - 89.7)	9,631	2,375,500	81.9	(76.1 - 87.7)
\$50,000-\$74,999	*	*	*	*	5,765	1,733,500	88.4	(84.9 - 91.9)
≥ \$75,000	112	9,200	84.8	(76.3 - 93.4)	8,547	3,907,000	87.0	(84.8 - 89.2)

<sup>1</sup> Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past 2 years

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have visited a doctor for a routine checkup within the past 2 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 13b. Visited a doctor for routine checkup within the past 2 years among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	657	79.8	(74.8 - 84.8)	38,026	85.7	(83.7 - 87.6)
<b>Gender</b>						
Male	241	75.8	(67.4 - 84.2)	14,592	81.9	(78.4 - 85.3)
Female	416	84.1	(78.8 - 89.3)	23,434	89.1	(87.4 - 90.9)
<b>Race/Ethnicity</b>						
White Non-Hispanic	636	80.5	(75.8 - 85.3)	35,721	86.2	(85.0 - 87.5)
Black Non-Hispanic	*	*	*	736	89.6	(83.0 - 96.3)
Hispanic	*	*	*	857	81.8	(73.8 - 89.8)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	313	77.7	(69.7 - 85.8)	15,640	84.6	(81.1 - 88.1)
Some College	*	*	*	9,880	84.4	(79.8 - 89.0)
≥ College Degree	182	83.0	(75.2 - 90.8)	12,428	87.3	(85.0 - 89.6)
<b>Income</b>						
≤ \$24,999	*	*	*	9,616	83.2	(77.8 - 88.5)
\$25,000-\$49,999	184	80.3	(70.9 - 89.8)	9,631	81.9	(76.3 - 87.4)
\$50,000-\$74,999	*	*	*	5,765	88.1	(84.4 - 91.8)
≥ \$75,000	112	84.3	(75.2 - 93.4)	8,547	86.4	(84.2 - 88.7)

<sup>1</sup> Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past 2 years

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 14a. Dentist visit within the past year among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	654	29,900	60.5	(54.5 - 66.4)	38,026	10,540,400	71.1	(68.6 - 73.6)
<b>Gender</b>								
Male	241	13,800	54.2	(44.4 - 64.0)	14,573	4,845,800	68.4	(64.5 - 72.2)
Female	413	16,100	67.2	(61.2 - 73.2)	23,453	5,694,600	73.6	(70.4 - 76.8)
<b>Age (years)</b>								
18-34	*	*	*	*	3,924	2,789,700	62.6	(56.3 - 69.0)
35-44	*	*	*	*	4,979	2,181,000	75.8	(70.9 - 80.7)
45-54	134	6,200	66.0	(56.8 - 75.1)	7,814	2,149,300	75.8	(71.9 - 79.7)
55-64	153	4,900	67.4	(59.4 - 75.4)	8,742	1,690,500	79.2	(75.9 - 82.6)
≥ 65	226	5,500	61.6	(54.8 - 68.5)	12,567	1,729,800	68.5	(64.7 - 72.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	633	27,900	60.5	(54.6 - 66.5)	35,721	6,959,900	74.7	(72.7 - 76.6)
Black Non-Hispanic	*	*	*	*	735	1,405,800	66.2	(57.9 - 74.5)
Hispanic	*	*	*	*	862	1,425,600	63.1	(53.7 - 72.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	310	12,000	53.3	(44.4 - 62.3)	15,613	3,620,700	65.1	(60.7 - 69.5)
Some College	*	*	*	*	9,880	2,504,700	70.8	(66.1 - 75.5)
≥ College Degree	*	*	*	*	12,453	4,402,400	78.2	(74.6 - 81.8)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,594	1,972,200	56.2	(49.8 - 62.6)
\$25,000-\$49,999	*	*	*	*	9,649	1,896,000	65.8	(59.5 - 72.1)
\$50,000-\$74,999	*	*	*	*	5,763	1,482,400	75.2	(69.4 - 81.0)
≥ \$75,000	*	*	*	*	8,564	3,707,900	82.4	(79.5 - 85.3)

<sup>1</sup> Defined as having visited a dentist or a dental clinic for any reason within the past year

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have visited a dentist or a dental clinic for any reason within the past year (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 14b. Dentist visit within the past year among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	654	61.0	(55.0 - 67.1)	38,026	71.1	(68.6 - 73.5)
<b>Gender</b>						
Male	241	55.3	(45.5 - 65.0)	14,573	68.3	(64.6 - 72.1)
Female	413	67.2	(60.6 - 73.8)	23,453	73.6	(70.4 - 76.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	633	61.0	(54.9 - 67.0)	35,721	74.4	(72.4 - 76.3)
Black Non-Hispanic	*	*	*	735	66.6	(58.3 - 74.9)
Hispanic	*	*	*	862	63.6	(54.4 - 72.9)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	310	53.8	(44.7 - 63.0)	15,613	65.3	(60.9 - 69.6)
Some College	*	*	*	9,880	70.4	(65.7 - 75.1)
≥ College Degree	*	*	*	12,453	78.1	(74.6 - 81.7)
<b>Income</b>						
≤ \$24,999	*	*	*	9,594	56.4	(50.1 - 62.8)
\$25,000-\$49,999	*	*	*	9,649	65.8	(59.6 - 72.0)
\$50,000-\$74,999	*	*	*	5,763	75.0	(69.1 - 80.8)
≥ \$75,000	*	*	*	8,564	82.1	(79.1 - 85.0)

<sup>1</sup> Defined as having visited a dentist or a dental clinic for any reason within the past year

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 15a. Had teeth cleaned within the past year among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	581	28,800	61.8	(55.6 - 68.1)	34,070	9,985,900	71.7	(69.1 - 74.3)
<b>Gender</b>								
Male	*	*	*	*	13,173	4,616,400	69.0	(65.0 - 73.1)
Female	368	15,200	68.3	(62.0 - 74.7)	20,897	5,369,500	74.2	(70.8 - 77.5)
<b>Age (years)</b>								
18-34	*	*	*	*	3,860	2,746,400	63.0	(56.5 - 69.5)
35-44	*	*	*	*	4,875	2,112,100	75.5	(70.7 - 80.3)
45-54	130	6,000	65.1	(55.7 - 74.4)	7,438	2,022,600	73.3	(69.1 - 77.4)
55-64	141	4,500	66.9	(58.7 - 75.2)	7,906	1,560,900	77.7	(73.8 - 81.6)
≥ 65	170	4,900	70.7	(63.4 - 78.0)	9,991	1,543,900	77.1	(73.7 - 80.5)
<b>Race/Ethnicity</b>								
White Non-Hispanic	560	27,600	64.0	(57.8 - 70.2)	31,982	6,601,500	75.4	(73.4 - 77.4)
Black Non-Hispanic	*	*	*	*	680	1,227,200	61.6	(52.1 - 71.0)
Hispanic	*	*	*	*	789	1,421,300	66.6	(57.1 - 76.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	253	11,300	56.3	(46.4 - 66.2)	12,740	3,388,600	67.7	(63.1 - 72.3)
Some College	*	*	*	*	9,145	2,286,800	68.5	(62.9 - 74.1)
≥ College Degree	*	*	*	*	12,128	4,298,500	78.3	(74.8 - 81.7)
<b>Income</b>								
≤ \$24,999	*	*	*	*	7,525	1,861,800	60.3	(53.5 - 67.1)
\$25,000-\$49,999	*	*	*	*	8,709	1,681,700	62.6	(55.4 - 69.8)
\$50,000-\$74,999	*	*	*	*	5,534	1,447,800	76.3	(70.7 - 81.8)
≥ \$75,000	*	*	*	*	8,448	3,617,500	80.8	(77.8 - 83.8)

<sup>1</sup> Defined as having had teeth cleaned by a dentist or dental hygienist within the past year

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have had their teeth cleaned by a dentist or dental hygienist within the past year (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 15b. Had teeth cleaned within the past year among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	581	62.3	(56.0 - 68.5)	34,070	71.7	(69.1 - 74.2)
<b>Gender</b>						
Male	*	*	*	13,173	69.2	(65.3 - 73.1)
Female	368	67.7	(60.7 - 74.6)	20,897	73.9	(70.5 - 77.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	560	64.3	(58.0 - 70.5)	31,982	75.1	(73.0 - 77.1)
Black Non-Hispanic	*	*	*	680	62.0	(52.7 - 71.4)
Hispanic	*	*	*	789	66.9	(57.5 - 76.2)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	253	56.2	(46.2 - 66.1)	12,740	67.7	(63.2 - 72.3)
Some College	*	*	*	9,145	68.1	(62.6 - 73.6)
≥ College Degree	*	*	*	12,128	78.3	(74.9 - 81.7)
<b>Income</b>						
≤ \$24,999	*	*	*	7,525	60.2	(53.4 - 67.0)
\$25,000-\$49,999	*	*	*	8,709	62.7	(55.7 - 69.7)
\$50,000-\$74,999	*	*	*	5,534	76.4	(70.9 - 81.9)
≥ \$75,000	*	*	*	8,448	80.4	(77.4 - 83.4)

<sup>1</sup> Defined as having had teeth cleaned by a dentist or dental hygienist within the past year

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 16a. Permanent teeth extracted due to decay or gum disease among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	650	24,800	50.2	(44.4 - 56.1)	37,652	7,426,500	50.4	(47.8 - 52.9)
<b>Gender</b>								
Male	239	11,500	45.1	(35.9 - 54.2)	14,457	3,367,000	47.8	(43.9 - 51.6)
Female	411	13,300	55.7	(49.4 - 61.9)	23,195	4,059,500	52.7	(49.5 - 56.0)
<b>Age (years)</b>								
18-34	58	2,600	17.3	(8.1 - 26.4)	3,921	1,121,600	25.4	(19.7 - 31.0)
35-44	*	*	*	*	4,964	1,250,000	43.6	(37.9 - 49.2)
45-54	135	6,000	63.1	(53.7 - 72.5)	7,780	1,595,300	56.1	(51.4 - 60.7)
55-64	151	5,100	71.1	(63.0 - 79.1)	8,648	1,382,800	65.5	(61.6 - 69.4)
≥ 65	223	7,700	85.8	(80.9 - 90.8)	12,339	2,076,900	83.0	(80.6 - 85.4)
<b>Race/Ethnicity</b>								
White Non-Hispanic	628	23,500	51.1	(45.3 - 56.9)	35,364	4,425,700	47.9	(46.0 - 49.9)
Black Non-Hispanic	*	*	*	*	727	1,244,700	58.2	(48.9 - 67.4)
Hispanic	*	*	*	*	853	1,220,500	54.2	(44.9 - 63.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	307	14,400	64.4	(54.5 - 74.4)	15,427	3,436,000	61.9	(57.2 - 66.6)
Some College	*	*	*	*	9,798	1,726,200	49.3	(44.3 - 54.2)
≥ College Degree	181	4,600	32.9	(24.5 - 41.3)	12,351	2,242,600	40.1	(36.5 - 43.6)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,483	2,171,900	62.3	(55.8 - 68.8)
\$25,000-\$49,999	*	*	*	*	9,552	1,577,400	54.7	(48.3 - 61.0)
\$50,000-\$74,999	*	*	*	*	5,744	1,000,700	51.5	(45.6 - 57.4)
≥ \$75,000	112	3,200	29.3	(19.4 - 39.1)	8,516	1,738,200	38.8	(35.3 - 42.2)

<sup>1</sup> Defined as having had at least one permanent tooth extracted because of tooth decay or gum disease

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have had at least one permanent tooth extracted because of tooth decay or gum disease (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 16b. Permanent teeth extracted due to decay or gum disease among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	650	48.7	(42.8 - 54.6)	37,652	49.4	(46.9 - 51.9)
<b>Gender</b>						
Male	239	45.6	(36.4 - 54.8)	14,457	47.9	(44.1 - 51.7)
Female	411	52.0	(45.2 - 58.8)	23,195	50.7	(47.4 - 54.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	628	49.5	(43.6 - 55.3)	35,364	46.5	(44.5 - 48.5)
Black Non-Hispanic	*	*	*	727	57.7	(48.6 - 66.9)
Hispanic	*	*	*	853	53.8	(44.6 - 62.9)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	307	63.7	(53.8 - 73.6)	15,427	61.3	(56.7 - 66.0)
Some College	*	*	*	9,798	47.9	(43.0 - 52.9)
≥ College Degree	181	31.3	(22.9 - 39.7)	12,351	39.0	(35.5 - 42.5)
<b>Income</b>						
≤ \$24,999	*	*	*	9,483	61.6	(55.2 - 68.1)
\$25,000-\$49,999	*	*	*	9,552	54.2	(48.0 - 60.3)
\$50,000-\$74,999	*	*	*	5,744	49.9	(44.0 - 55.9)
≥ \$75,000	112	27.6	(17.9 - 37.3)	8,516	37.7	(34.2 - 41.1)

<sup>1</sup> Defined as having had at least one permanent tooth extracted because of tooth decay or gum disease

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 17a. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	223	2,200	25.0	(18.9 - 31.2)	12,339	491,200	19.6	(16.0 - 23.3)
<b>Gender</b>								
Male	80	900	24.4	(14.5 - 34.3)	4,352	167,200	16.4	(11.8 - 20.9)
Female	143	1,300	25.5	(17.8 - 33.2)	7,987	324,000	21.8	(16.6 - 27.0)
<b>Age (years)</b>								
≥ 65	223	2,200	25.0	(18.9 - 31.2)	12,339	491,200	19.6	(16.0 - 23.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	220	2,200	25.5	(19.3 - 31.7)	11,885	322,600	16.8	(14.3 - 19.2)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	118	1,700	35.9	(26.5 - 45.3)	6,358	314,100	25.3	(19.9 - 30.7)
Some College	*	*	*	*	2,736	90,600	20.2	(12.1 - 28.2)
≥ College Degree	*	*	*	*	3,214	80,500	10.1	(3.6 - 16.5)
<b>Income</b>								
≤ \$24,999	*	*	*	*	4,558	199,600	26.9	(19.3 - 34.4)
\$25,000-\$49,999	*	*	*	*	3,365	123,900	21.4	(13.1 - 29.6)
\$50,000-\$74,999	*	*	*	*	1,179	21,400	7.3	(3.8 - 10.8)
≥ \$75,000	*	*	*	*	1,187	9,600	2.5	(1.0 - 4.0)

<sup>1</sup> Defined as having all permanent teeth extracted because of tooth decay or gum disease among adults aged 65 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 65 and older who have had all permanent teeth extracted because of tooth decay or gum disease (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 17b. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	223	25.0	(18.6 - 31.3)	12,339	19.4	(15.7 - 23.2)
<b>Gender</b>						
Male	80	24.4	(14.6 - 34.3)	4,352	16.5	(11.5 - 21.4)
Female	143	25.6	(17.9 - 33.2)	7,987	22.1	(16.6 - 27.7)
<b>Race/Ethnicity</b>						
White Non-Hispanic	220	25.4	(19.0 - 31.9)	11,885	16.6	(13.9 - 19.3)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	118	36.0	(26.2 - 45.7)	6,358	24.9	(19.3 - 30.5)
Some College	*	*	*	2,736	21.0	(12.1 - 29.9)
≥ College Degree	*	*	*	3,214	10.0	(3.6 - 16.3)
<b>Income</b>						
≤ \$24,999	*	*	*	4,558	27.3	(19.1 - 35.5)
\$25,000-\$49,999	*	*	*	3,365	21.0	(12.9 - 29.1)
\$50,000-\$74,999	*	*	*	1,179	7.6	(3.6 - 11.5)
≥ \$75,000	*	*	*	1,187	2.2	(0.9 - 3.5)

<sup>1</sup> Defined as having all permanent teeth extracted because of tooth decay or gum disease among adults aged 65 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 18a. Flu shot within the past 12 months<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	659	19,200	38.6	(33.1 - 44.1)	38,079	6,255,900	42.3	(39.9 - 44.8)
<b>Gender</b>								
Male	241	8,400	33.0	(24.0 - 41.9)	14,579	2,798,900	39.6	(35.8 - 43.5)
Female	418	10,800	44.6	(38.6 - 50.6)	23,500	3,456,900	44.8	(41.6 - 47.9)
<b>Age (years)</b>								
18-49	198	7,500	26.3	(17.7 - 34.9)	12,505	2,685,300	31.1	(27.3 - 34.9)
50-64	231	4,900	40.8	(33.8 - 47.8)	12,972	1,662,900	46.0	(42.6 - 49.5)
≥ 65	230	6,800	74.3	(68.3 - 80.2)	12,602	1,907,700	75.2	(72.0 - 78.4)
<b>Race/Ethnicity</b>								
White Non-Hispanic	637	17,900	38.6	(33.2 - 44.0)	35,770	4,029,000	43.3	(41.4 - 45.3)
Black Non-Hispanic	*	*	*	*	738	907,000	42.1	(33.1 - 51.1)
Hispanic	*	*	*	*	860	882,900	39.6	(30.4 - 48.9)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	313	8,900	39.4	(31.3 - 47.5)	15,663	2,457,800	44.3	(39.8 - 48.8)
Some College	161	3,700	28.9	(19.7 - 38.0)	9,898	1,377,700	38.9	(34.0 - 43.9)
≥ College Degree	*	*	*	*	12,436	2,404,400	43.0	(39.5 - 46.4)
<b>Income</b>								
≤ \$24,999	191	4,100	34.5	(24.9 - 44.0)	9,635	1,656,800	47.6	(41.2 - 54.0)
\$25,000-\$49,999	184	5,000	37.0	(27.9 - 46.2)	9,647	1,200,600	41.4	(35.6 - 47.3)
\$50,000-\$74,999	*	*	*	*	5,763	792,300	40.2	(34.3 - 46.2)
≥ \$75,000	*	*	*	*	8,548	1,776,100	39.6	(36.3 - 42.8)

<sup>1</sup> Defined as having had a flu shot within the past 12 months

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have had a flu shot within the past 12 months (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 18b. Flu shot within the past 12 months<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	659	37.3	(31.7 - 42.9)	38,079	41.7	(39.2 - 44.1)
<b>Gender</b>						
Male	241	33.6	(24.5 - 42.6)	14,579	40.0	(36.2 - 43.8)
Female	418	41.2	(34.8 - 47.7)	23,500	43.1	(40.0 - 46.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	637	37.0	(31.5 - 42.5)	35,770	42.4	(40.4 - 44.3)
Black Non-Hispanic	*	*	*	738	42.3	(33.3 - 51.3)
Hispanic	*	*	*	860	39.0	(29.9 - 48.1)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	313	37.5	(29.4 - 45.7)	15,663	43.7	(39.1 - 48.2)
Some College	161	27.8	(18.6 - 37.1)	9,898	37.8	(33.0 - 42.7)
≥ College Degree	*	*	*	12,436	42.5	(39.1 - 46.0)
<b>Income</b>						
≤ \$24,999	191	32.6	(23.3 - 42.0)	9,635	47.3	(41.0 - 53.7)
\$25,000-\$49,999	184	36.8	(27.2 - 46.3)	9,647	40.3	(34.5 - 46.0)
\$50,000-\$74,999	*	*	*	5,763	39.4	(33.4 - 45.4)
≥ \$75,000	*	*	*	8,548	39.1	(35.8 - 42.4)

<sup>1</sup> Defined as having had a flu shot within the past 12 months

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 19a. Flu vaccine that was sprayed in nose within the past 12 months<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	*	*	*	*	38,106	248,300	1.7	(0.6 - 2.8)
<b>Gender</b>								
Male	*	*	*	*	14,586	148,000	2.1	(0.0 - 4.2)
Female	*	*	*	*	23,520	100,300	1.3	(0.5 - 2.1)
<b>Age (years)</b>								
18-49	*	*	*	*	12,535	184,900	2.1	(0.3 - 3.9)
50-64	*	*	*	*	12,968	36,000	1.0	(0.2 - 1.8)
≥ 65	*	*	*	*	12,603	27,400	1.1	(0.4 - 1.8)
<b>Race/Ethnicity</b>								
White Non-Hispanic	*	*	*	*	35,796	66,000	0.7	(0.4 - 1.0)
Black Non-Hispanic	*	*	*	*	740	38,500	1.8	(0.0 - 4.1)
Hispanic	*	*	*	*	860	133,700	5.9	(0.0 - 12.3)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	*	*	*	*	15,679	172,500	3.1	(0.3 - 5.9)
Some College	*	*	*	*	9,897	43,500	1.2	(0.4 - 2.0)
≥ College Degree	*	*	*	*	12,449	31,600	0.6	(0.3 - 0.8)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,636	138,400	3.9	(0.0 - 8.1)
\$25,000-\$49,999	*	*	*	*	9,648	41,500	1.4	(0.0 - 3.1)
\$50,000-\$74,999	*	*	*	*	5,766	13,300	0.7	(0.2 - 1.2)
≥ \$75,000	*	*	*	*	8,561	35,300	0.8	(0.3 - 1.3)

<sup>1</sup> Defined as having had flu vaccine sprayed in nose within the past 12 months

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have had flu vaccine sprayed in nose within the past 12 months (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 19b. Flu vaccine that was sprayed in nose within the past 12 months<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	*	*	*	38,106	1.7	(0.6 - 2.7)
<b>Gender</b>						
Male	*	*	*	14,586	2.0	(0.1 - 3.9)
Female	*	*	*	23,520	1.4	(0.5 - 2.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	*	*	*	35,796	0.7	(0.4 - 1.0)
Black Non-Hispanic	*	*	*	740	1.9	(0.0 - 4.4)
Hispanic	*	*	*	860	5.9	(0.0 - 11.9)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	15,679	3.1	(0.4 - 5.8)
Some College	*	*	*	9,897	1.2	(0.4 - 2.0)
≥ College Degree	*	*	*	12,449	0.6	(0.3 - 0.8)
<b>Income</b>						
≤ \$24,999	*	*	*	9,636	3.8	(0.0 - 7.7)
\$25,000-\$49,999	*	*	*	9,648	1.5	(0.0 - 3.4)
\$50,000-\$74,999	*	*	*	5,766	0.7	(0.1 - 1.2)
≥ \$75,000	*	*	*	8,561	0.8	(0.3 - 1.3)

<sup>1</sup> Defined as having had flu vaccine sprayed in nose within the past 12 months

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 20a. Flu shot or flu vaccine sprayed in nose within the past 12 months<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	659	19,400	39.0	(33.5 - 44.5)	38,059	6,302,300	42.7	(40.2 - 45.1)
<b>Gender</b>								
Male	241	8,600	33.7	(24.7 - 42.7)	14,568	2,822,700	40.0	(36.1 - 43.9)
Female	418	10,800	44.6	(38.6 - 50.6)	23,491	3,479,500	45.1	(42.0 - 48.2)
<b>Age (years)</b>								
18-49	198	7,700	26.9	(18.3 - 35.5)	12,501	2,726,400	31.6	(27.8 - 35.4)
50-64	231	4,900	40.8	(33.8 - 47.8)	12,964	1,664,200	46.1	(42.7 - 49.5)
≥ 65	230	6,800	74.3	(68.3 - 80.2)	12,594	1,911,700	75.4	(72.2 - 78.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	637	18,100	39.0	(33.6 - 44.4)	35,750	4,048,700	43.6	(41.7 - 45.5)
Black Non-Hispanic	*	*	*	*	738	910,000	42.2	(33.2 - 51.2)
Hispanic	*	*	*	*	860	903,300	40.6	(31.3 - 49.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	313	8,900	39.4	(31.3 - 47.5)	15,651	2,480,100	44.7	(40.2 - 49.3)
Some College	161	3,700	28.9	(19.7 - 38.0)	9,895	1,387,300	39.2	(34.3 - 44.1)
≥ College Degree	*	*	*	*	12,431	2,418,300	43.2	(39.8 - 46.7)
<b>Income</b>								
≤ \$24,999	191	4,100	34.5	(24.9 - 44.0)	9,626	1,675,100	48.2	(41.8 - 54.6)
\$25,000-\$49,999	184	5,000	37.0	(27.9 - 46.2)	9,643	1,208,300	41.7	(35.8 - 47.6)
\$50,000-\$74,999	*	*	*	*	5,761	795,600	40.4	(34.5 - 46.3)
≥ \$75,000	*	*	*	*	8,548	1,791,200	39.9	(36.6 - 43.2)

<sup>1</sup> Defined as having had a flu shot or flu vaccine sprayed in nose within the past 12 months

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have had a flu shot or flu vaccine sprayed in nose within the past 12 months (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 20b. Flu shot or flu vaccine sprayed in nose within the past 12 months<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	659	37.7	(32.1 - 43.4)	38,059	42.0	(39.6 - 44.5)
<b>Gender</b>						
Male	241	34.4	(25.3 - 43.5)	14,568	40.4	(36.6 - 44.2)
Female	418	41.2	(34.8 - 47.7)	23,491	43.5	(40.3 - 46.7)
<b>Race/Ethnicity</b>						
White Non-Hispanic	637	37.5	(32.0 - 43.0)	35,750	42.6	(40.7 - 44.6)
Black Non-Hispanic	*	*	*	738	42.4	(33.4 - 51.4)
Hispanic	*	*	*	860	40.0	(30.9 - 49.1)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	313	37.5	(29.4 - 45.7)	15,651	44.1	(39.6 - 48.6)
Some College	161	27.8	(18.6 - 37.1)	9,895	38.1	(33.2 - 43.0)
≥ College Degree	*	*	*	12,431	42.8	(39.4 - 46.3)
<b>Income</b>						
≤ \$24,999	191	32.6	(23.3 - 42.0)	9,626	48.0	(41.6 - 54.3)
\$25,000-\$49,999	184	36.8	(27.2 - 46.3)	9,643	40.5	(34.8 - 46.3)
\$50,000-\$74,999	*	*	*	5,761	39.6	(33.6 - 45.6)
≥ \$75,000	*	*	*	8,548	39.4	(36.1 - 42.7)

<sup>1</sup> Defined as having had a flu shot or flu vaccine sprayed in nose within the past 12 months

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 21a. Flu shot among adults aged 65 and older<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	230	6,800	74.3	(68.3 - 80.2)	12,602	1,907,700	75.2	(72.0 - 78.4)
<b>Gender</b>								
Male	*	*	*	*	4,417	772,900	74.6	(69.3 - 79.8)
Female	149	4,000	75.7	(68.7 - 82.8)	8,185	1,134,800	75.6	(71.6 - 79.6)
<b>Age (years)</b>								
≥ 65	230	6,800	74.3	(68.3 - 80.2)	12,602	1,907,700	75.2	(72.0 - 78.4)
<b>Race/Ethnicity</b>								
White Non-Hispanic	227	6,700	74.6	(68.7 - 80.5)	12,137	1,487,100	76.0	(73.5 - 78.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	122	3,600	73.5	(65.3 - 81.7)	6,518	930,800	74.0	(68.9 - 79.2)
Some College	*	*	*	*	2,790	317,800	69.7	(62.7 - 76.7)
≥ College Degree	*	*	*	*	3,261	654,000	80.5	(76.4 - 84.6)
<b>Income</b>								
≤ \$24,999	98	2,600	75.7	(66.8 - 84.6)	4,665	545,300	72.4	(65.6 - 79.3)
\$25,000-\$49,999	*	*	*	*	3,422	430,600	72.8	(67.0 - 78.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	1,198	301,000	77.5	(70.9 - 84.2)

<sup>1</sup> Defined as having had a flu shot during the past 12 months among adults aged 65 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 65 and older who have had a flu shot within the past 12 months (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 21b. Flu shot among adults aged 65 and older<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	230	74.0	(67.8 - 80.2)	12,602	75.0	(71.5 - 78.5)
<b>Gender</b>						
Male	*	*	*	4,417	74.4	(68.7 - 80.1)
Female	149	75.7	(68.7 - 82.8)	8,185	75.5	(71.3 - 79.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	227	74.4	(68.2 - 80.6)	12,137	75.9	(73.0 - 78.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	122	73.2	(64.5 - 81.8)	6,518	73.3	(67.5 - 79.1)
Some College	*	*	*	2,790	70.0	(62.7 - 77.2)
≥ College Degree	*	*	*	3,261	80.7	(76.2 - 85.1)
<b>Income</b>						
≤ \$24,999	98	75.5	(66.1 - 84.8)	4,665	72.6	(65.1 - 80.1)
\$25,000-\$49,999	*	*	*	3,422	71.4	(65.0 - 77.9)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	1,198	78.0	(70.8 - 85.1)

<sup>1</sup> Defined as having had a flu shot during the past 12 months among adults aged 65 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 22a. Pneumonia shot or pneumococcal vaccine<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	625	12,600	27.7	(23.1 - 32.3)	35,303	3,456,800	26.1	(23.9 - 28.3)
<b>Gender</b>								
Male	220	5,400	23.7	(16.3 - 31.1)	13,052	1,542,200	25.2	(21.5 - 28.8)
Female	405	7,200	31.6	(26.4 - 36.8)	22,251	1,914,600	26.9	(24.2 - 29.6)
<b>Age (years)</b>								
18-49	184	3,300	13.1	(6.7 - 19.6)	10,943	1,097,000	14.6	(11.3 - 17.9)
50-64	218	2,900	25.1	(18.7 - 31.5)	12,179	758,000	23.0	(19.8 - 26.2)
≥ 65	223	6,400	72.5	(66.0 - 78.9)	12,181	1,601,700	65.7	(61.7 - 69.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	607	12,400	29.0	(24.3 - 33.7)	33,225	2,358,700	28.7	(26.9 - 30.5)
Black Non-Hispanic	*	*	*	*	685	584,400	28.4	(19.5 - 37.3)
Hispanic	*	*	*	*	769	408,000	19.5	(12.6 - 26.4)
Other Non-Hispanic	*	*	*	*	624	105,600	12.2	(2.3 - 22.2)
<b>Education</b>								
≤ High School	301	6,400	31.1	(23.9 - 38.4)	14,700	1,535,300	30.4	(26.4 - 34.3)
Some College	153	3,300	27.3	(18.1 - 36.5)	9,151	836,200	27.2	(21.9 - 32.5)
≥ College Degree	169	2,900	22.3	(15.2 - 29.4)	11,375	1,070,700	21.4	(18.6 - 24.1)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,082	1,050,900	33.1	(27.6 - 38.6)
\$25,000-\$49,999	172	3,100	25.1	(17.5 - 32.6)	9,033	797,500	30.3	(24.1 - 36.6)
\$50,000-\$74,999	*	*	*	*	5,320	378,800	21.8	(17.4 - 26.1)
≥ \$75,000	*	*	*	*	7,773	648,700	16.3	(14.1 - 18.5)

<sup>1</sup> Defined as having ever had a pneumonia shot or pneumococcal vaccine

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have ever had a pneumonia shot or pneumococcal vaccine (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 22b. Pneumonia shot or pneumococcal vaccine<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	625	26.0	(21.4 - 30.5)	35,303	25.4	(23.2 - 27.6)
<b>Gender</b>						
Male	220	24.1	(16.8 - 31.5)	13,052	25.8	(22.3 - 29.4)
Female	405	27.8	(22.5 - 33.1)	22,251	25.1	(22.4 - 27.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	607	27.1	(22.5 - 31.8)	33,225	27.7	(25.9 - 29.5)
Black Non-Hispanic	*	*	*	685	27.9	(19.1 - 36.7)
Hispanic	*	*	*	769	19.6	(12.8 - 26.5)
Other Non-Hispanic	*	*	*	624	11.3	(2.0 - 20.7)
<b>Education</b>						
≤ High School	301	28.9	(21.7 - 36.1)	14,700	29.5	(25.6 - 33.5)
Some College	153	25.3	(16.3 - 34.3)	9,151	26.2	(21.0 - 31.3)
≥ College Degree	169	21.8	(14.5 - 29.1)	11,375	21.0	(18.3 - 23.8)
<b>Income</b>						
≤ \$24,999	*	*	*	9,082	32.3	(26.9 - 37.8)
\$25,000-\$49,999	172	24.4	(16.8 - 32.1)	9,033	29.0	(23.0 - 35.0)
\$50,000-\$74,999	*	*	*	5,320	21.2	(16.8 - 25.7)
≥ \$75,000	*	*	*	7,773	16.2	(13.9 - 18.5)

<sup>1</sup> Defined as having ever had a pneumonia shot or pneumococcal vaccine

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 23a. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	223	6,400	72.5	(66.0 - 78.9)	12,181	1,601,700	65.7	(61.7 - 69.7)
<b>Gender</b>								
Male	*	*	*	*	4,180	606,400	62.1	(55.5 - 68.7)
Female	147	3,800	73.8	(66.0 - 81.6)	8,001	995,300	68.0	(63.1 - 73.0)
<b>Age (years)</b>								
≥ 65	223	6,400	72.5	(66.0 - 78.9)	12,181	1,601,700	65.7	(61.7 - 69.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	221	6,300	72.1	(65.6 - 78.6)	11,735	1,281,300	68.2	(64.9 - 71.5)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	120	3,500	74.1	(65.6 - 82.7)	6,345	806,200	65.6	(59.5 - 71.7)
Some College	*	*	*	*	2,694	289,500	66.8	(60.1 - 73.4)
≥ College Degree	*	*	*	*	3,111	501,100	65.6	(58.5 - 72.7)
<b>Income</b>								
≤ \$24,999	96	2,800	83.7	(76.3 - 91.2)	4,533	497,100	67.7	(59.9 - 75.5)
\$25,000-\$49,999	*	*	*	*	3,324	344,300	61.5	(53.3 - 69.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	1,138	236,300	64.5	(55.2 - 73.8)

<sup>1</sup> Defined as having ever had a pneumonia shot or pneumococcal vaccine among adults aged 65 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 65 and older who have ever had a pneumonia shot or pneumococcal vaccine (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 23b. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	223	72.2	(65.5 - 78.9)	12,181	64.7	(60.4 - 69.1)
<b>Gender</b>						
Male	*	*	*	4,180	61.5	(54.3 - 68.6)
Female	147	73.8	(66.0 - 81.6)	8,001	67.6	(62.4 - 72.9)
<b>Race/Ethnicity</b>						
White Non-Hispanic	221	71.8	(65.0 - 78.6)	11,735	67.1	(63.2 - 70.9)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	120	73.0	(63.8 - 82.2)	6,345	64.2	(57.4 - 71.0)
Some College	*	*	*	2,694	65.8	(58.4 - 73.3)
≥ College Degree	*	*	*	3,111	65.3	(57.7 - 72.9)
<b>Income</b>						
≤ \$24,999	96	83.2	(75.3 - 91.2)	4,533	66.0	(57.0 - 74.9)
\$25,000-\$49,999	*	*	*	3,324	60.3	(51.9 - 68.6)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	*	*

<sup>1</sup> Defined as having ever had a pneumonia shot or pneumococcal vaccine among adults aged 65 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 24a. Ever had a mammogram among women aged 40 and older<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	353	14,200	89.9	(86.0 - 93.8)	19,491	4,328,800	91.8	(89.7 - 94.0)
<b>Gender</b>								
Male	-	-	-	-	-	-	-	-
Female	353	14,200	89.9	(86.0 - 93.8)	19,491	4,328,800	91.8	(89.7 - 94.0)
<b>Age (years)</b>								
40-49	*	*	*	*	3,851	1,287,500	85.0	(79.7 - 90.3)
50-64	141	6,300	95.1	(90.9 - 99.2)	7,452	1,641,700	96.0	(94.1 - 97.9)
≥ 65	149	4,900	92.5	(88.2 - 96.8)	8,188	1,399,600	94.1	(90.5 - 97.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	345	13,900	90.6	(86.8 - 94.3)	18,548	2,976,100	93.9	(92.7 - 95.1)
Black Non-Hispanic	*	*	*	*	342	746,600	93.8	(87.3 - 100)
Hispanic	*	*	*	*	333	447,600	91.8	(84.6 - 99.0)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	177	6,600	86.1	(79.8 - 92.4)	8,266	1,698,700	91.0	(87.4 - 94.5)
Some College	92	3,900	89.3	(81.2 - 97.4)	5,322	1,055,500	93.6	(90.2 - 97.0)
≥ College Degree	84	3,700	98.1	(95.4 - 100)	5,872	1,560,700	91.6	(87.5 - 95.6)
<b>Income</b>								
≤ \$24,999	121	3,700	83.5	(74.6 - 92.5)	5,691	1,023,400	93.5	(89.7 - 97.4)
\$25,000-\$49,999	84	3,400	89.3	(80.3 - 98.2)	4,857	860,300	89.5	(83.6 - 95.5)
\$50,000-\$74,999	*	*	*	*	2,648	547,100	94.6	(92.2 - 96.9)
≥ \$75,000	53	3,000	95.3	(88.6 - 100)	3,582	1,185,100	92.2	(88.2 - 96.1)

<sup>1</sup> Defined as having ever had a mammogram among women aged 40 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of women aged 40 and older who have ever had a mammogram (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 24b. Ever had a mammogram among women aged 40 and older<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	353	88.6	(84.0 - 93.3)	19,491	91.1	(88.7 - 93.6)
<b>Gender</b>						
Male	-	-	-	-	-	-
Female	353	88.6	(84.0 - 93.3)	19,491	91.1	(88.7 - 93.6)
<b>Race/Ethnicity</b>						
White Non-Hispanic	345	89.5	(85.1 - 93.9)	18,548	93.3	(92.0 - 94.7)
Black Non-Hispanic	*	*	*	342	93.1	(85.7 - 100)
Hispanic	*	*	*	333	91.0	(82.8 - 99.2)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	177	84.5	(77.1 - 92.0)	8,266	90.2	(86.1 - 94.2)
Some College	*	*	*	5,322	92.9	(88.9 - 96.9)
≥ College Degree	84	98.3	(95.9 - 100)	5,872	90.9	(86.6 - 95.3)
<b>Income</b>						
≤ \$24,999	*	*	*	5,691	93.0	(88.5 - 97.4)
\$25,000-\$49,999	*	*	*	4,857	88.8	(82.5 - 95.1)
\$50,000-\$74,999	*	*	*	2,648	94.3	(91.7 - 96.8)
≥ \$75,000	53	94.4	(86.3 - 100)	3,582	91.2	(86.8 - 95.6)

<sup>1</sup> Defined as having ever had a mammogram among women aged 40 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 25a. Had mammogram within the past 2 years among women aged 40 and older<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	353	12,300	77.5	(72.4 - 82.7)	19,376	3,760,000	80.3	(77.8 - 82.9)
<b>Gender</b>								
Male	-	-	-	-	-	-	-	-
Female	353	12,300	77.5	(72.4 - 82.7)	19,376	3,760,000	80.3	(77.8 - 82.9)
<b>Age (years)</b>								
40-49	*	*	*	*	3,842	1,122,700	74.4	(68.5 - 80.2)
50-64	141	5,400	82.5	(75.2 - 89.8)	7,428	1,487,900	87.2	(84.5 - 89.8)
≥ 65	149	4,200	79.8	(73.3 - 86.3)	8,106	1,149,400	78.5	(74.2 - 82.9)
<b>Race/Ethnicity</b>								
White Non-Hispanic	345	12,000	78.1	(73.0 - 83.3)	18,438	2,531,500	80.2	(78.3 - 82.2)
Black Non-Hispanic	*	*	*	*	339	667,100	85.6	(77.5 - 93.8)
Hispanic	*	*	*	*	332	419,700	86.1	(77.5 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	177	5,500	70.8	(62.8 - 78.7)	8,204	1,486,900	80.5	(76.4 - 84.6)
Some College	*	*	*	*	5,297	929,400	83.0	(78.8 - 87.1)
≥ College Degree	84	3,500	93.0	(87.6 - 98.3)	5,845	1,334,800	78.6	(73.9 - 83.2)
<b>Income</b>								
≤ \$24,999	*	*	*	*	5,650	882,300	80.8	(75.9 - 85.7)
\$25,000-\$49,999	84	3,200	82.7	(72.9 - 92.6)	4,846	732,300	76.3	(69.7 - 82.9)
\$50,000-\$74,999	*	*	*	*	2,641	494,200	85.6	(81.9 - 89.4)
≥ \$75,000	*	*	*	*	3,572	1,025,900	80.0	(75.2 - 84.8)

<sup>1</sup> Defined as having had a mammogram within the past 2 years among women aged 40 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of women aged 40 and older who have had a mammogram within the past 2 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 25b. Had mammogram within the past 2 years among women aged 40 and older<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	353	76.3	(70.5 - 82.1)	19,376	79.7	(76.9 - 82.5)
<b>Gender</b>						
Male	-	-	-	-	-	-
Female	353	76.3	(70.5 - 82.1)	19,376	79.7	(76.9 - 82.5)
<b>Race/Ethnicity</b>						
White Non-Hispanic	345	77.0	(71.2 - 82.7)	18,438	79.7	(77.6 - 81.8)
Black Non-Hispanic	*	*	*	339	85.3	(76.5 - 94.2)
Hispanic	*	*	*	332	85.2	(75.6 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	177	67.9	(58.9 - 77.0)	8,204	80.1	(75.6 - 84.7)
Some College	*	*	*	5,297	82.3	(77.7 - 86.9)
≥ College Degree	84	93.3	(88.0 - 98.5)	5,845	77.8	(72.9 - 82.8)
<b>Income</b>						
≤ \$24,999	*	*	*	5,650	80.8	(75.4 - 86.2)
\$25,000-\$49,999	*	*	*	4,846	75.7	(68.7 - 82.6)
\$50,000-\$74,999	*	*	*	2,641	85.5	(81.6 - 89.4)
≥ \$75,000	*	*	*	3,572	79.0	(73.8 - 84.2)

<sup>1</sup> Defined as having had a mammogram within the past 2 years among women aged 40 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 26a. Had mammogram within the past 2 years among women aged 50 and older<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	290	9,600	81.3	(76.3 - 86.3)	15,534	2,637,300	83.2	(80.7 - 85.7)
<b>Gender</b>								
Male	-	-	-	-	-	-	-	-
Female	290	9,600	81.3	(76.3 - 86.3)	15,534	2,637,300	83.2	(80.7 - 85.7)
<b>Age (years)</b>								
50-64	141	5,400	82.5	(75.2 - 89.8)	7,428	1,487,900	87.2	(84.5 - 89.8)
≥ 65	149	4,200	79.8	(73.3 - 86.3)	8,106	1,149,400	78.5	(74.2 - 82.9)
<b>Race/Ethnicity</b>								
White Non-Hispanic	285	9,400	81.6	(76.6 - 86.7)	14,887	1,907,000	82.3	(80.3 - 84.4)
Black Non-Hispanic	*	*	*	*	249	385,600	89.9	(83.2 - 96.7)
Hispanic	*	*	*	*	220	286,700	93.9	(89.1 - 98.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	158	5,000	78.6	(71.7 - 85.5)	7,026	1,094,500	83.3	(80.0 - 86.6)
Some College	*	*	*	*	4,124	646,700	87.3	(84.2 - 90.3)
≥ College Degree	61	2,300	93.1	(87.5 - 98.8)	4,356	890,500	80.6	(75.1 - 86.1)
<b>Income</b>								
≤ \$24,999	110	2,800	77.4	(69.3 - 85.6)	4,991	649,100	81.0	(76.8 - 85.2)
\$25,000-\$49,999	79	2,900	84.5	(75.5 - 93.5)	3,913	533,300	78.6	(70.9 - 86.2)
\$50,000-\$74,999	*	*	*	*	1,943	359,300	89.6	(86.3 - 93.0)
≥ \$75,000	*	*	*	*	2,303	601,300	84.6	(79.3 - 89.9)

<sup>1</sup> Defined as having had a mammogram within the past 2 years among women aged 50 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of women aged 50 and older who have had a mammogram within the past 2 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 26b. Had mammogram within the past 2 years among women aged 50 and older<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	290	81.5	(76.5 - 86.5)	15,534	83.1	(80.3 - 85.8)
<b>Gender</b>						
Male	-	-	-	-	-	-
Female	290	81.5	(76.5 - 86.5)	15,534	83.1	(80.3 - 85.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	285	81.8	(76.8 - 86.8)	14,887	82.1	(79.9 - 84.3)
Black Non-Hispanic	*	*	*	249	90.4	(83.6 - 97.2)
Hispanic	*	*	*	220	94.1	(89.3 - 99.0)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	158	78.6	(71.6 - 85.6)	7,026	83.6	(80.1 - 87.0)
Some College	*	*	*	4,124	87.3	(84.1 - 90.5)
≥ College Degree	61	92.8	(86.9 - 98.7)	4,356	79.9	(73.9 - 86.0)
<b>Income</b>						
≤ \$24,999	110	77.6	(69.5 - 85.7)	4,991	81.3	(76.9 - 85.6)
\$25,000-\$49,999	79	84.6	(75.6 - 93.6)	3,913	78.1	(69.9 - 86.4)
\$50,000-\$74,999	*	*	*	1,943	90.2	(87.1 - 93.4)
≥ \$75,000	*	*	*	2,303	83.9	(77.9 - 89.9)

<sup>1</sup> Defined as having had a mammogram within the past 2 years among women aged 50 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 27a. Ever had a Pap test among women<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	416	23,500	97.5	(95.8 - 99.1)	23,381	7,045,300	92.1	(90.0 - 94.3)
<b>Gender</b>								
Male	-	-	-	-	-	-	-	-
Female	416	23,500	97.5	(95.8 - 99.1)	23,381	7,045,300	92.1	(90.0 - 94.3)
<b>Age (years)</b>								
18-34	*	*	*	*	2,377	1,896,600	85.2	(79.3 - 91.0)
35-44	58	4,200	100	(100 - 100)	3,023	1,301,500	92.6	(87.7 - 97.5)
45-54	86	4,600	98.9	(97.4 - 100)	4,657	1,420,300	98.3	(97.0 - 99.6)
55-64	87	3,500	96.6	(91.1 - 100)	5,187	1,071,200	97.2	(95.0 - 99.3)
≥ 65	147	4,900	94.4	(90.8 - 98.0)	8,137	1,355,700	92.3	(88.5 - 96.1)
<b>Race/Ethnicity</b>								
White Non-Hispanic	405	22,500	97.4	(95.6 - 99.1)	21,976	4,439,700	94.2	(92.1 - 96.3)
Black Non-Hispanic	*	*	*	*	481	1,231,600	91.2	(84.0 - 98.4)
Hispanic	*	*	*	*	533	984,500	90.1	(85.3 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	197	10,200	96.7	(94.2 - 99.2)	9,526	2,540,500	88.3	(83.8 - 92.7)
Some College	111	6,800	97.2	(93.0 - 100)	6,462	1,782,200	93.5	(90.4 - 96.6)
≥ College Degree	108	6,500	98.9	(97.7 - 100)	7,345	2,643,200	95.0	(92.2 - 97.9)
<b>Income</b>								
≤ \$24,999	129	5,200	95.7	(92.6 - 98.9)	6,526	1,717,200	93.7	(90.9 - 96.6)
\$25,000-\$49,999	103	6,000	98.0	(94.7 - 100)	5,787	1,378,400	91.0	(85.6 - 96.4)
\$50,000-\$74,999	65	4,500	99.5	(98.5 - 100)	3,310	933,900	96.0	(91.8 - 100)
≥ \$75,000	63	4,500	100	(100 - 100)	4,570	1,943,300	95.4	(92.6 - 98.2)

<sup>1</sup> Defined as having ever had a Pap test among women

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of women who have ever had a Pap test (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 27b. Ever had a Pap test among women<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	416	97.7	(96.1 - 99.4)	23,381	91.8	(89.6 - 94.0)
<b>Gender</b>						
Male	-	-	-	-	-	-
Female	416	97.7	(96.1 - 99.4)	23,381	91.8	(89.6 - 94.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	405	97.6	(95.9 - 99.4)	21,976	94.0	(91.8 - 96.2)
Black Non-Hispanic	*	*	*	481	90.8	(83.5 - 98.1)
Hispanic	*	*	*	533	89.7	(84.6 - 94.7)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	197	97.3	(95.2 - 99.4)	9,526	87.8	(83.2 - 92.4)
Some College	111	97.1	(92.3 - 100)	6,462	93.1	(89.7 - 96.4)
≥ College Degree	108	99.1	(98.1 - 100)	7,345	94.9	(92.0 - 97.7)
<b>Income</b>						
≤ \$24,999	129	96.3	(93.6 - 99.1)	6,526	93.3	(90.1 - 96.5)
\$25,000-\$49,999	103	98.4	(95.7 - 100)	5,787	91.1	(85.7 - 96.5)
\$50,000-\$74,999	65	99.6	(98.9 - 100)	3,310	96.0	(91.9 - 100)
≥ \$75,000	63	100	(100 - 100)	4,570	95.0	(91.9 - 98.1)

<sup>1</sup> Defined as having ever had a Pap test among women

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 28a. Had a Pap test within the past 3 years among women<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	411	19,300	80.6	(76.4 - 84.8)	23,109	6,205,600	82.2	(79.8 - 84.6)
<b>Gender</b>								
Male	-	-	-	-	-	-	-	-
Female	411	19,300	80.6	(76.4 - 84.8)	23,109	6,205,600	82.2	(79.8 - 84.6)
<b>Age (years)</b>								
18-34	*	*	*	*	2,371	1,818,400	81.8	(75.5 - 88.1)
35-44	*	*	*	*	3,012	1,220,300	86.9	(81.8 - 92.0)
45-54	85	4,000	87.0	(79.4 - 94.5)	4,640	1,311,500	92.0	(89.8 - 94.3)
55-64	87	2,800	77.8	(68.0 - 87.6)	5,152	924,300	84.5	(81.0 - 88.0)
≥ 65	143	3,000	58.8	(50.0 - 67.5)	7,934	931,100	66.4	(61.8 - 71.0)
<b>Race/Ethnicity</b>								
White Non-Hispanic	400	18,300	80.0	(75.6 - 84.3)	21,720	3,807,100	81.5	(79.2 - 83.8)
Black Non-Hispanic	*	*	*	*	473	1,142,800	86.7	(79.1 - 94.3)
Hispanic	*	*	*	*	527	894,900	83.6	(76.3 - 91.0)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	194	7,700	73.8	(66.8 - 80.7)	9,384	2,132,600	76.0	(71.1 - 81.0)
Some College	109	5,400	78.7	(69.7 - 87.8)	6,401	1,594,300	84.0	(80.4 - 87.5)
≥ College Degree	108	6,200	93.5	(89.6 - 97.4)	7,281	2,406,000	86.9	(83.7 - 90.1)
<b>Income</b>								
≤ \$24,999	*	*	*	*	6,441	1,453,900	82.0	(77.9 - 86.0)
\$25,000-\$49,999	102	4,900	81.2	(73.3 - 89.2)	5,740	1,182,000	78.3	(72.6 - 83.9)
\$50,000-\$74,999	64	4,100	89.7	(81.4 - 98.1)	3,296	825,000	85.0	(78.0 - 91.9)
≥ \$75,000	63	4,400	97.3	(92.9 - 100)	4,555	1,810,200	89.0	(85.7 - 92.2)

<sup>1</sup> Defined as having had a Pap test within the past 3 years among women

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of women who have had a Pap test within the past 3 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 28b. Had a Pap test within the past 3 years among women<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	411	82.3	(78.0 - 86.6)	23,109	82.7	(80.2 - 85.1)
<b>Gender</b>						
Male	-	-	-	-	-	-
Female	411	82.3	(78.0 - 86.6)	23,109	82.7	(80.2 - 85.1)
<b>Race/Ethnicity</b>						
White Non-Hispanic	400	81.6	(77.1 - 86.1)	21,720	82.3	(80.0 - 84.6)
Black Non-Hispanic	*	*	*	473	87.0	(79.3 - 94.6)
Hispanic	*	*	*	527	83.4	(75.9 - 90.9)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	194	75.9	(68.7 - 83.2)	9,384	76.4	(71.3 - 81.5)
Some College	109	79.2	(69.6 - 88.8)	6,401	84.2	(80.5 - 88.0)
≥ College Degree	108	94.8	(91.7 - 98.0)	7,281	87.5	(84.3 - 90.6)
<b>Income</b>						
≤ \$24,999	*	*	*	6,441	82.7	(78.5 - 86.9)
\$25,000-\$49,999	102	83.1	(75.2 - 91.0)	5,740	79.2	(73.5 - 84.8)
\$50,000-\$74,999	64	89.7	(80.9 - 98.5)	3,296	85.7	(78.7 - 92.8)
≥ \$75,000	63	97.6	(93.7 - 100)	4,555	88.9	(85.5 - 92.3)

<sup>1</sup> Defined as having had a Pap test within the past 3 years among women

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 29a. Ever had a digital rectal exam among men aged 40 and older<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	205	10,500	72.3	(63.9 - 80.8)	12,033	3,176,500	73.8	(69.7 - 77.8)
<b>Gender</b>								
Male	205	10,500	72.3	(63.9 - 80.8)	12,033	3,176,500	73.8	(69.7 - 77.8)
Female	-	-	-	-	-	-	-	-
<b>Age (years)</b>								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	3,098	922,800	66.8	(59.9 - 73.8)
55-64	64	3,100	86.9	(78.9 - 94.9)	3,504	865,600	87.1	(82.5 - 91.7)
≥ 65	80	3,700	94.9	(90.3 - 99.6)	4,348	953,900	92.9	(91.0 - 94.9)
<b>Race/Ethnicity</b>								
White Non-Hispanic	197	10,200	73.7	(65.0 - 82.3)	11,411	2,452,100	80.3	(77.7 - 82.9)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	*	*	*	*	5,013	931,600	64.1	(57.3 - 71.0)
Some College	*	*	*	*	2,695	674,200	75.5	(67.8 - 83.2)
≥ College Degree	*	*	*	*	4,302	1,568,000	80.5	(74.0 - 87.0)
<b>Income</b>								
≤ \$24,999	*	*	*	*	2,573	573,300	65.9	(56.0 - 75.9)
\$25,000-\$49,999	*	*	*	*	3,168	530,100	74.6	(67.2 - 82.1)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,309	1,351,100	78.0	(72.6 - 83.5)

<sup>1</sup> Defined as having ever had a digital rectal exam among men aged 40 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of men aged 40 and older who have ever had a digital rectal exam (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 29b. Ever had a digital rectal exam among men aged 40 and older<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	205	71.3	(62.3 - 80.3)	12,033	73.7	(69.6 - 77.8)
<b>Gender</b>						
Male	205	71.3	(62.3 - 80.3)	12,033	73.7	(69.6 - 77.8)
Female	-	-	-	-	-	-
<b>Race/Ethnicity</b>						
White Non-Hispanic	197	72.4	(63.1 - 81.7)	11,411	79.9	(77.2 - 82.5)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	5,013	64.7	(57.8 - 71.7)
Some College	*	*	*	2,695	75.0	(67.1 - 82.9)
≥ College Degree	*	*	*	4,302	80.3	(73.8 - 86.8)
<b>Income</b>						
≤ \$24,999	*	*	*	2,573	66.9	(56.9 - 76.9)
\$25,000-\$49,999	*	*	*	3,168	74.8	(67.3 - 82.2)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,309	77.5	(72.0 - 83.0)

<sup>1</sup> Defined as having ever had a digital rectal exam among men aged 40 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 30a. Had digital rectal exam within the past 2 years among men aged 40 and older<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	200	7,800	54.6	(46.1 - 63.2)	11,923	2,380,300	55.8	(51.8 - 59.8)
<b>Gender</b>								
Male	200	7,800	54.6	(46.1 - 63.2)	11,923	2,380,300	55.8	(51.8 - 59.8)
Female	-	-	-	-	-	-	-	-
<b>Age (years)</b>								
40-44	*	*	*	*	1,078	246,800	27.5	(18.5 - 36.5)
45-54	*	*	*	*	3,074	650,000	47.6	(40.8 - 54.4)
55-64	*	*	*	*	3,467	673,500	68.1	(62.4 - 73.8)
≥ 65	78	3,200	83.9	(75.7 - 92.1)	4,304	809,900	79.7	(75.0 - 84.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	192	7,600	56.2	(47.4 - 65.1)	11,305	1,875,600	62.2	(59.2 - 65.2)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	*	*	*	*	4,963	718,700	49.6	(43.2 - 56.1)
Some College	*	*	*	*	2,665	511,500	58.5	(50.8 - 66.3)
≥ College Degree	*	*	*	*	4,277	1,148,400	59.4	(52.8 - 66.0)
<b>Income</b>								
≤ \$24,999	*	*	*	*	2,553	388,000	45.0	(35.0 - 55.0)
\$25,000-\$49,999	*	*	*	*	3,139	422,400	60.5	(53.0 - 68.0)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,295	978,300	56.9	(51.1 - 62.7)

<sup>1</sup> Defined as having had a digital rectal exam within the past 2 years among men aged 40 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of men aged 40 and older who have had a digital rectal exam within the past 2 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 30b. Had digital rectal exam within the past 2 years among men aged 40 and older<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	200	54.2	(45.3 - 63.1)	11,923	55.9	(51.7 - 60.0)
<b>Gender</b>						
Male	200	54.2	(45.3 - 63.1)	11,923	55.9	(51.7 - 60.0)
Female	-	-	-	-	-	-
<b>Race/Ethnicity</b>						
White Non-Hispanic	192	55.6	(46.3 - 64.8)	11,305	62.0	(58.9 - 65.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	4,963	50.5	(43.7 - 57.3)
Some College	*	*	*	2,665	58.3	(50.4 - 66.2)
≥ College Degree	*	*	*	4,277	59.2	(52.4 - 65.9)
<b>Income</b>						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,139	60.8	(53.1 - 68.4)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,295	56.5	(50.5 - 62.4)

<sup>1</sup> Defined as having had a digital rectal exam within the past 2 years among men aged 40 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 31a. Ever had a prostate specific antigen test among men aged 40 and older<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	198	6,900	49.2	(40.8 - 57.6)	11,557	2,822,600	67.6	(63.9 - 71.2)
<b>Gender</b>								
Male	198	6,900	49.2	(40.8 - 57.6)	11,557	2,822,600	67.6	(63.9 - 71.2)
Female	-	-	-	-	-	-	-	-
<b>Age (years)</b>								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,938	760,600	57.4	(50.6 - 64.3)
55-64	*	*	*	*	3,373	811,400	83.4	(78.8 - 88.1)
≥ 65	78	3,400	89.4	(82.7 - 96.0)	4,215	905,400	90.4	(87.1 - 93.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	191	6,600	49.5	(40.8 - 58.2)	10,950	1,965,400	67.0	(63.9 - 70.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	*	*	*	*	4,775	829,200	59.0	(52.4 - 65.6)
Some College	*	*	*	*	2,584	535,000	61.9	(53.7 - 70.2)
≥ College Degree	*	*	*	*	4,176	1,455,700	76.8	(72.5 - 81.1)
<b>Income</b>								
≤ \$24,999	*	*	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	*	3,048	444,100	64.1	(56.0 - 72.3)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,188	1,211,100	72.3	(67.6 - 77.0)

<sup>1</sup> Defined as having ever had a prostate specific antigen test among men aged 40 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of men aged 40 and older who have ever had a prostate specific antigen test (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 31b. Ever had a prostate specific antigen test among men aged 40 and older<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	198	48.3	(39.6 - 57.0)	11,557	67.4	(63.7 - 71.1)
<b>Gender</b>						
Male	198	48.3	(39.6 - 57.0)	11,557	67.4	(63.7 - 71.1)
Female	-	-	-	-	-	-
<b>Race/Ethnicity</b>						
White Non-Hispanic	191	48.6	(39.7 - 57.5)	10,950	66.6	(63.4 - 69.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	4,775	59.7	(52.9 - 66.4)
Some College	*	*	*	2,584	61.4	(53.0 - 69.9)
≥ College Degree	*	*	*	4,176	76.3	(71.9 - 80.7)
<b>Income</b>						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,048	64.2	(56.1 - 72.4)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,188	71.5	(66.6 - 76.3)

<sup>1</sup> Defined as having ever had a prostate specific antigen test among men aged 40 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 32a. Had prostate specific antigen test within the past 2 years among men aged 40 and older<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	195	5,900	42.5	(34.5 - 50.6)	11,437	2,493,800	59.9	(56.0 - 63.9)
<b>Gender</b>								
Male	195	5,900	42.5	(34.5 - 50.6)	11,437	2,493,800	59.9	(56.0 - 63.9)
Female	-	-	-	-	-	-	-	-
<b>Age (years)</b>								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,917	659,400	49.9	(42.8 - 57.0)
55-64	*	*	*	*	3,326	719,700	74.5	(69.4 - 79.7)
≥ 65	76	3,100	82.7	(74.2 - 91.2)	4,166	830,000	83.3	(78.4 - 88.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	188	5,700	43.4	(35.1 - 51.6)	10,836	1,762,200	60.4	(57.3 - 63.5)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	*	*	*	*	4,727	734,200	52.4	(45.7 - 59.1)
Some College	*	*	*	*	2,558	483,800	56.3	(48.2 - 64.4)
≥ College Degree	*	*	*	*	4,133	1,273,200	67.4	(61.6 - 73.3)
<b>Income</b>								
≤ \$24,999	*	*	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	*	3,021	410,800	59.6	(51.5 - 67.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,170	1,080,100	64.6	(59.1 - 70.0)

<sup>1</sup> Defined as having had a prostate specific antigen test within the past 2 years among men aged 40 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of men aged 40 and older who have had a prostate specific antigen test within the past 2 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 32b. Had prostate specific antigen test within the past 2 years among men aged 40 and older<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	195	42.1	(33.8 - 50.3)	11,437	59.8	(55.8 - 63.9)
<b>Gender</b>						
Male	195	42.1	(33.8 - 50.3)	11,437	59.8	(55.8 - 63.9)
Female	-	-	-	-	-	-
<b>Race/Ethnicity</b>						
White Non-Hispanic	188	42.9	(34.4 - 51.4)	10,836	60.1	(56.9 - 63.4)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	4,727	52.9	(45.9 - 59.9)
Some College	*	*	*	2,558	56.0	(47.7 - 64.2)
≥ College Degree	*	*	*	4,133	67.1	(61.2 - 73.1)
<b>Income</b>						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,021	59.8	(51.7 - 67.9)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,170	63.9	(58.3 - 69.6)

<sup>1</sup> Defined as having had a prostate specific antigen test within the past 2 years among men aged 40 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 33a. Prostate cancer among men aged 40 and older<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	206	500	3.4	(1.1 - 5.7)	12,133	178,200	4.1	(3.0 - 5.2)
<b>Gender</b>								
Male	206	500	3.4	(1.1 - 5.7)	12,133	178,200	4.1	(3.0 - 5.2)
Female	-	-	-	-	-	-	-	-
<b>Age (years)</b>								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	3,120	6,100	0.4	(0.0 - 0.9)
55-64	*	*	*	*	3,532	31,600	3.1	(1.5 - 4.8)
≥ 65	*	*	*	*	4,388	140,400	13.6	(9.7 - 17.4)
<b>Race/Ethnicity</b>								
White Non-Hispanic	198	500	3.6	(1.2 - 6.0)	11,502	143,700	4.7	(3.5 - 5.8)
Black Non-Hispanic	*	*	*	*	184	23,500	5.1	(0.0 - 11.0)
Hispanic	*	*	*	*	197	7,700	1.6	(0.0 - 3.7)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	*	*	*	*	5,065	46,300	3.1	(2.2 - 4.1)
Some College	*	*	*	*	2,709	39,200	4.4	(1.3 - 7.5)
≥ College Degree	*	*	*	*	4,336	92,500	4.7	(3.0 - 6.5)
<b>Income</b>								
≤ \$24,999	*	*	*	*	2,597	35,800	4.1	(2.2 - 6.0)
\$25,000-\$49,999	*	*	*	*	3,196	29,600	4.1	(2.9 - 5.4)
\$50,000-\$74,999	*	*	*	*	2,007	20,300	2.8	(1.2 - 4.4)
≥ \$75,000	*	*	*	*	3,322	56,500	3.2	(1.7 - 4.8)

<sup>1</sup> Defined as having ever been told by a health professional that respondent had prostate cancer among men aged 40 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of men aged 40 and older who have ever been told they had prostate cancer (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 33b. Prostate cancer among men aged 40 and older<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	206	3.7	(1.2 - 6.1)	12,133	4.5	(3.2 - 5.7)
<b>Gender</b>						
Male	206	3.7	(1.2 - 6.1)	12,133	4.5	(3.2 - 5.7)
Female	-	-	-	-	-	-
<b>Race/Ethnicity</b>						
White Non-Hispanic	198	3.8	(1.3 - 6.4)	11,502	5.1	(3.8 - 6.4)
Black Non-Hispanic	*	*	*	184	5.7	(0.0 - 12.8)
Hispanic	*	*	*	197	1.8	(0.0 - 4.4)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	5,065	3.3	(2.3 - 4.3)
Some College	*	*	*	2,709	5.0	(1.1 - 9.0)
≥ College Degree	*	*	*	4,336	5.2	(3.1 - 7.2)
<b>Income</b>						
≤ \$24,999	*	*	*	2,597	4.5	(2.2 - 6.7)
\$25,000-\$49,999	*	*	*	3,196	4.3	(3.0 - 5.6)
\$50,000-\$74,999	*	*	*	2,007	3.0	(1.1 - 5.0)
≥ \$75,000	*	*	*	3,322	3.5	(1.6 - 5.4)

<sup>1</sup> Defined as having ever been told by a health professional that respondent had prostate cancer among men aged 40 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 34a. Home blood stool test ever used among adults aged 50 and older<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	456	8,400	40.2	(35.2 - 45.2)	25,322	2,288,500	38.0	(35.5 - 40.4)
<b>Gender</b>								
Male	167	3,600	39.3	(31.2 - 47.4)	9,571	1,081,900	38.9	(35.1 - 42.6)
Female	289	4,800	40.9	(34.6 - 47.1)	15,751	1,206,600	37.2	(34.0 - 40.4)
<b>Age (years)</b>								
50-64	227	3,600	30.4	(23.8 - 37.0)	12,834	1,192,000	34.0	(30.8 - 37.2)
≥ 65	229	4,800	52.7	(45.7 - 59.8)	12,472	1,093,400	43.6	(39.8 - 47.4)
<b>Race/Ethnicity</b>								
White Non-Hispanic	446	8,300	40.9	(35.8 - 46.0)	24,213	1,754,500	39.1	(37.1 - 41.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	238	4,200	38.7	(31.9 - 45.6)	11,146	823,500	34.9	(30.5 - 39.2)
Some College	*	*	*	*	6,261	480,300	37.0	(32.6 - 41.4)
≥ College Degree	115	2,400	44.5	(34.6 - 54.4)	7,851	973,700	41.6	(37.8 - 45.3)
<b>Income</b>								
≤ \$24,999	150	2,300	41.9	(33.2 - 50.5)	7,222	505,000	35.4	(29.7 - 41.2)
\$25,000-\$49,999	140	2,400	36.5	(27.8 - 45.1)	6,609	467,600	40.3	(35.2 - 45.4)
\$50,000-\$74,999	*	*	*	*	3,500	313,000	38.2	(32.6 - 43.9)
≥ \$75,000	*	*	*	*	4,647	657,800	37.3	(33.0 - 41.6)

<sup>1</sup> Defined as having ever used a home blood stool test among adults aged 50 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 50 and older who have ever used a home blood stool test (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 34b. Home blood stool test ever used among adults aged 50 and older<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	456	40.1	(35.0 - 45.2)	25,322	38.1	(35.5 - 40.8)
<b>Gender</b>						
Male	167	40.6	(32.4 - 48.8)	9,571	39.4	(35.3 - 43.5)
Female	289	39.7	(33.4 - 45.9)	15,751	36.9	(33.5 - 40.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	446	40.7	(35.6 - 45.9)	24,213	38.9	(36.7 - 41.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	238	38.3	(31.3 - 45.3)	11,146	35.1	(30.3 - 39.9)
Some College	*	*	*	6,261	36.6	(32.0 - 41.3)
≥ College Degree	*	*	*	7,851	41.9	(37.9 - 46.0)
<b>Income</b>						
≤ \$24,999	150	41.1	(32.3 - 50.0)	7,222	35.9	(29.5 - 42.4)
\$25,000-\$49,999	140	36.7	(27.9 - 45.5)	6,609	40.2	(34.8 - 45.6)
\$50,000-\$74,999	*	*	*	3,500	38.1	(31.9 - 44.3)
≥ \$75,000	*	*	*	4,647	37.5	(32.9 - 42.2)

<sup>1</sup> Defined as having ever used a home blood stool test among adults aged 50 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 35a. Home blood stool test used within the past year among adults aged 50 and older<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	446	2,600	12.8	(9.4 - 16.2)	25,074	863,300	14.4	(12.6 - 16.2)
<b>Gender</b>								
Male	166	1,300	14.2	(8.5 - 19.9)	9,508	444,400	16.0	(13.2 - 18.9)
Female	280	1,300	11.7	(7.5 - 15.8)	15,566	418,900	13.1	(10.8 - 15.4)
<b>Age (years)</b>								
50-64	227	1,100	9.6	(5.5 - 13.7)	12,759	454,400	13.0	(10.6 - 15.5)
≥ 65	219	1,500	17.1	(11.4 - 22.8)	12,299	406,400	16.4	(13.7 - 19.0)
<b>Race/Ethnicity</b>								
White Non-Hispanic	436	2,500	12.7	(9.2 - 16.1)	23,971	634,300	14.3	(12.8 - 15.7)
Black Non-Hispanic	*	*	*	*	396	136,000	18.6	(9.4 - 27.8)
Hispanic	*	*	*	*	349	53,400	9.9	(2.5 - 17.2)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	233	1,400	13.1	(8.3 - 17.9)	11,026	288,600	12.3	(9.4 - 15.3)
Some College	100	700	14.5	(6.4 - 22.6)	6,208	212,500	16.4	(12.5 - 20.4)
≥ College Degree	111	600	11.0	(5.3 - 16.8)	7,778	352,400	15.2	(12.5 - 17.9)
<b>Income</b>								
≤ \$24,999	146	900	17.2	(10.9 - 23.5)	7,157	196,300	13.9	(10.1 - 17.6)
\$25,000-\$49,999	138	600	9.0	(3.7 - 14.4)	6,562	221,500	19.2	(14.1 - 24.3)
\$50,000-\$74,999	*	*	*	*	3,469	111,900	13.7	(10.1 - 17.4)
≥ \$75,000	*	*	*	*	4,623	205,100	11.7	(8.8 - 14.6)

<sup>1</sup> Defined as having used a home blood stool test within the past year among adults aged 50 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 50 and older who have used a home blood stool test within the past year (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 35b. Home blood stool test used within the past year among adults aged 50 and older<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	446	13.0	(9.5 - 16.6)	25,074	14.6	(12.6 - 16.7)
<b>Gender</b>						
Male	166	14.8	(8.8 - 20.8)	9,508	16.4	(13.2 - 19.6)
Female	280	11.4	(7.3 - 15.4)	15,566	12.9	(10.4 - 15.4)
<b>Race/Ethnicity</b>						
White Non-Hispanic	436	12.8	(9.2 - 16.4)	23,971	14.2	(12.7 - 15.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	349	10.2	(1.7 - 18.7)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	233	13.6	(8.4 - 18.8)	11,026	12.7	(9.2 - 16.1)
Some College	100	14.6	(6.5 - 22.7)	6,208	16.6	(12.3 - 20.9)
≥ College Degree	111	10.8	(5.1 - 16.5)	7,778	15.3	(12.4 - 18.3)
<b>Income</b>						
≤ \$24,999	146	16.5	(10.2 - 22.7)	7,157	14.3	(9.8 - 18.7)
\$25,000-\$49,999	138	10.2	(4.1 - 16.4)	6,562	19.3	(13.8 - 24.8)
\$50,000-\$74,999	*	*	*	3,469	13.8	(9.8 - 17.7)
≥ \$75,000	*	*	*	4,623	11.8	(8.6 - 15.0)

<sup>1</sup> Defined as having used a home blood stool test within the past year among adults aged 50 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 36a. Home blood stool test used within the past 2 years among adults aged 50 and older<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	446	4,200	20.5	(16.3 - 24.6)	25,074	1,218,000	20.4	(18.3 - 22.5)
<b>Gender</b>								
Male	166	2,100	23.5	(16.6 - 30.5)	9,508	593,100	21.4	(18.3 - 24.5)
Female	280	2,100	18.1	(13.1 - 23.1)	15,566	624,900	19.5	(16.6 - 22.4)
<b>Age (years)</b>								
50-64	227	1,700	14.4	(9.3 - 19.4)	12,759	629,900	18.1	(15.4 - 20.7)
≥ 65	219	2,500	28.7	(22.0 - 35.4)	12,299	585,500	23.6	(20.2 - 27.0)
<b>Race/Ethnicity</b>								
White Non-Hispanic	436	4,100	20.6	(16.4 - 24.8)	23,971	877,500	19.7	(18.2 - 21.3)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	233	1,900	18.1	(12.8 - 23.4)	11,026	454,600	19.4	(15.6 - 23.3)
Some College	100	1,100	23.9	(14.0 - 33.9)	6,208	262,200	20.3	(16.3 - 24.2)
≥ College Degree	111	1,200	22.7	(14.3 - 31.2)	7,778	491,400	21.2	(18.1 - 24.2)
<b>Income</b>								
≤ \$24,999	146	1,500	27.3	(19.3 - 35.2)	7,157	263,900	18.6	(14.6 - 22.7)
\$25,000-\$49,999	138	1,000	15.3	(9.0 - 21.7)	6,562	285,900	24.8	(19.6 - 30.0)
\$50,000-\$74,999	*	*	*	*	3,469	167,500	20.6	(15.6 - 25.5)
≥ \$75,000	*	*	*	*	4,623	294,900	16.8	(13.7 - 20.0)

<sup>1</sup> Defined as having used a home blood stool test within the past 2 years among adults aged 50 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 50 and older who have used a home blood stool test within the past 2 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 36b. Home blood stool test used within the past 2 years among adults aged 50 and older<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	446	21.0	(16.7 - 25.4)	25,074	20.5	(18.2 - 22.8)
<b>Gender</b>						
Male	166	24.7	(17.4 - 31.9)	9,508	21.7	(18.3 - 25.1)
Female	280	17.6	(12.6 - 22.5)	15,566	19.4	(16.3 - 22.5)
<b>Race/Ethnicity</b>						
White Non-Hispanic	436	21.1	(16.7 - 25.5)	23,971	19.7	(18.0 - 21.3)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	233	18.6	(12.9 - 24.3)	11,026	19.5	(15.4 - 23.7)
Some College	*	*	*	6,208	20.4	(16.1 - 24.7)
≥ College Degree	111	23.2	(14.5 - 32.0)	7,778	21.4	(18.1 - 24.7)
<b>Income</b>						
≤ \$24,999	146	27.0	(19.0 - 35.1)	7,157	18.9	(14.2 - 23.6)
\$25,000-\$49,999	138	16.3	(9.4 - 23.2)	6,562	24.8	(19.3 - 30.3)
\$50,000-\$74,999	*	*	*	3,469	20.6	(15.3 - 25.9)
≥ \$75,000	*	*	*	4,623	16.9	(13.4 - 20.4)

<sup>1</sup> Defined as having used a home blood stool test within the past 2 years among adults aged 50 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 37a. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	456	14,500	68.9	(64.0 - 73.8)	25,357	4,019,100	66.6	(64.1 - 69.1)
<b>Gender</b>								
Male	165	6,300	69.3	(61.2 - 77.3)	9,582	1,894,100	68.0	(64.2 - 71.7)
Female	291	8,100	68.6	(62.5 - 74.7)	15,775	2,125,000	65.5	(62.1 - 68.8)
<b>Age (years)</b>								
50-54	*	*	*	*	4,179	718,900	50.6	(44.6 - 56.5)
55-64	150	4,800	66.8	(58.4 - 75.1)	8,695	1,493,500	71.0	(67.3 - 74.8)
≥ 65	228	7,300	80.3	(74.9 - 85.8)	12,466	1,802,000	72.1	(68.6 - 75.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	446	14,100	69.3	(64.4 - 74.3)	24,248	3,093,900	68.8	(66.8 - 70.8)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	236	7,300	68.3	(61.6 - 75.1)	11,121	1,445,100	61.2	(56.7 - 65.8)
Some College	*	*	*	*	6,287	835,300	64.3	(59.7 - 69.0)
≥ College Degree	117	4,100	74.2	(65.5 - 82.9)	7,884	1,732,400	73.7	(70.0 - 77.4)
<b>Income</b>								
≤ \$24,999	149	3,700	67.5	(58.9 - 76.1)	7,220	892,600	62.7	(56.8 - 68.6)
\$25,000-\$49,999	140	4,900	72.8	(64.8 - 80.8)	6,619	697,300	60.1	(54.6 - 65.7)
\$50,000-\$74,999	*	*	*	*	3,514	613,700	74.2	(68.9 - 79.6)
≥ \$75,000	*	*	*	*	4,665	1,294,400	73.3	(69.2 - 77.5)

<sup>1</sup> Defined as having ever had a sigmoidoscopy or colonoscopy among adults aged 50 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 50 and older who have ever had a sigmoidoscopy or colonoscopy (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 37b. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	456	69.3	(64.3 - 74.2)	25,357	66.6	(63.9 - 69.3)
<b>Gender</b>						
Male	165	70.6	(62.7 - 78.5)	9,582	68.5	(64.5 - 72.5)
Female	291	68.0	(61.8 - 74.2)	15,775	64.8	(61.2 - 68.4)
<b>Race/Ethnicity</b>						
White Non-Hispanic	446	69.7	(64.7 - 74.7)	24,248	68.7	(66.5 - 70.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	236	68.8	(62.0 - 75.6)	11,121	61.3	(56.4 - 66.2)
Some College	*	*	*	6,287	63.8	(58.8 - 68.8)
≥ College Degree	117	74.4	(65.5 - 83.2)	7,884	74.0	(70.2 - 77.9)
<b>Income</b>						
≤ \$24,999	149	67.5	(58.8 - 76.2)	7,220	63.0	(56.7 - 69.3)
\$25,000-\$49,999	140	73.6	(65.6 - 81.5)	6,619	59.7	(53.8 - 65.6)
\$50,000-\$74,999	*	*	*	3,514	74.4	(68.3 - 80.5)
≥ \$75,000	*	*	*	4,665	73.6	(69.3 - 78.0)

<sup>1</sup> Defined as having ever had a sigmoidoscopy or colonoscopy among adults aged 50 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 38a. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	454	13,800	66.1	(61.1 - 71.1)	25,169	3,855,800	64.6	(62.1 - 67.1)
<b>Gender</b>								
Male	165	6,100	67.2	(59.1 - 75.4)	9,521	1,835,400	66.4	(62.6 - 70.1)
Female	289	7,600	65.2	(59.0 - 71.4)	15,648	2,020,400	63.1	(59.8 - 66.4)
<b>Age (years)</b>								
50-54	*	*	*	*	4,169	700,500	49.3	(43.3 - 55.3)
55-64	150	4,500	63.5	(55.0 - 72.1)	8,652	1,455,100	69.4	(65.7 - 73.2)
≥ 65	226	6,900	77.1	(71.3 - 82.9)	12,331	1,695,700	69.5	(65.9 - 73.1)
<b>Race/Ethnicity</b>								
White Non-Hispanic	444	13,400	66.4	(61.4 - 71.5)	24,069	2,981,500	66.6	(64.6 - 68.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	236	6,900	64.9	(58.1 - 71.7)	11,017	1,360,400	59.0	(54.5 - 63.5)
Some College	*	*	*	*	6,251	807,100	62.4	(57.8 - 67.1)
≥ College Degree	116	4,000	71.2	(62.1 - 80.3)	7,841	1,682,500	71.7	(68.0 - 75.4)
<b>Income</b>								
≤ \$24,999	147	3,300	61.2	(52.4 - 70.1)	7,163	858,900	61.0	(55.1 - 67.0)
\$25,000-\$49,999	140	4,900	72.8	(64.8 - 80.8)	6,584	664,900	57.7	(52.2 - 63.2)
\$50,000-\$74,999	*	*	*	*	3,501	602,600	73.1	(67.7 - 78.5)
≥ \$75,000	*	*	*	*	4,655	1,260,400	71.5	(67.3 - 75.7)

<sup>1</sup> Defined as having had a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 50 and older who have had a sigmoidoscopy or colonoscopy within the past 10 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 38b. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	454	66.7	(61.6 - 71.7)	25,169	64.6	(61.9 - 67.3)
<b>Gender</b>						
Male	165	68.9	(60.9 - 76.9)	9,521	66.9	(62.9 - 70.9)
Female	289	64.7	(58.4 - 71.0)	15,648	62.4	(58.9 - 66.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	444	67.0	(62.0 - 72.1)	24,069	66.5	(64.3 - 68.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	236	65.6	(58.7 - 72.5)	11,017	59.0	(54.1 - 63.9)
Some College	*	*	*	6,251	61.9	(56.9 - 66.9)
≥ College Degree	116	71.7	(62.5 - 80.8)	7,841	72.1	(68.2 - 75.9)
<b>Income</b>						
≤ \$24,999	147	61.6	(52.7 - 70.6)	7,163	61.3	(54.9 - 67.6)
\$25,000-\$49,999	140	73.6	(65.6 - 81.5)	6,584	57.2	(51.4 - 63.0)
\$50,000-\$74,999	*	*	*	3,501	73.3	(67.2 - 79.4)
≥ \$75,000	*	*	*	4,655	71.8	(67.4 - 76.2)

<sup>1</sup> Defined as having had a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 39a. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	445	14,600	71.3	(66.5 - 76.2)	24,792	4,049,200	69.1	(66.7 - 71.6)
<b>Gender</b>								
Male	162	6,500	72.5	(64.5 - 80.6)	9,418	1,945,200	70.7	(67.0 - 74.3)
Female	283	8,100	70.4	(64.4 - 76.4)	15,374	2,104,100	67.8	(64.6 - 71.1)
<b>Age (years)</b>								
50-54	*	*	*	*	4,132	751,200	54.1	(48.0 - 60.1)
55-64	146	5,000	72.9	(65.0 - 80.7)	8,366	1,487,600	73.7	(70.0 - 77.4)
≥ 65	221	7,100	80.4	(74.8 - 85.9)	12,294	1,810,500	74.0	(70.6 - 77.4)
<b>Race/Ethnicity</b>								
White Non-Hispanic	435	14,200	71.7	(66.8 - 76.6)	23,726	3,091,400	70.4	(68.4 - 72.4)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	232	7,300	69.3	(62.6 - 76.0)	10,872	1,461,600	64.2	(59.8 - 68.7)
Some College	*	*	*	*	6,170	873,300	68.4	(63.9 - 72.8)
≥ College Degree	112	4,100	77.5	(68.9 - 86.2)	7,708	1,709,000	74.6	(70.9 - 78.3)
<b>Income</b>								
≤ \$24,999	145	3,600	67.9	(59.3 - 76.6)	7,092	923,300	66.0	(60.1 - 71.9)
\$25,000-\$49,999	140	5,000	74.3	(66.4 - 82.2)	6,526	730,000	64.2	(58.9 - 69.6)
\$50,000-\$74,999	*	*	*	*	3,467	621,500	76.6	(71.4 - 81.9)
≥ \$75,000	*	*	*	*	4,615	1,292,500	73.8	(69.8 - 77.9)

<sup>1</sup> Defined as having used a home blood stool test within the past year or a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 years and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 50 and older who have used a home blood stool test in the past year or sigmoidoscopy or colonoscopy within the past 10 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 39b. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	445	71.7	(66.8 - 76.6)	24,792	69.3	(66.7 - 71.9)
<b>Gender</b>						
Male	162	73.7	(65.8 - 81.6)	9,418	71.5	(67.6 - 75.4)
Female	283	69.9	(63.7 - 76.0)	15,374	67.2	(63.7 - 70.7)
<b>Race/Ethnicity</b>						
White Non-Hispanic	435	72.1	(67.1 - 77.0)	23,726	70.3	(68.1 - 72.4)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	232	69.8	(63.1 - 76.6)	10,872	64.5	(59.8 - 69.3)
Some College	*	*	*	6,170	68.2	(63.4 - 72.9)
≥ College Degree	112	77.5	(68.6 - 86.3)	7,708	75.0	(71.2 - 78.8)
<b>Income</b>						
≤ \$24,999	145	68.3	(59.6 - 77.0)	7,092	66.4	(60.2 - 72.6)
\$25,000-\$49,999	140	74.9	(67.1 - 82.8)	6,526	63.8	(58.2 - 69.5)
\$50,000-\$74,999	*	*	*	3,467	76.7	(70.6 - 82.7)
≥ \$75,000	*	*	*	4,615	74.4	(70.2 - 78.6)

<sup>1</sup> Defined as having used a home blood stool test within the past year or a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 years and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 40a. High blood pressure among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	657	15,400	31.0	(26.4 - 35.6)	38,125	3,974,800	26.8	(24.9 - 28.7)
<b>Gender</b>								
Male	240	7,500	29.5	(22.0 - 37.0)	14,611	1,960,500	27.7	(24.7 - 30.7)
Female	417	7,900	32.6	(27.5 - 37.7)	23,514	2,014,300	26.0	(23.6 - 28.3)
<b>Age (years)</b>								
18-34	*	*	*	*	3,935	312,200	7.0	(4.2 - 9.8)
35-44	83	1,800	20.4	(10.7 - 30.0)	4,984	426,200	14.8	(11.6 - 18.1)
45-54	135	2,900	30.7	(21.5 - 39.9)	7,825	862,400	30.3	(25.9 - 34.8)
55-64	153	3,300	45.1	(36.5 - 53.7)	8,765	890,800	41.6	(37.6 - 45.7)
≥ 65	228	6,200	68.0	(61.4 - 74.6)	12,616	1,483,200	58.5	(54.7 - 62.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	635	15,100	32.6	(27.9 - 37.4)	35,814	2,650,900	28.4	(26.9 - 30.0)
Black Non-Hispanic	*	*	*	*	736	558,500	26.2	(19.6 - 32.8)
Hispanic	*	*	*	*	863	538,100	23.8	(17.0 - 30.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	311	8,100	36.1	(28.5 - 43.8)	15,688	1,851,200	33.1	(29.4 - 36.9)
Some College	161	3,600	27.6	(19.5 - 35.7)	9,906	851,100	24.1	(20.9 - 27.3)
≥ College Degree	183	3,600	25.8	(18.2 - 33.3)	12,452	1,253,400	22.3	(19.7 - 24.9)
<b>Income</b>								
≤ \$24,999	190	4,200	34.9	(25.3 - 44.6)	9,640	1,153,000	32.8	(27.6 - 38.1)
\$25,000-\$49,999	184	4,600	34.3	(25.7 - 42.9)	9,657	807,800	27.9	(23.7 - 32.1)
\$50,000-\$74,999	*	*	*	*	5,773	507,600	25.8	(21.4 - 30.1)
≥ \$75,000	112	2,300	20.7	(12.6 - 28.8)	8,560	942,500	21.0	(18.4 - 23.6)

<sup>1</sup> Defined as having ever been told by a health professional that respondent has high blood pressure

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have ever been told by a health professional that they have high blood pressure (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 40b. High blood pressure among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	657	29.5	(24.9 - 34.1)	38,125	25.7	(23.8 - 27.5)
<b>Gender</b>						
Male	240	30.0	(22.5 - 37.6)	14,611	27.7	(24.7 - 30.7)
Female	417	28.9	(23.7 - 34.0)	23,514	23.8	(21.5 - 26.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	635	31.0	(26.2 - 35.7)	35,814	27.2	(25.6 - 28.7)
Black Non-Hispanic	*	*	*	736	24.9	(18.4 - 31.3)
Hispanic	*	*	*	863	23.3	(16.6 - 30.0)
Other Non-Hispanic	*	*	*	712	19.3	(9.5 - 29.1)
<b>Education</b>						
≤ High School	311	34.6	(27.0 - 42.3)	15,688	32.1	(28.4 - 35.8)
Some College	161	26.8	(18.6 - 35.1)	9,906	22.8	(19.7 - 25.9)
≥ College Degree	183	23.8	(16.5 - 31.2)	12,452	21.2	(18.6 - 23.8)
<b>Income</b>						
≤ \$24,999	190	34.2	(24.5 - 44.0)	9,640	31.8	(26.7 - 37.0)
\$25,000-\$49,999	184	33.3	(24.6 - 42.1)	9,657	26.9	(22.8 - 30.9)
\$50,000-\$74,999	*	*	*	5,773	24.8	(20.4 - 29.2)
≥ \$75,000	112	19.2	(11.4 - 26.9)	8,560	19.9	(17.3 - 22.5)

<sup>1</sup> Defined as having ever been told by a health professional that respondent has high blood pressure

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 41a. Blood pressure medication among adults with high blood pressure<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	295	11,400	74.0	(65.9 - 82.2)	15,288	3,145,800	79.2	(76.0 - 82.4)
<b>Gender</b>								
Male	*	*	*	*	6,080	1,432,100	73.1	(67.6 - 78.5)
Female	183	6,300	80.3	(73.0 - 87.7)	9,208	1,713,700	85.2	(82.1 - 88.2)
<b>Age (years)</b>								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,315	645,500	74.9	(66.8 - 82.9)
55-64	73	2,900	87.6	(79.5 - 95.6)	4,045	782,300	87.8	(84.5 - 91.1)
≥ 65	156	5,400	87.9	(82.0 - 93.8)	7,675	1,383,400	93.4	(91.1 - 95.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	290	11,100	73.5	(65.3 - 81.8)	14,442	2,117,300	79.9	(77.5 - 82.4)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	*	*	*	*	7,354	1,482,800	80.2	(74.7 - 85.6)
Some College	*	*	*	*	3,868	632,000	74.3	(67.8 - 80.7)
≥ College Degree	*	*	*	*	4,035	1,013,200	80.9	(76.8 - 84.9)
<b>Income</b>								
≤ \$24,999	*	*	*	*	4,898	948,200	82.3	(75.7 - 88.9)
\$25,000-\$49,999	*	*	*	*	4,088	668,300	82.8	(77.6 - 88.0)
\$50,000-\$74,999	*	*	*	*	1,971	391,500	77.2	(67.7 - 86.6)
≥ \$75,000	*	*	*	*	2,397	683,700	72.5	(66.1 - 78.9)

<sup>1</sup> Defined as currently taking medicine for high blood pressure among adults with high blood pressure

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults with high blood pressure who are taking medicine for high blood pressure (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 41b. Blood pressure medication among adults with high blood pressure<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	295	71.9	(63.3 - 80.4)	15,288	78.5	(75.2 - 81.7)
<b>Gender</b>						
Male	*	*	*	6,080	73.3	(67.9 - 78.7)
Female	183	77.3	(68.4 - 86.2)	9,208	84.0	(80.7 - 87.2)
<b>Race/Ethnicity</b>						
White Non-Hispanic	290	71.3	(62.6 - 79.9)	14,442	79.0	(76.5 - 81.6)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	7,354	79.7	(74.3 - 85.2)
Some College	*	*	*	3,868	72.5	(65.7 - 79.3)
≥ College Degree	*	*	*	4,035	80.4	(76.1 - 84.6)
<b>Income</b>						
≤ \$24,999	*	*	*	4,898	81.6	(75.0 - 88.3)
\$25,000-\$49,999	*	*	*	4,088	82.1	(76.7 - 87.5)
\$50,000-\$74,999	*	*	*	1,971	76.4	(66.7 - 86.1)
≥ \$75,000	*	*	*	2,397	71.6	(64.8 - 78.3)

<sup>1</sup> Defined as currently taking medicine for high blood pressure among adults with high blood pressure

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 42a. Ever had blood cholesterol checked among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	648	37,200	76.3	(70.3 - 82.4)	37,450	11,496,600	80.3	(77.7 - 83.0)
<b>Gender</b>								
Male	*	*	*	*	14,389	5,288,400	77.7	(73.3 - 82.0)
Female	409	18,300	78.3	(72.4 - 84.1)	23,061	6,208,200	82.7	(79.6 - 85.8)
<b>Age (years)</b>								
18-34	*	*	*	*	3,684	2,296,400	55.3	(48.7 - 61.9)
35-44	*	*	*	*	4,883	2,264,200	81.6	(76.6 - 86.5)
45-54	135	8,200	87.1	(80.8 - 93.4)	7,731	2,539,400	91.0	(87.0 - 94.9)
55-64	152	6,900	96.9	(94.4 - 99.4)	8,718	2,014,900	95.4	(93.5 - 97.2)
≥ 65	226	8,700	96.0	(93.7 - 98.3)	12,434	2,381,700	95.9	(94.3 - 97.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	628	35,300	77.2	(71.3 - 83.1)	35,198	7,832,600	86.3	(84.8 - 87.9)
Black Non-Hispanic	*	*	*	*	720	1,588,900	77.8	(68.4 - 87.2)
Hispanic	*	*	*	*	842	1,224,600	56.8	(46.8 - 66.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	*	*	*	*	15,342	3,795,700	70.6	(65.6 - 75.6)
Some College	160	10,700	82.5	(72.9 - 92.1)	9,732	2,599,800	77.4	(71.7 - 83.2)
≥ College Degree	182	11,900	84.8	(77.0 - 92.5)	12,296	5,008,300	91.3	(88.9 - 93.6)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,437	2,277,500	68.4	(61.5 - 75.3)
\$25,000-\$49,999	178	10,800	83.7	(74.8 - 92.6)	9,492	2,053,000	71.9	(64.5 - 79.3)
\$50,000-\$74,999	*	*	*	*	5,721	1,736,100	90.0	(87.7 - 92.3)
≥ \$75,000	*	*	*	*	8,460	4,019,300	92.2	(90.8 - 93.6)

<sup>1</sup> Defined as ever having blood cholesterol checked

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have ever had blood cholesterol checked (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 42b. Ever had blood cholesterol checked among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	648	75.3	(69.2 - 81.3)	37,450	79.9	(77.3 - 82.5)
<b>Gender</b>						
Male	239	75.2	(65.3 - 85.0)	14,389	78.0	(73.9 - 82.1)
Female	409	75.4	(68.7 - 82.1)	23,061	81.6	(78.4 - 84.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	628	76.0	(70.0 - 81.9)	35,198	85.6	(84.0 - 87.2)
Black Non-Hispanic	*	*	*	720	77.5	(68.2 - 86.7)
Hispanic	*	*	*	842	57.2	(47.4 - 67.0)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	15,342	70.2	(65.2 - 75.1)
Some College	160	81.7	(71.8 - 91.6)	9,732	76.7	(71.1 - 82.3)
≥ College Degree	182	83.5	(75.4 - 91.7)	12,296	90.9	(88.6 - 93.2)
<b>Income</b>						
≤ \$24,999	*	*	*	9,437	68.1	(61.3 - 74.9)
\$25,000-\$49,999	178	82.7	(73.6 - 91.9)	9,492	71.6	(64.4 - 78.7)
\$50,000-\$74,999	*	*	*	5,721	89.6	(87.2 - 92.0)
≥ \$75,000	*	*	*	8,460	91.7	(90.1 - 93.2)

<sup>1</sup> Defined as ever having blood cholesterol checked

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 43a. Blood cholesterol checked within the past 5 years among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	644	34,500	71.0	(64.9 - 77.1)	37,201	11,035,200	77.8	(75.1 - 80.4)
<b>Gender</b>								
Male	*	*	*	*	14,307	5,020,700	74.5	(70.2 - 78.9)
Female	406	17,600	75.8	(69.8 - 81.7)	22,894	6,014,600	80.7	(77.6 - 83.8)
<b>Age (years)</b>								
18-34	*	*	*	*	3,647	2,179,800	53.3	(46.7 - 59.9)
35-44	*	*	*	*	4,862	2,161,200	78.0	(72.8 - 83.2)
45-54	134	7,000	74.1	(65.1 - 83.1)	7,697	2,408,600	86.6	(82.4 - 90.8)
55-64	151	6,400	90.4	(85.5 - 95.3)	8,677	1,957,900	93.3	(91.4 - 95.3)
≥ 65	224	8,500	94.6	(91.8 - 97.4)	12,318	2,327,800	94.8	(93.1 - 96.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	625	32,600	71.6	(65.6 - 77.5)	34,960	7,486,600	83.4	(81.8 - 85.0)
Black Non-Hispanic	*	*	*	*	719	1,561,700	76.5	(67.1 - 85.9)
Hispanic	*	*	*	*	838	1,184,000	55.6	(45.6 - 65.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	*	*	*	*	15,215	3,612,500	68.3	(63.2 - 73.3)
Some College	*	*	*	*	9,673	2,524,200	75.4	(69.8 - 81.1)
≥ College Degree	181	11,200	79.8	(71.5 - 88.0)	12,236	4,806,500	88.1	(85.5 - 90.6)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,350	2,214,800	66.9	(60.1 - 73.8)
\$25,000-\$49,999	178	10,400	80.3	(71.1 - 89.5)	9,454	1,936,600	68.8	(61.5 - 76.1)
\$50,000-\$74,999	*	*	*	*	5,701	1,640,400	85.9	(82.4 - 89.5)
≥ \$75,000	*	*	*	*	8,435	3,872,400	89.2	(87.5 - 90.9)

<sup>1</sup> Defined as having had blood cholesterol checked within the past 5 years

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have had blood cholesterol checked within the past 5 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 43b. Blood cholesterol checked within the past 5 years among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	644	70.1	(63.9 - 76.2)	37,201	77.3	(74.7 - 79.9)
<b>Gender</b>						
Male	238	67.4	(57.6 - 77.3)	14,307	74.8	(70.7 - 79.0)
Female	406	72.9	(66.2 - 79.7)	22,894	79.5	(76.3 - 82.7)
<b>Race/Ethnicity</b>						
White Non-Hispanic	625	70.4	(64.4 - 76.5)	34,960	82.6	(81.0 - 84.3)
Black Non-Hispanic	*	*	*	719	76.2	(66.9 - 85.5)
Hispanic	*	*	*	838	55.9	(46.1 - 65.8)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	15,215	67.8	(62.9 - 72.8)
Some College	*	*	*	9,673	74.7	(69.1 - 80.2)
≥ College Degree	181	79.0	(70.5 - 87.5)	12,236	87.5	(85.0 - 90.1)
<b>Income</b>						
≤ \$24,999	*	*	*	9,350	66.6	(59.8 - 73.4)
\$25,000-\$49,999	178	79.4	(70.0 - 88.8)	9,454	68.5	(61.4 - 75.6)
\$50,000-\$74,999	*	*	*	5,701	85.4	(81.7 - 89.0)
≥ \$75,000	*	*	*	8,435	88.5	(86.7 - 90.3)

<sup>1</sup> Defined as having had blood cholesterol checked within the past 5 years

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 44a. Cardiovascular disease among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	655	4,500	9.1	(6.9 - 11.4)	37,810	1,135,000	7.7	(6.5 - 8.8)
<b>Gender</b>								
Male	239	2,200	8.6	(5.4 - 11.7)	14,477	664,600	9.4	(7.5 - 11.4)
Female	416	2,400	9.8	(6.7 - 12.8)	23,333	470,400	6.1	(4.8 - 7.4)
<b>Age (years)</b>								
18-34	*	*	*	*	3,930	100,800	2.3	(0.0 - 4.8)
35-44	*	*	*	*	4,971	103,400	3.6	(1.5 - 5.7)
45-54	*	*	*	*	7,768	131,300	4.6	(2.9 - 6.4)
55-64	151	900	11.9	(6.5 - 17.4)	8,699	224,200	10.6	(8.3 - 12.9)
≥ 65	228	2,300	25.2	(19.1 - 31.3)	12,442	575,300	23.0	(19.9 - 26.1)
<b>Race/Ethnicity</b>								
White Non-Hispanic	633	4,200	9.2	(6.9 - 11.4)	35,522	772,200	8.3	(7.4 - 9.2)
Black Non-Hispanic	*	*	*	*	728	140,700	6.6	(3.3 - 9.9)
Hispanic	*	*	*	*	854	163,300	7.3	(2.0 - 12.5)
Other Non-Hispanic	*	*	*	*	706	58,800	5.2	(0.9 - 9.5)
<b>Education</b>								
≤ High School	311	2,800	12.3	(8.5 - 16.1)	15,486	590,800	10.7	(8.2 - 13.2)
Some College	161	700	5.3	(2.4 - 8.1)	9,849	227,100	6.4	(4.7 - 8.1)
≥ College Degree	181	1,100	7.8	(3.6 - 12.0)	12,393	314,600	5.6	(4.3 - 6.9)
<b>Income</b>								
≤ \$24,999	190	1,900	16.4	(10.4 - 22.3)	9,505	448,100	12.9	(9.1 - 16.7)
\$25,000-\$49,999	184	1,200	8.9	(4.4 - 13.4)	9,571	179,300	6.2	(4.8 - 7.6)
\$50,000-\$74,999	*	*	*	*	5,751	166,900	8.5	(5.5 - 11.4)
≥ \$75,000	*	*	*	*	8,538	183,500	4.1	(2.9 - 5.3)

<sup>1</sup> Defined as having ever been told by a health professional that respondent had a heart attack or angina or a stroke

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults with cardiovascular disease (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 44b. Cardiovascular disease among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	655	8.9	(6.6 - 11.2)	37,810	7.6	(6.4 - 8.7)
<b>Gender</b>						
Male	239	8.8	(5.5 - 12.1)	14,477	9.7	(7.8 - 11.7)
Female	416	9.0	(5.8 - 12.2)	23,333	5.6	(4.3 - 6.9)
<b>Race/Ethnicity</b>						
White Non-Hispanic	633	9.0	(6.6 - 11.4)	35,522	8.2	(7.2 - 9.1)
Black Non-Hispanic	*	*	*	728	6.8	(3.2 - 10.4)
Hispanic	*	*	*	854	7.0	(2.0 - 12.0)
Other Non-Hispanic	*	*	*	706	5.5	(0.9 - 10.0)
<b>Education</b>						
≤ High School	311	11.9	(8.0 - 15.8)	15,486	10.5	(8.0 - 13.0)
Some College	161	5.1	(2.1 - 8.0)	9,849	6.4	(4.5 - 8.2)
≥ College Degree	181	7.9	(3.4 - 12.5)	12,393	5.5	(4.2 - 6.9)
<b>Income</b>						
≤ \$24,999	190	16.2	(10.1 - 22.2)	9,505	12.8	(9.0 - 16.6)
\$25,000-\$49,999	184	9.6	(4.6 - 14.7)	9,571	6.0	(4.7 - 7.4)
\$50,000-\$74,999	*	*	*	5,751	8.5	(5.4 - 11.6)
≥ \$75,000	*	*	*	8,538	4.0	(2.7 - 5.2)

<sup>1</sup> Defined as having ever been told by a health professional that respondent had a heart attack or angina or a stroke

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 45a. Coronary heart disease among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	656	3,900	7.9	(5.8 - 9.9)	37,838	929,400	6.3	(5.3 - 7.3)
<b>Gender</b>								
Male	239	2,000	7.8	(4.8 - 10.9)	14,479	598,600	8.5	(6.6 - 10.4)
Female	417	1,900	7.9	(5.1 - 10.6)	23,359	330,800	4.3	(3.4 - 5.2)
<b>Age (years)</b>								
18-34	*	*	*	*	3,932	63,400	1.4	(0.0 - 3.6)
35-44	*	*	*	*	4,973	95,300	3.3	(1.2 - 5.4)
45-54	*	*	*	*	7,778	108,300	3.8	(2.1 - 5.6)
55-64	152	800	10.9	(5.6 - 16.1)	8,703	188,600	8.9	(6.7 - 11.1)
≥ 65	228	1,900	21.2	(15.5 - 26.9)	12,452	473,800	18.9	(16.1 - 21.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	634	3,600	7.8	(5.7 - 9.9)	35,552	636,700	6.9	(6.0 - 7.7)
Black Non-Hispanic	*	*	*	*	728	113,100	5.3	(2.3 - 8.4)
Hispanic	*	*	*	*	853	128,000	5.7	(1.1 - 10.3)
Other Non-Hispanic	*	*	*	*	705	51,500	4.6	(0.5 - 8.7)
<b>Education</b>								
≤ High School	312	2,300	10.2	(6.8 - 13.6)	15,485	474,500	8.6	(6.3 - 10.8)
Some College	161	600	4.5	(1.8 - 7.2)	9,860	193,000	5.5	(3.9 - 7.1)
≥ College Degree	181	1,000	7.2	(3.0 - 11.3)	12,411	259,700	4.6	(3.4 - 5.8)
<b>Income</b>								
≤ \$24,999	190	1,600	13.5	(8.3 - 18.7)	9,507	372,300	10.7	(7.3 - 14.1)
\$25,000-\$49,999	184	1,000	7.7	(3.6 - 11.9)	9,585	136,300	4.7	(3.8 - 5.7)
\$50,000-\$74,999	*	*	*	*	5,755	154,600	7.8	(4.9 - 10.7)
≥ \$75,000	*	*	*	*	8,541	151,800	3.4	(2.3 - 4.5)

<sup>1</sup> Defined as having ever been told by a health professional that respondent had a heart attack or angina

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults with coronary heart disease (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 45b. Coronary heart disease among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	656	7.7	(5.5 - 9.8)	37,838	6.3	(5.2 - 7.3)
<b>Gender</b>						
Male	239	8.1	(4.9 - 11.2)	14,479	8.8	(6.9 - 10.8)
Female	417	7.2	(4.4 - 10.1)	23,359	3.9	(3.0 - 4.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	634	7.6	(5.4 - 9.8)	35,552	6.8	(5.9 - 7.7)
Black Non-Hispanic	*	*	*	728	5.6	(2.2 - 9.0)
Hispanic	*	*	*	853	5.4	(1.1 - 9.6)
Other Non-Hispanic	*	*	*	705	4.8	(0.4 - 9.2)
<b>Education</b>						
≤ High School	312	9.9	(6.4 - 13.4)	15,485	8.5	(6.2 - 10.7)
Some College	161	4.4	(1.6 - 7.3)	9,860	5.5	(3.7 - 7.2)
≥ College Degree	181	7.2	(2.8 - 11.6)	12,411	4.7	(3.4 - 5.9)
<b>Income</b>						
≤ \$24,999	190	13.4	(8.0 - 18.9)	9,507	10.7	(7.2 - 14.1)
\$25,000-\$49,999	184	8.2	(3.6 - 12.9)	9,585	4.7	(3.6 - 5.7)
\$50,000-\$74,999	*	*	*	5,755	8.0	(4.8 - 11.1)
≥ \$75,000	*	*	*	8,541	3.3	(2.1 - 4.5)

<sup>1</sup> Defined as having ever been told by a health professional that respondent had a heart attack or angina

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 46a. Ever diagnosed with diabetes among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	659	4,300	8.7	(6.5 - 10.9)	38,174	1,382,700	9.3	(8.0 - 10.5)
<b>Gender</b>								
Male	241	2,000	7.8	(4.5 - 11.1)	14,626	530,400	7.5	(6.2 - 8.8)
Female	418	2,300	9.6	(6.7 - 12.4)	23,548	852,300	11.0	(8.9 - 13.0)
<b>Age (years)</b>								
18-34	*	*	*	*	3,941	150,200	3.4	(1.0 - 5.8)
35-44	*	*	*	*	4,987	165,100	5.7	(3.6 - 7.9)
45-54	135	900	9.2	(2.9 - 15.6)	7,830	279,000	9.8	(6.7 - 12.8)
55-64	153	1,400	19.9	(13.0 - 26.8)	8,771	332,900	15.5	(12.3 - 18.8)
≥ 65	230	1,500	16.8	(11.3 - 22.2)	12,645	455,600	17.9	(15.0 - 20.8)
<b>Race/Ethnicity</b>								
White Non-Hispanic	637	4,100	8.9	(6.6 - 11.2)	35,862	809,300	8.7	(7.8 - 9.5)
Black Non-Hispanic	*	*	*	*	739	261,000	12.1	(6.6 - 17.6)
Hispanic	*	*	*	*	862	227,300	10.1	(5.4 - 14.8)
Other Non-Hispanic	*	*	*	*	711	85,100	7.6	(2.3 - 12.8)
<b>Education</b>								
≤ High School	313	2,700	11.9	(7.9 - 15.9)	15,709	730,000	13.0	(10.5 - 15.6)
Some College	161	700	5.2	(2.3 - 8.1)	9,918	303,500	8.6	(5.9 - 11.2)
≥ College Degree	183	900	6.7	(3.2 - 10.2)	12,464	337,500	6.0	(4.8 - 7.2)
<b>Income</b>								
≤ \$24,999	191	1,400	12.0	(7.1 - 16.9)	9,652	543,400	15.5	(11.4 - 19.5)
\$25,000-\$49,999	184	1,200	9.0	(4.0 - 14.0)	9,667	291,000	10.0	(7.5 - 12.6)
\$50,000-\$74,999	95	700	7.8	(2.6 - 12.9)	5,777	151,000	7.6	(5.0 - 10.3)
≥ \$75,000	*	*	*	*	8,565	220,800	4.9	(3.8 - 6.1)

<sup>1</sup> Defined as having ever been told by a doctor that respondent had diabetes, excluding pre-diabetes and women with diabetes only when pregnant

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have diabetes (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 46b. Ever diagnosed with diabetes among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	659	8.1	(6.0 - 10.3)	38,174	9.0	(7.8 - 10.3)
<b>Gender</b>						
Male	241	7.7	(4.4 - 10.9)	14,626	7.5	(6.1 - 8.9)
Female	418	8.6	(5.8 - 11.4)	23,548	10.4	(8.4 - 12.4)
<b>Race/Ethnicity</b>						
White Non-Hispanic	637	8.4	(6.1 - 10.7)	35,862	8.4	(7.5 - 9.2)
Black Non-Hispanic	*	*	*	739	11.4	(6.0 - 16.7)
Hispanic	*	*	*	862	10.1	(5.4 - 14.9)
Other Non-Hispanic	*	*	*	711	7.7	(2.3 - 13.0)
<b>Education</b>						
≤ High School	313	11.3	(7.3 - 15.2)	15,709	12.6	(10.0 - 15.2)
Some College	161	4.7	(2.0 - 7.5)	9,918	8.3	(5.6 - 10.9)
≥ College Degree	183	6.4	(3.0 - 9.9)	12,464	5.9	(4.7 - 7.2)
<b>Income</b>						
≤ \$24,999	191	11.8	(6.9 - 16.7)	9,652	15.2	(11.1 - 19.2)
\$25,000-\$49,999	184	8.4	(3.6 - 13.3)	9,667	9.7	(7.1 - 12.2)
\$50,000-\$74,999	95	7.1	(2.2 - 12.0)	5,777	7.6	(4.9 - 10.2)
≥ \$75,000	*	*	*	8,565	4.7	(3.6 - 5.9)

<sup>1</sup> Defined as having ever been told by a doctor that respondent had diabetes, excluding pre-diabetes and women with diabetes only when pregnant

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 47a. Ever diagnosed with asthma (lifetime) among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	656	6,100	12.3	(9.2 - 15.4)	38,096	2,225,100	15.0	(13.2 - 16.8)
<b>Gender</b>								
Male	239	1,700	6.6	(3.4 - 9.8)	14,601	875,800	12.4	(10.1 - 14.7)
Female	417	4,400	18.3	(13.4 - 23.2)	23,495	1,349,300	17.4	(14.6 - 20.1)
<b>Age (years)</b>								
18-34	58	1,500	10.4	(3.5 - 17.3)	3,931	825,100	18.5	(13.8 - 23.3)
35-44	81	1,000	11.2	(3.9 - 18.5)	4,972	431,700	15.1	(11.2 - 19.1)
45-54	135	1,700	18.4	(11.3 - 25.5)	7,819	365,900	12.8	(9.7 - 16.0)
55-64	153	900	11.8	(5.9 - 17.7)	8,756	316,700	14.8	(11.9 - 17.7)
≥ 65	229	1,000	10.6	(6.2 - 15.0)	12,618	285,700	11.2	(8.6 - 13.8)
<b>Race/Ethnicity</b>								
White Non-Hispanic	634	5,700	12.5	(9.3 - 15.6)	35,783	1,365,700	14.6	(13.0 - 16.3)
Black Non-Hispanic	*	*	*	*	737	367,000	17.2	(10.2 - 24.2)
Hispanic	*	*	*	*	863	317,100	14.1	(9.0 - 19.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	311	3,000	13.4	(8.4 - 18.4)	15,675	831,600	14.9	(12.0 - 17.8)
Some College	161	1,600	12.2	(6.5 - 17.9)	9,893	587,500	16.6	(13.1 - 20.1)
≥ College Degree	182	1,500	10.7	(5.7 - 15.8)	12,444	739,000	13.2	(10.4 - 15.9)
<b>Income</b>								
≤ \$24,999	189	1,500	13.0	(7.1 - 18.8)	9,643	708,900	20.2	(14.9 - 25.5)
\$25,000-\$49,999	184	1,400	10.6	(5.8 - 15.4)	9,641	389,800	13.4	(10.0 - 16.9)
\$50,000-\$74,999	94	900	10.4	(3.1 - 17.7)	5,762	257,800	13.2	(10.0 - 16.5)
≥ \$75,000	112	1,400	12.5	(5.6 - 19.5)	8,562	577,900	12.8	(10.4 - 15.3)

<sup>1</sup> Defined as adults having ever been told by a health professional that they had asthma

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have lifetime asthma (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 47b. Ever diagnosed with asthma (lifetime) among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	656	12.6	(9.3 - 15.9)	38,096	15.2	(13.3 - 17.0)
<b>Gender</b>						
Male	239	6.8	(3.4 - 10.2)	14,601	12.6	(10.3 - 14.9)
Female	417	18.6	(13.2 - 24.0)	23,495	17.5	(14.8 - 20.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	634	12.7	(9.2 - 16.1)	35,783	14.9	(13.2 - 16.6)
Black Non-Hispanic	*	*	*	737	17.1	(10.1 - 24.1)
Hispanic	*	*	*	863	14.5	(9.3 - 19.7)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	311	14.2	(8.7 - 19.8)	15,675	15.1	(12.1 - 18.0)
Some College	161	11.8	(6.0 - 17.6)	9,893	16.9	(13.4 - 20.4)
≥ College Degree	182	10.8	(5.4 - 16.2)	12,444	13.3	(10.6 - 16.0)
<b>Income</b>						
≤ \$24,999	189	13.5	(7.1 - 20.0)	9,643	20.3	(15.0 - 25.6)
\$25,000-\$49,999	184	10.8	(5.6 - 16.1)	9,641	13.7	(10.2 - 17.2)
\$50,000-\$74,999	94	10.7	(2.8 - 18.6)	5,762	13.5	(10.0 - 17.0)
≥ \$75,000	112	12.4	(5.3 - 19.6)	8,562	13.0	(10.6 - 15.4)

<sup>1</sup> Defined as adults having ever been told by a health professional that they had asthma

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 48a. Current asthma among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	654	4,600	9.4	(6.7 - 12.1)	37,933	1,415,000	9.6	(8.1 - 11.0)
<b>Gender</b>								
Male	238	900	3.6	(1.4 - 5.7)	14,546	440,700	6.3	(4.8 - 7.7)
Female	416	3,700	15.5	(10.9 - 20.1)	23,387	974,300	12.6	(10.2 - 15.0)
<b>Age (years)</b>								
18-34	*	*	*	*	3,909	467,200	10.5	(6.8 - 14.3)
35-44	*	*	*	*	4,947	246,800	8.7	(5.8 - 11.6)
45-54	135	1,500	15.7	(9.1 - 22.4)	7,787	240,000	8.4	(6.1 - 10.8)
55-64	152	400	6.2	(2.2 - 10.2)	8,726	231,100	10.8	(8.1 - 13.5)
≥ 65	229	900	9.5	(5.3 - 13.7)	12,564	229,900	9.1	(6.5 - 11.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	632	4,300	9.3	(6.6 - 12.1)	35,629	888,400	9.6	(8.2 - 11.0)
Black Non-Hispanic	*	*	*	*	736	246,900	11.6	(5.7 - 17.5)
Hispanic	*	*	*	*	859	186,800	8.3	(4.4 - 12.2)
Other Non-Hispanic	*	*	*	*	709	93,000	8.3	(1.4 - 15.2)
<b>Education</b>								
≤ High School	311	2,900	13.0	(8.0 - 17.9)	15,594	595,100	10.7	(8.2 - 13.2)
Some College	160	1,100	8.5	(3.8 - 13.2)	9,861	322,000	9.1	(7.0 - 11.3)
≥ College Degree	181	600	4.5	(2.0 - 7.1)	12,394	430,900	7.7	(5.8 - 9.7)
<b>Income</b>								
≤ \$24,999	189	1,500	12.6	(6.8 - 18.4)	9,600	489,200	14.0	(9.7 - 18.3)
\$25,000-\$49,999	182	1,200	9.2	(4.6 - 13.7)	9,590	268,400	9.3	(6.2 - 12.4)
\$50,000-\$74,999	*	*	*	*	5,741	143,400	7.4	(5.1 - 9.6)
≥ \$75,000	112	700	6.6	(2.0 - 11.2)	8,541	339,200	7.6	(5.8 - 9.4)

<sup>1</sup> Defined as adults having ever been told by a health professional that they had asthma and still have asthma

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who currently have asthma (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 48b. Current asthma among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	654	9.6	(6.7 - 12.5)	37,933	9.7	(8.2 - 11.2)
<b>Gender</b>						
Male	238	3.5	(1.4 - 5.6)	14,546	6.4	(4.9 - 7.9)
Female	416	15.9	(10.8 - 21.1)	23,387	12.7	(10.2 - 15.1)
<b>Race/Ethnicity</b>						
White Non-Hispanic	632	9.5	(6.5 - 12.4)	35,629	9.7	(8.3 - 11.2)
Black Non-Hispanic	*	*	*	736	11.5	(5.6 - 17.3)
Hispanic	*	*	*	859	8.6	(4.5 - 12.6)
Other Non-Hispanic	*	*	*	709	8.3	(1.3 - 15.3)
<b>Education</b>						
≤ High School	311	13.8	(8.3 - 19.3)	15,594	10.8	(8.3 - 13.3)
Some College	160	8.4	(3.6 - 13.2)	9,861	9.4	(7.2 - 11.6)
≥ College Degree	181	4.3	(1.8 - 6.8)	12,394	7.8	(5.8 - 9.7)
<b>Income</b>						
≤ \$24,999	189	13.2	(6.8 - 19.6)	9,600	14.1	(9.7 - 18.4)
\$25,000-\$49,999	182	9.5	(4.5 - 14.5)	9,590	9.5	(6.3 - 12.7)
\$50,000-\$74,999	*	*	*	5,741	7.5	(5.0 - 10.1)
≥ \$75,000	112	6.5	(1.8 - 11.1)	8,541	7.7	(5.9 - 9.5)

<sup>1</sup> Defined as adults having ever been told by a health professional that they had asthma and still have asthma

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 49a. Chronic joint symptoms among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	658	21,400	43.1	(37.7 - 48.6)	38,099	6,377,100	42.9	(40.5 - 45.3)
<b>Gender</b>								
Male	241	10,300	40.3	(31.4 - 49.1)	14,595	3,033,600	42.8	(39.0 - 46.6)
Female	417	11,200	46.2	(40.1 - 52.2)	23,504	3,343,500	43.0	(40.0 - 46.0)
<b>Age (years)</b>								
18-34	*	*	*	*	3,933	1,261,300	28.3	(22.6 - 34.0)
35-44	*	*	*	*	4,984	1,110,900	38.6	(33.1 - 44.0)
45-54	135	5,400	56.9	(47.3 - 66.6)	7,817	1,456,500	51.2	(46.4 - 55.9)
55-64	153	4,000	54.7	(46.0 - 63.4)	8,751	1,167,400	54.6	(50.4 - 58.8)
≥ 65	229	4,900	53.2	(46.2 - 60.3)	12,614	1,381,000	54.3	(50.5 - 58.1)
<b>Race/Ethnicity</b>								
White Non-Hispanic	636	20,500	44.3	(38.8 - 49.8)	35,785	4,267,600	45.8	(43.8 - 47.7)
Black Non-Hispanic	*	*	*	*	740	890,500	41.3	(32.5 - 50.0)
Hispanic	*	*	*	*	860	827,500	36.7	(27.7 - 45.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	312	10,700	47.5	(38.8 - 56.2)	15,676	2,450,000	43.8	(39.4 - 48.1)
Some College	161	5,100	38.9	(28.9 - 48.9)	9,900	1,598,300	45.1	(40.1 - 50.1)
≥ College Degree	183	5,700	40.4	(30.8 - 49.9)	12,440	2,313,100	41.1	(37.8 - 44.5)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,641	1,483,000	42.2	(36.3 - 48.1)
\$25,000-\$49,999	*	*	*	*	9,647	1,332,100	45.9	(39.8 - 52.1)
\$50,000-\$74,999	*	*	*	*	5,768	894,200	45.3	(39.5 - 51.1)
≥ \$75,000	*	*	*	*	8,549	1,874,200	41.7	(38.4 - 45.1)

<sup>1</sup> Defined as adults having symptoms of pain, aching or stiffness in or around a joint during the past 30 days and began more than 3 months ago

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have chronic joint symptoms (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 49b. Chronic joint symptoms among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	658	42.6	(37.0 - 48.2)	38,099	42.1	(39.7 - 44.5)
<b>Gender</b>						
Male	241	40.7	(31.8 - 49.6)	14,595	42.6	(38.8 - 46.3)
Female	417	44.5	(38.0 - 51.0)	23,504	41.6	(38.6 - 44.7)
<b>Race/Ethnicity</b>						
White Non-Hispanic	636	43.8	(38.1 - 49.4)	35,785	44.7	(42.7 - 46.6)
Black Non-Hispanic	*	*	*	740	40.2	(31.5 - 48.9)
Hispanic	*	*	*	860	36.6	(27.8 - 45.4)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	312	48.2	(39.3 - 57.1)	15,676	43.1	(38.8 - 47.5)
Some College	161	37.3	(27.4 - 47.3)	9,900	43.9	(39.0 - 48.8)
≥ College Degree	183	39.0	(29.3 - 48.7)	12,440	40.3	(37.0 - 43.7)
<b>Income</b>						
≤ \$24,999	*	*	*	9,641	41.6	(35.7 - 47.4)
\$25,000-\$49,999	*	*	*	9,647	45.0	(39.0 - 51.0)
\$50,000-\$74,999	*	*	*	5,768	44.3	(38.4 - 50.2)
≥ \$75,000	*	*	*	8,549	40.8	(37.4 - 44.2)

<sup>1</sup> Defined as adults having symptoms of pain, aching or stiffness in or around a joint during the past 30 days and began more than 3 months ago

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 50a. Arthritis among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	655	17,300	35.0	(30.1 - 39.9)	38,013	4,102,800	27.7	(25.9 - 29.6)
<b>Gender</b>								
Male	240	8,000	31.5	(23.7 - 39.2)	14,555	1,642,700	23.3	(20.7 - 25.8)
Female	415	9,300	38.7	(33.1 - 44.2)	23,458	2,460,100	31.8	(29.2 - 34.4)
<b>Age (years)</b>								
18-34	*	*	*	*	3,927	312,100	7.0	(4.9 - 9.1)
35-44	*	*	*	*	4,972	412,500	14.5	(11.3 - 17.6)
45-54	135	3,500	36.4	(27.1 - 45.8)	7,789	869,100	30.7	(26.6 - 34.8)
55-64	151	4,100	57.8	(49.1 - 66.5)	8,729	999,300	47.0	(42.9 - 51.1)
≥ 65	228	6,000	65.9	(59.2 - 72.6)	12,596	1,509,800	59.6	(55.8 - 63.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	633	17,000	36.8	(31.8 - 41.9)	35,710	2,967,700	31.9	(30.3 - 33.6)
Black Non-Hispanic	*	*	*	*	737	588,100	27.3	(20.2 - 34.4)
Hispanic	*	*	*	*	856	380,700	17.1	(11.8 - 22.5)
Other Non-Hispanic	*	*	*	*	710	166,300	14.8	(7.8 - 21.8)
<b>Education</b>								
≤ High School	311	9,700	43.1	(34.8 - 51.4)	15,652	1,742,700	31.3	(27.8 - 34.8)
Some College	159	3,400	26.1	(18.3 - 33.9)	9,874	902,000	25.5	(22.3 - 28.6)
≥ College Degree	183	4,300	30.5	(22.3 - 38.7)	12,407	1,437,800	25.6	(22.9 - 28.4)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,618	1,112,500	32.0	(27.1 - 36.9)
\$25,000-\$49,999	184	5,200	38.7	(29.4 - 48.1)	9,624	821,900	28.4	(24.2 - 32.6)
\$50,000-\$74,999	*	*	*	*	5,753	542,500	27.5	(23.2 - 31.9)
≥ \$75,000	111	2,300	20.9	(12.3 - 29.6)	8,536	1,012,800	22.6	(19.8 - 25.3)

<sup>1</sup> Defined as adults having ever been told by a health professional that they have some form of arthritis, rheumatoid arthritis, gout, lupus or fibromyalgia (excluding back or neck)

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have ever been told by a health professional they have some form of arthritis (excluding back or neck) (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 50b. Arthritis among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	655	33.2	(28.3 - 38.1)	38,013	26.6	(24.8 - 28.5)
<b>Gender</b>						
Male	240	31.6	(23.8 - 39.4)	14,555	23.5	(20.9 - 26.1)
Female	415	34.9	(29.2 - 40.6)	23,458	29.5	(27.0 - 32.1)
<b>Race/Ethnicity</b>						
White Non-Hispanic	633	35.0	(29.9 - 40.1)	35,710	30.4	(28.7 - 32.0)
Black Non-Hispanic	*	*	*	737	26.5	(19.3 - 33.6)
Hispanic	*	*	*	856	17.2	(11.8 - 22.6)
Other Non-Hispanic	*	*	*	710	14.9	(7.7 - 22.0)
<b>Education</b>						
≤ High School	311	41.4	(33.0 - 49.7)	15,652	30.4	(26.9 - 33.8)
Some College	159	24.9	(17.1 - 32.7)	9,874	24.1	(21.0 - 27.1)
≥ College Degree	183	28.5	(20.3 - 36.6)	12,407	24.7	(21.9 - 27.5)
<b>Income</b>						
≤ \$24,999	*	*	*	9,618	31.2	(26.3 - 36.1)
\$25,000-\$49,999	184	37.5	(27.9 - 47.1)	9,624	27.3	(23.2 - 31.3)
\$50,000-\$74,999	*	*	*	5,753	26.3	(21.9 - 30.7)
≥ \$75,000	111	20.3	(11.6 - 29.1)	8,536	21.6	(18.8 - 24.3)

<sup>1</sup> Defined as adults having ever been told by a health professional that they have some form of arthritis, rheumatoid arthritis, gout, lupus or fibromyalgia (excluding back or neck)

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 51a. Activity limitations because of physical, mental or emotional problems among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	656	9,300	18.8	(15.4 - 22.3)	37,998	3,053,000	20.6	(18.7 - 22.5)
<b>Gender</b>								
Male	238	3,500	13.7	(9.2 - 18.3)	14,566	1,318,400	18.6	(15.7 - 21.5)
Female	418	5,800	24.2	(19.3 - 29.0)	23,432	1,734,600	22.4	(19.8 - 25.0)
<b>Age (years)</b>								
18-34	*	*	*	*	3,929	580,000	13.0	(8.2 - 17.9)
35-44	83	1,900	21.3	(11.6 - 31.1)	4,973	418,600	14.5	(11.0 - 18.1)
45-54	135	2,100	22.4	(14.6 - 30.2)	7,812	620,200	21.7	(18.3 - 25.2)
55-64	152	1,900	26.8	(18.9 - 34.7)	8,739	651,600	30.5	(26.5 - 34.6)
≥ 65	228	2,700	29.4	(23.1 - 35.8)	12,545	782,700	31.1	(27.7 - 34.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	634	9,100	19.7	(16.1 - 23.3)	35,688	2,037,200	21.9	(20.3 - 23.5)
Black Non-Hispanic	*	*	*	*	735	429,700	20.1	(13.5 - 26.7)
Hispanic	*	*	*	*	863	448,200	19.8	(11.9 - 27.8)
Other Non-Hispanic	*	*	*	*	712	137,900	12.3	(5.2 - 19.3)
<b>Education</b>								
≤ High School	312	5,800	25.9	(19.6 - 32.2)	15,626	1,359,600	24.4	(20.6 - 28.1)
Some College	161	1,900	14.9	(9.4 - 20.4)	9,875	700,800	19.8	(16.5 - 23.2)
≥ College Degree	181	1,600	11.2	(6.7 - 15.6)	12,415	924,800	16.5	(14.2 - 18.7)
<b>Income</b>								
≤ \$24,999	190	3,800	32.3	(22.7 - 41.9)	9,603	1,055,800	30.1	(24.7 - 35.5)
\$25,000-\$49,999	183	2,100	15.3	(9.5 - 21.1)	9,628	630,900	21.8	(16.6 - 27.0)
\$50,000-\$74,999	95	1,100	12.7	(6.3 - 19.2)	5,760	408,100	20.7	(16.1 - 25.2)
≥ \$75,000	112	900	8.7	(3.6 - 13.7)	8,544	583,300	13.0	(11.1 - 14.9)

<sup>1</sup> Defined as activity limitations because of physical, mental or emotional problems

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who are limited in any activities because of physical, mental or emotional problems (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 51b. Activity limitations because of physical, mental or emotional problems among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	656	18.6	(15.0 - 22.2)	37,998	20.1	(18.2 - 22.0)
<b>Gender</b>						
Male	238	13.8	(9.1 - 18.4)	14,566	18.6	(15.8 - 21.4)
Female	418	23.6	(18.4 - 28.8)	23,432	21.5	(18.9 - 24.1)
<b>Race/Ethnicity</b>						
White Non-Hispanic	634	19.5	(15.7 - 23.2)	35,688	21.3	(19.7 - 22.9)
Black Non-Hispanic	*	*	*	735	19.9	(13.2 - 26.5)
Hispanic	*	*	*	863	19.1	(11.5 - 26.8)
Other Non-Hispanic	*	*	*	712	12.4	(5.3 - 19.6)
<b>Education</b>						
≤ High School	312	26.0	(19.3 - 32.7)	15,626	23.8	(20.1 - 27.4)
Some College	161	15.1	(9.3 - 20.9)	9,875	19.3	(16.0 - 22.6)
≥ College Degree	181	10.4	(6.1 - 14.8)	12,415	16.1	(13.9 - 18.3)
<b>Income</b>						
≤ \$24,999	190	32.7	(22.7 - 42.7)	9,603	29.7	(24.3 - 35.2)
\$25,000-\$49,999	183	15.2	(9.1 - 21.3)	9,628	21.2	(16.1 - 26.3)
\$50,000-\$74,999	95	12.3	(6.0 - 18.6)	5,760	20.1	(15.6 - 24.6)
≥ \$75,000	112	7.9	(3.2 - 12.6)	8,544	12.7	(10.8 - 14.6)

<sup>1</sup> Defined as activity limitations because of physical, mental or emotional problems

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 52a. Have health problem that requires the use of special equipment among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	658	3,700	7.4	(5.4 - 9.4)	38,171	1,169,400	7.9	(6.9 - 8.9)
<b>Gender</b>								
Male	240	1,400	5.4	(2.6 - 8.1)	14,617	460,700	6.5	(5.3 - 7.7)
Female	418	2,300	9.5	(6.6 - 12.3)	23,554	708,700	9.1	(7.6 - 10.7)
<b>Age (years)</b>								
18-34	*	*	*	*	3,940	68,800	1.5	(0.7 - 2.4)
35-44	*	*	*	*	4,989	122,500	4.3	(2.4 - 6.1)
45-54	135	500	5.3	(1.9 - 8.7)	7,836	182,500	6.4	(4.7 - 8.1)
55-64	153	800	10.5	(5.5 - 15.5)	8,773	219,600	10.3	(7.8 - 12.8)
≥ 65	230	1,600	17.8	(12.6 - 23.1)	12,633	576,100	22.7	(19.0 - 26.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	636	3,100	6.7	(4.9 - 8.4)	35,853	802,000	8.6	(7.6 - 9.6)
Black Non-Hispanic	*	*	*	*	739	235,800	10.9	(6.4 - 15.5)
Hispanic	*	*	*	*	864	104,100	4.6	(2.1 - 7.1)
Other Non-Hispanic	*	*	*	*	715	27,600	2.5	(0.8 - 4.1)
<b>Education</b>								
≤ High School	312	2,000	9.0	(5.8 - 12.1)	15,711	604,200	10.8	(8.7 - 12.8)
Some College	161	800	6.0	(1.6 - 10.3)	9,914	260,000	7.3	(5.5 - 9.1)
≥ College Degree	183	900	6.2	(3.3 - 9.1)	12,463	298,900	5.3	(4.1 - 6.5)
<b>Income</b>								
≤ \$24,999	190	1,900	15.9	(9.4 - 22.3)	9,656	430,500	12.2	(9.6 - 14.8)
\$25,000-\$49,999	184	700	5.5	(2.3 - 8.7)	9,663	238,900	8.2	(5.9 - 10.5)
\$50,000-\$74,999	*	*	*	*	5,775	103,000	5.2	(3.6 - 6.8)
≥ \$75,000	*	*	*	*	8,566	136,000	3.0	(2.1 - 4.0)

<sup>1</sup> Defined as having any health problem that requires the use of special equipment such as a cane, a wheelchair, a special bed, or a special telephone

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have any health problem that requires the use of special equipment (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 52b. Have health problem that requires the use of special equipment among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	658	6.9	(4.9 - 8.8)	38,171	7.6	(6.6 - 8.6)
<b>Gender</b>						
Male	240	5.3	(2.7 - 7.9)	14,617	6.8	(5.5 - 8.0)
Female	418	8.6	(5.7 - 11.4)	23,554	8.4	(6.9 - 9.9)
<b>Race/Ethnicity</b>						
White Non-Hispanic	636	6.2	(4.5 - 7.9)	35,853	8.3	(7.3 - 9.3)
Black Non-Hispanic	*	*	*	739	10.5	(6.1 - 15.0)
Hispanic	*	*	*	864	4.5	(2.1 - 6.9)
Other Non-Hispanic	*	*	*	715	2.2	(0.7 - 3.8)
<b>Education</b>						
≤ High School	312	8.5	(5.2 - 11.7)	15,711	10.3	(8.3 - 12.3)
Some College	161	5.5	(1.5 - 9.5)	9,914	7.1	(5.2 - 8.9)
≥ College Degree	183	5.7	(2.9 - 8.5)	12,463	5.3	(4.1 - 6.6)
<b>Income</b>						
≤ \$24,999	190	14.8	(8.6 - 21.0)	9,656	11.9	(9.4 - 14.5)
\$25,000-\$49,999	184	5.0	(1.9 - 8.1)	9,663	7.7	(5.5 - 10.0)
\$50,000-\$74,999	*	*	*	5,775	5.1	(3.4 - 6.8)
≥ \$75,000	*	*	*	8,566	2.9	(2.0 - 3.9)

<sup>1</sup> Defined as having any health problem that requires the use of special equipment such as a cane, a wheelchair, a special bed, or a special telephone

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 53a. Disability among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	657	10,700	21.5	(17.7 - 25.2)	38,016	3,394,200	22.9	(20.9 - 24.9)
<b>Gender</b>								
Male	239	4,200	16.4	(11.3 - 21.5)	14,568	1,469,700	20.7	(17.8 - 23.7)
Female	418	6,500	26.8	(21.8 - 31.8)	23,448	1,924,600	24.8	(22.1 - 27.5)
<b>Age (years)</b>								
18-34	*	*	*	*	3,931	590,300	13.3	(8.4 - 18.1)
35-44	83	1,900	21.3	(11.6 - 31.1)	4,974	447,900	15.6	(11.9 - 19.3)
45-54	135	2,200	23.2	(15.4 - 31.1)	7,811	643,900	22.6	(19.1 - 26.0)
55-64	152	2,200	30.6	(22.5 - 38.8)	8,740	702,100	33.0	(28.9 - 37.0)
≥ 65	229	3,300	35.9	(29.2 - 42.6)	12,560	1,010,000	39.9	(36.1 - 43.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	635	9,900	21.4	(17.7 - 25.1)	35,703	2,253,400	24.2	(22.6 - 25.9)
Black Non-Hispanic	*	*	*	*	736	511,700	23.8	(16.8 - 30.8)
Hispanic	*	*	*	*	864	484,300	21.4	(13.4 - 29.5)
Other Non-Hispanic	*	*	*	*	713	144,800	12.9	(5.8 - 19.9)
<b>Education</b>								
≤ High School	312	6,500	28.9	(22.3 - 35.6)	15,638	1,544,200	27.6	(23.7 - 31.4)
Some College	161	2,300	17.5	(11.0 - 24.1)	9,876	755,200	21.4	(17.9 - 24.8)
≥ College Degree	182	1,900	13.3	(8.4 - 18.1)	12,419	1,021,300	18.2	(15.9 - 20.5)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,613	1,179,100	33.6	(28.0 - 39.2)
\$25,000-\$49,999	184	2,300	17.0	(10.9 - 23.0)	9,629	699,000	24.2	(18.9 - 29.4)
\$50,000-\$74,999	95	1,200	13.6	(7.0 - 20.2)	5,757	430,100	21.8	(17.2 - 26.4)
≥ \$75,000	112	1,000	9.4	(4.2 - 14.5)	8,546	632,700	14.1	(12.1 - 16.1)

<sup>1</sup> Defined as having activity limitations due to physical, mental or emotional problems or have health problems that require the use of special equipment

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults with a disability (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 53b. Disability among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	657	21.0	(17.2 - 24.8)	38,016	22.3	(20.4 - 24.3)
<b>Gender</b>						
Male	239	16.3	(11.2 - 21.4)	14,568	20.9	(18.0 - 23.8)
Female	418	25.9	(20.6 - 31.2)	23,448	23.6	(21.0 - 26.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	635	21.0	(17.1 - 24.8)	35,703	23.6	(21.9 - 25.3)
Black Non-Hispanic	*	*	*	736	23.5	(16.5 - 30.5)
Hispanic	*	*	*	864	20.6	(12.9 - 28.3)
Other Non-Hispanic	*	*	*	713	12.9	(5.8 - 20.1)
<b>Education</b>						
≤ High School	312	28.9	(21.9 - 35.8)	15,638	26.8	(23.0 - 30.6)
Some College	161	17.4	(10.8 - 24.0)	9,876	20.9	(17.4 - 24.3)
≥ College Degree	182	12.3	(7.6 - 17.0)	12,419	17.9	(15.5 - 20.3)
<b>Income</b>						
≤ \$24,999	*	*	*	9,613	33.2	(27.6 - 38.8)
\$25,000-\$49,999	184	16.9	(10.5 - 23.3)	9,629	23.5	(18.3 - 28.6)
\$50,000-\$74,999	95	13.0	(6.6 - 19.5)	5,757	21.2	(16.6 - 25.8)
≥ \$75,000	112	8.5	(3.7 - 13.3)	8,546	13.8	(11.7 - 15.8)

<sup>1</sup> Defined as having activity limitations due to physical, mental or emotional problems or have health problems that require the use of special equipment

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 54a. Current smoking among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	659	11,500	23.2	(17.9 - 28.4)	38,029	2,483,500	16.8	(15.1 - 18.6)
<b>Gender</b>								
Male	241	6,200	24.2	(15.5 - 33.0)	14,559	1,356,400	19.3	(16.3 - 22.2)
Female	418	5,300	22.0	(16.5 - 27.5)	23,470	1,127,100	14.6	(12.7 - 16.6)
<b>Age (years)</b>								
18-34	*	*	*	*	3,929	993,500	22.3	(17.5 - 27.1)
35-44	83	1,900	20.9	(11.0 - 30.8)	4,977	421,100	14.8	(11.6 - 17.9)
45-54	135	2,500	26.9	(18.1 - 35.6)	7,807	508,800	18.3	(15.4 - 21.3)
55-64	153	1,500	21.0	(13.7 - 28.2)	8,744	314,500	14.7	(12.1 - 17.3)
≥ 65	230	600	6.9	(3.5 - 10.2)	12,572	245,600	9.7	(7.2 - 12.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	637	10,000	21.5	(16.6 - 26.4)	35,729	1,615,500	17.3	(15.9 - 18.8)
Black Non-Hispanic	*	*	*	*	735	284,100	13.6	(8.9 - 18.2)
Hispanic	*	*	*	*	858	350,900	15.8	(9.6 - 21.9)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	313	7,100	31.6	(22.7 - 40.6)	15,654	1,237,300	22.4	(19.1 - 25.6)
Some College	161	3,200	24.9	(15.4 - 34.4)	9,875	670,600	19.1	(15.5 - 22.8)
≥ College Degree	183	1,100	8.2	(3.0 - 13.3)	12,421	563,300	10.0	(7.5 - 12.5)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,626	835,100	24.1	(19.1 - 29.1)
\$25,000-\$49,999	184	3,300	24.6	(15.9 - 33.4)	9,629	558,100	19.4	(15.6 - 23.2)
\$50,000-\$74,999	*	*	*	*	5,756	338,100	17.3	(13.0 - 21.7)
≥ \$75,000	112	900	8.3	(2.6 - 14.0)	8,549	468,700	10.4	(8.6 - 12.3)

<sup>1</sup> Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day or some days

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adult current smokers (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 54b. Current smoking among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	659	23.5	(18.2 - 28.8)	38,029	17.0	(15.3 - 18.8)
<b>Gender</b>						
Male	241	23.4	(15.0 - 31.9)	14,559	19.2	(16.3 - 22.1)
Female	418	23.5	(17.4 - 29.7)	23,470	15.0	(13.0 - 17.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	637	22.1	(17.0 - 27.1)	35,729	17.7	(16.2 - 19.2)
Black Non-Hispanic	*	*	*	735	13.5	(8.8 - 18.2)
Hispanic	*	*	*	858	15.9	(9.8 - 22.0)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	313	32.2	(23.2 - 41.2)	15,654	22.8	(19.5 - 26.0)
Some College	161	25.4	(15.6 - 35.2)	9,875	19.4	(15.8 - 22.9)
≥ College Degree	183	8.7	(3.0 - 14.4)	12,421	10.0	(7.6 - 12.5)
<b>Income</b>						
≤ \$24,999	*	*	*	9,626	24.1	(19.2 - 29.1)
\$25,000-\$49,999	184	24.7	(15.8 - 33.7)	9,629	19.9	(16.0 - 23.7)
\$50,000-\$74,999	*	*	*	5,756	17.8	(13.3 - 22.3)
≥ \$75,000	112	8.2	(2.3 - 14.1)	8,549	10.6	(8.6 - 12.5)

<sup>1</sup> Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day or some days

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 55a. Everyday smoking among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	659	9,100	18.3	(13.6 - 23.0)	38,029	1,626,800	11.0	(9.8 - 12.2)
<b>Gender</b>								
Male	241	4,900	19.1	(11.2 - 27.0)	14,559	848,400	12.0	(10.0 - 14.1)
Female	418	4,200	17.4	(12.7 - 22.2)	23,470	778,400	10.1	(8.7 - 11.5)
<b>Age (years)</b>								
18-34	*	*	*	*	3,929	559,800	12.6	(9.6 - 15.6)
35-44	83	1,600	17.5	(8.4 - 26.6)	4,977	315,200	11.0	(8.4 - 13.7)
45-54	135	2,200	22.9	(14.4 - 31.4)	7,807	377,000	13.6	(11.1 - 16.0)
55-64	153	1,400	19.2	(12.1 - 26.3)	8,744	203,000	9.5	(7.5 - 11.5)
≥ 65	230	500	5.7	(2.7 - 8.8)	12,572	171,700	6.8	(4.8 - 8.8)
<b>Race/Ethnicity</b>								
White Non-Hispanic	637	7,900	17.0	(12.9 - 21.2)	35,729	1,158,100	12.4	(11.2 - 13.7)
Black Non-Hispanic	*	*	*	*	735	197,700	9.4	(5.6 - 13.3)
Hispanic	*	*	*	*	858	176,200	7.9	(4.2 - 11.6)
Other Non-Hispanic	*	*	*	*	707	94,900	8.5	(3.3 - 13.7)
<b>Education</b>								
≤ High School	313	5,800	25.9	(17.6 - 34.2)	15,654	881,900	15.9	(13.4 - 18.5)
Some College	161	2,200	16.8	(9.0 - 24.6)	9,875	410,200	11.7	(9.6 - 13.8)
≥ College Degree	183	1,100	7.6	(2.5 - 12.8)	12,421	323,500	5.8	(4.3 - 7.2)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,626	452,800	13.1	(10.3 - 15.8)
\$25,000-\$49,999	184	3,000	22.3	(13.8 - 30.8)	9,629	425,900	14.8	(11.9 - 17.7)
\$50,000-\$74,999	*	*	*	*	5,756	222,700	11.4	(8.5 - 14.4)
≥ \$75,000	*	*	*	*	8,549	320,200	7.1	(5.5 - 8.8)

<sup>1</sup> Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have smoked at least 100 cigarettes in lifetime and currently smoking everyday (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 55b. Everyday smoking among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	659	18.4	(13.7 - 23.1)	38,029	11.3	(10.0 - 12.5)
<b>Gender</b>						
Male	241	18.5	(10.8 - 26.1)	14,559	12.1	(10.1 - 14.1)
Female	418	18.4	(13.1 - 23.7)	23,470	10.5	(9.0 - 12.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	637	17.3	(13.0 - 21.7)	35,729	12.8	(11.5 - 14.1)
Black Non-Hispanic	*	*	*	735	9.4	(5.5 - 13.4)
Hispanic	*	*	*	858	8.0	(4.3 - 11.7)
Other Non-Hispanic	*	*	*	707	8.5	(3.2 - 13.8)
<b>Education</b>						
≤ High School	313	26.1	(17.8 - 34.4)	15,654	16.4	(13.8 - 19.0)
Some College	161	16.9	(9.1 - 24.8)	9,875	12.1	(9.9 - 14.3)
≥ College Degree	183	8.2	(2.5 - 13.9)	12,421	5.8	(4.3 - 7.3)
<b>Income</b>						
≤ \$24,999	*	*	*	9,626	13.4	(10.6 - 16.2)
\$25,000-\$49,999	184	22.4	(13.7 - 31.0)	9,629	15.3	(12.3 - 18.3)
\$50,000-\$74,999	*	*	*	5,756	11.7	(8.7 - 14.8)
≥ \$75,000	*	*	*	8,549	7.2	(5.5 - 8.9)

<sup>1</sup> Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 56a. Adults living in homes in which smoking is prohibited<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	656	35,800	72.1	(66.5 - 77.7)	38,045	11,997,100	80.8	(79.0 - 82.6)
<b>Gender</b>								
Male	240	17,900	70.1	(60.6 - 79.5)	14,577	5,634,400	79.4	(76.6 - 82.2)
Female	416	17,900	74.3	(68.7 - 79.9)	23,468	6,362,800	82.1	(79.9 - 84.3)
<b>Age (years)</b>								
18-34	*	*	*	*	3,934	3,655,200	82.1	(77.8 - 86.4)
35-44	*	*	*	*	4,985	2,479,700	86.1	(82.5 - 89.7)
45-54	135	6,700	71.0	(62.2 - 79.7)	7,808	2,194,400	77.2	(73.5 - 80.8)
55-64	152	5,700	79.4	(72.3 - 86.5)	8,753	1,732,500	81.0	(78.3 - 83.7)
≥ 65	228	7,000	76.6	(70.7 - 82.6)	12,565	1,935,400	76.5	(73.4 - 79.5)
<b>Race/Ethnicity</b>								
White Non-Hispanic	634	34,700	75.0	(70.0 - 80.0)	35,737	7,426,900	79.8	(78.3 - 81.2)
Black Non-Hispanic	*	*	*	*	738	1,692,100	78.6	(71.9 - 85.3)
Hispanic	*	*	*	*	863	2,003,900	88.7	(84.1 - 93.3)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	312	15,100	66.8	(58.0 - 75.6)	15,647	4,307,200	77.1	(74.0 - 80.2)
Some College	160	9,100	70.3	(60.3 - 80.2)	9,883	2,795,400	78.9	(75.0 - 82.8)
≥ College Degree	*	*	*	*	12,437	4,817,300	85.7	(83.3 - 88.2)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,608	2,627,700	74.9	(70.0 - 79.7)
\$25,000-\$49,999	184	10,100	75.3	(66.4 - 84.2)	9,640	2,314,500	79.8	(76.0 - 83.5)
\$50,000-\$74,999	*	*	*	*	5,766	1,551,300	78.7	(74.0 - 83.4)
≥ \$75,000	*	*	*	*	8,556	3,883,200	86.5	(84.3 - 88.7)

<sup>1</sup> Defined as having a rule that prohibits smoking anywhere inside the home

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who live in a home with a rule that prohibits smoking anywhere inside the home (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 56b. Adults living in homes in which smoking is prohibited<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	656	71.7	(65.9 - 77.4)	38,045	80.9	(79.2 - 82.7)
<b>Gender</b>						
Male	240	69.7	(60.1 - 79.3)	14,577	79.4	(76.7 - 82.2)
Female	416	73.7	(67.5 - 79.9)	23,468	82.3	(80.0 - 84.5)
<b>Race/Ethnicity</b>						
White Non-Hispanic	634	74.5	(69.3 - 79.8)	35,737	79.9	(78.4 - 81.4)
Black Non-Hispanic	*	*	*	738	78.6	(71.8 - 85.4)
Hispanic	*	*	*	863	88.9	(84.4 - 93.4)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	312	66.7	(57.7 - 75.6)	15,647	76.9	(73.8 - 80.0)
Some College	*	*	*	9,883	79.1	(75.2 - 83.0)
≥ College Degree	*	*	*	12,437	86.0	(83.5 - 88.5)
<b>Income</b>						
≤ \$24,999	*	*	*	9,608	74.9	(70.1 - 79.7)
\$25,000-\$49,999	184	75.2	(65.8 - 84.5)	9,640	79.8	(76.1 - 83.6)
\$50,000-\$74,999	*	*	*	5,766	78.6	(73.8 - 83.3)
≥ \$75,000	*	*	*	8,556	86.6	(84.3 - 88.9)

<sup>1</sup> Defined as having a rule that prohibits smoking anywhere inside the home

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 57a. Binge drinking within the past month among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	654	9,900	20.1	(15.1 - 25.0)	37,745	2,602,000	17.7	(15.7 - 19.8)
<b>Gender</b>								
Male	240	5,800	22.7	(14.7 - 30.8)	14,420	1,712,400	24.5	(20.9 - 28.1)
Female	414	4,100	17.2	(11.6 - 22.8)	23,325	889,600	11.6	(9.5 - 13.7)
<b>Age (years)</b>								
18-34	*	*	*	*	3,895	1,254,300	28.5	(22.8 - 34.2)
35-44	*	*	*	*	4,934	584,800	20.6	(16.5 - 24.7)
45-54	133	1,500	15.9	(8.7 - 23.0)	7,738	467,600	16.6	(13.3 - 19.9)
55-64	153	1,100	14.7	(8.3 - 21.1)	8,663	207,200	9.8	(7.9 - 11.8)
≥ 65	230	500	5.1	(2.2 - 8.0)	12,515	88,100	3.5	(2.5 - 4.5)
<b>Race/Ethnicity</b>								
White Non-Hispanic	632	9,300	20.2	(15.1 - 25.3)	35,453	1,761,900	19.2	(17.5 - 20.8)
Black Non-Hispanic	*	*	*	*	728	247,000	11.6	(6.3 - 17.0)
Hispanic	*	*	*	*	854	460,600	20.5	(11.3 - 29.8)
Other Non-Hispanic	*	*	*	*	710	132,500	12.1	(3.4 - 20.7)
<b>Education</b>								
≤ High School	311	4,800	21.4	(13.6 - 29.1)	15,491	945,800	17.4	(13.2 - 21.6)
Some College	*	*	*	*	9,819	636,600	18.1	(14.6 - 21.5)
≥ College Degree	182	2,100	14.8	(7.1 - 22.5)	12,358	1,018,700	18.2	(15.3 - 21.1)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,557	469,100	13.5	(8.5 - 18.5)
\$25,000-\$49,999	183	1,800	13.4	(7.7 - 19.0)	9,575	534,300	18.5	(13.0 - 24.0)
\$50,000-\$74,999	*	*	*	*	5,726	378,200	19.2	(14.1 - 24.3)
≥ \$75,000	*	*	*	*	8,501	1,011,400	22.6	(19.7 - 25.6)

<sup>1</sup> Defined as men having 5 or more drinks or women having 4 or more drinks on 1 or more occasion within the past month

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adult binge drinkers within the past month (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 57b. Binge drinking within the past month among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	654	21.1	(15.9 - 26.3)	37,745	18.1	(16.1 - 20.2)
<b>Gender</b>						
Male	240	23.1	(14.9 - 31.2)	14,420	24.5	(21.0 - 27.9)
Female	414	19.0	(12.6 - 25.4)	23,325	12.4	(10.2 - 14.5)
<b>Race/Ethnicity</b>						
White Non-Hispanic	632	21.5	(16.1 - 26.9)	35,453	19.8	(18.1 - 21.6)
Black Non-Hispanic	*	*	*	728	11.8	(6.4 - 17.1)
Hispanic	*	*	*	854	20.2	(11.3 - 29.1)
Other Non-Hispanic	*	*	*	710	12.1	(3.4 - 20.8)
<b>Education</b>						
≤ High School	311	22.0	(14.1 - 30.0)	15,491	17.5	(13.5 - 21.5)
Some College	*	*	*	9,819	18.6	(15.1 - 22.0)
≥ College Degree	182	16.1	(7.5 - 24.7)	12,358	18.8	(15.9 - 21.7)
<b>Income</b>						
≤ \$24,999	*	*	*	9,557	13.4	(8.6 - 18.2)
\$25,000-\$49,999	183	13.5	(7.5 - 19.4)	9,575	18.9	(13.5 - 24.2)
\$50,000-\$74,999	*	*	*	5,726	19.9	(14.8 - 25.1)
≥ \$75,000	*	*	*	8,501	23.2	(20.2 - 26.1)

<sup>1</sup> Defined as men having 5 or more drinks or women having 4 or more drinks on 1 or more occasion within the past month

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 58a. Heavy drinking within the past month among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	649	3,500	7.1	(3.7 - 10.5)	37,553	718,700	4.9	(4.2 - 5.6)
<b>Gender</b>								
Male	237	1,700	6.8	(1.4 - 12.3)	14,330	333,300	4.8	(3.9 - 5.7)
Female	412	1,800	7.4	(3.6 - 11.2)	23,223	385,400	5.0	(4.0 - 6.1)
<b>Age (years)</b>								
18-34	*	*	*	*	3,866	224,900	5.1	(3.5 - 6.8)
35-44	*	*	*	*	4,924	137,200	4.8	(3.0 - 6.7)
45-54	133	600	6.2	(2.4 - 10.0)	7,715	144,600	5.2	(4.1 - 6.2)
55-64	153	500	6.7	(2.3 - 11.1)	8,624	117,400	5.6	(3.9 - 7.3)
≥ 65	224	400	4.8	(1.9 - 7.8)	12,424	94,600	3.8	(2.7 - 4.9)
<b>Race/Ethnicity</b>								
White Non-Hispanic	627	3,400	7.5	(3.9 - 11.1)	35,279	572,400	6.3	(5.4 - 7.1)
Black Non-Hispanic	*	*	*	*	723	58,100	2.8	(0.6 - 4.9)
Hispanic	*	*	*	*	848	60,900	2.7	(0.5 - 4.9)
Other Non-Hispanic	*	*	*	*	703	27,200	2.5	(0.8 - 4.2)
<b>Education</b>								
≤ High School	309	2,300	10.4	(3.9 - 16.8)	15,411	228,100	4.2	(3.2 - 5.2)
Some College	158	800	6.3	(0.9 - 11.8)	9,759	163,700	4.7	(3.6 - 5.8)
≥ College Degree	*	*	*	*	12,306	326,800	5.9	(4.5 - 7.3)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,512	104,200	3.0	(1.8 - 4.1)
\$25,000-\$49,999	182	700	5.2	(2.2 - 8.2)	9,537	158,700	5.5	(3.9 - 7.2)
\$50,000-\$74,999	*	*	*	*	5,708	95,400	4.9	(3.1 - 6.7)
≥ \$75,000	110	600	5.6	(1.4 - 9.8)	8,481	313,900	7.0	(5.4 - 8.6)

<sup>1</sup> Defined as adult men averaging more than 2 alcoholic drinks per day and adult women averaging more than 1 alcoholic drink per day within the past month

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adult heavy drinkers within the past month (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 58b. Heavy drinking within the past month among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	649	7.2	(3.7 - 10.6)	37,553	5.0	(4.3 - 5.8)
<b>Gender</b>						
Male	237	6.8	(1.5 - 12.1)	14,330	4.9	(3.9 - 5.9)
Female	412	7.5	(3.2 - 11.9)	23,223	5.1	(4.0 - 6.2)
<b>Race/Ethnicity</b>						
White Non-Hispanic	627	7.6	(3.9 - 11.2)	35,279	6.4	(5.5 - 7.2)
Black Non-Hispanic	*	*	*	723	2.9	(0.6 - 5.1)
Hispanic	*	*	*	848	2.8	(0.5 - 5.1)
Other Non-Hispanic	*	*	*	703	2.5	(0.8 - 4.2)
<b>Education</b>						
≤ High School	309	10.6	(4.1 - 17.2)	15,411	4.3	(3.2 - 5.3)
Some College	158	6.6	(0.4 - 12.8)	9,759	4.8	(3.6 - 5.9)
≥ College Degree	*	*	*	12,306	6.0	(4.5 - 7.4)
<b>Income</b>						
≤ \$24,999	*	*	*	9,512	3.0	(1.9 - 4.1)
\$25,000-\$49,999	182	4.8	(2.0 - 7.7)	9,537	5.8	(4.0 - 7.5)
\$50,000-\$74,999	*	*	*	5,708	5.0	(3.1 - 7.0)
≥ \$75,000	110	5.3	(0.9 - 9.7)	8,481	7.0	(5.4 - 8.7)

<sup>1</sup> Defined as adult men averaging more than 2 alcoholic drinks per day and adult women averaging more than 1 alcoholic drink per day within the past month

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 59a. Never or rarely been asked about sexual history during routine check-up among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	592	27,200	62.9	(57.0 - 68.8)	35,193	8,480,600	61.9	(59.2 - 64.5)
<b>Gender</b>								
Male	212	14,500	69.3	(59.5 - 79.2)	13,299	4,357,400	67.7	(63.6 - 71.7)
Female	380	12,600	56.8	(50.3 - 63.4)	21,894	4,123,300	56.8	(53.3 - 60.2)
<b>Age (years)</b>								
18-34	*	*	*	*	3,507	1,804,600	44.3	(37.8 - 50.8)
35-44	*	*	*	*	4,531	1,635,300	62.2	(56.6 - 67.8)
45-54	*	*	*	*	7,215	1,715,600	65.4	(60.5 - 70.3)
55-64	134	4,200	66.7	(57.5 - 75.8)	8,168	1,416,200	70.7	(66.6 - 74.8)
≥ 65	212	7,200	83.6	(78.2 - 88.9)	11,772	1,908,900	80.3	(76.9 - 83.8)
<b>Race/Ethnicity</b>								
White Non-Hispanic	574	25,600	62.9	(56.9 - 68.9)	33,090	5,940,300	68.5	(66.3 - 70.7)
Black Non-Hispanic	*	*	*	*	683	831,800	40.8	(31.8 - 49.9)
Hispanic	*	*	*	*	784	1,071,700	53.4	(43.6 - 63.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	277	13,500	70.3	(62.2 - 78.4)	14,264	3,001,200	60.1	(55.4 - 64.9)
Some College	*	*	*	*	9,156	1,972,800	60.3	(54.9 - 65.7)
≥ College Degree	*	*	*	*	11,706	3,476,000	65.0	(61.2 - 68.7)
<b>Income</b>								
≤ \$24,999	*	*	*	*	8,772	1,529,400	49.3	(42.6 - 56.0)
\$25,000-\$49,999	*	*	*	*	8,932	1,705,000	64.4	(58.0 - 70.7)
\$50,000-\$74,999	*	*	*	*	5,433	1,217,000	64.5	(58.5 - 70.5)
≥ \$75,000	*	*	*	*	8,108	2,934,300	68.2	(64.7 - 71.7)

<sup>1</sup> Defined as respondents who never or rarely (hardly ever) were asked about their sexual history, such as about sexual partners and sexual practices, during a routine check-up at a doctor's office or clinic

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who were never or rarely asked about their sexual history during a routine check-up at a doctor's office or clinic (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 59b. Never or rarely been asked about sexual history during routine check-up among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	592	62.4	(56.4 - 68.4)	35,193	61.4	(58.7 - 64.0)
<b>Gender</b>						
Male	212	70.6	(61.0 - 80.1)	13,299	67.9	(64.0 - 71.9)
Female	380	54.6	(47.5 - 61.7)	21,894	55.5	(52.0 - 59.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	574	62.3	(56.2 - 68.4)	33,090	67.8	(65.6 - 70.1)
Black Non-Hispanic	*	*	*	683	41.1	(32.0 - 50.2)
Hispanic	*	*	*	784	53.0	(43.3 - 62.6)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	277	70.4	(62.1 - 78.7)	14,264	59.5	(54.7 - 64.3)
Some College	*	*	*	9,156	59.7	(54.4 - 65.0)
≥ College Degree	*	*	*	11,706	64.5	(60.8 - 68.2)
<b>Income</b>						
≤ \$24,999	*	*	*	8,772	48.7	(42.0 - 55.4)
\$25,000-\$49,999	*	*	*	8,932	63.7	(57.5 - 70.0)
\$50,000-\$74,999	*	*	*	5,433	64.2	(58.2 - 70.2)
≥ \$75,000	*	*	*	8,108	67.5	(64.0 - 71.0)

<sup>1</sup> Defined as respondents who never or rarely (hardly ever) were asked about their sexual history, such as about sexual partners and sexual practices, during a routine check-up at a doctor's office or clinic

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 60a. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	401	28,500	83.4	(77.7 - 89.0)	22,788	7,011,100	77.9	(75.2 - 80.5)
<b>Gender</b>								
Male	153	16,200	88.9	(80.6 - 97.3)	9,034	3,618,500	81.0	(77.2 - 84.8)
Female	248	12,300	77.0	(69.7 - 84.2)	13,754	3,392,600	74.8	(71.2 - 78.4)
<b>Age (years)</b>								
18-34	*	*	*	*	3,213	2,362,500	70.8	(65.1 - 76.6)
35-44	60	6,000	88.8	(81.6 - 96.0)	3,513	1,313,400	71.7	(65.6 - 77.7)
45-54	86	5,500	89.1	(82.8 - 95.3)	4,975	1,290,600	80.3	(75.9 - 84.8)
55-64	94	4,200	93.6	(88.3 - 98.8)	4,955	932,200	87.2	(84.1 - 90.3)
≥ 65	115	4,500	93.5	(88.4 - 98.6)	6,132	1,112,400	95.9	(94.4 - 97.5)
<b>Race/Ethnicity</b>								
White Non-Hispanic	388	26,300	83.0	(77.2 - 88.8)	21,540	4,875,200	78.3	(75.6 - 80.9)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	491	1,002,400	84.8	(77.8 - 91.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	189	13,900	86.2	(77.1 - 95.3)	8,879	2,581,700	82.1	(77.9 - 86.3)
Some College	*	*	*	*	5,953	1,599,500	77.5	(72.6 - 82.3)
≥ College Degree	118	8,500	83.7	(75.9 - 91.4)	7,928	2,824,800	75.6	(71.8 - 79.5)
<b>Income</b>								
≤ \$24,999	*	*	*	*	5,491	1,602,900	78.7	(71.9 - 85.6)
\$25,000-\$49,999	111	7,300	88.3	(80.8 - 95.8)	5,912	1,335,900	78.8	(73.8 - 83.9)
\$50,000-\$74,999	*	*	*	*	3,717	987,500	76.6	(70.4 - 82.8)
≥ \$75,000	76	7,300	90.9	(83.9 - 97.9)	5,771	2,321,600	77.6	(74.0 - 81.2)

<sup>1</sup> Defined as respondent estimating there were hardly any (0-1 out of 10) or a few (2-3 out of 10) persons their age in their community who have had a sexually transmitted disease such as gonorrhea, chlamydia, herpes or syphilis

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who believed hardly any or a few people their age in their community have had a sexually transmitted disease (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 60b. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	401	82.6	(76.8 - 88.3)	22,788	77.3	(74.7 - 79.9)
<b>Gender</b>						
Male	153	89.4	(81.4 - 97.4)	9,034	81.1	(77.4 - 84.8)
Female	248	75.0	(67.0 - 83.0)	13,754	73.6	(70.0 - 77.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	388	82.2	(76.3 - 88.1)	21,540	77.5	(74.9 - 80.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	491	84.6	(77.6 - 91.5)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	189	85.5	(76.3 - 94.7)	8,879	81.6	(77.4 - 85.7)
Some College	*	*	*	5,953	76.8	(72.0 - 81.6)
≥ College Degree	118	82.6	(74.2 - 91.0)	7,928	75.1	(71.3 - 79.0)
<b>Income</b>						
≤ \$24,999	*	*	*	5,491	77.9	(71.0 - 84.7)
\$25,000-\$49,999	111	87.1	(78.7 - 95.5)	5,912	77.9	(72.8 - 83.0)
\$50,000-\$74,999	*	*	*	3,717	76.1	(70.0 - 82.3)
≥ \$75,000	76	90.5	(83.0 - 98.0)	5,771	77.0	(73.4 - 80.6)

<sup>1</sup> Defined as respondent estimating there were hardly any (0-1 out of 10) or a few (2-3 out of 10) persons their age in their community who have had a sexually transmitted disease such as gonorrhea, chlamydia, herpes or syphilis

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 61a. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	622	41,600	86.9	(82.5 - 91.3)	36,287	12,738,300	89.7	(88.2 - 91.3)
<b>Gender</b>								
Male	230	21,100	85.4	(77.8 - 93.0)	13,893	6,104,600	89.4	(87.2 - 91.5)
Female	392	20,500	88.6	(84.7 - 92.6)	22,394	6,633,700	90.1	(87.9 - 92.2)
<b>Age (years)</b>								
18-34	*	*	*	*	3,850	4,007,400	91.3	(87.5 - 95.1)
35-44	80	8,400	96.8	(93.0 - 100)	4,875	2,578,600	93.2	(90.4 - 96.0)
45-54	133	8,200	88.4	(81.0 - 95.9)	7,607	2,416,800	90.5	(87.9 - 93.2)
55-64	147	6,300	90.1	(84.4 - 95.9)	8,411	1,876,500	92.2	(90.0 - 94.5)
≥ 65	205	6,200	74.6	(68.1 - 81.1)	11,544	1,858,900	79.7	(76.6 - 82.8)
<b>Race/Ethnicity</b>								
White Non-Hispanic	601	39,300	88.4	(84.9 - 92.0)	34,102	8,050,700	89.4	(87.7 - 91.0)
Black Non-Hispanic	*	*	*	*	702	1,822,200	91.9	(88.2 - 95.6)
Hispanic	*	*	*	*	820	1,967,200	89.9	(85.2 - 94.7)
Other Non-Hispanic	*	*	*	*	663	898,200	88.6	(79.5 - 97.6)
<b>Education</b>								
≤ High School	288	17,100	80.9	(73.1 - 88.6)	14,507	4,385,300	84.6	(81.9 - 87.2)
Some College	155	11,100	86.5	(78.0 - 95.1)	9,571	3,094,300	90.0	(86.7 - 93.4)
≥ College Degree	177	13,300	96.5	(92.7 - 100)	12,142	5,228,800	95.4	(94.0 - 96.7)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,021	2,833,500	86.5	(82.6 - 90.3)
\$25,000-\$49,999	178	10,900	84.4	(77.1 - 91.6)	9,261	2,529,900	89.5	(86.5 - 92.5)
\$50,000-\$74,999	*	*	*	*	5,644	1,740,000	91.2	(87.4 - 95.1)
≥ \$75,000	109	10,400	96.5	(91.7 - 100)	8,397	4,184,300	94.4	(92.5 - 96.3)

<sup>1</sup> Defined as believing it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who believed it is at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 61b. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	622	87.7	(83.5 - 91.9)	36,287	89.8	(88.3 - 91.4)
<b>Gender</b>						
Male	230	85.7	(78.5 - 92.8)	13,893	89.1	(86.9 - 91.3)
Female	392	89.9	(85.8 - 94.0)	22,394	90.5	(88.3 - 92.7)
<b>Race/Ethnicity</b>						
White Non-Hispanic	601	89.0	(85.5 - 92.5)	34,102	89.5	(87.8 - 91.2)
Black Non-Hispanic	*	*	*	702	91.7	(87.8 - 95.6)
Hispanic	*	*	*	820	89.6	(84.7 - 94.5)
Other Non-Hispanic	*	*	*	663	89.0	(80.5 - 97.6)
<b>Education</b>						
≤ High School	288	82.0	(74.6 - 89.5)	14,507	84.6	(81.8 - 87.4)
Some College	155	86.7	(78.0 - 95.4)	9,571	90.2	(86.9 - 93.5)
≥ College Degree	177	96.8	(93.2 - 100)	12,142	95.4	(94.0 - 96.7)
<b>Income</b>						
≤ \$24,999	*	*	*	9,021	86.5	(82.6 - 90.4)
\$25,000-\$49,999	178	84.6	(77.3 - 91.9)	9,261	89.6	(86.5 - 92.6)
\$50,000-\$74,999	*	*	*	5,644	91.1	(86.9 - 95.3)
≥ \$75,000	109	96.8	(92.4 - 100)	8,397	94.5	(92.6 - 96.4)

<sup>1</sup> Defined as believing it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 62a. Overweight among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	636	19,000	39.2	(33.2 - 45.1)	36,710	5,213,000	36.4	(34.0 - 38.8)
<b>Gender</b>								
Male	238	11,800	46.4	(36.7 - 56.2)	14,502	3,073,200	43.7	(39.9 - 47.6)
Female	398	7,200	31.1	(25.3 - 37.0)	22,208	2,139,800	29.3	(26.4 - 32.2)
<b>Age (years)</b>								
18-34	*	*	*	*	3,789	1,295,200	30.1	(24.7 - 35.6)
35-44	*	*	*	*	4,785	1,052,000	37.5	(31.7 - 43.3)
45-54	131	3,600	38.8	(29.4 - 48.3)	7,495	1,053,300	38.8	(34.1 - 43.4)
55-64	146	2,800	40.2	(31.4 - 49.0)	8,356	798,100	38.8	(34.8 - 42.7)
≥ 65	225	3,500	38.9	(31.9 - 45.9)	12,285	1,014,400	41.5	(37.6 - 45.4)
<b>Race/Ethnicity</b>								
White Non-Hispanic	614	17,700	39.3	(33.5 - 45.2)	34,501	3,282,200	36.4	(34.4 - 38.3)
Black Non-Hispanic	*	*	*	*	708	740,000	36.7	(28.3 - 45.1)
Hispanic	*	*	*	*	813	781,500	35.6	(27.1 - 44.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	300	8,100	36.8	(27.9 - 45.8)	15,079	1,889,700	35.4	(31.2 - 39.5)
Some College	*	*	*	*	9,515	1,314,500	38.7	(33.8 - 43.7)
≥ College Degree	*	*	*	*	12,062	1,996,900	36.3	(32.8 - 39.7)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,346	1,170,900	34.0	(28.2 - 39.8)
\$25,000-\$49,999	*	*	*	*	9,368	1,136,000	40.4	(34.6 - 46.2)
\$50,000-\$74,999	*	*	*	*	5,615	738,700	38.3	(32.3 - 44.3)
≥ \$75,000	*	*	*	*	8,337	1,659,100	37.5	(34.1 - 40.9)

<sup>1</sup> Defined as having a body mass index (BMI) more than 24.9 but less than 30.0. BMI is calculated as weight in kilograms divided by the square of height in meters.

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who are overweight (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 62b. Overweight among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	636	38.6	(32.5 - 44.6)	36,710	36.2	(33.8 - 38.6)
<b>Gender</b>						
Male	238	46.0	(36.3 - 55.7)	14,502	43.9	(40.1 - 47.6)
Female	398	30.3	(23.9 - 36.7)	22,208	28.9	(25.9 - 31.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	614	38.9	(33.0 - 44.9)	34,501	36.0	(34.0 - 38.0)
Black Non-Hispanic	*	*	*	708	37.0	(28.5 - 45.5)
Hispanic	*	*	*	813	36.0	(27.5 - 44.4)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	300	35.8	(26.8 - 44.8)	15,079	35.2	(31.1 - 39.4)
Some College	*	*	*	9,515	38.3	(33.3 - 43.2)
≥ College Degree	*	*	*	12,062	36.2	(32.7 - 39.8)
<b>Income</b>						
≤ \$24,999	*	*	*	9,346	33.8	(28.0 - 39.5)
\$25,000-\$49,999	*	*	*	9,368	40.6	(34.8 - 46.4)
\$50,000-\$74,999	*	*	*	5,615	38.6	(32.5 - 44.7)
≥ \$75,000	*	*	*	8,337	37.2	(33.7 - 40.6)

<sup>1</sup> Defined as having a body mass index (BMI) more than 24.9 but less than 30.0. BMI is calculated as weight in kilograms divided by the square of height in meters.

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 63a. Obesity among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	636	13,900	28.6	(23.4 - 33.9)	36,710	3,327,400	23.2	(21.2 - 25.3)
<b>Gender</b>								
Male	238	6,700	26.3	(17.9 - 34.7)	14,502	1,505,800	21.4	(19.0 - 23.9)
Female	398	7,200	31.2	(25.2 - 37.1)	22,208	1,821,600	25.0	(21.8 - 28.1)
<b>Age (years)</b>								
18-34	*	*	*	*	3,789	730,600	17.0	(12.4 - 21.6)
35-44	*	*	*	*	4,785	662,200	23.6	(19.4 - 27.9)
45-54	131	2,700	28.8	(19.2 - 38.4)	7,495	816,800	30.1	(25.3 - 34.8)
55-64	146	2,200	32.2	(24.0 - 40.4)	8,356	563,900	27.4	(23.7 - 31.1)
≥ 65	225	2,300	25.1	(18.9 - 31.3)	12,285	554,000	22.7	(19.2 - 26.1)
<b>Race/Ethnicity</b>								
White Non-Hispanic	614	12,800	28.4	(23.3 - 33.4)	34,501	2,062,700	22.9	(21.2 - 24.6)
Black Non-Hispanic	*	*	*	*	708	703,100	34.9	(26.1 - 43.7)
Hispanic	*	*	*	*	813	456,400	20.8	(14.3 - 27.2)
Other Non-Hispanic	*	*	*	*	688	105,200	9.6	(4.0 - 15.3)
<b>Education</b>								
≤ High School	300	7,100	32.3	(24.4 - 40.3)	15,079	1,561,600	29.2	(25.3 - 33.2)
Some College	157	3,400	26.8	(17.3 - 36.2)	9,515	792,800	23.4	(19.4 - 27.3)
≥ College Degree	*	*	*	*	12,062	915,000	16.6	(14.4 - 18.8)
<b>Income</b>								
≤ \$24,999	182	3,000	25.9	(16.6 - 35.2)	9,346	974,800	28.3	(22.6 - 34.0)
\$25,000-\$49,999	180	3,900	29.6	(20.9 - 38.3)	9,368	584,800	20.8	(17.1 - 24.5)
\$50,000-\$74,999	*	*	*	*	5,615	475,300	24.6	(20.0 - 29.3)
≥ \$75,000	*	*	*	*	8,337	867,700	19.6	(17.0 - 22.3)

<sup>1</sup> Defined as having a body mass index (BMI) of 30.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who are obese (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 63b. Obesity among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	636	29.3	(23.8 - 34.8)	36,710	23.1	(21.0 - 25.1)
<b>Gender</b>						
Male	238	26.9	(18.3 - 35.5)	14,502	21.5	(19.0 - 24.0)
Female	398	31.9	(25.3 - 38.5)	22,208	24.6	(21.5 - 27.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	614	28.8	(23.5 - 34.1)	34,501	22.8	(21.0 - 24.5)
Black Non-Hispanic	*	*	*	708	34.8	(25.9 - 43.6)
Hispanic	*	*	*	813	20.5	(14.1 - 26.9)
Other Non-Hispanic	*	*	*	688	9.5	(3.8 - 15.2)
<b>Education</b>						
≤ High School	300	33.2	(24.9 - 41.4)	15,079	29.4	(25.4 - 33.3)
Some College	157	27.4	(17.6 - 37.2)	9,515	23.1	(19.2 - 27.1)
≥ College Degree	*	*	*	12,062	16.4	(14.2 - 18.5)
<b>Income</b>						
≤ \$24,999	182	25.5	(16.0 - 35.0)	9,346	28.3	(22.6 - 34.1)
\$25,000-\$49,999	180	30.8	(21.5 - 40.1)	9,368	21.0	(17.2 - 24.8)
\$50,000-\$74,999	*	*	*	5,615	24.0	(19.4 - 28.7)
≥ \$75,000	*	*	*	8,337	19.3	(16.7 - 21.9)

<sup>1</sup> Defined as having a body mass index (BMI) of 30.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 64a. Overweight or obesity among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	636	32,800	67.8	(62.5 - 73.1)	36,710	8,540,500	59.6	(57.0 - 62.2)
<b>Gender</b>								
Male	238	18,500	72.7	(64.3 - 81.1)	14,502	4,579,100	65.2	(61.0 - 69.4)
Female	398	14,300	62.3	(56.3 - 68.3)	22,208	3,961,400	54.3	(51.0 - 57.6)
<b>Age (years)</b>								
18-34	*	*	*	*	3,789	2,025,800	47.1	(40.8 - 53.5)
35-44	*	*	*	*	4,785	1,714,200	61.1	(55.6 - 66.6)
45-54	131	6,300	67.6	(58.4 - 76.9)	7,495	1,870,100	68.8	(64.6 - 73.1)
55-64	146	5,000	72.4	(64.4 - 80.4)	8,356	1,362,000	66.2	(61.9 - 70.4)
≥ 65	225	5,800	64.0	(57.3 - 70.8)	12,285	1,568,400	64.1	(60.6 - 67.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	614	30,500	67.7	(62.3 - 73.0)	34,501	5,344,900	59.3	(57.2 - 61.3)
Black Non-Hispanic	*	*	*	*	708	1,443,100	71.6	(62.5 - 80.7)
Hispanic	*	*	*	*	813	1,237,900	56.4	(46.5 - 66.2)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	300	15,300	69.2	(60.9 - 77.5)	15,079	3,451,400	64.6	(59.8 - 69.4)
Some College	157	8,700	67.8	(58.0 - 77.5)	9,515	2,107,300	62.1	(57.0 - 67.2)
≥ College Degree	178	8,900	65.4	(55.8 - 75.0)	12,062	2,911,900	52.9	(49.2 - 56.6)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,346	2,145,700	62.3	(55.7 - 68.9)
\$25,000-\$49,999	180	9,000	67.4	(57.8 - 76.9)	9,368	1,720,700	61.2	(54.5 - 67.9)
\$50,000-\$74,999	*	*	*	*	5,615	1,214,000	62.9	(57.1 - 68.8)
≥ \$75,000	107	7,700	74.1	(64.2 - 84.0)	8,337	2,526,700	57.1	(53.6 - 60.6)

<sup>1</sup> Defined as having a body mass index (BMI) of 25.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who are overweight or obese (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 64b. Overweight or obesity among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	636	67.8	(62.4 - 73.3)	36,710	59.3	(56.8 - 61.9)
<b>Gender</b>						
Male	238	72.9	(64.6 - 81.3)	14,502	65.4	(61.3 - 69.4)
Female	398	62.2	(55.7 - 68.7)	22,208	53.5	(50.1 - 56.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	614	67.7	(62.2 - 73.2)	34,501	58.8	(56.7 - 60.8)
Black Non-Hispanic	*	*	*	708	71.8	(63.0 - 80.6)
Hispanic	*	*	*	813	56.5	(46.8 - 66.1)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	300	68.9	(60.3 - 77.5)	15,079	64.6	(59.9 - 69.3)
Some College	157	68.5	(58.7 - 78.2)	9,515	61.4	(56.4 - 66.5)
≥ College Degree	178	65.4	(55.5 - 75.3)	12,062	52.6	(48.9 - 56.3)
<b>Income</b>						
≤ \$24,999	*	*	*	9,346	62.1	(55.6 - 68.6)
\$25,000-\$49,999	180	67.2	(57.2 - 77.2)	9,368	61.6	(55.1 - 68.1)
\$50,000-\$74,999	*	*	*	5,615	62.6	(56.7 - 68.6)
≥ \$75,000	107	75.0	(65.2 - 84.9)	8,337	56.5	(52.9 - 60.0)

<sup>1</sup> Defined as having a body mass index (BMI) of 25.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 65a. Received advice about weight by a health professional among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	634	9,900	20.4	(16.6 - 24.3)	36,729	3,929,700	27.4	(25.1 - 29.7)
<b>Gender</b>								
Male	238	4,300	16.9	(11.6 - 22.3)	14,499	1,654,800	23.6	(20.6 - 26.6)
Female	396	5,600	24.3	(19.2 - 29.4)	22,230	2,274,900	31.0	(27.7 - 34.3)
<b>Age (years)</b>								
18-34	*	*	*	*	3,796	895,500	20.7	(15.3 - 26.2)
35-44	79	1,900	21.6	(11.7 - 31.5)	4,779	722,700	25.8	(20.9 - 30.6)
45-54	131	2,100	22.1	(13.8 - 30.4)	7,490	910,900	33.5	(29.0 - 38.1)
55-64	145	2,200	31.5	(23.4 - 39.5)	8,351	741,100	36.1	(32.0 - 40.3)
≥ 65	225	2,600	28.4	(21.9 - 35.0)	12,313	659,500	26.9	(23.0 - 30.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	612	9,600	21.5	(17.5 - 25.4)	34,502	2,386,200	26.5	(24.6 - 28.4)
Black Non-Hispanic	*	*	*	*	709	615,700	30.4	(22.0 - 38.8)
Hispanic	*	*	*	*	830	555,400	25.1	(17.8 - 32.4)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	300	4,300	19.7	(14.2 - 25.2)	15,116	1,469,700	27.5	(23.6 - 31.4)
Some College	157	2,800	21.7	(14.0 - 29.3)	9,515	946,800	27.8	(23.2 - 32.4)
≥ College Degree	176	2,700	20.1	(12.8 - 27.4)	12,041	1,445,500	26.2	(23.0 - 29.4)
<b>Income</b>								
≤ \$24,999	182	1,800	15.8	(9.7 - 21.9)	9,360	1,079,400	31.4	(25.3 - 37.5)
\$25,000-\$49,999	179	3,200	24.4	(16.8 - 32.1)	9,365	696,900	24.6	(19.8 - 29.4)
\$50,000-\$74,999	*	*	*	*	5,603	507,100	26.3	(21.6 - 31.0)
≥ \$75,000	107	1,800	16.9	(9.5 - 24.3)	8,334	1,207,100	27.3	(24.2 - 30.5)

<sup>1</sup> Defined as being advised by a doctor, nurse or other health professional about their weight (lose, gain or maintain)

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults provided advice by a health professional about their weight (lose, gain or maintain) (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 65b. Received advice about weight by a health professional among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	634	20.3	(16.3 - 24.2)	36,729	27.1	(24.8 - 29.4)
<b>Gender</b>						
Male	238	17.1	(11.6 - 22.5)	14,499	23.4	(20.5 - 26.3)
Female	396	23.8	(18.3 - 29.4)	22,230	30.6	(27.3 - 34.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	612	21.3	(17.1 - 25.4)	34,502	26.1	(24.2 - 28.0)
Black Non-Hispanic	*	*	*	709	30.4	(21.9 - 38.9)
Hispanic	*	*	*	830	24.9	(17.7 - 32.2)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	300	19.2	(13.5 - 24.8)	15,116	27.3	(23.4 - 31.2)
Some College	157	22.5	(14.2 - 30.7)	9,515	27.4	(22.9 - 32.0)
≥ College Degree	176	19.6	(12.2 - 27.0)	12,041	25.9	(22.7 - 29.0)
<b>Income</b>						
≤ \$24,999	182	15.4	(9.1 - 21.8)	9,360	31.0	(24.9 - 37.0)
\$25,000-\$49,999	179	24.3	(16.4 - 32.3)	9,365	24.9	(20.0 - 29.8)
\$50,000-\$74,999	*	*	*	5,603	25.9	(21.2 - 30.6)
≥ \$75,000	107	17.0	(9.3 - 24.6)	8,334	26.9	(23.7 - 30.0)

<sup>1</sup> Defined as being advised by a doctor, nurse or other health professional about their weight (lose, gain or maintain)

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 66a. Received advice to lose weight by a health professional among adults provided weight advice<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	166	8,600	87.7	(81.6 - 93.7)	9,724	3,412,900	86.8	(83.9 - 89.8)
<b>Gender</b>								
Male	62	3,800	87.7	(78.0 - 97.4)	3,807	1,468,700	88.8	(85.7 - 91.8)
Female	104	4,900	87.6	(80.0 - 95.3)	5,917	1,944,200	85.5	(80.8 - 90.1)
<b>Age (years)</b>								
18-34	*	*	*	*	681	779,700	87.1	(80.6 - 93.6)
35-44	*	*	*	*	1,215	622,100	86.1	(76.9 - 95.2)
45-54	*	*	*	*	2,238	813,300	89.3	(84.4 - 94.2)
55-64	50	2,100	96.4	(90.9 - 100)	2,697	645,500	87.1	(81.2 - 93.0)
≥ 65	*	*	*	*	2,893	552,300	83.7	(76.9 - 90.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	163	8,400	87.4	(81.2 - 93.5)	9,082	2,076,700	87.0	(84.5 - 89.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	251	483,600	87.1	(78.5 - 95.7)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	*	*	*	*	3,962	1,239,700	84.3	(78.6 - 90.1)
Some College	*	*	*	*	2,633	814,800	86.1	(80.1 - 92.0)
≥ College Degree	*	*	*	*	3,115	1,290,900	89.3	(85.3 - 93.3)
<b>Income</b>								
≤ \$24,999	*	*	*	*	2,539	922,100	85.4	(78.7 - 92.1)
\$25,000-\$49,999	52	3,000	93.4	(86.3 - 100)	2,482	594,100	85.2	(78.4 - 92.1)
\$50,000-\$74,999	*	*	*	*	1,521	446,100	88.0	(83.2 - 92.7)
≥ \$75,000	*	*	*	*	2,270	1,073,200	88.9	(83.7 - 94.1)

<sup>1</sup> Defined as provided advice by a health professional about their weight and advised to lose weight

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults provided advice by a health professional about their weight and advised to lose weight (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 66b. Received advice to lose weight by a health professional among adults provided weight advice<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	166	87.3	(80.7 - 93.9)	9,724	86.9	(83.8 - 89.9)
<b>Gender</b>						
Male	*	*	*	3,807	88.7	(85.7 - 91.8)
Female	104	87.9	(79.8 - 96.0)	5,917	85.5	(80.7 - 90.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	163	87.0	(80.2 - 93.7)	9,082	87.0	(84.4 - 89.6)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	251	87.6	(79.3 - 95.8)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	3,962	84.6	(78.7 - 90.5)
Some College	*	*	*	2,633	85.6	(79.5 - 91.7)
≥ College Degree	*	*	*	3,115	89.4	(85.4 - 93.3)
<b>Income</b>						
≤ \$24,999	*	*	*	2,539	85.7	(79.0 - 92.3)
\$25,000-\$49,999	52	93.2	(85.7 - 100)	2,482	85.3	(78.4 - 92.3)
\$50,000-\$74,999	*	*	*	1,521	88.0	(83.1 - 92.8)
≥ \$75,000	*	*	*	2,270	88.7	(83.1 - 94.2)

<sup>1</sup> Defined as provided advice by a health professional about their weight and advised to lose weight

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 67a. No leisure-time physical activity among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	657	12,100	24.4	(19.0 - 29.8)	38,128	3,541,700	23.9	(21.6 - 26.1)
<b>Gender</b>								
Male	241	7,200	28.3	(18.8 - 37.8)	14,602	1,482,500	20.9	(17.8 - 24.0)
Female	416	4,900	20.2	(16.1 - 24.3)	23,526	2,059,200	26.6	(23.4 - 29.7)
<b>Age (years)</b>								
18-34	*	*	*	*	3,938	982,900	22.0	(16.6 - 27.5)
35-44	*	*	*	*	4,988	556,000	19.3	(15.0 - 23.6)
45-54	135	2,000	21.4	(13.0 - 29.9)	7,827	759,000	26.8	(21.7 - 31.9)
55-64	153	1,900	26.6	(19.1 - 34.1)	8,765	464,900	21.7	(18.2 - 25.2)
≥ 65	228	3,000	32.5	(26.0 - 39.0)	12,610	778,800	30.7	(27.3 - 34.2)
<b>Race/Ethnicity</b>								
White Non-Hispanic	635	11,200	24.2	(19.1 - 29.4)	35,816	1,892,800	20.3	(18.6 - 22.0)
Black Non-Hispanic	*	*	*	*	739	690,000	32.0	(23.4 - 40.6)
Hispanic	*	*	*	*	860	600,800	26.7	(19.1 - 34.4)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	311	8,000	35.6	(26.5 - 44.6)	15,683	1,810,200	32.5	(28.4 - 36.5)
Some College	*	*	*	*	9,906	814,200	23.0	(18.6 - 27.3)
≥ College Degree	183	1,300	9.5	(5.5 - 13.5)	12,458	841,000	14.9	(11.9 - 18.0)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,640	1,332,200	38.0	(31.7 - 44.3)
\$25,000-\$49,999	*	*	*	*	9,658	648,000	22.3	(18.3 - 26.3)
\$50,000-\$74,999	*	*	*	*	5,773	430,700	21.8	(16.6 - 27.0)
≥ \$75,000	112	600	5.3	(1.8 - 8.9)	8,565	560,800	12.5	(10.3 - 14.6)

<sup>1</sup> Defined as having not participated in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have not participated in any physical activities or exercises (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)



**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 67b. No leisure-time physical activity among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	657	23.4	(18.1 - 28.7)	38,128	23.7	(21.5 - 25.9)
<b>Gender</b>						
Male	241	28.0	(18.7 - 37.3)	14,602	20.9	(17.8 - 23.9)
Female	416	18.5	(14.3 - 22.6)	23,526	26.3	(23.1 - 29.5)
<b>Race/Ethnicity</b>						
White Non-Hispanic	635	23.3	(18.2 - 28.5)	35,816	20.0	(18.3 - 21.7)
Black Non-Hispanic	*	*	*	739	32.0	(23.3 - 40.6)
Hispanic	*	*	*	860	26.9	(19.2 - 34.5)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	311	34.3	(25.1 - 43.4)	15,683	32.3	(28.3 - 36.4)
Some College	161	19.9	(10.1 - 29.8)	9,906	22.8	(18.5 - 27.2)
≥ College Degree	183	9.2	(5.2 - 13.2)	12,458	14.9	(11.8 - 17.9)
<b>Income</b>						
≤ \$24,999	*	*	*	9,640	37.9	(31.6 - 44.2)
\$25,000-\$49,999	*	*	*	9,658	22.0	(18.0 - 26.0)
\$50,000-\$74,999	*	*	*	5,773	21.6	(16.3 - 26.8)
≥ \$75,000	112	5.2	(1.7 - 8.8)	8,565	12.6	(10.3 - 14.9)

<sup>1</sup> Defined as having not participated in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 68a. Consumption of 5 or more servings of fruits and vegetables per day among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	659	12,600	25.3	(20.6 - 30.1)	38,203	4,023,400	27.0	(24.9 - 29.2)
<b>Gender</b>								
Male	241	5,500	21.7	(14.0 - 29.3)	14,634	1,603,900	22.6	(19.4 - 25.8)
Female	418	7,100	29.2	(23.9 - 34.6)	23,569	2,419,500	31.1	(28.2 - 34.0)
<b>Age (years)</b>								
18-34	*	*	*	*	3,940	1,261,700	28.3	(22.7 - 33.9)
35-44	83	1,900	21.5	(11.7 - 31.4)	4,989	688,000	23.9	(19.5 - 28.3)
45-54	135	2,600	27.2	(18.4 - 36.0)	7,837	746,800	26.2	(22.3 - 30.1)
55-64	153	1,800	24.8	(17.2 - 32.4)	8,777	579,100	27.0	(23.6 - 30.5)
≥ 65	230	2,700	29.7	(23.3 - 36.2)	12,660	747,800	29.4	(26.0 - 32.8)
<b>Race/Ethnicity</b>								
White Non-Hispanic	637	12,100	26.1	(21.2 - 30.9)	35,884	2,538,000	27.2	(25.4 - 28.9)
Black Non-Hispanic	*	*	*	*	740	572,400	26.5	(19.2 - 33.9)
Hispanic	*	*	*	*	864	672,200	29.8	(21.0 - 38.5)
Other Non-Hispanic	*	*	*	*	715	240,800	21.4	(12.0 - 30.8)
<b>Education</b>								
≤ High School	313	3,600	15.9	(10.8 - 21.1)	15,726	1,309,800	23.4	(19.3 - 27.4)
Some College	*	*	*	*	9,924	964,300	27.2	(23.0 - 31.3)
≥ College Degree	183	4,600	32.3	(22.6 - 42.1)	12,471	1,747,500	31.0	(27.9 - 34.1)
<b>Income</b>								
≤ \$24,999	191	2,300	19.1	(10.8 - 27.5)	9,667	831,500	23.7	(18.6 - 28.7)
\$25,000-\$49,999	184	3,200	23.9	(15.4 - 32.3)	9,669	917,200	31.6	(25.6 - 37.6)
\$50,000-\$74,999	*	*	*	*	5,777	501,000	25.3	(21.4 - 29.3)
≥ \$75,000	*	*	*	*	8,569	1,260,100	28.0	(25.0 - 31.0)

<sup>1</sup> Defined as having consumed 5 or more servings of fruits and vegetables in an average day

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who consume 5 or more servings of fruits and vegetables per day (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 68b. Consumption of 5 or more servings of fruits and vegetables per day among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	659	24.9	(20.1 - 29.7)	38,203	27.1	(24.9 - 29.2)
<b>Gender</b>						
Male	241	21.7	(14.1 - 29.2)	14,634	22.7	(19.6 - 25.9)
Female	418	28.4	(22.6 - 34.1)	23,569	31.0	(28.1 - 34.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	637	25.6	(20.6 - 30.5)	35,884	27.1	(25.4 - 28.9)
Black Non-Hispanic	*	*	*	740	26.8	(19.4 - 34.2)
Hispanic	*	*	*	864	29.5	(20.9 - 38.1)
Other Non-Hispanic	*	*	*	715	21.8	(12.2 - 31.5)
<b>Education</b>						
≤ High School	313	16.0	(10.5 - 21.6)	15,726	23.3	(19.3 - 27.3)
Some College	*	*	*	9,924	27.1	(22.9 - 31.2)
≥ College Degree	183	30.9	(21.1 - 40.6)	12,471	31.2	(28.0 - 34.3)
<b>Income</b>						
≤ \$24,999	191	18.3	(10.1 - 26.6)	9,667	23.9	(18.8 - 28.9)
\$25,000-\$49,999	184	23.7	(14.9 - 32.5)	9,669	32.0	(26.0 - 38.0)
\$50,000-\$74,999	*	*	*	5,777	25.0	(21.0 - 28.9)
≥ \$75,000	*	*	*	8,569	28.0	(24.9 - 31.1)

<sup>1</sup> Defined as having consumed 5 or more servings of fruits and vegetables in an average day

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 69a. Had a fall within the past 3 months among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	513	5,300	20.7	(16.6 - 24.8)	28,783	1,244,800	17.0	(14.9 - 19.0)
<b>Gender</b>								
Male	193	2,600	21.6	(14.7 - 28.6)	10,966	489,400	14.4	(11.9 - 16.8)
Female	320	2,700	19.8	(15.1 - 24.6)	17,817	755,300	19.2	(16.1 - 22.3)
<b>Age (years)</b>								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	134	2,400	26.3	(17.6 - 34.9)	7,741	445,500	16.0	(12.0 - 20.0)
55-64	150	1,500	20.6	(13.7 - 27.6)	8,454	336,000	16.5	(13.4 - 19.6)
≥ 65	229	1,400	15.1	(10.0 - 20.1)	12,588	463,300	18.4	(15.2 - 21.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	499	5,000	20.5	(16.4 - 24.7)	27,456	869,000	16.6	(15.1 - 18.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	424	76,100	11.0	(4.5 - 17.4)
Other Non-Hispanic	*	*	*	*	445	44,300	10.0	(3.1 - 16.8)
<b>Education</b>								
≤ High School	258	2,500	19.5	(13.8 - 25.2)	12,417	493,000	17.3	(13.3 - 21.3)
Some College	120	1,200	20.2	(11.1 - 29.3)	7,232	299,400	18.7	(14.6 - 22.7)
≥ College Degree	134	1,600	23.6	(15.7 - 31.5)	9,086	441,500	15.4	(12.9 - 17.9)
<b>Income</b>								
≤ \$24,999	165	1,900	28.8	(20.1 - 37.5)	7,824	423,900	25.0	(18.5 - 31.6)
\$25,000-\$49,999	150	1,200	16.0	(9.4 - 22.6)	7,445	223,600	16.0	(12.8 - 19.2)
\$50,000-\$74,999	*	*	*	*	4,141	151,200	14.6	(10.8 - 18.4)
≥ \$75,000	79	700	14.8	(6.9 - 22.7)	5,874	287,100	12.5	(10.0 - 15.1)

<sup>1</sup> Defined as respondent unintentionally coming to rest on the ground or another lower level one or more times within the past 3 months

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have had a fall within the past 3 months (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 69b. Had a fall within the past 3 months among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	513	20.7	(16.5 - 25.0)	28,783	17.3	(15.0 - 19.5)
<b>Gender</b>						
Male	193	21.4	(14.5 - 28.3)	10,966	15.0	(12.1 - 17.8)
Female	320	20.0	(15.2 - 24.9)	17,817	19.4	(16.0 - 22.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	499	20.6	(16.3 - 24.9)	27,456	16.9	(15.2 - 18.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	424	10.7	(4.2 - 17.1)
Other Non-Hispanic	*	*	*	445	10.0	(2.9 - 17.0)
<b>Education</b>						
≤ High School	258	19.7	(13.7 - 25.6)	12,417	17.4	(13.0 - 21.7)
Some College	120	20.2	(10.9 - 29.5)	7,232	19.0	(14.7 - 23.2)
≥ College Degree	134	23.4	(15.4 - 31.4)	9,086	16.0	(13.1 - 18.8)
<b>Income</b>						
≤ \$24,999	165	29.4	(20.3 - 38.5)	7,824	25.3	(18.3 - 32.4)
\$25,000-\$49,999	150	15.9	(9.3 - 22.6)	7,445	15.9	(12.6 - 19.2)
\$50,000-\$74,999	*	*	*	4,141	14.7	(10.6 - 18.8)
≥ \$75,000	79	14.0	(6.3 - 21.7)	5,874	12.9	(9.9 - 15.8)

<sup>1</sup> Defined as respondent unintentionally coming to rest on the ground or another lower level one or more times within the past 3 months

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Washington County**

**Table 70a. Had a fall that caused injury within the past 3 months among adults<sup>1</sup> in Washington County and New York State**

	Washington County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	513	1,300	5.1	(3.1 - 7.1)	28,760	335,700	4.6	(3.7 - 5.5)
<b>Gender</b>								
Male	*	*	*	*	10,957	120,000	3.5	(2.5 - 4.6)
Female	320	900	7.0	(4.0 - 10.0)	17,803	215,600	5.5	(4.1 - 6.9)
<b>Age (years)</b>								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	7,737	115,100	4.1	(2.9 - 5.4)
55-64	*	*	*	*	8,449	90,300	4.4	(2.7 - 6.1)
≥ 65	229	500	5.1	(2.0 - 8.3)	12,574	130,300	5.2	(3.4 - 6.9)
<b>Race/Ethnicity</b>								
White Non-Hispanic	499	1,300	5.2	(3.2 - 7.2)	27,433	270,800	5.2	(4.2 - 6.1)
Black Non-Hispanic	*	*	*	*	458	47,900	4.9	(0.9 - 9.0)
Hispanic	*	*	*	*	424	12,400	1.8	(0.2 - 3.3)
Other Non-Hispanic	*	*	*	*	445	4,600	1.0	(0.4 - 1.6)
<b>Education</b>								
≤ High School	258	700	5.3	(2.4 - 8.3)	12,404	148,900	5.2	(3.6 - 6.8)
Some College	*	*	*	*	7,228	85,400	5.3	(3.1 - 7.5)
≥ College Degree	*	*	*	*	9,081	101,300	3.5	(2.4 - 4.6)
<b>Income</b>								
≤ \$24,999	165	500	7.3	(2.8 - 11.8)	7,812	90,200	5.3	(3.6 - 7.1)
\$25,000-\$49,999	*	*	*	*	7,442	93,200	6.7	(3.9 - 9.4)
\$50,000-\$74,999	*	*	*	*	4,139	37,000	3.6	(1.8 - 5.3)
≥ \$75,000	*	*	*	*	5,872	54,700	2.4	(1.5 - 3.3)

<sup>1</sup> Defined as as respondent unintentionally coming to rest on the ground or another lower level and caused respondent to limit regular activities for at least a day or to go see a doctor one or more times within the past 3 months

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have had a fall that caused an injury within the past 3 months (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Washington County**

**Table 70b. Had a fall that caused injury within the past 3 months among adults<sup>1</sup> in Washington County and New York State - Age-Adjusted<sup>2</sup>**

	Washington County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	513	5.0	(3.1 - 7.0)	28,760	4.6	(3.6 - 5.5)
<b>Gender</b>						
Male	*	*	*	10,957	3.6	(2.4 - 4.7)
Female	320	7.0	(4.0 - 10.1)	17,803	5.5	(4.1 - 6.9)
<b>Race/Ethnicity</b>						
White Non-Hispanic	499	5.1	(3.1 - 7.2)	27,433	5.2	(4.2 - 6.2)
Black Non-Hispanic	*	*	*	458	4.7	(0.7 - 8.7)
Hispanic	*	*	*	424	1.8	(0.1 - 3.4)
Other Non-Hispanic	*	*	*	445	1.0	(0.4 - 1.6)
<b>Education</b>						
≤ High School	258	5.3	(2.3 - 8.2)	12,404	5.2	(3.5 - 6.8)
Some College	*	*	*	7,228	5.3	(3.1 - 7.5)
≥ College Degree	*	*	*	9,081	3.6	(2.4 - 4.7)
<b>Income</b>						
≤ \$24,999	165	7.3	(2.6 - 12.0)	7,812	5.1	(3.3 - 7.0)
\$25,000-\$49,999	*	*	*	7,442	6.6	(3.9 - 9.2)
\$50,000-\$74,999	*	*	*	4,139	3.7	(1.7 - 5.8)
≥ \$75,000	*	*	*	5,872	2.4	(1.4 - 3.3)

<sup>1</sup> Defined as as respondent unintentionally coming to rest on the ground or another lower level and caused respondent to limit regular activities for at least a day or to go see a doctor one or more times within the past 3 months

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)