

New York State Expanded Behavioral Risk Factor Surveillance System

Wyoming County

**Final Report
July 2008 - June 2009**

**New York State Department of Health
December 2009**

Contents

Introduction	12
Table 1. Wyoming County and New York State adult US Census Bureau population estimates, 2007	13
General Health	
Table 2a. Fair or poor general health among adults in Wyoming County and New York State	14
Table 2b. Fair or poor general health among adults in Wyoming County and New York State - Age-adjusted	15
Table 3a. Poor physical health 14 or more days within the past month among adults in Wyoming County and New York State	16
Table 3b. Poor physical health 14 or more days within the past month among adults in Wyoming County and New York State - Age-adjusted	17
Table 4a. Poor mental health 14 or more days within the past month among adults in Wyoming County and New York State	18
Table 4b. Poor mental health 14 or more days within the past month among adults in Wyoming County and New York State - Age-adjusted	19
Table 5a. Poor physical or mental health 14 or more days within the past month among adults in Wyoming County and New York State	20
Table 5b. Poor physical or mental health 14 or more days within the past month among adults in Wyoming County and New York State - Age-adjusted	21
Quality Health Care	
Table 6a. Health insurance among adults in Wyoming County and New York State	22
Table 6b. Health insurance among adults in Wyoming County and New York State - Age-adjusted	23
Table 7a. Health insurance among adults aged 18-64 in Wyoming County and New York State	24
Table 7b. Health insurance among adults aged 18-64 in Wyoming County and New York State - Age-adjusted	25

Contents

Table 8a. No health insurance among adults in Wyoming County and New York State	26
Table 8b. No health insurance among adults in Wyoming County and New York State - Age-adjusted	27
Table 9a. No health insurance among adults aged 18-64 in Wyoming County and New York State	28
Table 9b. No health insurance among adults aged 18-64 in Wyoming County and New York State - Age-adjusted	29
Table 10a. Adults with regular health care provider in Wyoming County and New York State	30
Table 10b. Adults with regular health care provider in Wyoming County and New York State - Age-adjusted	31
Table 11a. Cost prevented visit to doctor within the past year among adults in Wyoming County and New York State	32
Table 11b. Cost prevented visit to doctor within the past year among adults in Wyoming County and New York State - Age-adjusted	33
Table 12a. Visited a doctor for routine checkup within the past year among adults in Wyoming County and New York State	34
Table 12b. Visited a doctor for routine checkup within the past year among adults in Wyoming County and New York State - Age-adjusted	35
Table 13a. Visited a doctor for routine checkup within the past 2 years among adults in Wyoming County and New York State	36
Table 13b. Visited a doctor for routine checkup within the past 2 years among adults in Wyoming County and New York State - Age-adjusted	37
Table 14a. Dentist visit within the past year among adults in Wyoming County and New York State	38
Table 14b. Dentist visit within the past year among adults in Wyoming County and New York State - Age-adjusted	39
Table 15a. Had teeth cleaned within the past year among adults in Wyoming County and New York State	40
Table 15b. Had teeth cleaned within the past year among adults in Wyoming County and New York State - Age-adjusted	41

Contents

Table 16a. Permanent teeth extracted due to decay or gum disease among adults in Wyoming County and New York State	42
Table 16b. Permanent teeth extracted due to decay or gum disease among adults in Wyoming County and New York State - Age-adjusted	43
Table 17a. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older in Wyoming County and New York State	44
Table 17b. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older in Wyoming County and New York State - Age-adjusted	45
Table 18a. Flu shot within the past 12 months in Wyoming County and New York State	46
Table 18b. Flu shot within the past 12 months in Wyoming County and New York State - Age-adjusted	47
Table 19a. Flu vaccine that was sprayed in nose within the past 12 months in Wyoming County and New York State	48
Table 19b. Flu vaccine that was sprayed in nose within the past 12 months in Wyoming County and New York State - Age-adjusted	49
Table 20a. Flu shot or flu vaccine sprayed in nose within the past 12 months in Wyoming County and New York State	50
Table 20b. Flu shot or flu vaccine sprayed in nose within the past 12 months in Wyoming County and New York State - Age-adjusted	51
Table 21a. Flu shot among adults aged 65 and older in Wyoming County and New York State	52
Table 21b. Flu shot among adults aged 65 and older in Wyoming County and New York State - Age-adjusted	53
Table 22a. Pneumonia shot or pneumococcal vaccine in Wyoming County and New York State	54
Table 22b. Pneumonia shot or pneumococcal vaccine in Wyoming County and New York State - Age-adjusted	55
Table 23a. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older in Wyoming County and New York State	56
Table 23b. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older in Wyoming County and New York State - Age-adjusted	57

Contents

Cancer Screening

Table 24a. Ever had a mammogram among women aged 40 and older in Wyoming County and New York State	58
Table 24b. Ever had a mammogram among women aged 40 and older in Wyoming County and New York State - Age-adjusted	59
Table 25a. Had mammogram within the past 2 years among women aged 40 and older in Wyoming County and New York State	60
Table 25b. Had mammogram within the past 2 years among women aged 40 and older in Wyoming County and New York State - Age-adjusted	61
Table 26a. Had mammogram within the past 2 years among women aged 50 and older in Wyoming County and New York State	62
Table 26b. Had mammogram within the past 2 years among women aged 50 and older in Wyoming County and New York State - Age-adjusted	63
Table 27a. Ever had a Pap test among women in Wyoming County and New York State	64
Table 27b. Ever had a Pap test among women in Wyoming County and New York State - Age-adjusted	65
Table 28a. Had a Pap test within the past 3 years among women in Wyoming County and New York State	66
Table 28b. Had a Pap test within the past 3 years among women in Wyoming County and New York State - Age-adjusted	67
Table 29a. Ever had a digital rectal exam among men aged 40 and older in Wyoming County and New York State	68
Table 29b. Ever had a digital rectal exam among men aged 40 and older in Wyoming County and New York State - Age-adjusted	69
Table 30a. Had digital rectal exam within the past 2 years among men aged 40 and older in Wyoming County and New York State	70
Table 30b. Had digital rectal exam within the past 2 years among men aged 40 and older in Wyoming County and New York State - Age-adjusted	71

Contents

Table 31a. Ever had a prostate specific antigen test among men aged 40 and older in Wyoming County and New York State	72
Table 31b. Ever had a prostate specific antigen test among men aged 40 and older in Wyoming County and New York State - Age-adjusted	73
Table 32a. Had prostate specific antigen test within the past 2 years among men aged 40 and older in Wyoming County and New York State	74
Table 32b. Had prostate specific antigen test within the past 2 years among men aged 40 and older in Wyoming County and New York State - Age-adjusted	75
Table 33a. Prostate cancer among men aged 40 and older in Wyoming County and New York State	76
Table 33b. Prostate cancer among men aged 40 and older in Wyoming County and New York State - Age-adjusted	77
Table 34a. Home blood stool test ever used among adults aged 50 and older in Wyoming County and New York State	78
Table 34b. Home blood stool test ever used among adults aged 50 and older in Wyoming County and New York State - Age-adjusted	79
Table 35a. Home blood stool test used within the past year among adults aged 50 and older in Wyoming County and New York State	80
Table 35b. Home blood stool test used within the past year among adults aged 50 and older in Wyoming County and New York State - Age-adjusted	81
Table 36a. Home blood stool test used within the past 2 years among adults aged 50 and older in Wyoming County and New York State	82
Table 36b. Home blood stool test used within the past 2 years among adults aged 50 and older in Wyoming County and New York State - Age-adjusted	83
Table 37a. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in Wyoming County and New York State	84
Table 37b. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in Wyoming County and New York State - Age-adjusted	85
Table 38a. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Wyoming County and New York State	86

Contents

Table 38b. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Wyoming County and New York State - Age-adjusted	87
Table 39a. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Wyoming County and New York State	88
Table 39b. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Wyoming County and New York State - Age-adjusted	89
Chronic Diseases	
Table 40a. High blood pressure among adults in Wyoming County and New York State	90
Table 40b. High blood pressure among adults in Wyoming County and New York State - Age-adjusted	91
Table 41a. Blood pressure medication among adults with high blood pressure in Wyoming County and New York State	92
Table 41b. Blood pressure medication among adults with high blood pressure in Wyoming County and New York State - Age-adjusted	93
Table 42a. Ever had blood cholesterol checked among adults in Wyoming County and New York State	94
Table 42b. Ever had blood cholesterol checked among adults in Wyoming County and New York State - Age-adjusted	95
Table 43a. Blood cholesterol checked within the past 5 years among adults in Wyoming County and New York State	96
Table 43b. Blood cholesterol checked within the past 5 years among adults in Wyoming County and New York State - Age-adjusted	97
Table 44a. Cardiovascular disease among adults in Wyoming County and New York State	98
Table 44b. Cardiovascular disease among adults in Wyoming County and New York State - Age-adjusted	99
Table 45a. Coronary heart disease among adults in Wyoming County and New York State	100

Contents

Table 45b. Coronary heart disease among adults in Wyoming County and New York State - Age-adjusted	101
Table 46a. Ever diagnosed with diabetes among adults in Wyoming County and New York State	102
Table 46b. Ever diagnosed with diabetes among adults in Wyoming County and New York State - Age-adjusted	103
Table 47a. Ever diagnosed with asthma (lifetime) among adults in Wyoming County and New York State	104
Table 47b. Ever diagnosed with asthma (lifetime) among adults in Wyoming County and New York State - Age-adjusted	105
Table 48a. Current asthma among adults in Wyoming County and New York State	106
Table 48b. Current asthma among adults in Wyoming County and New York State - Age-adjusted	107
Table 49a. Chronic joint symptoms among adults in Wyoming County and New York State	108
Table 49b. Chronic joint symptoms among adults in Wyoming County and New York State - Age-adjusted	109
Table 50a. Arthritis among adults in Wyoming County and New York State	110
Table 50b. Arthritis among adults in Wyoming County and New York State - Age-adjusted	111
Table 51a. Activity limitations because of physical, mental or emotional problems among adults in Wyoming County and New York State	112
Table 51b. Activity limitations because of physical, mental or emotional problems among adults in Wyoming County and New York State - Age-adjusted	113
Table 52a. Have health problem that requires the use of special equipment among adults in Wyoming County and New York State	114
Table 52b. Have health problem that requires the use of special equipment among adults in Wyoming County and New York State - Age-adjusted	115
Table 53a. Disability among adults in Wyoming County and New York State	116
Table 53b. Disability among adults in Wyoming County and New York State - Age-adjusted	117

Contents

Health Behaviors

Table 54a. Current smoking among adults in Wyoming County and New York State	118
Table 54b. Current smoking among adults in Wyoming County and New York State - Age-adjusted	119
Table 55a. Everyday smoking among adults in Wyoming County and New York State	120
Table 55b. Everyday smoking among adults in Wyoming County and New York State - Age-adjusted	121
Table 56a. Adults living in homes in which smoking is prohibited in Wyoming County and New York State	122
Table 56b. Adults living in homes in which smoking is prohibited in Wyoming County and New York State - Age-adjusted	123
Table 57a. Binge drinking within the past month among adults in Wyoming County and New York State	124
Table 57b. Binge drinking within the past month among adults in Wyoming County and New York State - Age-adjusted	125
Table 58a. Heavy drinking within the past month among adults in Wyoming County and New York State	126
Table 58b. Heavy drinking within the past month among adults in Wyoming County and New York State - Age-adjusted	127
Table 59a. Never or rarely been asked about sexual history during routine check-up among adults in Wyoming County and New York State	128
Table 59b. Never or rarely been asked about sexual history during routine check-up among adults in Wyoming County and New York State - Age-adjusted	129
Table 60a. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults in Wyoming County and New York State	130
Table 60b. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults in Wyoming County and New York State - Age-adjusted	131

Contents

Table 61a. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults in Wyoming County and New York State	132
Table 61b. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults in Wyoming County and New York State - Age-adjusted	133

Overweight / Obesity

Table 62a. Overweight among adults in Wyoming County and New York State	134
Table 62b. Overweight among adults in Wyoming County and New York State - Age-adjusted	135
Table 63a. Obesity among adults in Wyoming County and New York State	136
Table 63b. Obesity among adults in Wyoming County and New York State - Age-adjusted	137
Table 64a. Overweight or obesity among adults in Wyoming County and New York State	138
Table 64b. Overweight or obesity among adults in Wyoming County and New York State - Age-adjusted	139
Table 65a. Received advice about weight by a health professional among adults in Wyoming County and New York State	140
Table 65b. Received advice about weight by a health professional among adults in Wyoming County and New York State - Age-adjusted	141
Table 66a. Received advice to lose weight by a health professional among adults provided weight advice in Wyoming County and New York State	142
Table 66b. Received advice to lose weight by a health professional among adults provided weight advice in Wyoming County and New York State - Age-adjusted	143

Physical Activity / Nutrition

Table 67a. No leisure-time physical activity among adults in Wyoming County and New York State	144
Table 67b. No leisure-time physical activity among adults in Wyoming County and New York State - Age-adjusted	145

Contents

Table 68a. Consumption of 5 or more servings of fruits and vegetables per day among adults in Wyoming County and New York State	146
Table 68b. Consumption of 5 or more servings of fruits and vegetables per day among adults in Wyoming County and New York State - Age-adjusted	147

Injury

Table 69a. Had a fall within the past 3 months among adults in Wyoming County and New York State	148
Table 69b. Had a fall within the past 3 months among adults in Wyoming County and New York State - Age-adjusted	149
Table 70a. Had a fall that caused injury within the past 3 months among adults in Wyoming County and New York State	150
Table 70b. Had a fall that caused injury within the past 3 months among adults in Wyoming County and New York State - Age-adjusted	151

Introduction

The New York State Expanded Behavioral Risk Factor Surveillance System (Expanded BRFSS) is a random digit dialed telephone survey of adults (18 years of age and older) that is representative of the non-institutionalized civilian population with landline telephones living in New York State. The goal of the Expanded BRFSS is to collect county-specific data on preventive health practices, risk behaviors, injuries and preventable chronic and infectious diseases. Topics assessed by the Expanded BRFSS include tobacco use, physical inactivity, diet, use of cancer screening services and other factors linked to the leading causes of morbidity and mortality. For each topic, health indicators are based on answers to one or more questions in the survey. The survey results are weighted to more accurately represent the adult population living in each county. Weighting procedures adjust for the differential selection probabilities by race, age and gender. More detailed information on weighting procedures is available in the technical brief.

Survey data collection occurred in 58 geographic areas of New York State, including each of the 57 counties outside of New York City and New York City (5 counties) as a single area. A standard questionnaire was used in all areas. Data collection for the 2009 Expanded BRFSS took place during the period of July 2008 through June 2009. The goal was to complete about 650 interviews in each area. The final number of interviews varies because of the sampling design of the survey and differential response rates. More detailed information on the sampling design is available in the technical brief.

The New York State Department of Health contracted with RTI International, which is responsible for the data collection, sampling, weighting and creation of the statistical reports. RTI International subcontracted with Clearwater Research Inc. for data collection and additional expertise in sample design and data quality control.

This report presents both crude estimates and estimates that have been age-adjusted to the Year 2000 US Standard Population when they represent the entire adult population of an area. Estimates for specific age groups are not age-adjusted. Age adjustment is a statistical process applied to rates that allows communities with different age structures to be compared. Counties can compare their data with results from New York State as a whole and with other counties. Confidence intervals at the 95% level associated with each health indicator give an estimated range of values which is likely to include the actual prevalence of the indicator. Confidence intervals are similar to margins of error. When the confidence intervals of two estimates of the same indicator from different areas do not overlap, they may be said to be statistically significantly different, i.e., these differences are unlikely related to chance and are considered true differences. If there is any value that is included in both intervals, the two estimates are not statistically significantly different.

The survey results for each of the 69 health indicators included in the survey are typically provided by gender, age, race and ethnicity, education and income in the Expanded BRFSS county reports. The estimates for some of these groupings may be suppressed in geographic areas with low proportions of adult respondents in these groupings.

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 1. Wyoming County and New York State adult US Census Bureau population estimates,¹ 2007

	Wyoming County		New York State	
	Adult Population ¹	Percent	Adult Population ¹	Percent
Total	33,496	100	14,884,315	100
Gender				
Male	18,572	55.4	7,103,791	47.7
Female	14,924	44.6	7,780,524	52.3
Age (years)				
18-34	10,157	30.3	4,497,479	30.2
35-44	6,402	19.1	2,844,115	19.1
45-54	6,728	20.1	2,854,316	19.2
55-64	4,766	14.2	2,142,000	14.4
≥ 65	5,443	16.2	2,546,405	17.1
Race/Ethnicity				
White Non-Hispanic	29,712	88.7	9,343,884	62.8
Black Non-Hispanic	2,290	6.8	2,156,941	14.5
Hispanic	1,243	3.7	2,258,717	15.2
Other Non-Hispanic	251	0.7	1,124,773	7.6

¹ Source: National Center for Health Statistics. Postcensal estimates of the resident population of the United States for July 1, 2000-July 1, 2007, by year, county, age, bridged race, Hispanic origin, and sex (Vintage 2007). Prepared under a collaborative arrangement with the U.S. Census Bureau; released August 7, 2008. Available from: http://www.cdc.gov/nchs/nvss/bridged_race.htm as of November 25, 2009.

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 2a. Fair or poor general health among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	671	4,500	13.5	(10.0 - 17.1)	38,124	2,531,400	17.1	(15.1 - 19.0)
Gender								
Male	248	2,100	11.3	(6.0 - 16.7)	14,599	1,069,700	15.1	(12.4 - 17.8)
Female	423	2,400	16.3	(11.9 - 20.6)	23,525	1,461,700	18.8	(16.1 - 21.6)
Age (years)								
18-34	*	*	*	*	3,940	535,800	12.0	(7.2 - 16.8)
35-44	92	600	9.9	(3.1 - 16.8)	4,985	289,200	10.0	(6.8 - 13.3)
45-54	154	1,100	16.2	(10.3 - 22.1)	7,819	507,800	17.9	(14.0 - 21.8)
55-64	169	800	16.4	(9.9 - 22.9)	8,757	424,700	19.9	(16.0 - 23.9)
≥ 65	196	1,200	21.5	(15.1 - 28.0)	12,623	773,900	30.5	(26.8 - 34.2)
Race/Ethnicity								
White Non-Hispanic	653	3,800	12.5	(9.7 - 15.4)	35,811	1,226,600	13.2	(11.7 - 14.6)
Black Non-Hispanic	*	*	*	*	739	457,300	21.3	(14.8 - 27.9)
Hispanic	*	*	*	*	861	684,100	30.4	(22.2 - 38.6)
Other Non-Hispanic	*	*	*	*	713	163,400	14.5	(5.1 - 24.0)
Education								
≤ High School	353	3,000	17.5	(11.3 - 23.7)	15,682	1,466,500	26.3	(22.6 - 30.0)
Some College	193	1,300	13.1	(7.5 - 18.8)	9,904	508,500	14.3	(10.9 - 17.8)
≥ College Degree	125	300	4.0	(1.3 - 6.6)	12,457	493,500	8.8	(6.5 - 11.1)
Income								
≤ \$24,999	179	1,600	21.8	(12.8 - 30.8)	9,633	1,269,200	36.3	(30.1 - 42.4)
\$25,000-\$49,999	186	1,000	12.1	(7.2 - 17.0)	9,649	447,600	15.4	(11.9 - 18.9)
\$50,000-\$74,999	117	400	5.0	(1.2 - 8.9)	5,769	237,800	12.1	(8.0 - 16.2)
≥ \$75,000	*	*	*	*	8,562	218,300	4.9	(3.6 - 6.1)

¹ Defined as reporting fair or poor general health

² Sample size from survey

³ Estimated number of adults who have fair or poor health (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 2b. Fair or poor general health among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	671	13.5	(9.9 - 17.1)	38,124	16.7	(14.8 - 18.6)
Gender						
Male	248	11.6	(6.5 - 16.8)	14,599	15.3	(12.6 - 18.0)
Female	423	15.8	(11.0 - 20.5)	23,525	18.0	(15.2 - 20.7)
Race/Ethnicity						
White Non-Hispanic	653	12.5	(9.5 - 15.4)	35,811	12.9	(11.4 - 14.4)
Black Non-Hispanic	*	*	*	739	20.8	(14.2 - 27.3)
Hispanic	*	*	*	861	29.9	(21.8 - 37.9)
Other Non-Hispanic	*	*	*	713	13.8	(4.8 - 22.8)
Education						
≤ High School	353	17.7	(11.5 - 23.8)	15,682	25.8	(22.1 - 29.5)
Some College	193	13.4	(7.5 - 19.3)	9,904	14.1	(10.7 - 17.6)
≥ College Degree	125	3.4	(1.1 - 5.7)	12,457	8.5	(6.3 - 10.8)
Income						
≤ \$24,999	179	22.2	(13.0 - 31.4)	9,633	35.9	(29.7 - 42.0)
\$25,000-\$49,999	186	12.2	(7.1 - 17.2)	9,649	15.0	(11.6 - 18.5)
\$50,000-\$74,999	117	4.9	(1.0 - 8.8)	5,769	11.8	(7.7 - 16.0)
≥ \$75,000	*	*	*	8,562	4.7	(3.5 - 6.0)

¹ Defined as reporting fair or poor general health

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wyoming County**

Table 3a. Poor physical health 14 or more days within the past month among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	663	3,700	11.4	(7.9 - 14.8)	37,327	1,472,900	10.1	(8.7 - 11.6)
Gender								
Male	245	1,800	10.0	(4.6 - 15.5)	14,325	604,200	8.7	(7.0 - 10.5)
Female	418	1,900	13.0	(9.0 - 17.0)	23,002	868,600	11.4	(9.2 - 13.6)
Age (years)								
18-34	*	*	*	*	3,891	268,100	6.1	(2.7 - 9.5)
35-44	91	1,000	15.2	(7.0 - 23.4)	4,948	232,100	8.1	(5.2 - 11.1)
45-54	153	800	12.3	(7.1 - 17.5)	7,764	350,900	12.5	(9.4 - 15.6)
55-64	168	400	8.7	(4.3 - 13.1)	8,621	271,200	13.0	(10.3 - 15.8)
≥ 65	193	800	15.7	(9.7 - 21.7)	12,103	350,600	14.5	(12.1 - 16.9)
Race/Ethnicity								
White Non-Hispanic	648	3,200	10.4	(7.7 - 13.1)	35,073	929,700	10.2	(8.8 - 11.6)
Black Non-Hispanic	*	*	*	*	724	208,200	9.9	(4.8 - 14.9)
Hispanic	*	*	*	*	835	250,300	11.5	(6.5 - 16.4)
Other Non-Hispanic	*	*	*	*	695	84,600	7.6	(1.3 - 13.9)
Education								
≤ High School	347	2,700	17.0	(10.8 - 23.3)	15,178	692,800	12.9	(10.6 - 15.3)
Some College	191	800	8.2	(3.9 - 12.5)	9,751	362,500	10.3	(7.4 - 13.2)
≥ College Degree	*	*	*	*	12,327	359,300	6.4	(4.8 - 8.1)
Income								
≤ \$24,999	176	1,500	23.0	(14.8 - 31.2)	9,294	625,600	18.4	(13.6 - 23.3)
\$25,000-\$49,999	185	600	7.7	(3.9 - 11.4)	9,495	279,100	9.8	(7.3 - 12.3)
\$50,000-\$74,999	*	*	*	*	5,733	186,400	9.5	(5.8 - 13.2)
≥ \$75,000	*	*	*	*	8,510	219,100	4.9	(3.6 - 6.2)

¹ Defined as 14 or more days within the past month when physical health was not good

² Sample size from survey

³ Estimated number of adults with poor physical health 14 or more days within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 3b. Poor physical health 14 or more days within the past month among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	663	11.8	(8.2 - 15.3)	37,327	9.9	(8.4 - 11.4)
Gender						
Male	245	10.6	(5.3 - 15.9)	14,325	8.7	(6.9 - 10.5)
Female	418	13.2	(8.7 - 17.7)	23,002	11.0	(8.7 - 13.2)
Race/Ethnicity						
White Non-Hispanic	648	10.9	(7.9 - 14.0)	35,073	9.9	(8.5 - 11.3)
Black Non-Hispanic	*	*	*	724	9.7	(4.6 - 14.9)
Hispanic	*	*	*	835	11.3	(6.3 - 16.2)
Other Non-Hispanic	*	*	*	695	7.6	(1.2 - 14.0)
Education						
≤ High School	347	18.3	(11.8 - 24.7)	15,178	12.7	(10.2 - 15.1)
Some College	191	8.0	(3.6 - 12.4)	9,751	10.1	(7.2 - 12.9)
≥ College Degree	*	*	*	12,327	6.3	(4.7 - 8.0)
Income						
≤ \$24,999	176	24.0	(15.2 - 32.9)	9,294	18.4	(13.5 - 23.4)
\$25,000-\$49,999	185	8.1	(4.0 - 12.1)	9,495	9.4	(7.0 - 11.9)
\$50,000-\$74,999	*	*	*	5,733	9.0	(5.3 - 12.6)
≥ \$75,000	*	*	*	8,510	4.9	(3.5 - 6.2)

¹ Defined as 14 or more days within the past month when physical health was not good

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 4a. Poor mental health 14 or more days within the past month among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	664	2,300	7.1	(4.8 - 9.4)	37,528	1,473,200	10.1	(8.6 - 11.5)
Gender								
Male	*	*	*	*	14,407	555,200	7.9	(6.3 - 9.6)
Female	418	1,900	13.1	(8.7 - 17.6)	23,121	918,000	12.0	(9.7 - 14.3)
Age (years)								
18-34	*	*	*	*	3,889	496,700	11.2	(7.4 - 15.1)
35-44	*	*	*	*	4,947	254,100	8.9	(6.7 - 11.1)
45-54	152	500	8.1	(3.6 - 12.7)	7,749	347,200	12.3	(9.0 - 15.6)
55-64	166	400	8.3	(3.6 - 12.9)	8,623	211,300	10.1	(7.9 - 12.3)
≥ 65	196	200	4.1	(1.5 - 6.7)	12,320	163,900	6.7	(4.7 - 8.7)
Race/Ethnicity								
White Non-Hispanic	647	2,100	6.8	(4.5 - 9.2)	35,266	899,200	9.8	(8.4 - 11.2)
Black Non-Hispanic	*	*	*	*	722	227,800	10.9	(5.4 - 16.4)
Hispanic	*	*	*	*	845	263,700	11.8	(6.8 - 16.8)
Other Non-Hispanic	*	*	*	*	695	82,500	7.4	(1.4 - 13.5)
Education								
≤ High School	347	1,500	9.4	(5.4 - 13.4)	15,335	665,000	12.2	(10.0 - 14.5)
Some College	192	500	4.8	(2.3 - 7.3)	9,771	404,500	11.5	(8.3 - 14.8)
≥ College Degree	*	*	*	*	12,342	346,600	6.2	(4.5 - 7.9)
Income								
≤ \$24,999	177	1,100	14.2	(7.0 - 21.4)	9,398	611,500	17.9	(13.0 - 22.8)
\$25,000-\$49,999	184	400	4.5	(1.4 - 7.7)	9,547	260,300	9.0	(6.7 - 11.4)
\$50,000-\$74,999	*	*	*	*	5,726	187,300	9.6	(6.0 - 13.1)
≥ \$75,000	*	*	*	*	8,520	276,500	6.2	(4.6 - 7.7)

¹ Defined as 14 or more days within the past month when mental health was not good

² Sample size from survey

³ Estimated number of adults with poor mental health 14 or more days within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 4b. Poor mental health 14 or more days within the past month among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	664	7.3	(4.7 - 9.8)	37,528	10.2	(8.7 - 11.7)
Gender						
Male	*	*	*	14,407	7.9	(6.3 - 9.6)
Female	418	13.6	(8.6 - 18.6)	23,121	12.2	(9.8 - 14.6)
Race/Ethnicity						
White Non-Hispanic	647	7.0	(4.5 - 9.6)	35,266	10.0	(8.5 - 11.4)
Black Non-Hispanic	*	*	*	722	11.1	(5.5 - 16.7)
Hispanic	*	*	*	845	11.5	(6.5 - 16.4)
Other Non-Hispanic	*	*	*	695	7.6	(1.5 - 13.7)
Education						
≤ High School	347	10.2	(5.7 - 14.6)	15,335	12.3	(10.0 - 14.6)
Some College	192	4.2	(2.0 - 6.4)	9,771	11.8	(8.5 - 15.1)
≥ College Degree	*	*	*	12,342	6.3	(4.6 - 8.0)
Income						
≤ \$24,999	177	14.5	(7.0 - 21.9)	9,398	18.2	(13.2 - 23.1)
\$25,000-\$49,999	184	4.7	(1.3 - 8.1)	9,547	9.4	(6.9 - 11.9)
\$50,000-\$74,999	*	*	*	5,726	9.5	(5.9 - 13.1)
≥ \$75,000	*	*	*	8,520	6.2	(4.6 - 7.7)

¹ Defined as 14 or more days within the past month when mental health was not good

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 5a. Poor physical or mental health 14 or more days within the past month among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	658	5,100	15.8	(11.9 - 19.6)	37,005	2,385,900	16.5	(14.8 - 18.2)
Gender								
Male	245	1,900	11.0	(5.5 - 16.5)	14,231	937,700	13.6	(11.4 - 15.8)
Female	413	3,100	21.5	(16.3 - 26.6)	22,774	1,448,200	19.1	(16.5 - 21.7)
Age (years)								
18-34	*	*	*	*	3,859	602,600	13.8	(9.7 - 17.9)
35-44	91	1,100	17.2	(8.8 - 25.7)	4,924	388,300	13.7	(10.4 - 17.0)
45-54	151	1,100	16.6	(10.5 - 22.6)	7,722	551,000	19.6	(15.8 - 23.4)
55-64	165	700	14.7	(8.9 - 20.5)	8,545	395,500	19.2	(16.1 - 22.3)
≥ 65	193	1,000	18.4	(12.1 - 24.7)	11,955	448,500	18.8	(15.9 - 21.7)
Race/Ethnicity								
White Non-Hispanic	642	4,400	14.5	(11.3 - 17.7)	34,778	1,485,700	16.3	(14.8 - 17.8)
Black Non-Hispanic	*	*	*	*	715	343,700	16.5	(10.3 - 22.7)
Hispanic	*	*	*	*	832	412,900	19.0	(12.6 - 25.3)
Other Non-Hispanic	*	*	*	*	680	143,500	13.0	(5.0 - 21.0)
Education								
≤ High School	344	3,500	22.3	(15.7 - 28.9)	15,018	1,077,200	20.3	(17.3 - 23.2)
Some College	189	1,100	11.3	(6.5 - 16.1)	9,672	615,500	17.6	(14.1 - 21.1)
≥ College Degree	125	500	6.9	(1.5 - 12.4)	12,244	634,200	11.4	(9.2 - 13.6)
Income								
≤ \$24,999	174	2,000	31.8	(22.7 - 41.0)	9,199	924,000	27.3	(21.9 - 32.8)
\$25,000-\$49,999	182	900	11.5	(6.7 - 16.3)	9,425	456,000	16.1	(13.0 - 19.2)
\$50,000-\$74,999	116	400	5.7	(2.1 - 9.4)	5,696	312,000	16.0	(11.4 - 20.5)
≥ \$75,000	*	*	*	*	8,475	447,600	10.0	(8.1 - 12.0)

¹ Defined as 14 or more days within the past month when physical or mental health prevented usual activities, such as self-care, work or recreation

² Sample size from survey

³ Estimated number of adults with poor physical or mental health 14 or more days within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 5b. Poor physical or mental health 14 or more days within the past month among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	658	16.2	(12.3 - 20.2)	37,005	16.4	(14.7 - 18.2)
Gender						
Male	245	11.6	(6.2 - 16.9)	14,231	13.6	(11.4 - 15.8)
Female	413	21.8	(16.1 - 27.5)	22,774	19.0	(16.3 - 21.6)
Race/Ethnicity						
White Non-Hispanic	642	15.1	(11.6 - 18.6)	34,778	16.3	(14.7 - 17.8)
Black Non-Hispanic	*	*	*	715	16.7	(10.3 - 23.0)
Hispanic	*	*	*	832	18.5	(12.2 - 24.8)
Other Non-Hispanic	*	*	*	680	13.1	(5.0 - 21.3)
Education						
≤ High School	344	23.8	(17.0 - 30.6)	15,018	20.1	(17.1 - 23.1)
Some College	189	10.7	(5.9 - 15.5)	9,672	17.6	(14.1 - 21.1)
≥ College Degree	125	6.9	(1.0 - 12.9)	12,244	11.4	(9.2 - 13.7)
Income						
≤ \$24,999	174	32.9	(23.1 - 42.8)	9,199	27.6	(22.1 - 33.1)
\$25,000-\$49,999	182	12.0	(6.8 - 17.1)	9,425	16.1	(12.9 - 19.3)
\$50,000-\$74,999	116	5.6	(1.8 - 9.4)	5,696	15.6	(11.0 - 20.1)
≥ \$75,000	*	*	*	8,475	10.0	(8.0 - 12.0)

¹ Defined as 14 or more days within the past month when physical or mental health prevented usual activities, such as self-care, work or recreation

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 6a. Health insurance among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	671	28,800	86.9	(80.4 - 93.3)	38,119	12,854,200	86.6	(84.2 - 89.0)
Gender								
Male	*	*	*	*	14,597	5,881,100	83.0	(78.9 - 87.1)
Female	424	13,800	92.9	(89.6 - 96.1)	23,522	6,973,100	89.9	(87.2 - 92.6)
Age (years)								
18-34	*	*	*	*	3,910	3,302,100	74.3	(67.8 - 80.7)
35-44	92	5,400	84.9	(75.7 - 94.2)	4,986	2,560,300	88.9	(84.7 - 93.1)
45-54	155	5,900	87.7	(81.8 - 93.7)	7,831	2,530,100	88.9	(84.6 - 93.2)
55-64	168	4,300	91.0	(86.1 - 96.0)	8,764	1,986,700	92.8	(90.2 - 95.5)
≥ 65	197	5,400	99.1	(97.8 - 100)	12,628	2,475,000	97.8	(96.7 - 98.8)
Race/Ethnicity								
White Non-Hispanic	653	27,600	90.7	(87.4 - 94.0)	35,811	8,591,200	92.2	(90.6 - 93.8)
Black Non-Hispanic	*	*	*	*	736	1,781,600	82.7	(75.3 - 90.1)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,673	4,448,900	79.7	(74.9 - 84.5)
Some College	194	9,200	92.9	(88.5 - 97.3)	9,906	3,004,600	85.0	(80.1 - 89.8)
≥ College Degree	124	6,400	96.6	(90.9 - 100)	12,459	5,361,700	95.3	(93.3 - 97.2)
Income								
≤ \$24,999	*	*	*	*	9,640	2,437,200	69.7	(62.5 - 76.8)
\$25,000-\$49,999	187	7,700	93.0	(87.4 - 98.7)	9,662	2,384,000	82.2	(76.1 - 88.3)
\$50,000-\$74,999	117	7,000	97.7	(95.1 - 100)	5,772	1,855,700	94.0	(91.2 - 96.7)
≥ \$75,000	111	5,900	91.2	(82.5 - 99.9)	8,567	4,388,400	97.5	(96.6 - 98.5)

¹ Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

² Sample size from survey

³ Estimated number of adults who have health insurance (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 6b. Health insurance among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	671	87.3	(81.3 - 93.4)	38,119	86.7	(84.4 - 89.1)
Gender						
Male	*	*	*	14,597	83.6	(79.7 - 87.5)
Female	424	92.6	(89.0 - 96.3)	23,522	89.6	(86.8 - 92.3)
Race/Ethnicity						
White Non-Hispanic	653	90.8	(87.5 - 94.1)	35,811	92.1	(90.5 - 93.7)
Black Non-Hispanic	*	*	*	736	82.5	(75.0 - 90.0)
Hispanic	*	*	*	860	68.7	(59.0 - 78.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	15,673	80.1	(75.4 - 84.7)
Some College	194	92.8	(88.3 - 97.4)	9,906	84.6	(79.7 - 89.4)
≥ College Degree	124	97.0	(91.9 - 100)	12,459	95.4	(93.6 - 97.2)
Income						
≤ \$24,999	*	*	*	9,640	70.2	(63.2 - 77.1)
\$25,000-\$49,999	187	93.2	(87.7 - 98.7)	9,662	81.9	(76.0 - 87.9)
\$50,000-\$74,999	117	97.6	(94.7 - 100)	5,772	94.0	(91.3 - 96.7)
≥ \$75,000	111	91.9	(83.7 - 100)	8,567	97.5	(96.6 - 98.5)

¹ Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 7a. Health insurance among adults aged 18-64¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	470	23,300	84.4	(76.8 - 92.0)	25,165	10,251,300	84.2	(81.3 - 87.1)
Gender								
Male	*	*	*	*	10,078	4,839,900	80.3	(75.5 - 85.0)
Female	288	10,800	91.2	(87.2 - 95.3)	15,087	5,411,400	88.0	(84.7 - 91.3)
Age (years)								
18-34	*	*	*	*	3,910	3,302,100	74.3	(67.8 - 80.7)
35-44	92	5,400	84.9	(75.7 - 94.2)	4,981	2,556,600	88.9	(84.7 - 93.1)
45-54	155	5,900	87.7	(81.8 - 93.7)	7,791	2,472,900	88.7	(84.3 - 93.0)
55-64	164	4,200	90.8	(85.7 - 95.9)	8,483	1,919,700	92.8	(90.1 - 95.5)
≥ 65	-	-	-	-	-	-	-	-
Race/Ethnicity								
White Non-Hispanic	455	22,200	88.9	(85.0 - 92.9)	23,354	6,582,200	90.3	(88.3 - 92.4)
Black Non-Hispanic	*	*	*	*	567	1,464,100	80.2	(71.6 - 88.7)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	9,037	3,191,500	74.6	(68.6 - 80.6)
Some College	143	7,700	91.6	(86.4 - 96.8)	7,031	2,540,400	82.9	(77.4 - 88.4)
≥ College Degree	97	5,600	96.1	(89.8 - 100)	9,068	4,502,400	94.5	(92.2 - 96.8)
Income								
≤ \$24,999	*	*	*	*	4,918	1,707,200	62.6	(54.1 - 71.2)
\$25,000-\$49,999	125	5,900	91.1	(83.9 - 98.3)	6,180	1,783,000	77.7	(70.3 - 85.1)
\$50,000-\$74,999	100	6,300	97.5	(94.6 - 100)	4,554	1,553,400	92.9	(89.7 - 96.1)
≥ \$75,000	100	5,500	91.1	(82.0 - 100)	7,324	3,987,600	97.3	(96.3 - 98.3)

¹ Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

² Sample size from survey

³ Estimated number of adults aged 18-64 who have health insurance (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 7b. Health insurance among adults aged 18-64¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	470	84.9	(77.7 - 92.0)	25,165	84.3	(81.5 - 87.2)
Gender						
Male	*	*	*	10,078	80.5	(75.9 - 85.1)
Female	288	91.3	(86.9 - 95.6)	15,087	87.9	(84.6 - 91.2)
Race/Ethnicity						
White Non-Hispanic	455	88.9	(84.9 - 93.0)	23,354	90.3	(88.3 - 92.3)
Black Non-Hispanic	*	*	*	567	80.0	(71.3 - 88.6)
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	9,037	75.1	(69.3 - 80.9)
Some College	143	91.5	(86.0 - 97.0)	7,031	82.6	(77.1 - 88.1)
≥ College Degree	97	96.6	(90.9 - 100)	9,068	94.6	(92.5 - 96.8)
Income						
≤ \$24,999	*	*	*	4,918	63.2	(54.9 - 71.6)
\$25,000-\$49,999	125	91.1	(84.1 - 98.2)	6,180	77.5	(70.2 - 84.7)
\$50,000-\$74,999	100	97.3	(94.1 - 100)	4,554	93.0	(89.8 - 96.2)
≥ \$75,000	100	91.8	(83.2 - 100)	7,324	97.3	(96.3 - 98.3)

¹ Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 8a. No health insurance among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	671	4,400	13.1	(6.7 - 19.6)	38,119	1,989,300	13.4	(11.0 - 15.8)
Gender								
Male	*	*	*	*	14,597	1,207,400	17.0	(12.9 - 21.1)
Female	424	1,100	7.1	(3.9 - 10.4)	23,522	781,900	10.1	(7.4 - 12.8)
Age (years)								
18-34	*	*	*	*	3,910	1,144,100	25.7	(19.3 - 32.2)
35-44	92	1,000	15.1	(5.8 - 24.3)	4,986	319,200	11.1	(6.9 - 15.3)
45-54	155	800	12.3	(6.3 - 18.2)	7,831	316,500	11.1	(6.8 - 15.4)
55-64	168	400	9.0	(4.0 - 13.9)	8,764	153,300	7.2	(4.5 - 9.8)
≥ 65	*	*	*	*	12,628	56,200	2.2	(1.2 - 3.3)
Race/Ethnicity								
White Non-Hispanic	653	2,800	9.3	(6.0 - 12.6)	35,811	727,100	7.8	(6.2 - 9.4)
Black Non-Hispanic	*	*	*	*	736	372,800	17.3	(9.9 - 24.7)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,673	1,134,400	20.3	(15.5 - 25.1)
Some College	194	700	7.1	(2.7 - 11.5)	9,906	531,400	15.0	(10.2 - 19.9)
≥ College Degree	*	*	*	*	12,459	266,300	4.7	(2.8 - 6.7)
Income								
≤ \$24,999	*	*	*	*	9,640	1,061,200	30.3	(23.2 - 37.5)
\$25,000-\$49,999	*	*	*	*	9,662	516,700	17.8	(11.7 - 23.9)
\$50,000-\$74,999	*	*	*	*	5,772	119,100	6.0	(3.3 - 8.8)
≥ \$75,000	*	*	*	*	8,567	111,100	2.5	(1.5 - 3.4)

¹ Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

² Sample size from survey

³ Estimated number of adults who have no health insurance (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 8b. No health insurance among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	671	12.7	(6.6 - 18.7)	38,119	13.3	(10.9 - 15.6)
Gender						
Male	*	*	*	14,597	16.4	(12.5 - 20.3)
Female	424	7.4	(3.7 - 11.0)	23,522	10.4	(7.7 - 13.2)
Race/Ethnicity						
White Non-Hispanic	653	9.2	(5.9 - 12.5)	35,811	7.9	(6.3 - 9.5)
Black Non-Hispanic	*	*	*	736	17.5	(10.0 - 25.0)
Hispanic	*	*	*	860	31.3	(21.6 - 41.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	15,673	19.9	(15.3 - 24.6)
Some College	194	7.2	(2.6 - 11.7)	9,906	15.4	(10.6 - 20.3)
≥ College Degree	*	*	*	12,459	4.6	(2.8 - 6.4)
Income						
≤ \$24,999	*	*	*	9,640	29.8	(22.9 - 36.8)
\$25,000-\$49,999	*	*	*	9,662	18.1	(12.1 - 24.0)
\$50,000-\$74,999	*	*	*	5,772	6.0	(3.3 - 8.7)
≥ \$75,000	*	*	*	8,567	2.5	(1.5 - 3.4)

¹ Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 9a. No health insurance among adults aged 18-64¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	470	4,300	15.6	(8.0 - 23.2)	25,165	1,928,400	15.8	(12.9 - 18.7)
Gender								
Male	*	*	*	*	10,078	1,190,200	19.7	(15.0 - 24.5)
Female	288	1,000	8.8	(4.7 - 12.8)	15,087	738,200	12.0	(8.7 - 15.3)
Age (years)								
18-34	*	*	*	*	3,910	1,144,100	25.7	(19.3 - 32.2)
35-44	92	1,000	15.1	(5.8 - 24.3)	4,981	319,200	11.1	(6.9 - 15.3)
45-54	155	800	12.3	(6.3 - 18.2)	7,791	316,100	11.3	(7.0 - 15.7)
55-64	164	400	9.2	(4.1 - 14.3)	8,483	148,900	7.2	(4.5 - 9.9)
≥ 65	-	-	-	-	-	-	-	-
Race/Ethnicity								
White Non-Hispanic	455	2,800	11.1	(7.1 - 15.0)	23,354	703,100	9.7	(7.6 - 11.7)
Black Non-Hispanic	*	*	*	*	567	361,800	19.8	(11.3 - 28.4)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	9,037	1,087,700	25.4	(19.4 - 31.4)
Some College	143	700	8.4	(3.2 - 13.6)	7,031	522,800	17.1	(11.6 - 22.6)
≥ College Degree	*	*	*	*	9,068	261,200	5.5	(3.2 - 7.8)
Income								
≤ \$24,999	*	*	*	*	4,918	1,018,000	37.4	(28.8 - 45.9)
\$25,000-\$49,999	*	*	*	*	6,180	511,100	22.3	(14.9 - 29.7)
\$50,000-\$74,999	*	*	*	*	4,554	118,800	7.1	(3.9 - 10.3)
≥ \$75,000	*	*	*	*	7,324	110,700	2.7	(1.7 - 3.7)

¹ Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

² Sample size from survey

³ Estimated number of adults aged 18-64 who have no health insurance (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 9b. No health insurance among adults aged 18-64¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	470	15.1	(8.0 - 22.3)	25,165	15.7	(12.8 - 18.5)
Gender						
Male	*	*	*	10,078	19.5	(14.9 - 24.1)
Female	288	8.7	(4.4 - 13.1)	15,087	12.1	(8.8 - 15.4)
Race/Ethnicity						
White Non-Hispanic	455	11.1	(7.0 - 15.1)	23,354	9.7	(7.7 - 11.7)
Black Non-Hispanic	*	*	*	567	20.0	(11.4 - 28.7)
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	9,037	24.9	(19.1 - 30.7)
Some College	143	8.5	(3.0 - 14.0)	7,031	17.4	(11.9 - 22.9)
≥ College Degree	*	*	*	9,068	5.4	(3.2 - 7.5)
Income						
≤ \$24,999	*	*	*	4,918	36.8	(28.4 - 45.1)
\$25,000-\$49,999	*	*	*	6,180	22.5	(15.3 - 29.8)
\$50,000-\$74,999	*	*	*	4,554	7.0	(3.8 - 10.2)
≥ \$75,000	*	*	*	7,324	2.7	(1.7 - 3.7)

¹ Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 10a. Adults with regular health care provider¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	672	29,000	86.8	(80.2 - 93.4)	38,135	12,324,300	83.0	(80.4 - 85.6)
Gender								
Male	*	*	*	*	14,593	5,494,300	77.6	(73.2 - 82.0)
Female	425	13,900	93.4	(89.4 - 97.4)	23,542	6,830,000	87.8	(85.1 - 90.5)
Age (years)								
18-34	*	*	*	*	3,929	2,931,600	65.8	(59.2 - 72.5)
35-44	92	5,300	82.6	(73.2 - 92.1)	4,988	2,471,000	86.1	(81.9 - 90.3)
45-54	155	6,200	92.8	(88.6 - 97.0)	7,823	2,529,300	88.6	(84.6 - 92.7)
55-64	167	4,500	97.1	(94.3 - 100)	8,770	2,002,000	93.6	(91.7 - 95.4)
≥ 65	198	5,100	93.4	(89.5 - 97.4)	12,625	2,390,400	94.1	(92.2 - 96.0)
Race/Ethnicity								
White Non-Hispanic	654	27,800	90.9	(87.2 - 94.6)	35,821	8,227,700	88.2	(86.4 - 89.9)
Black Non-Hispanic	*	*	*	*	740	1,753,200	81.3	(72.6 - 89.9)
Hispanic	*	*	*	*	860	1,453,100	64.7	(54.7 - 74.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,692	4,249,200	75.9	(70.9 - 80.8)
Some College	193	8,800	89.3	(81.3 - 97.3)	9,904	2,935,000	82.8	(77.2 - 88.4)
≥ College Degree	125	6,400	96.5	(92.7 - 100)	12,456	5,046,100	89.8	(87.1 - 92.6)
Income								
≤ \$24,999	*	*	*	*	9,644	2,554,200	72.7	(65.9 - 79.6)
\$25,000-\$49,999	186	7,200	87.7	(80.1 - 95.3)	9,655	2,243,100	77.3	(70.0 - 84.6)
\$50,000-\$74,999	117	6,700	94.5	(86.7 - 100)	5,771	1,767,500	89.5	(85.2 - 93.9)
≥ \$75,000	111	6,300	98.4	(96.6 - 100)	8,560	4,122,800	91.9	(90.2 - 93.7)

¹ Defined as having one or more persons respondent thinks of as a personal doctor or health care provider

² Sample size from survey

³ Estimated number of adults who have at least one person they think of as their personal doctor or health care provider (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 10b. Adults with regular health care provider¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	672	86.7	(80.4 - 92.9)	38,135	83.0	(80.4 - 85.5)
Gender						
Male	*	*	*	14,593	78.1	(73.9 - 82.3)
Female	425	92.6	(87.9 - 97.4)	23,542	87.4	(84.6 - 90.1)
Race/Ethnicity						
White Non-Hispanic	654	90.4	(86.5 - 94.2)	35,821	88.0	(86.2 - 89.7)
Black Non-Hispanic	*	*	*	740	81.4	(73.0 - 89.8)
Hispanic	*	*	*	860	65.2	(55.5 - 75.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	15,692	75.8	(71.0 - 80.6)
Some College	193	88.4	(80.1 - 96.8)	9,904	82.9	(77.5 - 88.3)
≥ College Degree	125	96.2	(92.2 - 100)	12,456	89.8	(87.1 - 92.4)
Income						
≤ \$24,999	*	*	*	9,644	72.9	(66.1 - 79.6)
\$25,000-\$49,999	186	86.7	(78.4 - 95.0)	9,655	77.4	(70.5 - 84.3)
\$50,000-\$74,999	117	94.4	(86.8 - 100)	5,771	89.6	(85.3 - 94.0)
≥ \$75,000	111	98.3	(96.3 - 100)	8,560	91.6	(89.8 - 93.4)

¹ Defined as having one or more persons respondent thinks of as a personal doctor or health care provider

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 11a. Cost prevented visit to doctor within the past year among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	674	3,100	9.2	(5.6 - 12.8)	38,143	2,034,800	13.7	(11.6 - 15.8)
Gender								
Male	248	1,800	9.5	(4.1 - 14.9)	14,607	903,300	12.7	(9.5 - 16.0)
Female	426	1,300	8.8	(4.4 - 13.2)	23,536	1,131,500	14.6	(11.8 - 17.4)
Age (years)								
18-34	*	*	*	*	3,939	947,300	21.2	(15.6 - 26.9)
35-44	92	900	13.6	(5.1 - 22.0)	4,987	347,900	12.1	(8.0 - 16.2)
45-54	155	700	10.8	(5.3 - 16.4)	7,825	382,200	13.5	(9.9 - 17.1)
55-64	*	*	*	*	8,766	236,800	11.1	(8.0 - 14.2)
≥ 65	*	*	*	*	12,626	120,600	4.8	(2.5 - 7.0)
Race/Ethnicity								
White Non-Hispanic	656	2,700	8.7	(5.7 - 11.7)	35,833	984,600	10.6	(8.9 - 12.2)
Black Non-Hispanic	*	*	*	*	738	280,000	13.1	(7.2 - 19.0)
Hispanic	*	*	*	*	859	597,900	26.5	(17.3 - 35.7)
Other Non-Hispanic	*	*	*	*	713	172,300	15.3	(5.7 - 24.9)
Education								
≤ High School	355	1,600	9.6	(4.1 - 15.1)	15,696	996,600	17.8	(13.8 - 21.9)
Some College	194	1,100	10.6	(4.2 - 17.0)	9,911	512,400	14.5	(10.4 - 18.6)
≥ College Degree	*	*	*	*	12,453	466,900	8.3	(6.0 - 10.6)
Income								
≤ \$24,999	180	800	11.3	(4.7 - 17.9)	9,639	854,500	24.3	(18.4 - 30.3)
\$25,000-\$49,999	187	1,000	12.5	(5.0 - 20.0)	9,656	532,900	18.4	(12.5 - 24.3)
\$50,000-\$74,999	*	*	*	*	5,776	202,700	10.3	(6.0 - 14.5)
≥ \$75,000	*	*	*	*	8,561	192,000	4.3	(2.9 - 5.6)

¹ Defined as needing to see a doctor within the past year but not being able to because of cost

² Sample size from survey

³ Estimated number of adults who did not see a doctor because of cost within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 11b. Cost prevented visit to doctor within the past year among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	674	9.5	(5.9 - 13.2)	38,143	13.8	(11.7 - 15.9)
Gender						
Male	248	9.5	(4.3 - 14.7)	14,607	12.5	(9.4 - 15.6)
Female	426	9.5	(4.4 - 14.6)	23,536	15.0	(12.1 - 17.9)
Race/Ethnicity						
White Non-Hispanic	656	9.2	(5.9 - 12.5)	35,833	10.8	(9.2 - 12.5)
Black Non-Hispanic	*	*	*	738	13.2	(7.2 - 19.3)
Hispanic	*	*	*	859	26.1	(17.2 - 35.1)
Other Non-Hispanic	*	*	*	713	14.8	(5.5 - 24.2)
Education						
≤ High School	355	9.8	(4.4 - 15.1)	15,696	17.8	(13.8 - 21.7)
Some College	194	11.3	(4.2 - 18.4)	9,911	15.3	(11.0 - 19.5)
≥ College Degree	*	*	*	12,453	8.2	(6.0 - 10.5)
Income						
≤ \$24,999	180	11.3	(4.5 - 18.0)	9,639	24.5	(18.6 - 30.4)
\$25,000-\$49,999	187	13.3	(5.1 - 21.5)	9,656	18.7	(12.9 - 24.6)
\$50,000-\$74,999	*	*	*	5,776	10.4	(6.1 - 14.7)
≥ \$75,000	*	*	*	8,561	4.4	(3.1 - 5.8)

¹ Defined as needing to see a doctor within the past year but not being able to because of cost

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 12a. Visited a doctor for routine checkup within the past year among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	669	23,500	70.5	(65.0 - 76.1)	38,026	10,809,300	72.9	(70.6 - 75.3)
Gender								
Male	246	12,600	67.9	(58.9 - 76.9)	14,592	4,738,400	66.9	(63.0 - 70.9)
Female	423	10,900	73.9	(68.1 - 79.7)	23,434	6,070,900	78.5	(75.9 - 81.0)
Age (years)								
18-34	*	*	*	*	3,921	2,747,400	61.7	(55.6 - 67.9)
35-44	*	*	*	*	4,973	1,883,100	65.5	(60.2 - 70.7)
45-54	154	4,500	67.9	(59.0 - 76.8)	7,808	2,174,100	76.6	(73.2 - 80.0)
55-64	168	3,500	74.6	(66.9 - 82.3)	8,751	1,728,600	81.4	(78.7 - 84.2)
≥ 65	197	4,900	90.0	(85.7 - 94.3)	12,573	2,276,000	90.0	(87.5 - 92.5)
Race/Ethnicity								
White Non-Hispanic	651	21,300	69.9	(64.5 - 75.3)	35,721	6,746,200	72.4	(70.6 - 74.2)
Black Non-Hispanic	*	*	*	*	736	1,688,900	78.8	(70.4 - 87.3)
Hispanic	*	*	*	*	857	1,550,900	69.3	(60.0 - 78.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	352	12,200	72.3	(64.4 - 80.2)	15,640	4,150,600	74.7	(70.7 - 78.7)
Some College	*	*	*	*	9,880	2,524,000	71.3	(66.1 - 76.5)
≥ College Degree	*	*	*	*	12,428	4,042,900	71.9	(68.5 - 75.3)
Income								
≤ \$24,999	178	5,900	78.6	(69.2 - 88.0)	9,616	2,469,200	70.4	(64.2 - 76.6)
\$25,000-\$49,999	187	5,600	67.8	(58.9 - 76.8)	9,631	2,068,400	71.3	(65.3 - 77.3)
\$50,000-\$74,999	*	*	*	*	5,765	1,466,000	74.8	(69.6 - 80.0)
≥ \$75,000	*	*	*	*	8,547	3,176,500	70.7	(67.3 - 74.2)

¹ Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past year

² Sample size from survey

³ Estimated number of adults who have visited a doctor for a routine checkup within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 12b. Visited a doctor for routine checkup within the past year among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	669	70.1	(64.5 - 75.7)	38,026	72.7	(70.3 - 75.0)
Gender						
Male	246	68.7	(60.2 - 77.3)	14,592	67.1	(63.3 - 70.9)
Female	423	71.9	(65.3 - 78.4)	23,434	77.7	(75.1 - 80.3)
Race/Ethnicity						
White Non-Hispanic	651	69.4	(63.9 - 74.8)	35,721	71.9	(70.1 - 73.7)
Black Non-Hispanic	*	*	*	736	79.3	(71.3 - 87.4)
Hispanic	*	*	*	857	69.2	(60.1 - 78.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	352	71.8	(64.0 - 79.7)	15,640	74.4	(70.4 - 78.4)
Some College	*	*	*	9,880	70.9	(65.9 - 76.0)
≥ College Degree	*	*	*	12,428	71.7	(68.3 - 75.0)
Income						
≤ \$24,999	178	78.6	(69.2 - 88.0)	9,616	70.2	(64.1 - 76.2)
\$25,000-\$49,999	187	67.4	(58.3 - 76.5)	9,631	71.1	(65.3 - 76.9)
\$50,000-\$74,999	*	*	*	5,765	74.5	(69.2 - 79.8)
≥ \$75,000	*	*	*	8,547	70.3	(66.9 - 73.8)

¹ Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past year

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 13a. Visited a doctor for routine checkup within the past 2 years among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	669	27,300	82.1	(77.5 - 86.7)	38,026	12,726,600	85.9	(83.9 - 87.9)
Gender								
Male	246	14,700	79.5	(72.2 - 86.8)	14,592	5,791,900	81.8	(78.2 - 85.4)
Female	423	12,600	85.3	(80.5 - 90.2)	23,434	6,934,700	89.6	(87.9 - 91.3)
Age (years)								
18-34	*	*	*	*	3,921	3,535,400	79.4	(73.9 - 85.0)
35-44	*	*	*	*	4,973	2,355,600	81.9	(78.0 - 85.7)
45-54	154	5,400	81.3	(73.4 - 89.1)	7,808	2,507,900	88.4	(85.6 - 91.2)
55-64	168	3,800	80.8	(73.7 - 88.0)	8,751	1,924,800	90.7	(88.6 - 92.7)
≥ 65	197	5,200	95.5	(92.5 - 98.5)	12,573	2,402,900	95.0	(93.5 - 96.5)
Race/Ethnicity								
White Non-Hispanic	651	25,000	82.0	(77.7 - 86.3)	35,721	8,070,000	86.6	(85.4 - 87.9)
Black Non-Hispanic	*	*	*	*	736	1,914,500	89.4	(82.3 - 96.5)
Hispanic	*	*	*	*	857	1,832,800	81.9	(73.7 - 90.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	352	13,300	79.0	(71.6 - 86.4)	15,640	4,713,700	84.8	(81.2 - 88.4)
Some College	193	8,700	87.4	(81.1 - 93.6)	9,880	2,990,500	84.5	(79.6 - 89.4)
≥ College Degree	124	5,400	82.1	(72.6 - 91.6)	12,428	4,928,300	87.6	(85.3 - 89.9)
Income								
≤ \$24,999	178	6,400	85.2	(77.5 - 93.0)	9,616	2,911,700	83.0	(77.4 - 88.6)
\$25,000-\$49,999	187	7,000	85.4	(79.3 - 91.4)	9,631	2,375,500	81.9	(76.1 - 87.7)
\$50,000-\$74,999	*	*	*	*	5,765	1,733,500	88.4	(84.9 - 91.9)
≥ \$75,000	109	5,300	83.4	(73.5 - 93.2)	8,547	3,907,000	87.0	(84.8 - 89.2)

¹ Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past 2 years

² Sample size from survey

³ Estimated number of adults who have visited a doctor for a routine checkup within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 13b. Visited a doctor for routine checkup within the past 2 years among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	669	81.9	(77.3 - 86.5)	38,026	85.7	(83.7 - 87.6)
Gender						
Male	246	80.1	(73.1 - 87.1)	14,592	81.9	(78.4 - 85.3)
Female	423	84.1	(78.5 - 89.7)	23,434	89.1	(87.4 - 90.9)
Race/Ethnicity						
White Non-Hispanic	651	81.7	(77.3 - 86.1)	35,721	86.2	(85.0 - 87.5)
Black Non-Hispanic	*	*	*	736	89.6	(83.0 - 96.3)
Hispanic	*	*	*	857	81.8	(73.8 - 89.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	352	78.6	(71.2 - 85.9)	15,640	84.6	(81.1 - 88.1)
Some College	193	87.1	(80.8 - 93.5)	9,880	84.4	(79.8 - 89.0)
≥ College Degree	124	82.4	(72.7 - 92.1)	12,428	87.3	(85.0 - 89.6)
Income						
≤ \$24,999	178	85.0	(77.2 - 92.9)	9,616	83.2	(77.8 - 88.5)
\$25,000-\$49,999	187	85.3	(79.2 - 91.5)	9,631	81.9	(76.3 - 87.4)
\$50,000-\$74,999	*	*	*	5,765	88.1	(84.4 - 91.8)
≥ \$75,000	*	*	*	8,547	86.4	(84.2 - 88.7)

¹ Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past 2 years

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wyoming County**

Table 14a. Dentist visit within the past year among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	669	23,600	70.9	(64.7 - 77.2)	38,026	10,540,400	71.1	(68.6 - 73.6)
Gender								
Male	*	*	*	*	14,573	4,845,800	68.4	(64.5 - 72.2)
Female	424	10,700	71.9	(66.1 - 77.7)	23,453	5,694,600	73.6	(70.4 - 76.8)
Age (years)								
18-34	*	*	*	*	3,924	2,789,700	62.6	(56.3 - 69.0)
35-44	*	*	*	*	4,979	2,181,000	75.8	(70.9 - 80.7)
45-54	155	5,100	76.4	(69.2 - 83.6)	7,814	2,149,300	75.8	(71.9 - 79.7)
55-64	169	3,200	67.7	(59.8 - 75.7)	8,742	1,690,500	79.2	(75.9 - 82.6)
≥ 65	195	3,700	68.8	(61.5 - 76.0)	12,567	1,729,800	68.5	(64.7 - 72.3)
Race/Ethnicity								
White Non-Hispanic	651	22,300	73.3	(68.6 - 77.9)	35,721	6,959,900	74.7	(72.7 - 76.6)
Black Non-Hispanic	*	*	*	*	735	1,405,800	66.2	(57.9 - 74.5)
Hispanic	*	*	*	*	862	1,425,600	63.1	(53.7 - 72.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,613	3,620,700	65.1	(60.7 - 69.5)
Some College	193	7,800	79.1	(71.5 - 86.6)	9,880	2,504,700	70.8	(66.1 - 75.5)
≥ College Degree	*	*	*	*	12,453	4,402,400	78.2	(74.6 - 81.8)
Income								
≤ \$24,999	*	*	*	*	9,594	1,972,200	56.2	(49.8 - 62.6)
\$25,000-\$49,999	186	5,900	72.2	(64.0 - 80.4)	9,649	1,896,000	65.8	(59.5 - 72.1)
\$50,000-\$74,999	117	5,900	83.6	(73.8 - 93.4)	5,763	1,482,400	75.2	(69.4 - 81.0)
≥ \$75,000	111	5,700	89.0	(81.5 - 96.5)	8,564	3,707,900	82.4	(79.5 - 85.3)

¹ Defined as having visited a dentist or a dental clinic for any reason within the past year

² Sample size from survey

³ Estimated number of adults who have visited a dentist or a dental clinic for any reason within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 14b. Dentist visit within the past year among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	669	71.3	(65.2 - 77.3)	38,026	71.1	(68.6 - 73.5)
Gender						
Male	245	70.4	(60.7 - 80.1)	14,573	68.3	(64.6 - 72.1)
Female	424	72.3	(65.9 - 78.7)	23,453	73.6	(70.4 - 76.8)
Race/Ethnicity						
White Non-Hispanic	651	73.4	(68.6 - 78.1)	35,721	74.4	(72.4 - 76.3)
Black Non-Hispanic	*	*	*	735	66.6	(58.3 - 74.9)
Hispanic	*	*	*	862	63.6	(54.4 - 72.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	351	65.2	(55.4 - 74.9)	15,613	65.3	(60.9 - 69.6)
Some College	193	78.7	(70.9 - 86.4)	9,880	70.4	(65.7 - 75.1)
≥ College Degree	*	*	*	12,453	78.1	(74.6 - 81.7)
Income						
≤ \$24,999	*	*	*	9,594	56.4	(50.1 - 62.8)
\$25,000-\$49,999	186	72.8	(64.6 - 81.0)	9,649	65.8	(59.6 - 72.0)
\$50,000-\$74,999	117	84.6	(75.2 - 94.0)	5,763	75.0	(69.1 - 80.8)
≥ \$75,000	111	88.9	(81.4 - 96.4)	8,564	82.1	(79.1 - 85.0)

¹ Defined as having visited a dentist or a dental clinic for any reason within the past year

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 15a. Had teeth cleaned within the past year among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	587	21,800	72.9	(67.8 - 78.1)	34,070	9,985,900	71.7	(69.1 - 74.3)
Gender								
Male	218	11,700	70.7	(62.7 - 78.7)	13,173	4,616,400	69.0	(65.0 - 73.1)
Female	369	10,100	75.7	(69.6 - 81.8)	20,897	5,369,500	74.2	(70.8 - 77.5)
Age (years)								
18-34	*	*	*	*	3,860	2,746,400	63.0	(56.5 - 69.5)
35-44	*	*	*	*	4,875	2,112,100	75.5	(70.7 - 80.3)
45-54	150	4,900	75.2	(67.6 - 82.8)	7,438	2,022,600	73.3	(69.1 - 77.4)
55-64	144	2,900	70.5	(62.0 - 79.0)	7,906	1,560,900	77.7	(73.8 - 81.6)
≥ 65	148	3,100	76.1	(68.3 - 83.9)	9,991	1,543,900	77.1	(73.7 - 80.5)
Race/Ethnicity								
White Non-Hispanic	571	20,500	72.8	(67.5 - 78.0)	31,982	6,601,500	75.4	(73.4 - 77.4)
Black Non-Hispanic	*	*	*	*	680	1,227,200	61.6	(52.1 - 71.0)
Hispanic	*	*	*	*	789	1,421,300	66.6	(57.1 - 76.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	282	9,700	70.0	(62.3 - 77.8)	12,740	3,388,600	67.7	(63.1 - 72.3)
Some College	183	7,300	76.3	(67.9 - 84.6)	9,145	2,286,800	68.5	(62.9 - 74.1)
≥ College Degree	*	*	*	*	12,128	4,298,500	78.3	(74.8 - 81.7)
Income								
≤ \$24,999	*	*	*	*	7,525	1,861,800	60.3	(53.5 - 67.1)
\$25,000-\$49,999	168	5,100	67.0	(57.5 - 76.5)	8,709	1,681,700	62.6	(55.4 - 69.8)
\$50,000-\$74,999	*	*	*	*	5,534	1,447,800	76.3	(70.7 - 81.8)
≥ \$75,000	109	5,700	89.9	(82.4 - 97.3)	8,448	3,617,500	80.8	(77.8 - 83.8)

¹ Defined as having had teeth cleaned by a dentist or dental hygienist within the past year

² Sample size from survey

³ Estimated number of adults who have had their teeth cleaned by a dentist or dental hygienist within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 15b. Had teeth cleaned within the past year among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	587	72.8	(67.6 - 78.1)	34,070	71.7	(69.1 - 74.2)
Gender						
Male	218	70.5	(62.7 - 78.3)	13,173	69.2	(65.3 - 73.1)
Female	369	75.6	(68.8 - 82.4)	20,897	73.9	(70.5 - 77.3)
Race/Ethnicity						
White Non-Hispanic	571	72.6	(67.3 - 77.9)	31,982	75.1	(73.0 - 77.1)
Black Non-Hispanic	*	*	*	680	62.0	(52.7 - 71.4)
Hispanic	*	*	*	789	66.9	(57.5 - 76.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	282	69.9	(62.1 - 77.7)	12,740	67.7	(63.2 - 72.3)
Some College	183	75.7	(67.1 - 84.3)	9,145	68.1	(62.6 - 73.6)
≥ College Degree	*	*	*	12,128	78.3	(74.9 - 81.7)
Income						
≤ \$24,999	*	*	*	7,525	60.2	(53.4 - 67.0)
\$25,000-\$49,999	168	67.1	(57.4 - 76.7)	8,709	62.7	(55.7 - 69.7)
\$50,000-\$74,999	*	*	*	5,534	76.4	(70.9 - 81.9)
≥ \$75,000	109	89.7	(82.2 - 97.2)	8,448	80.4	(77.4 - 83.4)

¹ Defined as having had teeth cleaned by a dentist or dental hygienist within the past year

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 16a. Permanent teeth extracted due to decay or gum disease among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	665	16,700	50.3	(44.0 - 56.6)	37,652	7,426,500	50.4	(47.8 - 52.9)
Gender								
Male	248	8,500	45.7	(36.1 - 55.4)	14,457	3,367,000	47.8	(43.9 - 51.6)
Female	417	8,200	56.1	(49.6 - 62.7)	23,195	4,059,500	52.7	(49.5 - 56.0)
Age (years)								
18-34	*	*	*	*	3,921	1,121,600	25.4	(19.7 - 31.0)
35-44	*	*	*	*	4,964	1,250,000	43.6	(37.9 - 49.2)
45-54	155	3,900	57.5	(48.4 - 66.7)	7,780	1,595,300	56.1	(51.4 - 60.7)
55-64	166	3,100	66.8	(58.5 - 75.1)	8,648	1,382,800	65.5	(61.6 - 69.4)
≥ 65	193	4,600	86.5	(80.8 - 92.2)	12,339	2,076,900	83.0	(80.6 - 85.4)
Race/Ethnicity								
White Non-Hispanic	647	15,300	50.2	(44.5 - 55.9)	35,364	4,425,700	47.9	(46.0 - 49.9)
Black Non-Hispanic	*	*	*	*	727	1,244,700	58.2	(48.9 - 67.4)
Hispanic	*	*	*	*	853	1,220,500	54.2	(44.9 - 63.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,427	3,436,000	61.9	(57.2 - 66.6)
Some College	189	4,100	41.9	(32.1 - 51.6)	9,798	1,726,200	49.3	(44.3 - 54.2)
≥ College Degree	125	1,700	25.8	(17.0 - 34.6)	12,351	2,242,600	40.1	(36.5 - 43.6)
Income								
≤ \$24,999	*	*	*	*	9,483	2,171,900	62.3	(55.8 - 68.8)
\$25,000-\$49,999	182	5,000	61.8	(52.3 - 71.4)	9,552	1,577,400	54.7	(48.3 - 61.0)
\$50,000-\$74,999	*	*	*	*	5,744	1,000,700	51.5	(45.6 - 57.4)
≥ \$75,000	*	*	*	*	8,516	1,738,200	38.8	(35.3 - 42.2)

¹ Defined as having had at least one permanent tooth extracted because of tooth decay or gum disease

² Sample size from survey

³ Estimated number of adults who have had at least one permanent tooth extracted because of tooth decay or gum disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 16b. Permanent teeth extracted due to decay or gum disease among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	665	49.7	(43.5 - 55.9)	37,652	49.4	(46.9 - 51.9)
Gender						
Male	248	47.3	(38.0 - 56.7)	14,457	47.9	(44.1 - 51.7)
Female	417	52.8	(45.6 - 59.9)	23,195	50.7	(47.4 - 54.0)
Race/Ethnicity						
White Non-Hispanic	647	49.5	(43.7 - 55.2)	35,364	46.5	(44.5 - 48.5)
Black Non-Hispanic	*	*	*	727	57.7	(48.6 - 66.9)
Hispanic	*	*	*	853	53.8	(44.6 - 62.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	15,427	61.3	(56.7 - 66.0)
Some College	189	42.1	(32.2 - 52.0)	9,798	47.9	(43.0 - 52.9)
≥ College Degree	125	24.9	(15.9 - 33.9)	12,351	39.0	(35.5 - 42.5)
Income						
≤ \$24,999	*	*	*	9,483	61.6	(55.2 - 68.1)
\$25,000-\$49,999	182	61.5	(51.7 - 71.3)	9,552	54.2	(48.0 - 60.3)
\$50,000-\$74,999	*	*	*	5,744	49.9	(44.0 - 55.9)
≥ \$75,000	*	*	*	8,516	37.7	(34.2 - 41.1)

¹ Defined as having had at least one permanent tooth extracted because of tooth decay or gum disease

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wyoming County**

Table 17a. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	193	1,300	23.9	(17.1 - 30.6)	12,339	491,200	19.6	(16.0 - 23.3)
Gender								
Male	*	*	*	*	4,352	167,200	16.4	(11.8 - 20.9)
Female	129	800	27.0	(18.3 - 35.7)	7,987	324,000	21.8	(16.6 - 27.0)
Age (years)								
≥ 65	193	1,300	23.9	(17.1 - 30.6)	12,339	491,200	19.6	(16.0 - 23.3)
Race/Ethnicity								
White Non-Hispanic	190	1,300	24.6	(17.7 - 31.4)	11,885	322,600	16.8	(14.3 - 19.2)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	119	1,100	36.0	(26.2 - 45.7)	6,358	314,100	25.3	(19.9 - 30.7)
Some College	*	*	*	*	2,736	90,600	20.2	(12.1 - 28.2)
≥ College Degree	*	*	*	*	3,214	80,500	10.1	(3.6 - 16.5)
Income								
≤ \$24,999	*	*	*	*	4,558	199,600	26.9	(19.3 - 34.4)
\$25,000-\$49,999	*	*	*	*	3,365	123,900	21.4	(13.1 - 29.6)
\$50,000-\$74,999	*	*	*	*	1,179	21,400	7.3	(3.8 - 10.8)
≥ \$75,000	*	*	*	*	1,187	9,600	2.5	(1.0 - 4.0)

¹ Defined as having all permanent teeth extracted because of tooth decay or gum disease among adults aged 65 and older

² Sample size from survey

³ Estimated number of adults aged 65 and older who have had all permanent teeth extracted because of tooth decay or gum disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 17b. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	193	23.1	(16.1 - 30.1)	12,339	19.4	(15.7 - 23.2)
Gender						
Male	*	*	*	4,352	16.5	(11.5 - 21.4)
Female	129	27.0	(18.3 - 35.7)	7,987	22.1	(16.6 - 27.7)
Race/Ethnicity						
White Non-Hispanic	190	23.9	(16.8 - 31.1)	11,885	16.6	(13.9 - 19.3)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	6,358	24.9	(19.3 - 30.5)
Some College	*	*	*	2,736	21.0	(12.1 - 29.9)
≥ College Degree	*	*	*	3,214	10.0	(3.6 - 16.3)
Income						
≤ \$24,999	*	*	*	4,558	27.3	(19.1 - 35.5)
\$25,000-\$49,999	*	*	*	3,365	21.0	(12.9 - 29.1)
\$50,000-\$74,999	*	*	*	1,179	7.6	(3.6 - 11.5)
≥ \$75,000	*	*	*	1,187	2.2	(0.9 - 3.5)

¹ Defined as having all permanent teeth extracted because of tooth decay or gum disease among adults aged 65 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 18a. Flu shot within the past 12 months¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	672	15,900	48.0	(41.7 - 54.2)	38,079	6,255,900	42.3	(39.9 - 44.8)
Gender								
Male	*	*	*	*	14,579	2,798,900	39.6	(35.8 - 43.5)
Female	425	7,500	50.4	(44.2 - 56.7)	23,500	3,456,900	44.8	(41.6 - 47.9)
Age (years)								
18-49	*	*	*	*	12,505	2,685,300	31.1	(27.3 - 34.9)
50-64	256	4,700	54.2	(46.8 - 61.5)	12,972	1,662,900	46.0	(42.6 - 49.5)
≥ 65	198	4,300	79.4	(73.0 - 85.9)	12,602	1,907,700	75.2	(72.0 - 78.4)
Race/Ethnicity								
White Non-Hispanic	654	13,800	45.5	(40.0 - 51.1)	35,770	4,029,000	43.3	(41.4 - 45.3)
Black Non-Hispanic	*	*	*	*	738	907,000	42.1	(33.1 - 51.1)
Hispanic	*	*	*	*	860	882,900	39.6	(30.4 - 48.9)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	353	8,400	50.5	(40.9 - 60.0)	15,663	2,457,800	44.3	(39.8 - 48.8)
Some College	*	*	*	*	9,898	1,377,700	38.9	(34.0 - 43.9)
≥ College Degree	*	*	*	*	12,436	2,404,400	43.0	(39.5 - 46.4)
Income								
≤ \$24,999	*	*	*	*	9,635	1,656,800	47.6	(41.2 - 54.0)
\$25,000-\$49,999	187	3,300	39.9	(31.3 - 48.5)	9,647	1,200,600	41.4	(35.6 - 47.3)
\$50,000-\$74,999	*	*	*	*	5,763	792,300	40.2	(34.3 - 46.2)
≥ \$75,000	*	*	*	*	8,548	1,776,100	39.6	(36.3 - 42.8)

¹ Defined as having had a flu shot within the past 12 months

² Sample size from survey

³ Estimated number of adults who have had a flu shot within the past 12 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 18b. Flu shot within the past 12 months¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	672	46.9	(40.7 - 53.1)	38,079	41.7	(39.2 - 44.1)
Gender						
Male	247	46.8	(37.1 - 56.6)	14,579	40.0	(36.2 - 43.8)
Female	425	47.0	(40.2 - 53.8)	23,500	43.1	(40.0 - 46.3)
Race/Ethnicity						
White Non-Hispanic	654	44.6	(39.1 - 50.2)	35,770	42.4	(40.4 - 44.3)
Black Non-Hispanic	*	*	*	738	42.3	(33.3 - 51.3)
Hispanic	*	*	*	860	39.0	(29.9 - 48.1)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	353	49.1	(39.7 - 58.6)	15,663	43.7	(39.1 - 48.2)
Some College	*	*	*	9,898	37.8	(33.0 - 42.7)
≥ College Degree	*	*	*	12,436	42.5	(39.1 - 46.0)
Income						
≤ \$24,999	*	*	*	9,635	47.3	(41.0 - 53.7)
\$25,000-\$49,999	187	39.4	(30.6 - 48.3)	9,647	40.3	(34.5 - 46.0)
\$50,000-\$74,999	*	*	*	5,763	39.4	(33.4 - 45.4)
≥ \$75,000	*	*	*	8,548	39.1	(35.8 - 42.4)

¹ Defined as having had a flu shot within the past 12 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 19a. Flu vaccine that was sprayed in nose within the past 12 months¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	*	*	*	*	38,106	248,300	1.7	(0.6 - 2.8)
Gender								
Male	*	*	*	*	14,586	148,000	2.1	(0.0 - 4.2)
Female	*	*	*	*	23,520	100,300	1.3	(0.5 - 2.1)
Age (years)								
18-49	*	*	*	*	12,535	184,900	2.1	(0.3 - 3.9)
50-64	*	*	*	*	12,968	36,000	1.0	(0.2 - 1.8)
≥ 65	*	*	*	*	12,603	27,400	1.1	(0.4 - 1.8)
Race/Ethnicity								
White Non-Hispanic	*	*	*	*	35,796	66,000	0.7	(0.4 - 1.0)
Black Non-Hispanic	*	*	*	*	740	38,500	1.8	(0.0 - 4.1)
Hispanic	*	*	*	*	860	133,700	5.9	(0.0 - 12.3)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,679	172,500	3.1	(0.3 - 5.9)
Some College	*	*	*	*	9,897	43,500	1.2	(0.4 - 2.0)
≥ College Degree	*	*	*	*	12,449	31,600	0.6	(0.3 - 0.8)
Income								
≤ \$24,999	*	*	*	*	9,636	138,400	3.9	(0.0 - 8.1)
\$25,000-\$49,999	*	*	*	*	9,648	41,500	1.4	(0.0 - 3.1)
\$50,000-\$74,999	*	*	*	*	5,766	13,300	0.7	(0.2 - 1.2)
≥ \$75,000	*	*	*	*	8,561	35,300	0.8	(0.3 - 1.3)

¹ Defined as having had flu vaccine sprayed in nose within the past 12 months

² Sample size from survey

³ Estimated number of adults who have had flu vaccine sprayed in nose within the past 12 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 19b. Flu vaccine that was sprayed in nose within the past 12 months¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	*	*	*	38,106	1.7	(0.6 - 2.7)
Gender						
Male	*	*	*	14,586	2.0	(0.1 - 3.9)
Female	*	*	*	23,520	1.4	(0.5 - 2.3)
Race/Ethnicity						
White Non-Hispanic	*	*	*	35,796	0.7	(0.4 - 1.0)
Black Non-Hispanic	*	*	*	740	1.9	(0.0 - 4.4)
Hispanic	*	*	*	860	5.9	(0.0 - 11.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	15,679	3.1	(0.4 - 5.8)
Some College	*	*	*	9,897	1.2	(0.4 - 2.0)
≥ College Degree	*	*	*	12,449	0.6	(0.3 - 0.8)
Income						
≤ \$24,999	*	*	*	9,636	3.8	(0.0 - 7.7)
\$25,000-\$49,999	*	*	*	9,648	1.5	(0.0 - 3.4)
\$50,000-\$74,999	*	*	*	5,766	0.7	(0.1 - 1.2)
≥ \$75,000	*	*	*	8,561	0.8	(0.3 - 1.3)

¹ Defined as having had flu vaccine sprayed in nose within the past 12 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 20a. Flu shot or flu vaccine sprayed in nose within the past 12 months¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	672	16,000	48.1	(41.8 - 54.4)	38,059	6,302,300	42.7	(40.2 - 45.1)
Gender								
Male	*	*	*	*	14,568	2,822,700	40.0	(36.1 - 43.9)
Female	425	7,500	50.5	(44.3 - 56.8)	23,491	3,479,500	45.1	(42.0 - 48.2)
Age (years)								
18-49	*	*	*	*	12,501	2,726,400	31.6	(27.8 - 35.4)
50-64	256	4,700	54.7	(47.4 - 62.1)	12,964	1,664,200	46.1	(42.7 - 49.5)
≥ 65	198	4,300	79.4	(73.0 - 85.9)	12,594	1,911,700	75.4	(72.2 - 78.6)
Race/Ethnicity								
White Non-Hispanic	654	13,900	45.7	(40.1 - 51.3)	35,750	4,048,700	43.6	(41.7 - 45.5)
Black Non-Hispanic	*	*	*	*	738	910,000	42.2	(33.2 - 51.2)
Hispanic	*	*	*	*	860	903,300	40.6	(31.3 - 49.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	353	8,400	50.8	(41.2 - 60.3)	15,651	2,480,100	44.7	(40.2 - 49.3)
Some College	*	*	*	*	9,895	1,387,300	39.2	(34.3 - 44.1)
≥ College Degree	*	*	*	*	12,431	2,418,300	43.2	(39.8 - 46.7)
Income								
≤ \$24,999	*	*	*	*	9,626	1,675,100	48.2	(41.8 - 54.6)
\$25,000-\$49,999	187	3,300	39.9	(31.3 - 48.5)	9,643	1,208,300	41.7	(35.8 - 47.6)
\$50,000-\$74,999	*	*	*	*	5,761	795,600	40.4	(34.5 - 46.3)
≥ \$75,000	*	*	*	*	8,548	1,791,200	39.9	(36.6 - 43.2)

¹ Defined as having had a flu shot or flu vaccine sprayed in nose within the past 12 months

² Sample size from survey

³ Estimated number of adults who have had a flu shot or flu vaccine sprayed in nose within the past 12 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 20b. Flu shot or flu vaccine sprayed in nose within the past 12 months¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	672	47.0	(40.9 - 53.2)	38,059	42.0	(39.6 - 44.5)
Gender						
Male	247	47.0	(37.2 - 56.7)	14,568	40.4	(36.6 - 44.2)
Female	425	47.1	(40.3 - 53.9)	23,491	43.5	(40.3 - 46.7)
Race/Ethnicity						
White Non-Hispanic	654	44.8	(39.2 - 50.4)	35,750	42.6	(40.7 - 44.6)
Black Non-Hispanic	*	*	*	738	42.4	(33.4 - 51.4)
Hispanic	*	*	*	860	40.0	(30.9 - 49.1)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	353	49.4	(40.0 - 58.8)	15,651	44.1	(39.6 - 48.6)
Some College	*	*	*	9,895	38.1	(33.2 - 43.0)
≥ College Degree	*	*	*	12,431	42.8	(39.4 - 46.3)
Income						
≤ \$24,999	*	*	*	9,626	48.0	(41.6 - 54.3)
\$25,000-\$49,999	187	39.4	(30.6 - 48.3)	9,643	40.5	(34.8 - 46.3)
\$50,000-\$74,999	*	*	*	5,761	39.6	(33.6 - 45.6)
≥ \$75,000	*	*	*	8,548	39.4	(36.1 - 42.7)

¹ Defined as having had a flu shot or flu vaccine sprayed in nose within the past 12 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wyoming County**

Table 21a. Flu shot among adults aged 65 and older¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	198	4,300	79.4	(73.0 - 85.9)	12,602	1,907,700	75.2	(72.0 - 78.4)
Gender								
Male	*	*	*	*	4,417	772,900	74.6	(69.3 - 79.8)
Female	134	2,500	83.6	(77.0 - 90.2)	8,185	1,134,800	75.6	(71.6 - 79.6)
Age (years)								
≥ 65	198	4,300	79.4	(73.0 - 85.9)	12,602	1,907,700	75.2	(72.0 - 78.4)
Race/Ethnicity								
White Non-Hispanic	195	4,200	79.4	(72.9 - 85.9)	12,137	1,487,100	76.0	(73.5 - 78.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	122	2,400	74.1	(65.5 - 82.8)	6,518	930,800	74.0	(68.9 - 79.2)
Some College	*	*	*	*	2,790	317,800	69.7	(62.7 - 76.7)
≥ College Degree	*	*	*	*	3,261	654,000	80.5	(76.4 - 84.6)
Income								
≤ \$24,999	76	1,400	81.0	(71.0 - 90.9)	4,665	545,300	72.4	(65.6 - 79.3)
\$25,000-\$49,999	*	*	*	*	3,422	430,600	72.8	(67.0 - 78.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	1,198	301,000	77.5	(70.9 - 84.2)

¹ Defined as having had a flu shot during the past 12 months among adults aged 65 and older

² Sample size from survey

³ Estimated number of adults aged 65 and older who have had a flu shot within the past 12 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 21b. Flu shot among adults aged 65 and older¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	198	78.4	(71.3 - 85.6)	12,602	75.0	(71.5 - 78.5)
Gender						
Male	*	*	*	4,417	74.4	(68.7 - 80.1)
Female	134	83.6	(77.0 - 90.2)	8,185	75.5	(71.3 - 79.8)
Race/Ethnicity						
White Non-Hispanic	195	78.4	(71.1 - 85.7)	12,137	75.9	(73.0 - 78.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	122	72.8	(63.2 - 82.3)	6,518	73.3	(67.5 - 79.1)
Some College	*	*	*	2,790	70.0	(62.7 - 77.2)
≥ College Degree	*	*	*	3,261	80.7	(76.2 - 85.1)
Income						
≤ \$24,999	*	*	*	4,665	72.6	(65.1 - 80.1)
\$25,000-\$49,999	*	*	*	3,422	71.4	(65.0 - 77.9)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	1,198	78.0	(70.8 - 85.1)

¹ Defined as having had a flu shot during the past 12 months among adults aged 65 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 22a. Pneumonia shot or pneumococcal vaccine¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	635	7,700	24.4	(20.1 - 28.7)	35,303	3,456,800	26.1	(23.9 - 28.3)
Gender								
Male	228	2,900	16.5	(10.9 - 22.1)	13,052	1,542,200	25.2	(21.5 - 28.8)
Female	407	4,800	34.1	(28.5 - 39.7)	22,251	1,914,600	26.9	(24.2 - 29.6)
Age (years)								
18-49	201	1,900	10.9	(5.8 - 15.9)	10,943	1,097,000	14.6	(11.3 - 17.9)
50-64	242	1,800	21.3	(16.0 - 26.7)	12,179	758,000	23.0	(19.8 - 26.2)
≥ 65	192	4,000	75.5	(68.5 - 82.4)	12,181	1,601,700	65.7	(61.7 - 69.7)
Race/Ethnicity								
White Non-Hispanic	617	7,200	25.1	(20.9 - 29.2)	33,225	2,358,700	28.7	(26.9 - 30.5)
Black Non-Hispanic	*	*	*	*	685	584,400	28.4	(19.5 - 37.3)
Hispanic	*	*	*	*	769	408,000	19.5	(12.6 - 26.4)
Other Non-Hispanic	*	*	*	*	624	105,600	12.2	(2.3 - 22.2)
Education								
≤ High School	337	4,200	26.3	(19.9 - 32.7)	14,700	1,535,300	30.4	(26.4 - 34.3)
Some College	178	2,300	24.5	(16.2 - 32.8)	9,151	836,200	27.2	(21.9 - 32.5)
≥ College Degree	120	1,200	19.6	(11.7 - 27.5)	11,375	1,070,700	21.4	(18.6 - 24.1)
Income								
≤ \$24,999	*	*	*	*	9,082	1,050,900	33.1	(27.6 - 38.6)
\$25,000-\$49,999	171	2,000	27.4	(19.8 - 34.9)	9,033	797,500	30.3	(24.1 - 36.6)
\$50,000-\$74,999	112	1,000	15.7	(7.6 - 23.8)	5,320	378,800	21.8	(17.4 - 26.1)
≥ \$75,000	102	1,000	16.3	(7.0 - 25.6)	7,773	648,700	16.3	(14.1 - 18.5)

¹ Defined as having ever had a pneumonia shot or pneumococcal vaccine

² Sample size from survey

³ Estimated number of adults who have ever had a pneumonia shot or pneumococcal vaccine (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 22b. Pneumonia shot or pneumococcal vaccine¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	635	24.5	(20.1 - 28.8)	35,303	25.4	(23.2 - 27.6)
Gender						
Male	228	18.9	(13.0 - 24.8)	13,052	25.8	(22.3 - 29.4)
Female	407	31.3	(25.4 - 37.2)	22,251	25.1	(22.4 - 27.8)
Race/Ethnicity						
White Non-Hispanic	617	25.0	(20.7 - 29.2)	33,225	27.7	(25.9 - 29.5)
Black Non-Hispanic	*	*	*	685	27.9	(19.1 - 36.7)
Hispanic	*	*	*	769	19.6	(12.8 - 26.5)
Other Non-Hispanic	*	*	*	624	11.3	(2.0 - 20.7)
Education						
≤ High School	337	26.7	(20.2 - 33.2)	14,700	29.5	(25.6 - 33.5)
Some College	178	24.0	(15.6 - 32.3)	9,151	26.2	(21.0 - 31.3)
≥ College Degree	120	19.9	(11.3 - 28.5)	11,375	21.0	(18.3 - 23.8)
Income						
≤ \$24,999	*	*	*	9,082	32.3	(26.9 - 37.8)
\$25,000-\$49,999	171	26.9	(19.2 - 34.5)	9,033	29.0	(23.0 - 35.0)
\$50,000-\$74,999	112	15.9	(7.6 - 24.2)	5,320	21.2	(16.8 - 25.7)
≥ \$75,000	102	16.5	(6.8 - 26.2)	7,773	16.2	(13.9 - 18.5)

¹ Defined as having ever had a pneumonia shot or pneumococcal vaccine

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wyoming County**

Table 23a. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	192	4,000	75.5	(68.5 - 82.4)	12,181	1,601,700	65.7	(61.7 - 69.7)
Gender								
Male	*	*	*	*	4,180	606,400	62.1	(55.5 - 68.7)
Female	130	2,400	81.9	(74.9 - 89.0)	8,001	995,300	68.0	(63.1 - 73.0)
Age (years)								
≥ 65	192	4,000	75.5	(68.5 - 82.4)	12,181	1,601,700	65.7	(61.7 - 69.7)
Race/Ethnicity								
White Non-Hispanic	189	3,800	74.7	(67.6 - 81.8)	11,735	1,281,300	68.2	(64.9 - 71.5)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	119	2,400	75.4	(66.7 - 84.0)	6,345	806,200	65.6	(59.5 - 71.7)
Some College	*	*	*	*	2,694	289,500	66.8	(60.1 - 73.4)
≥ College Degree	*	*	*	*	3,111	501,100	65.6	(58.5 - 72.7)
Income								
≤ \$24,999	75	1,400	84.8	(77.0 - 92.6)	4,533	497,100	67.7	(59.9 - 75.5)
\$25,000-\$49,999	*	*	*	*	3,324	344,300	61.5	(53.3 - 69.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	1,138	236,300	64.5	(55.2 - 73.8)

¹ Defined as having ever had a pneumonia shot or pneumococcal vaccine among adults aged 65 and older

² Sample size from survey

³ Estimated number of adults aged 65 and older who have ever had a pneumonia shot or pneumococcal vaccine (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 23b. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	192	73.9	(66.1 - 81.6)	12,181	64.7	(60.4 - 69.1)
Gender						
Male	*	*	*	4,180	61.5	(54.3 - 68.6)
Female	130	81.9	(74.9 - 89.0)	8,001	67.6	(62.4 - 72.9)
Race/Ethnicity						
White Non-Hispanic	189	73.0	(65.0 - 80.9)	11,735	67.1	(63.2 - 70.9)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	119	73.8	(64.2 - 83.4)	6,345	64.2	(57.4 - 71.0)
Some College	*	*	*	2,694	65.8	(58.4 - 73.3)
≥ College Degree	*	*	*	3,111	65.3	(57.7 - 72.9)
Income						
≤ \$24,999	75	85.8	(77.9 - 93.6)	4,533	66.0	(57.0 - 74.9)
\$25,000-\$49,999	*	*	*	3,324	60.3	(51.9 - 68.6)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	*	*

¹ Defined as having ever had a pneumonia shot or pneumococcal vaccine among adults aged 65 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 24a. Ever had a mammogram among women aged 40 and older¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	352	8,700	93.2	(90.1 - 96.3)	19,491	4,328,800	91.8	(89.7 - 94.0)
Gender								
Male	-	-	-	-	-	-	-	-
Female	352	8,700	93.2	(90.1 - 96.3)	19,491	4,328,800	91.8	(89.7 - 94.0)
Age (years)								
40-49	68	2,300	91.4	(84.1 - 98.7)	3,851	1,287,500	85.0	(79.7 - 90.3)
50-64	151	3,600	92.7	(87.5 - 98.0)	7,452	1,641,700	96.0	(94.1 - 97.9)
≥ 65	133	2,800	95.3	(91.7 - 98.9)	8,188	1,399,600	94.1	(90.5 - 97.6)
Race/Ethnicity								
White Non-Hispanic	348	8,600	93.8	(90.8 - 96.7)	18,548	2,976,100	93.9	(92.7 - 95.1)
Black Non-Hispanic	*	*	*	*	342	746,600	93.8	(87.3 - 100)
Hispanic	*	*	*	*	333	447,600	91.8	(84.6 - 99.0)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	186	4,600	94.0	(90.5 - 97.5)	8,266	1,698,700	91.0	(87.4 - 94.5)
Some College	108	2,800	93.4	(87.1 - 99.7)	5,322	1,055,500	93.6	(90.2 - 97.0)
≥ College Degree	58	1,300	90.2	(80.4 - 99.9)	5,872	1,560,700	91.6	(87.5 - 95.6)
Income								
≤ \$24,999	109	2,400	92.2	(86.1 - 98.2)	5,691	1,023,400	93.5	(89.7 - 97.4)
\$25,000-\$49,999	85	2,200	93.0	(86.8 - 99.2)	4,857	860,300	89.5	(83.6 - 95.5)
\$50,000-\$74,999	61	1,600	92.2	(84.0 - 100)	2,648	547,100	94.6	(92.2 - 96.9)
≥ \$75,000	*	*	*	*	3,582	1,185,100	92.2	(88.2 - 96.1)

¹ Defined as having ever had a mammogram among women aged 40 and older

² Sample size from survey

³ Estimated number of women aged 40 and older who have ever had a mammogram (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 24b. Ever had a mammogram among women aged 40 and older¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	352	93.1	(89.9 - 96.4)	19,491	91.1	(88.7 - 93.6)
Gender						
Male	-	-	-	-	-	-
Female	352	93.1	(89.9 - 96.4)	19,491	91.1	(88.7 - 93.6)
Race/Ethnicity						
White Non-Hispanic	348	93.7	(90.6 - 96.7)	18,548	93.3	(92.0 - 94.7)
Black Non-Hispanic	*	*	*	342	93.1	(85.7 - 100)
Hispanic	*	*	*	333	91.0	(82.8 - 99.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	186	93.9	(90.1 - 97.6)	8,266	90.2	(86.1 - 94.2)
Some College	108	93.4	(87.1 - 99.7)	5,322	92.9	(88.9 - 96.9)
≥ College Degree	*	*	*	5,872	90.9	(86.6 - 95.3)
Income						
≤ \$24,999	109	92.2	(86.0 - 98.3)	5,691	93.0	(88.5 - 97.4)
\$25,000-\$49,999	85	91.8	(84.2 - 99.4)	4,857	88.8	(82.5 - 95.1)
\$50,000-\$74,999	61	92.7	(84.6 - 100)	2,648	94.3	(91.7 - 96.8)
≥ \$75,000	*	*	*	3,582	91.2	(86.8 - 95.6)

¹ Defined as having ever had a mammogram among women aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 25a. Had mammogram within the past 2 years among women aged 40 and older¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	350	6,800	72.9	(67.6 - 78.3)	19,376	3,760,000	80.3	(77.8 - 82.9)
Gender								
Male	-	-	-	-	-	-	-	-
Female	350	6,800	72.9	(67.6 - 78.3)	19,376	3,760,000	80.3	(77.8 - 82.9)
Age (years)								
40-49	*	*	*	*	3,842	1,122,700	74.4	(68.5 - 80.2)
50-64	151	2,900	74.9	(66.8 - 83.0)	7,428	1,487,900	87.2	(84.5 - 89.8)
≥ 65	131	2,200	75.8	(68.0 - 83.6)	8,106	1,149,400	78.5	(74.2 - 82.9)
Race/Ethnicity								
White Non-Hispanic	346	6,600	73.0	(67.6 - 78.3)	18,438	2,531,500	80.2	(78.3 - 82.2)
Black Non-Hispanic	*	*	*	*	339	667,100	85.6	(77.5 - 93.8)
Hispanic	*	*	*	*	332	419,700	86.1	(77.5 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	184	3,400	70.1	(62.5 - 77.6)	8,204	1,486,900	80.5	(76.4 - 84.6)
Some College	108	2,300	76.6	(67.3 - 85.9)	5,297	929,400	83.0	(78.8 - 87.1)
≥ College Degree	*	*	*	*	5,845	1,334,800	78.6	(73.9 - 83.2)
Income								
≤ \$24,999	*	*	*	*	5,650	882,300	80.8	(75.9 - 85.7)
\$25,000-\$49,999	*	*	*	*	4,846	732,300	76.3	(69.7 - 82.9)
\$50,000-\$74,999	*	*	*	*	2,641	494,200	85.6	(81.9 - 89.4)
≥ \$75,000	*	*	*	*	3,572	1,025,900	80.0	(75.2 - 84.8)

¹ Defined as having had a mammogram within the past 2 years among women aged 40 and older

² Sample size from survey

³ Estimated number of women aged 40 and older who have had a mammogram within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 25b. Had mammogram within the past 2 years among women aged 40 and older¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	350	72.4	(66.6 - 78.1)	19,376	79.7	(76.9 - 82.5)
Gender						
Male	-	-	-	-	-	-
Female	350	72.4	(66.6 - 78.1)	19,376	79.7	(76.9 - 82.5)
Race/Ethnicity						
White Non-Hispanic	346	72.4	(66.7 - 78.2)	18,438	79.7	(77.6 - 81.8)
Black Non-Hispanic	*	*	*	339	85.3	(76.5 - 94.2)
Hispanic	*	*	*	332	85.2	(75.6 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	184	69.1	(60.9 - 77.3)	8,204	80.1	(75.6 - 84.7)
Some College	108	76.1	(66.2 - 86.0)	5,297	82.3	(77.7 - 86.9)
≥ College Degree	*	*	*	5,845	77.8	(72.9 - 82.8)
Income						
≤ \$24,999	*	*	*	5,650	80.8	(75.4 - 86.2)
\$25,000-\$49,999	*	*	*	4,846	75.7	(68.7 - 82.6)
\$50,000-\$74,999	*	*	*	2,641	85.5	(81.6 - 89.4)
≥ \$75,000	*	*	*	3,572	79.0	(73.8 - 84.2)

¹ Defined as having had a mammogram within the past 2 years among women aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 26a. Had mammogram within the past 2 years among women aged 50 and older¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	282	5,100	75.3	(69.6 - 81.0)	15,534	2,637,300	83.2	(80.7 - 85.7)
Gender								
Male	-	-	-	-	-	-	-	-
Female	282	5,100	75.3	(69.6 - 81.0)	15,534	2,637,300	83.2	(80.7 - 85.7)
Age (years)								
50-64	151	2,900	74.9	(66.8 - 83.0)	7,428	1,487,900	87.2	(84.5 - 89.8)
≥ 65	131	2,200	75.8	(68.0 - 83.6)	8,106	1,149,400	78.5	(74.2 - 82.9)
Race/Ethnicity								
White Non-Hispanic	278	5,000	75.4	(69.8 - 81.1)	14,887	1,907,000	82.3	(80.3 - 84.4)
Black Non-Hispanic	*	*	*	*	249	385,600	89.9	(83.2 - 96.7)
Hispanic	*	*	*	*	220	286,700	93.9	(89.1 - 98.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	152	2,700	73.0	(65.0 - 80.9)	7,026	1,094,500	83.3	(80.0 - 86.6)
Some College	*	*	*	*	4,124	646,700	87.3	(84.2 - 90.3)
≥ College Degree	*	*	*	*	4,356	890,500	80.6	(75.1 - 86.1)
Income								
≤ \$24,999	*	*	*	*	4,991	649,100	81.0	(76.8 - 85.2)
\$25,000-\$49,999	68	1,400	84.3	(74.6 - 94.0)	3,913	533,300	78.6	(70.9 - 86.2)
\$50,000-\$74,999	*	*	*	*	1,943	359,300	89.6	(86.3 - 93.0)
≥ \$75,000	*	*	*	*	2,303	601,300	84.6	(79.3 - 89.9)

¹ Defined as having had a mammogram within the past 2 years among women aged 50 and older

² Sample size from survey

³ Estimated number of women aged 50 and older who have had a mammogram within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 26b. Had mammogram within the past 2 years among women aged 50 and older¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	282	75.3	(69.6 - 81.1)	15,534	83.1	(80.3 - 85.8)
Gender						
Male	-	-	-	-	-	-
Female	282	75.3	(69.6 - 81.1)	15,534	83.1	(80.3 - 85.8)
Race/Ethnicity						
White Non-Hispanic	278	75.5	(69.8 - 81.2)	14,887	82.1	(79.9 - 84.3)
Black Non-Hispanic	*	*	*	249	90.4	(83.6 - 97.2)
Hispanic	*	*	*	220	94.1	(89.3 - 99.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	152	72.8	(64.7 - 80.9)	7,026	83.6	(80.1 - 87.0)
Some College	*	*	*	4,124	87.3	(84.1 - 90.5)
≥ College Degree	*	*	*	4,356	79.9	(73.9 - 86.0)
Income						
≤ \$24,999	*	*	*	4,991	81.3	(76.9 - 85.6)
\$25,000-\$49,999	68	84.6	(75.2 - 94.0)	3,913	78.1	(69.9 - 86.4)
\$50,000-\$74,999	*	*	*	1,943	90.2	(87.1 - 93.4)
≥ \$75,000	*	*	*	2,303	83.9	(77.9 - 89.9)

¹ Defined as having had a mammogram within the past 2 years among women aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 27a. Ever had a Pap test among women¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	423	13,500	90.9	(85.9 - 95.9)	23,381	7,045,300	92.1	(90.0 - 94.3)
Gender								
Male	-	-	-	-	-	-	-	-
Female	423	13,500	90.9	(85.9 - 95.9)	23,381	7,045,300	92.1	(90.0 - 94.3)
Age (years)								
18-34	*	*	*	*	2,377	1,896,600	85.2	(79.3 - 91.0)
35-44	*	*	*	*	3,023	1,301,500	92.6	(87.7 - 97.5)
45-54	96	3,000	97.0	(92.5 - 100)	4,657	1,420,300	98.3	(97.0 - 99.6)
55-64	102	2,300	98.1	(94.6 - 100)	5,187	1,071,200	97.2	(95.0 - 99.3)
≥ 65	133	2,800	94.8	(91.1 - 98.5)	8,137	1,355,700	92.3	(88.5 - 96.1)
Race/Ethnicity								
White Non-Hispanic	414	12,800	91.7	(86.6 - 96.8)	21,976	4,439,700	94.2	(92.1 - 96.3)
Black Non-Hispanic	*	*	*	*	481	1,231,600	91.2	(84.0 - 98.4)
Hispanic	*	*	*	*	533	984,500	90.1	(85.3 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	211	6,200	90.9	(84.5 - 97.4)	9,526	2,540,500	88.3	(83.8 - 92.7)
Some College	*	*	*	*	6,462	1,782,200	93.5	(90.4 - 96.6)
≥ College Degree	81	3,400	96.6	(90.2 - 100)	7,345	2,643,200	95.0	(92.2 - 97.9)
Income								
≤ \$24,999	124	3,200	88.5	(81.2 - 95.9)	6,526	1,717,200	93.7	(90.9 - 96.6)
\$25,000-\$49,999	*	*	*	*	5,787	1,378,400	91.0	(85.6 - 96.4)
\$50,000-\$74,999	*	*	*	*	3,310	933,900	96.0	(91.8 - 100)
≥ \$75,000	*	*	*	*	4,570	1,943,300	95.4	(92.6 - 98.2)

¹ Defined as having ever had a Pap test among women

² Sample size from survey

³ Estimated number of women who have ever had a Pap test (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wyoming County**

Table 27b. Ever had a Pap test among women¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	423	89.7	(83.9 - 95.6)	23,381	91.8	(89.6 - 94.0)
Gender						
Male	-	-	-	-	-	-
Female	423	89.7	(83.9 - 95.6)	23,381	91.8	(89.6 - 94.0)
Race/Ethnicity						
White Non-Hispanic	414	90.3	(84.3 - 96.4)	21,976	94.0	(91.8 - 96.2)
Black Non-Hispanic	*	*	*	481	90.8	(83.5 - 98.1)
Hispanic	*	*	*	533	89.7	(84.6 - 94.7)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	211	89.8	(82.0 - 97.5)	9,526	87.8	(83.2 - 92.4)
Some College	*	*	*	6,462	93.1	(89.7 - 96.4)
≥ College Degree	81	96.2	(89.0 - 100)	7,345	94.9	(92.0 - 97.7)
Income						
≤ \$24,999	124	87.9	(79.6 - 96.2)	6,526	93.3	(90.1 - 96.5)
\$25,000-\$49,999	*	*	*	5,787	91.1	(85.7 - 96.5)
\$50,000-\$74,999	*	*	*	3,310	96.0	(91.9 - 100)
≥ \$75,000	*	*	*	4,570	95.0	(91.9 - 98.1)

¹ Defined as having ever had a Pap test among women

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 28a. Had a Pap test within the past 3 years among women¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	421	10,800	73.4	(67.6 - 79.3)	23,109	6,205,600	82.2	(79.8 - 84.6)
Gender								
Male	-	-	-	-	-	-	-	-
Female	421	10,800	73.4	(67.6 - 79.3)	23,109	6,205,600	82.2	(79.8 - 84.6)
Age (years)								
18-34	*	*	*	*	2,371	1,818,400	81.8	(75.5 - 88.1)
35-44	*	*	*	*	3,012	1,220,300	86.9	(81.8 - 92.0)
45-54	96	2,500	83.1	(74.4 - 91.7)	4,640	1,311,500	92.0	(89.8 - 94.3)
55-64	102	1,800	75.4	(65.4 - 85.4)	5,152	924,300	84.5	(81.0 - 88.0)
≥ 65	131	1,700	58.9	(49.8 - 68.1)	7,934	931,100	66.4	(61.8 - 71.0)
Race/Ethnicity								
White Non-Hispanic	412	10,200	73.1	(67.2 - 79.1)	21,720	3,807,100	81.5	(79.2 - 83.8)
Black Non-Hispanic	*	*	*	*	473	1,142,800	86.7	(79.1 - 94.3)
Hispanic	*	*	*	*	527	894,900	83.6	(76.3 - 91.0)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	210	4,700	69.0	(60.9 - 77.0)	9,384	2,132,600	76.0	(71.1 - 81.0)
Some College	*	*	*	*	6,401	1,594,300	84.0	(80.4 - 87.5)
≥ College Degree	*	*	*	*	7,281	2,406,000	86.9	(83.7 - 90.1)
Income								
≤ \$24,999	*	*	*	*	6,441	1,453,900	82.0	(77.9 - 86.0)
\$25,000-\$49,999	*	*	*	*	5,740	1,182,000	78.3	(72.6 - 83.9)
\$50,000-\$74,999	*	*	*	*	3,296	825,000	85.0	(78.0 - 91.9)
≥ \$75,000	*	*	*	*	4,555	1,810,200	89.0	(85.7 - 92.2)

¹ Defined as having had a Pap test within the past 3 years among women

² Sample size from survey

³ Estimated number of women who have had a Pap test within the past 3 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 28b. Had a Pap test within the past 3 years among women¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	421	73.7	(67.2 - 80.3)	23,109	82.7	(80.2 - 85.1)
Gender						
Male	-	-	-	-	-	-
Female	421	73.7	(67.2 - 80.3)	23,109	82.7	(80.2 - 85.1)
Race/Ethnicity						
White Non-Hispanic	412	73.3	(66.5 - 80.0)	21,720	82.3	(80.0 - 84.6)
Black Non-Hispanic	*	*	*	473	87.0	(79.3 - 94.6)
Hispanic	*	*	*	527	83.4	(75.9 - 90.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	210	68.8	(59.7 - 77.8)	9,384	76.4	(71.3 - 81.5)
Some College	*	*	*	6,401	84.2	(80.5 - 88.0)
≥ College Degree	*	*	*	7,281	87.5	(84.3 - 90.6)
Income						
≤ \$24,999	*	*	*	6,441	82.7	(78.5 - 86.9)
\$25,000-\$49,999	*	*	*	5,740	79.2	(73.5 - 84.8)
\$50,000-\$74,999	*	*	*	3,296	85.7	(78.7 - 92.8)
≥ \$75,000	*	*	*	4,555	88.9	(85.5 - 92.3)

¹ Defined as having had a Pap test within the past 3 years among women

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 29a. Ever had a digital rectal exam among men aged 40 and older¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	204	7,700	75.3	(68.1 - 82.5)	12,033	3,176,500	73.8	(69.7 - 77.8)
Gender								
Male	204	7,700	75.3	(68.1 - 82.5)	12,033	3,176,500	73.8	(69.7 - 77.8)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	3,098	922,800	66.8	(59.9 - 73.8)
55-64	*	*	*	*	3,504	865,600	87.1	(82.5 - 91.7)
≥ 65	64	2,300	94.7	(89.9 - 99.6)	4,348	953,900	92.9	(91.0 - 94.9)
Race/Ethnicity								
White Non-Hispanic	198	7,300	74.7	(67.3 - 82.1)	11,411	2,452,100	80.3	(77.7 - 82.9)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	120	3,900	71.3	(61.8 - 80.8)	5,013	931,600	64.1	(57.3 - 71.0)
Some College	*	*	*	*	2,695	674,200	75.5	(67.8 - 83.2)
≥ College Degree	*	*	*	*	4,302	1,568,000	80.5	(74.0 - 87.0)
Income								
≤ \$24,999	*	*	*	*	2,573	573,300	65.9	(56.0 - 75.9)
\$25,000-\$49,999	*	*	*	*	3,168	530,100	74.6	(67.2 - 82.1)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,309	1,351,100	78.0	(72.6 - 83.5)

¹ Defined as having ever had a digital rectal exam among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have ever had a digital rectal exam (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 29b. Ever had a digital rectal exam among men aged 40 and older¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	204	76.3	(69.2 - 83.5)	12,033	73.7	(69.6 - 77.8)
Gender						
Male	204	76.3	(69.2 - 83.5)	12,033	73.7	(69.6 - 77.8)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	198	75.6	(68.3 - 83.0)	11,411	79.9	(77.2 - 82.5)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	120	73.4	(64.2 - 82.6)	5,013	64.7	(57.8 - 71.7)
Some College	*	*	*	2,695	75.0	(67.1 - 82.9)
≥ College Degree	*	*	*	4,302	80.3	(73.8 - 86.8)
Income						
≤ \$24,999	*	*	*	2,573	66.9	(56.9 - 76.9)
\$25,000-\$49,999	*	*	*	3,168	74.8	(67.3 - 82.2)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,309	77.5	(72.0 - 83.0)

¹ Defined as having ever had a digital rectal exam among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 30a. Had digital rectal exam within the past 2 years among men aged 40 and older¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	203	5,300	51.9	(43.8 - 60.0)	11,923	2,380,300	55.8	(51.8 - 59.8)
Gender								
Male	203	5,300	51.9	(43.8 - 60.0)	11,923	2,380,300	55.8	(51.8 - 59.8)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	1,078	246,800	27.5	(18.5 - 36.5)
45-54	*	*	*	*	3,074	650,000	47.6	(40.8 - 54.4)
55-64	*	*	*	*	3,467	673,500	68.1	(62.4 - 73.8)
≥ 65	63	2,000	83.1	(73.6 - 92.7)	4,304	809,900	79.7	(75.0 - 84.3)
Race/Ethnicity								
White Non-Hispanic	197	5,100	52.4	(44.2 - 60.7)	11,305	1,875,600	62.2	(59.2 - 65.2)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	4,963	718,700	49.6	(43.2 - 56.1)
Some College	*	*	*	*	2,665	511,500	58.5	(50.8 - 66.3)
≥ College Degree	*	*	*	*	4,277	1,148,400	59.4	(52.8 - 66.0)
Income								
≤ \$24,999	*	*	*	*	2,553	388,000	45.0	(35.0 - 55.0)
\$25,000-\$49,999	*	*	*	*	3,139	422,400	60.5	(53.0 - 68.0)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,295	978,300	56.9	(51.1 - 62.7)

¹ Defined as having had a digital rectal exam within the past 2 years among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have had a digital rectal exam within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 30b. Had digital rectal exam within the past 2 years among men aged 40 and older¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	203	53.7	(45.5 - 61.8)	11,923	55.9	(51.7 - 60.0)
Gender						
Male	203	53.7	(45.5 - 61.8)	11,923	55.9	(51.7 - 60.0)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	197	54.2	(45.9 - 62.4)	11,305	62.0	(58.9 - 65.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	4,963	50.5	(43.7 - 57.3)
Some College	*	*	*	2,665	58.3	(50.4 - 66.2)
≥ College Degree	*	*	*	4,277	59.2	(52.4 - 65.9)
Income						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,139	60.8	(53.1 - 68.4)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,295	56.5	(50.5 - 62.4)

¹ Defined as having had a digital rectal exam within the past 2 years among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 31a. Ever had a prostate specific antigen test among men aged 40 and older¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	196	5,900	59.8	(51.5 - 68.0)	11,557	2,822,600	67.6	(63.9 - 71.2)
Gender								
Male	196	5,900	59.8	(51.5 - 68.0)	11,557	2,822,600	67.6	(63.9 - 71.2)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,938	760,600	57.4	(50.6 - 64.3)
55-64	*	*	*	*	3,373	811,400	83.4	(78.8 - 88.1)
≥ 65	62	2,000	85.6	(76.6 - 94.7)	4,215	905,400	90.4	(87.1 - 93.6)
Race/Ethnicity								
White Non-Hispanic	190	5,600	59.2	(50.8 - 67.6)	10,950	1,965,400	67.0	(63.9 - 70.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	4,775	829,200	59.0	(52.4 - 65.6)
Some College	*	*	*	*	2,584	535,000	61.9	(53.7 - 70.2)
≥ College Degree	*	*	*	*	4,176	1,455,700	76.8	(72.5 - 81.1)
Income								
≤ \$24,999	*	*	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	*	3,048	444,100	64.1	(56.0 - 72.3)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,188	1,211,100	72.3	(67.6 - 77.0)

¹ Defined as having ever had a prostate specific antigen test among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have ever had a prostate specific antigen test (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 31b. Ever had a prostate specific antigen test among men aged 40 and older¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	196	61.0	(52.8 - 69.2)	11,557	67.4	(63.7 - 71.1)
Gender						
Male	196	61.0	(52.8 - 69.2)	11,557	67.4	(63.7 - 71.1)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	190	60.4	(52.0 - 68.8)	10,950	66.6	(63.4 - 69.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	4,775	59.7	(52.9 - 66.4)
Some College	*	*	*	2,584	61.4	(53.0 - 69.9)
≥ College Degree	*	*	*	4,176	76.3	(71.9 - 80.7)
Income						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,048	64.2	(56.1 - 72.4)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,188	71.5	(66.6 - 76.3)

¹ Defined as having ever had a prostate specific antigen test among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wyoming County**

Table 32a. Had prostate specific antigen test within the past 2 years among men aged 40 and older¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	195	4,700	48.1	(39.9 - 56.3)	11,437	2,493,800	59.9	(56.0 - 63.9)
Gender								
Male	195	4,700	48.1	(39.9 - 56.3)	11,437	2,493,800	59.9	(56.0 - 63.9)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,917	659,400	49.9	(42.8 - 57.0)
55-64	*	*	*	*	3,326	719,700	74.5	(69.4 - 79.7)
≥ 65	*	*	*	*	4,166	830,000	83.3	(78.4 - 88.3)
Race/Ethnicity								
White Non-Hispanic	189	4,600	49.0	(40.7 - 57.4)	10,836	1,762,200	60.4	(57.3 - 63.5)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	4,727	734,200	52.4	(45.7 - 59.1)
Some College	*	*	*	*	2,558	483,800	56.3	(48.2 - 64.4)
≥ College Degree	*	*	*	*	4,133	1,273,200	67.4	(61.6 - 73.3)
Income								
≤ \$24,999	*	*	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	*	3,021	410,800	59.6	(51.5 - 67.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,170	1,080,100	64.6	(59.1 - 70.0)

¹ Defined as having had a prostate specific antigen test within the past 2 years among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have had a prostate specific antigen test within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 32b. Had prostate specific antigen test within the past 2 years among men aged 40 and older¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	195	49.8	(41.6 - 58.0)	11,437	59.8	(55.8 - 63.9)
Gender						
Male	195	49.8	(41.6 - 58.0)	11,437	59.8	(55.8 - 63.9)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	189	50.6	(42.2 - 58.9)	10,836	60.1	(56.9 - 63.4)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	4,727	52.9	(45.9 - 59.9)
Some College	*	*	*	2,558	56.0	(47.7 - 64.2)
≥ College Degree	*	*	*	4,133	67.1	(61.2 - 73.1)
Income						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,021	59.8	(51.7 - 67.9)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,170	63.9	(58.3 - 69.6)

¹ Defined as having had a prostate specific antigen test within the past 2 years among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 33a. Prostate cancer among men aged 40 and older¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	208	500	5.0	(2.0 - 8.0)	12,133	178,200	4.1	(3.0 - 5.2)
Gender								
Male	208	500	5.0	(2.0 - 8.0)	12,133	178,200	4.1	(3.0 - 5.2)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	3,120	6,100	0.4	(0.0 - 0.9)
55-64	*	*	*	*	3,532	31,600	3.1	(1.5 - 4.8)
≥ 65	*	*	*	*	4,388	140,400	13.6	(9.7 - 17.4)
Race/Ethnicity								
White Non-Hispanic	201	500	5.3	(2.1 - 8.4)	11,502	143,700	4.7	(3.5 - 5.8)
Black Non-Hispanic	*	*	*	*	184	23,500	5.1	(0.0 - 11.0)
Hispanic	*	*	*	*	197	7,700	1.6	(0.0 - 3.7)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	5,065	46,300	3.1	(2.2 - 4.1)
Some College	*	*	*	*	2,709	39,200	4.4	(1.3 - 7.5)
≥ College Degree	*	*	*	*	4,336	92,500	4.7	(3.0 - 6.5)
Income								
≤ \$24,999	*	*	*	*	2,597	35,800	4.1	(2.2 - 6.0)
\$25,000-\$49,999	*	*	*	*	3,196	29,600	4.1	(2.9 - 5.4)
\$50,000-\$74,999	*	*	*	*	2,007	20,300	2.8	(1.2 - 4.4)
≥ \$75,000	*	*	*	*	3,322	56,500	3.2	(1.7 - 4.8)

¹ Defined as having ever been told by a health professional that respondent had prostate cancer among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have ever been told they had prostate cancer (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 33b. Prostate cancer among men aged 40 and older¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	208	6.0	(2.5 - 9.6)	12,133	4.5	(3.2 - 5.7)
Gender						
Male	208	6.0	(2.5 - 9.6)	12,133	4.5	(3.2 - 5.7)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	201	6.3	(2.6 - 10.1)	11,502	5.1	(3.8 - 6.4)
Black Non-Hispanic	*	*	*	184	5.7	(0.0 - 12.8)
Hispanic	*	*	*	197	1.8	(0.0 - 4.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	5,065	3.3	(2.3 - 4.3)
Some College	*	*	*	2,709	5.0	(1.1 - 9.0)
≥ College Degree	*	*	*	4,336	5.2	(3.1 - 7.2)
Income						
≤ \$24,999	*	*	*	2,597	4.5	(2.2 - 6.7)
\$25,000-\$49,999	*	*	*	3,196	4.3	(3.0 - 5.6)
\$50,000-\$74,999	*	*	*	2,007	3.0	(1.1 - 5.0)
≥ \$75,000	*	*	*	3,322	3.5	(1.6 - 5.4)

¹ Defined as having ever been told by a health professional that respondent had prostate cancer among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 34a. Home blood stool test ever used among adults aged 50 and older¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	447	6,500	46.9	(41.6 - 52.3)	25,322	2,288,500	38.0	(35.5 - 40.4)
Gender								
Male	163	3,000	44.3	(35.9 - 52.8)	9,571	1,081,900	38.9	(35.1 - 42.6)
Female	284	3,400	49.5	(43.0 - 56.0)	15,751	1,206,600	37.2	(34.0 - 40.4)
Age (years)								
50-64	251	3,000	36.1	(29.4 - 42.8)	12,834	1,192,000	34.0	(30.8 - 37.2)
≥ 65	196	3,400	63.6	(56.1 - 71.2)	12,472	1,093,400	43.6	(39.8 - 47.4)
Race/Ethnicity								
White Non-Hispanic	438	6,100	46.0	(40.6 - 51.4)	24,213	1,754,500	39.1	(37.1 - 41.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	248	3,500	47.5	(40.3 - 54.7)	11,146	823,500	34.9	(30.5 - 39.2)
Some College	124	1,800	46.2	(36.3 - 56.2)	6,261	480,300	37.0	(32.6 - 41.4)
≥ College Degree	*	*	*	*	7,851	973,700	41.6	(37.8 - 45.3)
Income								
≤ \$24,999	*	*	*	*	7,222	505,000	35.4	(29.7 - 41.2)
\$25,000-\$49,999	126	2,200	58.2	(48.6 - 67.7)	6,609	467,600	40.3	(35.2 - 45.4)
\$50,000-\$74,999	*	*	*	*	3,500	313,000	38.2	(32.6 - 43.9)
≥ \$75,000	*	*	*	*	4,647	657,800	37.3	(33.0 - 41.6)

¹ Defined as having ever used a home blood stool test among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have ever used a home blood stool test (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 34b. Home blood stool test ever used among adults aged 50 and older¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	447	48.1	(42.5 - 53.7)	25,322	38.1	(35.5 - 40.8)
Gender						
Male	163	46.9	(38.3 - 55.4)	9,571	39.4	(35.3 - 43.5)
Female	284	49.6	(43.1 - 56.2)	15,751	36.9	(33.5 - 40.3)
Race/Ethnicity						
White Non-Hispanic	438	47.1	(41.5 - 52.8)	24,213	38.9	(36.7 - 41.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	248	48.3	(40.8 - 55.8)	11,146	35.1	(30.3 - 39.9)
Some College	*	*	*	6,261	36.6	(32.0 - 41.3)
≥ College Degree	*	*	*	7,851	41.9	(37.9 - 46.0)
Income						
≤ \$24,999	*	*	*	7,222	35.9	(29.5 - 42.4)
\$25,000-\$49,999	126	60.1	(50.4 - 69.8)	6,609	40.2	(34.8 - 45.6)
\$50,000-\$74,999	*	*	*	3,500	38.1	(31.9 - 44.3)
≥ \$75,000	*	*	*	4,647	37.5	(32.9 - 42.2)

¹ Defined as having ever used a home blood stool test among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 35a. Home blood stool test used within the past year among adults aged 50 and older¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	441	2,700	19.6	(15.5 - 23.7)	25,074	863,300	14.4	(12.6 - 16.2)
Gender								
Male	163	1,300	18.6	(12.4 - 24.8)	9,508	444,400	16.0	(13.2 - 18.9)
Female	278	1,400	20.5	(15.2 - 25.9)	15,566	418,900	13.1	(10.8 - 15.4)
Age (years)								
50-64	251	1,200	14.9	(10.0 - 19.7)	12,759	454,400	13.0	(10.6 - 15.5)
≥ 65	190	1,400	27.1	(19.9 - 34.2)	12,299	406,400	16.4	(13.7 - 19.0)
Race/Ethnicity								
White Non-Hispanic	432	2,500	19.3	(15.2 - 23.4)	23,971	634,300	14.3	(12.8 - 15.7)
Black Non-Hispanic	*	*	*	*	396	136,000	18.6	(9.4 - 27.8)
Hispanic	*	*	*	*	349	53,400	9.9	(2.5 - 17.2)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	245	1,500	21.0	(15.2 - 26.7)	11,026	288,600	12.3	(9.4 - 15.3)
Some College	122	800	19.7	(12.1 - 27.2)	6,208	212,500	16.4	(12.5 - 20.4)
≥ College Degree	74	400	15.2	(5.8 - 24.5)	7,778	352,400	15.2	(12.5 - 17.9)
Income								
≤ \$24,999	132	900	24.4	(15.8 - 33.1)	7,157	196,300	13.9	(10.1 - 17.6)
\$25,000-\$49,999	126	1,000	26.2	(17.5 - 35.0)	6,562	221,500	19.2	(14.1 - 24.3)
\$50,000-\$74,999	*	*	*	*	3,469	111,900	13.7	(10.1 - 17.4)
≥ \$75,000	68	300	12.3	(4.7 - 20.0)	4,623	205,100	11.7	(8.8 - 14.6)

¹ Defined as having used a home blood stool test within the past year among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have used a home blood stool test within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 35b. Home blood stool test used within the past year among adults aged 50 and older¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	441	20.2	(15.8 - 24.6)	25,074	14.6	(12.6 - 16.7)
Gender						
Male	163	20.1	(13.5 - 26.7)	9,508	16.4	(13.2 - 19.6)
Female	278	20.4	(15.0 - 25.8)	15,566	12.9	(10.4 - 15.4)
Race/Ethnicity						
White Non-Hispanic	432	20.2	(15.7 - 24.6)	23,971	14.2	(12.7 - 15.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	349	10.2	(1.7 - 18.7)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	245	22.0	(15.8 - 28.2)	11,026	12.7	(9.2 - 16.1)
Some College	122	20.0	(12.0 - 28.0)	6,208	16.6	(12.3 - 20.9)
≥ College Degree	74	15.4	(5.7 - 25.1)	7,778	15.3	(12.4 - 18.3)
Income						
≤ \$24,999	132	25.4	(16.2 - 34.7)	7,157	14.3	(9.8 - 18.7)
\$25,000-\$49,999	126	27.2	(17.9 - 36.5)	6,562	19.3	(13.8 - 24.8)
\$50,000-\$74,999	*	*	*	3,469	13.8	(9.8 - 17.7)
≥ \$75,000	68	13.9	(4.9 - 22.8)	4,623	11.8	(8.6 - 15.0)

¹ Defined as having used a home blood stool test within the past year among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 36a. Home blood stool test used within the past 2 years among adults aged 50 and older¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	441	3,900	29.0	(24.2 - 33.8)	25,074	1,218,000	20.4	(18.3 - 22.5)
Gender								
Male	163	2,000	29.7	(22.2 - 37.2)	9,508	593,100	21.4	(18.3 - 24.5)
Female	278	1,900	28.3	(22.3 - 34.2)	15,566	624,900	19.5	(16.6 - 22.4)
Age (years)								
50-64	251	2,000	23.8	(17.9 - 29.7)	12,759	629,900	18.1	(15.4 - 20.7)
≥ 65	190	2,000	37.2	(29.4 - 45.0)	12,299	585,500	23.6	(20.2 - 27.0)
Race/Ethnicity								
White Non-Hispanic	432	3,800	28.8	(24.0 - 33.6)	23,971	877,500	19.7	(18.2 - 21.3)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	245	2,100	29.0	(22.6 - 35.4)	11,026	454,600	19.4	(15.6 - 23.3)
Some College	122	1,300	32.2	(22.7 - 41.7)	6,208	262,200	20.3	(16.3 - 24.2)
≥ College Degree	*	*	*	*	7,778	491,400	21.2	(18.1 - 24.2)
Income								
≤ \$24,999	132	1,300	36.1	(26.3 - 45.9)	7,157	263,900	18.6	(14.6 - 22.7)
\$25,000-\$49,999	126	1,400	35.8	(26.3 - 45.3)	6,562	285,900	24.8	(19.6 - 30.0)
\$50,000-\$74,999	*	*	*	*	3,469	167,500	20.6	(15.6 - 25.5)
≥ \$75,000	68	500	18.6	(9.3 - 27.9)	4,623	294,900	16.8	(13.7 - 20.0)

¹ Defined as having used a home blood stool test within the past 2 years among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have used a home blood stool test within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 36b. Home blood stool test used within the past 2 years among adults aged 50 and older¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	441	29.9	(24.8 - 35.0)	25,074	20.5	(18.2 - 22.8)
Gender						
Male	163	31.1	(23.4 - 38.8)	9,508	21.7	(18.3 - 25.1)
Female	278	28.3	(22.3 - 34.3)	15,566	19.4	(16.3 - 22.5)
Race/Ethnicity						
White Non-Hispanic	432	29.9	(24.7 - 35.0)	23,971	19.7	(18.0 - 21.3)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	245	29.8	(23.0 - 36.5)	11,026	19.5	(15.4 - 23.7)
Some College	*	*	*	6,208	20.4	(16.1 - 24.7)
≥ College Degree	*	*	*	7,778	21.4	(18.1 - 24.7)
Income						
≤ \$24,999	*	*	*	7,157	18.9	(14.2 - 23.6)
\$25,000-\$49,999	126	36.3	(26.4 - 46.1)	6,562	24.8	(19.3 - 30.3)
\$50,000-\$74,999	*	*	*	3,469	20.6	(15.3 - 25.9)
≥ \$75,000	*	*	*	4,623	16.9	(13.4 - 20.4)

¹ Defined as having used a home blood stool test within the past 2 years among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 37a. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	452	8,400	60.3	(55.0 - 65.6)	25,357	4,019,100	66.6	(64.1 - 69.1)
Gender								
Male	166	4,700	66.8	(58.7 - 75.0)	9,582	1,894,100	68.0	(64.2 - 71.7)
Female	286	3,700	53.6	(47.1 - 60.1)	15,775	2,125,000	65.5	(62.1 - 68.8)
Age (years)								
50-54	*	*	*	*	4,179	718,900	50.6	(44.6 - 56.5)
55-64	167	2,700	57.4	(49.0 - 65.9)	8,695	1,493,500	71.0	(67.3 - 74.8)
≥ 65	198	3,800	70.3	(63.4 - 77.3)	12,466	1,802,000	72.1	(68.6 - 75.7)
Race/Ethnicity								
White Non-Hispanic	443	8,200	60.6	(55.3 - 65.9)	24,248	3,093,900	68.8	(66.8 - 70.8)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	250	4,000	55.0	(47.8 - 62.2)	11,121	1,445,100	61.2	(56.7 - 65.8)
Some College	125	2,700	64.2	(54.4 - 74.0)	6,287	835,300	64.3	(59.7 - 69.0)
≥ College Degree	*	*	*	*	7,884	1,732,400	73.7	(70.0 - 77.4)
Income								
≤ \$24,999	*	*	*	*	7,220	892,600	62.7	(56.8 - 68.6)
\$25,000-\$49,999	128	2,400	62.2	(52.7 - 71.7)	6,619	697,300	60.1	(54.6 - 65.7)
\$50,000-\$74,999	*	*	*	*	3,514	613,700	74.2	(68.9 - 79.6)
≥ \$75,000	*	*	*	*	4,665	1,294,400	73.3	(69.2 - 77.5)

¹ Defined as having ever had a sigmoidoscopy or colonoscopy among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have ever had a sigmoidoscopy or colonoscopy (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 37b. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	452	61.8	(56.4 - 67.2)	25,357	66.6	(63.9 - 69.3)
Gender						
Male	166	68.1	(60.2 - 76.1)	9,582	68.5	(64.5 - 72.5)
Female	286	53.5	(47.0 - 60.1)	15,775	64.8	(61.2 - 68.4)
Race/Ethnicity						
White Non-Hispanic	443	62.0	(56.5 - 67.4)	24,248	68.7	(66.5 - 70.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	250	56.9	(49.5 - 64.3)	11,121	61.3	(56.4 - 66.2)
Some College	*	*	*	6,287	63.8	(58.8 - 68.8)
≥ College Degree	*	*	*	7,884	74.0	(70.2 - 77.9)
Income						
≤ \$24,999	*	*	*	7,220	63.0	(56.7 - 69.3)
\$25,000-\$49,999	128	63.8	(54.1 - 73.4)	6,619	59.7	(53.8 - 65.6)
\$50,000-\$74,999	*	*	*	3,514	74.4	(68.3 - 80.5)
≥ \$75,000	*	*	*	4,665	73.6	(69.3 - 78.0)

¹ Defined as having ever had a sigmoidoscopy or colonoscopy among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 38a. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	450	8,100	58.2	(52.8 - 63.5)	25,169	3,855,800	64.6	(62.1 - 67.1)
Gender								
Male	165	4,600	65.1	(56.8 - 73.4)	9,521	1,835,400	66.4	(62.6 - 70.1)
Female	285	3,500	51.1	(44.6 - 57.6)	15,648	2,020,400	63.1	(59.8 - 66.4)
Age (years)								
50-54	*	*	*	*	4,169	700,500	49.3	(43.3 - 55.3)
55-64	167	2,600	55.7	(47.2 - 64.2)	8,652	1,455,100	69.4	(65.7 - 73.2)
≥ 65	196	3,700	69.1	(62.0 - 76.2)	12,331	1,695,700	69.5	(65.9 - 73.1)
Race/Ethnicity								
White Non-Hispanic	441	7,800	58.4	(53.0 - 63.8)	24,069	2,981,500	66.6	(64.6 - 68.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	249	3,900	53.3	(46.0 - 60.6)	11,017	1,360,400	59.0	(54.5 - 63.5)
Some College	*	*	*	*	6,251	807,100	62.4	(57.8 - 67.1)
≥ College Degree	*	*	*	*	7,841	1,682,500	71.7	(68.0 - 75.4)
Income								
≤ \$24,999	*	*	*	*	7,163	858,900	61.0	(55.1 - 67.0)
\$25,000-\$49,999	127	2,200	58.0	(48.3 - 67.7)	6,584	664,900	57.7	(52.2 - 63.2)
\$50,000-\$74,999	*	*	*	*	3,501	602,600	73.1	(67.7 - 78.5)
≥ \$75,000	*	*	*	*	4,655	1,260,400	71.5	(67.3 - 75.7)

¹ Defined as having had a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have had a sigmoidoscopy or colonoscopy within the past 10 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 38b. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	450	59.8	(54.3 - 65.3)	25,169	64.6	(61.9 - 67.3)
Gender						
Male	165	66.6	(58.5 - 74.6)	9,521	66.9	(62.9 - 70.9)
Female	285	51.0	(44.5 - 57.6)	15,648	62.4	(58.9 - 66.0)
Race/Ethnicity						
White Non-Hispanic	441	59.9	(54.4 - 65.4)	24,069	66.5	(64.3 - 68.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	249	55.2	(47.8 - 62.7)	11,017	59.0	(54.1 - 63.9)
Some College	*	*	*	6,251	61.9	(56.9 - 66.9)
≥ College Degree	*	*	*	7,841	72.1	(68.2 - 75.9)
Income						
≤ \$24,999	*	*	*	7,163	61.3	(54.9 - 67.6)
\$25,000-\$49,999	127	59.9	(50.0 - 69.7)	6,584	57.2	(51.4 - 63.0)
\$50,000-\$74,999	*	*	*	3,501	73.3	(67.2 - 79.4)
≥ \$75,000	*	*	*	4,655	71.8	(67.4 - 76.2)

¹ Defined as having had a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 39a. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	442	9,000	65.7	(60.6 - 70.9)	24,792	4,049,200	69.1	(66.7 - 71.6)
Gender								
Male	163	5,000	71.2	(63.3 - 79.0)	9,418	1,945,200	70.7	(67.0 - 74.3)
Female	279	4,100	60.1	(53.7 - 66.6)	15,374	2,104,100	67.8	(64.6 - 71.1)
Age (years)								
50-54	*	*	*	*	4,132	751,200	54.1	(48.0 - 60.1)
55-64	162	3,000	65.0	(56.6 - 73.3)	8,366	1,487,600	73.7	(70.0 - 77.4)
≥ 65	193	4,100	76.5	(70.1 - 83.0)	12,294	1,810,500	74.0	(70.6 - 77.4)
Race/Ethnicity								
White Non-Hispanic	433	8,600	65.3	(60.1 - 70.5)	23,726	3,091,400	70.4	(68.4 - 72.4)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	244	4,400	61.6	(54.4 - 68.7)	10,872	1,461,600	64.2	(59.8 - 68.7)
Some College	122	2,700	67.6	(57.9 - 77.3)	6,170	873,300	68.4	(63.9 - 72.8)
≥ College Degree	*	*	*	*	7,708	1,709,000	74.6	(70.9 - 78.3)
Income								
≤ \$24,999	134	2,400	65.2	(55.7 - 74.6)	7,092	923,300	66.0	(60.1 - 71.9)
\$25,000-\$49,999	125	2,600	68.9	(59.8 - 78.0)	6,526	730,000	64.2	(58.9 - 69.6)
\$50,000-\$74,999	*	*	*	*	3,467	621,500	76.6	(71.4 - 81.9)
≥ \$75,000	*	*	*	*	4,615	1,292,500	73.8	(69.8 - 77.9)

¹ Defined as having used a home blood stool test within the past year or a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 years and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have used a home blood stool test in the past year or sigmoidoscopy or colonoscopy within the past 10 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 39b. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	442	67.3	(62.1 - 72.5)	24,792	69.3	(66.7 - 71.9)
Gender						
Male	163	72.9	(65.4 - 80.4)	9,418	71.5	(67.6 - 75.4)
Female	279	59.9	(53.4 - 66.4)	15,374	67.2	(63.7 - 70.7)
Race/Ethnicity						
White Non-Hispanic	433	66.8	(61.5 - 72.1)	23,726	70.3	(68.1 - 72.4)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	244	63.4	(56.2 - 70.6)	10,872	64.5	(59.8 - 69.3)
Some College	122	68.8	(58.9 - 78.7)	6,170	68.2	(63.4 - 72.9)
≥ College Degree	*	*	*	7,708	75.0	(71.2 - 78.8)
Income						
≤ \$24,999	134	67.8	(58.5 - 77.1)	7,092	66.4	(60.2 - 72.6)
\$25,000-\$49,999	125	70.9	(62.0 - 79.8)	6,526	63.8	(58.2 - 69.5)
\$50,000-\$74,999	*	*	*	3,467	76.7	(70.6 - 82.7)
≥ \$75,000	*	*	*	4,615	74.4	(70.2 - 78.6)

¹ Defined as having used a home blood stool test within the past year or a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 years and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 40a. High blood pressure among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	674	11,700	35.0	(29.6 - 40.3)	38,125	3,974,800	26.8	(24.9 - 28.7)
Gender								
Male	248	6,900	37.0	(28.2 - 45.7)	14,611	1,960,500	27.7	(24.7 - 30.7)
Female	426	4,800	32.5	(27.2 - 37.7)	23,514	2,014,300	26.0	(23.6 - 28.3)
Age (years)								
18-34	*	*	*	*	3,935	312,200	7.0	(4.2 - 9.8)
35-44	92	1,400	22.5	(12.5 - 32.5)	4,984	426,200	14.8	(11.6 - 18.1)
45-54	155	2,300	34.9	(26.1 - 43.7)	7,825	862,400	30.3	(25.9 - 34.8)
55-64	169	2,400	51.3	(42.8 - 59.7)	8,765	890,800	41.6	(37.6 - 45.7)
≥ 65	198	3,700	68.1	(60.7 - 75.5)	12,616	1,483,200	58.5	(54.7 - 62.3)
Race/Ethnicity								
White Non-Hispanic	656	10,900	35.5	(30.5 - 40.5)	35,814	2,650,900	28.4	(26.9 - 30.0)
Black Non-Hispanic	*	*	*	*	736	558,500	26.2	(19.6 - 32.8)
Hispanic	*	*	*	*	863	538,100	23.8	(17.0 - 30.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	355	7,400	43.6	(34.5 - 52.6)	15,688	1,851,200	33.1	(29.4 - 36.9)
Some College	194	2,900	29.2	(21.0 - 37.5)	9,906	851,100	24.1	(20.9 - 27.3)
≥ College Degree	125	1,400	21.5	(13.6 - 29.5)	12,452	1,253,400	22.3	(19.7 - 24.9)
Income								
≤ \$24,999	*	*	*	*	9,640	1,153,000	32.8	(27.6 - 38.1)
\$25,000-\$49,999	187	3,400	41.5	(32.5 - 50.6)	9,657	807,800	27.9	(23.7 - 32.1)
\$50,000-\$74,999	*	*	*	*	5,773	507,600	25.8	(21.4 - 30.1)
≥ \$75,000	*	*	*	*	8,560	942,500	21.0	(18.4 - 23.6)

¹ Defined as having ever been told by a health professional that respondent has high blood pressure

² Sample size from survey

³ Estimated number of adults who have ever been told by a health professional that they have high blood pressure (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 40b. High blood pressure among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	674	33.8	(28.6 - 39.0)	38,125	25.7	(23.8 - 27.5)
Gender						
Male	248	38.0	(29.4 - 46.5)	14,611	27.7	(24.7 - 30.7)
Female	426	28.6	(23.2 - 34.0)	23,514	23.8	(21.5 - 26.0)
Race/Ethnicity						
White Non-Hispanic	656	34.2	(29.2 - 39.1)	35,814	27.2	(25.6 - 28.7)
Black Non-Hispanic	*	*	*	736	24.9	(18.4 - 31.3)
Hispanic	*	*	*	863	23.3	(16.6 - 30.0)
Other Non-Hispanic	*	*	*	712	19.3	(9.5 - 29.1)
Education						
≤ High School	355	42.8	(34.1 - 51.6)	15,688	32.1	(28.4 - 35.8)
Some College	194	28.4	(20.1 - 36.6)	9,906	22.8	(19.7 - 25.9)
≥ College Degree	125	19.8	(11.9 - 27.7)	12,452	21.2	(18.6 - 23.8)
Income						
≤ \$24,999	*	*	*	9,640	31.8	(26.7 - 37.0)
\$25,000-\$49,999	187	40.3	(31.2 - 49.5)	9,657	26.9	(22.8 - 30.9)
\$50,000-\$74,999	*	*	*	5,773	24.8	(20.4 - 29.2)
≥ \$75,000	111	22.4	(12.7 - 32.1)	8,560	19.9	(17.3 - 22.5)

¹ Defined as having ever been told by a health professional that respondent has high blood pressure

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wyoming County**

Table 41a. Blood pressure medication among adults with high blood pressure¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	302	8,900	76.0	(67.2 - 84.8)	15,288	3,145,800	79.2	(76.0 - 82.4)
Gender								
Male	*	*	*	*	6,080	1,432,100	73.1	(67.6 - 78.5)
Female	184	4,100	84.4	(76.7 - 92.2)	9,208	1,713,700	85.2	(82.1 - 88.2)
Age (years)								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,315	645,500	74.9	(66.8 - 82.9)
55-64	86	2,000	83.0	(74.3 - 91.7)	4,045	782,300	87.8	(84.5 - 91.1)
≥ 65	136	3,500	93.8	(89.2 - 98.3)	7,675	1,383,400	93.4	(91.1 - 95.7)
Race/Ethnicity								
White Non-Hispanic	295	8,500	78.0	(70.3 - 85.6)	14,442	2,117,300	79.9	(77.5 - 82.4)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	7,354	1,482,800	80.2	(74.7 - 85.6)
Some College	79	2,600	91.0	(84.7 - 97.2)	3,868	632,000	74.3	(67.8 - 80.7)
≥ College Degree	*	*	*	*	4,035	1,013,200	80.9	(76.8 - 84.9)
Income								
≤ \$24,999	*	*	*	*	4,898	948,200	82.3	(75.7 - 88.9)
\$25,000-\$49,999	*	*	*	*	4,088	668,300	82.8	(77.6 - 88.0)
\$50,000-\$74,999	*	*	*	*	1,971	391,500	77.2	(67.7 - 86.6)
≥ \$75,000	*	*	*	*	2,397	683,700	72.5	(66.1 - 78.9)

¹ Defined as currently taking medicine for high blood pressure among adults with high blood pressure

² Sample size from survey

³ Estimated number of adults with high blood pressure who are taking medicine for high blood pressure (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 41b. Blood pressure medication among adults with high blood pressure¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	302	75.7	(66.9 - 84.5)	15,288	78.5	(75.2 - 81.7)
Gender						
Male	*	*	*	6,080	73.3	(67.9 - 78.7)
Female	184	82.6	(72.9 - 92.2)	9,208	84.0	(80.7 - 87.2)
Race/Ethnicity						
White Non-Hispanic	295	77.4	(69.5 - 85.3)	14,442	79.0	(76.5 - 81.6)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	7,354	79.7	(74.3 - 85.2)
Some College	79	91.4	(85.2 - 97.7)	3,868	72.5	(65.7 - 79.3)
≥ College Degree	*	*	*	4,035	80.4	(76.1 - 84.6)
Income						
≤ \$24,999	*	*	*	4,898	81.6	(75.0 - 88.3)
\$25,000-\$49,999	*	*	*	4,088	82.1	(76.7 - 87.5)
\$50,000-\$74,999	*	*	*	1,971	76.4	(66.7 - 86.1)
≥ \$75,000	*	*	*	2,397	71.6	(64.8 - 78.3)

¹ Defined as currently taking medicine for high blood pressure among adults with high blood pressure

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 42a. Ever had blood cholesterol checked among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	662	24,700	75.4	(68.1 - 82.7)	37,450	11,496,600	80.3	(77.7 - 83.0)
Gender								
Male	*	*	*	*	14,389	5,288,400	77.7	(73.3 - 82.0)
Female	420	11,700	80.4	(74.1 - 86.7)	23,061	6,208,200	82.7	(79.6 - 85.8)
Age (years)								
18-34	*	*	*	*	3,684	2,296,400	55.3	(48.7 - 61.9)
35-44	91	4,900	76.6	(66.9 - 86.2)	4,883	2,264,200	81.6	(76.6 - 86.5)
45-54	153	6,200	93.3	(88.9 - 97.7)	7,731	2,539,400	91.0	(87.0 - 94.9)
55-64	168	4,500	94.6	(91.0 - 98.3)	8,718	2,014,900	95.4	(93.5 - 97.2)
≥ 65	193	5,000	94.8	(91.2 - 98.4)	12,434	2,381,700	95.9	(94.3 - 97.6)
Race/Ethnicity								
White Non-Hispanic	645	23,600	78.7	(72.8 - 84.5)	35,198	7,832,600	86.3	(84.8 - 87.9)
Black Non-Hispanic	*	*	*	*	720	1,588,900	77.8	(68.4 - 87.2)
Hispanic	*	*	*	*	842	1,224,600	56.8	(46.8 - 66.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,342	3,795,700	70.6	(65.6 - 75.6)
Some College	*	*	*	*	9,732	2,599,800	77.4	(71.7 - 83.2)
≥ College Degree	125	5,700	86.8	(77.3 - 96.2)	12,296	5,008,300	91.3	(88.9 - 93.6)
Income								
≤ \$24,999	*	*	*	*	9,437	2,277,500	68.4	(61.5 - 75.3)
\$25,000-\$49,999	185	7,000	87.1	(79.7 - 94.4)	9,492	2,053,000	71.9	(64.5 - 79.3)
\$50,000-\$74,999	*	*	*	*	5,721	1,736,100	90.0	(87.7 - 92.3)
≥ \$75,000	109	5,400	85.3	(75.4 - 95.2)	8,460	4,019,300	92.2	(90.8 - 93.6)

¹ Defined as ever having blood cholesterol checked

² Sample size from survey

³ Estimated number of adults who have ever had blood cholesterol checked (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 42b. Ever had blood cholesterol checked among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	662	74.7	(67.7 - 81.8)	37,450	79.9	(77.3 - 82.5)
Gender						
Male	*	*	*	14,389	78.0	(73.9 - 82.1)
Female	420	77.3	(70.0 - 84.5)	23,061	81.6	(78.4 - 84.8)
Race/Ethnicity						
White Non-Hispanic	645	77.5	(71.7 - 83.4)	35,198	85.6	(84.0 - 87.2)
Black Non-Hispanic	*	*	*	720	77.5	(68.2 - 86.7)
Hispanic	*	*	*	842	57.2	(47.4 - 67.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	15,342	70.2	(65.2 - 75.1)
Some College	*	*	*	9,732	76.7	(71.1 - 82.3)
≥ College Degree	*	*	*	12,296	90.9	(88.6 - 93.2)
Income						
≤ \$24,999	*	*	*	9,437	68.1	(61.3 - 74.9)
\$25,000-\$49,999	185	85.4	(77.1 - 93.7)	9,492	71.6	(64.4 - 78.7)
\$50,000-\$74,999	*	*	*	5,721	89.6	(87.2 - 92.0)
≥ \$75,000	*	*	*	8,460	91.7	(90.1 - 93.2)

¹ Defined as ever having blood cholesterol checked

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 43a. Blood cholesterol checked within the past 5 years among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	660	23,200	71.2	(64.0 - 78.4)	37,201	11,035,200	77.8	(75.1 - 80.4)
Gender								
Male	*	*	*	*	14,307	5,020,700	74.5	(70.2 - 78.9)
Female	418	11,000	75.6	(69.0 - 82.2)	22,894	6,014,600	80.7	(77.6 - 83.8)
Age (years)								
18-34	*	*	*	*	3,647	2,179,800	53.3	(46.7 - 59.9)
35-44	*	*	*	*	4,862	2,161,200	78.0	(72.8 - 83.2)
45-54	153	6,000	90.3	(85.3 - 95.3)	7,697	2,408,600	86.6	(82.4 - 90.8)
55-64	167	4,300	92.2	(87.9 - 96.4)	8,677	1,957,900	93.3	(91.4 - 95.3)
≥ 65	192	4,900	93.0	(89.1 - 96.9)	12,318	2,327,800	94.8	(93.1 - 96.6)
Race/Ethnicity								
White Non-Hispanic	643	22,200	74.0	(68.1 - 80.0)	34,960	7,486,600	83.4	(81.8 - 85.0)
Black Non-Hispanic	*	*	*	*	719	1,561,700	76.5	(67.1 - 85.9)
Hispanic	*	*	*	*	838	1,184,000	55.6	(45.6 - 65.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,215	3,612,500	68.3	(63.2 - 73.3)
Some College	*	*	*	*	9,673	2,524,200	75.4	(69.8 - 81.1)
≥ College Degree	*	*	*	*	12,236	4,806,500	88.1	(85.5 - 90.6)
Income								
≤ \$24,999	*	*	*	*	9,350	2,214,800	66.9	(60.1 - 73.8)
\$25,000-\$49,999	185	6,800	85.2	(77.7 - 92.7)	9,454	1,936,600	68.8	(61.5 - 76.1)
\$50,000-\$74,999	*	*	*	*	5,701	1,640,400	85.9	(82.4 - 89.5)
≥ \$75,000	*	*	*	*	8,435	3,872,400	89.2	(87.5 - 90.9)

¹ Defined as having had blood cholesterol checked within the past 5 years

² Sample size from survey

³ Estimated number of adults who have had blood cholesterol checked within the past 5 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 43b. Blood cholesterol checked within the past 5 years among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	660	70.3	(63.3 - 77.2)	37,201	77.3	(74.7 - 79.9)
Gender						
Male	*	*	*	14,307	74.8	(70.7 - 79.0)
Female	418	72.0	(64.5 - 79.5)	22,894	79.5	(76.3 - 82.7)
Race/Ethnicity						
White Non-Hispanic	643	72.7	(66.6 - 78.7)	34,960	82.6	(81.0 - 84.3)
Black Non-Hispanic	*	*	*	719	76.2	(66.9 - 85.5)
Hispanic	*	*	*	838	55.9	(46.1 - 65.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	15,215	67.8	(62.9 - 72.8)
Some College	*	*	*	9,673	74.7	(69.1 - 80.2)
≥ College Degree	*	*	*	12,236	87.5	(85.0 - 90.1)
Income						
≤ \$24,999	*	*	*	9,350	66.6	(59.8 - 73.4)
\$25,000-\$49,999	185	83.4	(75.0 - 91.8)	9,454	68.5	(61.4 - 75.6)
\$50,000-\$74,999	*	*	*	5,701	85.4	(81.7 - 89.0)
≥ \$75,000	*	*	*	8,435	88.5	(86.7 - 90.3)

¹ Defined as having had blood cholesterol checked within the past 5 years

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 44a. Cardiovascular disease among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	666	3,200	9.6	(3.7 - 15.4)	37,810	1,135,000	7.7	(6.5 - 8.8)
Gender								
Male	*	*	*	*	14,477	664,600	9.4	(7.5 - 11.4)
Female	422	900	6.2	(3.9 - 8.4)	23,333	470,400	6.1	(4.8 - 7.4)
Age (years)								
18-34	*	*	*	*	3,930	100,800	2.3	(0.0 - 4.8)
35-44	*	*	*	*	4,971	103,400	3.6	(1.5 - 5.7)
45-54	*	*	*	*	7,768	131,300	4.6	(2.9 - 6.4)
55-64	*	*	*	*	8,699	224,200	10.6	(8.3 - 12.9)
≥ 65	194	1,300	24.2	(17.1 - 31.2)	12,442	575,300	23.0	(19.9 - 26.1)
Race/Ethnicity								
White Non-Hispanic	649	2,100	6.9	(4.9 - 8.9)	35,522	772,200	8.3	(7.4 - 9.2)
Black Non-Hispanic	*	*	*	*	728	140,700	6.6	(3.3 - 9.9)
Hispanic	*	*	*	*	854	163,300	7.3	(2.0 - 12.5)
Other Non-Hispanic	*	*	*	*	706	58,800	5.2	(0.9 - 9.5)
Education								
≤ High School	*	*	*	*	15,486	590,800	10.7	(8.2 - 13.2)
Some College	193	500	4.7	(1.9 - 7.6)	9,849	227,100	6.4	(4.7 - 8.1)
≥ College Degree	*	*	*	*	12,393	314,600	5.6	(4.3 - 6.9)
Income								
≤ \$24,999	*	*	*	*	9,505	448,100	12.9	(9.1 - 16.7)
\$25,000-\$49,999	184	800	10.2	(5.8 - 14.6)	9,571	179,300	6.2	(4.8 - 7.6)
\$50,000-\$74,999	*	*	*	*	5,751	166,900	8.5	(5.5 - 11.4)
≥ \$75,000	*	*	*	*	8,538	183,500	4.1	(2.9 - 5.3)

¹ Defined as having ever been told by a health professional that respondent had a heart attack or angina or a stroke

² Sample size from survey

³ Estimated number of adults with cardiovascular disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 44b. Cardiovascular disease among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	666	9.5	(4.1 - 14.9)	37,810	7.6	(6.4 - 8.7)
Gender						
Male	244	12.8	(3.5 - 22.2)	14,477	9.7	(7.8 - 11.7)
Female	422	5.4	(3.2 - 7.6)	23,333	5.6	(4.3 - 6.9)
Race/Ethnicity						
White Non-Hispanic	649	7.2	(5.0 - 9.3)	35,522	8.2	(7.2 - 9.1)
Black Non-Hispanic	*	*	*	728	6.8	(3.2 - 10.4)
Hispanic	*	*	*	854	7.0	(2.0 - 12.0)
Other Non-Hispanic	*	*	*	706	5.5	(0.9 - 10.0)
Education						
≤ High School	*	*	*	15,486	10.5	(8.0 - 13.0)
Some College	193	4.9	(1.8 - 8.1)	9,849	6.4	(4.5 - 8.2)
≥ College Degree	*	*	*	12,393	5.5	(4.2 - 6.9)
Income						
≤ \$24,999	*	*	*	9,505	12.8	(9.0 - 16.6)
\$25,000-\$49,999	184	10.7	(5.9 - 15.4)	9,571	6.0	(4.7 - 7.4)
\$50,000-\$74,999	*	*	*	5,751	8.5	(5.4 - 11.6)
≥ \$75,000	*	*	*	8,538	4.0	(2.7 - 5.2)

¹ Defined as having ever been told by a health professional that respondent had a heart attack or angina or a stroke

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wyoming County**

Table 45a. Coronary heart disease among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	667	3,000	9.1	(3.2 - 14.9)	37,838	929,400	6.3	(5.3 - 7.3)
Gender								
Male	*	*	*	*	14,479	598,600	8.5	(6.6 - 10.4)
Female	422	800	5.4	(3.3 - 7.6)	23,359	330,800	4.3	(3.4 - 5.2)
Age (years)								
18-34	*	*	*	*	3,932	63,400	1.4	(0.0 - 3.6)
35-44	*	*	*	*	4,973	95,300	3.3	(1.2 - 5.4)
45-54	*	*	*	*	7,778	108,300	3.8	(2.1 - 5.6)
55-64	*	*	*	*	8,703	188,600	8.9	(6.7 - 11.1)
≥ 65	194	1,200	22.4	(15.4 - 29.3)	12,452	473,800	18.9	(16.1 - 21.7)
Race/Ethnicity								
White Non-Hispanic	650	1,900	6.4	(4.4 - 8.3)	35,552	636,700	6.9	(6.0 - 7.7)
Black Non-Hispanic	*	*	*	*	728	113,100	5.3	(2.3 - 8.4)
Hispanic	*	*	*	*	853	128,000	5.7	(1.1 - 10.3)
Other Non-Hispanic	*	*	*	*	705	51,500	4.6	(0.5 - 8.7)
Education								
≤ High School	*	*	*	*	15,485	474,500	8.6	(6.3 - 10.8)
Some College	194	400	4.3	(1.6 - 7.0)	9,860	193,000	5.5	(3.9 - 7.1)
≥ College Degree	*	*	*	*	12,411	259,700	4.6	(3.4 - 5.8)
Income								
≤ \$24,999	*	*	*	*	9,507	372,300	10.7	(7.3 - 14.1)
\$25,000-\$49,999	184	800	9.8	(5.5 - 14.1)	9,585	136,300	4.7	(3.8 - 5.7)
\$50,000-\$74,999	*	*	*	*	5,755	154,600	7.8	(4.9 - 10.7)
≥ \$75,000	*	*	*	*	8,541	151,800	3.4	(2.3 - 4.5)

¹ Defined as having ever been told by a health professional that respondent had a heart attack or angina

² Sample size from survey

³ Estimated number of adults with coronary heart disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 45b. Coronary heart disease among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	667	9.1	(3.6 - 14.5)	37,838	6.3	(5.2 - 7.3)
Gender						
Male	245	12.5	(3.1 - 21.9)	14,479	8.8	(6.9 - 10.8)
Female	422	4.8	(2.7 - 6.9)	23,359	3.9	(3.0 - 4.8)
Race/Ethnicity						
White Non-Hispanic	650	6.7	(4.5 - 8.8)	35,552	6.8	(5.9 - 7.7)
Black Non-Hispanic	*	*	*	728	5.6	(2.2 - 9.0)
Hispanic	*	*	*	853	5.4	(1.1 - 9.6)
Other Non-Hispanic	*	*	*	705	4.8	(0.4 - 9.2)
Education						
≤ High School	*	*	*	15,485	8.5	(6.2 - 10.7)
Some College	194	4.6	(1.5 - 7.6)	9,860	5.5	(3.7 - 7.2)
≥ College Degree	*	*	*	12,411	4.7	(3.4 - 5.9)
Income						
≤ \$24,999	*	*	*	9,507	10.7	(7.2 - 14.1)
\$25,000-\$49,999	184	10.3	(5.6 - 14.9)	9,585	4.7	(3.6 - 5.7)
\$50,000-\$74,999	*	*	*	5,755	8.0	(4.8 - 11.1)
≥ \$75,000	*	*	*	8,541	3.3	(2.1 - 4.5)

¹ Defined as having ever been told by a health professional that respondent had a heart attack or angina

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wyoming County**

Table 46a. Ever diagnosed with diabetes among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	674	2,900	8.7	(6.5 - 10.9)	38,174	1,382,700	9.3	(8.0 - 10.5)
Gender								
Male	248	1,000	5.6	(3.3 - 8.0)	14,626	530,400	7.5	(6.2 - 8.8)
Female	426	1,900	12.5	(8.8 - 16.3)	23,548	852,300	11.0	(8.9 - 13.0)
Age (years)								
18-34	*	*	*	*	3,941	150,200	3.4	(1.0 - 5.8)
35-44	*	*	*	*	4,987	165,100	5.7	(3.6 - 7.9)
45-54	155	500	7.7	(3.6 - 11.7)	7,830	279,000	9.8	(6.7 - 12.8)
55-64	169	700	14.6	(8.8 - 20.4)	8,771	332,900	15.5	(12.3 - 18.8)
≥ 65	198	1,200	21.8	(15.5 - 28.2)	12,645	455,600	17.9	(15.0 - 20.8)
Race/Ethnicity								
White Non-Hispanic	656	2,900	9.5	(7.1 - 11.8)	35,862	809,300	8.7	(7.8 - 9.5)
Black Non-Hispanic	*	*	*	*	739	261,000	12.1	(6.6 - 17.6)
Hispanic	*	*	*	*	862	227,300	10.1	(5.4 - 14.8)
Other Non-Hispanic	*	*	*	*	711	85,100	7.6	(2.3 - 12.8)
Education								
≤ High School	355	1,900	11.2	(7.4 - 15.0)	15,709	730,000	13.0	(10.5 - 15.6)
Some College	194	700	7.2	(3.7 - 10.8)	9,918	303,500	8.6	(5.9 - 11.2)
≥ College Degree	125	300	4.5	(1.5 - 7.5)	12,464	337,500	6.0	(4.8 - 7.2)
Income								
≤ \$24,999	180	800	10.3	(5.4 - 15.1)	9,652	543,400	15.5	(11.4 - 19.5)
\$25,000-\$49,999	187	1,000	12.2	(7.2 - 17.3)	9,667	291,000	10.0	(7.5 - 12.6)
\$50,000-\$74,999	117	400	5.5	(1.7 - 9.2)	5,777	151,000	7.6	(5.0 - 10.3)
≥ \$75,000	*	*	*	*	8,565	220,800	4.9	(3.8 - 6.1)

¹ Defined as having ever been told by a doctor that respondent had diabetes, excluding pre-diabetes and women with diabetes only when pregnant

² Sample size from survey

³ Estimated number of adults who have diabetes (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 46b. Ever diagnosed with diabetes among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	674	8.7	(6.3 - 11.0)	38,174	9.0	(7.8 - 10.3)
Gender						
Male	248	6.2	(3.6 - 8.8)	14,626	7.5	(6.1 - 8.9)
Female	426	11.7	(7.7 - 15.8)	23,548	10.4	(8.4 - 12.4)
Race/Ethnicity						
White Non-Hispanic	656	9.4	(6.9 - 12.0)	35,862	8.4	(7.5 - 9.2)
Black Non-Hispanic	*	*	*	739	11.4	(6.0 - 16.7)
Hispanic	*	*	*	862	10.1	(5.4 - 14.9)
Other Non-Hispanic	*	*	*	711	7.7	(2.3 - 13.0)
Education						
≤ High School	355	11.4	(7.4 - 15.5)	15,709	12.6	(10.0 - 15.2)
Some College	194	7.2	(3.5 - 11.0)	9,918	8.3	(5.6 - 10.9)
≥ College Degree	125	4.0	(1.2 - 6.8)	12,464	5.9	(4.7 - 7.2)
Income						
≤ \$24,999	180	9.9	(5.1 - 14.8)	9,652	15.2	(11.1 - 19.2)
\$25,000-\$49,999	187	12.7	(7.2 - 18.2)	9,667	9.7	(7.1 - 12.2)
\$50,000-\$74,999	117	5.3	(1.5 - 9.1)	5,777	7.6	(4.9 - 10.2)
≥ \$75,000	*	*	*	8,565	4.7	(3.6 - 5.9)

¹ Defined as having ever been told by a doctor that respondent had diabetes, excluding pre-diabetes and women with diabetes only when pregnant

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 47a. Ever diagnosed with asthma (lifetime) among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	673	3,600	10.8	(7.6 - 14.0)	38,096	2,225,100	15.0	(13.2 - 16.8)
Gender								
Male	248	1,200	6.5	(2.6 - 10.4)	14,601	875,800	12.4	(10.1 - 14.7)
Female	425	2,400	16.2	(11.3 - 21.1)	23,495	1,349,300	17.4	(14.6 - 20.1)
Age (years)								
18-34	*	*	*	*	3,931	825,100	18.5	(13.8 - 23.3)
35-44	*	*	*	*	4,972	431,700	15.1	(11.2 - 19.1)
45-54	154	900	13.3	(7.1 - 19.5)	7,819	365,900	12.8	(9.7 - 16.0)
55-64	169	400	7.8	(3.8 - 11.8)	8,756	316,700	14.8	(11.9 - 17.7)
≥ 65	198	700	13.0	(7.3 - 18.8)	12,618	285,700	11.2	(8.6 - 13.8)
Race/Ethnicity								
White Non-Hispanic	655	3,400	11.2	(7.9 - 14.5)	35,783	1,365,700	14.6	(13.0 - 16.3)
Black Non-Hispanic	*	*	*	*	737	367,000	17.2	(10.2 - 24.2)
Hispanic	*	*	*	*	863	317,100	14.1	(9.0 - 19.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	355	1,600	9.6	(5.8 - 13.4)	15,675	831,600	14.9	(12.0 - 17.8)
Some College	193	1,300	13.1	(6.7 - 19.6)	9,893	587,500	16.6	(13.1 - 20.1)
≥ College Degree	125	700	10.5	(2.3 - 18.7)	12,444	739,000	13.2	(10.4 - 15.9)
Income								
≤ \$24,999	179	1,200	15.5	(7.8 - 23.2)	9,643	708,900	20.2	(14.9 - 25.5)
\$25,000-\$49,999	187	900	11.3	(5.5 - 17.1)	9,641	389,800	13.4	(10.0 - 16.9)
\$50,000-\$74,999	117	700	10.1	(2.9 - 17.3)	5,762	257,800	13.2	(10.0 - 16.5)
≥ \$75,000	*	*	*	*	8,562	577,900	12.8	(10.4 - 15.3)

¹ Defined as adults having ever been told by a health professional that they had asthma

² Sample size from survey

³ Estimated number of adults who have lifetime asthma (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 47b. Ever diagnosed with asthma (lifetime) among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	673	11.1	(7.7 - 14.5)	38,096	15.2	(13.3 - 17.0)
Gender						
Male	248	6.7	(2.8 - 10.6)	14,601	12.6	(10.3 - 14.9)
Female	425	16.6	(11.0 - 22.1)	23,495	17.5	(14.8 - 20.3)
Race/Ethnicity						
White Non-Hispanic	655	11.5	(8.0 - 15.0)	35,783	14.9	(13.2 - 16.6)
Black Non-Hispanic	*	*	*	737	17.1	(10.1 - 24.1)
Hispanic	*	*	*	863	14.5	(9.3 - 19.7)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	355	10.4	(6.1 - 14.6)	15,675	15.1	(12.1 - 18.0)
Some College	193	12.5	(6.2 - 18.8)	9,893	16.9	(13.4 - 20.4)
≥ College Degree	125	10.9	(1.9 - 19.9)	12,444	13.3	(10.6 - 16.0)
Income						
≤ \$24,999	179	16.3	(8.1 - 24.4)	9,643	20.3	(15.0 - 25.6)
\$25,000-\$49,999	187	11.3	(5.4 - 17.1)	9,641	13.7	(10.2 - 17.2)
\$50,000-\$74,999	117	10.5	(3.0 - 18.0)	5,762	13.5	(10.0 - 17.0)
≥ \$75,000	*	*	*	8,562	13.0	(10.6 - 15.4)

¹ Defined as adults having ever been told by a health professional that they had asthma

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 48a. Current asthma among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	671	2,500	7.5	(4.8 - 10.2)	37,933	1,415,000	9.6	(8.1 - 11.0)
Gender								
Male	247	700	3.6	(0.8 - 6.5)	14,546	440,700	6.3	(4.8 - 7.7)
Female	424	1,800	12.3	(7.7 - 16.9)	23,387	974,300	12.6	(10.2 - 15.0)
Age (years)								
18-34	*	*	*	*	3,909	467,200	10.5	(6.8 - 14.3)
35-44	*	*	*	*	4,947	246,800	8.7	(5.8 - 11.6)
45-54	152	600	9.0	(3.9 - 14.1)	7,787	240,000	8.4	(6.1 - 10.8)
55-64	169	300	5.6	(2.1 - 9.1)	8,726	231,100	10.8	(8.1 - 13.5)
≥ 65	198	600	10.2	(4.9 - 15.5)	12,564	229,900	9.1	(6.5 - 11.6)
Race/Ethnicity								
White Non-Hispanic	653	2,300	7.6	(4.8 - 10.4)	35,629	888,400	9.6	(8.2 - 11.0)
Black Non-Hispanic	*	*	*	*	736	246,900	11.6	(5.7 - 17.5)
Hispanic	*	*	*	*	859	186,800	8.3	(4.4 - 12.2)
Other Non-Hispanic	*	*	*	*	709	93,000	8.3	(1.4 - 15.2)
Education								
≤ High School	354	1,400	8.1	(4.5 - 11.7)	15,594	595,100	10.7	(8.2 - 13.2)
Some College	193	700	7.0	(2.4 - 11.6)	9,861	322,000	9.1	(7.0 - 11.3)
≥ College Degree	*	*	*	*	12,394	430,900	7.7	(5.8 - 9.7)
Income								
≤ \$24,999	179	1,000	13.9	(6.6 - 21.3)	9,600	489,200	14.0	(9.7 - 18.3)
\$25,000-\$49,999	185	400	4.7	(1.7 - 7.7)	9,590	268,400	9.3	(6.2 - 12.4)
\$50,000-\$74,999	*	*	*	*	5,741	143,400	7.4	(5.1 - 9.6)
≥ \$75,000	*	*	*	*	8,541	339,200	7.6	(5.8 - 9.4)

¹ Defined as adults having ever been told by a health professional that they had asthma and still have asthma

² Sample size from survey

³ Estimated number of adults who currently have asthma (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 48b. Current asthma among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	671	7.8	(4.8 - 10.7)	37,933	9.7	(8.2 - 11.2)
Gender						
Male	247	3.9	(0.9 - 6.8)	14,546	6.4	(4.9 - 7.9)
Female	424	12.6	(7.4 - 17.8)	23,387	12.7	(10.2 - 15.1)
Race/Ethnicity						
White Non-Hispanic	653	7.9	(4.8 - 10.9)	35,629	9.7	(8.3 - 11.2)
Black Non-Hispanic	*	*	*	736	11.5	(5.6 - 17.3)
Hispanic	*	*	*	859	8.6	(4.5 - 12.6)
Other Non-Hispanic	*	*	*	709	8.3	(1.3 - 15.3)
Education						
≤ High School	354	8.8	(4.8 - 12.8)	15,594	10.8	(8.3 - 13.3)
Some College	193	6.4	(2.1 - 10.8)	9,861	9.4	(7.2 - 11.6)
≥ College Degree	*	*	*	12,394	7.8	(5.8 - 9.7)
Income						
≤ \$24,999	179	14.7	(6.8 - 22.5)	9,600	14.1	(9.7 - 18.4)
\$25,000-\$49,999	185	4.9	(1.6 - 8.2)	9,590	9.5	(6.3 - 12.7)
\$50,000-\$74,999	*	*	*	5,741	7.5	(5.0 - 10.1)
≥ \$75,000	*	*	*	8,541	7.7	(5.9 - 9.5)

¹ Defined as adults having ever been told by a health professional that they had asthma and still have asthma

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 49a. Chronic joint symptoms among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	672	17,000	50.9	(44.6 - 57.2)	38,099	6,377,100	42.9	(40.5 - 45.3)
Gender								
Male	*	*	*	*	14,595	3,033,600	42.8	(39.0 - 46.6)
Female	424	7,400	49.9	(43.6 - 56.1)	23,504	3,343,500	43.0	(40.0 - 46.0)
Age (years)								
18-34	*	*	*	*	3,933	1,261,300	28.3	(22.6 - 34.0)
35-44	*	*	*	*	4,984	1,110,900	38.6	(33.1 - 44.0)
45-54	155	3,900	57.6	(48.5 - 66.8)	7,817	1,456,500	51.2	(46.4 - 55.9)
55-64	168	2,800	59.3	(51.0 - 67.6)	8,751	1,167,400	54.6	(50.4 - 58.8)
≥ 65	197	3,200	59.3	(51.5 - 67.0)	12,614	1,381,000	54.3	(50.5 - 58.1)
Race/Ethnicity								
White Non-Hispanic	654	16,100	52.4	(46.7 - 58.0)	35,785	4,267,600	45.8	(43.8 - 47.7)
Black Non-Hispanic	*	*	*	*	740	890,500	41.3	(32.5 - 50.0)
Hispanic	*	*	*	*	860	827,500	36.7	(27.7 - 45.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	354	9,100	53.8	(44.1 - 63.5)	15,676	2,450,000	43.8	(39.4 - 48.1)
Some College	*	*	*	*	9,900	1,598,300	45.1	(40.1 - 50.1)
≥ College Degree	*	*	*	*	12,440	2,313,100	41.1	(37.8 - 44.5)
Income								
≤ \$24,999	*	*	*	*	9,641	1,483,000	42.2	(36.3 - 48.1)
\$25,000-\$49,999	186	4,300	52.4	(43.0 - 61.9)	9,647	1,332,100	45.9	(39.8 - 52.1)
\$50,000-\$74,999	*	*	*	*	5,768	894,200	45.3	(39.5 - 51.1)
≥ \$75,000	*	*	*	*	8,549	1,874,200	41.7	(38.4 - 45.1)

¹ Defined as adults having symptoms of pain, aching or stiffness in or around a joint during the past 30 days and began more than 3 months ago

² Sample size from survey

³ Estimated number of adults who have chronic joint symptoms (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 49b. Chronic joint symptoms among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	672	49.9	(43.8 - 56.1)	38,099	42.1	(39.7 - 44.5)
Gender						
Male	248	51.4	(41.8 - 61.0)	14,595	42.6	(38.8 - 46.3)
Female	424	48.0	(41.2 - 54.9)	23,504	41.6	(38.6 - 44.7)
Race/Ethnicity						
White Non-Hispanic	654	51.1	(45.4 - 56.9)	35,785	44.7	(42.7 - 46.6)
Black Non-Hispanic	*	*	*	740	40.2	(31.5 - 48.9)
Hispanic	*	*	*	860	36.6	(27.8 - 45.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	354	53.4	(44.0 - 62.8)	15,676	43.1	(38.8 - 47.5)
Some College	*	*	*	9,900	43.9	(39.0 - 48.8)
≥ College Degree	*	*	*	12,440	40.3	(37.0 - 43.7)
Income						
≤ \$24,999	*	*	*	9,641	41.6	(35.7 - 47.4)
\$25,000-\$49,999	186	51.3	(41.6 - 60.9)	9,647	45.0	(39.0 - 51.0)
\$50,000-\$74,999	*	*	*	5,768	44.3	(38.4 - 50.2)
≥ \$75,000	*	*	*	8,549	40.8	(37.4 - 44.2)

¹ Defined as adults having symptoms of pain, aching or stiffness in or around a joint during the past 30 days and began more than 3 months ago

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 50a. Arthritis among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	673	9,900	29.4	(24.9 - 33.9)	38,013	4,102,800	27.7	(25.9 - 29.6)
Gender								
Male	247	4,200	22.5	(16.5 - 28.6)	14,555	1,642,700	23.3	(20.7 - 25.8)
Female	426	5,700	38.0	(32.4 - 43.6)	23,458	2,460,100	31.8	(29.2 - 34.4)
Age (years)								
18-34	*	*	*	*	3,927	312,100	7.0	(4.9 - 9.1)
35-44	92	1,200	18.1	(9.4 - 26.9)	4,972	412,500	14.5	(11.3 - 17.6)
45-54	155	2,500	37.6	(28.9 - 46.3)	7,789	869,100	30.7	(26.6 - 34.8)
55-64	168	2,100	44.8	(36.4 - 53.1)	8,729	999,300	47.0	(42.9 - 51.1)
≥ 65	198	3,300	61.1	(53.4 - 68.8)	12,596	1,509,800	59.6	(55.8 - 63.3)
Race/Ethnicity								
White Non-Hispanic	655	9,400	30.5	(26.2 - 34.9)	35,710	2,967,700	31.9	(30.3 - 33.6)
Black Non-Hispanic	*	*	*	*	737	588,100	27.3	(20.2 - 34.4)
Hispanic	*	*	*	*	856	380,700	17.1	(11.8 - 22.5)
Other Non-Hispanic	*	*	*	*	710	166,300	14.8	(7.8 - 21.8)
Education								
≤ High School	355	5,100	29.8	(23.0 - 36.6)	15,652	1,742,700	31.3	(27.8 - 34.8)
Some College	193	3,000	30.3	(22.4 - 38.3)	9,874	902,000	25.5	(22.3 - 28.6)
≥ College Degree	125	1,800	27.1	(18.4 - 35.7)	12,407	1,437,800	25.6	(22.9 - 28.4)
Income								
≤ \$24,999	*	*	*	*	9,618	1,112,500	32.0	(27.1 - 36.9)
\$25,000-\$49,999	186	3,000	36.2	(27.9 - 44.6)	9,624	821,900	28.4	(24.2 - 32.6)
\$50,000-\$74,999	117	1,800	25.7	(16.3 - 35.2)	5,753	542,500	27.5	(23.2 - 31.9)
≥ \$75,000	111	1,100	17.7	(10.6 - 24.8)	8,536	1,012,800	22.6	(19.8 - 25.3)

¹ Defined as adults having ever been told by a health professional that they have some form of arthritis, rheumatoid arthritis, gout, lupus or fibromyalgia (excluding back or neck)

² Sample size from survey

³ Estimated number of adults who have ever been told by a health professional they have some form of arthritis (excluding back or neck) (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 50b. Arthritis among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	673	28.6	(24.1 - 33.0)	38,013	26.6	(24.8 - 28.5)
Gender						
Male	247	24.1	(17.8 - 30.3)	14,555	23.5	(20.9 - 26.1)
Female	426	34.2	(28.4 - 40.0)	23,458	29.5	(27.0 - 32.1)
Race/Ethnicity						
White Non-Hispanic	655	29.4	(25.1 - 33.7)	35,710	30.4	(28.7 - 32.0)
Black Non-Hispanic	*	*	*	737	26.5	(19.3 - 33.6)
Hispanic	*	*	*	856	17.2	(11.8 - 22.6)
Other Non-Hispanic	*	*	*	710	14.9	(7.7 - 22.0)
Education						
≤ High School	355	29.6	(22.9 - 36.4)	15,652	30.4	(26.9 - 33.8)
Some College	193	28.7	(20.9 - 36.6)	9,874	24.1	(21.0 - 27.1)
≥ College Degree	125	25.8	(16.9 - 34.6)	12,407	24.7	(21.9 - 27.5)
Income						
≤ \$24,999	*	*	*	9,618	31.2	(26.3 - 36.1)
\$25,000-\$49,999	186	34.6	(26.2 - 43.0)	9,624	27.3	(23.2 - 31.3)
\$50,000-\$74,999	117	25.3	(15.7 - 34.8)	5,753	26.3	(21.9 - 30.7)
≥ \$75,000	111	16.5	(9.6 - 23.4)	8,536	21.6	(18.8 - 24.3)

¹ Defined as adults having ever been told by a health professional that they have some form of arthritis, rheumatoid arthritis, gout, lupus or fibromyalgia (excluding back or neck)

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 51a. Activity limitations because of physical, mental or emotional problems among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	670	6,300	18.9	(15.1 - 22.6)	37,998	3,053,000	20.6	(18.7 - 22.5)
Gender								
Male	246	2,300	12.6	(8.1 - 17.0)	14,566	1,318,400	18.6	(15.7 - 21.5)
Female	424	4,000	26.7	(21.2 - 32.2)	23,432	1,734,600	22.4	(19.8 - 25.0)
Age (years)								
18-34	*	*	*	*	3,929	580,000	13.0	(8.2 - 17.9)
35-44	92	1,600	24.8	(14.9 - 34.7)	4,973	418,600	14.5	(11.0 - 18.1)
45-54	155	1,400	21.5	(14.7 - 28.3)	7,812	620,200	21.7	(18.3 - 25.2)
55-64	167	1,000	20.3	(13.6 - 26.9)	8,739	651,600	30.5	(26.5 - 34.6)
≥ 65	196	1,500	28.2	(21.1 - 35.2)	12,545	782,700	31.1	(27.7 - 34.6)
Race/Ethnicity								
White Non-Hispanic	652	5,800	19.1	(15.4 - 22.8)	35,688	2,037,200	21.9	(20.3 - 23.5)
Black Non-Hispanic	*	*	*	*	735	429,700	20.1	(13.5 - 26.7)
Hispanic	*	*	*	*	863	448,200	19.8	(11.9 - 27.8)
Other Non-Hispanic	*	*	*	*	712	137,900	12.3	(5.2 - 19.3)
Education								
≤ High School	352	3,300	19.4	(14.1 - 24.7)	15,626	1,359,600	24.4	(20.6 - 28.1)
Some College	194	1,800	18.2	(11.9 - 24.4)	9,875	700,800	19.8	(16.5 - 23.2)
≥ College Degree	124	1,200	18.5	(9.1 - 27.9)	12,415	924,800	16.5	(14.2 - 18.7)
Income								
≤ \$24,999	*	*	*	*	9,603	1,055,800	30.1	(24.7 - 35.5)
\$25,000-\$49,999	187	1,500	18.2	(11.7 - 24.6)	9,628	630,900	21.8	(16.6 - 27.0)
\$50,000-\$74,999	117	900	12.2	(6.2 - 18.3)	5,760	408,100	20.7	(16.1 - 25.2)
≥ \$75,000	111	600	9.3	(2.9 - 15.7)	8,544	583,300	13.0	(11.1 - 14.9)

¹ Defined as activity limitations because of physical, mental or emotional problems

² Sample size from survey

³ Estimated number of adults who are limited in any activities because of physical, mental or emotional problems (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 51b. Activity limitations because of physical, mental or emotional problems among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	670	19.2	(15.3 - 23.2)	37,998	20.1	(18.2 - 22.0)
Gender						
Male	246	13.4	(8.7 - 18.0)	14,566	18.6	(15.8 - 21.4)
Female	424	26.5	(20.4 - 32.6)	23,432	21.5	(18.9 - 24.1)
Race/Ethnicity						
White Non-Hispanic	652	19.4	(15.5 - 23.4)	35,688	21.3	(19.7 - 22.9)
Black Non-Hispanic	*	*	*	735	19.9	(13.2 - 26.5)
Hispanic	*	*	*	863	19.1	(11.5 - 26.8)
Other Non-Hispanic	*	*	*	712	12.4	(5.3 - 19.6)
Education						
≤ High School	352	19.8	(14.3 - 25.2)	15,626	23.8	(20.1 - 27.4)
Some College	194	18.2	(11.7 - 24.6)	9,875	19.3	(16.0 - 22.6)
≥ College Degree	*	*	*	12,415	16.1	(13.9 - 18.3)
Income						
≤ \$24,999	*	*	*	9,603	29.7	(24.3 - 35.2)
\$25,000-\$49,999	187	18.0	(11.4 - 24.6)	9,628	21.2	(16.1 - 26.3)
\$50,000-\$74,999	117	11.9	(5.8 - 18.0)	5,760	20.1	(15.6 - 24.6)
≥ \$75,000	111	10.2	(2.9 - 17.6)	8,544	12.7	(10.8 - 14.6)

¹ Defined as activity limitations because of physical, mental or emotional problems

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wyoming County**

Table 52a. Have health problem that requires the use of special equipment among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	674	2,600	7.8	(5.6 - 10.0)	38,171	1,169,400	7.9	(6.9 - 8.9)
Gender								
Male	248	1,200	6.4	(3.3 - 9.5)	14,617	460,700	6.5	(5.3 - 7.7)
Female	426	1,400	9.5	(6.7 - 12.3)	23,554	708,700	9.1	(7.6 - 10.7)
Age (years)								
18-34	*	*	*	*	3,940	68,800	1.5	(0.7 - 2.4)
35-44	*	*	*	*	4,989	122,500	4.3	(2.4 - 6.1)
45-54	155	800	12.5	(6.6 - 18.5)	7,836	182,500	6.4	(4.7 - 8.1)
55-64	169	400	8.7	(3.8 - 13.7)	8,773	219,600	10.3	(7.8 - 12.8)
≥ 65	198	1,100	19.4	(13.2 - 25.7)	12,633	576,100	22.7	(19.0 - 26.3)
Race/Ethnicity								
White Non-Hispanic	656	2,500	8.1	(5.9 - 10.2)	35,853	802,000	8.6	(7.6 - 9.6)
Black Non-Hispanic	*	*	*	*	739	235,800	10.9	(6.4 - 15.5)
Hispanic	*	*	*	*	864	104,100	4.6	(2.1 - 7.1)
Other Non-Hispanic	*	*	*	*	715	27,600	2.5	(0.8 - 4.1)
Education								
≤ High School	355	1,600	9.7	(6.1 - 13.3)	15,711	604,200	10.8	(8.7 - 12.8)
Some College	194	700	7.3	(3.8 - 10.9)	9,914	260,000	7.3	(5.5 - 9.1)
≥ College Degree	*	*	*	*	12,463	298,900	5.3	(4.1 - 6.5)
Income								
≤ \$24,999	180	1,200	16.4	(9.3 - 23.6)	9,656	430,500	12.2	(9.6 - 14.8)
\$25,000-\$49,999	187	500	6.3	(2.0 - 10.7)	9,663	238,900	8.2	(5.9 - 10.5)
\$50,000-\$74,999	117	400	5.4	(1.3 - 9.5)	5,775	103,000	5.2	(3.6 - 6.8)
≥ \$75,000	*	*	*	*	8,566	136,000	3.0	(2.1 - 4.0)

¹ Defined as having any health problem that requires the use of special equipment such as a cane, a wheelchair, a special bed, or a special telephone

² Sample size from survey

³ Estimated number of adults who have any health problem that requires the use of special equipment (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 52b. Have health problem that requires the use of special equipment among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	674	7.6	(5.4 - 9.8)	38,171	7.6	(6.6 - 8.6)
Gender						
Male	248	6.9	(3.6 - 10.2)	14,617	6.8	(5.5 - 8.0)
Female	426	8.4	(5.7 - 11.1)	23,554	8.4	(6.9 - 9.9)
Race/Ethnicity						
White Non-Hispanic	656	7.8	(5.6 - 10.0)	35,853	8.3	(7.3 - 9.3)
Black Non-Hispanic	*	*	*	739	10.5	(6.1 - 15.0)
Hispanic	*	*	*	864	4.5	(2.1 - 6.9)
Other Non-Hispanic	*	*	*	715	2.2	(0.7 - 3.8)
Education						
≤ High School	355	9.9	(6.1 - 13.7)	15,711	10.3	(8.3 - 12.3)
Some College	194	6.7	(3.3 - 10.1)	9,914	7.1	(5.2 - 8.9)
≥ College Degree	*	*	*	12,463	5.3	(4.1 - 6.6)
Income						
≤ \$24,999	180	16.9	(9.4 - 24.4)	9,656	11.9	(9.4 - 14.5)
\$25,000-\$49,999	187	6.2	(1.8 - 10.5)	9,663	7.7	(5.5 - 10.0)
\$50,000-\$74,999	117	4.8	(1.0 - 8.6)	5,775	5.1	(3.4 - 6.8)
≥ \$75,000	*	*	*	8,566	2.9	(2.0 - 3.9)

¹ Defined as having any health problem that requires the use of special equipment such as a cane, a wheelchair, a special bed, or a special telephone

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 53a. Disability among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	671	7,100	21.2	(17.2 - 25.1)	38,016	3,394,200	22.9	(20.9 - 24.9)
Gender								
Male	246	2,900	15.7	(10.6 - 20.7)	14,568	1,469,700	20.7	(17.8 - 23.7)
Female	425	4,200	28.0	(22.5 - 33.6)	23,448	1,924,600	24.8	(22.1 - 27.5)
Age (years)								
18-34	*	*	*	*	3,931	590,300	13.3	(8.4 - 18.1)
35-44	92	1,600	24.8	(14.9 - 34.7)	4,974	447,900	15.6	(11.9 - 19.3)
45-54	155	1,800	26.2	(18.5 - 33.8)	7,811	643,900	22.6	(19.1 - 26.0)
55-64	167	1,000	20.3	(13.6 - 26.9)	8,740	702,100	33.0	(28.9 - 37.0)
≥ 65	197	2,000	36.7	(29.0 - 44.4)	12,560	1,010,000	39.9	(36.1 - 43.7)
Race/Ethnicity								
White Non-Hispanic	653	6,600	21.6	(17.7 - 25.5)	35,703	2,253,400	24.2	(22.6 - 25.9)
Black Non-Hispanic	*	*	*	*	736	511,700	23.8	(16.8 - 30.8)
Hispanic	*	*	*	*	864	484,300	21.4	(13.4 - 29.5)
Other Non-Hispanic	*	*	*	*	713	144,800	12.9	(5.8 - 19.9)
Education								
≤ High School	353	3,900	23.1	(17.2 - 28.9)	15,638	1,544,200	27.6	(23.7 - 31.4)
Some College	194	1,900	18.7	(12.4 - 25.0)	9,876	755,200	21.4	(17.9 - 24.8)
≥ College Degree	124	1,300	20.1	(10.4 - 29.7)	12,419	1,021,300	18.2	(15.9 - 20.5)
Income								
≤ \$24,999	*	*	*	*	9,613	1,179,100	33.6	(28.0 - 39.2)
\$25,000-\$49,999	187	1,600	19.6	(13.0 - 26.2)	9,629	699,000	24.2	(18.9 - 29.4)
\$50,000-\$74,999	117	1,000	13.5	(7.0 - 20.0)	5,757	430,100	21.8	(17.2 - 26.4)
≥ \$75,000	111	700	10.7	(3.7 - 17.6)	8,546	632,700	14.1	(12.1 - 16.1)

¹ Defined as having activity limitations due to physical, mental or emotional problems or have health problems that require the use of special equipment

² Sample size from survey

³ Estimated number of adults with a disability (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 53b. Disability among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	671	21.7	(17.5 - 25.9)	38,016	22.3	(20.4 - 24.3)
Gender						
Male	246	16.9	(11.7 - 22.2)	14,568	20.9	(18.0 - 23.8)
Female	425	27.6	(21.5 - 33.8)	23,448	23.6	(21.0 - 26.3)
Race/Ethnicity						
White Non-Hispanic	653	22.1	(18.0 - 26.3)	35,703	23.6	(21.9 - 25.3)
Black Non-Hispanic	*	*	*	736	23.5	(16.5 - 30.5)
Hispanic	*	*	*	864	20.6	(12.9 - 28.3)
Other Non-Hispanic	*	*	*	713	12.9	(5.8 - 20.1)
Education						
≤ High School	353	23.8	(17.7 - 29.9)	15,638	26.8	(23.0 - 30.6)
Some College	194	18.8	(12.3 - 25.4)	9,876	20.9	(17.4 - 24.3)
≥ College Degree	*	*	*	12,419	17.9	(15.5 - 20.3)
Income						
≤ \$24,999	*	*	*	9,613	33.2	(27.6 - 38.8)
\$25,000-\$49,999	187	19.4	(12.7 - 26.2)	9,629	23.5	(18.3 - 28.6)
\$50,000-\$74,999	117	13.1	(6.6 - 19.6)	5,757	21.2	(16.6 - 25.8)
≥ \$75,000	111	11.5	(3.8 - 19.2)	8,546	13.8	(11.7 - 15.8)

¹ Defined as having activity limitations due to physical, mental or emotional problems or have health problems that require the use of special equipment

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 54a. Current smoking among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	673	7,500	22.4	(16.2 - 28.7)	38,029	2,483,500	16.8	(15.1 - 18.6)
Gender								
Male	*	*	*	*	14,559	1,356,400	19.3	(16.3 - 22.2)
Female	425	3,300	22.0	(16.8 - 27.2)	23,470	1,127,100	14.6	(12.7 - 16.6)
Age (years)								
18-34	*	*	*	*	3,929	993,500	22.3	(17.5 - 27.1)
35-44	92	1,600	24.4	(14.5 - 34.3)	4,977	421,100	14.8	(11.6 - 17.9)
45-54	154	1,600	23.5	(15.8 - 31.2)	7,807	508,800	18.3	(15.4 - 21.3)
55-64	169	1,300	26.8	(19.1 - 34.4)	8,744	314,500	14.7	(12.1 - 17.3)
≥ 65	198	400	6.7	(3.4 - 10.1)	12,572	245,600	9.7	(7.2 - 12.3)
Race/Ethnicity								
White Non-Hispanic	655	5,900	19.3	(15.4 - 23.1)	35,729	1,615,500	17.3	(15.9 - 18.8)
Black Non-Hispanic	*	*	*	*	735	284,100	13.6	(8.9 - 18.2)
Hispanic	*	*	*	*	858	350,900	15.8	(9.6 - 21.9)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,654	1,237,300	22.4	(19.1 - 25.6)
Some College	193	1,900	19.4	(12.5 - 26.3)	9,875	670,600	19.1	(15.5 - 22.8)
≥ College Degree	125	600	8.9	(2.6 - 15.2)	12,421	563,300	10.0	(7.5 - 12.5)
Income								
≤ \$24,999	*	*	*	*	9,626	835,100	24.1	(19.1 - 29.1)
\$25,000-\$49,999	187	1,400	16.5	(10.2 - 22.9)	9,629	558,100	19.4	(15.6 - 23.2)
\$50,000-\$74,999	117	800	11.4	(5.3 - 17.4)	5,756	338,100	17.3	(13.0 - 21.7)
≥ \$75,000	110	900	13.7	(4.9 - 22.5)	8,549	468,700	10.4	(8.6 - 12.3)

¹ Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day or some days

² Sample size from survey

³ Estimated number of adult current smokers (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 54b. Current smoking among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	673	22.3	(16.3 - 28.3)	38,029	17.0	(15.3 - 18.8)
Gender						
Male	248	21.7	(11.9 - 31.4)	14,559	19.2	(16.3 - 22.1)
Female	425	23.1	(17.2 - 28.9)	23,470	15.0	(13.0 - 17.0)
Race/Ethnicity						
White Non-Hispanic	655	19.4	(15.4 - 23.5)	35,729	17.7	(16.2 - 19.2)
Black Non-Hispanic	*	*	*	735	13.5	(8.8 - 18.2)
Hispanic	*	*	*	858	15.9	(9.8 - 22.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	15,654	22.8	(19.5 - 26.0)
Some College	193	19.2	(12.2 - 26.2)	9,875	19.4	(15.8 - 22.9)
≥ College Degree	125	9.0	(2.1 - 15.9)	12,421	10.0	(7.6 - 12.5)
Income						
≤ \$24,999	*	*	*	9,626	24.1	(19.2 - 29.1)
\$25,000-\$49,999	187	16.6	(9.9 - 23.3)	9,629	19.9	(16.0 - 23.7)
\$50,000-\$74,999	117	11.4	(5.2 - 17.6)	5,756	17.8	(13.3 - 22.3)
≥ \$75,000	110	14.1	(4.7 - 23.5)	8,549	10.6	(8.6 - 12.5)

¹ Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day or some days

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 55a. Everyday smoking among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	673	4,700	13.9	(10.6 - 17.2)	38,029	1,626,800	11.0	(9.8 - 12.2)
Gender								
Male	248	2,200	11.7	(7.2 - 16.2)	14,559	848,400	12.0	(10.0 - 14.1)
Female	425	2,500	16.7	(12.0 - 21.3)	23,470	778,400	10.1	(8.7 - 11.5)
Age (years)								
18-34	60	900	8.6	(2.6 - 14.5)	3,929	559,800	12.6	(9.6 - 15.6)
35-44	92	1,300	19.6	(10.2 - 28.9)	4,977	315,200	11.0	(8.4 - 13.7)
45-54	154	1,300	19.9	(12.7 - 27.0)	7,807	377,000	13.6	(11.1 - 16.0)
55-64	169	900	19.6	(12.7 - 26.5)	8,744	203,000	9.5	(7.5 - 11.5)
≥ 65	198	300	5.0	(2.0 - 8.0)	12,572	171,700	6.8	(4.8 - 8.8)
Race/Ethnicity								
White Non-Hispanic	655	4,500	14.7	(11.3 - 18.1)	35,729	1,158,100	12.4	(11.2 - 13.7)
Black Non-Hispanic	*	*	*	*	735	197,700	9.4	(5.6 - 13.3)
Hispanic	*	*	*	*	858	176,200	7.9	(4.2 - 11.6)
Other Non-Hispanic	*	*	*	*	707	94,900	8.5	(3.3 - 13.7)
Education								
≤ High School	355	3,000	17.6	(12.2 - 23.0)	15,654	881,900	15.9	(13.4 - 18.5)
Some College	193	1,400	14.0	(8.2 - 19.8)	9,875	410,200	11.7	(9.6 - 13.8)
≥ College Degree	*	*	*	*	12,421	323,500	5.8	(4.3 - 7.2)
Income								
≤ \$24,999	180	1,800	24.3	(14.6 - 34.0)	9,626	452,800	13.1	(10.3 - 15.8)
\$25,000-\$49,999	187	800	9.2	(4.7 - 13.7)	9,629	425,900	14.8	(11.9 - 17.7)
\$50,000-\$74,999	117	700	9.8	(4.2 - 15.3)	5,756	222,700	11.4	(8.5 - 14.4)
≥ \$75,000	110	600	10.0	(2.1 - 17.9)	8,549	320,200	7.1	(5.5 - 8.8)

¹ Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day

² Sample size from survey

³ Estimated number of adults who have smoked at least 100 cigarettes in lifetime and currently smoking everyday (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 55b. Everyday smoking among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	673	14.0	(10.5 - 17.4)	38,029	11.3	(10.0 - 12.5)
Gender						
Male	248	11.4	(7.0 - 15.8)	14,559	12.1	(10.1 - 14.1)
Female	425	17.2	(12.0 - 22.3)	23,470	10.5	(9.0 - 12.0)
Race/Ethnicity						
White Non-Hispanic	655	14.6	(11.1 - 18.2)	35,729	12.8	(11.5 - 14.1)
Black Non-Hispanic	*	*	*	735	9.4	(5.5 - 13.4)
Hispanic	*	*	*	858	8.0	(4.3 - 11.7)
Other Non-Hispanic	*	*	*	707	8.5	(3.2 - 13.8)
Education						
≤ High School	355	18.0	(12.4 - 23.5)	15,654	16.4	(13.8 - 19.0)
Some College	193	13.9	(7.9 - 19.9)	9,875	12.1	(9.9 - 14.3)
≥ College Degree	*	*	*	12,421	5.8	(4.3 - 7.3)
Income						
≤ \$24,999	180	24.1	(14.5 - 33.7)	9,626	13.4	(10.6 - 16.2)
\$25,000-\$49,999	187	9.0	(4.4 - 13.6)	9,629	15.3	(12.3 - 18.3)
\$50,000-\$74,999	117	9.6	(4.0 - 15.1)	5,756	11.7	(8.7 - 14.8)
≥ \$75,000	110	10.2	(1.8 - 18.7)	8,549	7.2	(5.5 - 8.9)

¹ Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wyoming County**

Table 56a. Adults living in homes in which smoking is prohibited¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	670	24,500	73.4	(67.1 - 79.7)	38,045	11,997,100	80.8	(79.0 - 82.6)
Gender								
Male	*	*	*	*	14,577	5,634,400	79.4	(76.6 - 82.2)
Female	423	11,400	76.6	(71.5 - 81.7)	23,468	6,362,800	82.1	(79.9 - 84.3)
Age (years)								
18-34	*	*	*	*	3,934	3,655,200	82.1	(77.8 - 86.4)
35-44	*	*	*	*	4,985	2,479,700	86.1	(82.5 - 89.7)
45-54	154	4,700	69.4	(60.8 - 78.0)	7,808	2,194,400	77.2	(73.5 - 80.8)
55-64	169	3,300	69.9	(62.2 - 77.6)	8,753	1,732,500	81.0	(78.3 - 83.7)
≥ 65	195	4,200	79.3	(73.3 - 85.4)	12,565	1,935,400	76.5	(73.4 - 79.5)
Race/Ethnicity								
White Non-Hispanic	652	23,100	75.5	(70.7 - 80.2)	35,737	7,426,900	79.8	(78.3 - 81.2)
Black Non-Hispanic	*	*	*	*	738	1,692,100	78.6	(71.9 - 85.3)
Hispanic	*	*	*	*	863	2,003,900	88.7	(84.1 - 93.3)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,647	4,307,200	77.1	(74.0 - 80.2)
Some College	*	*	*	*	9,883	2,795,400	78.9	(75.0 - 82.8)
≥ College Degree	125	5,700	85.8	(78.6 - 93.0)	12,437	4,817,300	85.7	(83.3 - 88.2)
Income								
≤ \$24,999	*	*	*	*	9,608	2,627,700	74.9	(70.0 - 79.7)
\$25,000-\$49,999	184	6,600	81.3	(75.1 - 87.5)	9,640	2,314,500	79.8	(76.0 - 83.5)
\$50,000-\$74,999	*	*	*	*	5,766	1,551,300	78.7	(74.0 - 83.4)
≥ \$75,000	111	5,300	82.7	(72.8 - 92.6)	8,556	3,883,200	86.5	(84.3 - 88.7)

¹ Defined as having a rule that prohibits smoking anywhere inside the home

² Sample size from survey

³ Estimated number of adults who live in a home with a rule that prohibits smoking anywhere inside the home (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 56b. Adults living in homes in which smoking is prohibited¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	670	73.9	(67.8 - 80.0)	38,045	80.9	(79.2 - 82.7)
Gender						
Male	247	71.4	(61.6 - 81.3)	14,577	79.4	(76.7 - 82.2)
Female	423	77.0	(71.3 - 82.6)	23,468	82.3	(80.0 - 84.5)
Race/Ethnicity						
White Non-Hispanic	652	75.8	(71.0 - 80.6)	35,737	79.9	(78.4 - 81.4)
Black Non-Hispanic	*	*	*	738	78.6	(71.8 - 85.4)
Hispanic	*	*	*	863	88.9	(84.4 - 93.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	352	68.8	(59.1 - 78.6)	15,647	76.9	(73.8 - 80.0)
Some College	*	*	*	9,883	79.1	(75.2 - 83.0)
≥ College Degree	125	86.5	(79.0 - 93.9)	12,437	86.0	(83.5 - 88.5)
Income						
≤ \$24,999	*	*	*	9,608	74.9	(70.1 - 79.7)
\$25,000-\$49,999	184	81.8	(75.5 - 88.1)	9,640	79.8	(76.1 - 83.6)
\$50,000-\$74,999	*	*	*	5,766	78.6	(73.8 - 83.3)
≥ \$75,000	*	*	*	8,556	86.6	(84.3 - 88.9)

¹ Defined as having a rule that prohibits smoking anywhere inside the home

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 57a. Binge drinking within the past month among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	665	8,000	24.4	(17.3 - 31.5)	37,745	2,602,000	17.7	(15.7 - 19.8)
Gender								
Male	*	*	*	*	14,420	1,712,400	24.5	(20.9 - 28.1)
Female	423	2,300	15.3	(10.1 - 20.5)	23,325	889,600	11.6	(9.5 - 13.7)
Age (years)								
18-34	*	*	*	*	3,895	1,254,300	28.5	(22.8 - 34.2)
35-44	*	*	*	*	4,934	584,800	20.6	(16.5 - 24.7)
45-54	151	1,400	21.8	(14.3 - 29.2)	7,738	467,600	16.6	(13.3 - 19.9)
55-64	168	500	11.1	(6.0 - 16.2)	8,663	207,200	9.8	(7.9 - 11.8)
≥ 65	*	*	*	*	12,515	88,100	3.5	(2.5 - 4.5)
Race/Ethnicity								
White Non-Hispanic	648	6,500	21.4	(15.7 - 27.0)	35,453	1,761,900	19.2	(17.5 - 20.8)
Black Non-Hispanic	*	*	*	*	728	247,000	11.6	(6.3 - 17.0)
Hispanic	*	*	*	*	854	460,600	20.5	(11.3 - 29.8)
Other Non-Hispanic	*	*	*	*	710	132,500	12.1	(3.4 - 20.7)
Education								
≤ High School	*	*	*	*	15,491	945,800	17.4	(13.2 - 21.6)
Some College	188	1,900	19.5	(10.1 - 28.9)	9,819	636,600	18.1	(14.6 - 21.5)
≥ College Degree	*	*	*	*	12,358	1,018,700	18.2	(15.3 - 21.1)
Income								
≤ \$24,999	*	*	*	*	9,557	469,100	13.5	(8.5 - 18.5)
\$25,000-\$49,999	186	1,700	21.0	(13.5 - 28.6)	9,575	534,300	18.5	(13.0 - 24.0)
\$50,000-\$74,999	*	*	*	*	5,726	378,200	19.2	(14.1 - 24.3)
≥ \$75,000	*	*	*	*	8,501	1,011,400	22.6	(19.7 - 25.6)

¹ Defined as men having 5 or more drinks or women having 4 or more drinks on 1 or more occasion within the past month

² Sample size from survey

³ Estimated number of adult binge drinkers within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 57b. Binge drinking within the past month among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	665	24.3	(17.6 - 31.1)	37,745	18.1	(16.1 - 20.2)
Gender						
Male	*	*	*	14,420	24.5	(21.0 - 27.9)
Female	423	16.8	(10.9 - 22.7)	23,325	12.4	(10.2 - 14.5)
Race/Ethnicity						
White Non-Hispanic	648	21.6	(16.0 - 27.1)	35,453	19.8	(18.1 - 21.6)
Black Non-Hispanic	*	*	*	728	11.8	(6.4 - 17.1)
Hispanic	*	*	*	854	20.2	(11.3 - 29.1)
Other Non-Hispanic	*	*	*	710	12.1	(3.4 - 20.8)
Education						
≤ High School	*	*	*	15,491	17.5	(13.5 - 21.5)
Some College	188	20.0	(10.7 - 29.3)	9,819	18.6	(15.1 - 22.0)
≥ College Degree	*	*	*	12,358	18.8	(15.9 - 21.7)
Income						
≤ \$24,999	*	*	*	9,557	13.4	(8.6 - 18.2)
\$25,000-\$49,999	186	21.1	(13.3 - 28.8)	9,575	18.9	(13.5 - 24.2)
\$50,000-\$74,999	*	*	*	5,726	19.9	(14.8 - 25.1)
≥ \$75,000	*	*	*	8,501	23.2	(20.2 - 26.1)

¹ Defined as men having 5 or more drinks or women having 4 or more drinks on 1 or more occasion within the past month

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 58a. Heavy drinking within the past month among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	665	2,300	7.1	(4.5 - 9.8)	37,553	718,700	4.9	(4.2 - 5.6)
Gender								
Male	243	1,600	8.8	(4.3 - 13.2)	14,330	333,300	4.8	(3.9 - 5.7)
Female	422	800	5.1	(2.6 - 7.6)	23,223	385,400	5.0	(4.0 - 6.1)
Age (years)								
18-34	*	*	*	*	3,866	224,900	5.1	(3.5 - 6.8)
35-44	*	*	*	*	4,924	137,200	4.8	(3.0 - 6.7)
45-54	151	600	8.5	(3.6 - 13.3)	7,715	144,600	5.2	(4.1 - 6.2)
55-64	168	300	6.5	(2.5 - 10.5)	8,624	117,400	5.6	(3.9 - 7.3)
≥ 65	*	*	*	*	12,424	94,600	3.8	(2.7 - 4.9)
Race/Ethnicity								
White Non-Hispanic	648	2,300	7.5	(4.7 - 10.3)	35,279	572,400	6.3	(5.4 - 7.1)
Black Non-Hispanic	*	*	*	*	723	58,100	2.8	(0.6 - 4.9)
Hispanic	*	*	*	*	848	60,900	2.7	(0.5 - 4.9)
Other Non-Hispanic	*	*	*	*	703	27,200	2.5	(0.8 - 4.2)
Education								
≤ High School	350	1,200	7.1	(3.9 - 10.4)	15,411	228,100	4.2	(3.2 - 5.2)
Some College	190	800	8.6	(2.8 - 14.4)	9,759	163,700	4.7	(3.6 - 5.8)
≥ College Degree	*	*	*	*	12,306	326,800	5.9	(4.5 - 7.3)
Income								
≤ \$24,999	*	*	*	*	9,512	104,200	3.0	(1.8 - 4.1)
\$25,000-\$49,999	184	700	8.5	(3.0 - 14.0)	9,537	158,700	5.5	(3.9 - 7.2)
\$50,000-\$74,999	*	*	*	*	5,708	95,400	4.9	(3.1 - 6.7)
≥ \$75,000	*	*	*	*	8,481	313,900	7.0	(5.4 - 8.6)

¹ Defined as adult men averaging more than 2 alcoholic drinks per day and adult women averaging more than 1 alcoholic drink per day within the past month

² Sample size from survey

³ Estimated number of adult heavy drinkers within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 58b. Heavy drinking within the past month among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	665	7.1	(4.4 - 9.7)	37,553	5.0	(4.3 - 5.8)
Gender						
Male	243	8.7	(4.3 - 13.0)	14,330	4.9	(3.9 - 5.9)
Female	422	5.1	(2.4 - 7.9)	23,223	5.1	(4.0 - 6.2)
Race/Ethnicity						
White Non-Hispanic	648	7.5	(4.7 - 10.4)	35,279	6.4	(5.5 - 7.2)
Black Non-Hispanic	*	*	*	723	2.9	(0.6 - 5.1)
Hispanic	*	*	*	848	2.8	(0.5 - 5.1)
Other Non-Hispanic	*	*	*	703	2.5	(0.8 - 4.2)
Education						
≤ High School	350	7.3	(3.9 - 10.6)	15,411	4.3	(3.2 - 5.3)
Some College	190	8.8	(2.8 - 14.8)	9,759	4.8	(3.6 - 5.9)
≥ College Degree	*	*	*	12,306	6.0	(4.5 - 7.4)
Income						
≤ \$24,999	*	*	*	9,512	3.0	(1.9 - 4.1)
\$25,000-\$49,999	184	8.4	(2.8 - 14.1)	9,537	5.8	(4.0 - 7.5)
\$50,000-\$74,999	*	*	*	5,708	5.0	(3.1 - 7.0)
≥ \$75,000	*	*	*	8,481	7.0	(5.4 - 8.7)

¹ Defined as adult men averaging more than 2 alcoholic drinks per day and adult women averaging more than 1 alcoholic drink per day within the past month

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wyoming County**

Table 59a. Never or rarely been asked about sexual history during routine check-up among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	620	23,400	78.1	(72.7 - 83.5)	35,193	8,480,600	61.9	(59.2 - 64.5)
Gender								
Male	225	13,900	85.6	(77.1 - 94.0)	13,299	4,357,400	67.7	(63.6 - 71.7)
Female	395	9,600	69.3	(63.1 - 75.5)	21,894	4,123,300	56.8	(53.3 - 60.2)
Age (years)								
18-34	*	*	*	*	3,507	1,804,600	44.3	(37.8 - 50.8)
35-44	78	4,200	77.4	(67.7 - 87.2)	4,531	1,635,300	62.2	(56.6 - 67.8)
45-54	142	5,100	83.2	(77.2 - 89.1)	7,215	1,715,600	65.4	(60.5 - 70.3)
55-64	162	3,900	83.7	(77.5 - 89.9)	8,168	1,416,200	70.7	(66.6 - 74.8)
≥ 65	184	4,400	87.0	(81.3 - 92.6)	11,772	1,908,900	80.3	(76.9 - 83.8)
Race/Ethnicity								
White Non-Hispanic	605	21,200	77.5	(72.0 - 82.9)	33,090	5,940,300	68.5	(66.3 - 70.7)
Black Non-Hispanic	*	*	*	*	683	831,800	40.8	(31.8 - 49.9)
Hispanic	*	*	*	*	784	1,071,700	53.4	(43.6 - 63.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	325	12,400	80.5	(72.4 - 88.5)	14,264	3,001,200	60.1	(55.4 - 64.9)
Some College	177	6,200	74.7	(65.1 - 84.3)	9,156	1,972,800	60.3	(54.9 - 65.7)
≥ College Degree	*	*	*	*	11,706	3,476,000	65.0	(61.2 - 68.7)
Income								
≤ \$24,999	*	*	*	*	8,772	1,529,400	49.3	(42.6 - 56.0)
\$25,000-\$49,999	173	6,000	80.4	(73.4 - 87.4)	8,932	1,705,000	64.4	(58.0 - 70.7)
\$50,000-\$74,999	*	*	*	*	5,433	1,217,000	64.5	(58.5 - 70.5)
≥ \$75,000	*	*	*	*	8,108	2,934,300	68.2	(64.7 - 71.7)

¹ Defined as respondents who never or rarely (hardly ever) were asked about their sexual history, such as about sexual partners and sexual practices, during a routine check-up at a doctor's office or clinic

² Sample size from survey

³ Estimated number of adults who were never or rarely asked about their sexual history during a routine check-up at a doctor's office or clinic (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 59b. Never or rarely been asked about sexual history during routine check-up among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	620	77.0	(71.5 - 82.5)	35,193	61.4	(58.7 - 64.0)
Gender						
Male	225	85.6	(77.5 - 93.6)	13,299	67.9	(64.0 - 71.9)
Female	395	66.9	(59.9 - 74.0)	21,894	55.5	(52.0 - 59.0)
Race/Ethnicity						
White Non-Hispanic	605	76.6	(71.1 - 82.1)	33,090	67.8	(65.6 - 70.1)
Black Non-Hispanic	*	*	*	683	41.1	(32.0 - 50.2)
Hispanic	*	*	*	784	53.0	(43.3 - 62.6)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	325	78.9	(70.7 - 87.0)	14,264	59.5	(54.7 - 64.3)
Some College	177	74.5	(64.9 - 84.0)	9,156	59.7	(54.4 - 65.0)
≥ College Degree	*	*	*	11,706	64.5	(60.8 - 68.2)
Income						
≤ \$24,999	*	*	*	8,772	48.7	(42.0 - 55.4)
\$25,000-\$49,999	173	79.7	(72.1 - 87.2)	8,932	63.7	(57.5 - 70.0)
\$50,000-\$74,999	*	*	*	5,433	64.2	(58.2 - 70.2)
≥ \$75,000	*	*	*	8,108	67.5	(64.0 - 71.0)

¹ Defined as respondents who never or rarely (hardly ever) were asked about their sexual history, such as about sexual partners and sexual practices, during a routine check-up at a doctor's office or clinic

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wyoming County**

Table 60a. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	392	17,400	83.1	(76.1 - 90.1)	22,788	7,011,100	77.9	(75.2 - 80.5)
Gender								
Male	*	*	*	*	9,034	3,618,500	81.0	(77.2 - 84.8)
Female	243	7,300	78.5	(70.1 - 86.9)	13,754	3,392,600	74.8	(71.2 - 78.4)
Age (years)								
18-34	*	*	*	*	3,213	2,362,500	70.8	(65.1 - 76.6)
35-44	*	*	*	*	3,513	1,313,400	71.7	(65.6 - 77.7)
45-54	105	4,300	91.4	(85.7 - 97.1)	4,975	1,290,600	80.3	(75.9 - 84.8)
55-64	90	2,500	96.9	(93.1 - 100)	4,955	932,200	87.2	(84.1 - 90.3)
≥ 65	94	2,400	99.1	(97.4 - 100)	6,132	1,112,400	95.9	(94.4 - 97.5)
Race/Ethnicity								
White Non-Hispanic	380	15,700	82.5	(75.2 - 89.9)	21,540	4,875,200	78.3	(75.6 - 80.9)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	491	1,002,400	84.8	(77.8 - 91.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	8,879	2,581,700	82.1	(77.9 - 86.3)
Some College	106	5,300	87.3	(77.3 - 97.3)	5,953	1,599,500	77.5	(72.6 - 82.3)
≥ College Degree	*	*	*	*	7,928	2,824,800	75.6	(71.8 - 79.5)
Income								
≤ \$24,999	112	5,000	93.1	(86.8 - 99.3)	5,491	1,602,900	78.7	(71.9 - 85.6)
\$25,000-\$49,999	*	*	*	*	5,912	1,335,900	78.8	(73.8 - 83.9)
\$50,000-\$74,999	*	*	*	*	3,717	987,500	76.6	(70.4 - 82.8)
≥ \$75,000	*	*	*	*	5,771	2,321,600	77.6	(74.0 - 81.2)

¹ Defined as respondent estimating there were hardly any (0-1 out of 10) or a few (2-3 out of 10) persons their age in their community who have had a sexually transmitted disease such as gonorrhea, chlamydia, herpes or syphilis

² Sample size from survey

³ Estimated number of adults who believed hardly any or a few people their age in their community have had a sexually transmitted disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 60b. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	392	82.0	(75.0 - 89.1)	22,788	77.3	(74.7 - 79.9)
Gender						
Male	*	*	*	9,034	81.1	(77.4 - 84.8)
Female	243	75.6	(66.2 - 85.1)	13,754	73.6	(70.0 - 77.3)
Race/Ethnicity						
White Non-Hispanic	380	81.6	(74.2 - 88.9)	21,540	77.5	(74.9 - 80.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	491	84.6	(77.6 - 91.5)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	8,879	81.6	(77.4 - 85.7)
Some College	*	*	*	5,953	76.8	(72.0 - 81.6)
≥ College Degree	*	*	*	7,928	75.1	(71.3 - 79.0)
Income						
≤ \$24,999	112	91.9	(84.7 - 99.2)	5,491	77.9	(71.0 - 84.7)
\$25,000-\$49,999	*	*	*	5,912	77.9	(72.8 - 83.0)
\$50,000-\$74,999	*	*	*	3,717	76.1	(70.0 - 82.3)
≥ \$75,000	*	*	*	5,771	77.0	(73.4 - 80.6)

¹ Defined as respondent estimating there were hardly any (0-1 out of 10) or a few (2-3 out of 10) persons their age in their community who have had a sexually transmitted disease such as gonorrhea, chlamydia, herpes or syphilis

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 61a. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	640	27,800	86.7	(83.0 - 90.4)	36,287	12,738,300	89.7	(88.2 - 91.3)
Gender								
Male	239	15,400	86.6	(80.7 - 92.6)	13,893	6,104,600	89.4	(87.2 - 91.5)
Female	401	12,400	86.8	(82.9 - 90.6)	22,394	6,633,700	90.1	(87.9 - 92.2)
Age (years)								
18-34	59	8,800	89.4	(80.1 - 98.6)	3,850	4,007,400	91.3	(87.5 - 95.1)
35-44	89	5,700	92.2	(86.6 - 97.8)	4,875	2,578,600	93.2	(90.4 - 96.0)
45-54	146	5,700	89.1	(82.1 - 96.0)	7,607	2,416,800	90.5	(87.9 - 93.2)
55-64	162	3,700	81.5	(74.7 - 88.3)	8,411	1,876,500	92.2	(90.0 - 94.5)
≥ 65	184	3,900	76.4	(69.4 - 83.3)	11,544	1,858,900	79.7	(76.6 - 82.8)
Race/Ethnicity								
White Non-Hispanic	623	25,300	86.3	(82.4 - 90.2)	34,102	8,050,700	89.4	(87.7 - 91.0)
Black Non-Hispanic	*	*	*	*	702	1,822,200	91.9	(88.2 - 95.6)
Hispanic	*	*	*	*	820	1,967,200	89.9	(85.2 - 94.7)
Other Non-Hispanic	*	*	*	*	663	898,200	88.6	(79.5 - 97.6)
Education								
≤ High School	334	13,800	85.2	(79.4 - 91.0)	14,507	4,385,300	84.6	(81.9 - 87.2)
Some College	185	7,900	85.5	(78.6 - 92.3)	9,571	3,094,300	90.0	(86.7 - 93.4)
≥ College Degree	121	6,000	92.1	(86.6 - 97.6)	12,142	5,228,800	95.4	(94.0 - 96.7)
Income								
≤ \$24,999	166	5,800	83.1	(75.2 - 91.0)	9,021	2,833,500	86.5	(82.6 - 90.3)
\$25,000-\$49,999	179	6,700	84.2	(77.6 - 90.8)	9,261	2,529,900	89.5	(86.5 - 92.5)
\$50,000-\$74,999	114	6,300	90.4	(81.2 - 99.6)	5,644	1,740,000	91.2	(87.4 - 95.1)
≥ \$75,000	106	5,700	93.4	(88.3 - 98.5)	8,397	4,184,300	94.4	(92.5 - 96.3)

¹ Defined as believing it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community

² Sample size from survey

³ Estimated number of adults who believed it is at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 61b. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	640	87.0	(83.4 - 90.6)	36,287	89.8	(88.3 - 91.4)
Gender						
Male	239	86.4	(80.6 - 92.1)	13,893	89.1	(86.9 - 91.3)
Female	401	87.8	(83.9 - 91.7)	22,394	90.5	(88.3 - 92.7)
Race/Ethnicity						
White Non-Hispanic	623	86.6	(82.8 - 90.4)	34,102	89.5	(87.8 - 91.2)
Black Non-Hispanic	*	*	*	702	91.7	(87.8 - 95.6)
Hispanic	*	*	*	820	89.6	(84.7 - 94.5)
Other Non-Hispanic	*	*	*	663	89.0	(80.5 - 97.6)
Education						
≤ High School	334	85.8	(80.1 - 91.4)	14,507	84.6	(81.8 - 87.4)
Some College	185	85.6	(78.9 - 92.3)	9,571	90.2	(86.9 - 93.5)
≥ College Degree	121	91.8	(85.9 - 97.8)	12,142	95.4	(94.0 - 96.7)
Income						
≤ \$24,999	166	83.6	(75.8 - 91.3)	9,021	86.5	(82.6 - 90.4)
\$25,000-\$49,999	179	85.0	(78.6 - 91.4)	9,261	89.6	(86.5 - 92.6)
\$50,000-\$74,999	114	90.7	(81.9 - 99.5)	5,644	91.1	(86.9 - 95.3)
≥ \$75,000	106	92.9	(87.1 - 98.7)	8,397	94.5	(92.6 - 96.4)

¹ Defined as believing it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 62a. Overweight among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	644	11,400	35.5	(29.7 - 41.2)	36,710	5,213,000	36.4	(34.0 - 38.8)
Gender								
Male	245	7,400	40.7	(31.3 - 50.1)	14,502	3,073,200	43.7	(39.9 - 47.6)
Female	399	4,100	28.8	(22.9 - 34.8)	22,208	2,139,800	29.3	(26.4 - 32.2)
Age (years)								
18-34	*	*	*	*	3,789	1,295,200	30.1	(24.7 - 35.6)
35-44	*	*	*	*	4,785	1,052,000	37.5	(31.7 - 43.3)
45-54	151	2,900	43.2	(33.7 - 52.7)	7,495	1,053,300	38.8	(34.1 - 43.4)
55-64	157	1,800	38.8	(30.1 - 47.5)	8,356	798,100	38.8	(34.8 - 42.7)
≥ 65	188	1,900	37.7	(29.8 - 45.5)	12,285	1,014,400	41.5	(37.6 - 45.4)
Race/Ethnicity								
White Non-Hispanic	628	11,200	37.2	(31.6 - 42.9)	34,501	3,282,200	36.4	(34.4 - 38.3)
Black Non-Hispanic	*	*	*	*	708	740,000	36.7	(28.3 - 45.1)
Hispanic	*	*	*	*	813	781,500	35.6	(27.1 - 44.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	340	5,300	33.0	(24.5 - 41.6)	15,079	1,889,700	35.4	(31.2 - 39.5)
Some College	182	3,400	35.4	(25.5 - 45.4)	9,515	1,314,500	38.7	(33.8 - 43.7)
≥ College Degree	*	*	*	*	12,062	1,996,900	36.3	(32.8 - 39.7)
Income								
≤ \$24,999	*	*	*	*	9,346	1,170,900	34.0	(28.2 - 39.8)
\$25,000-\$49,999	182	3,400	42.4	(33.0 - 51.8)	9,368	1,136,000	40.4	(34.6 - 46.2)
\$50,000-\$74,999	*	*	*	*	5,615	738,700	38.3	(32.3 - 44.3)
≥ \$75,000	*	*	*	*	8,337	1,659,100	37.5	(34.1 - 40.9)

¹ Defined as having a body mass index (BMI) more than 24.9 but less than 30.0. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Sample size from survey

³ Estimated number of adults who are overweight (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 62b. Overweight among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	644	35.5	(29.7 - 41.2)	36,710	36.2	(33.8 - 38.6)
Gender						
Male	245	40.7	(31.7 - 49.7)	14,502	43.9	(40.1 - 47.6)
Female	399	28.8	(22.2 - 35.4)	22,208	28.9	(25.9 - 31.8)
Race/Ethnicity						
White Non-Hispanic	628	37.1	(31.5 - 42.8)	34,501	36.0	(34.0 - 38.0)
Black Non-Hispanic	*	*	*	708	37.0	(28.5 - 45.5)
Hispanic	*	*	*	813	36.0	(27.5 - 44.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	340	33.3	(24.9 - 41.6)	15,079	35.2	(31.1 - 39.4)
Some College	182	35.4	(25.5 - 45.3)	9,515	38.3	(33.3 - 43.2)
≥ College Degree	*	*	*	12,062	36.2	(32.7 - 39.8)
Income						
≤ \$24,999	*	*	*	9,346	33.8	(28.0 - 39.5)
\$25,000-\$49,999	182	42.2	(32.7 - 51.7)	9,368	40.6	(34.8 - 46.4)
\$50,000-\$74,999	*	*	*	5,615	38.6	(32.5 - 44.7)
≥ \$75,000	*	*	*	8,337	37.2	(33.7 - 40.6)

¹ Defined as having a body mass index (BMI) more than 24.9 but less than 30.0. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wyoming County**

Table 63a. Obesity among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	644	10,300	32.0	(26.5 - 37.5)	36,710	3,327,400	23.2	(21.2 - 25.3)
Gender								
Male	245	5,700	31.4	(22.8 - 40.0)	14,502	1,505,800	21.4	(19.0 - 23.9)
Female	399	4,600	32.7	(26.9 - 38.5)	22,208	1,821,600	25.0	(21.8 - 28.1)
Age (years)								
18-34	*	*	*	*	3,789	730,600	17.0	(12.4 - 21.6)
35-44	*	*	*	*	4,785	662,200	23.6	(19.4 - 27.9)
45-54	151	2,300	34.6	(25.8 - 43.5)	7,495	816,800	30.1	(25.3 - 34.8)
55-64	157	1,600	35.7	(27.5 - 44.0)	8,356	563,900	27.4	(23.7 - 31.1)
≥ 65	188	1,500	29.8	(22.4 - 37.3)	12,285	554,000	22.7	(19.2 - 26.1)
Race/Ethnicity								
White Non-Hispanic	628	9,900	33.0	(27.6 - 38.3)	34,501	2,062,700	22.9	(21.2 - 24.6)
Black Non-Hispanic	*	*	*	*	708	703,100	34.9	(26.1 - 43.7)
Hispanic	*	*	*	*	813	456,400	20.8	(14.3 - 27.2)
Other Non-Hispanic	*	*	*	*	688	105,200	9.6	(4.0 - 15.3)
Education								
≤ High School	340	5,400	33.7	(25.4 - 41.9)	15,079	1,561,600	29.2	(25.3 - 33.2)
Some College	*	*	*	*	9,515	792,800	23.4	(19.4 - 27.3)
≥ College Degree	122	1,700	25.9	(16.1 - 35.6)	12,062	915,000	16.6	(14.4 - 18.8)
Income								
≤ \$24,999	*	*	*	*	9,346	974,800	28.3	(22.6 - 34.0)
\$25,000-\$49,999	182	2,700	33.7	(24.8 - 42.7)	9,368	584,800	20.8	(17.1 - 24.5)
\$50,000-\$74,999	*	*	*	*	5,615	475,300	24.6	(20.0 - 29.3)
≥ \$75,000	*	*	*	*	8,337	867,700	19.6	(17.0 - 22.3)

¹ Defined as having a body mass index (BMI) of 30.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Sample size from survey

³ Estimated number of adults who are obese (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 63b. Obesity among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	644	31.6	(26.2 - 37.0)	36,710	23.1	(21.0 - 25.1)
Gender						
Male	245	31.3	(23.0 - 39.6)	14,502	21.5	(19.0 - 24.0)
Female	399	31.9	(25.6 - 38.3)	22,208	24.6	(21.5 - 27.8)
Race/Ethnicity						
White Non-Hispanic	628	32.4	(27.1 - 37.8)	34,501	22.8	(21.0 - 24.5)
Black Non-Hispanic	*	*	*	708	34.8	(25.9 - 43.6)
Hispanic	*	*	*	813	20.5	(14.1 - 26.9)
Other Non-Hispanic	*	*	*	688	9.5	(3.8 - 15.2)
Education						
≤ High School	340	33.2	(25.1 - 41.3)	15,079	29.4	(25.4 - 33.3)
Some College	*	*	*	9,515	23.1	(19.2 - 27.1)
≥ College Degree	122	24.8	(15.1 - 34.6)	12,062	16.4	(14.2 - 18.5)
Income						
≤ \$24,999	*	*	*	9,346	28.3	(22.6 - 34.1)
\$25,000-\$49,999	182	33.1	(24.1 - 42.0)	9,368	21.0	(17.2 - 24.8)
\$50,000-\$74,999	*	*	*	5,615	24.0	(19.4 - 28.7)
≥ \$75,000	*	*	*	8,337	19.3	(16.7 - 21.9)

¹ Defined as having a body mass index (BMI) of 30.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 64a. Overweight or obesity among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	644	21,800	67.4	(60.9 - 74.0)	36,710	8,540,500	59.6	(57.0 - 62.2)
Gender								
Male	*	*	*	*	14,502	4,579,100	65.2	(61.0 - 69.4)
Female	399	8,700	61.5	(55.1 - 67.9)	22,208	3,961,400	54.3	(51.0 - 57.6)
Age (years)								
18-34	*	*	*	*	3,789	2,025,800	47.1	(40.8 - 53.5)
35-44	*	*	*	*	4,785	1,714,200	61.1	(55.6 - 66.6)
45-54	151	5,200	77.8	(70.9 - 84.8)	7,495	1,870,100	68.8	(64.6 - 73.1)
55-64	157	3,400	74.5	(67.2 - 81.9)	8,356	1,362,000	66.2	(61.9 - 70.4)
≥ 65	188	3,500	67.5	(59.8 - 75.2)	12,285	1,568,400	64.1	(60.6 - 67.7)
Race/Ethnicity								
White Non-Hispanic	628	21,100	70.2	(64.9 - 75.5)	34,501	5,344,900	59.3	(57.2 - 61.3)
Black Non-Hispanic	*	*	*	*	708	1,443,100	71.6	(62.5 - 80.7)
Hispanic	*	*	*	*	813	1,237,900	56.4	(46.5 - 66.2)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,079	3,451,400	64.6	(59.8 - 69.4)
Some College	*	*	*	*	9,515	2,107,300	62.1	(57.0 - 67.2)
≥ College Degree	*	*	*	*	12,062	2,911,900	52.9	(49.2 - 56.6)
Income								
≤ \$24,999	*	*	*	*	9,346	2,145,700	62.3	(55.7 - 68.9)
\$25,000-\$49,999	182	6,200	76.1	(68.0 - 84.3)	9,368	1,720,700	61.2	(54.5 - 67.9)
\$50,000-\$74,999	*	*	*	*	5,615	1,214,000	62.9	(57.1 - 68.8)
≥ \$75,000	*	*	*	*	8,337	2,526,700	57.1	(53.6 - 60.6)

¹ Defined as having a body mass index (BMI) of 25.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Sample size from survey

³ Estimated number of adults who are overweight or obese (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 64b. Overweight or obesity among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	644	67.0	(60.6 - 73.5)	36,710	59.3	(56.8 - 61.9)
Gender						
Male	*	*	*	14,502	65.4	(61.3 - 69.4)
Female	399	60.7	(53.5 - 67.9)	22,208	53.5	(50.1 - 56.8)
Race/Ethnicity						
White Non-Hispanic	628	69.6	(64.1 - 75.0)	34,501	58.8	(56.7 - 60.8)
Black Non-Hispanic	*	*	*	708	71.8	(63.0 - 80.6)
Hispanic	*	*	*	813	56.5	(46.8 - 66.1)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	15,079	64.6	(59.9 - 69.3)
Some College	*	*	*	9,515	61.4	(56.4 - 66.5)
≥ College Degree	*	*	*	12,062	52.6	(48.9 - 56.3)
Income						
≤ \$24,999	*	*	*	9,346	62.1	(55.6 - 68.6)
\$25,000-\$49,999	182	75.2	(66.4 - 84.1)	9,368	61.6	(55.1 - 68.1)
\$50,000-\$74,999	*	*	*	5,615	62.6	(56.7 - 68.6)
≥ \$75,000	*	*	*	8,337	56.5	(52.9 - 60.0)

¹ Defined as having a body mass index (BMI) of 25.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 65a. Received advice about weight by a health professional among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	643	7,400	22.8	(18.6 - 27.1)	36,729	3,929,700	27.4	(25.1 - 29.7)
Gender								
Male	245	3,500	19.1	(13.2 - 25.0)	14,499	1,654,800	23.6	(20.6 - 26.6)
Female	398	3,900	27.6	(22.2 - 33.1)	22,230	2,274,900	31.0	(27.7 - 34.3)
Age (years)								
18-34	57	1,200	12.7	(4.4 - 21.1)	3,796	895,500	20.7	(15.3 - 26.2)
35-44	90	1,300	19.9	(10.9 - 28.8)	4,779	722,700	25.8	(20.9 - 30.6)
45-54	151	2,000	30.6	(22.2 - 39.0)	7,490	910,900	33.5	(29.0 - 38.1)
55-64	158	1,400	30.1	(22.0 - 38.2)	8,351	741,100	36.1	(32.0 - 40.3)
≥ 65	187	1,500	28.9	(21.4 - 36.3)	12,313	659,500	26.9	(23.0 - 30.7)
Race/Ethnicity								
White Non-Hispanic	627	7,000	23.4	(19.2 - 27.5)	34,502	2,386,200	26.5	(24.6 - 28.4)
Black Non-Hispanic	*	*	*	*	709	615,700	30.4	(22.0 - 38.8)
Hispanic	*	*	*	*	830	555,400	25.1	(17.8 - 32.4)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	339	3,800	23.3	(16.9 - 29.8)	15,116	1,469,700	27.5	(23.6 - 31.4)
Some College	182	2,100	21.7	(14.7 - 28.6)	9,515	946,800	27.8	(23.2 - 32.4)
≥ College Degree	122	1,500	23.3	(14.5 - 32.1)	12,041	1,445,500	26.2	(23.0 - 29.4)
Income								
≤ \$24,999	171	1,300	18.0	(9.9 - 26.1)	9,360	1,079,400	31.4	(25.3 - 37.5)
\$25,000-\$49,999	182	2,100	25.9	(17.8 - 34.0)	9,365	696,900	24.6	(19.8 - 29.4)
\$50,000-\$74,999	113	1,500	20.9	(11.9 - 29.8)	5,603	507,100	26.3	(21.6 - 31.0)
≥ \$75,000	107	1,500	24.1	(14.7 - 33.4)	8,334	1,207,100	27.3	(24.2 - 30.5)

¹ Defined as being advised by a doctor, nurse or other health professional about their weight (lose, gain or maintain)

² Sample size from survey

³ Estimated number of adults provided advice by a health professional about their weight (lose, gain or maintain) (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 65b. Received advice about weight by a health professional among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	643	22.7	(18.4 - 26.9)	36,729	27.1	(24.8 - 29.4)
Gender						
Male	245	19.5	(13.6 - 25.3)	14,499	23.4	(20.5 - 26.3)
Female	398	26.7	(20.9 - 32.6)	22,230	30.6	(27.3 - 34.0)
Race/Ethnicity						
White Non-Hispanic	627	23.1	(18.9 - 27.3)	34,502	26.1	(24.2 - 28.0)
Black Non-Hispanic	*	*	*	709	30.4	(21.9 - 38.9)
Hispanic	*	*	*	830	24.9	(17.7 - 32.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	339	23.6	(17.0 - 30.1)	15,116	27.3	(23.4 - 31.2)
Some College	182	21.9	(14.7 - 29.0)	9,515	27.4	(22.9 - 32.0)
≥ College Degree	122	21.6	(13.0 - 30.3)	12,041	25.9	(22.7 - 29.0)
Income						
≤ \$24,999	171	18.6	(10.2 - 26.9)	9,360	31.0	(24.9 - 37.0)
\$25,000-\$49,999	182	25.7	(17.6 - 33.9)	9,365	24.9	(20.0 - 29.8)
\$50,000-\$74,999	113	20.1	(11.4 - 28.9)	5,603	25.9	(21.2 - 30.6)
≥ \$75,000	107	22.9	(13.7 - 32.1)	8,334	26.9	(23.7 - 30.0)

¹ Defined as being advised by a doctor, nurse or other health professional about their weight (lose, gain or maintain)

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 66a. Received advice to lose weight by a health professional among adults provided weight advice¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	175	6,500	88.8	(82.6 - 95.0)	9,724	3,412,900	86.8	(83.9 - 89.8)
Gender								
Male	61	3,200	91.4	(83.6 - 99.1)	3,807	1,468,700	88.8	(85.7 - 91.8)
Female	114	3,400	86.5	(77.2 - 95.8)	5,917	1,944,200	85.5	(80.8 - 90.1)
Age (years)								
18-34	*	*	*	*	681	779,700	87.1	(80.6 - 93.6)
35-44	*	*	*	*	1,215	622,100	86.1	(76.9 - 95.2)
45-54	*	*	*	*	2,238	813,300	89.3	(84.4 - 94.2)
55-64	*	*	*	*	2,697	645,500	87.1	(81.2 - 93.0)
≥ 65	51	1,300	88.2	(78.6 - 97.8)	2,893	552,300	83.7	(76.9 - 90.6)
Race/Ethnicity								
White Non-Hispanic	171	6,200	88.2	(81.7 - 94.7)	9,082	2,076,700	87.0	(84.5 - 89.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	251	483,600	87.1	(78.5 - 95.7)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	3,962	1,239,700	84.3	(78.6 - 90.1)
Some College	53	2,000	98.6	(95.8 - 100)	2,633	814,800	86.1	(80.1 - 92.0)
≥ College Degree	*	*	*	*	3,115	1,290,900	89.3	(85.3 - 93.3)
Income								
≤ \$24,999	*	*	*	*	2,539	922,100	85.4	(78.7 - 92.1)
\$25,000-\$49,999	*	*	*	*	2,482	594,100	85.2	(78.4 - 92.1)
\$50,000-\$74,999	*	*	*	*	1,521	446,100	88.0	(83.2 - 92.7)
≥ \$75,000	*	*	*	*	2,270	1,073,200	88.9	(83.7 - 94.1)

¹ Defined as provided advice by a health professional about their weight and advised to lose weight

² Sample size from survey

³ Estimated number of adults provided advice by a health professional about their weight and advised to lose weight (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 66b. Received advice to lose weight by a health professional among adults provided weight advice¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	175	88.4	(81.7 - 95.1)	9,724	86.9	(83.8 - 89.9)
Gender						
Male	61	91.8	(84.6 - 99.1)	3,807	88.7	(85.7 - 91.8)
Female	*	*	*	5,917	85.5	(80.7 - 90.3)
Race/Ethnicity						
White Non-Hispanic	171	87.8	(80.7 - 94.8)	9,082	87.0	(84.4 - 89.6)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	251	87.6	(79.3 - 95.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	3,962	84.6	(78.7 - 90.5)
Some College	53	98.8	(96.5 - 100)	2,633	85.6	(79.5 - 91.7)
≥ College Degree	*	*	*	3,115	89.4	(85.4 - 93.3)
Income						
≤ \$24,999	*	*	*	2,539	85.7	(79.0 - 92.3)
\$25,000-\$49,999	*	*	*	2,482	85.3	(78.4 - 92.3)
\$50,000-\$74,999	*	*	*	1,521	88.0	(83.1 - 92.8)
≥ \$75,000	*	*	*	2,270	88.7	(83.1 - 94.2)

¹ Defined as provided advice by a health professional about their weight and advised to lose weight

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 67a. No leisure-time physical activity among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	673	8,800	26.3	(20.0 - 32.5)	38,128	3,541,700	23.9	(21.6 - 26.1)
Gender								
Male	*	*	*	*	14,602	1,482,500	20.9	(17.8 - 24.0)
Female	425	4,200	28.0	(22.8 - 33.3)	23,526	2,059,200	26.6	(23.4 - 29.7)
Age (years)								
18-34	*	*	*	*	3,938	982,900	22.0	(16.6 - 27.5)
35-44	*	*	*	*	4,988	556,000	19.3	(15.0 - 23.6)
45-54	155	1,700	25.1	(17.7 - 32.6)	7,827	759,000	26.8	(21.7 - 31.9)
55-64	169	1,300	26.7	(19.2 - 34.2)	8,765	464,900	21.7	(18.2 - 25.2)
≥ 65	198	1,800	32.5	(25.1 - 39.8)	12,610	778,800	30.7	(27.3 - 34.2)
Race/Ethnicity								
White Non-Hispanic	655	6,900	22.6	(18.4 - 26.7)	35,816	1,892,800	20.3	(18.6 - 22.0)
Black Non-Hispanic	*	*	*	*	739	690,000	32.0	(23.4 - 40.6)
Hispanic	*	*	*	*	860	600,800	26.7	(19.1 - 34.4)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,683	1,810,200	32.5	(28.4 - 36.5)
Some College	194	1,700	17.6	(10.6 - 24.5)	9,906	814,200	23.0	(18.6 - 27.3)
≥ College Degree	125	800	11.9	(5.6 - 18.2)	12,458	841,000	14.9	(11.9 - 18.0)
Income								
≤ \$24,999	*	*	*	*	9,640	1,332,200	38.0	(31.7 - 44.3)
\$25,000-\$49,999	187	2,200	26.4	(18.2 - 34.5)	9,658	648,000	22.3	(18.3 - 26.3)
\$50,000-\$74,999	117	900	12.3	(5.3 - 19.3)	5,773	430,700	21.8	(16.6 - 27.0)
≥ \$75,000	111	700	10.7	(3.5 - 18.0)	8,565	560,800	12.5	(10.3 - 14.6)

¹ Defined as having not participated in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise

² Sample size from survey

³ Estimated number of adults who have not participated in any physical activities or exercises (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 67b. No leisure-time physical activity among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	673	25.4	(19.4 - 31.4)	38,128	23.7	(21.5 - 25.9)
Gender						
Male	248	24.3	(14.5 - 34.2)	14,602	20.9	(17.8 - 23.9)
Female	425	26.7	(21.1 - 32.3)	23,526	26.3	(23.1 - 29.5)
Race/Ethnicity						
White Non-Hispanic	655	22.0	(17.8 - 26.1)	35,816	20.0	(18.3 - 21.7)
Black Non-Hispanic	*	*	*	739	32.0	(23.3 - 40.6)
Hispanic	*	*	*	860	26.9	(19.2 - 34.5)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	354	36.1	(26.2 - 46.0)	15,683	32.3	(28.3 - 36.4)
Some College	194	16.8	(9.9 - 23.8)	9,906	22.8	(18.5 - 27.2)
≥ College Degree	125	11.7	(5.0 - 18.5)	12,458	14.9	(11.8 - 17.9)
Income						
≤ \$24,999	*	*	*	9,640	37.9	(31.6 - 44.2)
\$25,000-\$49,999	187	24.8	(16.6 - 32.9)	9,658	22.0	(18.0 - 26.0)
\$50,000-\$74,999	117	11.2	(4.7 - 17.7)	5,773	21.6	(16.3 - 26.8)
≥ \$75,000	111	10.7	(3.4 - 17.9)	8,565	12.6	(10.3 - 14.9)

¹ Defined as having not participated in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 68a. Consumption of 5 or more servings of fruits and vegetables per day among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	674	6,600	19.8	(15.9 - 23.7)	38,203	4,023,400	27.0	(24.9 - 29.2)
Gender								
Male	248	2,800	14.8	(9.3 - 20.4)	14,634	1,603,900	22.6	(19.4 - 25.8)
Female	426	3,900	26.0	(21.1 - 31.0)	23,569	2,419,500	31.1	(28.2 - 34.0)
Age (years)								
18-34	60	1,000	9.6	(2.3 - 16.9)	3,940	1,261,700	28.3	(22.7 - 33.9)
35-44	92	1,500	23.4	(13.8 - 33.0)	4,989	688,000	23.9	(19.5 - 28.3)
45-54	155	1,500	21.7	(14.0 - 29.4)	7,837	746,800	26.2	(22.3 - 30.1)
55-64	169	1,000	21.3	(14.3 - 28.2)	8,777	579,100	27.0	(23.6 - 30.5)
≥ 65	198	1,700	31.1	(24.0 - 38.2)	12,660	747,800	29.4	(26.0 - 32.8)
Race/Ethnicity								
White Non-Hispanic	656	6,400	21.0	(17.0 - 24.9)	35,884	2,538,000	27.2	(25.4 - 28.9)
Black Non-Hispanic	*	*	*	*	740	572,400	26.5	(19.2 - 33.9)
Hispanic	*	*	*	*	864	672,200	29.8	(21.0 - 38.5)
Other Non-Hispanic	*	*	*	*	715	240,800	21.4	(12.0 - 30.8)
Education								
≤ High School	355	2,700	15.7	(11.0 - 20.3)	15,726	1,309,800	23.4	(19.3 - 27.4)
Some College	194	2,000	20.3	(12.6 - 27.9)	9,924	964,300	27.2	(23.0 - 31.3)
≥ College Degree	125	2,000	29.7	(19.9 - 39.5)	12,471	1,747,500	31.0	(27.9 - 34.1)
Income								
≤ \$24,999	180	1,100	14.3	(8.2 - 20.4)	9,667	831,500	23.7	(18.6 - 28.7)
\$25,000-\$49,999	187	1,900	23.0	(15.7 - 30.3)	9,669	917,200	31.6	(25.6 - 37.6)
\$50,000-\$74,999	117	1,500	21.6	(12.6 - 30.6)	5,777	501,000	25.3	(21.4 - 29.3)
≥ \$75,000	111	1,300	19.8	(11.0 - 28.5)	8,569	1,260,100	28.0	(25.0 - 31.0)

¹ Defined as having consumed 5 or more servings of fruits and vegetables in an average day

² Sample size from survey

³ Estimated number of adults who consume 5 or more servings of fruits and vegetables per day (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 68b. Consumption of 5 or more servings of fruits and vegetables per day among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	674	19.7	(15.7 - 23.6)	38,203	27.1	(24.9 - 29.2)
Gender						
Male	248	15.4	(9.9 - 21.0)	14,634	22.7	(19.6 - 25.9)
Female	426	24.9	(19.7 - 30.2)	23,569	31.0	(28.1 - 34.0)
Race/Ethnicity						
White Non-Hispanic	656	20.8	(16.8 - 24.8)	35,884	27.1	(25.4 - 28.9)
Black Non-Hispanic	*	*	*	740	26.8	(19.4 - 34.2)
Hispanic	*	*	*	864	29.5	(20.9 - 38.1)
Other Non-Hispanic	*	*	*	715	21.8	(12.2 - 31.5)
Education						
≤ High School	355	15.9	(11.1 - 20.7)	15,726	23.3	(19.3 - 27.3)
Some College	194	19.7	(12.1 - 27.3)	9,924	27.1	(22.9 - 31.2)
≥ College Degree	125	28.8	(18.9 - 38.7)	12,471	31.2	(28.0 - 34.3)
Income						
≤ \$24,999	180	12.8	(7.3 - 18.3)	9,667	23.9	(18.8 - 28.9)
\$25,000-\$49,999	187	23.4	(15.8 - 31.1)	9,669	32.0	(26.0 - 38.0)
\$50,000-\$74,999	117	22.8	(13.3 - 32.3)	5,777	25.0	(21.0 - 28.9)
≥ \$75,000	111	19.7	(10.8 - 28.6)	8,569	28.0	(24.9 - 31.1)

¹ Defined as having consumed 5 or more servings of fruits and vegetables in an average day

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wyoming County**

Table 69a. Had a fall within the past 3 months among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	515	3,000	18.2	(14.5 - 21.8)	28,783	1,244,800	17.0	(14.9 - 19.0)
Gender								
Male	187	1,400	16.2	(10.5 - 21.9)	10,966	489,400	14.4	(11.9 - 16.8)
Female	328	1,700	20.1	(15.5 - 24.7)	17,817	755,300	19.2	(16.1 - 22.3)
Age (years)								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	153	1,100	16.8	(10.6 - 23.1)	7,741	445,500	16.0	(12.0 - 20.0)
55-64	165	800	17.6	(11.1 - 24.1)	8,454	336,000	16.5	(13.4 - 19.6)
≥ 65	197	1,100	20.3	(14.0 - 26.5)	12,588	463,300	18.4	(15.2 - 21.6)
Race/Ethnicity								
White Non-Hispanic	506	2,900	17.9	(14.2 - 21.6)	27,456	869,000	16.6	(15.1 - 18.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	424	76,100	11.0	(4.5 - 17.4)
Other Non-Hispanic	*	*	*	*	445	44,300	10.0	(3.1 - 16.8)
Education								
≤ High School	284	1,800	19.7	(14.4 - 25.0)	12,417	493,000	17.3	(13.3 - 21.3)
Some College	144	800	16.2	(9.7 - 22.6)	7,232	299,400	18.7	(14.6 - 22.7)
≥ College Degree	87	500	16.7	(8.7 - 24.7)	9,086	441,500	15.4	(12.9 - 17.9)
Income								
≤ \$24,999	150	700	15.7	(9.2 - 22.2)	7,824	423,900	25.0	(18.5 - 31.6)
\$25,000-\$49,999	141	900	20.6	(13.1 - 28.2)	7,445	223,600	16.0	(12.8 - 19.2)
\$50,000-\$74,999	84	600	19.1	(9.6 - 28.6)	4,141	151,200	14.6	(10.8 - 18.4)
≥ \$75,000	80	500	15.8	(7.6 - 24.0)	5,874	287,100	12.5	(10.0 - 15.1)

¹ Defined as respondent unintentionally coming to rest on the ground or another lower level one or more times within the past 3 months

² Sample size from survey

³ Estimated number of adults who have had a fall within the past 3 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 69b. Had a fall within the past 3 months among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	515	18.1	(14.2 - 21.9)	28,783	17.3	(15.0 - 19.5)
Gender						
Male	187	16.4	(10.7 - 22.2)	10,966	15.0	(12.1 - 17.8)
Female	328	20.1	(15.4 - 24.7)	17,817	19.4	(16.0 - 22.8)
Race/Ethnicity						
White Non-Hispanic	506	17.8	(14.0 - 21.7)	27,456	16.9	(15.2 - 18.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	424	10.7	(4.2 - 17.1)
Other Non-Hispanic	*	*	*	445	10.0	(2.9 - 17.0)
Education						
≤ High School	284	19.9	(14.4 - 25.4)	12,417	17.4	(13.0 - 21.7)
Some College	144	16.0	(9.3 - 22.8)	7,232	19.0	(14.7 - 23.2)
≥ College Degree	87	15.8	(8.0 - 23.7)	9,086	16.0	(13.1 - 18.8)
Income						
≤ \$24,999	150	15.4	(8.6 - 22.2)	7,824	25.3	(18.3 - 32.4)
\$25,000-\$49,999	141	19.9	(12.4 - 27.3)	7,445	15.9	(12.6 - 19.2)
\$50,000-\$74,999	84	19.0	(9.1 - 28.9)	4,141	14.7	(10.6 - 18.8)
≥ \$75,000	80	15.9	(7.4 - 24.4)	5,874	12.9	(9.9 - 15.8)

¹ Defined as respondent unintentionally coming to rest on the ground or another lower level one or more times within the past 3 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wyoming County

Table 70a. Had a fall that caused injury within the past 3 months among adults¹ in Wyoming County and New York State

	Wyoming County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	515	800	4.8	(3.0 - 6.7)	28,760	335,700	4.6	(3.7 - 5.5)
Gender								
Male	*	*	*	*	10,957	120,000	3.5	(2.5 - 4.6)
Female	328	600	6.7	(3.8 - 9.6)	17,803	215,600	5.5	(4.1 - 6.9)
Age (years)								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	153	400	5.8	(2.5 - 9.1)	7,737	115,100	4.1	(2.9 - 5.4)
55-64	*	*	*	*	8,449	90,300	4.4	(2.7 - 6.1)
≥ 65	197	200	3.8	(1.2 - 6.4)	12,574	130,300	5.2	(3.4 - 6.9)
Race/Ethnicity								
White Non-Hispanic	506	800	4.7	(2.9 - 6.6)	27,433	270,800	5.2	(4.2 - 6.1)
Black Non-Hispanic	*	*	*	*	458	47,900	4.9	(0.9 - 9.0)
Hispanic	*	*	*	*	424	12,400	1.8	(0.2 - 3.3)
Other Non-Hispanic	*	*	*	*	445	4,600	1.0	(0.4 - 1.6)
Education								
≤ High School	284	400	4.9	(2.3 - 7.6)	12,404	148,900	5.2	(3.6 - 6.8)
Some College	144	200	5.2	(1.7 - 8.6)	7,228	85,400	5.3	(3.1 - 7.5)
≥ College Degree	*	*	*	*	9,081	101,300	3.5	(2.4 - 4.6)
Income								
≤ \$24,999	150	300	6.6	(2.6 - 10.7)	7,812	90,200	5.3	(3.6 - 7.1)
\$25,000-\$49,999	141	300	7.8	(2.9 - 12.7)	7,442	93,200	6.7	(3.9 - 9.4)
\$50,000-\$74,999	*	*	*	*	4,139	37,000	3.6	(1.8 - 5.3)
≥ \$75,000	*	*	*	*	5,872	54,700	2.4	(1.5 - 3.3)

¹ Defined as as respondent unintentionally coming to rest on the ground or another lower level and caused respondent to limit regular activities for at least a day or to go see a doctor one or more times within the past 3 months

² Sample size from survey

³ Estimated number of adults who have had a fall that caused an injury within the past 3 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wyoming County**

Table 70b. Had a fall that caused injury within the past 3 months among adults¹ in Wyoming County and New York State - Age-Adjusted²

	Wyoming County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	515	4.5	(2.8 - 6.3)	28,760	4.6	(3.6 - 5.5)
Gender						
Male	*	*	*	10,957	3.6	(2.4 - 4.7)
Female	328	6.7	(3.8 - 9.6)	17,803	5.5	(4.1 - 6.9)
Race/Ethnicity						
White Non-Hispanic	506	4.4	(2.7 - 6.1)	27,433	5.2	(4.2 - 6.2)
Black Non-Hispanic	*	*	*	458	4.7	(0.7 - 8.7)
Hispanic	*	*	*	424	1.8	(0.1 - 3.4)
Other Non-Hispanic	*	*	*	445	1.0	(0.4 - 1.6)
Education						
≤ High School	284	4.5	(2.0 - 6.9)	12,404	5.2	(3.5 - 6.8)
Some College	144	4.8	(1.6 - 8.1)	7,228	5.3	(3.1 - 7.5)
≥ College Degree	*	*	*	9,081	3.6	(2.4 - 4.7)
Income						
≤ \$24,999	150	5.9	(2.3 - 9.5)	7,812	5.1	(3.3 - 7.0)
\$25,000-\$49,999	141	7.3	(2.7 - 11.9)	7,442	6.6	(3.9 - 9.2)
\$50,000-\$74,999	*	*	*	4,139	3.7	(1.7 - 5.8)
≥ \$75,000	*	*	*	5,872	2.4	(1.4 - 3.3)

¹ Defined as as respondent unintentionally coming to rest on the ground or another lower level and caused respondent to limit regular activities for at least a day or to go see a doctor one or more times within the past 3 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)