### eBRFSS 2013-2014 Health Indicators: Mohawk Valley

General Description	2
Improve Health Status and Reduce Health Disparities	3
Percentage of adults who did not receive medical care because of cost	3
Percentage of adults who have a regular health care provider	4
Percentage of adults who had a dentist visit within the past year	
Percentage of adults living with a disability	
Percentage of adults aged 18-64 years with healthcare coverage	
Percentage of adults aged 18-64 years who saw a doctor for a routine checkup within the last year	
Percentage of adults experiencing housing insecurity in the past 12 months	
Percentage of adults experiencing food insecurity in the past 12 months	
Percentage of adults with poor self-reported health	
Percentage of adults who report 14 or more days of poor physical health	
Prevent Chronic Diseases	
Percentage of adults who are obese	
Percentage of adults with an annual household income less than \$25,000 who are obese	
Percentage of adults living with a disability who are obese	
Percentage of adults overweight or obese	
Percentage of adults who participated in leisure time physical activity in the past 30 days	
Percentage of adults who consume one or more sugary drinks daily	
Percentage of adults who consume fast-food three or more times per week	
Percentage of adults with physician-diagnosed diabetes	
Percentage of adults who are current smokers	
Percentage of adults with annual household income less than \$25,000 who are current smokers	
Percentage of adults aged 18-24 years who are current smokers	
Percentage of women aged 50-74 years receiving breast cancer screening based on recent guidelines	
Percentage of women aged 50-74 years with annual household income of less than \$25,000 receiving breast cancer screening based or guidelines	
Percentage of women aged 21-65 years receiving cervical cancer screening based on recent guidelines	
guidelines	
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines	28
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines	28 recent
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines  Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on guidelines	28 recent 29
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines	28 recent 29
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines	28 recent 29 30
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines	28 recent393131
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines  Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on guidelines  Percentage of adults who had a test for high blood sugar or diabetes within the past three years  Percentage of adults with physician-diagnosed prediabetes  Percentage of adults with physician-diagnosed high blood pressure  Percentage of adults with diagnosed high blood pressure taking high blood pressure medication	28 recent29303132
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines  Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on guidelines  Percentage of adults who had a test for high blood sugar or diabetes within the past three years  Percentage of adults with physician-diagnosed prediabetes  Percentage of adults with physician-diagnosed high blood pressure  Percentage of adults with diagnosed high blood pressure taking high blood pressure medication  Percentage of adults with current asthma	
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines  Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on guidelines  Percentage of adults who had a test for high blood sugar or diabetes within the past three years  Percentage of adults with physician-diagnosed prediabetes  Percentage of adults with physician-diagnosed high blood pressure  Percentage of adults with diagnosed high blood pressure taking high blood pressure medication  Percentage of adults with current asthma  Percent of adults with cholesterol checked	28 recent293031323334
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines  Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on guidelines  Percentage of adults who had a test for high blood sugar or diabetes within the past three years  Percentage of adults with physician-diagnosed prediabetes  Percentage of adults with physician-diagnosed high blood pressure  Percentage of adults with diagnosed high blood pressure taking high blood pressure medication  Percentage of adults with current asthma  Percentage of adults with cholesterol checked  Percentage of adults 18 years and older with elevated cholesterol	28 recent303132333435
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines  Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on guidelines  Percentage of adults who had a test for high blood sugar or diabetes within the past three years  Percentage of adults with physician-diagnosed prediabetes  Percentage of adults with physician-diagnosed high blood pressure  Percentage of adults with diagnosed high blood pressure taking high blood pressure medication  Percentage of adults with current asthma  Percentage of adults with cholesterol checked  Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition	28 recent30313233343536
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines  Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on guidelines  Percentage of adults who had a test for high blood sugar or diabetes within the past three years  Percentage of adults with physician-diagnosed prediabetes  Percentage of adults with physician-diagnosed high blood pressure  Percentage of adults with diagnosed high blood pressure taking high blood pressure medication  Percentage of adults with current asthma  Percent of adults with cholesterol checked  Percentage of adults 18 years and older with elevated cholesterol  Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition  Percentage of adults with arthritis	28 recent30313234353637
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines  Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on guidelines  Percentage of adults who had a test for high blood sugar or diabetes within the past three years  Percentage of adults with physician-diagnosed prediabetes  Percentage of adults with physician-diagnosed high blood pressure  Percentage of adults with diagnosed high blood pressure taking high blood pressure medication  Percentage of adults with current asthma  Percentage of adults with cholesterol checked  Percentage of adults 18 years and older with elevated cholesterol  Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition  Percentage of adults with arthritis  Promote a Healthy and Safe Environment	28 recent3031323435363738
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines  Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on guidelines  Percentage of adults who had a test for high blood sugar or diabetes within the past three years  Percentage of adults with physician-diagnosed prediabetes  Percentage of adults with physician-diagnosed high blood pressure  Percentage of adults with diagnosed high blood pressure taking high blood pressure medication  Percentage of adults with current asthma  Percentage of adults with cholesterol checked  Percentage of adults 18 years and older with elevated cholesterol  Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition  Percentage of adults with arthritis  Promote a Healthy and Safe Environment  Percentage of adults who consider their neighborhood suitable for walking and physical activity	28 recent303132343536373839
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines  Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on guidelines  Percentage of adults who had a test for high blood sugar or diabetes within the past three years.  Percentage of adults with physician-diagnosed prediabetes  Percentage of adults with physician-diagnosed high blood pressure.  Percentage of adults with diagnosed high blood pressure taking high blood pressure medication.  Percentage of adults with current asthma  Percent of adults with cholesterol checked.  Percentage of adults 18 years and older with elevated cholesterol.  Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition.  Percentage of adults with arthritis  Promote a Healthy and Safe Environment.  Percentage of adults who consider their neighborhood suitable for walking and physical activity  Percentage of adults aged 65+ years with at least one fall in the past 12 months.	28 recent293132343536373839
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines.  Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on guidelines.  Percentage of adults who had a test for high blood sugar or diabetes within the past three years.  Percentage of adults with physician-diagnosed prediabetes.  Percentage of adults with diagnosed high blood pressure.  Percentage of adults with diagnosed high blood pressure taking high blood pressure medication.  Percentage of adults with current asthma.  Percent of adults with cholesterol checked.  Percentage of adults 18 years and older with elevated cholesterol.  Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition.  Percentage of adults with arthritis.  Promote a Healthy and Safe Environment.  Percentage of adults who consider their neighborhood suitable for walking and physical activity.  Percentage of adults aged 65+ years with at least one fall in the past 12 months.  Among adults aged 65+ years with at least one fall in the past 12 months, percent that resulted in injury.	
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines  Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on guidelines  Percentage of adults who had a test for high blood sugar or diabetes within the past three years  Percentage of adults with physician-diagnosed prediabetes  Percentage of adults with physician-diagnosed high blood pressure.  Percentage of adults with diagnosed high blood pressure taking high blood pressure medication  Percentage of adults with current asthma  Percentage of adults with cholesterol checked  Percentage of adults 18 years and older with elevated cholesterol  Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition  Percentage of adults with arthritis  Promote a Healthy and Safe Environment  Percentage of adults who consider their neighborhood suitable for walking and physical activity  Percentage of adults aged 65+ years with at least one fall in the past 12 months  Among adults aged 65+ years with at least one fall in the past 12 months, percent that resulted in injury  Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections	28 recent29303132343536373839394142
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines	
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines.  Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on guidelines.  Percentage of adults who had a test for high blood sugar or diabetes within the past three years.  Percentage of adults with physician-diagnosed prediabetes  Percentage of adults with physician-diagnosed high blood pressure	
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines  Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on guidelines  Percentage of adults who had a test for high blood sugar or diabetes within the past three years.  Percentage of adults with physician-diagnosed prediabetes  Percentage of adults with physician-diagnosed high blood pressure  Percentage of adults with diagnosed high blood pressure taking high blood pressure medication  Percentage of adults with current asthma  Percent of adults with cholesterol checked  Percentage of adults 18 years and older with elevated cholesterol  Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition  Percentage of adults with arthritis  Promote a Healthy and Safe Environment.  Percentage of adults who consider their neighborhood suitable for walking and physical activity  Percentage of adults aged 65+ years with at least one fall in the past 12 months.  Among adults aged 65+ years with at least one fall in the past 12 months.  Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections.  Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV)  Percentage of adults aged 65+ years with flu immunization in the past year.	
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines  Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on guidelines  Percentage of adults who had a test for high blood sugar or diabetes within the past three years  Percentage of adults with physician-diagnosed prediabetes  Percentage of adults with diagnosed high blood pressure  Percentage of adults with diagnosed high blood pressure taking high blood pressure medication  Percentage of adults with current asthma  Percentage of adults 18 years and older with elevated cholesterol  Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition  Percentage of adults with arthritis.  Promote a Healthy and Safe Environment  Percentage of adults who consider their neighborhood suitable for walking and physical activity  Percentage of adults aged 65+ years with at least one fall in the past 12 months  Among adults aged 65+ years with at least one fall in the past 12 months, percent that resulted in injury.  Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections  Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV)  Percentage of adults aged 65+ years with flu immunization in the past year.  Percentage of adults aged 65+ years with flu immunization in the past year.  Percentage of adults aged 65+ years with flu immunization in the past year.	
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines	
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines.  Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on guidelines.  Percentage of adults who had a test for high blood sugar or diabetes within the past three years	
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines  Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on guidelines  Percentage of adults who had a test for high blood sugar or diabetes within the past three years  Percentage of adults with physician-diagnosed prediabetes  Percentage of adults with physician-diagnosed high blood pressure  Percentage of adults with diagnosed high blood pressure taking high blood pressure medication.  Percentage of adults with colesterol checked.  Percentage of adults with cholesterol checked.  Percentage of adults with nathritis.  Percentage of adults with arthritis.  Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition  Percentage of adults who consider their neighborhood suitable for walking and physical activity  Percentage of adults aged 65+ years with at least one fall in the past 12 months.  Among adults aged 65+ years with at least one fall in the past 12 months.  Among adults aged 65+ years with at least one fall in the past 12 months.  Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV).  Percentage of adults aged 48+ years with flu immunization in the past year.  Percentage of adults aged 65+ years with flu immunization in the past year.  Percentage of adults aged 65+ years with flu immunization in the past year.  Percentage of women aged 18-44 years who had dental visit within the last year.  Women aged 18-44 years who saw a doctor for a routine checkup within the last year.	
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines  Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on guidelines  Percentage of adults who had a test for high blood sugar or diabetes within the past three years  Percentage of adults with physician-diagnosed prediabetes  Percentage of adults with physician-diagnosed high blood pressure  Percentage of adults with colorected of the percentage of adults with cholesterol checked  Percentage of adults with cholesterol checked  Percentage of adults 18 years and older with elevated cholesterol  Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition  Percentage of adults with arthritis  Promote a Healthy and Safe Environment  Percentage of adults who consider their neighborhood suitable for walking and physical activity  Percentage of adults aged 65+ years with at least one fall in the past 12 months  Among adults aged 65+ years with at least one fall in the past 12 months  Among adults aged 47-68 years reporting ever tested for Hepatitis C (HCV)  Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV)  Percentage of adults aged 65+ years with flu immunization in the past year  Percentage of adults aged 65+ years with flu immunization in the past year  Percentage of adults aged 65+ years with pneumococcal immunization  Percentage of women aged 18-44 years who had dental visit within the last year  Women aged 18-64 years who saw a doctor for a routine checkup within the last year  Women aged 18-64 years who saw a doctor for a routine checkup within the last year  Percentage of women aged 18-64 years with health care coverage	28 recent293031333435363738394041424244454646
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years. Percentage of adults with physician-diagnosed prediabetes. Percentage of adults with physician-diagnosed high blood pressure. Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percent of adults with cholesterol checked Percentage of adults 18 years and older with elevated cholesterol. Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults who consider their neighborhood suitable for walking and physical activity Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months, percent that resulted in injury Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections. Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV). Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of women aged 18-44 years who had dental visit within the last year Percentage of women aged 18-64 years with health care coverage Percentage of women aged 18-64 years with health care coverage Percentage of women aged 18-64 years with health care coverage Percentage of women aged 18	
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines  Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on guidelines  Percentage of adults who had a test for high blood sugar or diabetes within the past three years.  Percentage of adults with physician-diagnosed prediabetes.  Percentage of adults with diagnosed high blood pressure.  Percentage of adults with diagnosed high blood pressure.  Percentage of adults with current asthma.  Percent of adults with cholesterol checked.  Percentage of adults with care accurse or class to learn how to manage their chronic disease or condition.  Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition.  Percentage of adults with arthritis.  Promote a Healthy and Safe Environment.  Percentage of adults aged 65+ years with at least one fall in the past 12 months.  Among adults aged 65+ years with at least one fall in the past 12 months.  Among adults aged 65+ years with at least one fall in the past 12 months.  Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV).  Percentage of adults aged 48+ years with flu immunization in the past year.  Percentage of adults aged 65+ years with flu immunization in the past year.  Percentage of adults aged 65+ years with pneumococcal immunization.  Promote Healthy Women, Infants, and Children  Percentage of women aged 18-44 years who had dental visit within the last year.  Percentage of women aged 18-44 years who had dental visit within the last year.  Percentage of women aged 18-44 years who had dental visit within the last year.  Percentage of women aged 18-44 years who had dental visit within the last year.  Percentage of women aged 18-44 years who had dental visit within the last year.  Percentage of women aged 18-64 years who had dental visit within the last year.  Percentage of women aged 18-44 years who had bental visi	
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines.  Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on guidelines.  Percentage of adults who had a test for high blood sugar or diabetes within the past three years.  Percentage of adults with physician-diagnosed prediabetes.  Percentage of adults with physician-diagnosed high blood pressure.  Percentage of adults with diagnosed high blood pressure taking high blood pressure medication.  Percentage of adults with current asthma.  Percent of adults with cholesterol checked.  Percentage of adults with cholesterol checked.  Percentage of adults with arthritis.  Promote a Healthy and Safe Environment.  Percentage of adults who consider their neighborhood suitable for walking and physical activity.  Percentage of adults aged 65+ years with at least one fall in the past 12 months.  Among adults aged 65+ years with at least one fall in the past 12 months, percent that resulted in injury.  Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections.  Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV).  Percentage of adults aged 45+ years with flu immunization in the past year.  Percentage of adults aged 65+ years with pneumococcal immunization in the past year.  Percentage of adults aged 65+ years with pneumococcal immunization.  Promote Healthy Momen, Infants, and Children  Percentage of women aged 18-64 years with health care coverage.  Percentage of women aged 18-64 years with health care coverage.  Percentage of women aged 18-64 years with sew hoad dental visit within the last year.  Percentage of women aged 18-64 years with sew hose health care coverage.  Percentage of women aged 18-64 years with sew hoad dental visit within the last year.  Percentage of women aged 18-64 years with health care coverage.  Percentage of women aged 18-64 years with ose health care provider discussed p	
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines  Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on guidelines  Percentage of adults who had a test for high blood sugar or diabetes within the past three years.  Percentage of adults with physician-diagnosed prediabetes.  Percentage of adults with diagnosed high blood pressure.  Percentage of adults with diagnosed high blood pressure.  Percentage of adults with current asthma.  Percent of adults with cholesterol checked.  Percentage of adults with care accurse or class to learn how to manage their chronic disease or condition.  Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition.  Percentage of adults with arthritis.  Promote a Healthy and Safe Environment.  Percentage of adults aged 65+ years with at least one fall in the past 12 months.  Among adults aged 65+ years with at least one fall in the past 12 months.  Among adults aged 65+ years with at least one fall in the past 12 months.  Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV).  Percentage of adults aged 48+ years with flu immunization in the past year.  Percentage of adults aged 65+ years with flu immunization in the past year.  Percentage of adults aged 65+ years with pneumococcal immunization.  Promote Healthy Women, Infants, and Children  Percentage of women aged 18-44 years who had dental visit within the last year.  Percentage of women aged 18-44 years who had dental visit within the last year.  Percentage of women aged 18-44 years who had dental visit within the last year.  Percentage of women aged 18-44 years who had dental visit within the last year.  Percentage of women aged 18-44 years who had dental visit within the last year.  Percentage of women aged 18-64 years who had dental visit within the last year.  Percentage of women aged 18-44 years who had bental visi	

#### General Description

The New York State Expanded Behavioral Risk Factor Surveillance System 2013-2014 (2014 eBRFSS) was design to supplement the Center for Disease Control and Prevention (CDC) Behavioral Risk Factor Surveillance System (BRFSS), which is conducted annually in New York State. The purpose of the 2014 eBRFSS was to produce local information on key public health issues. Data for the project were collected from April 15, 2013 to May 10, 2014. The 2014 eBRFSS reached both households with landline telephones and households which only had cell phones. Like the annual BRFSS, New York State's 2014 eBRFSS was designed to be representative of the non-institutionalized adult household population, aged 18 years and older.

#### **Reported Health Indicators**

There are a total of 50 health indicators included in the report. The health indicators include 10 measures included in the New York State Department of Health's Prevention Agenda, NYS Health Innovation Plan (SHIP), Delivery System Reform Incentive Payment (DSRIP) Program, or the NYS Community Health Indicator Reports (CHIRS). An additional 40 important health indicators are also included. The indicators are organized by the Priority Area of the Prevention Agenda and are reported statewide, by region, and by county.

#### **Data Collection Methodology, Sampling and Weighting**

The data collection methods for the 2014 eBRFSS aligned with the CDC's standard annual BRFSS methods. The 2014 eBRFSS questionnaire used standard questions from the BRFSS. Survey interviews were conducted in both English and Spanish. There were 31,690 completed interviews.

The sampling plan for the eBRFSS was developed to result in valid health indicator estimates for New York City (NYC) and each of the 57 counties outside NYC. The sampling approach stratified landlines by county and implemented a disproportionate stratified sampling design that accounted for both listed-households and unlisted numbers. Cell phone households were accounted for through additional strata that covered contiguous groups of counties.

Data were weighted to represent the population using a methodology known as iterative proportional fitting (raking). Raking adjusted the data so that groups underrepresented in the sample were more accurately represented in the estimated rates. Raking also enabled surveys conducted by cell phone to be included in the weighting and to accurately match sample distributions of specified demographic characteristics of the population. The raking categories included in the eBRFSS were: age, gender, race/ethnicity, education, marital status, owner/renter status, telephone usage (i.e. landline and cell phone combinations), and county of residence.


#### **Limitations of Data Use:**

During data analysis, criteria were applied to suppress rates based on too few surveys to be reliable and to identify rates that are highly variable. Rates in which less than 10 respondents were in the numerator or less than 50 respondents were in the denominator were considered unreliable and suppressed from reporting. Rates considered to have high variability due to having confidence limits greater than ±10% are noted.

### Mohawk Valley

### Improve Health Status and Reduce Health Disparities

Percentage of adults who did not receive medical care because of cost\*



		Mohawk	( Valley			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
_	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	25,495	12.2	9.7	15.3	1,994,210	13.1	11.2	15.3
Age								
18-24	4,963	20.3^	9.6	38.0	281,654	14.1^	7.5	25.1
25-34	6,352	19.9^	11.5	32.0	537,866	19.7	14.2	26.6
35-44	2,409	8.5	5.3	13.5	423,154	17.0	12.3	23.0
45-64	10,267	13.5	10.4	17.2	596,074	11.9	9.8	14.3
65+	1,314	2.9	1.8	4.4	139,733	5.1	3.0	8.4
Race								
White, NH	21,153	11.5	8.8	14.8	959,531	10.5	8.7	12.7
Black, NH		#			335,184	15.5	10.0	23.4
Hispanic	2,119	17.6^	8.3	33.5	465,114	18.4	14.5	23.0
Other, NH	1,640	18.3^	7.6	37.9	234,381	16.3	7.3	32.6
Sex								
Male	13,758	13.6	9.3	19.4	890,677	12.3	9.6	15.5
Female	11,736	10.9	8.5	13.9	1,103,533	13.8	11.2	16.9
Age-adjusted <sup>&amp;</sup>		12.6	10.0	15.9		13.6	11.6	15.8


<sup>\*</sup> Among people needing to see a doctor in the past 12 months

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

#### Percentage of adults who have a regular health care provider




		Mohawl	( Valley			State	wide	
_	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total	178,551	86.4	83.5	88.9	12,905,860	84.9	83.1	86.6
Age								
18-24	17,923	77.3^	64.8	86.3	1,486,954	76.3	69.8	81.8
25-34	24,155	75.8^	64.5	84.4	1,983,479	72.3	66.1	77.7
35-44	22,811	82.1^	71.0	89.6	2,091,488	84.7	79.6	88.7
45-64	68,580	89.9	87.0	92.2	4,450,345	88.6	85.5	91.1
65+	43,482	94.9	91.5	97.0	2,640,269	96.0	94.7	97.1
Race								
White, NH	159,236	87.4	84.6	89.7	7,996,007	88.0	86.6	89.3
Black, NH		#			1,926,090	89.0	83.6	92.7
Hispanic	9,218	72.8^	50.4	87.6	1,725,153	68.6	61.2	75.1
Other, NH	7,804	92.5^	82.1	97.1	1,258,610	87.9	79.9	93.0
Sex								
Male	79,609	79.7	74.3	84.1	5,804,831	80.1	77.1	82.8
Female	98,942	92.8	90.6	94.5	7,101,029	89.3	87.0	91.3
Age-adjusted <sup>&amp;</sup>		84.8	81.4	87.8		84.5	82.5	86.2

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

#### Percentage of adults who had a dentist visit within the past year




		Mohawl	( Valley			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
_	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	120,003	61.7	58.1	65.0	9,774,786	69.5	66.9	72.0
Age								
18-24	13,968	62.7^	47.0	76.1	1,100,494	62.2	51.0	72.2
25-34	19,077	62.4^	51.4	72.3	1,624,123	65.1	57.7	71.8
35-44	17,119	71.2	62.4	78.6	1,687,707	71.9	65.2	77.8
45-64	42,537	58.2	52.9	63.3	3,421,923	72.6	69.3	75.7
65+	26,653	61.4	56.5	66.1	1,743,883	69.8	66.0	73.4
Race								
White, NH	108,248	62.7	59.2	66.0	6,152,109	72.4	70.1	74.6
Black, NH		#			1,309,282	66.6	57.4	74.7
Hispanic	7,338	68.9^	51.9	82.0	1,435,933	62.9	55.3	70.0
Other, NH	3,087	38.3^	17.9	63.9	877,462	66.5	52.1	78.4
Sex								
Male	51,611	54.2	48.6	59.6	4,591,800	67.3	63.4	71.1
Female	68,392	68.8	65.0	72.5	5,182,986	71.6	68.1	74.8
Age-adjusted <sup>&amp;</sup>		62.9	59.1	66.6		69.3	66.6	71.8

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

### Percentage of adults living with a disability\*



		Mohawl	र Valley		Statewide			
	Est. # of	_	Lower	Upper	Est. # of	_	Lower	Upper
г	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	56,344	27.3	24.4	30.4	3,111,131	20.8	19.1	22.7
Age								
18-24		#			119,872	6.1	4.3	8.5
25-34	5,335	16.7	10.9	24.8	426,647	16.0	10.8	23.2
35-44	6,828	24.3^	15.9	35.4	294,471	12.2	9.7	15.3
45-64	24,816	32.9	28.0	38.2	1,195,176	24.2	21.6	27.1
65+	16,563	36.9	32.4	41.6	989,611	36.8	33.1	40.6
Race								
White, NH	46,724	25.7	23.2	28.5	2,187,495	24.4	22.4	26.6
Black, NH		#			332,469	15.8	11.9	20.7
Hispanic	5,344	42.2	25.0	61.5	420,342	17.0	12.7	22.5
Other, NH	3,225	35.9^	14.4	65.1	170,824	12.3	7.7	19.1
Sex								
Male	27,372	27.1	22.4	32.3	1,349,486	18.9	16.3	21.9
Female	28,972	27.6	24.1	31.3	1,761,645	22.6	20.4	24.9
Age-adjusted <sup>&amp;</sup>		25.8	22.7	29.2		19.9	18.3	21.6


<sup>\*</sup> Limited activities (physical, mental, or emotional problems); need of special equipment (e.g. cane, wheel chair).

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

#### Percentage of adults aged 18-64 years with healthcare coverage



		Mohawl	k Valley			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
г	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	135,591	84.7	80.9	87.9	10,275,389	85.2	82.8	87.3
Age⁺								
18-24								
25-34								
35-44								
45-64								
65+								
Race								
White, NH	119,366	85.5	81.6	88.8	6,192,415	90.1	88.8	91.3
Black, NH		#			1,606,586	87.9	82.0	92.1
Hispanic	7,974	72.7	48.5	88.3	1,626,540	73.2	66.2	79.2
Other, NH		#			849,849	74.5^	56.8	86.6
Sex								
Male	62,882	80.6	73.5	86.1	4,924,290	83.2	79.9	86.0
Female	72,708	88.7	85.3	91.4	5,351,099	87.2	83.4	90.2
Age-adjusted <sup>+,&amp;</sup>								

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>+</sup> Age-adjusted and age specific percent not generated for health indicators with age range limits.


<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

### Mohawk Valley

### Improve Health Status and Reduce Health Disparities

Percentage of adults aged 18-64 years who saw a doctor for a routine checkup within the last year



		Mohawl	( Valley			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	114,077	71.8	68.1	75.2	8,622,736	70.9	68.1	73.5
Age⁺								
18-24								
25-34								
35-44								
45-64								
65+								
Race								
White, NH	99,065	71.6	67.9	75.0	4,721,851	68.7	65.8	71.4
Black, NH		#			1,477,782	80.8	73.1	86.6
Hispanic	7,461	68.0^	44.9	84.7	1,583,308	71.1	63.6	77.6
Other, NH		#			839,795	68.4	53.1	80.5
Sex								
Male	51,586	67.1	61.0	72.8	3,938,722	65.6	61.4	69.6
Female	62,491	76.2	71.9	80.1	4,684,014	76.0	72.3	79.4
Age-adjusted+,&								

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>+</sup> Age-adjusted and age specific percent not generated for health indicators with age range limits.


<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

### Mohawk Valley

### Improve Health Status and Reduce Health Disparities

Percentage of adults experiencing housing insecurity in the past 12 months\*



		Mohawl	र Valley			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
<b>-</b>	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	68,431	36.6	33.3	40.0	5,967,540	43.4	40.8	46.1
Age								
18-24	5,154	29.2^	16.5	46.1	487,620	30.7	21.3	42.0
25-34	13,314	44.3^	34.2	55.0	1,403,277	57.7	50.8	64.3
35-44	12,783	50.7	40.9	60.5	1,180,494	50.7	43.4	57.9
45-64	28,632	40.5	35.7	45.4	2,100,265	44.9	41.4	48.4
65+	8,011	19.0	15.8	22.7	688,870	27.8	23.8	32.1
Race								
White, NH	60,465	36.6	33.3	40.0	3,030,019	36.7	34.3	39.2
Black, NH		#			1,078,432	55.5	46.2	64.3
Hispanic	4,317	40.8^	23.4	60.7	1,268,822	57.3	49.7	64.6
Other, NH	2,119	26.7	12.4	48.4	590,266	44.6^	31.2	58.9
Sex								
Male	29,853	33.1	28.1	38.5	2,630,128	40.1	36.1	44.2
Female	38,579	39.8	35.7	44.1	3,337,412	46.5	43.0	50.0
Age-adjusted <sup>&amp;</sup>		38.3	34.5	42.2		43.7	41.0	46.5


<sup>\*</sup> Worried or stressed about having enough money to pay for rent or mortgage.

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

Percentage of adults experiencing food insecurity in the past 12 months\*



		Mohawl	र Valley			State	wide	
_	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total Age	51,745	26.0	22.9	29.2	4,211,702	29.0	26.6	31.5
18-24	6,939	30.9^	18.6	46.7	339,128	17.9	13.5	23.3
25-34	10,887	35.0^	25.5	46.0	1,154,171	44.8	37.7	52.1
35-44	9,575	36.6	28.0	46.2	786,490	33.0	26.3	40.4
45-64	19,400	26.2	22.3	30.4	1,464,969	30.3	26.9	33.9
65+	4,715	10.7	8.2	13.7	405,259	15.6	12.5	19.3
Race								
White, NH	44,036	25.0	22.0	28.3	1,998,541	22.9	20.6	25.4
Black, NH		#			765,155	37.2	29.3	45.7
Hispanic	4,530	37.2	21.3	56.5	1,031,787	43.4	36.2	50.8
Other, NH	2,354	29.0^	13.7	51.3	416,220	30.4	20.2	42.9
Sex								
Male	23,314	23.9	19.2	29.2	1,903,904	27.4	23.6	31.5
Female	28,431	28.0	24.3	32.0	2,307,798	30.5	27.5	33.6
Age-adjusted <sup>&amp;</sup>		28.0	24.6	31.8		29.4	27.1	31.9


<sup>\*</sup> Worried or stressed about having enough money to buy nutritious meals.

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

### Percentage of adults with poor self-reported health



		Mohawl	र Valley			State	wide	
_	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total Age	10,550	5.1	3.9	6.5	680,314	4.5	3.7	5.4
18-24		#			20,778	1.0	0.5	2.3
25-34		#			38,067	1.4	0.8	2.5
35-44		#			58,460	2.4	1.5	3.9
45-64	6,176	8.2	5.8	11.3	300,529	6.1	4.5	8.1
65+	2,659	5.8	4.1	8.2	248,637	9.1	6.9	12.0
Race								
White, NH	9,037	4.9	3.8	6.3	365,138	4.0	3.5	4.7
Black, NH		#			58,105	2.7	1.6	4.7
Hispanic		#			213,536	8.4	5.3	13.2
Other, NH		#			43,535	3.0	1.4	6.4
Sex								
Male	3,898	3.9	2.6	5.6	298,899	4.1	3.1	5.5
Female	6,652	6.2	4.5	8.6	381,415	4.8	3.9	6.0
Age-adjusted <sup>&amp;</sup>		4.6	3.4	6.1		4.3	3.6	5.1


<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

### Mohawk Valley

### Improve Health Status and Reduce Health Disparities

Percentage of adults who report 14 or more days of poor physical health\*



		Mohawl	( Valley			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	29,861	14.8	12.7	17.2	1,717,941	11.8	10.5	13.1
Age								
18-24		#			59,108	3.1	1.9	5.1
25-34	3,109	9.9	5.3	17.8	184,980	7.0	4.8	10.0
35-44	4,086	15.3	9.4	23.8	176,691	7.2	5.2	9.9
45-64	13,912	18.9	15.3	23.1	763,670	15.8	13.5	18.5
65+	6,869	15.7	12.7	19.2	491,091	19.1	16.1	22.5
Race								
White, NH	25,670	14.4	12.3	16.8	1,043,432	11.8	10.7	13.0
Black, NH		#			188,315	9.3	6.3	13.5
Hispanic	3,000	24.1^	12.3	41.8	372,059	16.0	11.5	21.9
Other, NH	1,035	12.8^	4.1	33.4	114,135	8.1	4.7	13.7
Sex								
Male	11,588	12.0	9.4	15.1	722,008	10.3	8.7	12.3
Female	18,273	17.5	14.3	21.1	995,933	13.0	11.3	14.9
Age-adjusted <sup>&amp;</sup>		14.4	12.1	17.0		11.3	10.1	12.5


<sup>\*</sup> Out of the past 30 days.

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

#### Percentage of adults who are obese\*



		Mohawl	र Valley			State	wide	
_	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total Age	60,595	30.8	27.7	34.0	3,527,077	24.9	23.0	26.9
18-24	4,420	18.4^	8.7	34.7	318,556	17.8	11.7	26.0
25-34	9,122	31.3	22.1	42.3	509,101	20.8	16.8	25.4
35-44	9,682	36.9	28.3	46.4	573,976	24.7	20.0	30.1
45-64	24,732	34.4	29.8	39.3	1,452,755	30.5	27.5	33.7
65+	12,362	27.8	23.8	32.2	615,010	23.4	20.1	27.0
Race								
White, NH	53,932	31.0	27.9	34.4	2,139,878	25.0	23.2	26.9
Black, NH		#			638,079	30.7	24.2	38.1
Hispanic	2,996	27.1^	14.7	44.5	622,984	28.0	21.8	35.1
Other, NH	2,236	25.2	11.2	47.3	126,136	9.6	6.3	14.4
Sex								
Male	31,672	32.0	27.2	37.2	1,782,686	25.3	22.5	28.3
Female	28,923	29.5	25.7	33.6	1,744,390	24.5	22.0	27.1
Age-adjusted <sup>&amp;</sup>		31.2	27.7	34.9		24.6	22.7	26.6


<sup>\*</sup> Obesity defined as a Body Mass Index (BMI) >=30. BMI is calculated based on weight and height.

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

Percentage of adults with an annual household income less than \$25,000 who are obese\*



		Mohawl	र Valley			State	wide	
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total Age	17,765	32.2	26.8	38.3	1,029,079	28.7	24.7	33.0
18-24		#			88,578	21.3^	10.0	39.7
25-34		#			177,054	26.6	17.0	39.0
35-44	2,079	32.6^	17.9	51.7	165,333	28.8^	19.5	40.3
45-64	6,638	34.5	27.0	42.9	386,736	34.8	28.0	42.1
65+	4,972	30.4	23.7	38.1	190,291	24.4	18.8	31.1
Race								
White, NH	14,894	31.6	25.8	38.0	500,787	30.3	25.6	35.4
Black, NH		#			165,021	25.3^	16.7	36.3
Hispanic		#			287,948	30.1^	21.1	41.0
Other, NH		#			75,322	23.1	12.4	39.0
Sex								
Male	8,275	34.0^	24.5	45.0	386,764	23.8	18.1	30.5
Female	9,491	30.9	24.9	37.5	642,314	32.7	27.6	38.3
Age-adjusted <sup>&amp;</sup>		32.3	26.1	39.3		28.5	24.3	33.0


<sup>\*</sup> Obesity defined as a Body Mass Index (BMI) >=30. BMI is calculated based on weight and height.

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

#### Percentage of adults living with a disability who are obese\*



		Mohawl	( Valley			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
_	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	22,893	43.0	36.7	49.6	1,093,091	36.9	32.7	41.3
Age								
18-24		#			18,440	16.5^	9.8	26.5
25-34		#			122,752	31.0^	16.7	50.1
35-44		#			109,843	39.7	30.4	49.8
45-64	10,332	44.7^	34.6	55.3	535,031	45.9	39.5	52.5
65+	6,555	40.1	32.9	47.8	283,502	30.3	25.2	36.1
Race								
White, NH	19,729	44.7	38.9	50.8	698,173	33.3	29.0	37.8
Black, NH		#			188,077	58.3^	44.8	70.6
Hispanic		#			170,166	46.0^	30.4	62.5
Other, NH		#			36,674	21.7	12.1	35.9
Sex								
Male	10,802	40.2^	30.8	50.3	430,886	32.6	26.2	39.7
Female	12,091	45.9	38.0	54.0	662,205	40.4	35.1	45.8
Age-adjusted <sup>&amp;</sup>		43.3	33.9	53.1		35.4	31.0	40.0


<sup>\*</sup> Obesity defined as a Body Mass Index (BMI) >=30. BMI is calculated based on weight and height. Limited activities (physical, mental, or emotional problems); need of special equipment (e.g. cane, wheel chair).

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

#### Percentage of adults overweight or obese\*



		Mohawk	( Valley			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
<u>-</u>	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	131,535	66.8	63.3	70.1	8,631,360	60.9	58.4	63.3
Age								
18-24	10,728	44.7^	30.9	59.3	656,771	36.6	28.8	45.2
25-34	16,842	57.8^	47.2	67.8	1,528,245	62.4	55.9	68.4
35-44	20,315	77.4	70.0	83.4	1,362,541	58.6	50.8	66.0
45-64	51,948	72.3	67.1	77.0	3,335,153	70.0	66.8	73.1
65+	31,025	69.8	65.3	73.9	1,632,723	62.1	58.1	65.9
Race								
White, NH	115,938	66.7	63.3	70.0	5,260,942	61.5	59.3	63.5
Black, NH		#			1,520,031	73.1	63.8	80.7
Hispanic	8,112	73.4^	55.6	85.9	1,346,413	60.5	52.9	67.5
Other, NH	4,931	55.5	29.6	78.8	503,974	38.4	28.1	49.9
Sex								
Male	70,693	71.4	65.8	76.5	4,736,847	67.1	63.2	70.8
Female	60,843	62.1	57.8	66.2	3,894,512	54.7	51.4	57.9
Age-adjusted <sup>&amp;</sup>		66.8	63.2	70.2		60.5	57.9	63.0


<sup>\*</sup> Defined as a Body Mass Index (BMI) >=25. BMI is calculated based on weight and height.

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

Percentage of adults who participated in leisure time physical activity in the past 30 days\*



		Mohawl	( Valley			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	138,763	69.2	66.2	72.1	10,681,161	72.8	70.6	74.9
Age								
18-24	16,894	73.4^	58.5	84.5	1,601,630	84.1	78.5	88.5
25-34	26,053	82.7	75.0	88.4	1,900,064	73.1	67.1	78.4
35-44	16,488	63.4	53.8	72.0	1,722,858	71.5	63.8	78.1
45-64	49,363	66.6	61.9	71.0	3,443,881	70.8	67.5	73.8
65+	29,150	65.3	60.7	69.6	1,840,519	69.5	66.1	72.7
Race								
White, NH	122,921	69.5	66.4	72.4	6,662,678	75.8	74.0	77.4
Black, NH		#			1,525,317	73.8	66.8	79.7
Hispanic	7,241	59.4^	40.6	75.8	1,507,430	62.4	55.4	68.9
Other, NH	6,518	80.5	63.1	90.9	985,735	70.6^	56.9	81.3
Sex								
Male	69,524	70.6	65.7	75.1	5,439,883	77.3	74.6	79.8
Female	69,239	67.9	64.0	71.5	5,241,278	68.6	65.3	71.7
Age-adjusted <sup>&amp;</sup>		69.5	66.1	72.7		72.9	70.5	75.1


<sup>\*</sup> Any physical activities or exercises such as running or walking

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

#### Percentage of adults who consume one or more sugary drinks daily\*



		Mohawl	र Valley			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
<del>-</del>	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	52,428	26.6	23.4	30.0	3,402,766	23.8	21.5	26.3
Age								
18-24	9,776	43.6^	30.0	58.2	670,630	36.9	28.0	46.7
25-34	12,116	39.3^	29.4	50.1	733,289	28.9	22.5	36.4
35-44	9,558	36.9^	27.1	48.0	769,136	32.4	25.4	40.3
45-64	13,685	18.8	15.4	22.7	839,390	17.6	15.0	20.5
65+	7,223	16.4	13.1	20.4	348,734	13.7	11.1	16.9
Race								
White, NH	44,750	25.7	22.6	29.1	1,754,487	20.4	18.3	22.8
Black, NH		#			671,592	33.6	25.4	43.0
Hispanic	5,831	48.2^	30.0	66.9	700,337	29.8	23.7	36.8
Other, NH		#			276,350	20.4	10.6	35.5
Sex								
Male	30,568	31.7	26.6	37.3	1,782,648	26.0	22.5	29.8
Female	21,860	21.7	18.1	25.7	1,620,118	21.8	18.7	25.3
Age-adjusted <sup>&amp;</sup>		29.3	25.7	33.2		24.7	22.2	27.4


<sup>\*</sup> Sugary drinks include regular soda, and sugar-sweetened drinks.

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

 $<sup>^{\</sup>wedge}$  Highly variable rate (confidence interval with a half-width greater than 10).

#### Percentage of adults who consume fast-food three or more times per week




				Statewide				
Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	
8,472	4.3	3.2	5.8	817,815	5.8	4.8	6.9	
	#			172,464	9.7	6.9	13.6	
1,686	5.5	2.7	10.9	252,437	10.0	6.9	14.2	
946	3.7	1.9	6.9	106,397	4.5	3.2	6.3	
2,524	3.5	2.4	5.0	234,582	5.0	3.3	7.4	
1,145	2.6	1.5	4.5	47,255	1.9	1.3	2.7	
7,397	4.3	3.1	5.8	433,016	5.1	4.4	5.9	
	#			122,943	6.2	4.1	9.3	
	#			219,779	9.6	5.6	16.0	
	#			42,077	3.2	1.8	5.5	
6,106	6.4	4.4	9.2	560,514	8.2	6.6	10.3	
2,366	2.4	1.5	3.6	257,301	3.5	2.6	4.8	
	4.5	3.3	6.2		5.9	4.9	7.0	
	1,686 946 2,524 1,145 7,397	Adults     Percent       8,472     4.3       #     1,686       946     3.7       2,524     3.5       1,145     2.6       7,397     4.3       #     #       #     #       6,106     6.4	Adults         Percent         95% limit           8,472         4.3         3.2           #         1,686         5.5         2.7           946         3.7         1.9           2,524         3.5         2.4           1,145         2.6         1.5           7,397         4.3         3.1           #         #         #           6,106         6.4         4.4           2,366         2.4         1.5	Adults         Percent         95% limit         95% limit           8,472         4.3         3.2         5.8           #         1,686         5.5         2.7         10.9           946         3.7         1.9         6.9           2,524         3.5         2.4         5.0           1,145         2.6         1.5         4.5           7,397         4.3         3.1         5.8           #         #         #           6,106         6.4         4.4         9.2           2,366         2.4         1.5         3.6	Adults         Percent         95% limit         95% limit         Adults           8,472         4.3         3.2         5.8         817,815           #         172,464         172,464           1,686         5.5         2.7         10.9         252,437           946         3.7         1.9         6.9         106,397           2,524         3.5         2.4         5.0         234,582           1,145         2.6         1.5         4.5         47,255           7,397         4.3         3.1         5.8         433,016           #         122,943         219,779         42,077           6,106         6.4         4.4         9.2         560,514           2,366         2.4         1.5         3.6         257,301	Adults         Percent         95% limit         95% limit         Adults         Percent           8,472         4.3         3.2         5.8         817,815         5.8           #         172,464         9.7           1,686         5.5         2.7         10.9         252,437         10.0           946         3.7         1.9         6.9         106,397         4.5           2,524         3.5         2.4         5.0         234,582         5.0           1,145         2.6         1.5         4.5         47,255         1.9           7,397         4.3         3.1         5.8         433,016         5.1           #         122,943         6.2         219,779         9.6           #         219,779         9.6         42,077         3.2           6,106         6.4         4.4         9.2         560,514         8.2           2,366         2.4         1.5         3.6         257,301         3.5	Adults         Percent         95% limit         95% limit         Adults         Percent         95% limit           8,472         4.3         3.2         5.8         817,815         5.8         4.8           #         172,464         9.7         6.9         6.9         106,347         10.0         6.9           946         3.7         1.9         6.9         106,397         4.5         3.2         3.2           2,524         3.5         2.4         5.0         234,582         5.0         3.3           1,145         2.6         1.5         4.5         47,255         1.9         1.3           7,397         4.3         3.1         5.8         433,016         5.1         4.4           #         122,943         6.2         4.1         219,779         9.6         5.6           #         42,077         3.2         1.8           6,106         6.4         4.4         9.2         560,514         8.2         6.6           2,366         2.4         1.5         3.6         257,301         3.5         2.6	

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

#### Percentage of adults with physician-diagnosed diabetes\*



		Mohawl	र Valley			State	wide	
_	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total Age	26,908	12.9	11.0	15.0	1,456,783	9.5	8.4	10.8
18-24		#			8,338	0.4	0.2	0.8
25-34	1,776	5.6	2.7	11.2	58,319	2.1	0.8	5.5
35-44	2,255	8.0	4.0	15.5	111,205	4.5	2.5	7.7
45-64	11,420	15.0	11.9	18.6	714,501	14.2	11.9	16.9
65+	11,318	24.6	20.0	29.9	525,500	19.1	16.3	22.2
Race								
White, NH	21,832	11.8	10.2	13.8	781,028	8.6	7.6	9.6
Black, NH		#			250,921	11.6	8.2	16.2
Hispanic	1,646	13.0^	5.8	26.6	296,635	11.6	7.7	17.0
Other, NH	2,456	27.4	10.7	54.3	128,200	8.9	5.4	14.4
Sex								
Male	13,341	13.1	10.2	16.8	795,054	10.9	9.0	13.1
Female	13,567	12.6	10.4	15.3	661,729	8.3	7.0	9.7
Age-adjusted <sup>&amp;</sup>		11.4	9.7	13.5		8.9	7.9	10.1


<sup>\*</sup> As diagnosed by a health professional.

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

 $<sup>^{\</sup>wedge}$  Highly variable rate (confidence interval with a half-width greater than 10).

#### Percentage of adults who are current smokers




		Mohawl	( Valley			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
r	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	52,067	25.3	22.3	28.7	2,311,528	15.6	14.0	17.4
Age								
18-24	5,575	23.0^	12.8	37.9	254,033	13.0	9.8	17.0
25-34	12,670	39.8^	29.8	50.7	524,783	19.7	14.3	26.7
35-44	9,692	34.7	26.0	44.6	427,187	17.6	13.3	23.0
45-64	20,418	27.1	23.0	31.6	937,977	19.2	16.7	22.1
65+	3,513	7.9	5.8	10.8	149,820	5.7	4.7	6.8
Race								
White, NH	45,456	25.2	22.0	28.6	1,598,923	18.1	16.0	20.3
Black, NH		#			314,837	15.0	10.0	21.7
Hispanic	3,750	29.7^	15.5	49.2	303,908	12.4	8.8	17.1
Other, NH	2,323	25.9^	11.1	49.5	93,860	6.7	4.3	10.3
Sex								
Male	28,059	27.9	22.9	33.5	1,331,531	18.8	15.9	22.1
Female	24,008	22.9	19.5	26.7	979,997	12.7	11.1	14.4
Age-adjusted <sup>&amp;</sup>		27.3	23.9	30.9		15.9	14.1	17.7

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

#### Percentage of adults with annual household income less than \$25,000 who are current smokers




		Mohawl	( Valley			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
<u>-</u>	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	19,843	34.8	29.0	41.0	910,061	23.6	19.2	28.7
Age								
18-24		#			84,893	16.1	9.8	25.4
25-34	3,104	44.7^	29.0	61.5	254,993	32.7	18.8	50.6
35-44	4,248	66.0^	48.0	80.3	141,275	24.5	15.5	36.5
45-64	7,288	36.4	29.1	44.3	354,621	31.5	24.7	39.1
65+	1,892	11.7	7.6	17.6	63,044	7.9	5.6	11.0
Race								
White, NH	16,819	34.6	28.7	41.1	562,213	32.6	25.7	40.3
Black, NH		#			187,054	28.1^	17.8	41.4
Hispanic		#			119,918	11.3	7.1	17.6
Other, NH		#			40,876	10.0	5.0	19.2
Sex								
Male	9,513	39.1^	29.5	49.6	507,987	30.0	21.9	39.5
Female	10,330	31.5	24.7	39.2	402,073	18.6	14.7	23.3
Age-adjusted <sup>&amp;</sup>		41.4	35.2	47.9		24.2	20.0	28.9

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

#### Percentage of adults aged 18-24 years who are current smokers



		Mohawl	k Valley		Statewide			
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
F	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	5,575	23.0^	12.8	37.9	254,033	13.0	9.8	17.0
Age⁺								
18-24								
25-34								
35-44								
45-64								
65+								
Race								
White, NH	5,397	24.9^	13.6	41.1	188,030	19.4	14.8	25.2
Black, NH		#			21,086	6.1	2.8	13.1
Hispanic		#			33,121	9.8^	4.6	19.9
Other, NH		#			11,797	3.8	1.4	10.0
Sex								
Male		#			153,298	15.5	10.7	21.8
Female		#			100,736	10.4	6.6	16.0
Age-adjusted+,&								


<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>+</sup> Age-adjusted and age specific percent not generated for health indicators with age range limits.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

Percentage of women aged 50-74 years receiving breast cancer screening based on recent guidelines\*



		Mohawl	र Valley			State	wide	
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total Age+ 18-24 25-34 35-44 45-64 65+ Race	30,177	75.7	70.6	80.1	2,225,651	80.9	77.7	83.7
White, NH Black, NH Hispanic Other, NH Sex Male Female	27,509 30,177	78.0 # # # # 75.7	73.7 70.6	81.8 80.1	1,417,374 347,690 309,667 150,920 2,225,651	79.0 87.0^ 83.8^ 79.7^ # 80.9	75.9 73.4 70.9 61.2	81.8 94.3 91.6 90.7
Age-adjusted+,&								

<sup>\*</sup> Mammogram within past 2 years.

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>+</sup> Age-adjusted and age specific percent not generated for health indicators with age range limits.


<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

 $<sup>^{\</sup>wedge}$  Highly variable rate (confidence interval with a half-width greater than 10).

### Mohawk Valley

#### **Prevent Chronic Diseases**

Percentage of women aged 50-74 years with annual household income of less than \$25,000 receiving breast cancer screening based on recent guidelines\*



		Mohawl	ς Valley			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
-	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	7,622	63.8	54.2	72.4	554,880	76.7	69.6	82.6
Age <sup>+</sup>								
18-24								
25-34								
35-44								
45-64								
65+								
Race								
White, NH	6,482	66.1	56.5	74.5	232,138	66.5	57.9	74.1
Black, NH		#			139,053	95.4	89.6	98.1
Hispanic		#			134,524	82.0^	61.8	92.7
Other, NH		#			49,165	76.3^	44.1	93.0
Sex								
Male		#				#		
Female	7,622	63.8	54.2	72.4	554,880	76.7	69.6	82.6
Age-adjusted <sup>+,&amp;</sup>								

<sup>\*</sup> Mammogram within past 2 years.


<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>+</sup> Age-adjusted and age specific percent not generated for health indicators with age range limits.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

Percentage of women aged 21-65 years receiving cervical cancer screening based on recent guidelines\*



		Mohawl	k Valley		Statewide			
_	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total Age <sup>+</sup> 18-24	58,844	79.9	76.1	83.2	4,362,995	80.6	75.7	84.7
25-34 35-44 45-64 65+								
Race White, NH Black, NH Hispanic Other, NH Sex	52,741	80.0 # # #	76.1	83.4	2,640,598 711,417 684,308 326,671	84.5 84.5 <sup>^</sup> 78.0 <sup>^</sup> 57.1 <sup>^</sup>	82.2 72.6 66.2 31.0	86.6 91.8 86.6 79.8
Male Female	58,844	# 79.9	76.1	83.2	4,362,995	# 80.6	75.7	84.7

<sup>\*</sup> Pap test within past 3 years.

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>+</sup> Age-adjusted and age specific percent not generated for health indicators with age range limits.


<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

### Mohawk Valley

#### **Prevent Chronic Diseases**

Percentage of women aged 21-65 years with annual household income of less than \$25,000 receiving cervical cancer screening based on recent guidelines\*



		Mohawl	ς Valley		Statewide			
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
_	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	14,979	70.6	61.8	78.1	1,068,153	75.3^	63.2	84.4
Age <sup>+</sup>								
18-24								
25-34								
35-44								
45-64								
65+								
Race								
White, NH	12,453	71.0	61.8	78.6	433,489	74.8	68.6	80.2
Black, NH		#			283,693	93.7	87.8	96.8
Hispanic		#			292,551	77.4^	61.6	88.0
Other, NH		#			58,419	36.7	8.5	78.5
Sex								
Male		#				#		
Female	14,979	70.6	61.8	78.1	1,068,153	75.3^	63.2	84.4
Age-adjusted+,&								

<sup>\*</sup> Pap test within past 3 years.


<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>+</sup> Age-adjusted and age specific percent not generated for health indicators with age range limits.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines\*



		Mohawl	k Valley		Statewide				
	Est. # of		Lower	Upper	Est. # of		Lower	Upper	
Г	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit	
Total	50,394	64.0	59.5	68.2	3,471,945	69.3	66.3	72.2	
Age⁺									
18-24									
25-34									
35-44									
45-64									
65+									
Race									
White, NH	45,621	66.3	62.5	70.0	2,335,990	70.4	67.9	72.7	
Black, NH		#			514,345	69.6^	57.5	79.4	
Hispanic		#			416,087	70.6^	58.4	80.4	
Other, NH		#			205,524	57.4^	39.9	73.2	
Sex									
Male	24,385	63.8	56.4	70.5	1,545,645	66.8	62.1	71.2	
Female	26,009	64.2	58.7	69.3	1,926,301	71.5	67.5	75.2	
Age-adjusted+,&									

<sup>\*</sup> Blood stool test within 1 year, or sigmoidoscopy within 5 years with blood stool test within 3 years, or colonoscopy within 10 years.

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>+</sup> Age-adjusted and age specific percent not generated for health indicators with age range limits.


<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

 $<sup>^{\</sup>wedge}$  Highly variable rate (confidence interval with a half-width greater than 10).

### Mohawk Valley

#### **Prevent Chronic Diseases**

Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on recent guidelines\*



		Mohawl	ς Valley		Statewide			
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
F	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	11,737	54.0	47.0	60.7	736,005	61.4	54.7	67.7
Age⁺								
18-24								
25-34								
35-44								
45-64								
65+								
Race								
White, NH	10,335	56.2	49.1	63.0	345,064	57.6	51.4	63.7
Black, NH		#			155,775	60.0^	38.7	78.0
Hispanic		#			187,721	75.1^	60.9	85.4
Other, NH		#			47,444	52.8^	29.9	74.5
Sex								
Male	5,221	54.4	43.8	64.7	253,900	51.4	41.0	61.6
Female	6,516	53.6	44.5	62.5	482,104	68.5	60.2	75.7
Age-adjusted+,&								

<sup>\*</sup> Blood stool test within 1 year, or sigmoidoscopy within 5 years with blood stool test within 3 years, or colonoscopy within 10 years.


<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>+</sup> Age-adjusted and age specific percent not generated for health indicators with age range limits.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

### Percentage of adults who had a test for high blood sugar or diabetes within the past three years




		Mohawk	र Valley		Statewide			
_	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total	105,935	59.8	56.1	63.5	7,849,939	58.7	56.0	61.5
Age								
18-24	7,570	31.3^	19.1	46.8	503,420	26.8	20.2	34.6
25-34	16,024	55.2^	44.5	65.4	1,337,915	51.0	44.1	57.9
35-44	15,536	60.7^	50.3	70.2	1,475,455	63.3	56.2	69.9
45-64	43,443	68.2	63.4	72.7	2,822,324	68.0	64.3	71.4
65+	22,579	68.3	63.1	73.0	1,541,702	71.4	66.8	75.6
Race								
White, NH	94,138	59.8	56.1	63.5	4,823,563	61.0	58.6	63.4
Black, NH		#			1,240,630	65.1	55.4	73.7
Hispanic	5,108	46.7^	28.0	66.3	1,203,330	53.5	46.1	60.8
Other, NH		#			582,416	44.7	31.9	58.2
Sex								
Male	46,949	54.9	48.8	60.9	3,534,063	56.3	52.1	60.4
Female	58,985	64.4	60.0	68.6	4,315,875	60.9	57.2	64.5
Age-adjusted <sup>&amp;</sup>		59.4	55.6	63.2		59.1	56.6	61.6

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

#### Percentage of adults with physician-diagnosed prediabetes




		Mohawl	<ul><li>Valley</li></ul>		Statewide			
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	8,690	4.9	3.8	6.2	783,799	5.8	4.6	7.2
Age								
18-24		#			57,540	2.9	0.7	10.7
25-34		#			43,246	1.6	1.0	2.7
35-44		#			198,127	8.5	4.4	16.0
45-64	3,916	6.2	4.2	9.1	283,693	6.7	5.3	8.5
65+	2,933	8.7	6.3	11.8	171,261	7.8	6.2	9.9
Race								
White, NH	8,433	5.3	4.1	6.8	489,253	6.0	4.8	7.4
Black, NH		#			96,242	5.2	2.9	9.2
Hispanic		#			81,294	3.6	1.8	7.2
Other, NH		#			117,011	8.9^	3.2	22.3
Sex								
Male	3,986	4.6	3.0	6.8	377,968	5.9	4.1	8.5
Female	4,704	5.1	3.8	6.9	405,832	5.6	4.3	7.3
Age-adjusted <sup>&amp;</sup>		4.7	3.6	6.1		5.9	4.6	7.5

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

#### Percentage of adults with physician-diagnosed high blood pressure




	Mohawk Valley				Statewide			
_	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total Age	71,371	34.3	31.4	37.3	4,323,615	28.3	26.5	30.3
18-24		#			103,520	5.2	3.3	8.0
25-34	4,214	13.2	8.2	20.6	299,126	10.9	7.8	15.0
35-44	6,237	22.1	15.4	30.8	470,752	18.9	14.1	24.8
45-64	30,476	40.1	35.5	44.9	1,715,223	34.1	31.2	37.2
65+	28,845	62.9	58.2	67.3	1,654,516	60.4	56.5	64.2
Race								
White, NH	63,747	34.8	31.9	37.7	2,701,979	29.7	28.0	31.4
Black, NH		#			740,669	34.2	27.2	42.0
Hispanic	3,383	26.7^	14.8	43.5	609,542	23.8	18.7	29.9
Other, NH	3,562	39.7	19.3	64.5	271,424	18.9	12.7	27.1
Sex								
Male	35,130	34.8	30.3	39.6	2,116,294	29.0	26.2	32.1
Female	36,241	33.8	30.2	37.6	2,207,320	27.7	25.3	30.2
Age-adjusted <sup>&amp;</sup>		30.5	27.9	33.3		27.3	25.6	29.1

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

#### Percentage of adults with diagnosed high blood pressure taking high blood pressure medication




		Mohawl	( Valley		Statewide			
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
Ī	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	57,303	80.4	75.9	84.3	3,243,246	75.2	71.5	78.5
Age								
18-24		#			8,688	8.4	3.9	17.4
25-34		#			96,762	32.4^	18.9	49.5
35-44		#			186,971	39.7	26.7	54.3
45-64	24,611	80.8	74.4	85.8	1,395,152	81.6	77.8	84.9
65+	26,906	93.6	90.2	95.8	1,497,501	90.8	86.9	93.6
Race								
White, NH	51,214	80.5	75.8	84.5	2,081,796	77.4	74.1	80.3
Black, NH		#			552,688	74.6^	60.0	85.3
Hispanic		#			420,934	69.1^	56.3	79.5
Other, NH		#			187,828	69.2^	50.8	83.0
Sex								
Male	26,829	76.6	70.1	82.0	1,525,386	72.2	66.5	77.2
Female	30,474	84.2	77.4	89.3	1,717,859	78.1	73.2	82.4
Age-adjusted <sup>&amp;</sup>		59.6	53.7	65.3		55.6	51.1	60.0

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

#### Percentage of adults with current asthma\*



		Mohawl	( Valley		Statewide			
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
_	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	25,680	12.4	10.3	14.9	1,520,940	10.0	8.9	11.3
Age								
18-24	3,091	13.0^	6.2	25.2	170,340	8.6	6.2	11.8
25-34	3,952	12.5	7.8	19.4	322,447	11.7	8.2	16.4
35-44	5,118	18.2^	10.1	30.4	240,590	9.7	7.1	13.0
45-64	9,161	12.1	9.5	15.3	551,257	11.1	9.3	13.3
65+	4,273	9.3	6.9	12.4	222,952	8.1	6.5	10.1
Race								
White, NH	20,573	11.2	9.4	13.4	946,198	10.4	9.1	11.9
Black, NH		#			199,011	9.2	6.5	12.9
Hispanic	3,786	31.9^	15.4	54.6	283,809	11.3	7.9	16.0
Other, NH		#			91,921	6.4	4.0	10.0
Sex								
Male	8,771	8.7	5.8	12.8	504,039	7.0	5.6	8.6
Female	16,909	15.9	13.0	19.3	1,016,901	12.8	11.1	14.8
Age-adjusted <sup>&amp;</sup>		13.1	10.5	16.3		10.1	8.9	11.4


<sup>\*</sup> Physician diagnosed asthma.

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

 $<sup>^{\</sup>wedge}$  Highly variable rate (confidence interval with a half-width greater than 10).

#### Percent of adults with cholesterol checked\*



	Mohawk Valley				Statewide			
_	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total Age	172,485	84.3	81.1	87.0	12,572,076	84.2	81.7	86.3
18-24	8,988	37.6^	24.8	52.4	891,216	49.0	39.4	58.6
25-34	22,994	76.0	68.2	82.3	1,896,541	71.3	63.9	77.7
35-44	24,643	88.7	82.9	92.7	2,197,539	88.9	85.0	91.8
45-64	70,539	92.8	89.6	95.1	4,694,424	93.8	91.2	95.6
65+	44,081	97.7	95.9	98.8	2,641,873	97.2	95.8	98.2
Race								
White, NH	152,925	84.9	81.5	87.7	7,746,530	87.0	84.7	89.0
Black, NH		#			1,730,762	81.9	73.0	88.3
Hispanic	9,191	74.7	56.4	87.1	1,923,116	77.3	70.1	83.1
Other, NH	7,972	88.9^	75.7	95.3	1,171,667	81.9^	67.3	90.8
Sex								
Male	79,506	80.4	74.9	85.0	5,864,748	82.0	78.2	85.2
Female	92,979	87.9	84.6	90.6	6,707,327	86.2	82.9	88.9
Age-adjusted <sup>&amp;</sup>		82.6	79.8	85.0		83.4	81.2	85.4


<sup>\*</sup> As diagnosed by a health professional.

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

 $<sup>^{\</sup>wedge}$  Highly variable rate (confidence interval with a half-width greater than 10).

#### Percentage of adults 18 years and older with elevated cholesterol\*



		Mohawl	k Valley		Statewide			
_	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total Age	69,936	40.8	37.5	44.1	4,711,105	37.7	35.4	40.1
18-24		#			124,288	14.0	8.0	23.2
25-34	4,385	19.1^	11.6	29.9	435,898	23.0	16.9	30.5
35-44	6,830	27.7	19.8	37.3	620,994	28.3	22.0	35.6
45-64	32,804	47.0	42.0	52.0	2,009,889	43.1	39.8	46.4
65+	25,396	57.9	53.1	62.6	1,439,336	55.4	51.3	59.4
Race								
White, NH	61,740	40.6	37.5	43.8	3,102,928	40.3	38.2	42.4
Black, NH		#			561,511	33.1	25.6	41.6
Hispanic	4,290	46.7^	26.7	67.7	727,074	37.8	30.7	45.5
Other, NH		#			319,592	27.3	18.7	38.1
Sex								
Male	34,064	43.1	37.8	48.6	2,369,080	40.5	37.0	44.3
Female	35,873	38.8	34.8	42.8	2,342,025	35.2	32.4	38.2
Age-adjusted <sup>&amp;</sup>		33.6	30.6	36.7		34.5	32.1	37.0

<sup>\*</sup> As diagnosed by a health professional.


<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

# Mohawk Valley Prevent Chronic Diseases

Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition\*



		Mohawl	र Valley			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
_	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	12,472	10.8	8.8	13.2	713,905	10.2	8.5	12.2
Age								
18-24		#			11,969	2.9	1.4	6.1
25-34		#			137,115	17.9^	9.2	31.8
35-44		#			51,465	6.0	3.8	9.2
45-64	5,325	11.1	8.2	15.0	282,601	10.2	8.0	13.0
65+	4,606	12.6	9.6	16.3	208,847	10.1	8.0	12.7
Race								
White, NH	10,903	10.8	8.8	13.2	398,657	8.8	7.0	11.1
Black, NH		#			143,698	14.2	9.5	20.7
Hispanic		#			92,991	8.7	4.5	16.4
Other, NH		#			78,559	19.0^	10.4	32.2
Sex								
Male	5,467	10.2	7.5	13.8	324,668	10.2	7.7	13.4
Female	7,005	11.3	8.6	14.6	389,237	10.2	8.0	12.9
Age-adjusted <sup>&amp;</sup>		9.3	7.0	12.1		9.7	7.7	12.3

<sup>\*</sup> Courses for high blood pressure, arthritis, or diabetes.


<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

# Mohawk Valley Prevent Chronic Diseases

#### Percentage of adults with arthritis\*



		Mohawl	र Valley			State	wide	
_	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total Age	64,351	30.9	28.3	33.7	3,491,206	22.9	21.3	24.6
18-24		#			40,067	2.0	1.1	3.6
25-34	4,328	13.6	8.7	20.6	228,246	8.3	5.9	11.7
35-44	5,510	19.6	13.7	27.0	323,152	12.9	9.4	17.5
45-64	28,873	37.9	33.5	42.6	1,465,151	29.2	26.5	32.1
65+	24,702	53.9	49.0	58.8	1,353,867	49.3	45.4	53.2
Race								
White, NH	58,709	31.9	29.2	34.8	2,474,787	27.2	25.6	28.9
Black, NH		#			400,524	18.5	13.4	24.9
Hispanic	3,287	27.3^	15.1	44.2	460,563	18.0	13.6	23.4
Other, NH	1,541	17.4	8.3	33.0	155,331	10.8	7.2	15.8
Sex								
Male	27,291	27.1	23.2	31.3	1,258,898	17.3	15.2	19.6
Female	37,061	34.6	31.1	38.2	2,232,309	28.0	25.7	30.4
Age-adjusted <sup>&amp;</sup>		27.5	25.1	29.9		21.8	20.3	23.3

<sup>\*</sup> Includes rheumatoid arthritis, gout, lupus, or fibromyalgia.


<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

 $<sup>^{\</sup>wedge}$  Highly variable rate (confidence interval with a half-width greater than 10).

#### Promote a Healthy and Safe Environment

Percentage of adults who consider their neighborhood suitable for walking and physical activity



		Mohawl	( Valley			State	wide	95% limit  8 92.6  9 92.5  9 90.8  6 95.4  1 93.9  7 96.1  7 94.3  3 93.4  2 89.0  9 96.9	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper	
	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit	
Total	183,338	93.0	90.9	94.6	13,107,891	91.3	89.8	92.6	
Age									
18-24	19,425	87.4^	70.4	95.3	1,641,319	87.9	80.9	92.5	
25-34	28,290	91.3	86.1	94.7	2,218,379	86.6	80.9	90.8	
35-44	24,515	95.2	90.6	97.6	2,218,331	93.4	90.6	95.4	
45-64	68,724	93.4	90.8	95.3	4,372,743	92.2	90.1	93.9	
65+	41,118	94.8	92.7	96.4	2,447,810	94.7	92.7	96.1	
Race									
White, NH	162,230	93.0	90.8	94.7	8,055,698	93.1	91.7	94.3	
Black, NH		#			1,798,763	89.3	83.3	93.4	
Hispanic	10,248	89.2^	74.3	95.9	1,963,304	84.3	78.2	89.0	
Other, NH	7,643	94.2	84.8	98.0	1,290,125	94.0	88.9	96.9	
Sex									
Male	89,870	93.2	89.2	95.8	6,375,291	92.5	90.3	94.3	
Female	93,467	92.7	90.7	94.3	6,732,600	90.1	87.9	91.9	
Age-adjusted <sup>&amp;</sup>		92.9	90.6	94.6		91.3	89.8	92.6	


<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

#### Promote a Healthy and Safe Environment

#### Percentage of adults aged 65+ years with at least one fall in the past 12 months



		Mohawl	( Valley			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	13,908	31.5	27.3	36.1	802,383	30.8	27.3	34.5
Age <sup>+</sup>								
18-24								
25-34								
35-44								
45-64								
65+								
Race								
White, NH	13,643	33.7	29.4	38.3	600,059	31.3	28.3	34.5
Black, NH		#			61,485	22.9^	12.6	38.1
Hispanic		#			96,513	39.1^	21.8	59.7
Other, NH		#			44,326	25.2	12.2	45.0
Sex								
Male	6,493	31.5	24.6	39.3	324,764	30.4	25.5	35.7
Female	7,415	31.6	26.7	36.9	477,619	31.1	26.4	36.2
Age-adjusted+,&								

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.


<sup>+</sup> Age-adjusted and age specific percent not generated for health indicators with age range limits.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

#### Promote a Healthy and Safe Environments

Among adults aged 65+ years with at least one fall in the past 12 months, percent that resulted in injury\*



		Mohawl	ς Valley			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
<del>-</del>	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	5,044	36.4	28.7	44.9	304,376	38.3	32.2	44.9
Age⁺								
18-24								
25-34								
35-44								
45-64								
65+								
Race								
White, NH	4,935	36.3	28.5	44.9	223,708	37.6	32.1	43.4
Black, NH		#			33,030	53.7	28.5	77.2
Hispanic		#			31,933	34.1	11.7	66.8
Other, NH		#			15,705	35.4^	12.0	68.8
Sex								
Male	2,413	37.4	24.9	51.8	103,371	32.2	24.1	41.5
Female	2,632	35.6	26.9	45.3	201,005	42.5	33.9	51.5
Age-adjusted+,&								

<sup>\*</sup> Falls that limited regular activities or lead to doctor visits.

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.


<sup>+</sup> Age-adjusted and age specific percent not generated for health indicators with age range limits.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

 $<sup>^{\</sup>wedge}$  Highly variable rate (confidence interval with a half-width greater than 10).

#### Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections

#### Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV)



		Mohawl	( Valley			State	wide	Lower Upper	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper	
_	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit	
Total	24,095	33.8	29.3	38.6	1,662,747	37.3	34.0	40.8	
Age⁺									
18-24									
25-34									
35-44									
45-64									
65+									
Race									
White, NH	21,229	34.4	30.1	38.9	957,863	33.7	31.0	36.4	
Black, NH		#			347,642	52.3^	40.1	64.2	
Hispanic		#			270,194	44.2^	31.2	58.1	
Other, NH		#			87,049	26.2	15.7	40.4	
Sex									
Male	12,842	36.2	28.9	44.2	929,342	43.4	38.2	48.7	
Female	11,253	31.4	26.2	37.1	733,405	31.7	27.6	36.1	
Age-adjusted <sup>+,&amp;</sup>									

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.


<sup>+</sup> Age-adjusted and age specific percent not generated for health indicators with age range limits.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

#### Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections

Percentage of adults aged 18+ years with flu immunization in the past year\*



		Mohawl	र Valley			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
<b>-</b>	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	95,697	48.2	44.8	51.6	6,798,706	46.7	44.3	49.2
Age								
18-24	8,112	37.2	23.5	53.3	600,414	31.7	23.4	41.5
25-34	12,248	39.5^	30.2	49.6	907,897	35.3	29.0	42.1
35-44	10,337	39.7	30.7	49.5	981,422	41.1	33.9	48.6
45-64	33,478	45.3	40.5	50.2	2,282,268	47.4	44.1	50.8
65+	30,607	69.0	63.8	73.8	1,895,586	72.1	67.9	75.9
Race								
White, NH	86,474	49.2	45.8	52.6	4,296,522	49.3	47.1	51.5
Black, NH		#			718,122	35.0	27.6	43.3
Hispanic	5,931	52.2^	33.2	70.6	971,941	40.8	34.1	47.8
Other, NH	1,646	20.2^	9.4	38.3	812,120	58.5	45.1	70.7
Sex								
Male	40,862	42.2	37.0	47.6	3,045,064	43.8	40.1	47.5
Female	54,834	53.9	49.8	58.0	3,753,643	49.5	46.2	52.8
Age-adjusted <sup>&amp;</sup>		46.0	42.3	49.8		46.0	43.4	48.6

<sup>\*</sup> Includes flu shot or flu spray


<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

#### Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections

Percentage of adults aged 65+ years with flu immunization in the past year\*



		Mohawl	ς Valley			State	wide	79.6 55.5 84.3 89.7 79.3 76.0	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper	
-	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit	
Total	30,607	69.0	63.8	73.8	1,895,586	72.1	67.9	75.9	
Age⁺									
18-24									
25-34									
35-44									
45-64									
65+									
Race									
White, NH	28,875	71.0	66.7	74.9	1,484,360	76.9	73.9	79.6	
Black, NH		#			103,016	38.4^	23.8	55.5	
Hispanic		#			170,589	69.0^	47.8	84.3	
Other, NH		#			137,621	75.2^	51.4	89.7	
Sex									
Male	13,326	64.3	55.0	72.7	802,715	73.9	67.7	79.3	
Female	17,281	73.2	68.0	77.8	1,092,871	70.8	65.0	76.0	
Age-adjusted+,&									

<sup>\*</sup> Includes flu shot or flu spray

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.


<sup>+</sup> Age-adjusted and age specific percent not generated for health indicators with age range limits.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

 $<sup>^{\</sup>wedge}$  Highly variable rate (confidence interval with a half-width greater than 10).

#### Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections

#### Percentage of adults aged 65+ years with pneumococcal immunization



		Mohawl	ς Valley			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
_	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	30,090	69.6	64.2	74.5	1,617,276	65.1	60.7	69.2
Age⁺								
18-24								
25-34								
35-44								
45-64								
65+								
Race								
White, NH	28,484	71.6	67.2	75.6	1,303,462	71.4	68.1	74.4
Black, NH		#			123,843	48.9^	30.2	67.8
Hispanic		#			104,466	45.7	27.5	65.3
Other, NH		#			85,505	48.1^	27.7	69.2
Sex								
Male	12,646	62.5	53.1	71.1	631,549	62.4	56.7	67.7
Female	17,444	75.7	70.5	80.2	985,727	66.9	60.5	72.7
Age-adjusted+,&								

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.


<sup>+</sup> Age-adjusted and age specific percent not generated for health indicators with age range limits.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

#### Promote Healthy Women, Infants, and Children

#### Percentage of women aged 18-44 years who had dental visit within the last year



		Mohawl	ς Valley			State	wide	Section   Sect		
	Est. # of		Lower	Upper	Est. # of		Lower	Upper		
-	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit		
Total	28,581	72.5	64.5	79.2	2,268,176	70.7	64.0	76.5		
Age⁺										
18-24										
25-34										
35-44										
45-64										
65+										
Race										
White, NH	25,670	73.8	65.6	80.6	1,155,302	73.5	68.1	78.3		
Black, NH		#			426,771	78.8^	61.7	89.5		
Hispanic		#			412,421	62.2^	48.3	74.4		
Other, NH		#			273,683	63.0^	30.7	86.8		
Sex										
Male		#				#				
Female	28,581	72.5	64.5	79.2	2,268,176	70.7	64.0	76.5		
Age-adjusted+,&										

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.


<sup>+</sup> Age-adjusted and age specific percent not generated for health indicators with age range limits.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

#### Promote Healthy Women, Infants, and Children

Women aged 18-44 years who saw a doctor for a routine checkup within the last year



		Mohawl	k Valley			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
<b>-</b>	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	32,547	74.6	67.3	80.7	2,615,375	72.5	66.5	77.8
Age⁺								
18-24								
25-34								
35-44								
45-64								
65+								
Race								
White, NH	28,057	72.6	64.7	79.3	1,220,281	70.1	65.1	74.6
Black, NH		#			512,270	83.8^	68.8	92.4
Hispanic		#			606,739	76.3^	64.1	85.3
Other, NH		#			276,085	60.1^	30.5	83.8
Sex								
Male		#				#		
Female	32,547	74.6	67.3	80.7	2,615,375	72.5	66.5	77.8
Age-adjusted+,&								

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.


<sup>+</sup> Age-adjusted and age specific percent not generated for health indicators with age range limits.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

#### Promote Healthy Women, Infants, and Children

#### Percentage of women aged 18-64 years with health care coverage



		Mohawl	<ul><li>Valley</li></ul>			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	72,708	88.7	85.3	91.4	5,351,099	87.2	83.4	90.2
Age⁺								
18-24								
25-34								
35-44								
45-64								
65+								
Race								
White, NH	65,234	89.6	86.0	92.4	3,178,979	91.6	90.0	93.0
Black, NH		#			916,108	93.3	88.7	96.1
Hispanic		#			807,553	76.0	66.5	83.5
Other, NH		#			448,460	71.8^	43.3	89.5
Sex								
Male		#				#		
Female	72,708	88.7	85.3	91.4	5,351,099	87.2	83.4	90.2
Age-adjusted+,&								

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.


<sup>+</sup> Age-adjusted and age specific percent not generated for health indicators with age range limits.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

#### Promote Healthy Women, Infants, and Children

Percentage of women aged 18-44 years whose health care provider discussed planning a healthy pregnancy



	Mohawk Valley				Statewide			
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
<del>-</del>	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	19,970	50.4	42.2	58.6	1,268,971	39.3	33.7	45.2
Age⁺								
18-24								
25-34								
35-44								
45-64								
65+								
Race								
White, NH	17,541	50.2	41.6	58.8	723,302	45.9	41.1	50.9
Black, NH		#			291,237	53.7	35.5	71.0
Hispanic		#			192,904	28.5	19.0	40.3
Other, NH		#			61,528	14.1^	6.5	27.9
Sex								
Male		#				#		
Female	19,970	50.4	42.2	58.6	1,268,971	39.3	33.7	45.2
Age-adjusted+,&								

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.


<sup>+</sup> Age-adjusted and age specific percent not generated for health indicators with age range limits.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

### Promote Mental Health and Prevention Substance Abuse

#### Percentage of adults with poor mental health for 14 or more days in the last month



	Mohawk Valley				Statewide			
_	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total	25,228	12.4	10.4	14.8	1,672,677	11.3	9.7	13.0
Age								
18-24	3,129	13.4^	6.6	25.3	186,862	9.5	6.7	13.1
25-34	4,401	13.9	8.9	21.2	418,716	15.7	10.2	23.4
35-44	3,323	12.0	7.5	18.7	259,463	10.5	7.0	15.4
45-64	11,249	15.1	11.6	19.4	587,164	12.1	10.3	14.2
65+	3,115	7.1	5.3	9.3	175,961	6.7	4.9	9.0
Race								
White, NH	19,994	11.1	9.2	13.4	1,072,012	12.0	10.2	14.2
Black, NH		#			237,227	11.2	7.0	17.5
Hispanic	2,924	25.2	12.2	44.8	264,437	11.0	7.1	16.6
Other, NH	1,611	18.0^	7.6	36.9	99,001	7.0	3.9	12.2
Sex								
Male	10,364	10.5	7.9	13.8	734,315	10.4	7.9	13.5
Female	14,864	14.2	11.3	17.8	938,362	12.1	10.2	14.2
Age-adjusted <sup>&amp;</sup>		12.6	10.4	15.2		11.1	9.6	13.0


<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

#### Promote Mental Health and Prevention Substance Abuse

#### Percentage of adult binge drinking during the past month\*



	Mohawk Valley				Statewide			
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total Age	33,798	16.9	14.2	20.0	2,413,447	16.9	14.9	19.1
18-24	7,077	29.2^	18.0	43.7	515,466	27.1	19.4	36.6
25-34	9,514	30.1^	20.7	41.4	633,963	25.0	18.9	32.3
35-44	5,720	21.9	15.6	29.9	509,131	22.6	17.2	29.1
45-64	9,475	13.1	10.3	16.4	656,578	13.7	11.7	16.0
65+	2,002	4.6	2.8	7.2	86,636	3.4	2.7	4.2
Race								
White, NH	32,077	18.2	15.2	21.5	1,649,594	19.1	17.0	21.5
Black, NH		#			324,364	16.3^	9.4	26.8
Hispanic		#			277,966	11.6	7.7	17.1
Other, NH		#			161,523	12.4	6.8	21.7
Sex								
Male	21,816	22.5	17.9	27.8	1,567,117	22.8	19.3	26.8
Female	11,982	11.7	9.0	15.0	846,330	11.4	9.6	13.4
Age-adjusted <sup>&amp;</sup>		18.7	15.8	22.1		17.7	15.6	20.0

 $<sup>^{</sup>st}$  Men who have five or more drinks , women having four or more drinks on one occasion


<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).

### Promote Mental Health and Prevention Substance Abuse

#### Percentage of cigarette smoking among adults who report poor mental health



	Mohawk Valley				Statewide			
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
<del>-</del>	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	11,899	49.3	40.1	58.4	519,245	31.8	24.1	40.6
Age								
18-24		#			64,468	35.6^	22.4	51.5
25-34		#			192,926	47.3^	24.8	70.8
35-44		#			70,340	27.6^	16.6	42.1
45-64	5,940	53.3^	38.9	67.2	177,977	31.2	24.0	39.5
65+		#			13,368	7.7	4.1	13.9
Race								
White, NH	9,319	49.2	40.0	58.6	423,246	39.9^	29.9	50.7
Black, NH		#			33,694	14.4^	7.7	25.3
Hispanic		#			45,968	18.1^	9.4	31.9
Other, NH		#			16,336	19.6^	9.4	36.5
Sex								
Male	4,770	46.4^	32.6	60.7	272,857	38.4^	24.1	55.0
Female	7,129	51.4	39.5	63.1	246,388	26.7	20.7	33.8
Age-adjusted <sup>&amp;</sup>		47.5	39.2	56.0		29.9	24.1	36.4

<sup>&</sup>amp; Percent is age adjusted to the 2000 United States Population.

<sup>#</sup> Rates with <10 in the numerator or <50 in the denominator are suppressed.

<sup>^</sup> Highly variable rate (confidence interval with a half-width greater than 10).