eBRFSS 2013-2014 Health Indicators: New York City

General Description	
Improve Health Status and Reduce Health Disparities	
Percentage of adults who did not receive medical care because of cost	
Percentage of adults who have a regular health care provider	
Percentage of adults who had a dentist visit within the past year	
Percentage of adults living with a disability	
Percentage of adults aged 18-64 years with healthcare coverage	
Percentage of adults aged 18-64 years who saw a doctor for a routine checkup within the last year	
Percentage of adults experiencing housing insecurity in the past 12 months	
Percentage of adults experiencing food insecurity in the past 12 months	
Percentage of adults with poor self-reported health	
Percentage of adults who report 14 or more days of poor physical health	
Prevent Chronic Diseases	
Percentage of adults who are obese	
Percentage of adults with an annual household income less than \$25,000 who are obese	
Percentage of adults living with a disability who are obese	
Percentage of adults overweight or obese	
Percentage of adults who participated in leisure time physical activity in the past 30 days	
Percentage of adults who consume one or more sugary drinks daily	
Percentage of adults who consume fast-food three or more times per week	
Percentage of adults with physician-diagnosed diabetes	
Percentage of adults who are current smokers	
Percentage of adults with annual household income less than \$25,000 who are current smokers	
Percentage of adults aged 18-24 years who are current smokers	
Percentage of women aged 50-74 years receiving breast cancer screening based on recent guidelines	
Percentage of women aged 50-74 years with annual household income of less than \$25,000 receiving breast cancer screer	
guidelines	
Percentage of women aged 21-65 years receiving cervical cancer screening based on recent guidelines	
Percentage of women aged 21-65 years with annual household income of less than \$25,000 receiving cervical cancer scree	
guidelines	
Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines	20
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen	ning based on recent
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines	ning based on recent 29
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years	ning based on recent
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes	ning based on recent 29
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure	ning based on recent 29 30 31 32
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication	ning based on recent 29 30 31 32 33
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma	ning based on recent 29 30 31 32 33 34
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percent of adults with cholesterol checked	ning based on recent 29 30 31 32 33 34 34 35
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percent of adults with cholesterol checked Percentage of adults 18 years and older with elevated cholesterol	ning based on recent 29 30 31 32 33 34 34 35 36
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked Percentage of adults 18 years and older with elevated cholesterol Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition	ning based on recent 29 30 31 32 33 34 34 35 36 37
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked Percentage of adults 18 years and older with elevated cholesterol Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults with arthritis	ning based on recent 29 30 31 32 33 34 35 36 37 38
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked Percentage of adults 18 years and older with elevated cholesterol Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults with arthritis Promote a Healthy and Safe Environment	ning based on recent 29 30 31 32 33 34 34 35 36 37 38 39
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked Percentage of adults 18 years and older with elevated cholesterol Percentage of adults with arthritis Percentage of adults with arthritis Promote a Healthy and Safe Environment Percentage of adults who consider their neighborhood suitable for walking and physical activity	ning based on recent 29 30 31 32 33 34 34 35 36 37 38 39 39 39
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked Percentage of adults 18 years and older with elevated cholesterol Percentage of adults with atken a course or class to learn how to manage their chronic disease or condition Percentage of adults with arthritis Promote a Healthy and Safe Environment Percentage of adults who consider their neighborhood suitable for walking and physical activity Percentage of adults aged 65+ years with at least one fall in the past 12 months	ning based on recent 29 30 31 32 33 34 35 36 37 38 39 39 39 40
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked Percentage of adults 18 years and older with elevated cholesterol Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults with arthritis Promote a Healthy and Safe Environment Percentage of adults who consider their neighborhood suitable for walking and physical activity Percentage of adults aged 65+ years with at least one fall in the past 12 months, percent that resulted in injury	ning based on recent 29 30 31 32 33 34 34 35 36 37 38 39 39 40 40 41
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with choiseterol checked Percentage of adults with choiseterol checked Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults with arthritis Promote a Healthy and Safe Environment Percentage of adults who consider their neighborhood suitable for walking and physical activity Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months. Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections.	ning based on recent 29 30 31 32 33 34 34 35 36 37 38 39 39 40 40 41
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines	ning based on recent 29 30 31 32 33 34 34 35 36 37 38 39 39 40 40 41 42 42
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with choiseterol checked Percentage of adults with choiseterol checked Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults with arthritis Promote a Healthy and Safe Environment Percentage of adults who consider their neighborhood suitable for walking and physical activity Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months. Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections.	ning based on recent 29 30 31 32 33 34 34 35 36 37 38 39 39 40 40 41 42 42
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked Percentage of adults with cholesterol checked Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults with arthritis Promote a Healthy and Safe Environment Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 47-68 years reporting ever tested for Hepatitis C (HCV) Percentage of adults aged 18+ years with flu immunization in the past year	ning based on recent 29 30 31 32 33 34 35 36 37 38 39 39 40 40 41 42 42 43 44
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with constant adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults 18 years and older with elevated cholesterol Percentage of adults 18 years and older with elevated cholesterol Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults with arthritis Promote a Healthy and Safe Environment Percentage of adults who consider their neighborhood suitable for walking and physical activity Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months. Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV). Percentage of adults aged 18+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year	ning based on recent 29 30 31 32 33 34 34 35 36 36 37 38 39 40 40 41 42 42 42 43 44 45
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with colesterol checked Percentage of adults 18 years and older with elevated cholesterol Percentage of adults with arthritis Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months, percent that resulted in injury Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV). Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in	ning based on recent 29 30 31 32 33 34 34 35 36 36 37 38 39 40 40 41 41 42 42 42 43 44 45 46
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines. Percentage of adults who had a test for high blood sugar or diabetes within the past three years. Percentage of adults with physician-diagnosed prediabetes Percentage of adults with diagnosed high blood pressure. Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percent of adults with cholesterol checked. Percentage of adults 18 years and older with elevated cholesterol. Percentage of adults with attritis Percentage of adults with arthritis Percentage of adults with arthritis Percentage of adults with arthritis Percentage of adults with arthritis Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults who consider their neighborhood suitable for walking and physical activity. Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months. Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV). Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV). Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization. Promote Healthy Women, Infants, and Children Percentage of women aged 18-44 years who had dental visit within the last year	ning based on recent 29 30 31 32 33 34 34 35 36 37 38 39 40 40 41 42 42 42 42 43 44 44 45 46 46
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked Percentage of adults with cholesterol checked Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults who consider their neighborhood suitable for walking and physical activity Percentage of adults who consider their neighborhood suitable for walking and physical activity Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months, percent that resulted in injury Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV) Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu mounization in the past year Percentage of adults aged 65+ years with flu mounization in the past year Percentage of adults aged 65+ years with flu mounization in the past year Percentage of adults aged 65+ years with flu mounization in the past year Percentage of women aged 18-44 years who had dental visit within the last year Women aged 18-44 years who saw a doctor for a rout	ning based on recent 29 30 31 32 33 34 34 35 36 37 38 39 40 40 41 42 42 42 42 42 42 43 44 44 45 46 46 46 47
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with hysician-diagnosed nigh blood pressure Percentage of adults with current asthma Percentage of adults with current asthma Percentage of adults 18 years and older with elevated cholesterol Percentage of adults with arthritis Percentage of adults with arthritis Promote a Healthy and Safe Environment Percentage of adults who consider their neighborhood suitable for walking and physical activity Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months. Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV). Percentage of adults aged 18+ years with flu immunization in the past year Percentage of adults aged 18+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of women aged 18-44 years who had dental visit within the last year Percentage of women aged 18-44 years who had dental visit within the last year Percentage of women aged 18-64 years with health care coverage	ning based on recent 29 30 31 32 33 34 34 35 36 37 38 39 39 40 40 41 42 42 42 42 42 42 42 43 44 44 45 46 46 47 48
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked Percentage of adults with cholesterol checked Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults who consider their neighborhood suitable for walking and physical activity Percentage of adults who consider their neighborhood suitable for walking and physical activity Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months, percent that resulted in injury Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV) Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu mounization in the past year Percentage of adults aged 65+ years with flu mounization in the past year Percentage of adults aged 65+ years with flu mounization in the past year Percentage of adults aged 65+ years with flu mounization in the past year Percentage of women aged 18-44 years who had dental visit within the last year Women aged 18-44 years who saw a doctor for a rout	ning based on recent 29 30 31 32 33 34 34 35 36 37 38 39 39 40 40 41 42 42 42 42 42 42 42 43 44 44 45 46 46 47 48
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with current asthma Percentage of adults with colesterol checked Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults who consider their neighborhood suitable for walking and physical activity Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months, percent that resulted in injury Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV). Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with peumococcal immunization. Promote Healthy Women, Infants, and Children Percentage of women aged 18-44 years who had dental visit within the last year Percentage of women aged 18-44 years who sheal the care coverage Percentage of women aged 18-44 years who had dental visit within the last year Percentage of women aged 18-44 years who had dental visit within the last year Percentage of women aged 18-44 years who sheal the care coverage Percentage of women aged 18-44 years who had dental visit within the last year Perce	ning based on recent 29 30 31 32 33 34 34 35 36 37 38 39 39 40 40 41 42 42 42 42 42 42 42 42 43 44 44 45 46 46 47 48 49 50
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines. Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed prediabetes Percentage of adults with diagnosed high blood pressure aking high blood pressure medication Percentage of adults with current astmma Percentage of adults with collesterol checked. Percentage of adults with arthritis Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults with arthritis Percentage of adults who nove taken a course or class to learn how to manage their chronic disease or condition Percentage of adults who nove taken a course or class to learn how to manage their chronic disease or condition Percentage of adults who consider their neighborhood suitable for walking and physical activity. Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months, percent that resulted in injury. Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections. Percentage of adults aged 45+ years with flu immunization in the past year Percentage of adults aged 45+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of women aged 18-64 years who had dental visit within the last year Percentage of women aged 18-64 years who had dental visit within the last year Percentage of women aged 18-64 years who had dental visit within the last year Percentage of women aged 18-64 years whose health care coverage Percentage of wome	ning based on recent 29 30 31 32 33 34 34 35 36 37 38 39 39 40 40 41 41 42 42 43 40 41 42 42 43 40 41 42 43 40 41 42 43 44 45 46 46 47 48 49 50 50 50
Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screen guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with current asthma Percentage of adults with colesterol checked Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults who consider their neighborhood suitable for walking and physical activity Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months, percent that resulted in injury Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV). Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with peumococcal immunization. Promote Healthy Women, Infants, and Children Percentage of women aged 18-44 years who had dental visit within the last year Percentage of women aged 18-44 years who sheal the care coverage Percentage of women aged 18-44 years who had dental visit within the last year Percentage of women aged 18-44 years who had dental visit within the last year Percentage of women aged 18-44 years who sheal the care coverage Percentage of women aged 18-44 years who sheal the care coverage Percentage of wome	ning based on recent 29 30 31 32 33 34 34 35 36 37 38 39 39 40 40 41 42 42 42 42 43 40 41 42 42 43 40 41 42 42 43 40 41 42 43 40 41 42 50 50 50 50 50

General Description

The New York State Expanded Behavioral Risk Factor Surveillance System 2013-2014 (2014 eBRFSS) was design to supplement the Center for Disease Control and Prevention (CDC) Behavioral Risk Factor Surveillance System (BRFSS), which is conducted annually in New York State. The purpose of the 2014 eBRFSS was to produce local information on key public health issues. Data for the project were collected from April 15, 2013 to May 10, 2014. The 2014 eBRFSS reached both households with landline telephones and households which only had cell phones. Like the annual BRFSS, New York State's 2014 eBRFSS was designed to be representative of the non-institutionalized adult household population, aged 18 years and older.

Reported Health Indicators

There are a total of 50 health indicators included in the report. The health indicators include 10 measures included in the New York State Department of Health's Prevention Agenda, NYS Health Innovation Plan (SHIP), Delivery System Reform Incentive Payment (DSRIP) Program, or the NYS Community Health Indicator Reports (CHIRS). An additional 40 important health indicators are also included. The indicators are organized by the Priority Area of the Prevention Agenda and are reported statewide, by region, and by county.

Data Collection Methodology, Sampling and Weighting

The data collection methods for the 2014 eBRFSS aligned with the CDC's standard annual BRFSS methods. The 2014 eBRFSS questionnaire used standard questions from the BRFSS. Survey interviews were conducted in both English and Spanish. There were 31,690 completed interviews.


The sampling plan for the eBRFSS was developed to result in valid health indicator estimates for New York City (NYC) and each of the 57 counties outside NYC. The sampling approach stratified landlines by county and implemented a disproportionate stratified sampling design that accounted for both listed-households and unlisted numbers. Cell phone households were accounted for through additional strata that covered contiguous groups of counties.

Data were weighted to represent the population using a methodology known as iterative proportional fitting (raking). Raking adjusted the data so that groups underrepresented in the sample were more accurately represented in the estimated rates. Raking also enabled surveys conducted by cell phone to be included in the weighting and to accurately match sample distributions of specified demographic characteristics of the population. The raking categories included in the eBRFSS were: age, gender, race/ethnicity, education, marital status, owner/renter status, telephone usage (i.e. landline and cell phone combinations), and county of residence.

Limitations of Data Use:

During data analysis, criteria were applied to suppress rates based on too few surveys to be reliable and to identify rates that are highly variable. Rates in which less than 10 respondents were in the numerator or less than 50 respondents were in the denominator were considered unreliable and suppressed from reporting. Rates considered to have high variability due to having confidence limits greater than $\pm 10\%$ are noted.

Percentage of adults who did not receive medical care because of cost^{*}




		New Yo	rk City			State	wide	
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total	1,016,292	15.6	11.6	20.7	1,994,210	13.1	11.2	15.3
Age								
18-24	145,946	17.1^	5.0	44.9	281,654	14.1^	7.5	25.1
25-34	281,073	19.5^	10.5	33.4	537,866	19.7	14.2	26.6
35-44	237,777	20.8^	11.7	34.2	423,154	17.0	12.3	23.0
45-64	259,718	13.6	9.0	20.0	596,074	11.9	9.8	14.3
65+		#			139,733	5.1	3.0	8.4
Race								
White, NH	294,197	12.5	7.0	21.5	959,531	10.5	8.7	12.7
Black, NH	256,235	16.0^	9.0	27.0	335,184	15.5	10.0	23.4
Hispanic	283,109	16.5	11.5	23.2	465,114	18.4	14.5	23.0
Other, NH	182,751	21.5^	7.6	47.6	234,381	16.3^	7.3	32.6
Sex								
Male	434,900	14.3	8.8	22.4	890,677	12.3	9.6	15.5
Female	581,392	16.7	11.4	23.9	1,103,533	13.8	11.2	16.9
Age-adjusted ^{&}		15.7	11.7	20.8		13.6	11.6	15.8

* Among people needing to see a doctor in the past 12 months

& Percent is age adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.


Percentage of adults who have a regular health care provider



		New Yo	rk City			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	5,438,543	83.8	79.5	87.3	12,905,860	84.9	83.1	86.6
Age								
18-24	676,933	82.3^	67.4	91.2	1,486,954	76.3	69.8	81.8
25-34	1,019,141	70.3^	58.6	79.8	1,983,479	72.3	66.1	77.7
35-44	950,005	84.9^	72.8	92.2	2,091,488	84.7	79.6	88.7
45-64	1,662,644	86.2	78.0	91.6	4,450,345	88.6	85.5	91.1
65+	1,008,026	96.4	92.5	98.3	2,640,269	96.0	94.7	97.1
Race								
White, NH	2,064,563	88.4	83.4	92.1	7,996,007	88.0	86.6	89.3
Black, NH	1,459,286	90.8	83.1	95.2	1,926,090	89.0	83.6	92.7
Hispanic	1,165,816	68.7^	57.9	77.7	1,725,153	68.6	61.2	75.1
Other, NH	748,879	87.9^	73.1	95.1	1,258,610	87.9	79.9	93.0
Sex								
Male	2,441,393	80.4	73.2	86.1	5,804,831	80.1	77.1	82.8
Female	2,997,151	86.7	81.3	90.7	7,101,029	89.3	87.0	91.3
Age-adjusted ^{&}		84.2	80.0	87.7		84.5	82.5	86.2

& Percent is age adjusted to the 2000 United States Population.


Percentage of adults who had a dentist visit within the past year



		New Yo	rk City			State	wide	
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total	4,015,108	67.1	61.2	72.5	9,774,786	69.5	66.9	72.0
Age								
18-24	358,306	48.4^	27.1	70.3	1,100,494	62.2^	51.0	72.2
25-34	881,381	66.6^	52.7	78.2	1,624,123	65.1	57.7	71.8
35-44	772,628	70.8^	56.3	82.0	1,687,707	71.9	65.2	77.8
45-64	1,290,999	71.6	63.2	78.6	3,421,923	72.6	69.3	75.7
65+	602,728	66.7^	56.6	75.4	1,743,883	69.8	66.0	73.4
Race								
White, NH	1,545,904	71.5	62.9	78.8	6,152,109	72.4	70.1	74.6
Black, NH	986,146	66.6^	54.4	77.0	1,309,282	66.6	57.4	74.7
Hispanic	961,412	61.6^	50.7	71.5	1,435,933	62.9	55.3	70.0
Other, NH	521,645	66.7^	42.7	84.4	877,462	66.5^	52.1	78.4
Sex								
Male	1,917,419	66.0	56.9	74.0	4,591,800	67.3	63.4	71.1
Female	2,097,689	68.2	60.4	75.1	5,182,986	71.6	68.1	74.8
Age-adjusted ^{&}		66.7	60.9	72.0		69.3	66.6	71.8

& Percent is age adjusted to the 2000 United States Population.

Percentage of adults living with a disability^{*}




		New Yo	rk City			State	wide	
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total	1,189,911	18.7	15.2	22.9	3,111,131	20.8	19.1	22.7
Age								
18-24		#			119,872	6.1	4.3	8.5
25-34	211,835	15.2^	6.9	30.3	426,647	16.0	10.8	23.2
35-44	76,132	7.0	3.5	13.5	294,471	12.2	9.7	15.3
45-64	445,926	23.6	17.7	30.8	1,195,176	24.2	21.6	27.1
65+	380,351	37.7	29.0	47.3	989,611	36.8	33.1	40.6
Race								
White, NH	575,166	24.9	18.3	32.9	2,187,495	24.4	22.4	26.6
Black, NH	215,117	13.7	9.1	20.3	332,469	15.8	11.9	20.7
Hispanic	305,162	18.3	12.2	26.5	420,342	17.0	12.7	22.5
Other, NH	94,466	11.7^	5.2	24.2	170,824	12.3	7.7	19.1
Sex								
Male	452,400	15.3	10.0	22.6	1,349,486	18.9	16.3	21.9
Female	737,512	21.8	17.3	27.0	1,761,645	22.6	20.4	24.9
Age-adjusted ^{&}		18.0	14.8	21.7		19.9	18.3	21.6

* Limited activities (physical, mental, or emotional problems); need of special equipment (e.g. cane, wheel chair).

& Percent is age adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults aged 18-64 years with healthcare coverage




		New Yo	ork City		Statewide			
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total	4,394,441	84.1	78.5	88.5	10,275,389	85.2	82.8	87.3
Age ⁺								
18-24								
25-34								
35-44								
45-64								
65+								
Race								
White, NH	1,590,086	91.3	86.7	94.4	6,192,415	90.1	88.8	91.3
Black, NH	1,206,115	89.9	81.1	94.9	1,606,586	87.9	82.0	92.1
Hispanic	1,176,018	79.3^	68.4	87.1	1,626,540	73.2	66.2	79.2
Other, NH	422,222	64.3^	38.1	84.0	849,849	74.5^	56.8	86.6
Sex								
Male	2,137,361	83.8	76.1	89.3	4,924,290	83.2	79.9	86.0
Female	2,257,081	84.5	75.4	90.6	5,351,099	87.2	83.4	90.2
Age-adjusted ^{+,&}								

& Percent is age adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Percentage of adults aged 18-64 years who saw a doctor for a routine checkup within the last year




		New Yo	rk City			State	wide	
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total	3,862,124	72.7	66.3	78.2	8,622,736	70.9	68.1	73.5
Age ⁺								
18-24								
25-34								
35-44								
45-64								
65+								
Race								
White, NH	1,150,429	66.0^	55.7	75.0	4,721,851	68.7	65.8	71.4
Black, NH	1,086,240	81.0^	70.3	88.5	1,477,782	80.8	73.1	86.6
Hispanic	1,110,968	74.7^	63.2	83.6	1,583,308	71.1	63.6	77.6
Other, NH	514,487	69.5^	43.9	86.8	839,795	68.4^	53.1	80.5
Sex								
Male	1,753,441	67.1	57.6	75.4	3,938,722	65.6	61.4	69.6
Female	2,108,683	78.1	69.3	84.9	4,684,014	76.0	72.3	79.4
Age-adjusted ^{+,&}								

& Percent is age adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Percentage of adults experiencing housing insecurity in the past 12 months*




		New Yo	rk City		Statewide			
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total	3,123,072	52.4	46.6	58.2	5,967,540	43.4	40.8	46.1
Age								
18-24	219,055	29.2^	12.7	53.9	487,620	30.7^	21.3	42.0
25-34	843,314	65.3^	52.9	75.9	1,403,277	57.7	50.8	64.3
35-44	608,717	55.6^	40.6	69.7	1,180,494	50.7	43.4	57.9
45-64	1,013,525	56.7	48.6	64.5	2,100,265	44.9	41.4	48.4
65+	366,521	39.7^	30.1	50.1	688,870	27.8	23.8	32.1
Race								
White, NH	985,685	45.7	37.9	53.8	3,030,019	36.7	34.3	39.2
Black, NH	828,366	56.4^	44.3	67.8	1,078,432	55.5	46.2	64.3
Hispanic	893,217	58.9^	48.0	68.9	1,268,822	57.3	49.7	64.6
Other, NH	415,804	50.9^	29.7	71.8	590,266	44.6^	31.2	58.9
Sex								
Male	1,381,454	48.9	40.1	57.7	2,630,128	40.1	36.1	44.2
Female	1,741,618	55.6	48.0	63.0	3,337,412	46.5	43.0	50.0
Age-adjusted ^{&}		51.6	45.9	57.2		43.7	41.0	46.5

* Worried or stressed about having enough money to pay for rent or mortgage.

& Percent is age adjusted to the 2000 United States Population.

Percentage of adults experiencing food insecurity in the past 12 months*




		New Yo	rk City			State	wide	
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
[
Total	2,322,404	37.4	32.1	43.0	4,211,702	29.0	26.6	31.5
Age								
18-24	95,066	11.5^	5.3	23.2	339,128	17.9	13.5	23.3
25-34	733,080	54.3^	41.7	66.4	1,154,171	44.8	37.7	52.1
35-44	422,905	38.5^	25.1	53.9	786,490	33.0	26.3	40.4
45-64	790,129	42.4	34.5	50.6	1,464,969	30.3	26.9	33.9
65+	239,789	25.1	17.5	34.8	405,259	15.6	12.5	19.3
Race								
White, NH	687,170	30.7	23.2	39.3	1,998,541	22.9	20.6	25.4
Black, NH	578,910	37.5^	27.3	48.8	765,155	37.2	29.3	45.7
Hispanic	753,522	46.7^	36.5	57.2	1,031,787	43.4	36.2	50.8
Other, NH	302,803	37.2^	20.3	58.1	416,220	30.4^	20.2	42.9
Sex								
Male	1,087,316	36.9	28.7	45.9	1,903,904	27.4	23.6	31.5
Female	1,235,088	37.8	31.3	44.9	2,307,798	30.5	27.5	33.6
Age-adjusted ^{&}		36.8	31.9	41.9		29.4	27.1	31.9

* Worried or stressed about having enough money to buy nutritious meals.

& Percent is age adjusted to the 2000 United States Population.

Percentage of adults with poor self-reported health




		New Yo	ork City		Statewide			
-	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total	299,000	4.6	3.1	6.7	680,314	4.5	3.7	5.4
Age								
18-24		#			20,778	1.0	0.5	2.3
25-34		#			38,067	1.4	0.8	2.5
35-44		#			58,460	2.4	1.5	3.9
45-64	132,629	7.0	3.7	12.9	300,529	6.1	4.5	8.1
65+	116,277	11.3	6.5	18.8	248,637	9.1	6.9	12.0
Race								
White, NH	75,286	3.2	1.9	5.5	365,138	4.0	3.5	4.7
Black, NH		#			58,105	2.7	1.6	4.7
Hispanic	170,026	9.9	5.6	17.0	213,536	8.4	5.3	13.2
Other, NH		#			43,535	3.0	1.4	6.4
Sex								
Male	99,384	3.3	1.5	7.1	298,899	4.1	3.1	5.5
Female	199,616	5.8	3.8	8.7	381,415	4.8	3.9	6.0
Age-adjusted ^{&}		4.8	3.3	6.9		4.3	3.6	5.1

& Percent is age adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults who report 14 or more days of poor physical health*




		New Yo	ork City			State	Statewide			
F	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit		
Total	676,652	11.1	8.6	14.1	1,717,941	11.8	10.5	13.1		
Age										
18-24		#			59,108	3.1	1.9	5.1		
25-34	71,459	5.1	2.3	10.9	184,980	7.0	4.8	10.0		
35-44		#			176,691	7.2	5.2	9.9		
45-64	314,891	17.5	12.1	24.8	763,670	15.8	13.5	18.5		
65+	207,554	22.4	15.3	31.5	491,091	19.1	16.1	22.5		
Race										
White, NH	213,107	9.4	6.5	13.5	1,043,432	11.8	10.7	13.0		
Black, NH	123,949	8.3	4.7	14.3	188,315	9.3	6.3	13.5		
Hispanic	278,199	18.1	11.7	26.9	372,059	16.0	11.5	21.9		
Other, NH		#			114,135	8.1	4.7	13.7		
Sex										
Male	241,944	8.5	5.4	13.3	722,008	10.3	8.7	12.3		
Female	434,708	13.2	9.8	17.7	995,933	13.0	11.3	14.9		
Age-adjusted ^{&}		11.0	8.6	13.9		11.3	10.1	12.5		

* Out of the past 30 days.

& Percent is age adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults who are obese*




		New Yo	ork City		Statewide			
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total	1,318,207	22.0	18.1	26.5	3,527,077	24.9	23.0	26.9
Age								
18-24		#			318,556	17.8	11.7	26.0
25-34	178,169	14.1	8.4	22.9	509,101	20.8	16.8	25.4
35-44	184,062	17.0 [^]	9.4	28.7	573,976	24.7	20.0	30.1
45-64	578,996	31.2	24.3	39.0	1,452,755	30.5	27.5	33.7
65+	214,044	21.6	14.3	31.1	615,010	23.4	20.1	27.0
Race								
White, NH	440,302	20.0	14.7	26.6	2,139,878	25.0	23.2	26.9
Black, NH	435,968	28.1	19.9	37.9	638,079	30.7	24.2	38.1
Hispanic	395,691	26.6^	18.0	37.4	622,984	28.0	21.8	35.1
Other, NH	46,245	6.1	2.5	14.3	126,136	9.6	6.3	14.4
Sex								
Male	598,759	20.5	14.8	27.6	1,782,686	25.3	22.5	28.3
Female	719,448	23.4	18.3	29.3	1,744,390	24.5	22.0	27.1
Age-adjusted ^{&}		21.6	17.7	26.1		24.6	22.7	26.6

* Obesity defined as a Body Mass Index (BMI) >=30. BMI is calculated based on weight and height.

& Percent is age adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults with an annual household income less than \$25,000 who are obese*




		New Yo	ork City			State	wide	
-	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total	463,087	25.4	18.5	33.7	1,029,079	28.7	24.7	33.0
Age								
18-24		#			88,578	21.3^	10.0	39.7
25-34		#			177,054	26.6^	17.0	39.0
35-44		#			165,333	28.8^	19.5	40.3
45-64	187,615	31.2^	20.2	44.9	386,736	34.8	28.0	42.1
65+		#			190,291	24.4	18.8	31.1
Race								
White, NH	124,996	26.7^	14.0	44.9	500,787	30.3	25.6	35.4
Black, NH	105,664	22.1^	11.8	37.5	165,021	25.3^	16.7	36.3
Hispanic	196,200	29.3^	17.5	44.7	287,948	30.1^	21.1	41.0
Other, NH		#			75,322	23.1^	12.4	39.0
Sex								
Male	144,457	18.2^	9.3	32.6	386,764	23.8	18.1	30.5
Female	318,630	30.9^	22.1	41.3	642,314	32.7	27.6	38.3
Age-adjusted ^{&}		24.5	17.5	33.1		28.5	24.3	33.0

* Obesity defined as a Body Mass Index (BMI) >=30. BMI is calculated based on weight and height.

& Percent is age adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults living with a disability who are obese*




		New Yo	rk City			State	wide	
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total	404,395	35.7^	25.9	46.8	1,093,091	36.9	32.7	41.3
Age								
18-24		#			18,440	16.5^	9.8	26.5
25-34		#			122,752	31.0^	16.7	50.1
35-44		#			109,843	39.7^	30.4	49.8
45-64	223,413	50.1^	34.9	65.3	535,031	45.9	39.5	52.5
65+	88,569	25.5^	14.5	40.7	283,502	30.3	25.2	36.1
Race								
White, NH	144,544	25.9^	14.3	42.2	698,173	33.3	29.0	37.8
Black, NH		#			188,077	58.3^	44.8	70.6
Hispanic		#			170,166	46.0^	30.4	62.5
Other, NH		#			36,674	21.7^	12.1	35.9
Sex								
Male	119,028	26.6^	12.5	48.0	430,886	32.6	26.2	39.7
Female	285,367	41.5^	30.2	53.8	662,205	40.4	35.1	45.8
Age-adjusted ^{&}		30.5^	21.5	41.3		35.4	31.0	40.0

* Obesity defined as a Body Mass Index (BMI) >=30. BMI is calculated based on weight and height. Limited activities (physical, mental, or emotional problems); need of special equipment (e.g. cane, wheel chair).

& Percent is age adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults overweight or obese*




		New Yo	rk City			State	wide	
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total	3,543,794	59.0	53.4	64.5	8,631,360	60.9	58.4	63.3
Age								
18-24	249,054	35.4^	19.0	56.2	656,771	36.6	28.8	45.2
25-34	818,604	65.0^	52.9	75.4	1,528,245	62.4	55.9	68.4
35-44	527,506	48.6^	34.3	63.1	1,362,541	58.6	50.8	66.0
45-64	1,314,112	70.8	62.9	77.6	3,335,153	70.0	66.8	73.1
65+	567,753	57.2	47.4	66.4	1,632,723	62.1	58.1	65.9
Race								
White, NH	1,328,856	60.4	53.2	67.1	5,260,942	61.5	59.3	63.5
Black, NH	1,138,694	73.3^	60.6	83.0	1,520,031	73.1	63.8	80.7
Hispanic	838,571	56.4^	45.7	66.5	1,346,413	60.5	52.9	67.5
Other, NH	237,673	31.3^	17.2	49.9	503,974	38.4	28.1	49.9
Sex								
Male	1,837,525	62.8	53.9	70.9	4,736,847	67.1	63.2	70.8
Female	1,706,269	55.5	48.3	62.4	3,894,512	54.7	51.4	57.9
Age-adjusted ^{&}		58.0	52.6	63.2		60.5	57.9	63.0

* Defined as a Body Mass Index (BMI) >=25. BMI is calculated based on weight and height.

& Percent is age adjusted to the 2000 United States Population.

Percentage of adults who participated in leisure time physical activity in the past 30 days*




		New Yo	rk City			State	wide	
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total	4,481,891	71.6	66.6	76.1	10,681,161	72.8	70.6	74.9
Age								
18-24	694,203	84.3^	70.5	92.4	1,601,630	84.1	78.5	88.5
25-34	1,002,189	73.6^	62.2	82.6	1,900,064	73.1	67.1	78.4
35-44	764,298	69.4^	52.7	82.1	1,722,858	71.5	63.8	78.1
45-64	1,220,635	65.3	57.4	72.4	3,443,881	70.8	67.5	73.8
65+	721,195	73.4	64.9	80.6	1,840,519	69.5	66.1	72.7
Race								
White, NH	1,743,865	77.0	70.9	82.1	6,662,678	75.8	74.0	77.4
Black, NH	1,163,722	75.3	66.0	82.7	1,525,317	73.8	66.8	79.7
Hispanic	1,009,996	62.3^	52.1	71.5	1,507,430	62.4	55.4	68.9
Other, NH	564,307	68.1^	45.5	84.5	985,735	70.6^	56.9	81.3
Sex								
Male	2,415,145	81.3	75.2	86.2	5,439,883	77.3	74.6	79.8
Female	2,066,746	62.8	55.4	69.6	5,241,278	68.6	65.3	71.7
Age-adjusted ^{&}		71.5	66.5	76.1		72.9	70.5	75.1

* Any physical activities or exercises such as running or walking

& Percent is age adjusted to the 2000 United States Population.

Percentage of adults who consume one or more sugary drinks daily*




		New Yo	rk City			State	wide	
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total	1,495,929	24.6	19.5	30.4	3,402,766	23.8	21.5	26.3
Age								
18-24	258,019	34.0^	16.5	57.2	670,630	36.9	28.0	46.7
25-34	343,173	25.7^	14.9	40.6	733,289	28.9	22.5	36.4
35-44	390,979	35.8^	22.1	52.2	769,136	32.4	25.4	40.3
45-64	343,525	18.6	12.9	26.1	839,390	17.6	15.0	20.5
65+	142,871	15.3	9.2	24.4	348,734	13.7	11.1	16.9
Race								
White, NH	411,964	18.8	12.0	28.3	1,754,487	20.4	18.3	22.8
Black, NH	486,679	32.5^	22.0	45.1	671,592	33.6	25.4	43.0
Hispanic	439,189	27.6^	19.3	37.8	700,337	29.8	23.7	36.8
Other, NH	158,098	19.6^	6.3	46.9	276,350	20.4^	10.6	35.5
Sex								
Male	631,827	21.6	14.5	31.0	1,782,648	26.0	22.5	29.8
Female	864,102	27.3	20.7	35.2	1,620,118	21.8	18.7	25.3
Age-adjusted ^{&}		25.1	20.0	31.0		24.7	22.2	27.4

* Sugary drinks include regular soda, and sugar-sweetened drinks.

& Percent is age adjusted to the 2000 United States Population.

Percentage of adults who consume fast-food three or more times per week




		New Yo	ork City			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
г	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	301,967	5.1	3.2	7.9	817,815	5.8	4.8	6.9
Age								
18-24	54,000	7.5^	3.0	17.9	172,464	9.7	6.9	13.6
25-34	128,251	9.7	4.9	18.1	252,437	10.0	6.9	14.2
35-44		#			106,397	4.5	3.2	6.3
45-64	96,308	5.4	2.1	13.2	234,582	5.0	3.3	7.4
65+		#			47,255	1.9	1.3	2.7
Race								
White, NH	94,579	4.4	2.6	7.5	433,016	5.1	4.4	5.9
Black, NH		#			122,943	6.2	4.1	9.3
Hispanic	142,628	9.2^	4.1	19.5	219,779	9.6	5.6	16.0
Other, NH		#			42,077	3.2	1.8	5.5
Sex								
Male	181,707	6.3	3.3	11.6	560,514	8.2	6.6	10.3
Female	120,260	3.9	2.1	7.1	257,301	3.5	2.6	4.8
Age-adjusted ^{&}		4.8	3.0	7.5		5.9	4.9	7.0

& Percent is age adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults with physician-diagnosed diabetes*




		New Yo	ork City			State	wide	
-	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total	653,301	10.0	7.7	12.9	1,456,783	9.5	8.4	10.8
Age								
18-24		#			8,338	0.4	0.2	0.8
25-34		#			58,319	2.1	0.8	5.5
35-44		#			111,205	4.5	2.5	7.7
45-64	351,344	18.2	13.1	24.8	714,501	14.2	11.9	16.9
65+	181,994	17.4	11.3	25.8	525,500	19.1	16.3	22.2
Race								
White, NH	193,890	8.3	5.5	12.1	781,028	8.6	7.6	9.6
Black, NH	165,796	10.3	6.2	16.8	250,921	11.6	8.2	16.2
Hispanic	246,202	14.1	8.7	22.2	296,635	11.6	7.7	17.0
Other, NH	47,413	5.6	2.2	13.2	128,200	8.9	5.4	14.4
Sex								
Male	361,788	11.8	8.0	17.0	795,054	10.9	9.0	13.1
Female	291,513	8.4	5.9	11.8	661,729	8.3	7.0	9.7
Age-adjusted ^{&}		10.0	7.7	12.8		8.9	7.9	10.1

* As diagnosed by a health professional.

& Percent is age adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.


Percentage of adults who are current smokers



		New Yo	ork City			State	wide	
-	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total Age	836,449	13.3	9.9	17.6	2,311,528	15.6	14.0	17.4
18-24	31,258	3.7	1.5	8.5	254,033	13.0	9.8	17.0
25-34	224,069	16.0^	7.4	31.2	524,783	19.7	14.3	26.7
35-44	171,907	15.5^	7.8	28.4	427,187	17.6	13.3	23.0
45-64	377,632	20.3	14.5	27.8	937,977	19.2	16.7	22.1
65+	28,750	3.0	1.5	6.1	149,820	5.7	4.7	6.8
Race								
White, NH	430,870	19.2	12.6	28.2	1,598,923	18.1	16.0	20.3
Black, NH	186,650	11.9	6.1	21.8	314,837	15.0	10.0	21.7
Hispanic	177,985	10.8	6.2	18.3	303,908	12.4	8.8	17.1
Other, NH	40,944	4.9	2.0	11.4	93,860	6.7	4.3	10.3
Sex								
Male	520,119	17.6	11.6	25.9	1,331,531	18.8	15.9	22.1
Female	316,330	9.5	6.5	13.5	979,997	12.7	11.1	14.4
Age-adjusted ^{&}		13.4	10.1	17.5		15.9	14.1	17.7

& Percent is age adjusted to the 2000 United States Population.

Percentage of adults with annual household income less than \$25,000 who are current smokers




		New Yo	ork City			State	wide	
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total	389,522	19.6^	12.1	30.2	910,061	23.6	19.2	28.7
Age								
18-24		#			84,893	16.1	9.8	25.4
25-34		#			254,993	32.7^	18.8	50.6
35-44		#			141,275	24.5	15.5	36.5
45-64	157,351	27.1^	15.8	42.4	354,621	31.5	24.7	39.1
65+		#			63,044	7.9	5.6	11.0
Race								
White, NH	193,369	40.1^	20.3	63.8	562,213	32.6	25.7	40.3
Black, NH	107,914	22.6^	10.4	42.3	187,054	28.1^	17.8	41.4
Hispanic	72,201	9.8	4.6	19.4	119,918	11.3	7.1	17.6
Other, NH		#			40,876	10.0	5.0	19.2
Sex								
Male	234,120	28.1^	14.0	48.5	507,987	30.0	21.9	39.5
Female	155,402	13.5	7.6	22.8	402,073	18.6	14.7	23.3
Age-adjusted ^{&}		19.5	12.9	28.4		24.2	20.0	28.9

& Percent is age adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Prevent Chronic Diseases

Percentage of adults aged 18-24 years who are current smokers




		New Yo	ork City			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
F	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	31,258	3.7	1.5	8.5	254,033	13.0	9.8	17.0
Age⁺								
18-24								
25-34								
35-44								
45-64								
65+								
Race								
White, NH		#			188,030	19.4	14.8	25.2
Black, NH		#			21,086	6.1	2.8	13.1
Hispanic		#			33,121	9.8^	4.6	19.9
Other, NH		#			11,797	3.8	1.4	10.0
Sex								
Male	27,012	6.8^	2.4	17.8	153,298	15.5	10.7	21.8
Female		#			100,736	10.4	6.6	16.0
Age-adjusted ^{+,&}								

& Percent is age adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of women aged 50-74 years receiving breast cancer screening based on recent guidelines^{*}



		New Yo	ork City			State	wide	
F	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total Age ⁺ 18-24 25-34 35-44	912,405	81.3	73.7	87.1	2,225,651	80.9	77.7	83.7
45-64 65+ Race White, NH Black, NH Hispanic Other, NH Sex	289,125	71.6 [^] # # #	59.2	81.4	1,417,374 347,690 309,667 150,920	79.0 87.0 [°] 83.8 [°] 79.7 [°]	75.9 73.4 70.9 61.2	81.8 94.3 91.6 90.7
Male Female Age-adjusted ^{+,&}	912,405	# 81.3	73.7	87.1	2,225,651	# 80.9	77.7	83.7

* Mammogram within past 2 years.


& Percent is age adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Prevent Chronic Diseases

Percentage of women aged 50-74 years with annual household income of less than \$25,000 receiving breast cancer screening based on recent guidelines^{*}



		New Yo	ork City			Adults Percent 95% limit 95% limit 554,880 76.7 69.6 82.6 232,138 66.5 57.9 74.1 139,053 95.4 89.6 98.1 134,524 82.0^ 61.8 92.7		
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	313,114	80.2^	66.6	89.1	554,880	76.7	69.6	82.6
Age⁺								
18-24								
25-34								
35-44								
45-64								
65+								
Race								
White, NH		#			232,138	66.5	57.9	74.1
Black, NH		#			139,053	95.4	89.6	98.1
Hispanic		#			134,524	82.0^	61.8	92.7
Other, NH		#			49,165	76.3^	44.1	93.0
Sex								
Male		#				#		
Female	313,114	80.2^	66.6	89.1	554,880	76.7	69.6	82.6
Age-adjusted ^{+,&}								


* Mammogram within past 2 years.

& Percent is age adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of women aged 21-65 years receiving cervical cancer screening based on recent guidelines^{*}



		New Yo	ork City		Statewide			
	Est. # of	_	Lower	Upper	Est. # of		Lower	Upper
	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	1,775,281	76.3^	65.0	84.8	4,362,995	80.6	75.7	84.7
Age ⁺								
18-24								
25-34								
35-44								
45-64								
65+								
Race								
White, NH	597,667	84.4	74.8	90.8	2,640,598	84.5	82.2	86.6
Black, NH	544,993	85.5^	68.5	94.1	711,417	84.5^	72.6	91.8
Hispanic	468,713	76.3^	59.6	87.5	684,308	78.0^	66.2	86.6
Other, NH		#			326,671	57.1^	31.0	79.8
Sex								
Male		#				#		
Female	1,775,281	76.3^	65.0	84.8	4,362,995	80.6	75.7	84.7
Age-adjusted ^{+,&}								

* Pap test within past 3 years.


& Percent is age adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Prevent Chronic Diseases

Percentage of women aged 21-65 years with annual household income of less than \$25,000 receiving cervical cancer screening based on recent guidelines^{*}



		New York City				Statewide			
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	
Total Age⁺	584,197	75.1^	52.0	89.3	1,068,153	75.3^	63.2	84.4	
18-24									
25-34									
35-44									
45-64									
65+									
Race									
White, NH		#			433,489	74.8	68.6	80.2	
Black, NH		#			283,693	93.7	87.8	96.8	
Hispanic		#			292,551	77.4^	61.6	88.0	
Other, NH		#			58,419	36.7^	8.5	78.5	
Sex									
Male		#				#			
Female	584,197	75.1^	52.0	89.3	1,068,153	75.3^	63.2	84.4	
Age-adjusted ^{+,&}									

* Pap test within past 3 years.


& Percent is age adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Prevent Chronic Diseases

Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines^{*}



		New Yo	ork City		Statewide			
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	1,292,027	68.2	60.7	74.9	3,471,945	69.3	66.3	72.2
Age⁺								
18-24								
25-34								
35-44								
45-64								
65+								
Race								
White, NH	517,970	70.5	61.4	78.2	2,335,990	70.4	67.9	72.7
Black, NH	391,561	67.6^	52.3	79.9	514,345	69.6^	57.5	79.4
Hispanic	330,041	74.6^	57.8	86.3	416,087	70.6^	58.4	80.4
Other, NH		#			205,524	57.4^	39.9	73.2
Sex								
Male	529,863	62.6^	50.7	73.2	1,545,645	66.8	62.1	71.2
Female	762,163	72.7^	62.6	80.9	1,926,301	71.5	67.5	75.2
Age-adjusted ^{+,&}								

* Blood stool test within 1 year, or sigmoidoscopy within 5 years with blood stool test within 3 years, or colonoscopy within 10 years.


& Percent is age adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Prevent Chronic Diseases

Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on recent guidelines^{*}



		New Yo	ork City		Statewide				
	Est. # of		Lower	Upper	Est. # of	_	Lower	Upper	
r	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit	
Total	368,639	62.9^	49.6	74.5	736,005	61.4	54.7	67.7	
Age⁺									
18-24									
25-34									
35-44									
45-64									
65+									
Race									
White, NH		#			345,064	57.6	51.4	63.7	
Black, NH		#			155,775	60.0^	38.7	78.0	
Hispanic		#			187,721	75.1^	60.9	85.4	
Other, NH		#			47,444	52.8^	29.9	74.5	
Sex									
Male		#			253,900	51.4^	41.0	61.6	
Female	254,514	69.9^	53.7	82.2	482,104	68.5	60.2	75.7	
Age-adjusted ^{+,&}									


* Blood stool test within 1 year, or sigmoidoscopy within 5 years with blood stool test within 3 years, or colonoscopy within 10 years.

& Percent is age adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.


Percentage of adults who had a test for high blood sugar or diabetes within the past three years



		New Yo	rk City		Statewide			
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	3,444,625	59.9	53.7	65.8	7,849,939	58.7	56.0	61.5
Age								
18-24	204,001	25.2^	13.0	43.1	503,420	26.8	20.2	34.6
25-34	743,677	53.0^	40.5	65.1	1,337,915	51.0	44.1	57.9
35-44	771,763	72.5^	55.7	84.7	1,475,455	63.3	56.2	69.9
45-64	1,055,744	69.1	59.5	77.3	2,822,324	68.0	64.3	71.4
65+	588,774	69.5^	58.2	78.9	1,541,702	71.4	66.8	75.6
Race								
White, NH	1,282,389	63.7	55.1	71.5	4,823,563	61.0	58.6	63.4
Black, NH	951,629	66.0^	53.1	77.0	1,240,630	65.1	55.4	73.7
Hispanic	859,304	57.6^	46.6	67.9	1,203,330	53.5	46.1	60.8
Other, NH	351,303	43.7^	24.8	64.7	582,416	44.7^	31.9	58.2
Sex								
Male	1,535,477	58.2	48.5	67.2	3,534,063	56.3	52.1	60.4
Female	1,909,148	61.4	53.3	68.9	4,315,875	60.9	57.2	64.5
Age-adjusted ^{&}		61.3	55.7	66.6		59.1	56.6	61.6

& Percent is age adjusted to the 2000 United States Population.

Percentage of adults with physician-diagnosed prediabetes




		New Yo	ork City		Statewide			
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total	391,603	6.8	4.4	10.4	783,799	5.8	4.6	7.2
Age								
18-24		#			57,540	2.9	0.7	10.7
25-34		#			43,246	1.6	1.0	2.7
35-44		#			198,127	8.5	4.4	16.0
45-64	113,167	7.3	4.3	12.3	283,693	6.7	5.3	8.5
65+	62,557	7.3	4.0	13.0	171,261	7.8	6.2	9.9
Race								
White, NH	174,254	8.2	4.6	14.1	489,253	6.0	4.8	7.4
Black, NH	75,444	5.5	2.6	11.2	96,242	5.2	2.9	9.2
Hispanic		#			81,294	3.6	1.8	7.2
Other, NH		#			117,011	8.9^	3.2	22.3
Sex								
Male	185,072	7.1	3.4	14.3	377,968	5.9	4.1	8.5
Female	206,531	6.5	3.9	10.7	405,832	5.6	4.3	7.3
Age-adjusted ^{&}		7.3	4.6	11.2		5.9	4.6	7.5

& Percent is age adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults with physician-diagnosed high blood pressure




		New Yo	ork City		Statewide			
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total	1,688,419	25.8	21.9	30.2	4,323,615	28.3	26.5	30.3
Age								
18-24		#			103,520	5.2	3.3	8.0
25-34	127,335	8.8	4.2	17.5	299,126	10.9	7.8	15.0
35-44	233,767	20.5	11.4	34.1	470,752	18.9	14.1	24.8
45-64	616,201	31.9	25.3	39.4	1,715,223	34.1	31.2	37.2
65+	639,414	61.9	52.0	70.9	1,654,516	60.4	56.5	64.2
Race								
White, NH	578,870	24.8	19.7	30.7	2,701,979	29.7	28.0	31.4
Black, NH	525,071	32.7^	23.7	43.1	740,669	34.2	27.2	42.0
Hispanic	441,996	25.4	18.1	34.3	609,542	23.8	18.7	29.9
Other, NH	142,482	16.7^	8.3	30.8	271,424	18.9	12.7	27.1
Sex								
Male	758,744	24.8	19.0	31.8	2,116,294	29.0	26.2	32.1
Female	929,676	26.7	21.7	32.4	2,207,320	27.7	25.3	30.2
Age-adjusted ^{&}		26.7	23.0	30.7		27.3	25.6	29.1

& Percent is age adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults with diagnosed high blood pressure taking high blood pressure medication




		New Yo	rk City		Statewide			
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
r	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	1,239,506	73.4	64.3	80.9	3,243,246	75.2	71.5	78.5
Age								
18-24		#			8,688	8.4	3.9	17.4
25-34		#			96,762	32.4^	18.9	49.5
35-44		#			186,971	39.7^	26.7	54.3
45-64	541,074	87.8	78.2	93.5	1,395,152	81.6	77.8	84.9
65+	560,300	87.6^	76.8	93.8	1,497,501	90.8	86.9	93.6
Race								
White, NH	449,043	77.6^	63.2	87.5	2,081,796	77.4	74.1	80.3
Black, NH	378,585	72.1^	51.9	86.1	552,688	74.6^	60.0	85.3
Hispanic	325,588	73.7^	55.4	86.3	420,934	69.1^	56.3	79.5
Other, NH		#			187,828	69.2^	50.8	83.0
Sex								
Male	551,194	72.6^	56.9	84.2	1,525,386	72.2	66.5	77.2
Female	688,312	74.0^	62.7	82.9	1,717,859	78.1	73.2	82.4
Age-adjusted ^{&}		54.7	45.7	63.4		55.6	51.1	60.0

& Percent is age adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults with current asthma*




		New Yo	ork City		Statewide				
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	
Total	635,303	9.8	7.5	12.7	1,520,940	10.0	8.9	11.3	
Age									
18-24	40,154	4.7	1.6	12.9	170,340	8.6	6.2	11.8	
25-34	153,618	10.6	5.3	19.9	322,447	11.7	8.2	16.4	
35-44	98,600	8.6	4.3	16.6	240,590	9.7	7.1	13.0	
45-64	248,033	13.3	9.0	19.2	551,257	11.1	9.3	13.3	
65+	88,532	8.5	4.9	14.2	222,952	8.1	6.5	10.1	
Race									
White, NH	254,666	10.9	7.0	16.7	946,198	10.4	9.1	11.9	
Black, NH	122,641	7.6	4.5	12.7	199,011	9.2	6.5	12.9	
Hispanic	211,613	12.5	7.7	19.6	283,809	11.3	7.9	16.0	
Other, NH	46,383	5.4	2.4	11.8	91,921	6.4	4.0	10.0	
Sex									
Male	161,853	5.4	3.0	9.4	504,039	7.0	5.6	8.6	
Female	473,450	13.6	10.1	18.2	1,016,901	12.8	11.1	14.8	
Age-adjusted ^{&}		9.8	7.6	12.7		10.1	8.9	11.4	

* Physician diagnosed asthma.

& Percent is age adjusted to the 2000 United States Population.

Percent of adults with cholesterol checked*




		New Yo	rk City		Statewide			
	Est. # of	- .	Lower	Upper	Est. # of	- .	Lower	Upper
	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	5,341,000	83.3	77.5	87.8	12,572,076	84.2	81.7	86.3
Age								
18-24	386,451	50.2^	29.5	70.9	891,216	49.0	39.4	58.6
25-34	1,005,315	70.6^	56.7	81.5	1,896,541	71.3	63.9	77.7
35-44	1,032,479	90.6	81.3	95.5	2,197,539	88.9	85.0	91.8
45-64	1,787,414	92.8	85.3	96.6	4,694,424	93.8	91.2	95.6
65+	1,007,322	98.0	92.3	99.5	2,641,873	97.2	95.8	98.2
Race								
White, NH	2,024,384	87.3	77.6	93.2	7,746,530	87.0	84.7	89.0
Black, NH	1,279,746	81.8^	69.4	89.9	1,730,762	81.9	73.0	88.3
Hispanic	1,332,706	79.4^	68.4	87.3	1,923,116	77.3	70.1	83.1
Other, NH	704,165	82.7^	55.6	94.8	1,171,667	81.9^	67.3	90.8
Sex								
Male	2,446,287	81.4	72.2	88.0	5,864,748	82.0	78.2	85.2
Female	2,894,713	85.0	77.1	90.5	6,707,327	86.2	82.9	88.9
Age-adjusted ^{&}		83.7	78.9	87.5		83.4	81.2	85.4

* As diagnosed by a health professional.

& Percent is age adjusted to the 2000 United States Population.

Percentage of adults 18 years and older with elevated cholesterol*



		New Yo	rk City		Statewide				
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	
Total	1,955,143	36.8	31.8	42.1	4,711,105	37.7	35.4	40.1	
Age									
18-24		#			124,288	14.0	8.0	23.2	
25-34	272,447	27.1^	16.6	40.9	435,898	23.0	16.9	30.5	
35-44	302,584	29.3^	17.4	44.9	620,994	28.3	22.0	35.6	
45-64	747,531	42.0	34.5	49.9	2,009,889	43.1	39.8	46.4	
65+	534,890	54.3	44.3	64.0	1,439,336	55.4	51.3	59.4	
Race									
White, NH	857,215	42.4	35.6	49.5	3,102,928	40.3	38.2	42.4	
Black, NH	424,255	34.0^	24.1	45.5	561,511	33.1	25.6	41.6	
Hispanic	529,437	39.7^	29.9	50.4	727,074	37.8	30.7	45.5	
Other, NH	144,237	20.5^	9.9	37.6	319,592	27.3^	18.7	38.1	
Sex									
Male	947,507	38.7	30.9	47.2	2,369,080	40.5	37.0	44.3	
Female	1,007,637	35.2	29.2	41.8	2,342,025	35.2	32.4	38.2	
Age-adjusted ^{&}		34.9	29.9	40.3		34.5	32.1	37.0	


* As diagnosed by a health professional.

& Percent is age adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

New York City Prevent Chronic Diseases

Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition^{*}



		New Yo	ork City			State	wide	
-	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total	343,576	12.6	8.7	18.0	713,905	10.2	8.5	12.2
Age								
18-24		#			11,969	2.9	1.4	6.1
25-34		#			137,115	17.9^	9.2	31.8
35-44		#			51,465	6.0	3.8	9.2
45-64	129,039	12.5	7.4	20.3	282,601	10.2	8.0	13.0
65+	79,228	10.5	5.9	18.0	208,847	10.1	8.0	12.7
Race								
White, NH	122,839	11.7^	5.7	22.5	398,657	8.8	7.0	11.1
Black, NH	99,503	13.8	7.9	23.2	143,698	14.2	9.5	20.7
Hispanic	72,771	9.5^	4.0	20.7	92,991	8.7	4.5	16.4
Other, NH		#			78,559	19.0^	10.4	32.2
Sex								
Male	153,407	13.0	7.2	22.4	324,668	10.2	7.7	13.4
Female	190,169	12.4	7.6	19.4	389,237	10.2	8.0	12.9
Age-adjusted ^{&}		11.9	7.9	17.6		9.7	7.7	12.3


* Courses for high blood pressure, arthritis, or diabetes.

& Percent is age adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

New York City Prevent Chronic Diseases

Percentage of adults with arthritis*



		New Yo	rk City			State	wide	
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total	1,233,424	18.9	15.7	22.5	3,491,206	22.9	21.3	24.6
Age								
18-24		#			40,067	2.0	1.1	3.6
25-34	103,325	7.1	3.6	13.7	228,246	8.3	5.9	11.7
35-44	102,372	9.0^	3.5	21.0	323,152	12.9	9.4	17.5
45-64	500,875	26.1	20.0	33.3	1,465,151	29.2	26.5	32.1
65+	483,293	46.4	37.1	55.9	1,353,867	49.3	45.4	53.2
Race								
White, NH	566,758	24.3	19.4	30.0	2,474,787	27.2	25.6	28.9
Black, NH	278,480	17.3	11.0	26.2	400,524	18.5	13.4	24.9
Hispanic	325,868	18.7	12.7	26.7	460,563	18.0	13.6	23.4
Other, NH	62,319	7.3	3.3	15.6	155,331	10.8	7.2	15.8
Sex								
Male	377,747	12.4	8.4	17.8	1,258,898	17.3	15.2	19.6
Female	855,677	24.6	20.0	29.8	2,232,309	28.0	25.7	30.4
Age-adjusted ^{&}		19.0	15.9	22.4		21.8	20.3	23.3


* Includes rheumatoid arthritis, gout, lupus, or fibromyalgia.

& Percent is age adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Promote a Healthy and Safe Environment

Percentage of adults who consider their neighborhood suitable for walking and physical activity




		New Yo	rk City			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	5,472,566	89.8	86.2	92.5	13,107,891	91.3	89.8	92.6
Age								
18-24	702,702	87.8^	70.0	95.7	1,641,319	87.9	80.9	92.5
25-34	1,118,322	83.5^	72.2	90.8	2,218,379	86.6	80.9	90.8
35-44	1,018,215	94.0	87.0	97.4	2,218,331	93.4	90.6	95.4
45-64	1,627,236	90.6	85.1	94.2	4,372,743	92.2	90.1	93.9
65+	912,043	94.3	88.1	97.4	2,447,810	94.7	92.7	96.1
Race								
White, NH	2,048,097	92.5	86.7	95.9	8,055,698	93.1	91.7	94.3
Black, NH	1,349,120	89.7	81.2	94.7	1,798,763	89.3	83.3	93.4
Hispanic	1,272,494	81.5	72.3	88.1	1,963,304	84.3	78.2	89.0
Other, NH	802,854	98.5	93.1	99.7	1,290,125	94.0	88.9	96.9
Sex								
Male	2,679,026	92.1	86.6	95.5	6,375,291	92.5	90.3	94.3
Female	2,793,539	87.7	82.5	91.5	6,732,600	90.1	87.9	91.9
Age-adjusted ^{&}		90.3	87.0	92.9		91.3	89.8	92.6

& Percent is age adjusted to the 2000 United States Population.

Promote a Healthy and Safe Environment

Percentage of adults aged 65+ years with at least one fall in the past 12 months



		New Yo	ork City			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
г	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	315,428	32.6	24.3	42.1	802,383	30.8	27.3	34.5
Age⁺								
18-24								
25-34								
35-44								
45-64								
65+								
Race								
White, NH	163,129	34.6^	24.5	46.3	600,059	31.3	28.3	34.5
Black, NH		#			61,485	22.9^	12.6	38.1
Hispanic		#			96,513	39.1^	21.8	59.7
Other, NH		#			44,326	25.2^	12.2	45.0
Sex								
Male	106,522	33.1^	19.7	50.0	324,764	30.4	25.5	35.7
Female	208,907	32.3^	22.4	44.2	477,619	31.1	26.4	36.2
Age-adjusted ^{+,&}								


& Percent is age adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Promote a Healthy and Safe Environments

Among adults aged 65+ years with at least one fall in the past 12 months, percent that resulted in injury^{*}



		New Yo	ork City			State	wide	Upper 95% limit 44.9 43.4 77.2 66.8 68.8 41.5 51.5	
	Est. # of	_	Lower	Upper	Est. # of	_	Lower		
г	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit	
Total	114,386	36.3^	23.1	51.9	304,376	38.3	32.2	44.9	
Age⁺									
18-24									
25-34									
35-44									
45-64									
65+									
Race									
White, NH		#			223,708	37.6	32.1	43.4	
Black, NH		#			33,030	53.7^	28.5	77.2	
Hispanic		#			31,933	34.1^	11.7	66.8	
Other, NH		#			15,705	35.4^	12.0	68.8	
Sex									
Male		#			103,371	32.2	24.1	41.5	
Female		#			201,005	42.5	33.9	51.5	
Age-adjusted ^{+,&}									


* Falls that limited regular activities or lead to doctor visits.

& Percent is age adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV)



		New Yo	ork City			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
г	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	747,301	43.8	35.8	52.1	1,662,747	37.3	34.0	40.8
Age⁺								
18-24								
25-34								
35-44								
45-64								
65+								
Race								
White, NH	241,118	39.3^	29.8	49.8	957,863	33.7	31.0	36.4
Black, NH	265,641	52.5^	36.8	67.7	347,642	52.3^	40.1	64.2
Hispanic	198,922	45.3^	27.9	64.0	270,194	44.2^	31.2	58.1
Other, NH		#			87,049	26.2^	15.7	40.4
Sex								
Male	414,080	50.1^	38.0	62.3	929,342	43.4	38.2	48.7
Female	333,221	37.8^	28.1	48.6	733,405	31.7	27.6	36.1
Age-adjusted ^{+,&}								


& Percent is age adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections

Percentage of adults aged 18+ years with flu immunization in the past year*




		New Yo	rk City			State	wide	
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	2,685,739	43.2	37.8	48.8	6,798,706	46.7	44.3	49.2
Age								
18-24	253,400	30.8^	14.8	53.3	600,414	31.7	23.4	41.5
25-34	491,661	36.4^	25.3	49.1	907,897	35.3	29.0	42.1
35-44	476,999	43.7	29.3	59.2	981,422	41.1	33.9	48.6
45-64	771,810	41.7	34.2	49.7	2,282,268	47.4	44.1	50.8
65+	630,200	64.2^	53.7	73.4	1,895,586	72.1	67.9	75.9
Race								
White, NH	1,010,401	45.2	38.0	52.5	4,296,522	49.3	47.1	51.5
Black, NH	494,107	32.0^	22.7	43.1	718,122	35.0	27.6	43.3
Hispanic	675,057	41.9^	32.4	52.0	971,941	40.8	34.1	47.8
Other, NH	506,174	61.1^	39.2	79.3	812,120	58.5^	45.1	70.7
Sex								
Male	1,184,611	40.2	32.3	48.7	3,045,064	43.8	40.1	47.5
Female	1,501,128	45.9	38.7	53.2	3,753,643	49.5	46.2	52.8
Age-adjusted ^{&}		43.6	38.1	49.3		46.0	43.4	48.6

* Includes flu shot or flu spray

& Percent is age adjusted to the 2000 United States Population.

Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections

Percentage of adults aged 65+ years with flu immunization in the past year*



		New Yo	ork City			State	wide	iii 95% limit 9 75.9 9 75.9 9 79.6 8 55.5 3 84.3 4 89.7 7 79.3	
-	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit		
Total	630,200	64.2^	53.7	73.4	1,895,586	72.1	67.9	75.9	
Age⁺									
18-24									
25-34									
35-44									
45-64									
65+									
Race									
White, NH	366,619	76.6^	65.1	85.2	1,484,360	76.9	73.9	79.6	
Black, NH		#			103,016	38.4^	23.8	55.5	
Hispanic		#			170,589	69.0^	47.8	84.3	
Other, NH		#			137,621	75.2^	51.4	89.7	
Sex									
Male	215,815	64.8^	46.6	79.5	802,715	73.9	67.7	79.3	
Female	414,385	63.9^	50.9	75.0	1,092,871	70.8	65.0	76.0	
Age-adjusted ^{+,&}									

* Includes flu shot or flu spray


& Percent is age adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections

Percentage of adults aged 65+ years with pneumococcal immunization



		New Yo	ork City			State	wide	
	Est. # of	_	Lower	Upper	Est. # of	_	Lower	Upper
г	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	504,961	55.3^	44.8	65.3	1,617,276	65.1	60.7	69.2
Age⁺								
18-24								
25-34								
35-44								
45-64								
65+								
Race								
White, NH	290,588	66.2^	54.0	76.5	1,303,462	71.4	68.1	74.4
Black, NH		#			123,843	48.9^	30.2	67.8
Hispanic		#			104,466	45.7^	27.5	65.3
Other, NH		#			85,505	48.1^	27.7	69.2
Sex								
Male	158,971	52.4^	35.6	68.6	631,549	62.4	56.7	67.7
Female	345,990	56.8^	43.5	69.2	985,727	66.9	60.5	72.7
Age-adjusted ^{+,&}								


& Percent is age adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

New York City Promote Healthy Women, Infants, and Children

Percentage of women aged 18-44 years who had dental visit within the last year



		New Yo	ork City			State	wide	Upper 95% limit 76.5 78.3 89.5 74.4 86.8	
	Est. # of		Lower	Upper	Est. # of		Lower		
	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit	
Total	1,014,206	67.9^	54.0	79.3	2,268,176	70.7	64.0	76.5	
Age ⁺									
18-24									
25-34									
35-44									
45-64									
65+									
Race									
White, NH	202,098	63.6^	43.0	80.1	1,155,302	73.5	68.1	78.3	
Black, NH		#			426,771	78.8^	61.7	89.5	
Hispanic		#			412,421	62.2^	48.3	74.4	
Other, NH		#			273,683	63.0^	30.7	86.8	
Sex									
Male		#				#			
Female	1,014,206	67.9^	54.0	79.3	2,268,176	70.7	64.0	76.5	
Age-adjusted ^{+,&}									


& Percent is age adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Promote Healthy Women, Infants, and Children

Women aged 18-44 years who saw a doctor for a routine checkup within the last year



		New Yo	ork City			State	wide	it 95% limit 5 77.8 77.6 92.4 85.3 5 83.8	
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit		
Tatal									
Total	1,296,418	74.6^	61.9	84.2	2,615,375	72.5	66.5	//.8	
Age⁺									
18-24									
25-34									
35-44									
45-64									
65+									
Race									
White, NH	278,185	67.5	49.0	81.8	1,220,281	70.1	65.1	74.6	
Black, NH		#			512,270	83.8^	68.8	92.4	
Hispanic	463,185	81.5^	63.2	91.8	606,739	76.3^	64.1	85.3	
Other, NH		#			276,085	60.1^	30.5	83.8	
Sex									
Male		#				#			
Female	1,296,418	74.6^	61.9	84.2	2,615,375	72.5	66.5	77.8	
Age-adjusted ^{+,&}									


& Percent is age adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

New York City Promote Healthy Women, Infants, and Children

Percentage of women aged 18-64 years with health care coverage



		New Yo	ork City			State	wide	90.2 93.0 96.1 83.5 89.5	
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit		
Total	2,257,081	84.5	75.4	90.6	5,351,099	87.2	83.4	90.2	
Age⁺									
18-24									
25-34									
35-44									
45-64									
65+									
Race									
White, NH	737,040	91.0	84.3	95.0	3,178,979	91.6	90.0	93.0	
Black, NH	698,411	95.3	88.2	98.2	916,108	93.3	88.7	96.1	
Hispanic	606,604	80.4^	66.6	89.4	807,553	76.0	66.5	83.5	
Other, NH		#			448,460	71.8^	43.3	89.5	
Sex									
Male		#				#			
Female	2,257,081	84.5	75.4	90.6	5,351,099	87.2	83.4	90.2	
Age-adjusted ^{+,&}									


& Percent is age adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

New York City Promote Healthy Women, Infants, and Children

Percentage of women aged 18-44 years whose health care provider discussed planning a healthy pregnancy



	New York City				Statewide			
F	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total	494,242	33.0^	22.7	45.3	1,268,971	39.3	33.7	45.2
Age⁺								
18-24								
25-34								
35-44								
45-64								
65+								
Race								
White, NH	136,662	43.0^	25.9	61.9	723,302	45.9	41.1	50.9
Black, NH		#			291,237	53.7^	35.5	71.0
Hispanic		#			192,904	28.5^	19.0	40.3
Other, NH		#			61,528	14.1^	6.5	27.9
Sex								
Male		#				#		
Female	494,242	33.0^	22.7	45.3	1,268,971	39.3	33.7	45.2
Age-adjusted ^{+,&}								


& Percent is age adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Promote Mental Health and Prevention Substance Abuse

Percentage of adults with poor mental health for 14 or more days in the last month




	New York City				Statewide			
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
F	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	688,361	10.9	7.7	15.2	1,672,677	11.3	9.7	13.0
Age								
18-24		#			186,862	9.5	6.7	13.1
25-34	251,196	17.9^	8.7	33.3	418,716	15.7	10.2	23.4
35-44	96,372	8.5^	3.0	22.0	259,463	10.5	7.0	15.4
45-64	209,902	11.5	7.6	17.1	587,164	12.1	10.3	14.2
65+	74,766	7.6	3.8	14.6	175,961	6.7	4.9	9.0
Race								
White, NH	301,904	13.2	7.6	22.1	1,072,012	12.0	10.2	14.2
Black, NH	154,557	9.8	4.8	19.0	237,227	11.2	7.0	17.5
Hispanic	173,941	10.8	5.7	19.7	264,437	11.0	7.1	16.6
Other, NH	57,959	7.0^	2.7	17.0	99,001	7.0	3.9	12.2
Sex								
Male	295,017	10.1	5.3	18.5	734,315	10.4	7.9	13.5
Female	393,344	11.6	8.0	16.5	938,362	12.1	10.2	14.2
Age-adjusted ^{&}		10.1	7.2	14.0		11.1	9.6	13.0

& Percent is age adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Promote Mental Health and Prevention Substance Abuse

Percentage of adult binge drinking during the past month*



	New York City				Statewide			
	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit	Est. # of Adults	Percent	Lower 95% limit	Upper 95% limit
Total	1,075,670	17.8	13.5	23.1	2,413,447	16.9	14.9	19.1
Age								
18-24	224,877	26.7^	12.2	48.7	515,466	27.1	19.4	36.6
25-34	332,669	25.2^	14.8	39.5	633,963	25.0	18.9	32.3
35-44	263,743	27.2^	16.1	42.3	509,131	22.6	17.2	29.1
45-64	236,027	12.7	8.5	18.6	656,578	13.7	11.7	16.0
65+		#			86,636	3.4	2.7	4.2
Race								
White, NH	540,817	24.6	17.7	33.2	1,649,594	19.1	17.0	21.5
Black, NH	250,029	16.8^	8.2	31.4	324,364	16.3^	9.4	26.8
Hispanic	178,777	11.0	5.9	19.7	277,966	11.6	7.7	17.1
Other, NH	106,048	14.3^	5.6	31.9	161,523	12.4	6.8	21.7
Sex								
Male	723,874	25.0	17.5	34.4	1,567,117	22.8	19.3	26.8
Female	351,797	11.2	7.7	15.9	846,330	11.4	9.6	13.4
Age-adjusted ^{&}		18.1	13.9	23.3		17.7	15.6	20.0
Age-aujusteu		10.1	13.9	25.5		1/./	13.0	20.0


* Men who have five or more drinks , women having four or more drinks on one occasion

& Percent is age adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

New York City Promote Mental Health and Prevention Substance Abuse

Percentage of cigarette smoking among adults who report poor mental health



	New York City				Statewide			
	Est. # of		Lower	Upper	Est. # of		Lower	Upper
г	Adults	Percent	95% limit	95% limit	Adults	Percent	95% limit	95% limit
Total	188,875	28.4^	12.9	51.4	519,245	31.8	24.1	40.6
Age								
18-24		#			64,468	35.6^	22.4	51.5
25-34		#			192,926	47.3^	24.8	70.8
35-44		#			70,340	27.6^	16.6	42.1
45-64		#			177,977	31.2	24.0	39.5
65+		#			13,368	7.7	4.1	13.9
Race								
White, NH		#			423,246	39.9^	29.9	50.7
Black, NH		#			33,694	14.4^	7.7	25.3
Hispanic		#			45,968	18.1^	9.4	31.9
Other, NH		#			16,336	19.6^	9.4	36.5
Sex								
Male		#			272,857	38.4^	24.1	55.0
Female	82,113	21.3^	10.5	38.4	246,388	26.7	20.7	33.8
Age-adjusted ^{&}		20.6^	12.2	32.5		29.9	24.1	36.4

& Percent is age adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.