eBRFSS 2013-2014 Health Indicators: Schuyler

| General Description | 2 |
|---|---|
| Improve Health Status and Reduce Health Disparities | |
| Percentage of adults who did not receive medical care because of cost | |
| Percentage of adults who have a regular health care provider | 4 |
| Percentage of adults who had a dentist visit within the past year | 5 |
| Percentage of adults living with a disability | |
| Percentage of adults aged 18-64 years with healthcare coverage | 7 |
| Percentage of adults aged 18-64 years who saw a doctor for a routine checkup within the last year | |
| Percentage of adults experiencing housing insecurity in the past 12 months | |
| Percentage of adults experiencing food insecurity in the past 12 months | |
| Percentage of adults with poor self-reported health | |
| Percentage of adults who report 14 or more days of poor physical health | |
| Prevent Chronic Diseases | |
| Percentage of adults who are obese | |
| Percentage of adults with an annual household income less than \$25,000 who are obese | |
| Percentage of adults living with a disability who are obese | |
| Percentage of adults overweight or obese | |
| Percentage of adults who participated in leisure time physical activity in the past 30 days | |
| Percentage of adults who consume one or more sugary drinks daily | |
| Percentage of adults who consume fast-food three or more times per week | |
| Percentage of adults with physician diagnosed diabetes | |
| Percentage of adults who are current smokers | |
| Percentage of adults with annual household income less than \$25,000 who are current smokers | |
| Percentage of adults with annual household income less than \$25,000 who are current shokers | |
| Percentage of women aged 50-74 years receiving breast cancer screening based on recent guidelines | |
| Percentage of women aged 50-74 years with annual household income of less than \$25,000 receiving breast cancer scre | |
| guidelines | |
| Percentage of women aged 21-65 years receiving cervical cancer screening based on recent guidelines | |
| Percentage of women aged 21-65 years with annual household income of less than \$25,000 receiving cervical cancer sc | |
| guidelines | |
| guidelines | |
| Percentage of adults aged E0.75 years receiving colorectal cancer screening based on recent guidelines | 20 |
| Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines | |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree | ening based on recent |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines | ening based on recent 29 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years | ening based on recent 29 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes | ening based on recent 29 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure | ening based on recent 29 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication | ening based on recent 29 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma | ening based on recent 29 30 31 32 33 34 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percent of adults with cholesterol checked. | ening based on recent 29 30 31 32 33 34 34 35 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked Percentage of adults 18 years and older with elevated cholesterol | ening based on recent 29 30 31 32 33 34 34 35 36 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked Percentage of adults 18 years and older with elevated cholesterol Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition | ening based on recent 29 30 31 32 33 34 35 36 37 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked Percentage of adults 18 years and older with elevated cholesterol Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults with arthritis | ening based on recent 29 30 31 32 33 34 35 36 37 38 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked Percentage of adults 18 years and older with elevated cholesterol Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults with arthritis Promote a Healthy and Safe Environment | ening based on recent 29 30 31 32 33 34 34 35 36 37 38 39 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked Percentage of adults 18 years and older with elevated cholesterol Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults with arthritis Promote a Healthy and Safe Environment Percentage of adults who consider their neighborhood suitable for walking and physical activity | ening based on recent 29 30 31 32 33 34 34 35 36 37 38 39 39 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked Percentage of adults 18 years and older with elevated cholesterol Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults with arthritis Promote a Healthy and Safe Environment Percentage of adults who consider their neighborhood suitable for walking and physical activity Percentage of adults aged 65+ years with at least one fall in the past 12 months | ening based on recent 29 30 31 32 33 34 34 35 36 37 38 39 39 39 40 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with current asthma Percentage of adults with cholesterol checked Percentage of adults 18 years and older with elevated cholesterol Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults with arthritis Promote a Healthy and Safe Environment Percentage of adults who consider their neighborhood suitable for walking and physical activity Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months, percent that resulted in injury | ening based on recent 29 30 31 32 33 34 34 35 36 37 38 39 39 40 41 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked Percentage of adults 18 years and older with elevated cholesterol Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults with arthritis Promote a Healthy and Safe Environment Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months, percent that resulted in injury | ening based on recent 29 30 31 32 33 34 34 35 36 37 38 39 39 40 41 42 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines | ening based on recent 29 30 31 32 33 34 34 35 36 37 38 39 39 40 40 41 42 42 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked Percentage of adults with cholesterol checked Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults with arthritis Promote a Healthy and Safe Environment Percentage of adults who consider their neighborhood suitable for walking and physical activity Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months. Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV) Percentage of adults aged 18+ years with flu immunization in the past year | ening based on recent 29 30 31 32 33 34 35 36 37 38 39 39 40 40 41 42 42 42 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked Percentage of adults with cholesterol checked Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults with arthritis Promote a Healthy and Safe Environment Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 47-68 years reporting ever tested for Hepatitis C (HCV) Percentage of adults aged 18+ years with flu immunization in the past year | ening based on recent 29 30 31 32 33 34 35 36 37 38 39 39 40 40 41 42 42 43 44 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked Percentage of adults with cholesterol checked Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults with affe Environment Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months. Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV) Percentage of adults aged 18+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year | ening based on recent 29 30 31 32 33 34 34 35 36 36 37 38 39 40 40 41 42 42 42 42 43 44 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked. Percentage of adults 18 years and older with elevated cholesterol Percentage of adults with arthritis Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months. Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV). Percentage of adults aged 18+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year | ening based on recent 29 30 31 32 33 34 34 35 36 36 37 38 39 40 40 41 42 42 42 42 43 44 45 46 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked. Percentage of adults 18 years and older with elevated cholesterol Percentage of adults with arthritis Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months, percent that resulted in injury Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections. Percentage of adults aged 65+ years with flu immunization in the past year. Percentage of adults aged 65+ years with flu immunization in the past year. Percentage of adults aged 65+ years with flu immunization in the past year. Percentage of adults aged 65+ years with flu immunization in the past year. Percentage of adults aged 65+ years with flu immunization in the past year. Percentage of adults aged 65+ years with flu immunization in the past year. Percentage of adults aged 65+ years with flu immunization in the past year. Percentage of adults aged 65+ years with pneumococcal immunization. Promote Healthy Women, Infants, and Children Percentage of women aged 18-44 years who had dental visit within the last year | ening based on recent 29 30 31 32 33 34 34 35 36 37 38 39 39 40 40 41 41 42 42 42 42 43 44 44 45 46 46 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults who consider their neighborhood suitable for walking and physical activity Percentage of adults who consider their neighborhood suitable for walking and physical activity Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months. Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV) Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ yea | ening based on recent 29 30 31 32 33 34 34 35 36 37 38 39 39 40 41 41 42 42 42 42 43 44 44 45 46 46 46 47 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with physician-diagnosed high blood pressure medication Percentage of adults with current asthma Percentage of adults with colesterol checked. Percentage of adults 18 years and older with elevated cholesterol Percentage of adults with arthritis Percentage of adults with arthritis. Promote a Healthy and Safe Environment Percentage of adults who consider their neighborhood suitable for walking and physical activity Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months. Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV). Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV). Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of women aged 18-44 years who had dental visit within the last year Women aged 18-44 years who saw a doctor for a routine checkup within the last year Percentage of women | ening based on recent 29 30 31 32 33 34 34 35 36 37 38 39 39 40 40 41 42 42 42 42 42 42 42 43 44 44 45 46 46 47 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with physician-diagnosed high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked. Percentage of adults the years and older with elevated cholesterol Percentage of adults with arthritis Percentage of adults with arthritis Promote a Healthy and Safe Environment Percentage of adults aged 65+ years with at least one fall in the past 12 months, percent that resulted in injury Prevent HIV/STDS, Vaccine Preventable Diseases and Healthcare-Associated Infections. Percentage of adults aged 65+ years with at least one fall in the past 12 months, percent that resulted in injury Prevent HIV/STDS, Vaccine Preventable Diseases and Healthcare-Associated Infections. Percentage of adults aged 18+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of women aged 18-44 years who had dental visit within the last year Women aged 18-44 years who saw a doctor for a routine checkup within the last year Percentage of women aged 18-44 years whose health care coverage Percentage of women aged 18-44 years wh | ening based on recent 29 30 31 32 33 34 34 35 36 37 38 39 39 40 40 41 42 42 42 42 42 43 44 44 45 46 46 47 48 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with current asthma Percentage of adults with current asthma Percentage of adults with aver taken a course or class to learn how to manage their chronic disease or condition Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults who consider their neighborhood suitable for walking and physical activity Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months. Percentage of adults aged 45+ years with at least one fall in the past 12 months. Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV). Percentage of adults aged 18+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with pueumococcal immunization Promote Healthy Women, Infants, and Children Percentage of women aged 18-44 years who had dental visit within the last year. Percentage of women aged 18-44 years who had dental visit within the last year. Percentage of women aged 18-44 years who had dental visit within the last year. Percentage of women aged 18-44 years who saw a doctor for a routine checkup within the last year. Percentage of women aged 18-44 years who saw a doctor for a routine checkup within the last year. Percentage of w | ening based on recent 29 30 31 32 33 34 34 35 36 37 38 39 39 40 40 41 42 42 42 42 42 43 44 45 46 46 46 47 48 49 50 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with clurrent asthma Percentage of adults with clurrent asthma Percentage of adults with current asthma Percentage of adults with current asthma Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults who consider their neighborhood suitable for walking and physical activity Percentage of adults who consider their neighborhood suitable for walking and physical activity Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months. Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV). Percentage of adults aged 18+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 65+ years with flu immunization in the past year Percentage of adults aged 18+ 4years with flu immunization in the past year Percentage of women aged 18-44 years who had dental visit within the last year Percentage of women aged 18-44 years who had dental visit within the last year Percentage of women aged 18-44 years who had dental visit within the last year Percentage of women aged 18-44 years who had dental visit within the last year Percentage of women aged 18-44 years who had dental visit within the last year Percentage of women aged 18-44 years who had dental visit within the last year Percentage of women aged 18-44 years who had dental visit within the last year Perc | ening based on recent 29 30 31 32 33 34 34 35 36 37 38 39 39 40 40 41 41 42 42 43 44 44 45 45 46 46 47 48 49 50 50 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines | ening based on recent 29 30 31 32 33 34 34 35 36 37 38 39 39 40 40 41 42 42 42 43 40 41 42 42 43 40 41 42 42 43 40 41 42 43 40 41 42 43 40 41 42 43 50 50 50 50 50 50 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer scree guidelines | ening based on recent 29 30 31 32 33 34 34 35 36 37 38 39 39 40 40 41 42 42 42 43 40 41 42 42 43 40 41 42 42 43 40 41 42 43 40 41 42 43 40 41 42 43 50 50 50 50 50 50 |

General Description

The New York State Expanded Behavioral Risk Factor Surveillance System 2013-2014 (2014 eBRFSS) was design to supplement the Center for Disease Control and Prevention (CDC) Behavioral Risk Factor Surveillance System (BRFSS), which is conducted annually in New York State. The purpose of the 2014 eBRFSS was to produce local information on key public health issues. Data for the project were collected from April 15, 2013 to May 10, 2014. The 2014 eBRFSS reached both households with landline telephones and households which only had cell phones. Like the annual BRFSS, New York State's 2014 eBRFSS was designed to be representative of the non-institutionalized adult household population, aged 18 years and older.

Reported Health Indicators

There are a total of 50 health indicators included in the report. The health indicators include 10 measures included in the New York State Department of Health's Prevention Agenda, NYS Health Innovation Plan (SHIP), Delivery System Reform Incentive Payment (DSRIP) Program, or the NYS Community Health Indicator Reports (CHIRS). An additional 40 important health indicators are also included. The indicators are organized by the Priority Area of the Prevention Agenda and are reported statewide, by region, and by county.

Data Collection Methodology, Sampling and Weighting

The data collection methods for the 2014 eBRFSS aligned with the CDC's standard annual BRFSS methods. The 2014 eBRFSS questionnaire used standard questions from the BRFSS. Survey interviews were conducted in both English and Spanish. There were 31,690 completed interviews.

The sampling plan for the eBRFSS was developed to result in valid health indicator estimates for New York City (NYC) and each of the 57 counties outside NYC. The sampling approach stratified landlines by county and implemented a disproportionate stratified sampling design that accounted for both listed-households and unlisted numbers. Cell phone households were accounted for through additional strata that covered contiguous groups of counties.


Data were weighted to represent the population using a methodology known as iterative proportional fitting (raking). Raking adjusted the data so that groups underrepresented in the sample were more accurately represented in the estimated rates. Raking also enabled surveys conducted by cell phone to be included in the weighting and to accurately match sample distributions of specified demographic characteristics of the population. The raking categories included in the eBRFSS were: age, gender, race/ethnicity, education, marital status, owner/renter status, telephone usage (i.e. landline and cell phone combinations), and county of residence.

Limitations of Data Use:

During data analysis, criteria were applied to suppress rates based on too few surveys to be reliable and to identify rates that are highly variable. Rates in which less than 10 respondents were in the numerator or less than 50 respondents were in the denominator were considered unreliable and suppressed from reporting. Rates considered to have high variability due to having confidence limits greater than $\pm 10\%$ are noted.

Improve Health Status and Reduce Health Disparities

Percentage of adults who did not receive medical care because of cost*



| Schuyler County | |
|-----------------|--|
| | |

Finger Lakes

| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|------------|-----------|---------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 1,173 | 7.9 | (4.7-13.1) | 105,514 | 10.5 | (9.0-12.3) |
| Age | | | | | | |
| 18-24 | | # | | 18,069 | 13.6 | (8.4-21.2) |
| 25-34 | | # | | 23,656 | 15.2 | (10.8-21.1) |
| 35-44 | | # | | 18,598 | 12.5 | (8.5-17.9) |
| 45-64 | 338 | 6.0 | (2.9-12.3) | 35,930 | 10.1 | (7.9-12.8) |
| 65+ | | # | | 7,633 | 3.9 | (2.5-6.1) |
| Race | | | | | | |
| White, NH | 998 | 7.3 | (4.1-12.5) | 78,405 | 9.9 | (8.3-11.7) |
| Black, NH | | # | | 11,314 | 14.5 | (8.2-24.2) |
| Hispanic | | # | | 8,090 | 15.4 | (9.2-24.8) |
| Other, NH | | # | | 7,706 | 10.2^ | (4.8-20.4) |
| Sex | | | | | | |
| male | 649 | 8.8 | (4.6-16.4) | 46,595 | 9.7 | (7.5-12.3) |
| female | 524 | 7.1 | (3.1-15.4) | 58,919 | 11.4 | (9.3-13.9) |
| Age-adjusted ^{&} | | 12.6 | (7.3-20.9) | | 11.0 | (9.3-12.9) |


* Among people needing to see a doctor in the past 12 months

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Schuyler Improve Health Status and Reduce Health Disparities

Percentage of adults who have a regular health care provider



Schuyler County

Finger Lakes

| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 13,121 | 89.1 | (82.9-93.2) | 891,951 | 89.5 | (87.7-91.0) |
| Age | | | | | | |
| 18-24 | | # | | 98,637 | 75.1 | (65.5-82.8) |
| 25-34 | | # | | 124,962 | 80.4 | (74.7-85.1) |
| 35-44 | | # | | 130,769 | 88.1 | (82.8-91.9) |
| 45-64 | 4,954 | 89.1 | (81.6-93.8) | 336,849 | 94.7 | (92.7-96.1) |
| 65+ | 3,245 | 98.6 | (96.5-99.5) | 190,189 | 97.8 | (96.5-98.6) |
| Race | | | | | | |
| White, NH | 12,242 | 89.7 | (83.8-93.6) | 717,170 | 90.5 | (88.8-92.0) |
| Black, NH | | # | | 64,548 | 83.7 | (73.7-90.4) |
| Hispanic | | # | | 40,023 | 77.2^ | (63.5-86.8) |
| Other, NH | | # | | 70,210 | 93.3 | (84.6-97.2) |
| Sex | | | | | | |
| male | 6,479 | 88.8 | (79.1-94.3) | 407,470 | 84.7 | (81.6-87.3) |
| female | 6,642 | 89.4 | (80.0-94.7) | 484,481 | 94.0 | (92.0-95.5) |
| Age-adjusted ^{&} | | 86.2 | (77.8-91.8) | | 88.6 | (86.7-90.4) |


1

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Schuyler Improve Health Status and Reduce Health Disparities

Percentage of adults who had a dentist visit within the past year



Schuyler County

Finger Lakes

| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 9,280 | 65.7 | (57.4-73.1) | 684,119 | 72.7 | (69.7-75.6) |
| Age | | | | | | |
| 18-24 | | # | | 90,313 | 71.2 | (57.6-81.8) |
| 25-34 | | # | | 89,852 | 61.3 | (52.1-69.7) |
| 35-44 | | # | | 107,877 | 77.2 | (69.7-83.3) |
| 45-64 | 3,673 | 67.5^ | (57.2-76.4) | 254,354 | 76.5 | (72.6-79.9) |
| 65+ | 1,818 | 58.7^ | (47.1-69.3) | 136,480 | 72.9 | (67.8-77.5) |
| Race | | | | | | |
| White, NH | 8,937 | 68.4 | (59.8-75.9) | 562,395 | 74.8 | (72.3-77.1) |
| Black, NH | | # | | 45,366 | 66.5 | (56.6-75.2) |
| Hispanic | | # | | 38,236 | 76.0^ | (62.5-85.8) |
| Other, NH | | # | | 38,122 | 54.6^ | (32.4-75.0) |
| Sex | | | | | | |
| male | 4,017 | 58.3^ | (44.8-70.7) | 317,427 | 70.0 | (65.6-74.1) |
| female | 5,262 | 72.7 | (63.5-80.3) | 366,693 | 75.2 | (70.9-79.1) |
| Age-adjusted ^{&} | | 61.3 | (52.6-69.4) | | 72.6 | (69.4-75.5) |


Í

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Improve Health Status and Reduce Health Disparities

Percentage of adults living with a disability $\!\!\!\!\!\!^*$



| Schuyler County | |
|-----------------|--|
| | |

Finger Lakes

| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 3,516 | 24.2 | (18.6-30.8) | 248,993 | 25.3 | (22.9-27.8) |
| Age | | | | | | |
| 18-24 | | # | | 13,165 | 9.9 | (5.7-16.7) |
| 25-34 | | # | | 24,795 | 16.2 | (11.5-22.3) |
| 35-44 | | # | | 30,187 | 21.0 | (15.4-28.0) |
| 45-64 | 1,675 | 30.3^ | (21.6-40.8) | 103,044 | 29.1 | (25.3-33.2) |
| 65+ | 1,342 | 40.7 | (30.6-51.6) | 74,769 | 39.3 | (34.1-44.7) |
| Race | | | | | | |
| White, NH | 3,120 | 23.0 | (17.4-29.8) | 193,463 | 24.7 | (22.3-27.3) |
| Black, NH | | # | | 23,963 | 31.7^ | (22.9-41.9) |
| Hispanic | | # | | 17,435 | 34.3^ | (23.4-47.2) |
| Other, NH | | # | | 14,132 | 18.8^ | (9.4-34.1) |
| Sex | | | | | | |
| male | 1,680 | 23.3^ | (15.3-33.8) | 112,778 | 23.7 | (20.7-27.1) |
| female | 1,835 | 25.0 | (17.8-34.0) | 136,215 | 26.7 | (23.3-30.5) |
| Age-adjusted ^{&} | | 22.3 | (16.6-29.3) | | 24.2 | (22.0-26.7) |


* Limited activities (physical, mental, or emotional problems); need of special equipment (e.g. cane, wheel chair).

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Schuyler Improve Health Status and Reduce Health Disparities

Percentage of adults aged 18-64 years with healthcare coverage



| | Schuyler County | | | Finger Lakes | | |
|---------------------------------|---------------------|---------|-------------|---------------------|---------|-------------|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 8,574 | 81.1^ | (69.6-88.9) | 694,148 | 87.8 | (85.7-89.6) |
| Age⁺ | | | | | | |
| 18-24 | | | | | | |
| 25-34 | | | | | | |
| 35-44 | | | | | | |
| 45-64 | | | | | | |
| 65+ | | | | | | |
| Race | | | | | | |
| White, NH | 8,224 | 80.6^ | (68.8-88.7) | 546,972 | 89.1 | (87.0-90.9) |
| Black, NH | | # | | 54,874 | 83.7 | (75.1-89.8) |
| Hispanic | | # | | 37,833 | 76.4^ | (61.7-86.6) |
| Other, NH | | # | | 54,469 | 88.5^ | (75.0-95.2) |
| Sex | | | | | | |
| male | 3,769 | 76.2 | (56.4-88.8) | 329,824 | 84.5 | (80.9-87.5) |
| female | 4,806 | 85.3^ | (72.6-92.8) | 364,324 | 91.0 | (88.6-93.0) |
| Age-adjusted ^{+,&} | | | | | | |


& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Schuyler Improve Health Status and Reduce Health Disparities

Percentage of adults aged 18-64 years who saw a doctor for a routine checkup within the last year



| | Schuyler County | | | Finger Lakes | | |
|---------------------------------|---------------------|---------|-------------|---------------------|---------|-------------|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 8,190 | 72.8 | (63.6-80.5) | 501,658 | 63.5 | (59.9-66.9) |
| Age⁺ | | | | | | |
| 18-24 | | | | | | |
| 25-34 | | | | | | |
| 35-44 | | | | | | |
| 45-64 | | | | | | |
| 65+ | | | | | | |
| Race | | | | | | |
| White, NH | 7,673 | 71.9 | (62.3-79.9) | 387,058 | 63.1 | (59.7-66.4) |
| Black, NH | | # | | 52,686 | 79.2^ | (68.3-87.1) |
| Hispanic | | # | | 32,661 | 66.3^ | (51.9-78.3) |
| Other, NH | | # | | 29,253 | 47.7^ | (25.7-70.6) |
| Sex | | | | | | |
| male | 4,198 | 76.7^ | (63.3-86.2) | 228,555 | 58.5 | (53.4-63.5) |
| female | 3,992 | 69.2^ | (56.5-79.5) | 273,103 | 68.3 | (63.6-72.6) |
| Age-adjusted ^{+,&} | | | | | | |


& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Improve Health Status and Reduce Health Disparities

Percentage of adults experiencing housing insecurity in the past 12 months*



Schuyler County

Finger Lakes

| | | layier coa | incy. | | inger Luk | 65 |
|-------------------------------|---------------------|------------|-------------|---------------------|-----------|-------------|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 4,576 | 33.9 | (26.6-42.0) | 314,889 | 34.4 | (31.5-37.4) |
| Age | | | | | | |
| 18-24 | | # | | 32,879 | 32.4 | (22.8-43.7) |
| 25-34 | | # | | 72,916 | 51.0 | (42.6-59.4) |
| 35-44 | | # | | 49,765 | 35.8 | (29.2-43.0) |
| 45-64 | 1,907 | 36.5^ | (27.1-47.0) | 111,638 | 33.0 | (28.9-37.4) |
| 65+ | 638 | 19.8^ | (12.1-30.7) | 42,474 | 23.1 | (17.9-29.3) |
| Race | | | | | | |
| White, NH | 4,026 | 32.4 | (25.1-40.7) | 232,180 | 31.9 | (29.2-34.7) |
| Black, NH | | # | | 34,856 | 49.6^ | (39.3-60.0) |
| Hispanic | | # | | 18,970 | 38.4^ | (26.4-51.9) |
| Other, NH | | # | | 28,883 | 42.5 | (23.1-64.4) |
| Sex | | | | | | |
| male | 1,872 | 27.4^ | (17.8-39.6) | 141,970 | 32.0 | (27.8-36.6) |
| female | 2,704 | 40.6^ | (31.0-51.0) | 172,920 | 36.6 | (32.7-40.7) |
| Age-adjusted ^{&} | | 41.6 | (33.3-50.4) | | 35.1 | (32.2-38.2) |


* Worried or stressed about having enough money to pay for rent or mortgage.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Improve Health Status and Reduce Health Disparities

Percentage of adults experiencing food insecurity in the past 12 months*



Schuyler County

Finger Lakes

| | Senayier county | | | | | |
|-------------------------------|---------------------|----------------|-------------|---------------------|---------|-------------|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 3,475 | 24.0 | (17.5-32.1) | 213,556 | 22.2 | (19.7-24.8) |
| Age | | | | | | |
| 18-24 | | # | | 27,370 | 21.4 | (15.0-29.6) |
| 25-34 | | # | | 52,899 | 35.3 | (27.0-44.7) |
| 35-44 | | # | | 36,748 | 26.0 | (20.2-32.8) |
| 45-64 | 1,229 | 22.3^ | (14.7-32.3) | 70,256 | 20.4 | (17.2-24.1) |
| 65+ | 539 | 16.5^ | (9.3-27.6) | 23,573 | 12.4 | (9.0-17.0) |
| Race | | | | | | |
| White, NH | 2,919 | 21.8 | (15.3-30.1) | 154,900 | 20.2 | (18.0-22.7) |
| Black, NH | | # | | 27,276 | 37.2^ | (28.1-47.4) |
| Hispanic | | # | | 14,323 | 28.0^ | (18.4-40.2) |
| Other, NH | | # | | 17,057 | 23.4 | (9.5-47.1) |
| Sex | | | | | | |
| male | 1,280 | 18.0° | (10.5-29.2) | 97,708 | 21.1 | (17.2-25.4) |
| female | 2,195 | 29.8^ | (19.9-42.1) | 115,848 | 23.2 | (20.2-26.5) |
| Age-adjusted ^{&} | | 28.1 | (20.7-37.0) | | 23.1 | (20.6-25.9) |


* Worried or stressed about having enough money to buy nutritious meals.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Schuyler Improve Health Status and Reduce Health Disparities

Percentage of adults with poor self-reported health




| | Scl | huyler Cou | nty | Finger Lakes | | |
|-------------------------------|---------------------|------------|------------|---------------------|---------|------------|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 633 | 4.3 | (2.6-7.2) | 35,254 | 3.5 | (2.8-4.5) |
| Age | | | | | | |
| 18-24 | | # | | | # | |
| 25-34 | | # | | | # | |
| 35-44 | | # | | | # | |
| 45-64 | | # | | 16,753 | 4.7 | (3.4-6.4) |
| 65+ | 455 | 14.0^ | (7.7-24.0) | 11,921 | 6.1 | (4.4-8.5) |
| Race | | | | | | |
| White, NH | 539 | 4.0 | (2.2-6.9) | 23,699 | 3.0 | (2.3-3.8) |
| Black, NH | | # | | 4,892 | 6.3 | (3.1-12.1) |
| Hispanic | | # | | 4,142 | 7.9^ | (3.2-18.0) |
| Other, NH | | # | | 2,522 | 3.4 | (1.3-8.5) |
| Sex | | | | | | |
| male | 503 | 6.9 | (3.6-12.7) | 19,535 | 4.1 | (2.9-5.6) |
| female | | # | | 15,719 | 3.1 | (2.2-4.2) |
| Age-adjusted ^{&} | | 3.3 | (2.1-5.4) | | 3.3 | (2.6-4.3) |

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Improve Health Status and Reduce Health Disparities

Percentage of adults who report 14 or more days of poor physical health*



| Schuyler County | |
|-----------------|--|
| | |

Finger Lakes


| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| F | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 1,832 | 12.6 | (7.6-20.3) | 113,479 | 11.6 | (10.0-13.5) |
| Age | | | | | | |
| 18-24 | | # | | 6,519 | 4.9 | (1.9-12.4) |
| 25-34 | | # | | 11,848 | 7.9 | (5.0-12.3) |
| 35-44 | | # | | 13,334 | 9.1 | (5.5-14.7) |
| 45-64 | 342 | 6.2 | (3.0-12.4) | 53,981 | 15.3 | (12.4-18.8) |
| 65+ | 744 | 23.7 | (15.3-34.7) | 26,922 | 14.7 | (11.9-18.1) |
| Race | | | | | | |
| White, NH | 1,477 | 11.0 | (6.0-19.2) | 91,151 | 11.7 | (10.0-13.8) |
| Black, NH | | # | | 10,140 | 13.4 | (8.2-21.4) |
| Hispanic | | # | | 8,112 | 16.1 | (9.8-25.5) |
| Other, NH | | # | | 4,076 | 5.6 | (2.7-11.2) |
| Sex | | | | | | |
| male | 869 | 12.0 | (7.0-19.9) | 53,813 | 11.3 | (9.0-14.1) |
| female | 963 | 13.2 | (5.7-27.6) | 59,667 | 11.9 | (9.8-14.4) |
| Age-adjusted ^{&} | | 11.8 | (7.3-18.4) | | 11.2 | (9.6-13.0) |

* Out of the past 30 days.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults who are obese*



Schuyler County

Finger Lakes


| | ••• | | , | | | |
|-------------------------------|---------------------|---------|-------------|---------------------|---------|-------------|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 3,260 | 24.1 | (18.6-30.5) | 254,425 | 27.0 | (24.4-29.8) |
| Age | | | | | | |
| 18-24 | | # | | 16,808 | 13.5 | (7.9-22.1) |
| 25-34 | | # | | 43,074 | 29.7 | (21.3-39.6) |
| 35-44 | | # | | 41,076 | 28.7 | (22.4-36.0) |
| 45-64 | 1,963 | 37.6^ | (28.3-47.8) | 100,327 | 30.0 | (26.3-34.1) |
| 65+ | 735 | 22.6^ | (14.8-32.9) | 51,798 | 27.9 | (23.3-33.1) |
| Race | | | | | | |
| White, NH | 2,877 | 23.1 | (17.6-29.7) | 195,881 | 26.2 | (23.8-28.8) |
| Black, NH | | # | | 25,307 | 34.7 | (26.0-44.5) |
| Hispanic | | # | | 17,493 | 34.1^ | (22.9-47.5) |
| Other, NH | | # | | 15,744 | 22.5^ | (8.5-47.8) |
| Sex | | | | | | |
| male | 1,799 | 26.3^ | (18.1-36.5) | 131,663 | 27.7 | (23.8-32.0) |
| female | 1,461 | 21.9 | (15.3-30.2) | 122,762 | 26.3 | (23.0-29.9) |
| Age-adjusted ^{&} | | 22.9 | (16.9-30.2) | | 27.2 | (24.5-30.1) |

* Obesity defined as a Body Mass Index (BMI) >=30. BMI is calculated based on weight and height.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults with an annual household income less than \$25,000 who are obese*




| | Schuyler County | | | Finger Lakes | | |
|-------------------------------|---------------------|---------|-------------|---------------------|---------|-------------|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 974 | 27.0^ | (16.5-40.9) | 83,889 | 36.9 | (30.6-43.7) |
| Age | | | | - | | |
| 18-24 | | # | | 5,671 | 17.5^ | (7.9-34.3) |
| 25-34 | | # | | 19,305 | 46.4^ | (25.9-68.2) |
| 35-44 | | # | | 9,968 | 34.8^ | (22.0-50.3) |
| 45-64 | | # | | 28,074 | 40.1 | (31.3-49.6) |
| 65+ | 381 | 34.4 | (18.5-54.8) | 20,012 | 37.5 | (28.1-48.0) |
| Race | | | | | | |
| White, NH | 708 | 23.8^ | (13.7-38.0) | 56,366 | 36.0 | (30.2-42.2) |
| Black, NH | | # | | 10,712 | 30.8^ | (20.0-44.3) |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 10,722 | 55.3^ | (20.0-85.9) |
| Sex | | | | | | |
| male | 631 | 36.6 | (19.7-57.7) | 38,883 | 32.8^ | (23.0-44.5) |
| female | 343 | 18.2^ | (8.8-33.9) | 45,006 | 41.3 | (34.4-48.7) |
| Age-adjusted ^{&} | | 21.4^ | (13.0-33.3) | | 36.7 | (30.6-43.3) |

* Obesity defined as a Body Mass Index (BMI) >=30. BMI is calculated based on weight and height.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults living with a disability who are obese*



| Schuyler County | |
|-----------------|--|
| | |

Finger Lakes


| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 1,382 | 41.1^ | (29.0-54.3) | 90,664 | 38.5 | (33.5-43.7) |
| Age | | | | | | |
| 18-24 | | # | | | # | |
| 25-34 | | # | | 7,827 | 35.1^ | (21.1-52.2) |
| 35-44 | | # | | 10,209 | 34.6^ | (21.2-50.8) |
| 45-64 | | # | | 43,860 | 45.1 | (37.3-53.1) |
| 65+ | 555 | 41.9^ | (25.8-60.0) | 25,854 | 36.6 | (28.1-46.1) |
| Race | | | | | | |
| White, NH | 1,131 | 38.1^ | (25.7-52.2) | 68,651 | 37.7 | (32.3-43.4) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | | # | |
| Sex | | | | | | |
| male | | # | | 40,686 | 36.9 | (30.2-44.1) |
| female | 718 | 42.6^ | (26.9-60.0) | 49,978 | 40.0 | (32.9-47.5) |
| Age-adjusted ^{&} | | 25.3 | (18.4-33.7) | | 35.8 | (30.4-41.6) |

* Obesity defined as a Body Mass Index (BMI) >=30. BMI is calculated based on weight and height. Limited activities (physical, mental, or emotional problems); need of special equipment (e.g. cane, wheel chair).

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults overweight or obese*



Schuyler County

Finger Lakes


| | Senayler county | | | i inger zakes | | |
|-------------------------------|---------------------|---------|-------------|---------------------|---------|-------------|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 8,011 | 59.2 | (50.3-67.5) | 581,065 | 61.8 | (58.6-64.8) |
| Age | | | | | | |
| 18-24 | | # | | 41,425 | 33.2 | (23.5-44.5) |
| 25-34 | | # | | 83,868 | 57.7 | (49.6-65.5) |
| 35-44 | | # | | 83,850 | 58.6 | (50.5-66.3) |
| 45-64 | 3,806 | 72.9 | (63.3-80.7) | 241,793 | 72.4 | (68.0-76.4) |
| 65+ | 2,219 | 68.2 | (58.3-76.6) | 125,252 | 67.6 | (62.1-72.6) |
| Race | | | | | | |
| White, NH | 7,356 | 59.0 | (49.8-67.7) | 471,523 | 63.1 | (60.2-66.0) |
| Black, NH | | # | | 49,324 | 67.6^ | (57.0-76.7) |
| Hispanic | | # | | 33,809 | 65.9^ | (50.0-78.9) |
| Other, NH | | # | | 26,409 | 37.8^ | (20.1-59.5) |
| Sex | | | | | | |
| male | 3,961 | 57.8^ | (44.8-69.8) | 329,186 | 69.4 | (65.4-73.1) |
| female | 4,050 | 60.6^ | (48.0-71.8) | 251,879 | 54.0 | (49.3-58.6) |
| Age-adjusted ^{&} | | 61.1 | (53.4-68.3) | | 60.8 | (57.7-63.8) |

* Defined as a Body Mass Index (BMI) >=25. BMI is calculated based on weight and height.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults who participated in leisure time physical activity in the past 30 days*



Schuyler County

Finger Lakes


| | Senariei Councy | | | | Tinger Luk | |
|-------------------------------|---------------------|---------|-------------|---------------------|------------|-------------|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 9,792 | 67.6 | (58.3-75.8) | 703,242 | 72.6 | (69.5-75.6) |
| Age | | | | | | |
| 18-24 | | # | | 97,421 | 75.5^ | (60.0-86.3) |
| 25-34 | | # | | 106,989 | 71.1^ | (60.7-79.7) |
| 35-44 | | # | | 105,857 | 74.8 | (66.9-81.3) |
| 45-64 | 4,311 | 78.1^ | (68.0-85.7) | 256,126 | 73.8 | (69.9-77.4) |
| 65+ | 2,053 | 62.5 | (51.9-72.1) | 128,648 | 67.9 | (62.3-73.0) |
| Race | | | | | | |
| White, NH | 9,074 | 67.7 | (57.8-76.2) | 588,978 | 76.4 | (73.9-78.8) |
| Black, NH | | # | | 53 <i>,</i> 055 | 72.5 | (63.6-79.9) |
| Hispanic | | # | | 33,495 | 65.3^ | (52.7-76.0) |
| Other, NH | | # | | 27,713 | 38.0^ | (22.0-57.0) |
| Sex | | | | | | |
| male | 4,514 | 63.5 | (49.4-75.7) | 346,071 | 74.1 | (69.7-78.0) |
| female | 5,278 | 71.6^ | (59.3-81.3) | 357,171 | 71.3 | (66.6-75.6) |
| Age-adjusted ^{&} | | 68.7 | (60.0-76.3) | | 72.7 | (69.4-75.9) |

* Any physical activities or exercises such as running or walking

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults who consume one or more sugary drinks daily*



Schuyler County

Finger Lakes


| | Senayler County | | | Tinger Eakes | | |
|-------------------------------|---------------------|---------|-------------|---------------------|---------|-------------|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 2,686 | 18.9 | (13.2-26.2) | 214,679 | 22.5 | (19.9-25.3) |
| Age | | | | | | |
| 18-24 | | # | | 55,277 | 43.6^ | (31.8-56.0) |
| 25-34 | | # | | 35,087 | 23.6 | (18.5-29.7) |
| 35-44 | | # | | 34,865 | 24.9 | (19.2-31.6) |
| 45-64 | 1,078 | 19.8^ | (11.6-31.6) | 57,660 | 16.9 | (13.8-20.6) |
| 65+ | 646 | 20.7 | (12.8-31.6) | 30,644 | 16.3 | (12.8-20.6) |
| Race | | | | | | |
| White, NH | 2,334 | 17.7 | (12.2-25.1) | 150,493 | 19.8 | (17.6-22.2) |
| Black, NH | | # | | 22,275 | 31.3^ | (22.6-41.4) |
| Hispanic | | # | | 15,272 | 30.5^ | (19.3-44.5) |
| Other, NH | | # | | 26,640 | 36.6^ | (18.8-59.1) |
| Sex | | | | | | |
| male | 1,369 | 19.9^ | (11.9-31.3) | 120,224 | 26.1 | (22.7-29.8) |
| female | 1,317 | 17.9^ | (10.7-28.5) | 94,456 | 19.2 | (15.4-23.6) |
| Age-adjusted ^{&} | | 18.5 | (12.7-26.3) | | 23.2 | (20.7-26.0) |

* Sugary drinks include regular soda, and sugar-sweetened drinks.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults who consume fast-food three or more times per week




| | Schuyler County | | | Finger Lakes | | |
|-------------------------------|---------------------|---------|------------|---------------------|---------|------------|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 782 | 5.5 | (2.6-11.3) | 58,515 | 6.2 | (5.0-7.7) |
| Age | | | | | | |
| 18-24 | | # | | 16,427 | 12.9 | (7.6-20.9) |
| 25-34 | | # | | 9,670 | 6.5 | (3.9-10.9) |
| 35-44 | | # | | 5,131 | 3.7 | (2.0-6.7) |
| 45-64 | | # | | 20,550 | 6.2 | (4.2-8.9) |
| 65+ | | # | | 6,737 | 3.6 | (2.4-5.6) |
| Race | | | | | | |
| White, NH | 736 | 5.6 | (2.5-12.0) | 39,988 | 5.3 | (4.1-6.8) |
| Black, NH | | # | | 7,235 | 10.4 | (5.5-18.9) |
| Hispanic | | # | | 6,059 | 12.1^ | (5.1-25.9) |
| Other, NH | | # | | 5,233 | 7.4^ | (2.8-18.0) |
| Sex | | | | | | |
| male | | # | | 43,869 | 9.7 | (7.5-12.5) |
| female | | # | | 14,646 | 3.0 | (2.0-4.5) |
| Age-adjusted ^{&} | | 4.3 | (2.1-8.6) | | 6.1 | (4.9-7.6) |

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults with physician diagnosed diabetes*



| Schuyler County | |
|-----------------|--|
| | |

Finger Lakes


| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| _ | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 1,570 | 10.6 | (7.4-14.9) | 100,031 | 10.0 | (8.7-11.5) |
| Age | | | | | | |
| 18-24 | | # | | | # | |
| 25-34 | | # | | 7,103 | 4.6 | (2.5-8.4) |
| 35-44 | | # | | 6,332 | 4.3 | (2.4-7.6) |
| 45-64 | 464 | 8.3 | (4.8-13.8) | 47,349 | 13.3 | (10.8-16.3) |
| 65+ | 1,065 | 32.4 | (22.4-44.3) | 38,439 | 19.6 | (16.4-23.3) |
| Race | | | | | | |
| White, NH | 1,363 | 9.9 | (7.0-13.9) | 70,994 | 8.9 | (7.7-10.3) |
| Black, NH | | # | | 17,328 | 22.6 | (15.2-32.3) |
| Hispanic | | # | | 6,956 | 13.5 | (8.0-21.9) |
| Other, NH | | # | | 4,753 | 6.3 | (2.5-15.0) |
| Sex | | | | | | |
| male | 853 | 11.6 | (6.7-19.2) | 50,941 | 10.6 | (8.7-12.8) |
| female | 717 | 9.7 | (6.2-14.9) | 49,090 | 9.5 | (7.8-11.5) |
| Age-adjusted ^{&} | | 8.0 | (6.0-10.6) | | 9.1 | (8.0-10.5) |

* As diagnosed by a health professional.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults who are current smokers



Schuyler County


Finger Lakes

| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 3,233 | 22.3 | (15.6-30.9) | 170,861 | 17.4 | (15.5-19.6) |
| Age | | | | | | |
| 18-24 | | # | | 25,688 | 19.3 | (12.8-28.2) |
| 25-34 | | # | | 40,770 | 26.8 | (21.3-33.1) |
| 35-44 | | # | | 27,819 | 19.5 | (14.0-26.5) |
| 45-64 | 1,319 | 23.9^ | (16.1-34.1) | 61,479 | 17.5 | (14.6-20.8) |
| 65+ | 232 | 7.0 | (3.9-12.4) | 14,504 | 7.6 | (5.5-10.5) |
| Race | | | | | | |
| White, NH | 2,862 | 21.3 | (14.5-30.3) | 134,960 | 17.4 | (15.3-19.6) |
| Black, NH | | # | | 20,478 | 26.8^ | (18.6-37.0) |
| Hispanic | | # | | 6,643 | 12.9 | (7.7-20.9) |
| Other, NH | | # | | 8,780 | 11.8^ | (5.6-23.2) |
| Sex | | | | | | |
| male | 1,354 | 19.0^ | (11.3-30.1) | 95,560 | 20.3 | (17.2-23.8) |
| female | 1,880 | 25.5^ | (15.7-38.6) | 75,301 | 14.8 | (12.6-17.4) |
| Age-adjusted ^{&} | | 23.3 | (16.3-32.2) | | 18.2 | (16.1-20.5) |

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults with annual household income less than \$25,000 who are current smokers




| | Schuyler County | | | | Finger Lak | es |
|-------------------------------|---------------------|---------|-------------|---------------------|------------|-------------|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 1,379 | 31.7^ | (19.0-48.0) | 69,503 | 29.6 | (24.7-34.9) |
| Age | | | | | | |
| 18-24 | | # | | 14,247 | 40.5 | (26.5-56.2) |
| 25-34 | | # | | 16,608 | 37.7^ | (22.9-55.2) |
| 35-44 | | # | | 9,252 | 33.3^ | (19.5-50.8) |
| 45-64 | | # | | 23,452 | 32.4 | (24.6-41.5) |
| 65+ | | # | | 5,766 | 10.6 | (6.7-16.2) |
| Race | | | | | | |
| White, NH | 1,065 | 28.6^ | (16.1-45.6) | 48,760 | 29.6 | (24.4-35.3) |
| Black, NH | | # | | 15,001 | 43.1^ | (29.3-58.0) |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 3,255 | 16.8^ | (5.4-41.5) |
| Sex | | | | | | |
| male | 680 | 30.7^ | (13.8-55.0) | 36,273 | 31.2 | (23.4-40.3) |
| female | 699 | 32.9^ | (16.5-54.9) | 33,230 | 27.9 | (22.4-34.1) |
| Age-adjusted ^{&} | | 34.6^ | (20.3-52.3) | | 30.9 | (25.5-36.9) |

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults aged 18-24 years who are current smokers




| | Schuyler County | | | Finger Lakes | | | |
|---|---------------------|-------------|--------|---------------------|--|---------------------------|--|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI | |
| Total Age⁺ 18-24 | | # | | 25,688 | 19.3 | (12.8-28.2) | |
| 25-34 35-44 45-64 65+ | | | | | | | |
| Race | | | | | | | |
| White, NH Black, NH Hispanic Other, NH | | # # # | | 18,243 | 21.5 # # # | (14.1-31.5) | |
| Sex male female | | # # | | 13,906 11,783 | 22.2 [^] 16.8 [^] | (13.2-34.9) (8.6-30.1) | |
| Age-adjusted ^{+,&} | | | | | | | |

& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of women aged 50-74 years receiving breast cancer screening based on recent guidelines^{*}



| | Schuyler County | | | Finger Lakes | | |
|--|---------------------|---|----------------------------|---------------------|-----------------------------|----------------------------|
| F | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total Age ⁺ 18-24 25-34 35-44 45-64 65+ | 2,051 | 69.9^ | (57.1-80.2) | 156,901 | 82.4 | (78.1-86.0) |
| Race White, NH Black, NH Hispanic Other, NH Sex male female | 2,035 2,051 | 70.0 [^] # # # 69.9 [^] | (57.0-80.4) (57.1-80.2) | 136,370 156,901 | 83.2 # # # 82.4 | (78.8-86.8) (78.1-86.0) |
| Age-adjusted ^{+,&} | | | . , | | | . , |

* Mammogram within past 2 years.


& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Prevent Chronic Diseases

Percentage of women aged 50-74 years with annual household income of less than \$25,000 receiving breast cancer screening based on recent guidelines^{*}



| | Schuyler County | | | 1 | Finger Lak | es |
|---------------------------------|---------------------|---------|--------|---------------------|------------|-------------|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | | # | | 32,858 | 75.7^ | (65.5-83.6) |
| Age⁺ | | | | | | |
| 18-24 | | | | | | |
| 25-34 | | | | | | |
| 35-44 | | | | | | |
| 45-64 | | | | | | |
| 65+ | | | | | | |
| Race | | | | | | |
| White, NH | | # | | 24,468 | 77.6 | (68.5-84.7) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | | # | |
| Sex | | | | | | |
| male | | # | | | # | |
| female | | # | | 32,858 | 75.7^ | (65.5-83.6) |
| Age-adjusted ^{+,&} | | | | | | |


* Mammogram within past 2 years.

& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of women aged 21-65 years receiving cervical cancer screening based on recent guidelines^{*}



| | Schuyler County | | | | Finger Lak | es |
|---------------------------------|---------------------|---------|-------------|---------------------|------------|-------------|
| - | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 3,491 | 70.5^ | (58.5-80.2) | 287,036 | 83.6 | (80.3-86.4) |
| Age ⁺ | | | | | | |
| 18-24 | | | | | | |
| 25-34 | | | | | | |
| 35-44 | | | | | | |
| 45-64 | | | | | | |
| 65+ | | | | | | |
| Race | | | | | | |
| White, NH | 3,420 | 71.1^ | (59.0-80.9) | 236,785 | 84.1 | (80.7-87.0) |
| Black, NH | | # | | 23,521 | 86.1^ | (71.0-94.0) |
| Hispanic | | # | | 12,743 | 79.7^ | (63.4-89.9) |
| Other, NH | | # | | | # | |
| Sex | | | | | | |
| male | | # | | | # | |
| female | 3,491 | 70.5^ | (58.5-80.2) | 287,036 | 83.6 | (80.3-86.4) |
| Age-adjusted ^{+,&} | | | | | | |

* Pap test within past 3 years.


& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Prevent Chronic Diseases

Percentage of women aged 21-65 years with annual household income of less than \$25,000 receiving cervical cancer screening based on recent guidelines^{*}



| | Schuyler County | | | Finger Lakes | | |
|---------------------------------|---------------------|---------|--------|---------------------|---------|-------------|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | | # | | 55,266 | 73.1 | (65.6-79.5) |
| Age⁺ | | | | | | |
| 18-24 | | | | | | |
| 25-34 | | | | | | |
| 35-44 | | | | | | |
| 45-64 | | | | | | |
| 65+ | | | | | | |
| Race | | | | | | |
| White, NH | | # | | 37,702 | 73.2 | (64.7-80.2) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | | # | |
| Sex | | | | | | |
| male | | # | | | # | |
| female | | # | | 55,266 | 73.1 | (65.6-79.5) |
| Age-adjusted ^{+,&} | | | | | | |


* Pap test within past 3 years.

& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines^{*}



| | Schuyler County | | Finger Lakes | | | |
|---------------------------------|---------------------|---------|--------------|---------------------|---------|-------------|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 3,549 | 63.6 | (53.7-72.5) | 250,507 | 69.3 | (65.3-73.0) |
| Age⁺ | | | | | | |
| 18-24 | | | | | | |
| 25-34 | | | | | | |
| 35-44 | | | | | | |
| 45-64 | | | | | | |
| 65+ | | | | | | |
| Race | | | | | | |
| White, NH | 3,314 | 62.3^ | (52.3-71.4) | 218,863 | 70.7 | (66.5-74.6) |
| Black, NH | | # | | 18,387 | 75.8^ | (61.6-86.0) |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 9,650 | 51.4^ | (30.2-72.2) |
| Sex | | | | | | |
| male | 1,811 | 68.0^ | (51.8-80.7) | 116,224 | 69.1 | (62.9-74.7) |
| female | 1,738 | 59.6^ | (47.3-70.9) | 134,284 | 69.5 | (64.2-74.3) |
| Age-adjusted ^{+,&} | | | | | | |

* Blood stool test within 1 year, or sigmoidoscopy within 5 years with blood stool test within 3 years, or colonoscopy within 10 years.


& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Prevent Chronic Diseases

Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on recent guidelines^{*}



| | Schuyler County | | | Finger Lakes | | |
|---------------------------------|---------------------|---------|-------------|---------------------|---------|-------------|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 842 | 59.5^ | (42.6-74.4) | 51,562 | 66.4 | (59.0-73.2) |
| Age⁺ | | | | | | |
| 18-24 | | | | | | |
| 25-34 | | | | | | |
| 35-44 | | | | | | |
| 45-64 | | | | | | |
| 65+ | | | | | | |
| Race | | | | | | |
| White, NH | 761 | 57.0^ | (39.7-72.8) | 38,437 | 68.1 | (60.3-74.9) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | | # | |
| Sex | | | | | | |
| male | | # | | 21,576 | 64.5^ | (53.7-74.1) |
| female | | # | | 29,986 | 67.9^ | (57.5-76.7) |
| Age-adjusted ^{+,&} | | | | | | |


* Blood stool test within 1 year, or sigmoidoscopy within 5 years with blood stool test within 3 years, or colonoscopy within 10 years.

& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults who had a test for high blood sugar or diabetes within the past three years



Finger Lakes


| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| _ | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 6,337 | 49.0 | (40.1-58.0) | 473,695 | 54.6 | (51.3-57.9) |
| Age | | | | | | |
| 18-24 | | # | | 26,476 | 20.7 | (14.1-29.4) |
| 25-34 | | # | | 60,826 | 42.3 | (34.5-50.4) |
| 35-44 | | # | | 68,543 | 50.5 | (42.7-58.3) |
| 45-64 | 3,056 | 60.9^ | (50.2-70.6) | 205,917 | 68.5 | (63.9-72.9) |
| 65+ | 1,567 | 75.2 | (65.0-83.2) | 106,367 | 71.4 | (66.0-76.3) |
| Race | | | | | | |
| White, NH | 5,847 | 48.2 | (39.1-57.4) | 397,203 | 57.1 | (54.1-60.2) |
| Black, NH | | # | | 32,135 | 55.7^ | (44.5-66.3) |
| Hispanic | | # | | 21,736 | 48.1^ | (33.4-63.1) |
| Other, NH | | # | | 22,622 | 32.5 | (17.6-52.2) |
| Sex | | | | | | |
| male | 2,843 | 45.2 | (32.2-59.0) | 220,518 | 52.8 | (48.1-57.4) |
| female | 3,494 | 52.6^ | (40.9-64.0) | 253,177 | 56.3 | (51.5-61.0) |
| Age-adjusted ^{&} | | 52.2 | (45.0-59.4) | | 54.1 | (51.2-57.0) |

Ĩ

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults with physician-diagnosed prediabetes




| Schuyler County | Finger Lakes |
|-----------------|--------------|
| | |

| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
|-------------------------------|---------------------|---------|------------|---------------------|---------|------------|
| Total | 821 | 6.4 | (3.8-10.6) | 49,926 | 5.6 | (4.5-7.0) |
| Age | 021 | 0.4 | (3.8-10.0) | 49,920 | 5.0 | (4.5-7.0) |
| 18-24 | | # | | | # | |
| _ | | | | | | |
| 25-34 | | # | | 2,644 | 1.8 | (0.9-3.7) |
| 35-44 | | # | | 5,077 | 3.6 | (1.7-7.7) |
| 45-64 | 574 | 11.4 | (6.2-20.2) | 23,774 | 7.9 | (5.9-10.5) |
| 65+ | 232 | 10.5^ | (4.5-22.7) | 15,665 | 10.3 | (7.2-14.4) |
| Race | | | | | | |
| White, NH | 727 | 6.1 | (3.6-10.2) | 42,109 | 5.9 | (4.7-7.5) |
| Black, NH | | # | | 4,034 | 7.0 | (3.0-15.7) |
| Hispanic | | # | | | # | |
| Other, NH | | # | | | # | |
| Sex | | | | | | |
| male | 348 | 5.4 | (2.4-11.6) | 25,274 | 6.0 | (4.3-8.2) |
| female | 473 | 7.5 | (3.8-14.3) | 24,651 | 5.3 | (3.9-7.2) |
| Age-adjusted ^{&} | | 5.4 | (3.4-8.6) | | 5.5 | (4.4-6.9) |

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults with physician-diagnosed high blood pressure



| Schuyler County | |
|-----------------|--|
| | |


Finger Lakes

| | Schuyler County | | | Filiger Lakes | | |
|-------------------------------|---------------------|---------|-------------|---------------------|---------|-------------|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 4,676 | 32.5 | (26.2-39.5) | 314,920 | 31.7 | (29.2-34.2) |
| Age | - | | . , | - | | |
| 18-24 | | # | | 8,315 | 6.4 | (3.1-12.6) |
| 25-34 | | # | | 20,783 | 13.5 | (9.7-18.4) |
| 35-44 | | # | | 29,430 | 19.8 | (14.6-26.2) |
| 45-64 | 1,940 | 34.6^ | (25.7-44.7) | 134,360 | 37.8 | (33.7-42.1) |
| 65+ | 2,266 | 68.7 | (59.4-76.7) | 118,288 | 60.8 | (55.4-66.0) |
| Race | | | | | | |
| White, NH | 4,024 | 30.3 | (24.1-37.2) | 256,263 | 32.3 | (29.8-34.9) |
| Black, NH | | # | | 32,880 | 44.2 | (34.8-54.0) |
| Hispanic | | # | | 14,927 | 28.5^ | (19.0-40.3) |
| Other, NH | | # | | 10,850 | 14.4^ | (7.9-25.0) |
| Sex | | | | | | |
| male | 2,453 | 33.3^ | (23.6-44.7) | 158,361 | 33.0 | (29.5-36.6) |
| female | 2,223 | 31.7 | (24.4-40.1) | 156,559 | 30.4 | (27.1-33.9) |
| Age-adjusted ^{&} | | 27.6 | (22.0-34.0) | | 29.3 | (27.1-31.5) |

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults with diagnosed high blood pressure taking high blood pressure medication



Schuyler County


Finger Lakes

| | Schayler county | | | Tinger Eakes | | |
|-------------------------------|---------------------|---------|-------------|---------------------|---------|-------------|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 3,533 | 76.9^ | (64.6-85.8) | 243,314 | 77.3 | (73.4-80.8) |
| Age | | | | | | |
| 18-24 | | # | | | # | |
| 25-34 | | # | | 5,688 | 27.4^ | (15.6-43.5) |
| 35-44 | | # | | 16,627 | 56.5^ | (40.3-71.4) |
| 45-64 | 1,181 | 62.4^ | (42.9-78.6) | 108,563 | 80.8 | (74.8-85.7) |
| 65+ | 2,094 | 93.7^ | (80.5-98.2) | 108,613 | 91.9 | (88.6-94.3) |
| Race | | | | | | |
| White, NH | 2,992 | 75.8^ | (62.6-85.5) | 196,305 | 76.7 | (72.2-80.6) |
| Black, NH | | # | | 27,437 | 83.4^ | (71.8-90.9) |
| Hispanic | | # | | | # | |
| Other, NH | | # | | | # | |
| Sex | | | | | | |
| male | 1,743 | 72.5^ | (52.4-86.3) | 113,745 | 71.9 | (66.0-77.1) |
| female | 1,790 | 81.7^ | (68.7-90.0) | 129,569 | 82.8 | (77.4-87.2) |
| Age-adjusted ^{&} | | 57.0^ | (44.9-68.3) | | 58.7 | (53.6-63.6) |

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults with current asthma*



| Schuyler County | |
|-----------------|--|
|-----------------|--|

Finger Lakes


| | Schayler county | | | Tinger Lakes | | |
|-------------------------------|---------------------|---------|------------|---------------------|---------|-------------|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 1,106 | 7.5 | (4.7-11.8) | 112,310 | 11.3 | (9.6-13.3) |
| Age | | | | | | . , |
| 18-24 | | # | | 20,356 | 15.3 | (8.9-25.2) |
| 25-34 | | # | | 20,425 | 13.3 | (9.3-18.6) |
| 35-44 | | # | | 13,324 | 9.1 | (5.4-15.0) |
| 45-64 | 762 | 13.7 | (7.9-22.7) | 40,947 | 11.6 | (9.0-14.7) |
| 65+ | | # | | 15,560 | 8.0 | (5.9-10.6) |
| Race | | | | | | |
| White, NH | 1,000 | 7.4 | (4.5-11.8) | 81,926 | 10.4 | (8.7-12.4) |
| Black, NH | | # | | 11,764 | 15.4 | (9.3-24.3) |
| Hispanic | | # | | 7,513 | 14.3 | (8.6-23.0) |
| Other, NH | | # | | 11,106 | 14.8^ | (6.3-30.8) |
| Sex | | | | | | |
| male | 454 | 6.2 | (2.8-13.4) | 41,563 | 8.7 | (6.6-11.3) |
| female | 653 | 8.9 | (5.1-15.0) | 70,747 | 13.8 | (11.2-16.8) |
| Age-adjusted ^{&} | | 8.1 | (4.6-13.7) | | 11.2 | (9.4-13.3) |

* Physician diagnosed asthma.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percent of adults with cholesterol checked*



Schuyler County

Finger Lakes


| | Schayler county | | | Thiger Eakes | | |
|-------------------------------|---------------------|---------|-------------|---------------------|---------|-------------|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 10,249 | 74.0^ | (63.6-82.3) | 798,681 | 82.4 | (79.3-85.0) |
| Age | - | | | - | | |
| 18-24 | | # | | 49,911 | 42.6^ | (31.0-55.0) |
| 25-34 | | # | | 95,361 | 64.6 | (55.3-72.9) |
| 35-44 | | # | | 117,490 | 80.4 | (73.0-86.1) |
| 45-64 | 4,842 | 86.6 | (77.4-92.4) | 336,574 | 95.0 | (93.5-96.3) |
| 65+ | 3,132 | 97.2 | (88.8-99.3) | 188,822 | 97.7 | (96.4-98.5) |
| Race | | | | | | |
| White, NH | 9,445 | 73.7^ | (62.8-82.3) | 646,121 | 83.9 | (81.6-86.0) |
| Black, NH | | # | | 59,974 | 80.5^ | (70.4-87.8) |
| Hispanic | | # | | 42,285 | 84.3^ | (74.0-91.0) |
| Other, NH | | # | | 50,300 | 67.0^ | (41.4-85.4) |
| Sex | | | | | | |
| male | 4,795 | 65.6^ | (49.8-78.6) | 375,301 | 80.0 | (75.5-83.8) |
| female | 5,454 | 83.5^ | (72.4-90.7) | 423,380 | 84.6 | (80.3-88.1) |
| Age-adjusted ^{&} | | 71.3 | (63.9-77.8) | | 80.0 | (77.1-82.5) |

* As diagnosed by a health professional.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults 18 years and older with elevated cholesterol*



Schuyler County

Finger Lakes

| | Solidyier County | | | i inger zakes | | |
|-------------------------------|---------------------|---------|-------------|---------------------|---------|-------------|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 4,132 | 40.6 | (33.9-47.8) | 300,637 | 37.9 | (35.0-40.9) |
| Age | | | | | | |
| 18-24 | | # | | | # | |
| 25-34 | | # | | 17,828 | 18.8 | (12.5-27.2) |
| 35-44 | | # | | 30,289 | 25.8 | (19.3-33.5) |
| 45-64 | 1,891 | 39.6^ | (30.0-50.1) | 139,330 | 41.7 | (37.3-46.4) |
| 65+ | 1,719 | 55.0^ | (44.3-65.3) | 102,175 | 54.6 | (49.2-60.0) |
| Race | | | | | | |
| White, NH | 3,772 | 40.3 | (33.5-47.5) | 241,155 | 37.6 | (34.7-40.6) |
| Black, NH | | # | | 23,513 | 39.5^ | (29.4-50.5) |
| Hispanic | | # | | 15,023 | 35.5^ | (23.3-50.0) |
| Other, NH | | # | | 20,947 | 41.7^ | (24.0-61.8) |
| Sex | | | | | | |
| male | 2,289 | 48.3^ | (37.3-59.5) | 151,064 | 40.6 | (36.4-44.9) |
| female | 1,843 | 33.9 | (26.2-42.6) | 149,573 | 35.6 | (31.6-39.7) |
| Age-adjusted ^{&} | | 27.8 | (22.2-34.1) | | 32.7 | (29.1-36.6) |


* As diagnosed by a health professional.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Schuyler Prevent Chronic Diseases

Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition^{*}



| | Schuyler County | | | Finger Lakes | | |
|-------------------------------|---------------------|---------|------------|---------------------|---------|-------------|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 431 | 5.2 | (3.2-8.3) | 51,667 | 10.0 | (7.8-12.8) |
| Age | | | | | | |
| 18-24 | | # | | | # | |
| 25-34 | | # | | | # | |
| 35-44 | | # | | 3,726 | 6.8 | (3.1-14.2) |
| 45-64 | | # | | 26,449 | 12.7 | (9.1-17.5) |
| 65+ | 255 | 9.1 | (5.2-15.3) | 12,705 | 8.1 | (5.8-11.2) |
| Race | | | | | | |
| White, NH | 431 | 5.8 | (3.6-9.3) | 34,619 | 8.2 | (6.3-10.7) |
| Black, NH | | # | | 6,746 | 15.1^ | (8.2-26.3) |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 8,215 | 31.5^ | (12.6-59.6) |
| Sex | | | | | | |
| male | 212 | 5.8 | (2.8-11.8) | 23,976 | 9.8 | (6.9-13.9) |
| female | 219 | 4.7 | (2.6-8.6) | 27,691 | 10.2 | (7.2-14.2) |
| Age-adjusted ^{&} | | 3.0 | (1.8-4.8) | | 9.5 | (6.8-13.3) |


* Courses for high blood pressure, arthritis, or diabetes.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Schuyler Prevent Chronic Diseases

Percentage of adults with arthritis*



Schuyler County

Finger Lakes

| | Senayier County | | | Tinger Eakes | | |
|-------------------------------|---------------------|---------|-------------|---------------------|---------|-------------|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 4,389 | 29.8 | (23.9-36.5) | 273,908 | 27.5 | (25.1-30.0) |
| Age | | | | | | |
| 18-24 | | # | | | # | |
| 25-34 | | # | | 11,753 | 7.6 | (4.9-11.6) |
| 35-44 | | # | | 27,015 | 18.2 | (12.9-25.1) |
| 45-64 | 2,391 | 42.6^ | (33.1-52.7) | 122,234 | 34.4 | (30.2-38.9) |
| 65+ | 1,618 | 49.1^ | (38.7-59.5) | 107,901 | 55.4 | (50.3-60.3) |
| Race | | | | | | |
| White, NH | 4,071 | 29.8 | (23.7-36.8) | 230,014 | 29.0 | (26.4-31.7) |
| Black, NH | | # | | 21,965 | 28.9 | (21.1-38.3) |
| Hispanic | | # | | 10,718 | 20.5 | (13.4-30.1) |
| Other, NH | | # | | 11,212 | 14.9^ | (6.7-29.9) |
| Sex | | | | | | |
| male | 2,044 | 27.8^ | (19.4-38.1) | 118,312 | 24.6 | (21.3-28.2) |
| female | 2,345 | 31.9 | (24.0-40.9) | 155,596 | 30.1 | (26.7-33.8) |
| Age-adjusted ^{&} | | 25.8 | (20.9-31.3) | | 25.3 | (23.2-27.5) |


* Includes rheumatoid arthritis, gout, lupus, or fibromyalgia.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Promote a Healthy and Safe Environment

Percentage of adults who consider their neighborhood suitable for walking and physical activity



Schuyler County

Finger Lakes


| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
|-------------------------------|---------------------|---------|-------------|---------------------|---------|-------------|
| Total | 13,447 | 94.7 | (89.9-97.3) | 897,106 | 93.4 | (92.0-94.6) |
| Age | | | | | | |
| 18-24 | | # | | 114,680 | 89.2 | (81.1-94.1) |
| 25-34 | | # | | 134,928 | 90.1 | (85.9-93.2) |
| 35-44 | | # | | 134,641 | 95.3 | (92.3-97.2) |
| 45-64 | 5,352 | 96.9 | (92.6-98.8) | 326,991 | 94.9 | (93.0-96.3) |
| 65+ | 3,021 | 94.5 | (88.6-97.4) | 176,297 | 94.4 | (92.1-96.1) |
| Race | | | | | | |
| White, NH | 12,370 | 94.3 | (89.1-97.1) | 723,466 | 94.7 | (93.4-95.8) |
| Black, NH | | # | | 58,381 | 80.4^ | (70.2-87.8) |
| Hispanic | | # | | 45,612 | 89.3 | (79.9-94.6) |
| Other, NH | | # | | 69,648 | 95.6 | (90.2-98.0) |
| Sex | | | | | | |
| male | 6,746 | 97.0 | (90.4-99.1) | 435,275 | 93.9 | (91.7-95.5) |
| female | 6,701 | 92.6 | (84.2-96.7) | 461,832 | 92.9 | (91.0-94.5) |
| Age-adjusted ^{&} | | 92.2 | (82.5-96.7) | | 93.3 | (91.9-94.5) |

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Promote a Health and Safe Environment

Percentage of adults aged 65+ years with at least one fall in the past 12 months



| | Scl | huyler Cou | nty | Finger Lakes | | |
|---|-----------------------|---|---|----------------------------|--------------------------------|---|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total Age ⁺ 18-24 25-34 35-44 45-64 65+ Race | 1,607 | 50.6^ | (40.1-61.1) | 57,486 | 30.5 | (25.8-35.7) |
| White, NH Black, NH Hispanic Other, NH Sex male female | 1,181 1,058 549 | 43.9 [^] # # 62.2 [^] 37.3 [^] | (34.0-54.2) (47.5-75.0) (24.1-52.7) | 43,381 26,845 30,641 | 26.6 # # 31.6 29.6 | (22.9-30.7) (25.8-38.1) (22.7-37.6) |
| Age-adjusted ^{+,&} | 545 | 57.5 | (27:2 32.7) | 50,041 | 23.0 | (22.7 37.0) |


& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Promote a Health and Safe Environment

Among adults aged 65+ years with at least one fall in the past 12 months, percent that resulted in injury^{*}



| | Scl | huyler Cou | nty | Finger Lakes | | |
|---|---------------------|----------------------------------|-------------|---------------------------|--|---|
| - | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total Age ⁺ 18-24 25-34 35-44 45-64 65+ Race | 523 | 32.6^ | (18.1-51.3) | 19,362 | 33.9 | (26.1-42.8) |
| White, NH Black, NH Hispanic Other, NH Sex male female | 350 | 29.6 [^] # # # | (15.8-48.7) | 16,678 8,530 10,832 | 38.8 # # 32.0 [^] 35.6 [^] | (31.4-46.7) (22.0-43.9) (24.2-49.0) |
| Age-adjusted ^{+,&} | | | | | | |


* Falls that limited regular activities or lead to doctor visits.

& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV)




| | Scl | huyler Cou | nty | Finger Lakes | | |
|---|---------------------|--|---|--------------------------------------|--|--|
| - | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total Age ⁺ 18-24 25-34 35-44 45-64 65+ Race | 1,271 | 24.4 | (17.7-32.6) | 106,026 | 32.9 | (28.9-37.2) |
| White, NH Black, NH Hispanic Other, NH Sex male female | 1,242 606 666 | 24.3 # # 25.8 [^] 23.3 [^] | (17.5-32.6) (16.0-38.7) (14.8-34.6) | 83,817 14,865 56,263 49,763 | 31.1 57.1 [^] # 36.6 29.6 | (26.9-35.6) (40.3-72.4) (30.6-43.1) (24.4-35.3) |
| Age-adjusted ^{+,&} | | | | | | |

& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections Percentage of adults aged 18+ years with flu immunization in the past year^{*}



Schuyler County

Finger Lakes


| | ochayler county | | | | | |
|-------------------------------|---------------------|---------|-------------|---------------------|---------|-------------|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 5,412 | 37.4 | (30.5-44.8) | 556,353 | 57.7 | (54.7-60.6) |
| Age | | | | | | |
| 18-24 | | # | | 52,329 | 40.8^ | (30.1-52.4) |
| 25-34 | | # | | 65,604 | 43.6 | (35.2-52.4) |
| 35-44 | | # | | 69,933 | 49.5 | (41.8-57.1) |
| 45-64 | 1,938 | 35.1 | (26.7-44.5) | 205,087 | 59.5 | (55.2-63.7) |
| 65+ | 2,256 | 68.7^ | (58.1-77.7) | 157,801 | 83.2 | (80.0-85.9) |
| Race | | | | | | |
| White, NH | 4,943 | 36.9 | (29.8-44.6) | 443,049 | 57.6 | (54.7-60.4) |
| Black, NH | | # | | 40,970 | 57.1^ | (47.0-66.7) |
| Hispanic | | # | | 30,516 | 59.7^ | (46.2-71.9) |
| Other, NH | | # | | 41,817 | 57.3^ | (35.8-76.4) |
| Sex | | | | | | |
| male | 2,697 | 38.0^ | (27.4-49.8) | 260,048 | 55.9 | (51.8-60.0) |
| female | 2,715 | 36.8 | (28.1-46.4) | 296,305 | 59.3 | (54.9-63.5) |
| Age-adjusted ^{&} | | 36.4 | (29.4-43.9) | | 56.0 | (52.9-59.1) |

* Includes flu shot or flu spray

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections Percentage of adults aged 65+ years with flu immunization in the past year^{*}



| | Scl | huyler Cou | nty | Finger Lakes | | |
|--|-------------------------|--|---|-----------------------------|--------------------------------|---|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total Age ⁺ 18-24 25-34 35-44 45-64 65+ Race | 2,256 | 68.7^ | (58.1-77.7) | 157,801 | 83.2 | (80.0-85.9) |
| White, NH Black, NH Hispanic Other, NH Sex male female | 1,926 1,242 1,014 | 69.3 # # 70.2 [^] 67.0 [^] | (59.4-77.7) (54.8-82.1) (52.0-79.2) | 136,667 72,669 85,131 | 83.2 # # 85.0 81.7 | (80.1-85.9) (80.9-88.3) (76.8-85.8) |
| Age-adjusted ^{+,&} | 1,014 | 67.0 | (32.075.2) | 55,151 | 51.7 | (70.0 05.0) |

* Includes flu shot or flu spray


& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections

Percentage of adults aged 65+ years with pneumococcal immunization




| | Scl | huyler Cou | nty | Finger Lakes | | |
|---|---------------------|--|----------------------------|---------------------|---------------------|----------------------------|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total Age⁺ 18-24 | 2,203 | 73.9 | (64.1-81.7) | 135,060 | 74.0 | (68.5-78.8) |
| 25-34 35-44 45-64 65+ Race | | | | | | |
| White, NH Black, NH Hispanic Other, NH Sex | 1,969 | 74.2 # # # | (65.2-81.5) | 122,920 | 78.0 # # # | (74.0-81.6) |
| male female Age-adjusted ^{+,&} | 1,095 1,108 | 72.8 [^] 74.9 [^] | (59.5-83.0) (59.5-85.9) | 56,221 78,839 | 68.3 78.7 | (61.1-74.7) (70.1-85.3) |

& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of women aged 18-44 years who had dental visit within the last year




| | Schuyler County | | | Finger Lakes | | |
|--|---------------------|-------------|--------|---------------------|---------------------|-------------|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total Age ⁺ 18-24 25-34 35-44 | | # | | 158,478 | 76.1 | (67.5-83.1) |
| 45-64 65+ Race White, NH Black, NH Hispanic Other, NH Sex | | # # # | | 120,797 | 78.0 # # # | (71.5-83.4) |
| male female Age-adjusted ^{+,&} | | # # | | 158,478 | # 76.1 | (67.5-83.1) |

& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Women aged 18-44 years who saw a doctor for a routine checkup within the last year




| | Schuyler County | | | Finger Lakes | | |
|---------------------------------|---------------------|---------|--------|---------------------|---------|-------------|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | | # | | 135,013 | 61.9 | (54.3-68.9) |
| Age⁺ | | | | | | |
| 18-24 | | | | | | |
| 25-34 | | | | | | |
| 35-44 | | | | | | |
| 45-64 | | | | | | |
| 65+ | | | | | | |
| Race | | | | | | |
| White, NH | | # | | 99,001 | 61.6 | (54.7-68.1) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | | # | |
| Sex | | | | | | |
| male | | # | | | # | |
| female | | # | | 135,013 | 61.9 | (54.3-68.9) |
| Age-adjusted ^{+,&} | | | | | | |

& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of women aged 18-64 years with health care coverage




| | Scl | nuyler Cou | nty | Finger Lakes | | |
|---|---------------------|---|----------------------------|--|--|---|
| - | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total Age ⁺ 18-24 25-34 35-44 45-64 65+ Race | 4,806 | 85.3^ | (72.6-92.8) | 364,324 | 91.0 | (88.6-93.0) |
| White, NH Black, NH Hispanic Other, NH Sex male female | 4,661 4,806 | 85.0 [^] # # # 85.3 [^] | (72.0-92.6) (72.6-92.8) | 282,703 28,417 16,002 37,203 364,324 | 91.1 85.7 [^] 91.5 [^] 94.7 [^] # 91.0 | (88.4-93.2) (74.1-92.6) (74.4-97.5) (84.6-98.3) (88.6-93.0) |
| Age-adjusted ^{+,&} | | | | | | |

& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of women aged 18-44 years whose health care provider discussed planning a healthy pregnancy



| | Schuyler County | | | Finger Lakes | | |
|---|---------------------|-------------|--------|---------------------|---------------------|-------------|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total Age ⁺ 18-24 25-34 35-44 45-64 | | # | | 89,890 | 43.1 | (35.9-50.6) |
| 65+ Race White, NH Black, NH Hispanic Other, NH Sex | | # # # | | 72,684 | 46.8 # # # | (39.7-54.0) |
| male female Age-adjusted ^{+,&} | | # # | | 89,890 | # 43.1 | (35.9-50.6) |


& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Promote Mental Health and Prevention Substance Abuse

Percentage of adults with poor mental health for 14 or more days in the last month



Finger Lakes

| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|----------------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 2,454 | 16.9 | (10.3-26.4) | 120,784 | 12.3 | (10.6-14.2) |
| Age | | | | | | |
| 18-24 | | # | | 18,248 | 13.9 | (8.2-22.4) |
| 25-34 | | # | | 21,945 | 14.4 | (10.5-19.4) |
| 35-44 | | # | | 17,847 | 12.2 | (8.1-18.0) |
| 45-64 | 886 | 16.0^ | (9.3-26.1) | 50,391 | 14.4 | (11.5-17.7) |
| 65+ | 311 | 10.0^ | (4.6-20.5) | 11,819 | 6.3 | (4.5-8.7) |
| Race | | | | | | |
| White, NH | 2,175 | 16.2^ | (9.4-26.5) | 90,447 | 11.6 | (9.9-13.6) |
| Black, NH | | # | | 12,561 | 16.5 | (10.8-24.5) |
| Hispanic | | # | | 9,450 | 18.5^ | (11.2-29.0) |
| Other, NH | | # | | 8,325 | 11.1° | (4.0-27.3) |
| Sex | | | | | | |
| male | 1,209 | 16.9^ | (8.4-31.3) | 49,351 | 10.4 | (8.2-13.2) |
| female | 1,245 | 16.9^ | (8.4-31.0) | 71,433 | 14.1 | (11.7-16.9) |
| Age-adjusted ^{&} | | 16.0 | (10.1-24.6) | | 12.5 | (10.7-14.5) |


Ĩ

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Promote Mental Health and Prevention Substance Abuse

Percentage of adult binge drinking during the past month*



Schuyler County

Finger Lakes

| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|------------|-----------|---------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 1,463 | 10.2 | (5.8-17.3) | 154,347 | 16.1 | (14.1-18.4) |
| Age | | | | | | |
| 18-24 | | # | | 32,152 | 24.8 | (17.7-33.6) |
| 25-34 | | # | | 37,099 | 24.8 | (19.2-31.4) |
| 35-44 | | # | | 26,167 | 18.7 | (12.6-26.7) |
| 45-64 | 461 | 8.4 | (4.2-16.0) | 52,235 | 15.2 | (12.0-19.1) |
| 65+ | | # | | 5,899 | 3.2 | (2.2-4.5) |
| Race | | | | | | |
| White, NH | 1,383 | 10.4 | (5.8-18.1) | 129,788 | 17.1 | (14.7-19.6) |
| Black, NH | | # | | 8,750 | 12.2 | (7.4-19.5) |
| Hispanic | | # | | 10,236 | 19.8^ | (12.0-30.9) |
| Other, NH | | # | | 5,572 | 7.5 | (3.2-16.9) |
| Sex | | | | | | |
| male | 890 | 12.7 | (5.6-26.1) | 99,542 | 21.7 | (18.3-25.4) |
| female | 573 | 7.9 | (3.8-15.5) | 54,805 | 11.0 | (8.8-13.7) |
| Age-adjusted ^{&} | | 12.8 | (7.3-21.6) | | 16.9 | (14.6-19.5) |

Í


* Men who have five or more drinks , women having four or more drinks on one occasion

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Schuyler Promote Mental Health and Prevention Substance Abuse

Percentage of cigarette smoking among adults who report poor mental health



_.

. .

| | Schuyler County | | | Finger Lakes | | | |
|-------------------------------|---------------------|---------|--------|---------------------|---------|-------------|--|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI | |
| Total | | # | | 41,696 | 34.8 | (27.8-42.5) | |
| Age | | | | | | | |
| 18-24 | | # | | | # | | |
| 25-34 | | # | | 10,400 | 48.1^ | (33.0-63.5) | |
| 35-44 | | # | | 7,009 | 39.7^ | (20.7-62.3) | |
| 45-64 | | # | | 19,075 | 38.1^ | (27.3-50.2) | |
| 65+ | | # | | 740 | 6.3 | (3.3-11.9) | |
| Race | | | | | | | |
| White, NH | | # | | 32,992 | 36.8 | (28.9-45.4) | |
| Black, NH | | # | | | # | | |
| Hispanic | | # | | | # | | |
| Other, NH | | # | | | # | | |
| Sex | | | | | | | |
| male | | # | | 19,261 | 39.2^ | (27.0-53.0) | |
| female | | # | | 22,435 | 31.7 | (24.4-40.1) | |
| Age-adjusted ^{&} | | # | | | 33.1 | (26.5-40.5) | |

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

.