eBRFSS 2013-2014 Health Indicators: Tioga

| General Description | |
|---|---|
| Improve Health Status and Reduce Health Disparities | |
| Percentage of adults who did not receive medical care because of cost | 3 |
| Percentage of adults who have a regular health care provider | |
| Percentage of adults who had a dentist visit within the past year | 5 |
| Percentage of adults living with a disability | |
| Percentage of adults aged 18-64 years with healthcare coverage | |
| Percentage of adults aged 18-64 years who saw a doctor for a routine checkup within the last year | 8 |
| Percentage of adults experiencing housing insecurity in the past 12 months | 9 |
| Percentage of adults experiencing food insecurity in the past 12 months | |
| Percentage of adults with poor self-reported health | |
| Percentage of adults who report 14 or more days of poor physical health | |
| Prevent Chronic Diseases | |
| Percentage of adults who are obese | |
| Percentage of adults with an annual household income less than \$25,000 who are obese | |
| Percentage of adults living with a disability who are obese | |
| Percentage of adults overweight or obese | |
| Percentage of adults who participated in leisure time physical activity in the past 30 days | 17 |
| Percentage of adults who consume one or more sugary drinks daily | |
| Percentage of adults who consume fast-food three or more times per week | 19 |
| Percentage of adults with physician diagnosed diabetes | |
| Percentage of adults who are current smokers | 21 |
| Percentage of adults with annual household income less than \$25,000 who are current smokers | |
| Percentage of adults aged 18-24 years who are current smokers | |
| Percentage of women aged 50-74 years receiving breast cancer screening based on recent guidelines | |
| Percentage of women aged 50-74 years with annual household income of less than \$25,000 receiving breast cancer screening | ng based on recent |
| guidelines | |
| Percentage of women aged 21-65 years receiving cervical cancer screening based on recent guidelines | |
| Percentage of women aged 21-65 years with annual household income of less than \$25,000 receiving cervical cancer screen | |
| guidelines | |
| Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines | 20 |
| | |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening | g based on recent |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines | g based on recent 29 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years | g based on recent 29 30 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes | g based on recent 29 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure | g based on recent 29 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication | g based on recent 29 30 31 32 33 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma | g based on recent 29 30 31 32 33 34 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines | g based on recent 29 30 31 32 33 34 34 35 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked Percentage of adults 18 years and older with elevated cholesterol | g based on recent 29 30 31 32 33 34 35 36 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked Percentage of adults 18 years and older with elevated cholesterol Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition | g based on recent 29 30 31 32 33 34 34 35 36 37 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines | g based on recent 29 30 31 32 33 34 35 36 37 38 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines | g based on recent 29 30 31 32 33 34 35 36 37 38 39 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines | g based on recent 29 30 31 32 33 34 35 36 37 38 39 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines | g based on recent 29 30 31 32 33 34 35 36 37 38 39 39 39 40 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines | g based on recent 29 30 31 32 33 34 35 36 37 38 39 39 40 40 41 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines | g based on recent 29 30 31 32 33 34 35 36 37 38 39 39 40 40 41 42 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines | g based on recent 29 30 31 32 33 34 35 36 37 38 39 40 40 41 42 42 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years Percentage of adults with physician-diagnosed prediabetes Percentage of adults with physician-diagnosed high blood pressure Percentage of adults with diagnosed high blood pressure taking high blood pressure medication Percentage of adults with current asthma Percentage of adults with cholesterol checked Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition Percentage of adults with arthritis Promote a Healthy and Safe Environment Percentage of adults aged 65+ years with at least one fall in the past 12 months. Among adults aged 65+ years with at least one fall in the past 12 months. Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV) Percentage of adults aged 18+ years with flu immunization in the past year | g based on recent 29 30 31 32 33 34 35 36 37 38 39 40 40 41 42 42 42 43 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines | g based on recent 29 30 31 32 33 34 35 36 37 38 39 40 40 41 42 42 42 43 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines | g based on recent 29 30 31 32 33 34 35 36 36 37 38 39 39 40 40 41 42 42 42 42 43 44 45 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines | g based on recent 29 30 31 32 33 34 35 36 37 38 39 40 40 41 42 42 42 42 43 44 45 46 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines. Percentage of adults who had a test for high blood sugar or diabetes within the past three years | g based on recent 29 30 31 32 33 34 34 35 36 37 38 39 40 40 41 42 42 42 42 42 43 44 44 45 46 46 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines. Percentage of adults who had a test for high blood sugar or diabetes within the past three years | g based on recent 29 30 31 32 33 34 34 35 36 37 38 39 40 40 41 42 42 42 42 42 43 44 44 45 46 46 47 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines Percentage of adults who had a test for high blood sugar or diabetes within the past three years | g based on recent 29 30 31 32 33 34 35 36 37 38 39 40 40 41 42 42 42 42 42 43 44 44 45 46 46 47 48 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines | g based on recent 29 30 31 32 33 34 35 36 37 38 39 39 40 40 41 42 42 42 42 42 43 44 44 45 46 46 47 48 49 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines | g based on recent 29 30 31 32 33 34 35 36 37 38 39 39 40 40 41 42 42 42 42 43 40 41 42 42 42 43 44 45 46 46 47 48 49 50 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines | g based on recent 29 30 31 32 33 34 35 36 37 38 39 39 40 40 41 41 42 42 42 43 44 45 44 45 46 46 47 48 49 50 50 |
| Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening guidelines | g based on recent 29 30 31 32 33 34 35 36 37 38 39 40 41 42 42 42 42 43 44 45 46 46 47 48 49 50 50 50 50 |

General Description

The New York State Expanded Behavioral Risk Factor Surveillance System 2013-2014 (2014 eBRFSS) was design to supplement the Center for Disease Control and Prevention (CDC) Behavioral Risk Factor Surveillance System (BRFSS), which is conducted annually in New York State. The purpose of the 2014 eBRFSS was to produce local information on key public health issues. Data for the project were collected from April 15, 2013 to May 10, 2014. The 2014 eBRFSS reached both households with landline telephones and households which only had cell phones. Like the annual BRFSS, New York State's 2014 eBRFSS was designed to be representative of the non-institutionalized adult household population, aged 18 years and older.

Reported Health Indicators

There are a total of 50 health indicators included in the report. The health indicators include 10 measures included in the New York State Department of Health's Prevention Agenda, NYS Health Innovation Plan (SHIP), Delivery System Reform Incentive Payment (DSRIP) Program, or the NYS Community Health Indicator Reports (CHIRS). An additional 40 important health indicators are also included. The indicators are organized by the Priority Area of the Prevention Agenda and are reported statewide, by region, and by county.

Data Collection Methodology, Sampling and Weighting

The data collection methods for the 2014 eBRFSS aligned with the CDC's standard annual BRFSS methods. The 2014 eBRFSS questionnaire used standard questions from the BRFSS. Survey interviews were conducted in both English and Spanish. There were 31,690 completed interviews.


The sampling plan for the eBRFSS was developed to result in valid health indicator estimates for New York City (NYC) and each of the 57 counties outside NYC. The sampling approach stratified landlines by county and implemented a disproportionate stratified sampling design that accounted for both listed-households and unlisted numbers. Cell phone households were accounted for through additional strata that covered contiguous groups of counties.

Data were weighted to represent the population using a methodology known as iterative proportional fitting (raking). Raking adjusted the data so that groups underrepresented in the sample were more accurately represented in the estimated rates. Raking also enabled surveys conducted by cell phone to be included in the weighting and to accurately match sample distributions of specified demographic characteristics of the population. The raking categories included in the eBRFSS were: age, gender, race/ethnicity, education, marital status, owner/renter status, telephone usage (i.e. landline and cell phone combinations), and county of residence.

Limitations of Data Use:

During data analysis, criteria were applied to suppress rates based on too few surveys to be reliable and to identify rates that are highly variable. Rates in which less than 10 respondents were in the numerator or less than 50 respondents were in the denominator were considered unreliable and suppressed from reporting. Rates considered to have high variability due to having confidence limits greater than $\pm 10\%$ are noted.

Percentage of adults who did not receive medical care because of ${\rm cost}^*$



| Tioga County | |
|--------------|--|
| | |

Southern Tier


| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|------------|-----------|---------|-------------|
| _ | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 3,844 | 9.8 | (6.5-14.7) | 36,229 | 10.0 | (8.2-12.2) |
| Age | | | | | | |
| 18-24 | | # | | 3,485 | 6.2 | (3.1-11.9) |
| 25-34 | | # | | 7,509 | 12.3 | (7.7-19.0) |
| 35-44 | | # | | 6,772 | 14.2 | (8.3-23.3) |
| 45-64 | 2,208 | 14.6 | (8.3-24.6) | 16,052 | 13.5 | (10.3-17.5) |
| 65+ | | # | | 2,128 | 2.9 | (1.8-4.6) |
| Race | | | | | | |
| White, NH | 3,688 | 10.5 | (6.9-15.7) | 28,746 | 9.5 | (7.7-11.8) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 1,904 | 7.0 | (3.2-14.7) |
| Sex | | | | | | |
| male | 2,339 | 12.2 | (6.6-21.5) | 16,682 | 9.4 | (7.2-12.3) |
| female | 1,505 | 7.6 | (4.5-12.4) | 19,547 | 10.6 | (7.9-13.9) |
| Age-adjusted ^{&} | | 10.3 | (6.6-15.8) | | 10.7 | (8.6-13.2) |

* Among people needing to see a doctor in the past 12 months

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults who have a regular health care provider



Tioga County

Southern Tier


| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 32,448 | 83.4 | (75.3-89.3) | 305,033 | 85.0 | (82.4-87.2) |
| Age | | | | | | |
| 18-24 | | # | | 42,341 | 75.8^ | (64.6-84.3) |
| 25-34 | | # | | 47,402 | 77.7 | (69.2-84.4) |
| 35-44 | | # | | 37,697 | 79.2 | (71.3-85.4) |
| 45-64 | 12,910 | 86.0 | (76.4-92.1) | 105,441 | 89.5 | (86.3-92.0) |
| 65+ | 8,020 | 94.4 | (87.8-97.6) | 67,547 | 93.9 | (90.9-95.9) |
| Race | | | | | | |
| White, NH | 29,254 | 84.0 | (76.4-89.4) | 253,550 | 84.9 | (82.3-87.2) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 22,327 | 81.9^ | (68.4-90.4) |
| Sex | | | | | | |
| male | 13,766 | 72.1 | (57.6-83.0) | 136,951 | 77.5 | (73.5-81.1) |
| female | 18,682 | 94.4 | (89.5-97.1) | 168,081 | 92.2 | (88.9-94.6) |
| Age-adjusted ^{&} | | 81.5 | (72.3-88.2) | | 84.1 | (81.3-86.5) |

1

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults who had a dentist visit within the past year



| Tioga | County | |
|-------|--------|--|
| TIUga | County | |


Southern Tier

| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 19,673 | 56.2 | (49.0-63.1) | 220,423 | 66.7 | (63.4-69.8) |
| Age | | | | | | |
| 18-24 | | # | | 29,851 | 66.4 | (54.2-76.8) |
| 25-34 | | # | | 36,594 | 65.9^ | (55.2-75.2) |
| 35-44 | | # | | 28,479 | 64.5 | (54.6-73.3) |
| 45-64 | 8,073 | 57.4 | (48.0-66.3) | 77,932 | 68.5 | (63.7-73.0) |
| 65+ | 4,424 | 54.0 | (44.8-63.0) | 44,963 | 65.7 | (60.9-70.2) |
| Race | | | | | | |
| White, NH | 19,228 | 58.6 | (51.4-65.5) | 188,016 | 66.6 | (63.2-69.7) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 16,234 | 73.3^ | (57.0-85.0) |
| Sex | | | | | | |
| male | 8,667 | 52.4 | (40.6-64.0) | 102,550 | 62.3 | (57.5-67.0) |
| female | 11,005 | 59.5 | (51.0-67.5) | 117,873 | 71.0 | (66.6-75.0) |
| Age-adjusted ^{&} | | 55.6 | (46.3-64.5) | | 66.4 | (62.8-69.8) |

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults living with a disability^{*}



| Tioga County | |
|--------------|--|
| | |

Southern Tier


| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 10,532 | 27.3 | (21.7-33.8) | 89,618 | 25.2 | (22.5-28.1) |
| Age | | | | | | |
| 18-24 | | # | | 4,554 | 8.2 | (4.3-15.1) |
| 25-34 | | # | | 12,577 | 21.1^ | (13.6-31.4) |
| 35-44 | | # | | 8,889 | 19.4 | (13.5-27.2) |
| 45-64 | 4,819 | 32.5 | (24.0-42.2) | 36,485 | 30.8 | (26.7-35.3) |
| 65+ | 3,777 | 44.8 | (36.3-53.7) | 26,446 | 36.6 | (31.9-41.5) |
| Race | | | | | | |
| White, NH | 9,574 | 27.8 | (22.5-33.7) | 77,336 | 26.0 | (23.3-28.9) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 5,878 | 23.1 | (12.8-38.2) |
| Sex | | | | | | |
| male | 4,513 | 23.8 | (15.9-33.9) | 43,459 | 24.9 | (21.0-29.2) |
| female | 6,020 | 30.8 | (23.8-38.7) | 46,159 | 25.5 | (21.9-29.5) |
| Age-adjusted ^{&} | | 25.8 | (20.2-32.4) | | 24.6 | (22.0-27.5) |

* Limited activities (physical, mental, or emotional problems); need of special equipment (e.g. cane, wheel chair).

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults aged 18-64 years with healthcare coverage




| | Tioga County | | | Southern Tier | | |
|---|---------------------|-----------------------------|----------------------------|------------------------------|--|---|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total Age ⁺ 18-24 25-34 35-44 45-64 65+ Race | 26,898 | 91.0 | (86.2-94.3) | 245,502 | 88.7 | (86.2-90.8) |
| White, NH Black, NH Hispanic Other, NH Sex male | 23,029 14,180 | 90.2 # # # 90.1 | (85.0-93.8) (82.1-94.7) | 203,392 19,259 119,966 | 88.5 # 91.4 [^] 86.4 | (85.9-90.7) (78.8-96.8) (82.8-89.3) |
| female Age-adjusted ^{+,&} | 12,719 | 92.1 | (85.4-95.9) | 125,536 | 91.0 | (87.3-93.7) |

& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults aged 18-64 years who saw a doctor for a routine checkup within the last year



| | Tioga County | | | Southern Tier | | |
|--|---------------------|-------------------------------------|----------------------------|-----------------------------|--|---|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total Age ⁺ 18-24 25-34 35-44 45-64 65+ | 21,603 | 72.3 | (63.8-79.5) | 182,948 | 64.9 | (60.8-68.8) |
| Race White, NH Black, NH Hispanic Other, NH Sex male | 17,771 | 68.7 # # 70.9 [^] | (60.2-76.1) (57.1-81.7) | 143,786 13,771 78,883 | 62.5 # 65.4 [^] 56.8 | (58.3-66.5) (47.0-80.2) (51.1-62.4) |
| female Age-adjusted ^{+,&} | 10,470 | 73.9 | (63.8-82.0) | 104,066 | 72.8 | (67.0-77.9) |

& Percent is age-adjusted to the 2000 United States Population.


+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Tioga

Improve Health Status and Reduce Health Disparities

Percentage of adults experiencing housing insecurity in the past 12 months*



Tioga County

Southern Tier

| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| r | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 12,275 | 33.4 | (26.4-41.3) | 110,120 | 33.7 | (30.4-37.2) |
| Age | | | | | | |
| 18-24 | | # | | 13,414 | 30.9^ | (19.6-45.0) |
| 25-34 | | # | | 20,168 | 36.6^ | (27.2-47.0) |
| 35-44 | | # | | 21,029 | 46.7 | (37.1-56.5) |
| 45-64 | 5,773 | 40.4^ | (31.1-50.5) | 42,480 | 38.1 | (33.2-43.2) |
| 65+ | 2,180 | 27.4 | (19.4-37.1) | 12,309 | 18.1 | (14.2-22.8) |
| Race | | | | | | |
| White, NH | 12,223 | 37.1 | (30.2-44.5) | 94,835 | 34.9 | (31.5-38.3) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 5,731 | 24.3 | (13.0-40.7) |
| Sex | | | | | | |
| male | 5,886 | 32.0^ | (21.0-45.4) | 47,371 | 30.5 | (26.3-35.0) |
| female | 6,388 | 34.9 | (27.2-43.4) | 62,749 | 36.6 | (31.6-42.0) |
| Age-adjusted ^{&} | | 33.5 | (25.9-42.1) | | 35.4 | (31.8-39.1) |


1

* Worried or stressed about having enough money to pay for rent or mortgage.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults experiencing food insecurity in the past 12 months*



| Tioga | County | |
|-------|--------|--|
| 11064 | county | |

Southern Tier

| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| _ | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 8,501 | 22.5 | (16.5-29.8) | 82,315 | 23.7 | (20.6-27.1) |
| Age | | | | | | |
| 18-24 | | # | | 12,360 | 23.2 | (14.6-34.8) |
| 25-34 | | # | | 13,426 | 23.0 | (16.2-31.4) |
| 35-44 | | # | | 16,225 | 35.4^ | (26.0-46.1) |
| 45-64 | 3,037 | 20.7 | (13.8-29.9) | 30,573 | 26.5 | (21.5-32.2) |
| 65+ | 1,635 | 20.2 | (13.8-28.5) | 9,530 | 13.5 | (9.7-18.6) |
| Race | | | | | | |
| White, NH | 8,412 | 24.9 | (18.7-32.3) | 66,830 | 23.0 | (20.0-26.3) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 7,161 | 29.6^ | (14.1-51.9) |
| Sex | | | | | | |
| male | 3,705 | 19.6 | (11.2-32.2) | 34,087 | 20.1 | (16.1-24.9) |
| female | 4,797 | 25.3 | (18.2-34.0) | 48,228 | 27.1 | (22.6-32.1) |
| Age-adjusted ^{&} | | 24.2 | (17.3-32.7) | | 25.2 | (21.9-28.8) |


Í

* Worried or stressed about having enough money to buy nutritious meals.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults with poor self-reported health




| | Tioga County | | | Southern Tier | | |
|-------------------------------|---------------------|---------|------------|---------------------|---------|------------|
| - | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 2,298 | 5.9 | (3.7-9.4) | 18,143 | 5.1 | (3.7-6.9) |
| Age | | | | | | |
| 18-24 | | # | | | # | |
| 25-34 | | # | | | # | |
| 35-44 | | # | | | # | |
| 45-64 | 1,323 | 8.8 | (4.5-16.5) | 7,509 | 6.4 | (4.5-9.0) |
| 65+ | 684 | 8.2 | (4.1-15.5) | 5,843 | 8.1 | (5.8-11.2) |
| Race | | | | | | |
| White, NH | 1,637 | 4.7 | (2.9-7.7) | 14,079 | 4.7 | (3.2-6.8) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 2,274 | 8.6 | (4.0-17.6) |
| Sex | | | | | | |
| male | 833 | 4.4 | (2.1-8.7) | 10,260 | 5.8 | (3.6-9.4) |
| female | 1,465 | 7.4 | (4.0-13.6) | 7,882 | 4.3 | (3.1-6.1) |
| Age-adjusted ^{&} | | 5.3 | (3.2-8.6) | | 4.8 | (3.4-6.6) |

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults who report 14 or more days of poor physical health*



| Tioga Co | unty |
|----------|------|
|----------|------|

Southern Tier

| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 5,698 | 15.2 | (11.0-20.6) | 43,428 | 12.4 | (10.4-14.8) |
| Age | | | | | | |
| 18-24 | | # | | | # | |
| 25-34 | | # | | 5,741 | 9.6^ | (4.3-20.1) |
| 35-44 | | # | | 5,563 | 12.0 | (6.8-20.5) |
| 45-64 | 2,771 | 19.4 | (12.2-29.3) | 17,383 | 15.3 | (12.1-19.0) |
| 65+ | 1,783 | 21.8 | (15.0-30.6) | 12,114 | 17.3 | (13.5-21.9) |
| Race | | | | | | |
| White, NH | 5,075 | 15.2 | (11.1-20.4) | 34,674 | 11.9 | (9.9-14.3) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 4,940 | 19.1^ | (9.7-34.2) |
| Sex | | | | | | |
| male | 2,516 | 13.4 | (7.8-22.1) | 22,786 | 13.2 | (10.1-17.2) |
| female | 3,183 | 17.1 | (11.8-24.1) | 20,642 | 11.6 | (9.2-14.5) |
| Age-adjusted ^{&} | | 14.6 | (10.4-20.3) | | 12.3 | (10.1-14.9) |


Ĩ

* Out of the past 30 days.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults who are obese*



| Tioga | County |
|-------|--------|
| 11050 | county |

Southern Tier

| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 11,610 | 32.6 | (25.7-40.4) | 95,241 | 28.5 | (25.4-31.9) |
| Age | | | | | | |
| 18-24 | | # | | 7,926 | 14.8^ | (8.1-25.3) |
| 25-34 | | # | | 16,121 | 30.6^ | (21.4-41.6) |
| 35-44 | | # | | 15,806 | 36.4^ | (27.1-46.9) |
| 45-64 | 4,702 | 34.0 | (25.3-43.9) | 37,057 | 32.8 | (28.3-37.6) |
| 65+ | 2,536 | 32.9 | (24.9-41.9) | 18,039 | 26.4 | (21.6-31.9) |
| Race | | | | | | |
| White, NH | 9,656 | 30.6 | (24.6-37.3) | 82,917 | 29.7 | (26.5-33.1) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 4,994 | 19.2^ | (9.3-35.7) |
| Sex | | | | | | |
| male | 5,385 | 28.6 | (18.8-40.9) | 49,087 | 28.2 | (23.8-33.0) |
| female | 6,226 | 37.1 | (28.9-46.1) | 46,154 | 28.9 | (24.5-33.7) |
| Age-adjusted ^{&} | | 36.5 | (29.1-44.7) | | 29.8 | (26.4-33.4) |


Ĩ

* Obesity defined as a Body Mass Index (BMI) >=30. BMI is calculated based on weight and height.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults with an annual household income less than \$25,000 who are obese*



| Tioga County | |
|--------------|--|
|--------------|--|

Southern Tier

| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 4,313 | 36.4^ | (25.3-49.1) | 29,061 | 32.0 | (26.9-37.5) |
| Age | | | | | | |
| 18-24 | | # | | | # | |
| 25-34 | | # | | 3,472 | 20.1^ | (11.0-33.8) |
| 35-44 | | # | | 4,258 | 44.8^ | (26.9-64.2) |
| 45-64 | 1,910 | 37.2 | (22.1-55.4) | 12,769 | 40.5 | (31.9-49.7) |
| 65+ | 1,007 | 34.3^ | (21.6-49.7) | 6,552 | 28.8 | (22.1-36.6) |
| Race | | | | | | |
| White, NH | 3,471 | 32.1^ | (21.3-45.3) | 24,568 | 32.3 | (27.0-38.0) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | | # | |
| Sex | | | | | | |
| male | 1,070 | 20.0^ | (9.8-36.6) | 11,743 | 25.7 | (19.2-33.5) |
| female | 3,243 | 49.8^ | (35.5-64.0) | 17,318 | 38.3 | (31.1-46.1) |
| Age-adjusted ^{&} | | 35.3^ | (25.7-46.4) | | 33.2 | (27.4-39.4) |


Í

* Obesity defined as a Body Mass Index (BMI) >=30. BMI is calculated based on weight and height.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults living with a disability who are obese*



| Tioga County | |
|--------------|--|
|--------------|--|

Southern Tier

| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 3,912 | 41.9^ | (31.5-53.1) | 32,685 | 38.8 | (32.9-45.1) |
| Age | | | | | | |
| 18-24 | | # | | | # | |
| 25-34 | | # | | | # | |
| 35-44 | | # | | | # | |
| 45-64 | 1,814 | 40.8^ | (25.1-58.8) | 15,081 | 43.9 | (36.0-52.2) |
| 65+ | 1,374 | 37.8^ | (26.2-51.1) | 7,075 | 27.6 | (21.3-35.0) |
| Race | | | | | | |
| White, NH | 3,070 | 36.7^ | (27.2-47.3) | 26,719 | 37.0 | (30.9-43.7) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | | # | |
| Sex | | | | | | |
| male | 1,638 | 38.2^ | (23.5-55.5) | 15,797 | 36.6^ | (27.5-46.8) |
| female | 2,274 | 45.1^ | (31.9-59.1) | 16,888 | 41.2 | (34.0-48.7) |
| Age-adjusted ^{&} | | 47.9^ | (32.2-64.0) | | 41.7 | (34.0-49.9) |


Ĩ

* Obesity defined as a Body Mass Index (BMI) >=30. BMI is calculated based on weight and height. Limited activities (physical, mental, or emotional problems); need of special equipment (e.g. cane, wheel chair).

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults overweight or obese*



Tioga County

Southern Tier

| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 23,661 | 66.4 | (58.1-73.8) | 199,985 | 59.9 | (56.1-63.7) |
| Age | | | | | | |
| 18-24 | | # | | 19,724 | 36.7 | (25.1-50.2) |
| 25-34 | | # | | 26,653 | 50.6^ | (40.5-60.7) |
| 35-44 | | # | | 30,381 | 70.0 | (60.6-78.1) |
| 45-64 | 9,728 | 70.3 | (60.4-78.6) | 76,054 | 67.3 | (62.0-72.3) |
| 65+ | 5,058 | 65.5 | (56.4-73.7) | 45,261 | 66.4 | (61.6-70.8) |
| Race | | | | | | |
| White, NH | 19,830 | 62.9 | (54.9-70.1) | 174,723 | 62.5 | (59.1-65.9) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 11,547 | 44.5^ | (28.3-61.9) |
| Sex | | | | | | |
| male | 12,811 | 67.9^ | (54.1-79.2) | 112,653 | 64.7 | (59.7-69.4) |
| female | 10,850 | 64.7 | (56.3-72.3) | 87,331 | 54.7 | (49.0-60.3) |
| Age-adjusted ^{&} | | 68.1 | (59.0-76.0) | | 60.8 | (57.1-64.3) |


Í

* Defined as a Body Mass Index (BMI) >=25. BMI is calculated based on weight and height.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults who participated in leisure time physical activity in the past 30 days*



Tioga County

Southern Tier

| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 26,429 | 69.5 | (62.1-76.0) | 259,735 | 74.2 | (70.9-77.4) |
| Age | | | | | | |
| 18-24 | | # | | 41,971 | 78.8^ | (66.5-87.4) |
| 25-34 | | # | | 42,562 | 71.7^ | (58.8-81.7) |
| 35-44 | | # | | 36,839 | 80.3 | (72.6-86.3) |
| 45-64 | 10,658 | 72.9 | (63.8-80.5) | 85,632 | 73.9 | (69.3-78.1) |
| 65+ | 4,637 | 55.8 | (46.9-64.5) | 48,976 | 68.8 | (64.2-73.1) |
| Race | | | | | | |
| White, NH | 22,685 | 66.8 | (59.7-73.1) | 213,124 | 72.9 | (69.7-75.9) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 22,139 | 88.7^ | (78.0-94.6) |
| Sex | | | | | | |
| male | 15,049 | 79.3^ | (67.5-87.6) | 133,344 | 78.2 | (74.0-81.9) |
| female | 11,380 | 59.7 | (51.0-67.8) | 126,391 | 70.5 | (65.2-75.2) |
| Age-adjusted ^{&} | | 69.9 | (61.8-77.0) | | 74.7 | (71.2-77.8) |


Í

* Any physical activities or exercises such as running or walking

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults who consume one or more sugary drinks daily*



| Tinga | County | |
|-------|--------|--|
| Tiuga | County | |

Southern Tier

| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 9,429 | 25.1 | (18.8-32.7) | 88,098 | 25.6 | (22.3-29.3) |
| Age | | | | | | |
| 18-24 | | # | | 20,979 | 40.1^ | (27.3-54.3) |
| 25-34 | | # | | 23,280 | 40.2 | (29.5-52.0) |
| 35-44 | | # | | 11,061 | 24.2 | (18.0-31.8) |
| 45-64 | 3,707 | 25.6 | (18.2-34.7) | 23,746 | 20.8 | (17.0-25.1) |
| 65+ | 1,649 | 20.0 | (13.5-28.5) | 8,799 | 12.6 | (10.0-15.8) |
| Race | | | | | | |
| White, NH | 8,128 | 24.3 | (18.7-31.0) | 71,236 | 24.8 | (21.6-28.3) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 4,913 | 20.3^ | (11.0-34.3) |
| Sex | | | | | | |
| male | 5,967 | 32.1 | (21.0-45.6) | 44,773 | 26.7 | (22.6-31.2) |
| female | 3,461 | 18.3 | (12.3-26.3) | 43,325 | 24.6 | (19.6-30.4) |
| Age-adjusted ^{&} | | 26.6 | (19.7-34.9) | | 26.2 | (22.9-29.7) |


Ĩ

* Sugary drinks include regular soda, and sugar-sweetened drinks.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults who consume fast-food three or more times per week




| | Tioga County | | | Southern Tier | | |
|-------------------------------|---------------------|---------|------------|---------------------|---------|------------|
| - | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 1,481 | 4.2 | (1.5-10.9) | 20,347 | 6.0 | (4.5-8.0) |
| Age | | | | | | |
| 18-24 | | # | | | # | |
| 25-34 | | # | | 6,344 | 11.0 | (6.3-18.6) |
| 35-44 | | # | | 2,414 | 5.3 | (2.5-10.8) |
| 45-64 | | # | | 5,371 | 4.7 | (2.9-7.5) |
| 65+ | | # | | 1,671 | 2.4 | (1.4-4.1) |
| Race | | | | | | |
| White, NH | 1,481 | 4.5 | (1.6-11.5) | 15,486 | 5.4 | (3.9-7.5) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | | # | |
| Sex | | | | | | |
| male | | # | | 12,975 | 7.9 | (5.5-11.2) |
| female | | # | | 7,372 | 4.2 | (2.6-6.9) |
| Age-adjusted ^{&} | | 4.9 | (1.8-12.7) | | 6.1 | (4.5-8.2) |

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults with physician diagnosed diabetes*



Southern Tier

| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
|-------------------------------|---------------------|---------|-------------|---------------------|---------|-------------|
| Total | 4,220 | 10.8 | (7.6-15.1) | 36,788 | 10.2 | (8.1-12.7) |
| Age | .)==0 | 2010 | (710 2012) | 00,700 | 2012 | (012 22.7) |
| 18-24 | | # | | | # | |
| 25-34 | | # | | | # | |
| 35-44 | | # | | | # | |
| 45-64 | 1,749 | 11.6 | (6.7-19.4) | 14,046 | 11.8 | (9.1-15.1) |
| 65+ | 1,641 | 19.4 | (13.5-26.9) | 16,124 | 22.1 | (17.8-27.1) |
| Race | | | | | | |
| White, NH | 3,429 | 9.8 | (7.1-13.4) | 27,473 | 9.1 | (7.8-10.7) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 4,025 | 14.8^ | (6.9-28.9) |
| Sex | | | | | | |
| male | 1,525 | 8.0 | (4.5-13.7) | 16,397 | 9.3 | (7.2-11.9) |
| female | 2,695 | 13.6 | (8.9-20.1) | 20,391 | 11.1 | (7.7-15.6) |
| Age-adjusted ^{&} | | 10.4 | (7.0-15.2) | | 9.1 | (7.1-11.7) |


I

* As diagnosed by a health professional.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults who are current smokers



| Tioga County | |
|--------------|--|
| | |


Southern Tier

| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 8,732 | 22.8 | (16.8-30.0) | 71,044 | 20.1 | (17.4-23.0) |
| Age | | | | | | |
| 18-24 | | # | | 9,973 | 18.3^ | (10.8-29.1) |
| 25-34 | | # | | 15,353 | 26.0 | (18.4-35.4) |
| 35-44 | | # | | 12,078 | 26.2 | (17.7-36.9) |
| 45-64 | 4,731 | 31.9 | (23.6-41.4) | 28,406 | 24.0 | (20.1-28.4) |
| 65+ | 758 | 9.0 | (5.1-15.5) | 5,115 | 7.1 | (5.2-9.7) |
| Race | | | | | | |
| White, NH | 8,323 | 24.3 | (18.2-31.6) | 63,096 | 21.3 | (18.5-24.5) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 3,338 | 13.4^ | (7.1-23.9) |
| Sex | | | | | | |
| male | 4,809 | 25.2 | (15.6-38.0) | 36,695 | 21.1 | (17.4-25.4) |
| female | 3,924 | 20.3 | (14.3-28.2) | 34,348 | 19.1 | (15.4-23.3) |
| Age-adjusted ^{&} | | 22.7 | (16.4-30.5) | | 21.2 | (18.3-24.6) |

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults with annual household income less than \$25,000 who are current smokers



#

35.6

31.7

35.6

(27.7 - 44.3)

(24.1 - 40.3)

(29.4 - 42.4)

15,897

17,185

| | Tioga County | | | Southern Tier | | |
|-----------|---------------------|---------|-------------|---------------------|---------|-------------|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 4,927 | 40.6^ | (28.0-54.5) | 33,082 | 33.4 | (27.8-39.6) |
| Age | | | | | | |
| 18-24 | | # | | | # | |
| 25-34 | | # | | 8,325 | 38.6^ | (22.8-57.1) |
| 35-44 | | # | | 4,330 | 37.6^ | (21.9-56.6) |
| 45-64 | 2,739 | 50.5^ | (33.3-67.5) | 14,058 | 43.6 | (35.1-52.4) |
| 65+ | 411 | 13.1 | (6.1-25.7) | 2,371 | 9.7 | (5.9-15.5) |
| Race | | | | | | |
| White, NH | 4,640 | 41.8^ | (28.6-56.2) | 28,558 | 35.6 | (29.9-41.7) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |

(20.4-48.3)

(28.5-57.3)

& Percent is age-adjusted to the 2000 United States Population.

Other, NH

Sex male

female

Age-adjusted[&]

Rates with <10 in the numerator or <50 in the denominator are suppressed.

2,251


#

#

32.9^

42.2

Percentage of adults aged 18-24 years who are current smokers




| | Tioga County | | | Southern Tier | | |
|--|---------------------|-------------|--------|---------------------|-----------------------------|-------------|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total Age ⁺ 18-24 25-34 35-44 | | # | | 9,973 | 18.3 [^] | (10.8-29.1) |
| 45-64 65+ Race | | | | | | |
| White, NH Black, NH Hispanic Other, NH Sex | | # # # | | 9,651 | 23.3 [^] # # | (14.1-36.0) |
| male female Age-adjusted ^{+,&} | | # # | | 6,557 | 23.4 [^] # | (12.3-40.0) |

& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of women aged 50-74 years receiving breast cancer screening based on recent guidelines^{*}



| | Tioga County | | | Southern Tier | | |
|--|---------------------|----------------------------------|-------------|---------------------|---------------------|-------------|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total Age ⁺ 18-24 25-34 35-44 45-64 65+ | 5,443 | 59.3^ | (47.8-69.9) | 51,498 | 79.0 | (74.1-83.2) |
| Race White, NH Black, NH Hispanic Other, NH Sex male | 5,172 | 61.5 [^] # # # | (49.7-72.1) | 45,195 | 78.4 # # # | (73.2-82.8) |
| female Age-adjusted ^{+,&} | 5,443 | 59.3^ | (47.8-69.9) | 51,498 | 79.0 | (74.1-83.2) |

* Mammogram within past 2 years.

& Percent is age-adjusted to the 2000 United States Population.


+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Tioga

Prevent Chronic Diseases

Percentage of women aged 50-74 years with annual household income of less than \$25,000 receiving breast cancer screening based on recent guidelines^{*}



| | Tioga County | | | Southern Tier | | |
|---------------------------------|---------------------|---------|--------|---------------------|---------|-------------|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | | # | | 12,012 | 69.1^ | (57.8-78.5) |
| Age⁺ | | | | | | |
| 18-24 | | | | | | |
| 25-34 | | | | | | |
| 35-44 | | | | | | |
| 45-64 | | | | | | |
| 65+ | | | | | | |
| Race | | | | | | |
| White, NH | | # | | 10,516 | 68.8^ | (56.7-78.7) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | | # | |
| Sex | | | | | | |
| male | | # | | | # | |
| female | | # | | 12,012 | 69.1^ | (57.8-78.5) |
| Age-adjusted ^{+,&} | | | | | | |


* Mammogram within past 2 years.

& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of women aged 21-65 years receiving cervical cancer screening based on recent guidelines^{*}



| | Tioga County | | | Southern Tier | | |
|--|---------------------|-----------------------------|-------------|---------------------|---------------------|-------------|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total Age ⁺ 18-24 25-34 35-44 | 10,421 | 80.2^ | (70.1-87.5) | 100,500 | 81.9 | (77.2-85.8) |
| 45-64 65+ Race White, NH Black, NH Hispanic Other, NH | 9,402 | 78.5 [^] # # | (67.7-86.4) | 84,356 | 81.9 # # # | (77.5-85.6) |
| Sex male female Age-adjusted ^{+,&} | 10,421 | # 80.2 [^] | (70.1-87.5) | 100,500 | # 81.9 | (77.2-85.8) |

* Pap test within past 3 years.

& Percent is age-adjusted to the 2000 United States Population.


+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Tioga

Prevent Chronic Diseases

Percentage of women aged 21-65 years with annual household income of less than \$25,000 receiving cervical cancer screening based on recent guidelines^{*}



| | Tioga County | | | Southern Tier | | | |
|---------------------------------|---------------------|---------|--------|---------------------|---------|-------------|--|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI | |
| Total | | # | | 29,790 | 79.7 | (71.0-86.4) | |
| Age⁺ | | | | | | | |
| 18-24 | | | | | | | |
| 25-34 | | | | | | | |
| 35-44 | | | | | | | |
| 45-64 | | | | | | | |
| 65+ | | | | | | | |
| Race | | | | | | | |
| White, NH | | # | | 21,880 | 75.4 | (66.0-82.8) | |
| Black, NH | | # | | | # | | |
| Hispanic | | # | | | # | | |
| Other, NH | | # | | | # | | |
| Sex | | | | | | | |
| male | | # | | | # | | |
| female | | # | | 29,790 | 79.7 | (71.0-86.4) | |
| Age-adjusted ^{+,&} | | | | | | | |


* Pap test within past 3 years.

& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines^{*}



| | Tioga County | | | Southern Tier | | |
|---|-------------------------|--|---|----------------------------|--------------------------------|---|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total Age ⁺ 18-24 25-34 35-44 45-64 65+ Race | 8,559 | 53.9 | (45.3-62.2) | 82,061 | 63.4 | (58.5-68.1) |
| White, NH Black, NH Hispanic Other, NH Sex male female | 8,250 4,018 4,540 | 54.6 # # 56.9 [^] 51.5 [^] | (45.8-63.1) (43.4-69.4) (40.4-62.4) | 72,849 38,092 43,969 | 65.4 # # 61.3 65.4 | (61.3-69.3) (53.2-68.7) (59.3-71.0) |
| Age-adjusted ^{+,&} | | | | | | |

* Blood stool test within 1 year, or sigmoidoscopy within 5 years with blood stool test within 3 years, or colonoscopy within 10 years.

& Percent is age-adjusted to the 2000 United States Population.


+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Tioga

Prevent Chronic Diseases

Percentage of adults aged 50-75 years with annual household income less than \$25,000 receiving colorectal cancer screening based on recent guidelines^{*}



| | Tioga County | | | Southern Tier | | | |
|---------------------------------|---------------------|---------|-------------|---------------------|---------|-------------|--|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI | |
| Total | 2,044 | 39.1^ | (25.6-54.6) | 19,541 | 56.0 | (47.4-64.3) | |
| Age⁺ | | | | | | | |
| 18-24 | | | | | | | |
| 25-34 | | | | | | | |
| 35-44 | | | | | | | |
| 45-64 | | | | | | | |
| 65+ | | | | | | | |
| Race | | | | | | | |
| White, NH | 2,044 | 42.9^ | (28.0-59.3) | 16,747 | 55.0 | (46.2-63.4) | |
| Black, NH | | # | | | # | | |
| Hispanic | | # | | | # | | |
| Other, NH | | # | | | # | | |
| Sex | | | | | | | |
| male | | # | | 9,923 | 60.7^ | (49.2-71.2) | |
| female | | # | | 9,618 | 51.9^ | (39.9-63.6) | |
| Age-adjusted ^{+,&} | | | | | | | |


* Blood stool test within 1 year, or sigmoidoscopy within 5 years with blood stool test within 3 years, or colonoscopy within 10 years.

& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults who had a test for high blood sugar or diabetes within the past three years



Southern Tier


| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 19,303 | 56.6 | (47.0-65.8) | 164,536 | 53.6 | (49.7-57.5) |
| Age | | | | | | |
| 18-24 | | # | | 13,960 | 28.0^ | (18.0-40.8) |
| 25-34 | | # | | 25,224 | 46.6^ | (36.7-56.7) |
| 35-44 | | # | | 24,844 | 54.5 | (44.6-64.1) |
| 45-64 | 8,305 | 63.1^ | (52.3-72.8) | 60,606 | 60.3 | (54.6-65.7) |
| 65+ | 4,885 | 74.0 | (64.4-81.8) | 37,837 | 71.0 | (65.6-75.8) |
| Race | | | | | | |
| White, NH | 17,852 | 57.9 | (49.8-65.6) | 142,752 | 55.3 | (51.7-59.0) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 8,459 | 39.2^ | (23.2-57.8) |
| Sex | | | | | | |
| male | 8,264 | 47.3^ | (33.2-61.8) | 78,714 | 51.4 | (46.1-56.7) |
| female | 11,039 | 66.5 | (56.5-75.1) | 85,822 | 55.8 | (49.9-61.6) |
| Age-adjusted ^{&} | | 58.9 | (51.6-65.9) | | 54.2 | (50.4-57.9) |

Í

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults with physician-diagnosed prediabetes




| | Tioga County | | | Southern Tier | | |
|-------------------------------|---------------------|---------|------------|---------------------|---------|------------|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 1,700 | 5.0 | (3.0-8.1) | 16,763 | 5.3 | (3.8-7.3) |
| Age | | | | | | |
| 18-24 | | # | | | # | |
| 25-34 | | # | | | # | |
| 35-44 | | # | | | # | |
| 45-64 | 924 | 7.1 | (3.3-14.5) | 6,771 | 6.6 | (4.8-9.1) |
| 65+ | 531 | 7.9 | (4.2-14.3) | 3,770 | 6.8 | (4.5-10.1) |
| Race | | | | | | |
| White, NH | 1,700 | 5.5 | (3.3-8.9) | 15,289 | 5.7 | (4.0-8.0) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | | # | |
| Sex | | | | | | |
| male | 477 | 2.8 | (1.3-5.8) | 8,956 | 5.7 | (3.3-9.7) |
| female | 1,223 | 7.2 | (3.9-13.0) | 7,807 | 4.8 | (3.4-6.7) |
| Age-adjusted ^{&} | | 4.1 | (2.5-6.7) | | 5.4 | (3.8-7.5) |

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults with physician-diagnosed high blood pressure



| Tioga County | |
|--------------|--|
| | |


Southern Tier

| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 13,231 | 33.9 | (27.3-41.3) | 108,205 | 30.1 | (27.2-33.1) |
| Age | | | | | | |
| 18-24 | | # | | 5,198 | 9.2 | (4.7-17.4) |
| 25-34 | | # | | 8,421 | 13.8 | (8.3-22.1) |
| 35-44 | | # | | 6,819 | 14.3 | (9.4-21.3) |
| 45-64 | 4,425 | 29.2 | (21.9-37.9) | 41,371 | 35.0 | (30.6-39.7) |
| 65+ | 5,308 | 62.5 | (53.6-70.6) | 44,299 | 61.4 | (56.5-66.1) |
| Race | | | | | | |
| White, NH | 12,367 | 35.4 | (29.1-42.3) | 93,946 | 31.4 | (28.6-34.3) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 7,589 | 27.8^ | (15.4-44.8) |
| Sex | | | | | | |
| male | 6,934 | 36.1 | (24.8-49.1) | 57,850 | 33.0 | (28.8-37.5) |
| female | 6,297 | 31.8 | (25.0-39.5) | 50,354 | 27.3 | (23.7-31.3) |
| Age-adjusted ^{&} | | 29.9 | (23.8-36.8) | | 27.8 | (25.3-30.4) |

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults with diagnosed high blood pressure taking high blood pressure medication



| HUYA CUUIILV | Tioga | County | |
|--------------|-------|--------|--|
|--------------|-------|--------|--|

Southern Tier


| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 9,546 | 72.2^ | (56.7-83.7) | 81,095 | 75.3 | (70.3-79.7) |
| Age | | | | | | |
| 18-24 | | # | | | # | |
| 25-34 | | # | | | # | |
| 35-44 | | # | | | # | |
| 45-64 | 3,361 | 76.0^ | (60.0-86.9) | 32,070 | 77.5 | (70.3-83.4) |
| 65+ | 4,803 | 90.5 | (81.6-95.3) | 40,246 | 91.8 | (88.4-94.3) |
| Race | | | | | | |
| White, NH | 8,966 | 72.5^ | (56.2-84.4) | 69,969 | 74.9 | (69.7-79.4) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | | # | |
| Sex | | | | | | |
| male | 4,092 | 59.0^ | (37.7-77.4) | 38,709 | 67.5 | (59.6-74.5) |
| female | 5,454 | 86.6^ | (75.1-93.3) | 42,386 | 84.2 | (78.8-88.4) |
| Age-adjusted ^{&} | | 62.6^ | (49.4-74.2) | | 57.9 | (50.4-65.1) |

Í

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults with current asthma*



Southern Tier


| | noga county | | | | Southernin | |
|-------------------------------|---------------------|---------|------------|---------------------|------------|-------------|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 4,063 | 10.5 | (7.0-15.5) | 39,362 | 11.1 | (8.7-13.9) |
| Age | - | | | - | | . , |
| 18-24 | | # | | 7,986 | 15.4 | (9.2-24.5) |
| 25-34 | | # | | 9,891 | 16.2^ | (8.1-29.8) |
| 35-44 | | # | | 3,554 | 7.5 | (4.4-12.4) |
| 45-64 | 1,628 | 11.0 | (6.1-18.9) | 11,753 | 9.9 | (7.7-12.8) |
| 65+ | 855 | 10.1 | (6.2-16.1) | 6,076 | 8.4 | (4.9-13.8) |
| Race | | | | | | |
| White, NH | 3,529 | 10.2 | (6.7-15.2) | 28,448 | 9.5 | (7.8-11.5) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 3,373 | 12.4 | (4.7-28.7) |
| Sex | | | | | | |
| male | 1,337 | 7.0 | (3.3-14.2) | 14,033 | 7.9 | (5.9-10.6) |
| female | 2,726 | 13.9 | (8.7-21.5) | 25,329 | 14.1 | (10.2-19.1) |
| Age-adjusted ^{&} | | 11.9 | (7.4-18.7) | | 11.0 | (8.7-13.8) |

* Physician diagnosed asthma.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percent of adults with cholesterol checked*



Tioga County

Southern Tier

| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 33,180 | 86.1 | (79.7-90.8) | 280,906 | 81.0 | (76.9-84.5) |
| Age | | | | | | |
| 18-24 | | # | | 25,093 | 50.3^ | (36.0-64.5) |
| 25-34 | | # | | 37,573 | 65.7^ | (53.6-76.1) |
| 35-44 | | # | | 36,981 | 79.8 | (70.1-87.0) |
| 45-64 | 13,640 | 90.2 | (82.7-94.6) | 107,406 | 91.5 | (88.6-93.6) |
| 65+ | 8,247 | 97.3 | (92.8-99.1) | 69,502 | 97.5 | (95.6-98.6) |
| Race | | | | | | |
| White, NH | 29,141 | 84.6 | (77.7-89.6) | 238,055 | 82.8 | (79.7-85.6) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 21,553 | 80.6^ | (64.6-90.4) |
| Sex | | | | | | |
| male | 15,336 | 80.1^ | (68.0-88.4) | 138,828 | 81.3 | (77.1-84.9) |
| female | 17,844 | 92.2 | (86.5-95.6) | 142,078 | 80.7 | (73.5-86.3) |
| Age-adjusted ^{&} | | 84.5 | (76.4-90.1) | | 79.9 | (76.3-83.2) |


Í

* As diagnosed by a health professional.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of adults 18 years and older with elevated cholesterol*



Tioga County

Southern Tier

| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 11,899 | 36.2 | (29.2-43.8) | 107,760 | 38.9 | (35.5-42.5) |
| Age | | | | | | |
| 18-24 | | # | | | # | |
| 25-34 | | # | | 8,162 | 21.9^ | (11.8-36.9) |
| 35-44 | | # | | 9,810 | 26.7 | (18.9-36.3) |
| 45-64 | 5,863 | 43.0 | (33.7-52.8) | 49,298 | 46.7 | (41.4-52.2) |
| 65+ | 4,478 | 55.5 | (46.3-64.4) | 37,715 | 55.7 | (50.5-60.7) |
| Race | | | | | | |
| White, NH | 11,065 | 38.3 | (32.0-45.1) | 89,873 | 38.3 | (35.0-41.7) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 11,862 | 56.1^ | (36.5-74.0) |
| Sex | | | | | | |
| male | 5,064 | 33.2 | (22.4-46.0) | 56,113 | 40.9 | (35.6-46.5) |
| female | 6,835 | 38.8 | (30.9-47.2) | 51,646 | 37.0 | (32.7-41.6) |
| Age-adjusted ^{&} | | 28.4 | (23.3-34.2) | | 33.6 | (30.1-37.2) |

Í


* As diagnosed by a health professional.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Tioga Prevent Chronic Diseases

Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition^{*}



| | т | Tioga Count | ty | Southern Tier | | | |
|-----------|---------------------|-------------|------------|---------------------|---------|------------|--|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI | |
| Total | 2,354 | 10.8 | (6.9-16.6) | 20,040 | 10.6 | (7.3-15.1) | |
| Age | | | | | | | |
| 18-24 | | # | | | # | | |
| 25-34 | | # | | | # | | |
| 35-44 | | # | | | # | | |
| 45-64 | 1,116 | 12.9^ | (6.6-23.6) | 6,755 | 9.8 | (7.3-13.0) | |
| 65+ | | # | | 6,303 | 10.7 | (7.6-14.9) | |
| Race | | | | | | | |
| White, NH | 2,030 | 9.9 | (6.2-15.3) | 14,247 | 9.0 | (7.2-11.2) | |
| Black, NH | | # | | | # | | |
| Hispanic | | # | | | # | | |
| Other, NH | | # | | | # | | |
| Sex | | | | | | | |
| male | 937 | 9.5 | (4.9-17.7) | 6,930 | 7.9 | (5.4-11.5) | |
| female | 1,416 | 11.9 | (6.5-20.8) | 13,110 | 12.9 | (7.6-20.9) | |
| | | | | | | | |

(7.0-20.3)

10.5

(6.4 - 16.7)

* Courses for high blood pressure, arthritis, or diabetes.

Age-adjusted[&]


& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

12.2

Tioga Prevent Chronic Diseases

Percentage of adults with arthritis*



Tioga County

Southern Tier

| | nogu county | | | Southern ner | | |
|-------------------------------|---------------------|---------|-------------|---------------------|---------|-------------|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 14,386 | 37.1 | (30.5-44.2) | 102,167 | 28.3 | (25.6-31.1) |
| Age | | | | | | |
| 18-24 | | # | | | # | |
| 25-34 | | # | | 5,390 | 8.9 | (5.4-14.2) |
| 35-44 | | # | | 7,770 | 16.4 | (11.2-23.3) |
| 45-64 | 6,571 | 43.4 | (34.4-53.0) | 44,194 | 37.1 | (32.6-41.8) |
| 65+ | 5,630 | 67.5 | (58.9-75.0) | 40,660 | 56.0 | (50.9-60.9) |
| Race | | | | | | |
| White, NH | 13,567 | 39.1 | (33.0-45.6) | 91,598 | 30.4 | (27.7-33.3) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 5,619 | 20.6^ | (10.7-35.9) |
| Sex | | | | | | |
| male | 5,879 | 30.8^ | (21.4-42.0) | 38,769 | 21.9 | (18.9-25.3) |
| female | 8,506 | 43.2 | (35.5-51.3) | 63,397 | 34.3 | (30.0-39.0) |
| Age-adjusted ^{&} | | 33.0 | (27.5-39.0) | | 26.3 | (24.1-28.6) |


* Includes rheumatoid arthritis, gout, lupus, or fibromyalgia.

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Promote a Healthy and Safe Environment

Percentage of adults who consider their neighborhood suitable for walking and physical activity



Tioga County

Southern Tier

| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 34,590 | 92.1 | (88.1-94.9) | 315,627 | 91.6 | (89.7-93.2) |
| Age | | | | | | |
| 18-24 | | # | | 47,540 | 89.3 | (79.4-94.7) |
| 25-34 | | # | | 53,444 | 91.4 | (85.0-95.2) |
| 35-44 | | # | | 40,753 | 90.0 | (84.4-93.7) |
| 45-64 | 13,746 | 94.6 | (89.0-97.4) | 106,162 | 93.3 | (90.2-95.4) |
| 65+ | 6,937 | 86.7 | (77.4-92.5) | 63,978 | 91.6 | (88.9-93.7) |
| Race | | | | | | |
| White, NH | 30,712 | 91.7 | (87.3-94.6) | 261,499 | 90.9 | (88.7-92.7) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 23,022 | 95.9 | (89.7-98.4) |
| Sex | | | | | | |
| male | 18,512 | 98.5 | (95.2-99.6) | 156,230 | 92.9 | (89.8-95.1) |
| female | 16,078 | 85.7 | (78.7-90.8) | 159,397 | 90.4 | (87.5-92.7) |
| Age-adjusted ^{&} | | 91.4 | (86.3-94.7) | | 91.4 | (89.4-93.1) |


1

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Promote a Health and Safe Environment

Percentage of adults aged 65+ years with at least one fall in the past 12 months



| | Tioga County | | | Southern Tier | | |
|---------------------------------|---------------------|---------|-------------|---------------------|---------|-------------|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 3,242 | 39.7 | (31.2-48.9) | 23,488 | 33.4 | (28.7-38.5) |
| Age⁺ | | | | | | |
| 18-24 | | | | | | |
| 25-34 | | | | | | |
| 35-44 | | | | | | |
| 45-64 | | | | | | |
| 65+ | | | | | | |
| Race | | | | | | |
| White, NH | 3,191 | 39.3 | (30.8-48.6) | 20,224 | 32.3 | (28.2-36.8) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | | # | |
| Sex | | | | | | |
| male | 1,385 | 43.1^ | (29.0-58.4) | 10,838 | 32.5 | (25.0-40.9) |
| female | 1,857 | 37.5^ | (27.3-48.9) | 12,650 | 34.2 | (28.3-40.6) |
| Age-adjusted ^{+,&} | | | | | | |


& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Promote a Health and Safe Environment

Among adults aged 65+ years with at least one fall in the past 12 months, percent that resulted in injury^{*}



| | Tioga County | | | Southern Tier | | |
|---------------------------------|---------------------|---------|-------------|---------------------|---------|-------------|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 1,365 | 42.5^ | (28.8-57.4) | 8,522 | 36.4 | (28.5-45.0) |
| Age ⁺ | | | | | | |
| 18-24 | | | | | | |
| 25-34 | | | | | | |
| 35-44 | | | | | | |
| 45-64 | | | | | | |
| 65+ | | | | | | |
| Race | | | | | | |
| White, NH | 1,313 | 41.6^ | (27.9-56.7) | 7,465 | 37.0 | (29.7-45.0) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | | # | |
| Sex | | | | | | |
| male | | # | | 3,078 | 28.4^ | (17.1-43.3) |
| female | | # | | 5,444 | 43.2 | (34.0-52.9) |
| Age-adjusted ^{+,&} | | | | | | |


* Falls that limited regular activities or lead to doctor visits.

& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections Percentage of adults aged 47-68 years reporting ever tested for Hepatitis C (HCV)




| | Tioga County | | | Southern Tier | | |
|---|---------------------|--|----------------------------|---------------------|---------------------|----------------------------|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total Age⁺ 18-24 | 3,486 | 23.5 | (17.1-31.3) | 28,278 | 25.7 | (21.9-29.9) |
| 25-34 35-44 45-64 | | | | | | |
| 65+ Race | | | (1= 0, 00, 0) | 25 4 22 | | |
| White, NH Black, NH Hispanic Other, NH | 3,408 | 24.6 # # # | (17.9-32.9) | 25,192 | 27.0 # # # | (23.1-31.2) |
| Sex male female | 1,665 1,821 | 26.2 [^] 21.4 [^] | (16.8-38.5) (13.4-32.6) | 14,697 13,581 | 28.1 23.5 | (22.3-34.8) (18.7-29.0) |
| Age-adjusted ^{+,&} | | | | | | |

& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections Percentage of adults aged 18+ years with flu immunization in the past year^{*}



Tioga County

Southern Tier

| | Est. # of | | | Est. # of | | |
|-------------------------------|-----------|---------|-------------|-----------|---------|-------------|
| г | Adults | Percent | 95% CI | Adults | Percent | 95% CI |
| Total | 20,000 | 52.7 | (44.6-60.7) | 169,235 | 48.7 | (44.9-52.4) |
| Age | | | | | | |
| 18-24 | | # | | 22,678 | 42.8^ | (28.8-58.0) |
| 25-34 | | # | | 21,280 | 36.4^ | (25.5-48.8) |
| 35-44 | | # | | 13,516 | 29.5 | (22.3-37.9) |
| 45-64 | 6,512 | 44.5 | (35.7-53.7) | 58,844 | 50.9 | (45.9-55.9) |
| 65+ | 5,896 | 71.4 | (62.5-78.9) | 49,932 | 70.4 | (65.0-75.3) |
| Race | | | | | | |
| White, NH | 17,880 | 52.8 | (45.7-59.7) | 135,586 | 46.6 | (43.2-50.1) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 12,106 | 50.0^ | (32.3-67.7) |
| Sex | | | | | | |
| male | 9,990 | 52.8^ | (39.0-66.2) | 76,549 | 45.2 | (40.3-50.2) |
| female | 10,010 | 52.6 | (44.1-61.0) | 92,686 | 52.0 | (46.5-57.4) |
| Age-adjusted ^{&} | | 48.5 | (40.8-56.2) | | 45.8 | (42.1-49.7) |


1

* Includes flu shot or flu spray

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections Percentage of adults aged 65+ years with flu immunization in the past year^{*}



| | Tioga County | | | Southern Tier | | |
|---------------------------------|---------------------|---------|-------------|---------------------|---------|-------------|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total | 5,896 | 71.4 | (62.5-78.9) | 49,932 | 70.4 | (65.0-75.3) |
| Age⁺ | | | | | | |
| 18-24 | | | | | | |
| 25-34 | | | | | | |
| 35-44 | | | | | | |
| 45-64 | | | | | | |
| 65+ | | | | | | |
| Race | | | | | | |
| White, NH | 5,845 | 71.2 | (62.3-78.7) | 45,116 | 71.5 | (66.6-75.9) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | | # | |
| Sex | | | | | | |
| male | 2,714 | 83.6^ | (70.8-91.5) | 22,538 | 66.9 | (58.4-74.4) |
| female | 3,182 | 63.5^ | (51.7-73.8) | 27,394 | 73.6 | (67.0-79.3) |
| Age-adjusted ^{+,&} | | | | | | |

* Includes flu shot or flu spray


& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Prevent HIV/STDs, Vaccine Preventable Diseases and Healthcare-Associated Infections

Percentage of adults aged 65+ years with pneumococcal immunization




| | т | ioga Count | ty | Southern Tier | | |
|---|-------------------------|--|---|----------------------------|--------------------------------|---|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total Age ⁺ 18-24 25-34 35-44 45-64 65+ Race | 5,203 | 64.2 | (54.8-72.6) | 46,831 | 68.2 | (62.5-73.3) |
| White, NH Black, NH Hispanic Other, NH Sex male female | 5,151 2,355 2,847 | 64.0 # # 74.6 [^] 57.5 [^] | (54.5-72.4) (60.3-85.1) (45.7-68.6) | 42,016 20,119 26,713 | 68.9 # # 62.6 73.1 | (63.7-73.6) (54.0-70.5) (66.0-79.1) |
| Age-adjusted ^{+,&} | | | | | | |

& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of women aged 18-44 years who had dental visit within the last year




| | Tioga County | | | Southern Tier | | | |
|---------------------------------|---------------------|---------|--------|---------------------|---------|-------------|--|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI | |
| Total | | # | | 52,014 | 74.0 | (65.7-80.9) | |
| Age⁺ | | | | | | | |
| 18-24 | | | | | | | |
| 25-34 | | | | | | | |
| 35-44 | | | | | | | |
| 45-64 | | | | | | | |
| 65+ | | | | | | | |
| Race | | | | | | | |
| White, NH | | # | | 41,836 | 72.1 | (63.2-79.6) | |
| Black, NH | | # | | | # | | |
| Hispanic | | # | | | # | | |
| Other, NH | | # | | | # | | |
| Sex | | | | | | | |
| male | | # | | | # | | |
| female | | # | | 52,014 | 74.0 | (65.7-80.9) | |
| Age-adjusted ^{+,&} | | | | | | | |

& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Women aged 18-44 years who saw a doctor for a routine checkup within the last year




| | Tioga County | | | Southern Tier | | | |
|---------------------------------|---------------------|---------|--------|---------------------|---------|-------------|--|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI | |
| Total | | # | | 57,821 | 69.8 | (60.6-77.7) | |
| Age⁺ | | | | | | | |
| 18-24 | | | | | | | |
| 25-34 | | | | | | | |
| 35-44 | | | | | | | |
| 45-64 | | | | | | | |
| 65+ | | | | | | | |
| Race | | | | | | | |
| White, NH | | # | | 42,375 | 69.2 | (59.7-77.2) | |
| Black, NH | | # | | | # | | |
| Hispanic | | # | | | # | | |
| Other, NH | | # | | | # | | |
| Sex | | | | | | | |
| male | | # | | | # | | |
| female | | # | | 57,821 | 69.8 | (60.6-77.7) | |
| Age-adjusted ^{+,&} | | | | | | | |

& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of women aged 18-64 years with health care coverage




| | Tioga County | | | Southern Tier | | |
|--|---------------------|---------------------|-------------|---------------------|---------------------|-------------|
| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total Age ⁺ 18-24 25-34 35-44 45-64 65+ | 12,719 | 92.1 | (85.4-95.9) | 125,536 | 91.0 | (87.3-93.7) |
| Race White, NH Black, NH Hispanic Other, NH Sex | 11,854 | 92.7 # # # | (85.5-96.5) | 103,783 | 90.9 # # # | (87.0-93.7) |
| male female Age-adjusted ^{+,&} | 12,719 | # 92.1 | (85.4-95.9) | 125,536 | # 91.0 | (87.3-93.7) |

& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Percentage of women aged 18-44 years whose health care provider discussed planning a healthy pregnancy



| | Tioga County | | | Southern Tier | | |
|--|---------------------|-------------|--------|---------------------|---------------------|-------------|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
| Total Age ⁺ 18-24 | | # | | 28,913 | 38.5^ | (29.1-48.7) |
| 25-34 35-44 45-64 65+ | | | | | | |
| Race White, NH Black, NH Hispanic Other, NH Sex | | # # # | | 26,118 | 45.0 # # # | (35.5-55.0) |
| male female Age-adjusted ^{+,&} | | # # | | 28,913 | # 38.5^ | (29.1-48.7) |


& Percent is age-adjusted to the 2000 United States Population.

+ Age-adjusted and age specific percent not generated for health indicators with age range limits.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Tioga Promote Mental Health and Prevention Substance Abuse

Percentage of adults with poor mental health for 14 or more days in the last month



| Tioga | County |
|-------|--------|
| Tiuga | County |

Southern Tier

| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
|-------------------------------|---------------------|---------|-------------|---------------------|---------|------------|
| Total | 4,370 | 11.5 | (8.0-16.4) | 36,482 | 10.3 | (8.6-12.4) |
| Age | | | . , | | | . , |
| 18-24 | | # | | 4,996 | 8.9 | (5.0-15.3) |
| 25-34 | | # | | 8,024 | 13.5 | (7.8-22.5) |
| 35-44 | | # | | 5,170 | 11.0 | (7.3-16.3) |
| 45-64 | 2,557 | 17.8 | (11.2-27.0) | 13,514 | 11.7 | (9.1-14.9) |
| 65+ | 559 | 6.9 | (3.4-13.3) | 4,696 | 6.6 | (4.5-9.7) |
| Race | | | | | | |
| White, NH | 3,774 | 11.1 | (7.6-16.1) | 30,947 | 10.5 | (8.6-12.8) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 2,326 | 8.8^ | (3.8-19.1) |
| Sex | | | | | | |
| male | 1,420 | 7.5 | (3.9-14.2) | 16,352 | 9.5 | (6.9-12.9) |
| female | 2,950 | 15.5 | (10.1-23.0) | 20,130 | 11.2 | (8.9-13.8) |
| Age-adjusted ^{&} | | 11.6 | (7.9-16.8) | | 10.6 | (8.8-12.8) |


I

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Promote Mental Health and Prevention Substance Abuse

Percentage of adult binge drinking during the past month*



Tioga County

Southern Tier

| _ | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI |
|-------------------------------|---------------------|---------|-------------|---------------------|---------|-------------|
| Total | 7,789 | 21.7 | (15.3-29.7) | 61,605 | 18.1 | (15.2-21.4) |
| Age | | | | | | |
| 18-24 | | # | | 15,284 | 30.1^ | (19.4-43.6) |
| 25-34 | | # | | 16,238 | 28.9^ | (18.7-41.7) |
| 35-44 | | # | | 10,075 | 22.3 | (15.8-30.6) |
| 45-64 | 2,826 | 19.5 | (12.5-29.1) | 16,863 | 14.8 | (12.0-18.2) |
| 65+ | 514 | 6.3 | (2.9-13.2) | 3,145 | 4.5 | (3.1-6.4) |
| Race | | | | | | |
| White, NH | 7,666 | 22.7 | (16.0-31.1) | 52,768 | 18.2 | (15.5-21.3) |
| Black, NH | | # | | | # | |
| Hispanic | | # | | | # | |
| Other, NH | | # | | 3,495 | 15.9^ | (7.6-30.2) |
| Sex | | | | | | |
| male | 5,244 | 30.7^ | (19.8-44.3) | 37,622 | 23.0 | (19.2-27.4) |
| female | 2,545 | 13.5 | (8.0-21.8) | 23,983 | 13.5 | (9.5-19.0) |
| Age-adjusted ^{&} | | 24.0 | (17.6-31.7) | | 19.2 | (16.2-22.7) |


* Men who have five or more drinks , women having four or more drinks on one occasion

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.

Tioga Promote Mental Health and Prevention Substance Abuse

Percentage of cigarette smoking among adults who report poor mental health



| | Tioga County | | | Southern Tier | | | |
|-------------------------------|---------------------|---------|--------|---------------------|---------|-------------|--|
| | Est. # of Adults | Percent | 95% CI | Est. # of Adults | Percent | 95% CI | |
| Total | | # | | 14,908 | 42.2 | (33.1-51.8) | |
| Age | | | | | | | |
| 18-24 | | # | | | # | | |
| 25-34 | | # | | | # | | |
| 35-44 | | # | | | # | | |
| 45-64 | | # | | 5,646 | 42.7^ | (30.4-55.9) | |
| 65+ | | # | | 718 | 15.5^ | (7.1-30.8) | |
| Race | | | | | | | |
| White, NH | | # | | 12,617 | 42.3^ | (32.5-52.7) | |
| Black, NH | | # | | | # | | |
| Hispanic | | # | | | # | | |
| Other, NH | | # | | | # | | |
| Sex | | | | | | | |
| male | | # | | 5,524 | 34.7 | (21.2-51.2) | |
| female | | # | | 9,384 | 48.2^ | (37.5-59.2) | |
| Age-adjusted ^{&} | | # | | | 40.4 | (31.9-49.6) | |

& Percent is age-adjusted to the 2000 United States Population.

Rates with <10 in the numerator or <50 in the denominator are suppressed.