

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
All Invasive Malignant Tumors**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	66,297	397.8	394.7-400.9	32,169	469.1	463.7-474.6	34,128	360.1	356.2-364.0	36,530	221.8	219.5-224.2	18,967	283.7	279.4-288.0	17,563	184.0	181.2-186.8
1977	66,426	397.1	394.0-400.2	32,353	471.2	465.8-476.7	34,073	357.5	353.7-361.4	36,701	221.4	219.1-223.8	19,118	284.7	280.4-289.0	17,583	182.6	179.9-185.4
1978	67,165	400.8	397.7-403.9	32,752	477.2	471.8-482.7	34,413	359.9	356.0-363.8	36,470	218.7	216.4-221.0	18,875	281.1	276.9-285.3	17,595	181.2	178.4-183.9
1979	68,544	406.8	403.7-409.9	33,383	485.0	479.6-490.5	35,161	365.3	361.4-369.2	36,511	218.3	216.1-220.6	18,815	279.8	275.6-284.0	17,696	181.3	178.6-184.1
1980	70,887	417.6	414.4-420.7	35,127	506.4	500.9-512.0	35,760	369.0	365.1-372.9	37,475	222.0	219.7-224.3	19,366	287.0	282.8-291.3	18,109	183.3	180.6-186.0
1981	70,846	416.0	412.8-419.1	34,720	500.9	495.4-506.5	36,126	370.6	366.7-374.5	37,169	219.2	216.9-221.5	19,084	282.0	277.8-286.2	18,085	182.1	179.4-184.8
1982	72,146	420.6	417.5-423.7	35,175	504.5	499.0-510.1	36,971	376.5	372.6-380.4	37,109	217.6	215.3-219.8	19,066	281.3	277.1-285.5	18,043	180.3	177.6-183.0
1983	73,570	425.8	422.7-428.9	35,890	510.0	504.5-515.5	37,680	381.7	377.7-385.6	37,487	217.9	215.7-220.2	19,199	281.6	277.5-285.8	18,288	180.7	178.1-183.4
1984	74,608	429.5	426.4-432.7	36,203	510.7	505.2-516.2	38,405	386.8	382.8-390.7	37,797	218.4	216.2-220.7	19,201	279.7	275.6-283.9	18,596	182.8	180.1-185.5
1985	76,570	438.8	435.7-442.0	37,455	524.3	518.8-529.9	39,115	392.5	388.5-396.4	37,977	218.3	216.1-220.5	19,374	280.2	276.1-284.4	18,603	182.1	179.5-184.8
1986	76,739	437.9	434.8-441.1	37,491	524.2	518.7-529.8	39,248	392.1	388.2-396.1	38,311	219.3	217.1-221.5	19,437	280.3	276.2-284.4	18,874	183.7	181.1-186.4
1987	77,199	437.5	434.4-440.7	37,421	519.4	514.0-524.9	39,778	394.0	390.1-398.0	38,189	217.4	215.2-219.6	19,400	279.0	274.9-283.1	18,789	181.3	178.7-184.0
1988	76,968	434.7	431.6-437.8	36,994	512.7	507.3-518.2	39,974	394.8	390.9-398.8	38,597	218.8	216.6-221.0	19,615	281.5	277.4-285.7	18,982	182.4	179.8-185.0
1989	77,635	437.0	433.9-440.1	37,905	521.0	515.5-526.4	39,730	392.0	388.0-395.9	37,829	213.9	211.7-216.1	19,094	272.5	268.5-276.6	18,735	179.7	177.1-182.4
1990	81,426	455.8	452.6-458.9	39,812	542.2	536.7-547.8	41,614	408.9	404.9-412.9	38,109	214.4	212.2-216.6	19,088	271.7	267.8-275.8	19,021	181.3	178.7-183.9
1991	84,129	468.5	465.4-471.8	41,665	563.9	558.4-569.5	42,464	415.9	411.8-419.9	38,348	214.5	212.4-216.7	19,267	272.4	268.4-276.4	19,081	181.0	178.4-183.6
1992	86,866	479.6	476.4-482.8	44,210	591.6	585.9-597.3	42,656	415.1	411.1-419.1	38,499	213.8	211.6-215.9	19,080	268.7	264.8-272.6	19,419	182.9	180.4-185.6
1993	86,321	473.2	470.0-476.4	43,691	576.6	571.1-582.2	42,630	411.6	407.7-415.6	38,243	210.7	208.6-212.8	19,070	265.8	261.9-269.7	19,173	179.2	176.6-181.8
1994	88,997	485.8	482.6-489.0	44,450	582.2	576.7-587.8	44,547	428.1	424.1-432.1	39,129	214.2	212.1-216.3	19,691	272.2	268.3-276.2	19,438	180.3	177.8-182.9
1995	87,522	475.1	472.0-478.3	43,200	560.3	554.9-565.7	44,322	424.1	420.1-428.1	38,654	210.4	208.3-212.5	19,130	262.7	258.9-266.6	19,524	180.1	177.6-182.7
1996	90,009	486.4	483.2-489.6	44,689	575.8	570.4-581.3	45,320	432.1	428.1-436.1	38,112	206.2	204.1-208.3	18,844	255.7	252.0-259.5	19,268	177.0	174.5-179.6
1997	91,384	490.1	486.9-493.3	45,338	579.0	573.6-584.5	46,046	434.9	430.9-439.0	37,289	200.0	198.0-202.1	18,519	248.8	245.2-252.5	18,770	170.9	168.4-173.4
1998	93,644	497.0	493.8-500.2	46,266	580.8	575.5-586.2	47,378	444.8	440.8-448.9	36,821	195.2	193.3-197.2	18,154	238.2	234.7-241.8	18,667	168.5	166.0-170.9
1999	94,557	495.0	491.8-498.1	46,991	582.8	577.4-588.1	47,566	439.8	435.8-443.8	37,335	195.0	193.0-196.9	18,411	239.3	235.8-242.9	18,924	168.3	165.9-170.7
2000	97,320	502.5	499.4-505.7	48,949	599.2	593.9-604.6	48,371	441.1	437.2-445.1	37,222	191.4	189.4-193.3	18,305	234.9	231.5-238.4	18,917	165.2	162.8-167.6
2001	96,709	494.3	491.2-497.4	48,668	587.2	581.9-592.5	48,041	434.8	430.8-438.7	36,804	187.3	185.3-189.2	18,113	229.4	226.0-232.8	18,691	162.0	159.6-164.3
2002	97,860	494.2	491.1-497.3	49,858	592.0	586.8-597.3	48,002	430.2	426.3-434.1	36,488	183.4	181.5-185.3	18,004	224.4	221.0-227.7	18,484	158.6	156.3-160.9
2003	97,840	489.0	485.9-492.1	49,165	576.4	571.3-581.6	48,675	432.5	428.6-436.4	36,051	179.2	177.3-181.0	17,559	215.3	212.1-218.6	18,492	157.1	154.8-159.4
2004	98,643	488.5	485.4-491.5	49,534	572.4	567.3-577.5	49,109	433.9	430.1-437.8	35,826	176.3	174.5-178.2	17,634	213.0	209.8-216.2	18,192	153.3	151.1-155.6
2005	100,018	491.5	488.4-494.6	50,358	574.6	569.5-579.7	49,660	436.3	432.5-440.2	35,272	172.1	170.4-174.0	17,322	206.9	203.8-210.0	17,950	150.4	148.2-152.7
2006	102,891	502.6	499.5-505.7	52,424	592.2	587.1-597.4	50,467	442.3	438.4-446.2	35,041	169.9	168.1-171.7	17,174	202.8	199.8-205.9	17,867	149.1	146.9-151.3
2007	104,541	504.1	501.1-507.2	53,629	594.3	589.2-599.4	50,912	443.1	439.2-447.0	35,192	168.2	166.5-170.0	17,453	202.1	199.1-205.2	17,739	146.7	144.6-149.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
All Invasive Malignant Tumors**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	106,285	505.9	502.9-509.0	53,868	586.9	581.9-591.9	52,417	451.8	447.9-455.8	35,099	165.4	163.7-167.2	17,490	199.4	196.4-202.4	17,609	143.6	141.5-145.8
2009	106,934	501.7	498.7-504.7	54,095	577.5	572.6-582.5	52,839	450.0	446.1-453.9	34,816	162.0	160.3-163.8	17,494	196.4	193.4-199.3	17,322	139.8	137.7-142.0
2010	104,820	484.8	481.8-487.8	52,754	552.4	547.6-557.2	52,066	439.3	435.5-443.2	35,024	160.1	158.4-161.8	17,338	190.5	187.6-193.4	17,686	140.5	138.4-142.6
2011	108,837	495.3	492.3-498.3	54,989	563.9	559.1-568.8	53,848	448.6	444.8-452.5	35,030	157.5	155.9-159.2	17,315	186.1	183.3-189.0	17,715	138.6	136.5-140.7
2012	106,034	475.0	472.1-477.9	52,723	530.2	525.6-534.8	53,311	439.4	435.6-443.3	35,591	157.7	156.0-159.3	17,796	187.8	185.0-190.7	17,795	137.7	135.6-139.8
2013	109,632	484.1	481.1-487.0	54,065	533.4	528.8-538.0	55,567	453.2	449.4-457.1	35,082	152.2	150.6-153.9	17,605	181.1	178.4-183.8	17,477	132.7	130.7-134.7
2014	109,505	475.6	472.8-478.5	53,806	519.3	514.8-523.8	55,699	448.7	444.9-452.6	34,679	147.8	146.2-149.4	17,480	175.9	173.3-178.6	17,199	128.8	126.9-130.8
2015	111,039	475.8	473.0-478.7	54,567	517.1	512.7-521.6	56,472	451.0	447.2-454.9	34,464	144.6	143.0-146.1	17,155	169.5	167.0-172.2	17,309	127.7	125.8-129.7
2016	112,357	474.8	472.0-477.7	56,050	521.5	517.1-526.0	56,307	445.3	441.5-449.1	34,589	142.9	141.4-144.5	17,327	167.7	165.1-170.2	17,262	126.3	124.3-128.2
2017	114,325	477.3	474.5-480.2	56,885	520.8	516.4-525.2	57,440	450.1	446.3-453.9	34,229	138.9	137.4-140.4	17,214	163.5	161.0-166.0	17,015	122.2	120.3-124.1
2018	113,661	468.8	466.0-471.6	56,760	510.3	506.0-514.6	56,901	441.9	438.1-445.7	33,714	134.6	133.2-136.1	16,797	156.4	154.0-158.8	16,917	120.0	118.1-121.8
2019	117,487	479.4	476.6-482.2	58,751	519.8	515.5-524.1	58,736	453.6	449.7-457.4	33,497	131.8	130.4-133.3	16,718	152.8	150.5-155.2	16,779	117.2	115.4-119.0
2020	103,544	419.1	416.5-421.7	51,769	453.8	449.8-457.9	51,775	397.0	393.4-400.5	32,145	125.2	123.8-126.6	16,018	144.8	142.5-147.1	16,127	111.8	110.1-113.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Oral cavity and pharynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,933	11.3	10.8-11.9	1,313	18.0	17.0-19.1	620	6.4	5.9-7.0	838	4.9	4.6-5.3	599	8.3	7.6-9.1	239	2.4	2.1-2.8
1977	1,901	11.0	10.5-11.5	1,281	17.4	16.5-18.5	620	6.3	5.8-6.8	843	4.9	4.6-5.3	591	8.2	7.5-8.9	252	2.5	2.2-2.9
1978	1,924	11.2	10.7-11.7	1,276	17.3	16.3-18.3	648	6.7	6.2-7.2	837	4.9	4.5-5.2	581	8.0	7.4-8.8	256	2.6	2.3-3.0
1979	1,969	11.4	10.9-11.9	1,314	17.8	16.8-18.8	655	6.7	6.2-7.2	808	4.7	4.4-5.0	542	7.4	6.7-8.0	266	2.7	2.4-3.1
1980	1,932	11.2	10.7-11.8	1,275	17.4	16.4-18.4	657	6.7	6.2-7.3	793	4.6	4.3-4.9	546	7.6	6.9-8.3	247	2.4	2.1-2.8
1981	2,043	11.8	11.3-12.4	1,420	19.1	18.1-20.1	623	6.3	5.8-6.9	768	4.4	4.1-4.7	523	7.1	6.5-7.7	245	2.4	2.1-2.8
1982	2,103	12.2	11.7-12.8	1,415	19.3	18.3-20.4	688	7.1	6.5-7.6	796	4.6	4.3-5.0	566	7.9	7.2-8.6	230	2.3	2.0-2.6
1983	2,008	11.5	11.0-12.0	1,357	18.1	17.1-19.2	651	6.6	6.1-7.1	774	4.4	4.1-4.7	545	7.3	6.7-8.0	229	2.3	2.0-2.6
1984	2,048	11.8	11.2-12.3	1,384	18.7	17.7-19.7	664	6.7	6.2-7.2	739	4.2	3.9-4.5	508	7.0	6.3-7.6	231	2.3	2.0-2.6
1985	2,076	11.9	11.4-12.5	1,365	18.4	17.4-19.4	711	7.1	6.6-7.7	715	4.1	3.8-4.4	478	6.5	6.0-7.2	237	2.3	2.0-2.7
1986	1,984	11.4	10.9-11.9	1,330	18.0	17.0-19.0	654	6.5	6.0-7.0	742	4.2	3.9-4.5	498	6.7	6.1-7.4	244	2.4	2.1-2.7
1987	2,003	11.5	11.0-12.0	1,303	17.4	16.4-18.4	700	7.0	6.5-7.5	679	3.9	3.6-4.2	453	6.3	5.7-6.9	226	2.2	1.9-2.5
1988	1,910	10.8	10.4-11.3	1,229	16.2	15.3-17.2	681	6.7	6.2-7.3	755	4.4	4.0-4.7	501	7.0	6.4-7.7	254	2.5	2.2-2.8
1989	1,911	10.8	10.3-11.3	1,264	16.6	15.7-17.6	647	6.5	6.0-7.0	652	3.7	3.4-4.0	420	5.7	5.2-6.3	232	2.3	2.0-2.6
1990	1,921	10.9	10.4-11.4	1,254	16.5	15.6-17.5	667	6.6	6.1-7.1	626	3.5	3.3-3.8	403	5.5	4.9-6.0	223	2.1	1.9-2.4
1991	2,073	11.8	11.3-12.3	1,354	17.8	16.8-18.8	719	7.1	6.6-7.7	654	3.7	3.4-4.0	428	5.8	5.2-6.4	226	2.2	1.9-2.5
1992	2,085	11.7	11.2-12.2	1,356	17.5	16.6-18.5	729	7.2	6.7-7.8	594	3.3	3.0-3.6	380	5.0	4.5-5.5	214	2.0	1.7-2.3
1993	2,000	11.2	10.7-11.7	1,321	17.2	16.3-18.2	679	6.6	6.1-7.1	573	3.2	2.9-3.4	382	5.0	4.5-5.5	191	1.8	1.6-2.1
1994	1,957	10.8	10.3-11.3	1,290	16.5	15.6-17.4	667	6.5	6.0-7.0	577	3.2	2.9-3.4	413	5.4	4.9-5.9	164	1.6	1.3-1.8
1995	1,770	9.7	9.3-10.2	1,145	14.4	13.6-15.3	625	6.0	5.5-6.5	604	3.3	3.1-3.6	407	5.3	4.8-5.9	197	1.8	1.6-2.1
1996	1,955	10.7	10.2-11.1	1,285	15.9	15.1-16.8	670	6.5	6.0-7.0	543	3.0	2.7-3.2	352	4.5	4.0-5.0	191	1.8	1.5-2.1
1997	1,963	10.6	10.1-11.1	1,268	15.6	14.7-16.5	695	6.6	6.1-7.1	517	2.8	2.5-3.0	358	4.5	4.1-5.0	159	1.5	1.2-1.7
1998	1,933	10.3	9.9-10.8	1,226	14.9	14.1-15.8	707	6.7	6.2-7.2	533	2.8	2.6-3.1	346	4.4	3.9-4.8	187	1.7	1.4-1.9
1999	1,928	10.1	9.7-10.6	1,325	15.8	14.9-16.7	603	5.6	5.1-6.0	502	2.6	2.4-2.9	335	4.1	3.7-4.6	167	1.5	1.3-1.7
2000	1,847	9.6	9.1-10.0	1,215	14.4	13.6-15.2	632	5.8	5.4-6.3	468	2.4	2.2-2.6	303	3.7	3.3-4.1	165	1.5	1.2-1.7
2001	1,909	9.8	9.3-10.2	1,250	14.3	13.6-15.2	659	5.9	5.5-6.4	503	2.6	2.3-2.8	334	4.0	3.5-4.4	169	1.4	1.2-1.7
2002	1,982	10.0	9.6-10.4	1,291	14.6	13.8-15.5	691	6.2	5.7-6.7	454	2.3	2.1-2.5	297	3.5	3.1-3.9	157	1.3	1.1-1.6
2003	1,955	9.7	9.3-10.2	1,288	14.4	13.7-15.3	667	5.9	5.5-6.4	455	2.3	2.1-2.5	301	3.5	3.1-3.9	154	1.3	1.1-1.5
2004	2,056	10.1	9.7-10.6	1,369	15.2	14.4-16.0	687	6.1	5.6-6.6	529	2.6	2.4-2.8	355	4.0	3.6-4.4	174	1.5	1.3-1.7
2005	1,978	9.6	9.2-10.1	1,314	14.2	13.5-15.0	664	5.9	5.4-6.3	459	2.2	2.0-2.4	305	3.4	3.0-3.8	154	1.3	1.1-1.5
2006	2,075	10.0	9.6-10.5	1,414	15.2	14.4-16.0	661	5.8	5.4-6.3	439	2.1	1.9-2.3	306	3.4	3.0-3.8	133	1.1	0.9-1.3
2007	2,235	10.7	10.2-11.1	1,521	16.1	15.3-17.0	714	6.2	5.8-6.7	453	2.1	1.9-2.3	313	3.4	3.0-3.8	140	1.2	1.0-1.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Oral cavity and pharynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	2,211	10.4	10.0-10.9	1,464	15.3	14.5-16.1	747	6.4	5.9-6.9	448	2.1	1.9-2.3	320	3.4	3.0-3.8	128	1.0	0.9-1.2
2009	2,298	10.6	10.2-11.1	1,574	16.0	15.2-16.8	724	6.1	5.7-6.6	469	2.1	2.0-2.4	320	3.4	3.0-3.8	149	1.2	1.0-1.4
2010	2,326	10.6	10.1-11.0	1,543	15.5	14.7-16.3	783	6.6	6.1-7.0	488	2.2	2.0-2.4	317	3.3	2.9-3.7	171	1.4	1.2-1.6
2011	2,359	10.6	10.1-11.0	1,638	16.0	15.2-16.8	721	6.0	5.5-6.4	486	2.1	2.0-2.4	327	3.3	2.9-3.7	159	1.2	1.0-1.5
2012	2,397	10.5	10.1-11.0	1,609	15.5	14.8-16.3	788	6.4	5.9-6.8	501	2.2	2.0-2.4	345	3.4	3.1-3.8	156	1.2	1.0-1.4
2013	2,444	10.6	10.2-11.1	1,704	16.3	15.5-17.1	740	5.9	5.5-6.3	505	2.1	2.0-2.3	354	3.4	3.0-3.8	151	1.1	0.9-1.3
2014	2,550	10.9	10.5-11.3	1,696	15.8	15.0-16.6	854	6.8	6.3-7.3	439	1.9	1.7-2.0	303	2.9	2.6-3.3	136	1.0	0.9-1.2
2015	2,675	11.3	10.9-11.8	1,857	17.2	16.4-18.0	818	6.5	6.0-7.0	525	2.2	2.0-2.4	363	3.4	3.1-3.8	162	1.2	1.0-1.4
2016	2,799	11.7	11.2-12.1	1,926	17.5	16.7-18.4	873	6.8	6.3-7.3	536	2.2	2.0-2.4	373	3.4	3.1-3.8	163	1.2	1.0-1.4
2017	2,750	11.4	10.9-11.8	1,874	16.9	16.2-17.7	876	6.6	6.2-7.1	539	2.2	2.0-2.4	378	3.4	3.1-3.8	161	1.2	1.0-1.4
2018	2,761	11.2	10.8-11.7	1,880	16.6	15.9-17.4	881	6.7	6.2-7.1	531	2.1	1.9-2.3	360	3.2	2.9-3.6	171	1.2	1.0-1.4
2019	2,896	11.7	11.3-12.2	1,941	16.9	16.1-17.7	955	7.3	6.8-7.8	550	2.1	2.0-2.3	392	3.5	3.1-3.8	158	1.1	0.9-1.2
2020	2,658	10.5	10.1-11.0	1,822	15.6	14.9-16.4	836	6.2	5.8-6.6	550	2.1	2.0-2.3	367	3.2	2.8-3.5	183	1.3	1.1-1.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Esophagus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	856	5.0	4.7-5.4	616	8.5	7.8-9.2	240	2.5	2.2-2.8	813	4.8	4.4-5.1	599	8.3	7.6-9.0	214	2.2	1.9-2.6
1977	854	5.1	4.7-5.4	585	8.2	7.5-8.9	269	2.8	2.5-3.2	778	4.6	4.2-4.9	539	7.6	6.9-8.3	239	2.4	2.1-2.8
1978	879	5.2	4.8-5.5	607	8.4	7.8-9.2	272	2.8	2.5-3.2	811	4.8	4.4-5.1	561	8.0	7.3-8.7	250	2.6	2.2-2.9
1979	884	5.1	4.8-5.4	618	8.5	7.8-9.2	266	2.7	2.4-3.0	783	4.6	4.3-4.9	547	7.5	6.9-8.2	236	2.4	2.1-2.8
1980	938	5.4	5.1-5.8	654	9.2	8.4-9.9	284	2.8	2.5-3.2	785	4.5	4.2-4.9	538	7.5	6.9-8.2	247	2.4	2.1-2.8
1981	901	5.2	4.9-5.6	614	8.4	7.7-9.1	287	2.9	2.6-3.2	823	4.7	4.4-5.1	593	8.2	7.5-8.9	230	2.3	2.0-2.6
1982	882	5.0	4.7-5.4	591	8.1	7.4-8.8	291	2.8	2.5-3.2	777	4.4	4.1-4.8	540	7.5	6.9-8.2	237	2.3	2.0-2.6
1983	896	5.1	4.7-5.4	628	8.5	7.8-9.2	268	2.6	2.3-3.0	793	4.6	4.2-4.9	571	8.0	7.3-8.7	222	2.2	1.9-2.5
1984	914	5.1	4.8-5.5	626	8.5	7.8-9.2	288	2.8	2.5-3.1	802	4.5	4.2-4.8	549	7.5	6.8-8.2	253	2.5	2.2-2.8
1985	863	4.9	4.6-5.3	617	8.4	7.8-9.1	246	2.4	2.1-2.7	786	4.4	4.1-4.8	582	7.8	7.2-8.5	204	2.0	1.7-2.2
1986	937	5.3	5.0-5.7	651	8.9	8.2-9.6	286	2.8	2.5-3.2	806	4.6	4.3-4.9	583	8.1	7.4-8.8	223	2.2	1.9-2.5
1987	855	4.8	4.5-5.2	562	7.6	6.9-8.2	293	2.8	2.5-3.2	777	4.4	4.1-4.7	545	7.4	6.8-8.1	232	2.2	2.0-2.6
1988	870	4.9	4.6-5.3	604	8.1	7.5-8.8	266	2.5	2.2-2.9	841	4.8	4.4-5.1	619	8.5	7.8-9.2	222	2.1	1.8-2.4
1989	930	5.2	4.9-5.6	650	8.7	8.0-9.4	280	2.7	2.4-3.0	803	4.5	4.2-4.9	575	7.7	7.1-8.4	228	2.2	1.9-2.5
1990	945	5.3	5.0-5.6	630	8.3	7.7-9.0	315	3.0	2.7-3.4	812	4.5	4.2-4.8	560	7.5	6.9-8.1	252	2.4	2.1-2.7
1991	902	5.0	4.7-5.3	615	8.1	7.5-8.8	287	2.7	2.4-3.1	804	4.4	4.1-4.8	560	7.4	6.8-8.1	244	2.3	2.0-2.6
1992	985	5.5	5.1-5.8	662	8.7	8.0-9.4	323	3.0	2.7-3.4	828	4.6	4.3-4.9	594	7.8	7.2-8.5	234	2.2	1.9-2.5
1993	966	5.3	5.0-5.6	678	8.7	8.1-9.4	288	2.7	2.4-3.0	842	4.6	4.3-5.0	617	8.1	7.5-8.8	225	2.1	1.8-2.4
1994	1,014	5.5	5.2-5.9	712	9.2	8.6-10.0	302	2.8	2.5-3.1	843	4.6	4.3-4.9	597	7.7	7.1-8.4	246	2.3	2.0-2.6
1995	926	5.1	4.7-5.4	626	8.1	7.5-8.8	300	2.8	2.5-3.1	871	4.8	4.4-5.1	637	8.3	7.7-9.0	234	2.1	1.9-2.4
1996	973	5.3	4.9-5.6	662	8.4	7.8-9.1	311	2.8	2.5-3.2	849	4.6	4.3-4.9	588	7.6	7.0-8.2	261	2.4	2.1-2.7
1997	909	4.9	4.6-5.2	623	8.0	7.4-8.6	286	2.6	2.3-2.9	820	4.4	4.1-4.7	602	7.8	7.1-8.4	218	1.9	1.7-2.2
1998	988	5.2	4.9-5.6	718	9.0	8.3-9.7	270	2.5	2.2-2.8	843	4.5	4.2-4.8	609	7.7	7.1-8.3	234	2.1	1.8-2.4
1999	964	5.0	4.7-5.4	702	8.6	8.0-9.3	262	2.3	2.1-2.6	829	4.3	4.0-4.6	610	7.6	7.0-8.2	219	1.9	1.7-2.2
2000	995	5.1	4.8-5.5	681	8.3	7.7-9.0	314	2.7	2.4-3.0	886	4.6	4.3-4.9	616	7.6	7.0-8.2	270	2.3	2.1-2.6
2001	945	4.8	4.5-5.1	687	8.2	7.6-8.9	258	2.2	2.0-2.5	846	4.3	4.0-4.6	645	7.8	7.2-8.4	201	1.7	1.5-2.0
2002	986	5.0	4.7-5.3	719	8.5	7.9-9.1	267	2.3	2.0-2.6	828	4.2	3.9-4.5	606	7.2	6.7-7.8	222	1.9	1.6-2.1
2003	1,037	5.2	4.9-5.5	749	8.7	8.1-9.3	288	2.4	2.1-2.7	891	4.4	4.1-4.7	637	7.5	6.9-8.1	254	2.1	1.9-2.4
2004	1,045	5.1	4.8-5.5	756	8.7	8.1-9.4	289	2.4	2.1-2.7	913	4.5	4.2-4.8	678	7.8	7.3-8.5	235	1.9	1.7-2.2
2005	1,092	5.3	5.0-5.7	790	9.0	8.4-9.6	302	2.6	2.3-2.9	941	4.6	4.3-4.9	707	8.2	7.6-8.8	234	1.9	1.7-2.2
2006	1,089	5.3	5.0-5.6	833	9.4	8.8-10.1	256	2.2	1.9-2.5	869	4.2	4.0-4.5	633	7.2	6.7-7.8	236	2.0	1.7-2.3
2007	1,048	5.0	4.7-5.3	778	8.5	8.0-9.2	270	2.3	2.0-2.6	900	4.3	4.0-4.6	667	7.5	6.9-8.1	233	1.9	1.7-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Esophagus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	1,128	5.3	5.0-5.6	844	9.2	8.6-9.8	284	2.3	2.1-2.6	889	4.2	3.9-4.5	674	7.3	6.8-7.9	215	1.8	1.6-2.0
2009	1,116	5.2	4.9-5.5	854	9.0	8.4-9.7	262	2.1	1.9-2.4	883	4.1	3.8-4.4	692	7.5	6.9-8.0	191	1.5	1.3-1.7
2010	1,020	4.6	4.4-4.9	750	7.8	7.3-8.4	270	2.1	1.9-2.4	880	4.0	3.7-4.3	680	7.2	6.6-7.8	200	1.6	1.3-1.8
2011	1,042	4.7	4.4-5.0	774	7.8	7.3-8.4	268	2.1	1.9-2.4	896	4.0	3.8-4.3	674	7.0	6.5-7.5	222	1.7	1.5-2.0
2012	975	4.3	4.0-4.5	748	7.5	6.9-8.0	227	1.7	1.5-2.0	829	3.6	3.4-3.9	634	6.5	6.0-7.0	195	1.4	1.2-1.7
2013	1,075	4.6	4.4-4.9	815	8.0	7.4-8.6	260	2.0	1.7-2.2	871	3.7	3.5-4.0	661	6.5	6.0-7.0	210	1.6	1.4-1.8
2014	1,080	4.5	4.2-4.8	818	7.7	7.2-8.3	262	1.9	1.7-2.2	895	3.8	3.6-4.1	703	6.9	6.4-7.5	192	1.4	1.2-1.6
2015	1,106	4.6	4.3-4.9	815	7.6	7.1-8.2	291	2.2	2.0-2.5	817	3.4	3.2-3.6	609	5.8	5.3-6.3	208	1.5	1.3-1.7
2016	1,133	4.6	4.4-4.9	855	7.9	7.3-8.4	278	2.1	1.8-2.3	872	3.6	3.3-3.8	675	6.3	5.8-6.8	197	1.4	1.2-1.7
2017	1,041	4.2	4.0-4.5	785	7.1	6.6-7.6	256	1.9	1.7-2.2	852	3.4	3.2-3.6	676	6.1	5.6-6.6	176	1.3	1.1-1.5
2018	1,094	4.4	4.1-4.6	834	7.5	7.0-8.0	260	1.8	1.6-2.1	808	3.2	3.0-3.4	619	5.5	5.1-6.0	189	1.3	1.1-1.5
2019	1,127	4.4	4.1-4.6	852	7.4	6.9-7.9	275	1.9	1.7-2.1	816	3.1	2.9-3.4	638	5.6	5.2-6.1	178	1.2	1.0-1.4
2020	1,005	3.9	3.6-4.1	759	6.6	6.1-7.1	246	1.7	1.5-1.9	796	3.1	2.9-3.3	615	5.4	5.0-5.9	181	1.2	1.0-1.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Stomach**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	2,107	13.1	12.5-13.6	1,211	18.3	17.2-19.4	896	9.5	8.9-10.1	1,658	10.4	9.9-10.9	935	14.5	13.5-15.5	723	7.7	7.1-8.3
1977	2,106	13.1	12.5-13.7	1,218	18.6	17.5-19.8	888	9.4	8.7-10.0	1,632	10.2	9.7-10.7	935	14.7	13.8-15.8	697	7.3	6.8-7.9
1978	2,066	12.6	12.1-13.2	1,160	17.6	16.6-18.7	906	9.3	8.7-10.0	1,530	9.5	9.0-10.0	843	13.1	12.1-14.0	687	7.2	6.6-7.7
1979	2,109	12.8	12.2-13.3	1,198	17.9	16.9-19.0	911	9.3	8.7-10.0	1,522	9.3	8.8-9.8	859	13.1	12.2-14.1	663	6.7	6.2-7.3
1980	2,128	12.9	12.3-13.4	1,246	18.6	17.6-19.7	882	9.0	8.4-9.7	1,551	9.5	9.0-10.0	881	13.5	12.6-14.5	670	6.8	6.3-7.4
1981	2,147	12.8	12.3-13.4	1,260	18.7	17.6-19.8	887	8.9	8.3-9.5	1,547	9.3	8.8-9.8	915	13.9	12.9-14.8	632	6.3	5.9-6.9
1982	2,058	12.1	11.6-12.7	1,186	17.6	16.5-18.6	872	8.6	8.1-9.2	1,423	8.5	8.1-9.0	843	13.0	12.1-13.9	580	5.7	5.3-6.2
1983	2,126	12.5	12.0-13.1	1,189	17.5	16.5-18.5	937	9.2	8.6-9.8	1,443	8.5	8.1-9.0	830	12.5	11.6-13.4	613	6.0	5.5-6.5
1984	2,008	11.6	11.1-12.1	1,166	16.7	15.7-17.7	842	8.1	7.6-8.7	1,429	8.3	7.9-8.8	799	11.6	10.8-12.5	630	6.1	5.6-6.6
1985	2,043	11.9	11.3-12.4	1,210	17.6	16.5-18.6	833	8.1	7.5-8.6	1,374	8.0	7.6-8.4	783	11.6	10.7-12.4	591	5.7	5.2-6.2
1986	1,999	11.5	11.0-12.0	1,182	17.1	16.1-18.1	817	7.8	7.3-8.4	1,364	7.9	7.5-8.3	804	11.8	11.0-12.7	560	5.3	4.9-5.8
1987	2,046	11.7	11.2-12.2	1,203	17.1	16.1-18.1	843	7.9	7.4-8.5	1,359	7.8	7.4-8.2	791	11.5	10.7-12.4	568	5.4	4.9-5.8
1988	1,954	11.2	10.7-11.7	1,109	15.8	14.8-16.8	845	8.0	7.5-8.6	1,337	7.7	7.3-8.1	738	10.6	9.9-11.5	599	5.7	5.2-6.2
1989	1,889	10.7	10.2-11.2	1,136	16.0	15.0-17.0	753	7.0	6.5-7.6	1,381	7.8	7.4-8.3	799	11.6	10.7-12.4	582	5.4	5.0-5.8
1990	2,100	11.9	11.4-12.4	1,248	17.6	16.6-18.6	852	8.0	7.5-8.6	1,282	7.3	6.9-7.7	762	11.0	10.2-11.8	520	4.8	4.4-5.3
1991	1,988	11.2	10.7-11.7	1,190	16.7	15.7-17.7	798	7.5	6.9-8.0	1,288	7.3	6.9-7.7	788	11.4	10.6-12.2	500	4.6	4.2-5.1
1992	2,009	11.2	10.7-11.7	1,173	16.3	15.3-17.2	836	7.7	7.2-8.3	1,350	7.6	7.2-8.0	772	11.1	10.3-11.9	578	5.3	4.9-5.8
1993	2,080	11.5	11.0-12.0	1,224	16.9	16.0-17.9	856	7.8	7.3-8.3	1,274	7.1	6.7-7.5	737	10.5	9.7-11.3	537	4.9	4.5-5.3
1994	2,091	11.5	11.0-12.0	1,196	16.2	15.3-17.2	895	8.2	7.6-8.7	1,349	7.4	7.0-7.8	787	11.0	10.2-11.8	562	5.1	4.7-5.5
1995	1,964	10.7	10.2-11.2	1,127	15.2	14.3-16.1	837	7.6	7.1-8.1	1,267	6.9	6.6-7.3	694	9.7	9.0-10.5	573	5.1	4.7-5.6
1996	2,000	10.8	10.4-11.3	1,166	15.6	14.7-16.6	834	7.5	7.0-8.0	1,220	6.6	6.3-7.0	679	9.3	8.6-10.1	541	4.8	4.4-5.2
1997	1,977	10.6	10.1-11.1	1,165	15.4	14.5-16.4	812	7.3	6.8-7.8	1,148	6.2	5.8-6.5	636	8.5	7.8-9.2	512	4.6	4.2-5.0
1998	2,068	11.0	10.5-11.4	1,226	15.8	14.9-16.7	842	7.5	7.0-8.1	1,155	6.1	5.8-6.5	645	8.5	7.9-9.2	510	4.4	4.1-4.9
1999	1,942	10.1	9.7-10.6	1,177	15.0	14.2-15.9	765	6.7	6.2-7.2	1,109	5.8	5.4-6.1	636	8.3	7.7-9.0	473	4.1	3.7-4.5
2000	1,942	10.0	9.5-10.4	1,113	14.1	13.3-14.9	829	7.1	6.6-7.6	1,076	5.5	5.2-5.9	588	7.6	7.0-8.3	488	4.1	3.8-4.5
2001	1,918	9.8	9.3-10.2	1,130	14.0	13.1-14.8	788	6.8	6.3-7.3	1,021	5.2	4.9-5.5	572	7.2	6.6-7.8	449	3.8	3.4-4.2
2002	1,874	9.4	9.0-9.9	1,088	13.2	12.4-14.0	786	6.7	6.3-7.2	989	5.0	4.7-5.3	559	7.0	6.4-7.6	430	3.6	3.3-3.9
2003	1,994	9.9	9.5-10.3	1,173	14.0	13.2-14.8	821	7.0	6.5-7.5	1,003	5.0	4.7-5.3	558	6.8	6.3-7.4	445	3.7	3.4-4.1
2004	1,908	9.4	9.0-9.9	1,107	13.2	12.4-14.0	801	6.8	6.3-7.3	1,053	5.2	4.9-5.5	573	7.0	6.4-7.6	480	4.0	3.6-4.4
2005	1,930	9.5	9.0-9.9	1,118	13.0	12.3-13.8	812	6.8	6.4-7.3	1,011	4.9	4.6-5.2	580	7.0	6.4-7.6	431	3.5	3.2-3.8
2006	1,861	9.1	8.7-9.5	1,094	12.6	11.8-13.3	767	6.5	6.0-7.0	905	4.4	4.1-4.7	512	6.0	5.5-6.5	393	3.2	2.9-3.6
2007	1,928	9.3	8.9-9.7	1,160	13.1	12.3-13.8	768	6.5	6.0-6.9	872	4.2	3.9-4.5	490	5.6	5.1-6.1	382	3.1	2.8-3.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Stomach**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	1,775	8.4	8.0-8.8	1,091	12.3	11.5-13.0	684	5.7	5.2-6.1	904	4.2	4.0-4.5	544	6.2	5.6-6.7	360	2.9	2.6-3.2
2009	1,808	8.5	8.1-8.9	1,101	12.0	11.3-12.8	707	5.8	5.3-6.2	859	4.0	3.8-4.3	505	5.7	5.2-6.2	354	2.8	2.5-3.1
2010	1,872	8.6	8.3-9.1	1,121	11.9	11.2-12.6	751	6.1	5.7-6.6	811	3.7	3.5-4.0	453	5.0	4.6-5.5	358	2.8	2.5-3.1
2011	1,924	8.7	8.3-9.1	1,139	12.0	11.3-12.7	785	6.3	5.9-6.8	817	3.7	3.4-3.9	470	5.0	4.6-5.5	347	2.7	2.4-3.0
2012	1,955	8.8	8.4-9.2	1,168	12.0	11.3-12.7	787	6.3	5.9-6.8	873	3.9	3.6-4.2	500	5.3	4.8-5.7	373	2.9	2.6-3.2
2013	2,101	9.3	8.9-9.7	1,270	12.8	12.1-13.6	831	6.5	6.1-7.0	853	3.7	3.4-4.0	512	5.2	4.8-5.7	341	2.6	2.3-2.9
2014	1,979	8.6	8.2-9.0	1,238	12.3	11.6-13.0	741	5.8	5.4-6.3	875	3.8	3.5-4.1	529	5.3	4.8-5.8	346	2.7	2.4-3.0
2015	2,006	8.5	8.2-8.9	1,182	11.3	10.7-12.0	824	6.4	5.9-6.8	778	3.3	3.1-3.6	457	4.6	4.1-5.0	321	2.4	2.1-2.7
2016	2,082	8.8	8.4-9.2	1,255	11.9	11.2-12.6	827	6.4	6.0-6.9	849	3.5	3.3-3.8	542	5.2	4.7-5.6	307	2.3	2.0-2.5
2017	2,095	8.8	8.4-9.2	1,216	11.4	10.8-12.1	879	6.7	6.3-7.2	791	3.3	3.1-3.5	447	4.2	3.8-4.7	344	2.6	2.3-2.9
2018	1,979	8.2	7.8-8.6	1,162	10.7	10.1-11.3	817	6.3	5.8-6.7	765	3.1	2.9-3.4	447	4.2	3.8-4.6	318	2.3	2.0-2.6
2019	2,079	8.4	8.1-8.8	1,236	11.1	10.5-11.8	843	6.3	5.9-6.8	822	3.4	3.1-3.6	491	4.6	4.2-5.0	331	2.4	2.1-2.7
2020	1,811	7.4	7.0-7.7	1,088	9.8	9.2-10.4	723	5.5	5.1-5.9	710	2.8	2.6-3.1	417	3.8	3.4-4.2	293	2.1	1.9-2.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Colorectal**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	10,933	66.7	65.5-68.0	5,305	79.5	77.3-81.8	5,628	58.5	57.0-60.1	5,720	35.8	34.9-36.8	2,789	43.8	42.1-45.5	2,931	30.9	29.8-32.0
1977	11,133	67.5	66.2-68.8	5,391	81.0	78.7-83.3	5,742	59.0	57.5-60.6	5,548	34.5	33.6-35.4	2,689	42.3	40.6-44.1	2,859	29.7	28.6-30.8
1978	11,121	67.2	66.0-68.5	5,396	81.0	78.7-83.3	5,725	58.5	57.0-60.1	5,670	35.0	34.1-36.0	2,795	43.6	41.9-45.4	2,875	29.7	28.6-30.8
1979	11,293	67.7	66.4-69.0	5,426	80.9	78.6-83.2	5,867	59.3	57.8-60.9	5,639	34.4	33.5-35.3	2,676	41.7	40.1-43.4	2,963	30.0	28.9-31.1
1980	11,686	69.2	68.0-70.5	5,808	85.6	83.3-87.9	5,878	58.7	57.2-60.3	5,629	34.1	33.2-35.1	2,758	42.9	41.2-44.6	2,871	28.9	27.8-29.9
1981	11,695	69.3	68.0-70.5	5,749	85.8	83.5-88.1	5,946	58.9	57.4-60.4	5,526	33.3	32.4-34.2	2,629	40.6	39.0-42.3	2,897	28.8	27.8-29.9
1982	11,707	68.6	67.4-69.9	5,708	83.9	81.6-86.2	5,999	58.9	57.4-60.4	5,446	32.5	31.6-33.4	2,677	41.2	39.5-42.8	2,769	27.3	26.3-28.3
1983	11,754	68.3	67.1-69.6	5,771	84.3	82.0-86.6	5,983	58.3	56.8-59.8	5,396	31.9	31.0-32.8	2,615	39.8	38.3-41.5	2,781	27.0	26.0-28.0
1984	11,764	68.1	66.9-69.4	5,686	82.4	80.2-84.6	6,078	58.9	57.4-60.4	5,588	32.7	31.9-33.6	2,756	41.7	40.1-43.3	2,832	27.4	26.3-28.4
1985	12,091	69.3	68.0-70.5	6,023	86.6	84.3-88.9	6,068	58.3	56.8-59.8	5,402	31.5	30.6-32.3	2,602	38.8	37.3-40.4	2,800	26.8	25.8-27.8
1986	11,909	68.0	66.8-69.3	5,868	84.4	82.2-86.7	6,041	57.8	56.3-59.2	5,353	31.0	30.1-31.8	2,635	39.4	37.9-41.0	2,718	25.9	24.9-26.9
1987	11,392	64.6	63.4-65.8	5,622	80.4	78.3-82.7	5,770	54.6	53.2-56.0	5,326	30.6	29.8-31.4	2,630	39.2	37.6-40.7	2,696	25.3	24.4-26.3
1988	11,058	62.5	61.3-63.7	5,343	75.8	73.7-78.0	5,715	53.8	52.4-55.2	5,211	29.8	29.0-30.6	2,616	38.7	37.2-40.3	2,595	24.3	23.4-25.3
1989	10,852	61.3	60.1-62.4	5,372	76.0	73.9-78.2	5,480	51.7	50.4-53.1	4,890	27.8	27.0-28.6	2,353	34.7	33.3-36.2	2,537	23.6	22.7-24.6
1990	10,858	60.7	59.6-61.9	5,223	72.9	70.9-75.0	5,635	52.4	51.0-53.8	4,837	27.4	26.6-28.1	2,394	35.0	33.6-36.5	2,443	22.6	21.7-23.5
1991	10,912	60.9	59.8-62.1	5,372	75.4	73.4-77.6	5,540	51.6	50.2-53.0	4,819	27.1	26.3-27.9	2,323	34.0	32.6-35.5	2,496	22.9	22.0-23.8
1992	10,675	59.1	58.0-60.3	5,195	72.1	70.1-74.2	5,480	50.8	49.4-52.1	4,601	25.6	24.9-26.4	2,198	31.7	30.3-33.1	2,403	21.9	21.0-22.8
1993	10,806	59.3	58.2-60.4	5,201	71.0	69.0-73.0	5,605	51.4	50.1-52.8	4,644	25.6	24.9-26.4	2,242	31.9	30.5-33.3	2,402	21.6	20.8-22.5
1994	11,392	62.2	61.1-63.4	5,491	74.8	72.8-76.8	5,901	54.0	52.6-55.4	4,799	26.3	25.6-27.1	2,300	32.8	31.4-34.2	2,499	22.3	21.5-23.2
1995	11,021	59.8	58.7-60.9	5,327	72.0	70.0-74.0	5,694	51.7	50.3-53.0	4,760	25.9	25.2-26.7	2,263	32.1	30.7-33.4	2,497	22.2	21.3-23.1
1996	10,830	58.5	57.4-59.6	5,347	71.5	69.5-73.4	5,483	49.7	48.3-51.0	4,417	23.8	23.1-24.6	2,168	30.0	28.8-31.3	2,249	19.8	19.0-20.7
1997	11,363	60.9	59.7-62.0	5,503	72.5	70.5-74.4	5,860	52.6	51.3-54.0	4,415	23.7	23.0-24.4	2,178	29.8	28.5-31.1	2,237	19.6	18.8-20.4
1998	11,763	62.3	61.2-63.5	5,694	73.8	71.9-75.8	6,069	54.3	53.0-55.7	4,291	22.7	22.0-23.4	2,023	27.0	25.8-28.2	2,268	19.7	18.9-20.5
1999	11,830	61.8	60.7-62.9	5,732	73.2	71.3-75.1	6,098	53.9	52.5-55.3	4,269	22.2	21.6-22.9	2,036	27.0	25.8-28.2	2,233	19.2	18.4-20.0
2000	11,806	60.7	59.7-61.9	5,794	72.7	70.8-74.6	6,012	52.3	51.0-53.7	4,280	21.9	21.3-22.6	2,115	27.5	26.3-28.7	2,165	18.1	17.4-18.9
2001	11,406	58.1	57.0-59.1	5,586	69.2	67.4-71.1	5,820	50.3	49.0-51.6	4,097	20.8	20.1-21.4	1,946	25.1	24.0-26.2	2,151	18.0	17.2-18.8
2002	11,309	56.9	55.9-58.0	5,543	67.0	65.3-68.9	5,766	49.3	48.1-50.6	4,106	20.6	19.9-21.2	2,000	25.2	24.1-26.4	2,106	17.4	16.7-18.2
2003	11,043	54.9	53.9-56.0	5,387	64.4	62.7-66.2	5,656	48.1	46.8-49.4	3,938	19.5	18.9-20.1	1,904	23.6	22.5-24.7	2,034	16.7	16.0-17.5
2004	10,829	53.3	52.3-54.3	5,189	61.1	59.5-62.9	5,640	47.6	46.4-48.9	3,762	18.4	17.9-19.0	1,815	22.0	21.0-23.1	1,947	15.7	15.0-16.5
2005	10,362	50.5	49.6-51.5	5,127	59.7	58.1-61.4	5,235	43.9	42.7-45.1	3,492	16.9	16.4-17.5	1,673	20.3	19.4-21.3	1,819	14.6	13.9-15.3
2006	10,063	48.9	47.9-49.8	4,897	56.5	54.9-58.1	5,166	43.3	42.1-44.5	3,504	16.9	16.3-17.5	1,686	20.3	19.3-21.3	1,818	14.6	14.0-15.3
2007	10,172	48.8	47.8-49.7	5,004	56.3	54.7-57.9	5,168	43.1	41.9-44.3	3,436	16.3	15.8-16.9	1,660	19.4	18.5-20.4	1,776	14.1	13.5-14.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Colorectal**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	9,951	47.1	46.2-48.1	4,890	54.2	52.7-55.8	5,061	41.8	40.7-43.0	3,398	16.0	15.4-16.5	1,658	19.1	18.2-20.0	1,740	13.7	13.0-14.4
2009	9,546	44.5	43.6-45.4	4,710	51.1	49.6-52.6	4,836	39.3	38.2-40.5	3,318	15.4	14.9-15.9	1,614	18.3	17.4-19.3	1,704	13.3	12.6-13.9
2010	9,441	43.5	42.6-44.4	4,664	49.8	48.4-51.3	4,777	38.7	37.5-39.8	3,277	14.9	14.3-15.4	1,648	18.2	17.3-19.1	1,629	12.4	11.8-13.0
2011	9,196	41.9	41.0-42.8	4,558	48.0	46.6-49.4	4,638	37.2	36.1-38.3	3,243	14.4	13.9-14.9	1,590	17.0	16.2-17.9	1,653	12.4	11.8-13.1
2012	8,817	39.4	38.6-40.3	4,506	46.2	44.8-47.6	4,311	34.2	33.1-35.2	3,190	14.0	13.5-14.5	1,534	16.2	15.4-17.1	1,656	12.3	11.7-13.0
2013	9,089	40.0	39.2-40.8	4,620	46.5	45.1-47.8	4,469	35.0	33.9-36.0	3,118	13.4	13.0-13.9	1,590	16.4	15.6-17.2	1,528	11.1	10.6-11.7
2014	9,070	39.3	38.5-40.2	4,627	45.5	44.2-46.9	4,443	34.5	33.4-35.5	3,056	12.9	12.5-13.4	1,550	15.6	14.8-16.4	1,506	10.9	10.4-11.5
2015	8,997	38.6	37.8-39.5	4,571	44.2	42.9-45.5	4,426	34.2	33.1-35.2	3,056	12.8	12.3-13.2	1,536	15.2	14.5-16.0	1,520	10.9	10.4-11.5
2016	9,077	38.6	37.8-39.4	4,693	44.8	43.5-46.1	4,384	33.6	32.6-34.7	2,938	12.1	11.6-12.5	1,454	14.0	13.3-14.8	1,484	10.6	10.0-11.1
2017	9,076	38.1	37.3-39.0	4,658	43.7	42.4-45.0	4,418	33.7	32.7-34.8	3,093	12.6	12.1-13.0	1,530	14.6	13.8-15.3	1,563	11.1	10.5-11.6
2018	8,828	36.7	35.9-37.5	4,552	42.3	41.0-43.6	4,276	32.1	31.1-33.1	2,800	11.4	10.9-11.8	1,408	13.3	12.6-14.0	1,392	9.9	9.3-10.4
2019	9,002	37.2	36.4-38.0	4,666	42.6	41.4-43.9	4,336	32.7	31.7-33.7	2,811	11.1	10.7-11.6	1,451	13.4	12.7-14.2	1,360	9.2	8.7-9.8
2020	7,981	32.8	32.1-33.6	4,195	38.2	37.0-39.4	3,786	28.4	27.5-29.4	2,801	11.0	10.6-11.4	1,462	13.3	12.6-14.0	1,339	9.2	8.7-9.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Colon excluding rectum**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	7,658	47.0	46.0-48.1	3,530	53.7	51.9-55.6	4,128	42.9	41.6-44.3	4,517	28.3	27.5-29.2	2,111	33.3	31.8-34.8	2,406	25.4	24.4-26.4
1977	7,840	47.9	46.9-49.0	3,621	55.6	53.7-57.5	4,219	43.5	42.2-44.8	4,508	28.1	27.2-28.9	2,125	33.6	32.1-35.1	2,383	24.8	23.8-25.8
1978	7,900	48.1	47.0-49.2	3,676	56.0	54.1-57.9	4,224	43.3	42.0-44.6	4,613	28.5	27.6-29.3	2,202	34.6	33.0-36.1	2,411	24.8	23.8-25.8
1979	8,113	49.0	48.0-50.1	3,732	56.6	54.7-58.6	4,381	44.4	43.1-45.8	4,681	28.5	27.7-29.3	2,155	33.5	32.0-35.0	2,526	25.6	24.6-26.6
1980	8,307	49.6	48.5-50.7	3,984	59.7	57.8-61.7	4,323	43.2	41.9-44.5	4,669	28.3	27.5-29.1	2,249	35.1	33.6-36.7	2,420	24.3	23.3-25.3
1981	8,341	49.6	48.6-50.7	3,950	59.7	57.8-61.7	4,391	43.6	42.3-44.9	4,698	28.4	27.6-29.2	2,197	34.1	32.6-35.6	2,501	24.9	24.0-25.9
1982	8,295	48.9	47.9-50.0	3,852	57.6	55.7-59.5	4,443	43.6	42.3-44.9	4,575	27.3	26.5-28.1	2,220	34.2	32.8-35.8	2,355	23.2	22.3-24.2
1983	8,379	49.0	48.0-50.1	3,915	58.2	56.3-60.1	4,464	43.5	42.2-44.8	4,531	26.8	26.0-27.6	2,156	32.9	31.5-34.4	2,375	23.0	22.1-23.9
1984	8,488	49.4	48.3-50.4	3,910	57.5	55.7-59.4	4,578	44.2	43.0-45.6	4,742	27.8	27.0-28.6	2,301	34.8	33.4-36.4	2,441	23.6	22.6-24.5
1985	8,782	50.6	49.5-51.6	4,182	61.0	59.1-62.9	4,600	44.1	42.9-45.4	4,622	26.9	26.1-27.7	2,187	32.6	31.2-34.1	2,435	23.3	22.4-24.3
1986	8,687	49.8	48.8-50.9	4,083	59.5	57.6-61.4	4,604	44.0	42.7-45.3	4,580	26.5	25.7-27.3	2,216	33.2	31.7-34.6	2,364	22.5	21.6-23.4
1987	8,209	46.6	45.6-47.6	3,895	56.4	54.6-58.3	4,314	40.6	39.4-41.8	4,568	26.2	25.5-27.0	2,257	33.6	32.2-35.1	2,311	21.8	20.9-22.7
1988	8,018	45.4	44.4-46.4	3,687	52.9	51.1-54.7	4,331	40.6	39.4-41.9	4,515	25.8	25.1-26.6	2,251	33.3	31.9-34.7	2,264	21.2	20.3-22.1
1989	7,800	44.2	43.2-45.2	3,734	53.8	52.0-55.6	4,066	38.3	37.1-39.5	4,236	24.1	23.4-24.8	2,000	29.6	28.3-31.0	2,236	20.8	19.9-21.7
1990	7,853	44.0	43.1-45.0	3,622	51.4	49.7-53.1	4,231	39.3	38.1-40.5	4,226	23.9	23.2-24.6	2,084	30.6	29.2-32.0	2,142	19.8	19.0-20.7
1991	7,989	44.7	43.7-45.7	3,801	54.0	52.2-55.8	4,188	38.8	37.6-40.0	4,126	23.2	22.5-24.0	1,967	28.9	27.6-30.3	2,159	19.8	18.9-20.6
1992	7,755	43.0	42.1-44.0	3,609	50.8	49.1-52.5	4,146	38.3	37.1-39.5	3,944	22.0	21.3-22.7	1,848	26.8	25.5-28.0	2,096	19.1	18.3-19.9
1993	7,909	43.5	42.5-44.4	3,626	50.0	48.3-51.7	4,283	39.1	38.0-40.3	4,013	22.2	21.5-22.9	1,916	27.4	26.1-28.7	2,097	18.9	18.1-19.7
1994	8,378	45.8	44.8-46.8	3,903	53.7	52.0-55.5	4,475	40.7	39.5-41.9	4,169	22.8	22.2-23.6	1,971	28.2	26.9-29.5	2,198	19.6	18.8-20.5
1995	8,088	43.9	42.9-44.9	3,776	51.6	49.9-53.3	4,312	38.9	37.7-40.0	4,087	22.2	21.6-22.9	1,917	27.3	26.1-28.6	2,170	19.3	18.4-20.1
1996	7,844	42.3	41.4-43.3	3,704	50.1	48.5-51.8	4,140	37.2	36.1-38.4	3,787	20.4	19.8-21.1	1,845	25.7	24.5-26.9	1,942	17.1	16.3-17.9
1997	8,439	45.2	44.2-46.2	3,912	52.0	50.4-53.7	4,527	40.5	39.3-41.7	3,823	20.5	19.8-21.1	1,847	25.3	24.2-26.5	1,976	17.3	16.5-18.1
1998	8,605	45.6	44.6-46.5	4,032	52.9	51.2-54.6	4,573	40.7	39.6-42.0	3,710	19.6	19.0-20.3	1,728	23.2	22.1-24.3	1,982	17.2	16.4-18.0
1999	8,614	44.9	44.0-45.9	3,997	51.6	50.0-53.2	4,617	40.5	39.4-41.7	3,637	18.9	18.3-19.6	1,702	22.6	21.6-23.8	1,935	16.6	15.9-17.4
2000	8,646	44.5	43.5-45.4	4,042	51.3	49.7-52.9	4,604	39.8	38.7-41.0	3,671	18.8	18.2-19.4	1,792	23.5	22.4-24.6	1,879	15.7	15.0-16.5
2001	8,313	42.3	41.4-43.2	3,926	49.1	47.6-50.7	4,387	37.6	36.5-38.7	3,503	17.8	17.2-18.4	1,628	21.1	20.0-22.1	1,875	15.7	14.9-16.4
2002	8,275	41.6	40.7-42.5	3,880	47.5	46.0-49.1	4,395	37.4	36.3-38.6	3,466	17.3	16.8-17.9	1,669	21.2	20.1-22.2	1,797	14.8	14.2-15.5
2003	8,032	39.9	39.1-40.8	3,761	45.5	44.1-47.0	4,271	36.1	35.0-37.2	3,366	16.7	16.1-17.2	1,589	19.8	18.8-20.8	1,777	14.6	13.9-15.3
2004	7,835	38.6	37.7-39.4	3,612	43.1	41.6-44.5	4,223	35.4	34.3-36.5	3,177	15.6	15.0-16.1	1,499	18.3	17.3-19.2	1,678	13.5	12.9-14.2
2005	7,600	37.1	36.2-37.9	3,611	42.5	41.1-43.9	3,989	33.3	32.2-34.3	2,940	14.3	13.7-14.8	1,374	16.8	15.9-17.7	1,566	12.5	11.9-13.2
2006	7,316	35.5	34.7-36.3	3,436	40.1	38.8-41.5	3,880	32.2	31.2-33.3	2,972	14.3	13.8-14.9	1,412	17.0	16.1-17.9	1,560	12.5	11.9-13.2
2007	7,363	35.3	34.5-36.2	3,494	39.8	38.4-41.1	3,869	32.0	31.0-33.1	2,911	13.8	13.3-14.3	1,377	16.2	15.3-17.0	1,534	12.2	11.6-12.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Colon excluding rectum**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	7,209	34.1	33.3-34.9	3,386	38.0	36.7-39.3	3,823	31.4	30.4-32.5	2,810	13.2	12.7-13.7	1,347	15.6	14.8-16.4	1,463	11.5	10.9-12.1
2009	6,848	31.9	31.2-32.7	3,206	35.2	34.0-36.5	3,642	29.3	28.4-30.3	2,781	12.9	12.4-13.4	1,332	15.2	14.4-16.1	1,449	11.3	10.7-11.9
2010	6,762	31.2	30.4-32.0	3,189	34.5	33.3-35.7	3,573	28.6	27.7-29.6	2,729	12.3	11.9-12.8	1,336	14.8	14.0-15.6	1,393	10.5	10.0-11.1
2011	6,563	29.9	29.2-30.6	3,113	33.2	32.0-34.4	3,450	27.4	26.4-28.3	2,687	12.0	11.5-12.4	1,282	13.8	13.0-14.6	1,405	10.5	10.0-11.1
2012	6,338	28.3	27.6-29.1	3,063	31.8	30.6-32.9	3,275	25.7	24.8-26.6	2,649	11.6	11.2-12.1	1,228	13.0	12.3-13.8	1,421	10.6	10.0-11.1
2013	6,400	28.1	27.4-28.8	3,115	31.7	30.5-32.8	3,285	25.4	24.5-26.3	2,578	11.1	10.7-11.6	1,273	13.2	12.5-14.0	1,305	9.5	9.0-10.0
2014	6,443	27.9	27.2-28.6	3,127	31.1	30.0-32.3	3,316	25.5	24.6-26.4	2,482	10.5	10.1-10.9	1,226	12.4	11.7-13.1	1,256	9.1	8.6-9.6
2015	6,315	27.1	26.4-27.8	3,048	29.9	28.8-31.0	3,267	24.9	24.1-25.8	2,505	10.4	10.0-10.9	1,228	12.2	11.5-12.9	1,277	9.1	8.6-9.7
2016	6,468	27.4	26.8-28.1	3,186	30.6	29.5-31.7	3,282	25.0	24.1-25.9	2,331	9.6	9.2-10.0	1,115	10.8	10.1-11.4	1,216	8.6	8.1-9.1
2017	6,400	26.8	26.1-27.5	3,118	29.4	28.4-30.5	3,282	24.7	23.9-25.6	2,475	10.0	9.6-10.5	1,187	11.4	10.7-12.0	1,288	9.1	8.6-9.6
2018	6,208	25.7	25.1-26.4	3,058	28.5	27.5-29.6	3,150	23.3	22.5-24.2	2,231	9.0	8.6-9.4	1,085	10.3	9.7-10.9	1,146	8.1	7.6-8.6
2019	6,369	26.1	25.4-26.8	3,143	28.7	27.7-29.8	3,226	23.9	23.1-24.8	2,229	8.8	8.5-9.2	1,110	10.4	9.7-11.0	1,119	7.6	7.1-8.0
2020	5,618	23.0	22.3-23.6	2,800	25.5	24.6-26.5	2,818	20.8	20.0-21.7	2,172	8.5	8.1-8.9	1,102	10.0	9.4-10.7	1,070	7.3	6.9-7.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Rectum & rectosigmoid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	3,275	19.7	19.0-20.4	1,775	25.8	24.6-27.1	1,500	15.6	14.8-16.4	1,203	7.5	7.1-7.9	678	10.5	9.7-11.4	525	5.5	5.1-6.0
1977	3,293	19.5	18.9-20.2	1,770	25.4	24.2-26.7	1,523	15.6	14.8-16.4	1,040	6.4	6.0-6.8	564	8.7	8.0-9.5	476	4.9	4.5-5.4
1978	3,221	19.2	18.5-19.8	1,720	25.0	23.7-26.2	1,501	15.3	14.5-16.1	1,057	6.5	6.1-6.9	593	9.1	8.3-9.9	464	4.9	4.4-5.3
1979	3,180	18.7	18.0-19.3	1,694	24.2	23.0-25.5	1,486	14.9	14.1-15.7	958	5.9	5.5-6.3	521	8.2	7.5-9.0	437	4.4	4.0-4.9
1980	3,379	19.7	19.0-20.4	1,824	25.8	24.6-27.1	1,555	15.5	14.7-16.3	960	5.8	5.5-6.2	509	7.8	7.1-8.5	451	4.6	4.2-5.0
1981	3,354	19.6	18.9-20.3	1,799	26.1	24.9-27.4	1,555	15.3	14.6-16.1	828	4.9	4.6-5.3	432	6.6	5.9-7.3	396	3.9	3.5-4.3
1982	3,412	19.7	19.0-20.4	1,856	26.3	25.0-27.5	1,556	15.3	14.5-16.1	871	5.2	4.9-5.6	457	6.9	6.3-7.6	414	4.1	3.7-4.5
1983	3,375	19.3	18.6-20.0	1,856	26.1	24.9-27.4	1,519	14.7	14.0-15.5	865	5.1	4.8-5.5	459	6.9	6.3-7.6	406	4.0	3.6-4.4
1984	3,276	18.8	18.1-19.4	1,776	24.9	23.7-26.1	1,500	14.6	13.9-15.4	846	5.0	4.6-5.3	455	6.8	6.2-7.5	391	3.8	3.4-4.2
1985	3,309	18.7	18.1-19.4	1,841	25.6	24.4-26.9	1,468	14.1	13.4-14.9	780	4.5	4.2-4.9	415	6.2	5.6-6.9	365	3.5	3.1-3.8
1986	3,222	18.2	17.6-18.9	1,785	24.9	23.7-26.1	1,437	13.8	13.1-14.5	773	4.5	4.2-4.8	419	6.3	5.7-6.9	354	3.4	3.0-3.8
1987	3,183	18.0	17.4-18.6	1,727	24.0	22.9-25.2	1,456	14.0	13.3-14.8	758	4.4	4.0-4.7	373	5.6	5.0-6.2	385	3.6	3.2-4.0
1988	3,040	17.1	16.4-17.7	1,656	22.9	21.8-24.1	1,384	13.1	12.4-13.9	696	4.0	3.7-4.3	365	5.4	4.9-6.0	331	3.1	2.8-3.5
1989	3,052	17.1	16.5-17.7	1,638	22.3	21.2-23.4	1,414	13.4	12.7-14.2	654	3.7	3.4-4.0	353	5.1	4.6-5.7	301	2.8	2.5-3.1
1990	3,005	16.7	16.1-17.3	1,601	21.5	20.5-22.7	1,404	13.1	12.4-13.8	611	3.5	3.2-3.7	310	4.4	3.9-5.0	301	2.8	2.5-3.1
1991	2,923	16.3	15.7-16.9	1,571	21.5	20.4-22.6	1,352	12.8	12.1-13.5	693	3.9	3.6-4.2	356	5.1	4.6-5.7	337	3.1	2.8-3.5
1992	2,920	16.1	15.5-16.7	1,586	21.4	20.3-22.5	1,334	12.5	11.8-13.2	657	3.7	3.4-4.0	350	4.9	4.4-5.5	307	2.8	2.5-3.1
1993	2,897	15.8	15.3-16.4	1,575	21.0	19.9-22.0	1,322	12.3	11.6-12.9	631	3.4	3.2-3.7	326	4.5	4.0-5.0	305	2.7	2.4-3.1
1994	3,014	16.4	15.8-17.0	1,588	21.1	20.0-22.1	1,426	13.2	12.5-13.9	630	3.5	3.2-3.8	329	4.6	4.1-5.1	301	2.7	2.4-3.0
1995	2,933	15.9	15.3-16.5	1,551	20.4	19.4-21.5	1,382	12.8	12.1-13.5	673	3.7	3.4-4.0	346	4.7	4.2-5.3	327	3.0	2.6-3.3
1996	2,986	16.2	15.6-16.8	1,643	21.3	20.3-22.4	1,343	12.4	11.8-13.1	630	3.4	3.1-3.7	323	4.4	3.9-4.9	307	2.8	2.5-3.1
1997	2,924	15.7	15.1-16.2	1,591	20.4	19.4-21.5	1,333	12.2	11.5-12.8	592	3.2	2.9-3.4	331	4.5	4.0-5.0	261	2.3	2.0-2.6
1998	3,158	16.7	16.2-17.3	1,662	20.9	19.9-22.0	1,496	13.6	12.9-14.3	581	3.1	2.8-3.3	295	3.8	3.4-4.3	286	2.5	2.2-2.8
1999	3,216	16.8	16.2-17.4	1,735	21.6	20.6-22.7	1,481	13.4	12.7-14.1	632	3.3	3.0-3.6	334	4.3	3.9-4.8	298	2.6	2.3-2.9
2000	3,160	16.3	15.7-16.9	1,752	21.4	20.4-22.4	1,408	12.5	11.9-13.2	609	3.1	2.9-3.4	323	4.1	3.6-4.5	286	2.4	2.1-2.7
2001	3,093	15.8	15.2-16.3	1,660	20.1	19.1-21.1	1,433	12.7	12.1-13.4	594	3.0	2.8-3.3	318	4.0	3.6-4.5	276	2.3	2.1-2.6
2002	3,034	15.3	14.7-15.8	1,663	19.5	18.6-20.5	1,371	11.9	11.3-12.6	640	3.2	3.0-3.5	331	4.1	3.6-4.5	309	2.6	2.3-2.9
2003	3,011	15.0	14.5-15.5	1,626	18.9	18.0-19.9	1,385	12.0	11.4-12.7	572	2.8	2.6-3.1	315	3.8	3.4-4.3	257	2.2	1.9-2.4
2004	2,994	14.7	14.2-15.3	1,577	18.1	17.2-19.0	1,417	12.3	11.6-12.9	585	2.9	2.6-3.1	316	3.8	3.3-4.2	269	2.2	1.9-2.5
2005	2,762	13.5	13.0-14.0	1,516	17.3	16.4-18.2	1,246	10.6	10.1-11.3	552	2.7	2.5-2.9	299	3.5	3.1-4.0	253	2.0	1.8-2.3
2006	2,747	13.3	12.8-13.8	1,461	16.4	15.5-17.2	1,286	11.0	10.4-11.7	532	2.6	2.4-2.8	274	3.2	2.9-3.6	258	2.1	1.9-2.4
2007	2,809	13.4	12.9-13.9	1,510	16.5	15.7-17.4	1,299	11.1	10.5-11.7	525	2.5	2.3-2.7	283	3.2	2.9-3.6	242	2.0	1.7-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Rectum & rectosigmoid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	2,742	13.0	12.5-13.5	1,504	16.3	15.4-17.1	1,238	10.4	9.8-11.0	588	2.7	2.5-3.0	311	3.5	3.1-3.9	277	2.2	1.9-2.4
2009	2,698	12.6	12.1-13.1	1,504	15.9	15.1-16.7	1,194	10.0	9.4-10.6	537	2.5	2.3-2.7	282	3.1	2.7-3.5	255	2.0	1.8-2.3
2010	2,679	12.3	11.8-12.8	1,475	15.3	14.5-16.2	1,204	10.0	9.4-10.6	548	2.5	2.3-2.7	312	3.3	3.0-3.7	236	1.9	1.6-2.1
2011	2,633	12.0	11.5-12.5	1,445	14.8	14.0-15.6	1,188	9.8	9.2-10.4	556	2.5	2.3-2.7	308	3.2	2.9-3.6	248	1.9	1.7-2.2
2012	2,479	11.1	10.7-11.5	1,443	14.4	13.7-15.2	1,036	8.5	7.9-9.0	541	2.4	2.2-2.6	306	3.2	2.8-3.6	235	1.8	1.6-2.0
2013	2,689	11.9	11.4-12.4	1,505	14.8	14.0-15.6	1,184	9.6	9.0-10.1	540	2.3	2.1-2.5	317	3.2	2.8-3.5	223	1.7	1.4-1.9
2014	2,627	11.4	11.0-11.9	1,500	14.4	13.6-15.1	1,127	9.0	8.4-9.5	574	2.5	2.3-2.7	324	3.2	2.9-3.6	250	1.9	1.6-2.1
2015	2,682	11.5	11.1-12.0	1,523	14.3	13.6-15.1	1,159	9.2	8.7-9.8	551	2.3	2.1-2.5	308	3.1	2.7-3.4	243	1.8	1.6-2.0
2016	2,609	11.1	10.7-11.6	1,507	14.2	13.4-14.9	1,102	8.7	8.1-9.2	607	2.5	2.3-2.7	339	3.3	2.9-3.7	268	1.9	1.7-2.2
2017	2,676	11.3	10.9-11.8	1,540	14.2	13.5-15.0	1,136	9.0	8.5-9.5	618	2.5	2.3-2.7	343	3.2	2.9-3.6	275	2.0	1.8-2.2
2018	2,620	11.0	10.6-11.5	1,494	13.8	13.0-14.5	1,126	8.7	8.2-9.3	569	2.3	2.1-2.5	323	3.0	2.7-3.4	246	1.8	1.6-2.0
2019	2,633	11.1	10.6-11.5	1,523	13.9	13.2-14.7	1,110	8.7	8.2-9.3	582	2.3	2.1-2.5	341	3.1	2.8-3.4	241	1.7	1.5-1.9
2020	2,363	9.9	9.5-10.3	1,395	12.6	11.9-13.3	968	7.6	7.1-8.1	629	2.5	2.3-2.7	360	3.3	2.9-3.6	269	1.9	1.6-2.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Liver / intrahepatic bile duct**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	561	3.4	3.1-3.7	335	4.8	4.3-5.3	226	2.3	2.0-2.7	515	3.1	2.8-3.4	308	4.4	3.9-4.9	207	2.2	1.9-2.5
1977	592	3.4	3.2-3.7	390	5.4	4.8-6.0	202	2.0	1.8-2.4	605	3.5	3.3-3.8	358	4.9	4.4-5.5	247	2.5	2.2-2.9
1978	593	3.5	3.2-3.8	354	5.0	4.5-5.6	239	2.4	2.1-2.7	577	3.4	3.1-3.7	336	4.8	4.3-5.3	241	2.5	2.2-2.8
1979	545	3.3	3.0-3.5	311	4.4	3.9-4.9	234	2.4	2.1-2.8	560	3.4	3.1-3.7	305	4.5	4.0-5.0	255	2.6	2.3-3.0
1980	567	3.4	3.1-3.6	340	4.9	4.4-5.5	227	2.3	2.0-2.6	538	3.2	2.9-3.4	295	4.3	3.8-4.8	243	2.4	2.1-2.8
1981	595	3.5	3.2-3.8	356	5.1	4.5-5.7	239	2.4	2.1-2.7	614	3.6	3.3-3.9	349	5.2	4.6-5.7	265	2.6	2.3-3.0
1982	565	3.3	3.0-3.6	379	5.4	4.8-6.0	186	1.8	1.6-2.1	582	3.4	3.1-3.7	364	5.3	4.7-5.9	218	2.1	1.9-2.5
1983	604	3.5	3.2-3.8	394	5.5	4.9-6.1	210	2.1	1.8-2.4	637	3.7	3.4-4.0	391	5.5	4.9-6.1	246	2.4	2.1-2.8
1984	613	3.5	3.2-3.8	375	5.2	4.7-5.8	238	2.3	2.0-2.6	631	3.6	3.3-3.9	365	5.2	4.6-5.8	266	2.6	2.3-2.9
1985	605	3.5	3.2-3.8	379	5.4	4.9-6.0	226	2.2	1.9-2.5	647	3.8	3.5-4.1	389	5.6	5.1-6.2	258	2.5	2.2-2.9
1986	646	3.7	3.4-4.0	421	5.9	5.3-6.5	225	2.2	1.9-2.5	671	3.9	3.6-4.2	398	5.6	5.1-6.2	273	2.7	2.3-3.0
1987	626	3.5	3.3-3.8	389	5.3	4.8-5.9	237	2.3	2.0-2.6	706	4.1	3.8-4.4	411	5.9	5.3-6.5	295	2.9	2.5-3.2
1988	712	4.1	3.8-4.4	456	6.3	5.7-6.9	256	2.5	2.2-2.8	718	4.1	3.8-4.4	442	6.1	5.5-6.7	276	2.6	2.3-3.0
1989	669	3.8	3.5-4.1	426	5.7	5.2-6.3	243	2.3	2.0-2.6	647	3.7	3.4-3.9	390	5.3	4.8-5.9	257	2.5	2.2-2.8
1990	791	4.4	4.1-4.8	504	6.8	6.2-7.5	287	2.7	2.4-3.0	769	4.3	4.0-4.6	462	6.5	5.9-7.1	307	2.9	2.5-3.2
1991	824	4.6	4.3-4.9	534	7.0	6.4-7.7	290	2.7	2.4-3.1	804	4.4	4.1-4.7	490	6.6	6.0-7.2	314	2.9	2.6-3.2
1992	872	4.9	4.5-5.2	531	7.2	6.6-7.8	341	3.2	2.9-3.6	835	4.6	4.3-5.0	504	6.9	6.3-7.5	331	3.1	2.8-3.5
1993	873	4.8	4.5-5.1	548	7.2	6.6-7.9	325	3.0	2.7-3.3	877	4.8	4.5-5.1	511	6.9	6.3-7.5	366	3.3	3.0-3.7
1994	899	4.9	4.6-5.3	584	7.6	7.0-8.3	315	2.9	2.6-3.3	789	4.3	4.0-4.6	490	6.6	6.0-7.2	299	2.8	2.4-3.1
1995	993	5.4	5.1-5.8	643	8.4	7.7-9.0	350	3.2	2.9-3.6	864	4.7	4.4-5.0	535	7.1	6.5-7.7	329	3.0	2.7-3.3
1996	1,003	5.4	5.1-5.8	670	8.6	8.0-9.3	333	3.1	2.7-3.4	861	4.6	4.3-5.0	532	6.9	6.3-7.6	329	3.0	2.6-3.3
1997	1,100	5.9	5.6-6.3	717	9.1	8.4-9.7	383	3.5	3.1-3.8	845	4.6	4.3-4.9	531	6.8	6.3-7.4	314	2.8	2.5-3.1
1998	1,215	6.5	6.1-6.8	818	10.1	9.4-10.8	397	3.6	3.3-4.0	892	4.7	4.4-5.1	562	7.0	6.5-7.6	330	3.0	2.6-3.3
1999	1,180	6.2	5.8-6.6	774	9.4	8.7-10.1	406	3.6	3.3-4.0	879	4.6	4.3-4.9	541	6.7	6.1-7.3	338	3.0	2.6-3.3
2000	1,203	6.2	5.9-6.6	806	9.6	9.0-10.3	397	3.5	3.2-3.9	941	4.9	4.5-5.2	592	7.2	6.6-7.8	349	3.0	2.7-3.4
2001	1,240	6.3	6.0-6.7	882	10.4	9.7-11.1	358	3.2	2.8-3.5	929	4.7	4.4-5.0	600	7.2	6.6-7.8	329	2.8	2.5-3.2
2002	1,267	6.4	6.0-6.7	901	10.4	9.7-11.1	366	3.2	2.9-3.6	998	5.0	4.7-5.3	654	7.8	7.2-8.4	344	2.9	2.6-3.2
2003	1,360	6.8	6.4-7.2	910	10.3	9.6-11.0	450	3.9	3.6-4.3	966	4.8	4.5-5.1	625	7.3	6.7-7.9	341	2.9	2.6-3.2
2004	1,467	7.2	6.9-7.6	1,051	11.7	11.0-12.4	416	3.6	3.3-4.0	1,082	5.3	5.0-5.7	681	7.7	7.1-8.3	401	3.4	3.1-3.7
2005	1,553	7.6	7.2-8.0	1,128	12.3	11.6-13.1	425	3.7	3.3-4.1	1,105	5.4	5.1-5.7	735	8.3	7.7-8.9	370	3.1	2.8-3.4
2006	1,573	7.6	7.2-7.9	1,130	12.2	11.5-12.9	443	3.8	3.4-4.2	1,145	5.5	5.2-5.9	738	8.3	7.7-8.9	407	3.4	3.1-3.8
2007	1,735	8.2	7.8-8.6	1,249	13.2	12.4-13.9	486	4.1	3.8-4.5	1,167	5.5	5.2-5.9	787	8.6	8.0-9.2	380	3.1	2.8-3.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Liver / intrahepatic bile duct**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	1,777	8.3	7.9-8.7	1,291	13.3	12.6-14.0	486	4.1	3.7-4.5	1,200	5.6	5.3-5.9	814	8.7	8.1-9.3	386	3.1	2.8-3.4
2009	1,829	8.3	8.0-8.7	1,304	13.2	12.5-13.9	525	4.3	3.9-4.7	1,251	5.8	5.5-6.1	843	8.9	8.3-9.5	408	3.3	3.0-3.7
2010	1,821	8.2	7.8-8.6	1,338	13.3	12.6-14.0	483	4.0	3.6-4.3	1,285	5.8	5.5-6.1	855	8.8	8.2-9.4	430	3.4	3.1-3.7
2011	1,912	8.4	8.0-8.8	1,357	13.1	12.4-13.8	555	4.5	4.1-4.9	1,308	5.8	5.5-6.1	839	8.3	7.8-8.9	469	3.7	3.3-4.0
2012	2,059	8.8	8.5-9.2	1,487	14.0	13.3-14.8	572	4.5	4.2-4.9	1,431	6.2	5.9-6.6	951	9.3	8.7-9.9	480	3.7	3.4-4.1
2013	2,128	9.0	8.6-9.4	1,492	13.9	13.1-14.6	636	4.9	4.5-5.3	1,450	6.2	5.8-6.5	981	9.5	8.9-10.1	469	3.5	3.2-3.9
2014	2,221	9.2	8.8-9.6	1,580	14.3	13.6-15.0	641	4.9	4.5-5.3	1,411	5.9	5.6-6.2	953	8.8	8.3-9.4	458	3.5	3.2-3.8
2015	2,214	9.0	8.7-9.4	1,561	14.0	13.3-14.7	653	4.9	4.5-5.3	1,462	6.0	5.7-6.3	966	9.0	8.4-9.6	496	3.6	3.3-4.0
2016	2,360	9.4	9.0-9.8	1,709	14.9	14.2-15.7	651	4.8	4.5-5.2	1,505	6.1	5.8-6.4	1,020	9.2	8.6-9.8	485	3.5	3.2-3.8
2017	2,324	9.2	8.9-9.6	1,661	14.5	13.8-15.3	663	4.8	4.4-5.2	1,404	5.7	5.4-6.0	962	8.7	8.1-9.3	442	3.2	2.9-3.6
2018	2,141	8.4	8.0-8.8	1,503	13.0	12.3-13.7	638	4.6	4.3-5.0	1,417	5.5	5.2-5.8	937	8.2	7.7-8.8	480	3.4	3.1-3.7
2019	2,208	8.6	8.2-8.9	1,525	13.0	12.4-13.7	683	4.9	4.5-5.3	1,400	5.4	5.1-5.7	921	8.0	7.5-8.6	479	3.3	3.0-3.7
2020	1,987	7.6	7.3-8.0	1,338	11.4	10.8-12.0	649	4.5	4.2-4.9	1,343	5.1	4.8-5.4	865	7.4	6.9-8.0	478	3.3	3.0-3.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Pancreas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	2,105	12.7	12.2-13.3	1,080	15.8	14.8-16.8	1,025	10.6	10.0-11.3	1,979	12.0	11.5-12.6	994	14.7	13.7-15.7	985	10.2	9.6-10.9
1977	2,148	13.1	12.5-13.7	1,093	16.5	15.5-17.5	1,055	10.8	10.2-11.5	2,048	12.5	11.9-13.0	1,038	15.5	14.5-16.5	1,010	10.4	9.8-11.1
1978	2,138	12.8	12.2-13.3	1,078	15.5	14.6-16.5	1,060	10.8	10.1-11.4	2,000	11.9	11.3-12.4	1,008	14.7	13.7-15.6	992	10.0	9.4-10.7
1979	2,221	13.2	12.7-13.8	1,133	16.5	15.5-17.5	1,088	10.9	10.3-11.6	2,109	12.6	12.0-13.1	1,061	15.6	14.6-16.6	1,048	10.5	9.9-11.1
1980	2,197	13.0	12.4-13.5	1,091	15.9	14.9-16.9	1,106	11.0	10.3-11.6	2,016	11.9	11.4-12.5	1,024	15.0	14.1-16.0	992	9.8	9.2-10.5
1981	2,139	12.6	12.1-13.2	1,083	15.7	14.8-16.8	1,056	10.5	9.9-11.2	2,019	11.9	11.4-12.5	1,002	14.6	13.7-15.6	1,017	10.0	9.4-10.7
1982	2,182	12.8	12.2-13.3	1,021	15.1	14.1-16.1	1,161	11.3	10.6-12.0	1,943	11.4	10.9-11.9	947	14.0	13.0-14.9	996	9.7	9.1-10.3
1983	2,113	12.2	11.7-12.8	1,028	14.6	13.7-15.6	1,085	10.5	9.9-11.2	2,047	11.8	11.3-12.3	1,009	14.2	13.3-15.2	1,038	10.1	9.5-10.7
1984	2,181	12.6	12.0-13.1	1,031	15.0	14.1-16.0	1,150	11.0	10.4-11.7	2,016	11.6	11.1-12.2	959	13.9	13.0-14.9	1,057	10.1	9.5-10.8
1985	2,206	12.6	12.1-13.1	1,037	14.5	13.6-15.5	1,169	11.1	10.5-11.8	1,955	11.2	10.7-11.7	930	13.4	12.5-14.3	1,025	9.8	9.2-10.4
1986	2,181	12.5	12.0-13.1	1,025	14.6	13.7-15.5	1,156	11.0	10.4-11.7	2,005	11.4	10.9-12.0	946	13.3	12.4-14.2	1,059	10.0	9.4-10.6
1987	2,201	12.5	12.0-13.0	1,013	14.4	13.5-15.4	1,188	11.1	10.5-11.8	1,988	11.3	10.8-11.8	946	13.5	12.6-14.4	1,042	9.8	9.2-10.4
1988	2,239	12.7	12.2-13.3	1,044	14.8	13.9-15.8	1,195	11.2	10.5-11.8	1,996	11.3	10.8-11.8	960	13.7	12.8-14.7	1,036	9.6	9.0-10.2
1989	2,121	11.9	11.4-12.5	999	14.1	13.2-15.0	1,122	10.5	9.9-11.2	1,961	11.1	10.6-11.6	936	13.3	12.4-14.2	1,025	9.6	9.0-10.2
1990	2,226	12.5	12.0-13.0	1,029	14.4	13.5-15.4	1,197	11.2	10.5-11.8	1,998	11.2	10.7-11.7	915	13.0	12.1-13.9	1,083	10.1	9.5-10.7
1991	2,256	12.6	12.1-13.1	1,059	14.6	13.7-15.5	1,197	11.1	10.4-11.7	1,975	11.0	10.6-11.5	928	12.9	12.1-13.8	1,047	9.7	9.1-10.3
1992	2,243	12.5	12.0-13.0	1,044	14.6	13.7-15.5	1,199	11.0	10.4-11.7	1,966	10.9	10.4-11.4	887	12.5	11.6-13.3	1,079	9.8	9.2-10.4
1993	2,251	12.4	11.9-12.9	1,049	14.2	13.4-15.1	1,202	11.0	10.4-11.6	2,081	11.4	11.0-11.9	980	13.5	12.6-14.4	1,101	10.0	9.4-10.6
1994	2,403	13.2	12.6-13.7	1,132	15.3	14.4-16.2	1,271	11.5	10.9-12.2	2,159	11.8	11.3-12.3	999	13.6	12.8-14.5	1,160	10.4	9.8-11.1
1995	2,279	12.4	11.9-12.9	1,020	13.6	12.8-14.5	1,259	11.4	10.7-12.0	2,072	11.2	10.8-11.7	947	12.6	11.8-13.5	1,125	10.1	9.5-10.7
1996	2,288	12.4	11.9-12.9	1,093	14.6	13.7-15.5	1,195	10.8	10.2-11.5	2,101	11.4	10.9-11.9	965	13.1	12.3-14.0	1,136	10.2	9.6-10.8
1997	2,333	12.5	12.0-13.0	1,095	14.3	13.4-15.1	1,238	11.1	10.5-11.7	2,087	11.2	10.7-11.7	995	13.3	12.5-14.2	1,092	9.7	9.1-10.2
1998	2,439	12.9	12.4-13.5	1,148	14.9	14.0-15.7	1,291	11.5	10.9-12.2	2,093	11.1	10.6-11.6	988	13.0	12.2-13.9	1,105	9.7	9.2-10.3
1999	2,454	12.8	12.3-13.3	1,145	14.5	13.7-15.4	1,309	11.5	10.9-12.1	2,202	11.5	11.0-12.0	1,056	13.5	12.7-14.3	1,146	10.0	9.4-10.6
2000	2,516	12.9	12.4-13.5	1,135	14.2	13.4-15.1	1,381	12.0	11.3-12.6	2,134	11.0	10.5-11.4	967	12.3	11.5-13.1	1,167	10.0	9.4-10.5
2001	2,502	12.7	12.2-13.2	1,112	13.7	12.9-14.5	1,390	11.9	11.3-12.6	2,206	11.2	10.7-11.7	981	12.4	11.6-13.2	1,225	10.4	9.8-11.0
2002	2,533	12.7	12.2-13.2	1,178	14.2	13.4-15.1	1,355	11.5	10.9-12.1	2,182	11.0	10.5-11.4	1,013	12.3	11.6-13.1	1,169	9.8	9.3-10.4
2003	2,611	13.0	12.5-13.5	1,249	15.0	14.2-15.9	1,362	11.4	10.8-12.1	2,168	10.8	10.3-11.2	1,007	12.2	11.4-13.0	1,161	9.6	9.1-10.2
2004	2,600	12.8	12.3-13.3	1,210	14.3	13.5-15.1	1,390	11.6	11.0-12.3	2,247	11.0	10.6-11.5	1,086	12.9	12.2-13.7	1,161	9.6	9.1-10.2
2005	2,609	12.7	12.2-13.2	1,251	14.5	13.7-15.3	1,358	11.3	10.7-11.9	2,249	10.9	10.5-11.4	1,063	12.4	11.7-13.2	1,186	9.7	9.2-10.3
2006	2,732	13.2	12.7-13.7	1,278	14.7	13.9-15.5	1,454	12.0	11.4-12.7	2,246	10.8	10.4-11.3	1,064	12.3	11.6-13.1	1,182	9.7	9.1-10.2
2007	2,806	13.4	12.9-13.9	1,354	15.3	14.5-16.1	1,452	11.8	11.2-12.5	2,289	10.9	10.4-11.3	1,097	12.5	11.8-13.3	1,192	9.7	9.1-10.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Pancreas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	2,927	13.8	13.3-14.3	1,399	15.6	14.8-16.4	1,528	12.4	11.7-13.0	2,392	11.3	10.8-11.7	1,114	12.6	11.8-13.3	1,278	10.2	9.6-10.7
2009	2,950	13.7	13.2-14.2	1,500	16.4	15.5-17.2	1,450	11.6	11.0-12.2	2,427	11.2	10.8-11.7	1,211	13.4	12.6-14.1	1,216	9.6	9.1-10.2
2010	3,016	13.8	13.3-14.4	1,454	15.6	14.8-16.4	1,562	12.5	11.8-13.1	2,429	11.0	10.6-11.5	1,176	12.6	11.9-13.3	1,253	9.7	9.2-10.3
2011	3,114	14.0	13.5-14.5	1,532	16.0	15.2-16.8	1,582	12.4	11.8-13.0	2,474	11.1	10.7-11.6	1,192	12.7	12.0-13.5	1,282	9.9	9.3-10.4
2012	3,147	13.9	13.4-14.4	1,551	16.0	15.2-16.8	1,596	12.4	11.8-13.0	2,568	11.3	10.9-11.8	1,271	13.1	12.4-13.9	1,297	10.0	9.4-10.5
2013	3,227	13.9	13.4-14.4	1,592	15.8	15.0-16.6	1,635	12.3	11.7-13.0	2,542	11.0	10.6-11.4	1,253	12.7	12.0-13.5	1,289	9.6	9.1-10.2
2014	3,320	14.2	13.7-14.7	1,666	16.2	15.4-17.0	1,654	12.5	11.9-13.1	2,597	11.0	10.6-11.4	1,291	12.8	12.1-13.5	1,306	9.6	9.1-10.1
2015	3,358	14.0	13.6-14.5	1,651	15.7	15.0-16.5	1,707	12.6	12.0-13.3	2,521	10.5	10.1-10.9	1,244	12.0	11.3-12.7	1,277	9.2	8.7-9.7
2016	3,464	14.4	13.9-14.9	1,738	16.4	15.6-17.2	1,726	12.7	12.1-13.4	2,675	11.0	10.6-11.4	1,328	12.6	11.9-13.3	1,347	9.7	9.2-10.3
2017	3,651	14.8	14.4-15.3	1,856	17.2	16.4-18.0	1,795	13.0	12.4-13.6	2,727	11.0	10.6-11.4	1,346	12.6	11.9-13.3	1,381	9.7	9.2-10.3
2018	3,616	14.4	14.0-14.9	1,785	16.1	15.4-16.9	1,831	13.0	12.4-13.6	2,688	10.6	10.2-11.0	1,327	12.1	11.4-12.8	1,361	9.4	8.9-9.9
2019	3,702	14.6	14.1-15.1	1,836	16.5	15.7-17.3	1,866	13.0	12.4-13.7	2,772	10.8	10.4-11.2	1,390	12.4	11.7-13.1	1,382	9.5	9.0-10.0
2020	3,583	13.9	13.4-14.4	1,832	15.9	15.2-16.7	1,751	12.2	11.6-12.8	2,721	10.5	10.1-10.9	1,361	12.0	11.4-12.7	1,360	9.3	8.8-9.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State

Larynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	985	5.5	5.2-5.9	825	10.9	10.1-11.7	160	1.6	1.4-1.9	385	2.2	2.0-2.5	312	4.4	3.9-4.9	73	0.8	0.6-1.0
1977	970	5.5	5.2-5.9	804	10.7	10.0-11.5	166	1.7	1.4-2.0	367	2.1	1.9-2.3	309	4.3	3.8-4.8	58	0.6	0.5-0.8
1978	998	5.7	5.3-6.1	818	11.0	10.3-11.9	180	1.8	1.6-2.1	395	2.3	2.0-2.5	318	4.4	3.9-4.9	77	0.8	0.6-1.0
1979	1,087	6.1	5.7-6.5	866	11.3	10.6-12.1	221	2.2	1.9-2.6	408	2.3	2.1-2.5	333	4.6	4.1-5.1	75	0.7	0.6-0.9
1980	1,092	6.2	5.8-6.6	891	11.9	11.1-12.7	201	2.1	1.8-2.4	374	2.1	1.9-2.4	299	4.2	3.7-4.7	75	0.8	0.6-1.0
1981	1,108	6.3	5.9-6.7	898	12.0	11.2-12.8	210	2.1	1.8-2.4	375	2.1	1.9-2.3	301	4.1	3.7-4.7	74	0.7	0.6-0.9
1982	1,029	5.8	5.5-6.2	835	11.1	10.4-11.9	194	1.9	1.7-2.3	374	2.1	1.9-2.3	299	4.0	3.6-4.5	75	0.7	0.6-0.9
1983	1,032	5.8	5.5-6.2	821	10.7	10.0-11.5	211	2.2	1.9-2.5	358	2.0	1.8-2.3	277	3.8	3.4-4.3	81	0.8	0.6-1.0
1984	1,088	6.1	5.8-6.5	846	11.1	10.4-11.9	242	2.5	2.2-2.8	348	1.9	1.7-2.2	275	3.7	3.3-4.2	73	0.7	0.6-0.9
1985	1,131	6.3	6.0-6.7	887	11.5	10.8-12.3	244	2.5	2.2-2.8	368	2.0	1.8-2.2	291	3.9	3.4-4.4	77	0.7	0.6-0.9
1986	1,068	6.0	5.6-6.4	828	11.0	10.2-11.8	240	2.4	2.1-2.8	339	1.9	1.7-2.1	270	3.7	3.2-4.2	69	0.7	0.5-0.9
1987	1,064	6.0	5.6-6.4	847	11.2	10.5-12.0	217	2.2	1.9-2.5	361	2.0	1.8-2.2	279	3.8	3.4-4.3	82	0.8	0.6-1.0
1988	1,051	5.9	5.5-6.2	799	10.3	9.6-11.1	252	2.6	2.2-2.9	368	2.1	1.9-2.3	298	4.1	3.6-4.6	70	0.7	0.5-0.9
1989	1,144	6.4	6.1-6.8	835	10.9	10.2-11.7	309	3.2	2.8-3.6	347	2.0	1.8-2.2	247	3.3	2.9-3.7	100	1.0	0.8-1.2
1990	1,120	6.2	5.8-6.6	864	11.1	10.4-11.9	256	2.6	2.3-2.9	312	1.8	1.6-2.0	256	3.5	3.1-4.0	56	0.5	0.4-0.7
1991	1,011	5.6	5.3-6.0	801	10.4	9.7-11.2	210	2.1	1.8-2.4	335	1.9	1.7-2.1	259	3.4	3.0-3.9	76	0.7	0.6-0.9
1992	1,077	6.0	5.6-6.3	821	10.5	9.8-11.3	256	2.6	2.2-2.9	401	2.2	2.0-2.4	305	4.1	3.7-4.6	96	0.9	0.7-1.1
1993	991	5.4	5.1-5.8	746	9.6	8.9-10.3	245	2.4	2.1-2.7	364	2.0	1.8-2.2	284	3.7	3.3-4.2	80	0.8	0.6-1.0
1994	959	5.3	4.9-5.6	744	9.6	8.9-10.3	215	2.1	1.8-2.4	349	1.9	1.7-2.1	265	3.5	3.1-4.0	84	0.8	0.6-1.0
1995	931	5.1	4.8-5.4	710	8.9	8.3-9.6	221	2.2	1.9-2.5	351	1.9	1.7-2.1	270	3.5	3.1-3.9	81	0.8	0.6-0.9
1996	936	5.1	4.8-5.4	736	9.3	8.6-10.0	200	2.0	1.7-2.2	349	1.9	1.7-2.1	279	3.6	3.2-4.1	70	0.7	0.5-0.8
1997	903	4.9	4.6-5.2	712	8.9	8.2-9.5	191	1.9	1.6-2.1	290	1.6	1.4-1.8	220	2.8	2.5-3.2	70	0.7	0.5-0.8
1998	886	4.7	4.4-5.0	650	8.0	7.4-8.7	236	2.3	2.0-2.6	309	1.6	1.5-1.8	242	3.0	2.6-3.4	67	0.6	0.5-0.8
1999	889	4.7	4.4-5.0	707	8.5	7.9-9.1	182	1.7	1.5-2.0	302	1.6	1.4-1.8	231	2.8	2.5-3.2	71	0.7	0.5-0.8
2000	881	4.6	4.3-4.9	679	8.1	7.5-8.8	202	1.9	1.6-2.2	300	1.5	1.4-1.7	238	2.9	2.6-3.3	62	0.6	0.4-0.7
2001	840	4.3	4.0-4.6	647	7.6	7.0-8.2	193	1.8	1.5-2.0	249	1.3	1.1-1.4	204	2.5	2.2-2.9	45	0.4	0.3-0.5
2002	822	4.1	3.9-4.4	647	7.5	7.0-8.2	175	1.6	1.4-1.9	296	1.5	1.3-1.7	231	2.8	2.4-3.2	65	0.6	0.4-0.7
2003	845	4.2	3.9-4.5	652	7.4	6.9-8.0	193	1.7	1.5-2.0	270	1.3	1.2-1.5	207	2.4	2.1-2.8	63	0.5	0.4-0.7
2004	833	4.1	3.8-4.4	682	7.6	7.0-8.1	151	1.3	1.1-1.6	267	1.3	1.2-1.5	216	2.5	2.2-2.9	51	0.4	0.3-0.6
2005	863	4.2	3.9-4.5	673	7.4	6.9-8.0	190	1.7	1.4-1.9	270	1.3	1.2-1.5	208	2.4	2.1-2.7	62	0.5	0.4-0.7
2006	819	4.0	3.7-4.2	632	7.0	6.4-7.5	187	1.6	1.4-1.9	261	1.3	1.1-1.4	206	2.4	2.0-2.7	55	0.5	0.3-0.6
2007	801	3.8	3.5-4.1	629	6.8	6.3-7.4	172	1.5	1.2-1.7	249	1.2	1.0-1.3	202	2.2	1.9-2.5	47	0.4	0.3-0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State

Larynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	898	4.2	3.9-4.5	692	7.4	6.8-8.0	206	1.8	1.5-2.0	236	1.1	1.0-1.3	183	2.0	1.7-2.3	53	0.4	0.3-0.6
2009	801	3.7	3.4-4.0	619	6.5	6.0-7.0	182	1.5	1.3-1.8	263	1.2	1.1-1.4	196	2.1	1.8-2.5	67	0.5	0.4-0.7
2010	785	3.6	3.3-3.8	622	6.4	5.9-6.9	163	1.4	1.2-1.6	257	1.2	1.0-1.3	198	2.1	1.8-2.5	59	0.5	0.3-0.6
2011	757	3.3	3.1-3.5	607	5.9	5.4-6.4	150	1.2	1.0-1.4	248	1.1	1.0-1.3	185	1.9	1.6-2.2	63	0.5	0.4-0.6
2012	786	3.4	3.2-3.7	621	6.1	5.6-6.6	165	1.3	1.1-1.5	236	1.0	0.9-1.2	188	1.9	1.6-2.2	48	0.4	0.3-0.5
2013	800	3.4	3.2-3.6	643	6.2	5.7-6.7	157	1.2	1.1-1.5	269	1.1	1.0-1.3	218	2.1	1.9-2.5	51	0.4	0.3-0.5
2014	780	3.3	3.1-3.5	612	5.7	5.3-6.2	168	1.3	1.1-1.5	249	1.0	0.9-1.2	202	1.9	1.7-2.2	47	0.4	0.3-0.5
2015	808	3.3	3.1-3.6	647	6.0	5.5-6.5	161	1.2	1.0-1.4	213	0.9	0.8-1.0	163	1.5	1.3-1.8	50	0.4	0.3-0.5
2016	725	3.0	2.8-3.2	582	5.4	4.9-5.8	143	1.1	0.9-1.3	237	1.0	0.9-1.1	193	1.8	1.6-2.1	44	0.3	0.2-0.4
2017	765	3.1	2.9-3.3	585	5.2	4.8-5.7	180	1.4	1.2-1.6	194	0.8	0.6-0.9	152	1.4	1.1-1.6	42	0.3	0.2-0.4
2018	702	2.7	2.5-3.0	551	4.7	4.4-5.2	151	1.1	0.9-1.3	211	0.8	0.7-0.9	161	1.4	1.2-1.7	50	0.4	0.3-0.5
2019	721	2.8	2.6-3.0	574	4.9	4.5-5.4	147	1.1	0.9-1.3	191	0.7	0.6-0.8	155	1.3	1.1-1.6	36	0.2	0.2-0.3
2020	616	2.4	2.2-2.6	493	4.2	3.8-4.6	123	0.9	0.7-1.1	159	0.6	0.5-0.7	127	1.1	0.9-1.3	32	0.2	0.1-0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Lung and bronchus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	9,533	55.1	53.9-56.2	6,841	94.8	92.4-97.2	2,692	27.5	26.5-28.6	7,648	44.5	43.4-45.5	5,597	79.0	76.8-81.2	2,051	20.9	20.0-21.9
1977	9,585	55.1	54.0-56.2	6,779	94.2	91.9-96.6	2,806	28.4	27.4-29.5	7,988	46.2	45.2-47.2	5,818	81.7	79.5-83.9	2,170	22.0	21.1-23.0
1978	9,774	56.3	55.1-57.4	6,980	96.9	94.6-99.3	2,794	28.4	27.4-29.5	8,014	46.3	45.3-47.3	5,763	81.1	78.9-83.3	2,251	22.7	21.8-23.7
1979	10,050	57.7	56.6-58.9	6,933	96.7	94.3-99.1	3,117	31.5	30.4-32.7	8,005	46.1	45.1-47.2	5,655	79.8	77.6-81.9	2,350	23.6	22.6-24.6
1980	10,404	59.4	58.3-60.6	7,213	99.6	97.3-102.1	3,191	32.0	30.9-33.2	8,537	48.9	47.8-49.9	5,988	83.8	81.6-86.1	2,549	25.4	24.4-26.4
1981	10,358	59.0	57.8-60.2	6,991	97.1	94.7-99.5	3,367	33.6	32.4-34.8	8,463	48.3	47.3-49.4	5,900	82.6	80.5-84.9	2,563	25.5	24.5-26.5
1982	10,799	61.3	60.1-62.5	7,155	99.0	96.6-101.4	3,644	36.2	35.0-37.4	8,651	49.3	48.2-50.4	5,824	81.6	79.4-83.8	2,827	28.0	27.0-29.1
1983	11,085	62.4	61.3-63.6	7,243	99.3	97.0-101.7	3,842	37.7	36.5-39.0	8,914	50.4	49.4-51.5	5,882	82.4	80.2-84.6	3,032	29.6	28.5-30.7
1984	11,083	62.4	61.2-63.6	7,217	98.5	96.1-100.8	3,866	38.2	37.0-39.5	8,828	49.7	48.7-50.8	5,863	81.2	79.0-83.4	2,965	28.9	27.8-30.0
1985	11,427	64.0	62.8-65.2	7,326	99.7	97.3-102.1	4,101	40.2	39.0-41.5	9,260	51.9	50.9-53.0	6,004	83.0	80.8-85.2	3,256	31.6	30.5-32.7
1986	11,365	63.7	62.5-64.9	7,140	97.4	95.0-99.7	4,225	41.3	40.0-42.6	9,263	51.8	50.7-52.9	5,875	81.1	79.0-83.3	3,388	32.9	31.8-34.1
1987	11,653	64.8	63.6-66.0	7,274	98.3	96.0-100.6	4,379	42.4	41.2-43.7	9,396	52.5	51.4-53.6	6,006	82.7	80.6-84.9	3,390	32.6	31.5-33.8
1988	11,746	65.3	64.1-66.5	7,172	97.6	95.2-99.9	4,574	44.2	42.9-45.5	9,503	52.9	51.8-53.9	5,991	82.3	80.1-84.5	3,512	33.8	32.6-34.9
1989	11,968	66.4	65.2-67.7	7,218	97.0	94.7-99.4	4,750	46.1	44.8-47.4	9,613	53.4	52.3-54.5	5,968	81.6	79.5-83.8	3,645	35.0	33.8-36.2
1990	12,031	66.5	65.3-67.7	7,251	97.6	95.3-99.9	4,780	45.8	44.5-47.2	9,794	54.2	53.1-55.3	5,955	81.1	79.0-83.2	3,839	36.5	35.4-37.7
1991	12,196	67.5	66.3-68.7	7,134	95.8	93.5-98.1	5,062	48.6	47.3-50.0	10,006	55.3	54.2-56.4	5,999	81.7	79.6-83.8	4,007	38.0	36.9-39.3
1992	12,429	68.3	67.1-69.5	7,272	96.8	94.6-99.1	5,157	49.2	47.9-50.6	9,966	54.8	53.7-55.9	5,860	79.5	77.4-81.6	4,106	38.7	37.5-39.9
1993	12,333	67.4	66.2-68.6	7,023	93.0	90.8-95.2	5,310	50.5	49.1-51.9	9,899	54.1	53.0-55.2	5,787	77.9	75.8-80.0	4,112	38.7	37.5-39.9
1994	12,634	68.7	67.5-70.0	7,090	93.2	91.0-95.4	5,544	52.6	51.2-54.0	9,989	54.3	53.3-55.4	5,951	79.5	77.5-81.6	4,038	37.7	36.5-38.9
1995	12,436	67.3	66.1-68.5	7,033	91.4	89.2-93.6	5,403	50.7	49.4-52.1	9,936	53.8	52.7-54.8	5,717	75.7	73.7-77.7	4,219	39.2	38.0-40.4
1996	12,656	68.3	67.1-69.5	7,097	92.1	89.9-94.3	5,559	52.2	50.8-53.6	9,799	52.8	51.8-53.9	5,628	73.8	71.9-75.8	4,171	38.6	37.5-39.8
1997	12,700	68.0	66.9-69.2	6,982	89.7	87.6-91.9	5,718	53.3	52.0-54.8	9,536	51.1	50.0-52.1	5,412	70.7	68.8-72.6	4,124	37.9	36.8-39.1
1998	12,828	68.0	66.8-69.2	6,955	87.6	85.6-89.7	5,873	54.3	53.0-55.8	9,613	50.9	49.9-52.0	5,401	69.0	67.1-70.9	4,212	38.4	37.3-39.6
1999	12,674	66.3	65.1-67.4	6,816	85.2	83.2-87.3	5,858	53.4	52.0-54.8	9,505	49.7	48.7-50.7	5,232	66.3	64.5-68.2	4,273	38.5	37.4-39.7
2000	13,054	67.4	66.2-68.5	6,982	86.1	84.1-88.2	6,072	54.6	53.2-56.0	9,639	49.7	48.7-50.7	5,314	66.5	64.7-68.3	4,325	38.2	37.1-39.4
2001	12,809	65.4	64.3-66.6	6,728	82.2	80.3-84.2	6,081	54.2	52.8-55.6	9,512	48.5	47.6-49.5	5,206	64.8	63.0-66.6	4,306	37.8	36.7-39.0
2002	13,023	65.7	64.6-66.9	6,905	83.1	81.2-85.1	6,118	54.0	52.6-55.4	9,625	48.5	47.6-49.5	5,193	63.3	61.6-65.1	4,432	38.6	37.4-39.7
2003	13,011	65.0	63.9-66.2	6,890	81.8	79.9-83.8	6,121	53.5	52.2-54.9	9,462	47.2	46.3-48.2	5,106	61.4	59.7-63.1	4,356	37.5	36.4-38.7
2004	13,123	64.9	63.8-66.1	6,812	79.8	77.9-81.8	6,311	54.6	53.3-56.0	9,360	46.2	45.3-47.2	4,969	59.1	57.4-60.7	4,391	37.5	36.4-38.6
2005	13,188	64.8	63.7-65.9	6,871	79.8	77.9-81.7	6,317	54.5	53.1-55.9	9,273	45.6	44.7-46.5	4,943	58.3	56.7-60.0	4,330	36.8	35.7-38.0
2006	13,349	65.3	64.2-66.5	6,728	77.5	75.6-79.4	6,621	57.1	55.7-58.5	9,181	44.8	43.9-45.7	4,912	57.3	55.7-58.9	4,269	36.1	35.1-37.3
2007	13,543	65.4	64.3-66.5	6,929	78.7	76.8-80.5	6,614	56.3	55.0-57.7	9,317	44.8	43.9-45.8	4,883	55.9	54.4-57.5	4,434	37.2	36.1-38.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Lung and bronchus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	13,751	65.5	64.4-66.6	6,932	77.5	75.7-79.4	6,819	57.4	56.0-58.8	9,180	43.5	42.6-44.4	4,811	54.4	52.9-56.0	4,369	36.1	35.0-37.2
2009	13,708	64.4	63.3-65.5	7,037	77.3	75.5-79.2	6,671	55.3	53.9-56.6	8,919	41.8	40.9-42.7	4,697	52.2	50.7-53.7	4,222	34.5	33.5-35.6
2010	13,495	62.4	61.3-63.4	6,777	72.7	70.9-74.5	6,718	55.3	53.9-56.6	9,041	41.8	40.9-42.6	4,675	51.0	49.5-52.5	4,366	35.4	34.3-36.4
2011	13,822	62.7	61.6-63.8	6,928	73.0	71.3-74.8	6,894	55.5	54.2-56.9	8,940	40.5	39.6-41.3	4,630	49.5	48.0-50.9	4,310	34.1	33.0-35.1
2012	13,736	61.1	60.1-62.2	6,842	70.6	68.9-72.3	6,894	54.8	53.4-56.1	9,131	40.7	39.9-41.6	4,838	50.7	49.2-52.2	4,293	33.7	32.7-34.8
2013	13,805	60.4	59.4-61.4	6,824	68.8	67.1-70.5	6,981	54.5	53.2-55.8	8,900	38.7	37.9-39.6	4,605	46.9	45.5-48.3	4,295	32.9	31.9-33.9
2014	13,595	58.1	57.1-59.1	6,736	66.2	64.6-67.8	6,859	52.3	51.0-53.6	8,411	36.0	35.2-36.8	4,396	43.9	42.6-45.3	4,015	30.2	29.3-31.2
2015	14,142	59.4	58.4-60.4	6,974	67.3	65.7-69.0	7,168	53.9	52.6-55.2	8,420	35.3	34.5-36.1	4,351	42.5	41.3-43.9	4,069	30.1	29.1-31.0
2016	14,064	58.0	57.0-59.0	6,880	64.9	63.3-66.5	7,184	53.2	52.0-54.5	7,939	32.7	32.0-33.5	4,097	39.4	38.2-40.6	3,842	27.9	27.0-28.8
2017	14,565	59.0	58.1-60.0	7,070	65.4	63.9-67.0	7,495	54.6	53.3-55.8	7,670	30.9	30.2-31.6	3,916	36.6	35.5-37.8	3,754	26.7	25.9-27.6
2018	14,310	56.9	55.9-57.8	6,874	62.0	60.5-63.5	7,436	53.3	52.1-54.5	7,578	30.0	29.3-30.7	3,889	35.7	34.5-36.8	3,689	25.9	25.0-26.7
2019	14,304	55.6	54.7-56.6	6,814	60.1	58.7-61.6	7,490	52.6	51.4-53.8	7,299	28.2	27.6-28.9	3,694	33.1	32.1-34.2	3,605	24.7	23.9-25.5
2020	12,478	47.8	47.0-48.7	5,978	52.0	50.7-53.4	6,500	44.9	43.8-46.0	6,755	25.8	25.2-26.5	3,471	30.6	29.6-31.7	3,284	22.3	21.6-23.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Melanoma of the skin**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	954	5.7	5.3-6.1	486	6.6	6.0-7.2	468	5.1	4.6-5.6	389	2.4	2.1-2.6	216	3.0	2.6-3.4	173	1.9	1.6-2.2
1977	1,106	6.6	6.2-7.0	564	7.6	7.0-8.3	542	5.9	5.4-6.4	398	2.4	2.2-2.7	236	3.3	2.9-3.8	162	1.8	1.5-2.1
1978	1,043	6.3	5.9-6.7	518	7.1	6.5-7.8	525	5.8	5.3-6.4	341	2.1	1.8-2.3	210	3.0	2.6-3.4	131	1.4	1.2-1.7
1979	1,106	6.7	6.3-7.1	571	7.7	7.1-8.4	535	5.9	5.4-6.4	381	2.3	2.1-2.6	211	2.9	2.5-3.4	170	1.8	1.6-2.1
1980	1,286	7.7	7.3-8.1	680	9.2	8.5-10.0	606	6.6	6.1-7.2	403	2.4	2.2-2.7	231	3.1	2.7-3.6	172	1.8	1.6-2.2
1981	1,239	7.4	7.0-7.8	626	8.4	7.8-9.2	613	6.6	6.1-7.2	394	2.3	2.1-2.6	226	3.1	2.7-3.5	168	1.8	1.5-2.1
1982	1,280	7.6	7.2-8.0	640	8.7	8.0-9.4	640	6.9	6.3-7.4	392	2.3	2.1-2.6	236	3.3	2.8-3.7	156	1.6	1.4-1.9
1983	1,396	8.2	7.8-8.7	741	10.1	9.3-10.8	655	7.1	6.5-7.7	400	2.4	2.1-2.6	230	3.2	2.8-3.6	170	1.8	1.5-2.1
1984	1,335	7.8	7.4-8.2	702	9.3	8.6-10.1	633	6.8	6.2-7.3	425	2.5	2.2-2.7	248	3.3	2.9-3.8	177	1.8	1.6-2.1
1985	1,383	8.0	7.6-8.5	752	10.0	9.2-10.7	631	6.7	6.2-7.3	435	2.5	2.3-2.8	252	3.5	3.1-4.0	183	1.9	1.6-2.2
1986	1,441	8.3	7.9-8.7	753	10.0	9.3-10.8	688	7.3	6.7-7.8	442	2.6	2.3-2.8	250	3.4	3.0-3.8	192	2.0	1.7-2.3
1987	1,366	7.8	7.4-8.3	741	9.8	9.1-10.6	625	6.5	6.0-7.0	450	2.6	2.4-2.9	267	3.6	3.2-4.1	183	1.9	1.6-2.2
1988	1,306	7.4	7.0-7.9	726	9.7	9.0-10.4	580	5.9	5.4-6.4	434	2.5	2.3-2.7	272	3.7	3.3-4.2	162	1.6	1.4-1.9
1989	1,358	7.7	7.3-8.2	742	9.7	9.0-10.5	616	6.3	5.8-6.8	444	2.6	2.3-2.8	278	3.8	3.4-4.3	166	1.7	1.4-1.9
1990	1,413	7.9	7.5-8.3	755	9.8	9.1-10.5	658	6.7	6.2-7.2	422	2.4	2.2-2.6	252	3.4	3.0-3.8	170	1.7	1.5-2.0
1991	1,380	7.7	7.3-8.1	784	10.1	9.4-10.9	596	5.9	5.4-6.4	426	2.4	2.2-2.6	252	3.4	3.0-3.9	174	1.7	1.4-2.0
1992	1,311	7.3	6.9-7.7	739	9.5	8.8-10.2	572	5.6	5.2-6.1	402	2.2	2.0-2.5	235	3.1	2.7-3.5	167	1.6	1.4-1.9
1993	1,357	7.4	7.0-7.8	766	9.8	9.1-10.5	591	5.8	5.3-6.2	400	2.2	2.0-2.4	242	3.3	2.8-3.7	158	1.5	1.3-1.8
1994	1,492	8.1	7.7-8.6	856	10.8	10.1-11.6	636	6.2	5.8-6.8	416	2.3	2.1-2.5	261	3.4	3.0-3.9	155	1.5	1.2-1.7
1995	1,606	8.7	8.3-9.2	938	11.7	11.0-12.5	668	6.6	6.1-7.1	434	2.4	2.2-2.6	249	3.2	2.9-3.7	185	1.7	1.5-2.0
1996	1,697	9.2	8.7-9.6	945	11.8	11.1-12.6	752	7.3	6.8-7.9	448	2.4	2.2-2.7	260	3.3	2.9-3.7	188	1.8	1.5-2.1
1997	1,844	9.9	9.4-10.4	1,047	13.1	12.3-14.0	797	7.7	7.2-8.3	421	2.3	2.0-2.5	268	3.5	3.1-3.9	153	1.4	1.2-1.7
1998	1,946	10.3	9.9-10.8	1,084	13.2	12.5-14.1	862	8.3	7.8-8.9	431	2.3	2.1-2.5	264	3.3	2.9-3.8	167	1.5	1.3-1.8
1999	1,941	10.2	9.7-10.6	1,105	13.4	12.6-14.2	836	7.9	7.4-8.5	401	2.1	1.9-2.3	238	3.0	2.6-3.4	163	1.5	1.3-1.7
2000	2,211	11.4	11.0-11.9	1,235	14.8	14.0-15.7	976	9.2	8.6-9.8	446	2.3	2.1-2.5	278	3.4	3.0-3.9	168	1.5	1.3-1.8
2001	2,471	12.7	12.2-13.2	1,358	16.0	15.1-16.8	1,113	10.4	9.8-11.1	479	2.4	2.2-2.7	296	3.6	3.2-4.1	183	1.6	1.4-1.9
2002	2,589	13.1	12.6-13.7	1,443	16.8	16.0-17.7	1,146	10.7	10.1-11.3	425	2.2	2.0-2.4	222	2.6	2.3-3.0	203	1.8	1.6-2.1
2003	2,439	12.2	11.8-12.7	1,365	15.8	15.0-16.7	1,074	9.8	9.3-10.5	437	2.2	2.0-2.4	251	3.0	2.6-3.4	186	1.6	1.4-1.9
2004	2,957	14.7	14.2-15.3	1,659	18.9	18.0-19.8	1,298	11.9	11.3-12.6	430	2.1	1.9-2.3	253	3.0	2.6-3.4	177	1.5	1.3-1.8
2005	3,236	16.1	15.6-16.7	1,838	20.9	19.9-21.9	1,398	12.8	12.1-13.5	443	2.2	2.0-2.4	265	3.1	2.7-3.5	178	1.5	1.3-1.8
2006	3,600	17.8	17.2-18.4	2,033	22.8	21.8-23.8	1,567	14.3	13.6-15.0	466	2.3	2.1-2.5	296	3.4	3.0-3.8	170	1.5	1.2-1.7
2007	3,307	16.1	15.6-16.7	1,851	20.6	19.6-21.5	1,456	13.2	12.5-13.9	454	2.2	2.0-2.4	308	3.6	3.2-4.0	146	1.2	1.0-1.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Melanoma of the skin**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	3,584	17.3	16.8-17.9	2,033	22.5	21.5-23.5	1,551	13.9	13.2-14.6	459	2.2	2.0-2.4	292	3.2	2.9-3.7	167	1.4	1.2-1.6
2009	3,819	18.2	17.6-18.8	2,190	23.7	22.7-24.8	1,629	14.4	13.7-15.2	460	2.2	2.0-2.4	277	3.1	2.7-3.5	183	1.5	1.3-1.8
2010	3,567	16.8	16.3-17.4	2,011	21.5	20.5-22.5	1,556	13.6	12.9-14.3	467	2.2	2.0-2.4	277	3.0	2.6-3.3	190	1.6	1.3-1.8
2011	4,069	18.8	18.2-19.4	2,320	24.2	23.2-25.2	1,749	15.0	14.3-15.8	470	2.1	1.9-2.3	293	3.1	2.8-3.5	177	1.4	1.2-1.6
2012	3,847	17.5	16.9-18.1	2,161	22.1	21.2-23.1	1,686	14.4	13.7-15.1	536	2.4	2.2-2.6	324	3.4	3.1-3.8	212	1.6	1.4-1.9
2013	4,290	19.3	18.7-19.9	2,443	24.5	23.5-25.5	1,847	15.6	14.9-16.4	503	2.2	2.0-2.4	327	3.4	3.0-3.8	176	1.4	1.2-1.6
2014	4,192	18.6	18.1-19.2	2,399	23.9	22.9-24.9	1,793	15.0	14.3-15.7	450	1.9	1.8-2.1	290	2.9	2.6-3.3	160	1.2	1.0-1.4
2015	4,143	18.0	17.5-18.6	2,408	23.6	22.6-24.6	1,735	14.2	13.5-14.9	378	1.6	1.4-1.8	241	2.4	2.1-2.7	137	1.1	0.9-1.3
2016	4,188	18.0	17.5-18.6	2,442	23.4	22.5-24.4	1,746	14.2	13.6-15.0	417	1.7	1.6-1.9	260	2.5	2.2-2.9	157	1.1	1.0-1.3
2017	4,244	18.2	17.6-18.7	2,498	23.8	22.9-24.8	1,746	14.2	13.5-14.9	394	1.6	1.5-1.8	250	2.4	2.1-2.8	144	1.0	0.9-1.2
2018	4,317	18.0	17.5-18.6	2,472	22.8	21.9-23.8	1,845	14.6	13.9-15.3	391	1.5	1.4-1.7	239	2.2	1.9-2.5	152	1.1	0.9-1.2
2019	4,488	18.6	18.1-19.2	2,584	23.6	22.7-24.6	1,904	15.2	14.5-16.0	356	1.4	1.3-1.6	224	2.1	1.8-2.4	132	0.9	0.8-1.1
2020	3,843	15.8	15.3-16.3	2,154	19.5	18.7-20.4	1,689	13.2	12.5-13.8	381	1.5	1.4-1.7	243	2.3	2.0-2.6	138	1.0	0.8-1.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Female breast**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	9,439	102.0	99.9-104.1	3,677	38.8	37.6-40.1
1977	9,282	100.2	98.1-102.3	3,674	38.8	37.5-40.1
1978	9,458	101.8	99.7-103.9	3,688	38.7	37.4-39.9
1979	9,353	100.2	98.2-102.3	3,503	36.8	35.6-38.1
1980	9,553	102.1	100.0-104.2	3,674	38.1	36.8-39.3
1981	9,700	103.5	101.4-105.6	3,648	37.7	36.4-39.0
1982	9,847	104.1	102.0-106.2	3,596	36.9	35.7-38.1
1983	10,239	107.6	105.4-109.7	3,576	36.5	35.3-37.8
1984	10,683	111.4	109.3-113.6	3,756	38.0	36.8-39.3
1985	11,062	115.3	113.1-117.5	3,707	37.4	36.2-38.7
1986	11,125	115.2	113.0-117.4	3,753	37.6	36.4-38.9
1987	11,508	118.2	116.0-120.4	3,731	37.1	35.9-38.3
1988	11,752	120.1	117.9-122.3	3,842	38.1	36.9-39.4
1989	11,541	117.8	115.6-120.0	3,776	37.5	36.3-38.8
1990	12,321	125.6	123.3-127.8	3,701	36.5	35.3-37.7
1991	12,434	125.6	123.3-127.8	3,592	35.3	34.1-36.4
1992	12,454	125.4	123.2-127.6	3,614	35.2	34.1-36.4
1993	12,266	122.5	120.3-124.7	3,582	34.5	33.3-35.6
1994	12,823	127.4	125.2-129.6	3,564	34.1	33.0-35.3
1995	13,042	128.9	126.7-131.1	3,544	33.8	32.7-34.9
1996	13,841	135.9	133.6-138.2	3,553	33.5	32.4-34.6
1997	13,911	135.2	133.0-137.5	3,309	30.9	29.9-32.0
1998	14,411	138.6	136.3-140.9	3,261	30.1	29.1-31.2
1999	14,213	134.6	132.4-136.9	3,082	27.9	27.0-29.0
2000	14,216	132.7	130.5-134.9	3,088	27.5	26.5-28.5
2001	13,938	128.8	126.7-131.0	3,124	27.6	26.7-28.6
2002	13,939	127.5	125.4-129.7	2,975	25.9	25.0-26.8
2003	14,028	127.2	125.1-129.3	2,988	25.8	24.9-26.8
2004	13,820	124.5	122.4-126.6	2,880	24.6	23.7-25.5
2005	13,922	124.4	122.3-126.5	2,842	24.2	23.3-25.1
2006	14,309	127.8	125.7-129.9	2,714	23.0	22.1-23.8
2007	14,467	127.8	125.7-129.9	2,717	22.7	21.8-23.6
2008	14,860	130.2	128.1-132.4	2,639	21.8	20.9-22.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Female breast**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	15,247	131.7	129.5-133.8	2,654	21.7	20.9-22.6
2010	14,565	124.9	122.8-127.0	2,666	21.3	20.5-22.1
2011	15,508	131.5	129.4-133.7	2,655	21.1	20.3-21.9
2012	15,145	127.2	125.1-129.3	2,607	20.4	19.6-21.2
2013	15,901	132.3	130.2-134.4	2,576	20.0	19.2-20.8
2014	16,250	133.8	131.7-136.0	2,448	18.7	17.9-19.5
2015	16,456	134.6	132.5-136.8	2,483	18.8	18.0-19.6
2016	16,155	130.9	128.8-133.0	2,561	19.3	18.5-20.1
2017	16,796	135.7	133.6-137.9	2,414	17.8	17.0-18.5
2018	17,029	136.6	134.5-138.8	2,576	18.8	18.1-19.6
2019	17,855	142.3	140.1-144.5	2,492	18.2	17.4-18.9
2020	15,606	124.7	122.6-126.7	2,387	17.3	16.5-18.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Cervix uteri**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,317	14.7	13.9-15.6	502	5.4	4.9-5.9
1977	1,203	13.4	12.6-14.2	456	4.9	4.4-5.4
1978	1,192	13.5	12.7-14.3	464	5.0	4.5-5.5
1979	1,215	13.7	12.9-14.5	427	4.7	4.3-5.2
1980	1,192	13.2	12.4-14.0	439	4.7	4.3-5.2
1981	1,035	11.3	10.6-12.0	443	4.7	4.3-5.2
1982	1,071	11.9	11.2-12.7	407	4.3	3.9-4.8
1983	1,090	12.0	11.2-12.7	363	3.8	3.4-4.2
1984	1,131	12.4	11.7-13.2	325	3.4	3.1-3.9
1985	1,046	11.3	10.6-12.0	344	3.6	3.3-4.0
1986	1,013	11.1	10.4-11.8	345	3.7	3.3-4.1
1987	1,120	11.9	11.2-12.7	336	3.5	3.1-3.9
1988	1,076	11.5	10.8-12.3	366	3.8	3.4-4.2
1989	1,045	11.0	10.4-11.7	297	3.0	2.7-3.4
1990	1,192	12.6	11.9-13.4	350	3.6	3.2-4.0
1991	1,155	12.1	11.4-12.9	306	3.1	2.8-3.5
1992	1,107	11.4	10.7-12.1	316	3.2	2.9-3.6
1993	1,105	11.3	10.7-12.0	332	3.4	3.0-3.8
1994	1,157	11.8	11.1-12.5	340	3.4	3.1-3.8
1995	1,238	12.5	11.8-13.2	315	3.1	2.8-3.5
1996	1,247	12.5	11.9-13.3	331	3.3	2.9-3.7
1997	1,115	11.1	10.5-11.8	347	3.4	3.0-3.8
1998	1,054	10.5	9.8-11.1	290	2.8	2.5-3.2
1999	1,036	10.1	9.5-10.8	339	3.2	2.9-3.6
2000	1,025	9.9	9.3-10.6	303	2.9	2.5-3.2
2001	953	9.1	8.6-9.7	280	2.6	2.3-2.9
2002	936	8.9	8.4-9.5	292	2.7	2.4-3.0
2003	948	9.0	8.5-9.6	297	2.7	2.4-3.0
2004	863	8.2	7.6-8.7	267	2.4	2.1-2.7
2005	961	9.1	8.6-9.7	287	2.6	2.3-2.9
2006	904	8.5	8.0-9.1	275	2.5	2.2-2.8
2007	917	8.6	8.1-9.2	295	2.6	2.3-3.0
2008	901	8.5	7.9-9.0	250	2.1	1.9-2.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Cervix uteri**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	896	8.3	7.8-8.9	252	2.2	1.9-2.5
2010	866	8.2	7.6-8.7	249	2.1	1.8-2.4
2011	793	7.3	6.8-7.9	276	2.4	2.1-2.7
2012	849	7.8	7.2-8.3	294	2.5	2.3-2.9
2013	889	8.1	7.6-8.7	286	2.4	2.1-2.7
2014	837	7.6	7.0-8.1	288	2.4	2.1-2.7
2015	866	7.9	7.3-8.4	240	1.9	1.7-2.2
2016	827	7.6	7.1-8.1	260	2.2	1.9-2.4
2017	903	8.2	7.7-8.8	255	2.1	1.8-2.4
2018	798	7.2	6.7-7.8	226	1.8	1.6-2.1
2019	817	7.6	7.0-8.1	216	1.8	1.5-2.1
2020	698	6.3	5.8-6.8	209	1.6	1.4-1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Corpus uteris and uterus, not otherwise specified**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	2,568	26.4	25.4-27.5	606	6.2	5.7-6.7
1977	2,321	23.7	22.7-24.7	552	5.6	5.1-6.1
1978	2,095	21.1	20.2-22.1	514	5.2	4.7-5.6
1979	2,293	23.3	22.3-24.3	526	5.2	4.8-5.7
1980	2,359	24.0	23.0-25.0	574	5.6	5.2-6.1
1981	2,372	24.0	23.0-25.0	529	5.2	4.8-5.7
1982	2,347	23.5	22.6-24.5	527	5.1	4.7-5.6
1983	2,301	23.0	22.1-24.0	522	5.0	4.5-5.4
1984	2,301	23.0	22.0-23.9	532	5.1	4.7-5.5
1985	2,270	22.5	21.5-23.4	500	4.8	4.4-5.2
1986	2,192	21.8	20.9-22.8	484	4.6	4.2-5.0
1987	2,341	23.2	22.2-24.2	486	4.6	4.2-5.0
1988	2,219	21.9	20.9-22.8	491	4.6	4.2-5.0
1989	2,177	21.3	20.4-22.3	496	4.7	4.3-5.1
1990	2,236	22.1	21.2-23.0	461	4.3	3.9-4.7
1991	2,401	24.0	23.1-25.0	502	4.7	4.3-5.1
1992	2,461	24.3	23.4-25.3	522	4.9	4.5-5.3
1993	2,399	23.5	22.6-24.5	546	5.0	4.6-5.4
1994	2,607	25.3	24.4-26.3	536	4.9	4.5-5.4
1995	2,572	25.1	24.1-26.1	539	4.9	4.5-5.3
1996	2,717	26.3	25.3-27.3	538	4.9	4.5-5.3
1997	2,726	26.3	25.3-27.3	541	4.9	4.5-5.4
1998	2,935	28.1	27.1-29.2	550	5.0	4.6-5.4
1999	2,923	27.5	26.5-28.5	521	4.6	4.2-5.1
2000	2,855	26.5	25.5-27.4	534	4.7	4.3-5.1
2001	3,097	28.6	27.6-29.6	538	4.7	4.3-5.1
2002	3,106	28.2	27.2-29.2	544	4.7	4.3-5.2
2003	3,139	28.2	27.2-29.2	599	5.1	4.7-5.6
2004	3,261	29.0	28.0-30.0	558	4.7	4.4-5.2
2005	3,311	29.1	28.1-30.1	595	5.1	4.7-5.5
2006	3,306	28.8	27.8-29.8	626	5.3	4.8-5.7
2007	3,350	28.9	27.9-29.9	609	5.1	4.7-5.5
2008	3,723	31.8	30.7-32.8	626	5.2	4.8-5.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Corpus uteris and uterus, not otherwise specified**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	3,547	29.8	28.9-30.9	605	4.9	4.5-5.4
2010	3,600	29.8	28.9-30.9	665	5.3	4.9-5.7
2011	3,768	30.7	29.7-31.7	696	5.5	5.1-6.0
2012	3,851	30.7	29.7-31.7	703	5.5	5.1-5.9
2013	4,107	32.4	31.4-33.4	740	5.6	5.2-6.0
2014	4,101	31.9	30.9-32.9	773	5.8	5.4-6.2
2015	4,151	31.9	30.9-32.9	754	5.5	5.1-6.0
2016	4,292	32.9	31.9-33.9	830	6.1	5.6-6.5
2017	4,205	31.7	30.8-32.7	808	5.8	5.4-6.2
2018	4,147	31.0	30.0-31.9	835	5.9	5.5-6.3
2019	4,241	31.3	30.4-32.3	893	6.3	5.9-6.8
2020	3,959	28.9	28.0-29.8	847	5.8	5.4-6.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State

Ovary

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,572	16.5	15.7-17.4	1,114	11.5	10.8-12.2
1977	1,507	15.8	15.0-16.7	1,031	10.6	9.9-11.3
1978	1,492	15.6	14.8-16.4	1,009	10.2	9.6-10.9
1979	1,558	16.4	15.6-17.3	985	10.1	9.4-10.7
1980	1,541	16.0	15.2-16.8	1,038	10.5	9.9-11.2
1981	1,634	17.0	16.2-17.9	1,029	10.3	9.7-10.9
1982	1,589	16.5	15.7-17.4	1,010	10.1	9.5-10.7
1983	1,675	17.3	16.4-18.1	1,023	10.1	9.5-10.7
1984	1,598	16.4	15.6-17.3	1,029	10.2	9.6-10.9
1985	1,663	17.1	16.3-18.0	987	9.8	9.2-10.5
1986	1,596	16.4	15.6-17.2	984	9.7	9.1-10.3
1987	1,599	16.3	15.5-17.1	1,021	10.0	9.4-10.6
1988	1,602	16.2	15.4-17.1	1,058	10.2	9.6-10.9
1989	1,602	16.1	15.3-16.9	994	9.6	9.0-10.2
1990	1,686	16.9	16.1-17.7	1,042	10.0	9.4-10.6
1991	1,716	17.2	16.4-18.1	1,095	10.6	9.9-11.2
1992	1,729	17.3	16.5-18.1	1,105	10.7	10.1-11.4
1993	1,678	16.6	15.8-17.4	1,038	9.8	9.2-10.5
1994	1,760	17.2	16.4-18.1	1,022	9.7	9.1-10.3
1995	1,807	17.7	16.9-18.5	1,061	10.0	9.4-10.6
1996	1,736	16.8	16.0-17.7	983	9.2	8.7-9.8
1997	1,742	16.8	16.1-17.7	1,061	9.9	9.3-10.5
1998	1,584	15.2	14.5-16.0	944	8.7	8.2-9.3
1999	1,633	15.4	14.6-16.2	977	8.9	8.3-9.5
2000	1,637	15.1	14.4-15.9	1,027	9.2	8.6-9.7
2001	1,686	15.5	14.7-16.2	1,002	8.8	8.3-9.4
2002	1,659	15.0	14.3-15.8	1,012	8.9	8.3-9.4
2003	1,595	14.3	13.6-15.0	1,038	9.0	8.4-9.5
2004	1,615	14.4	13.7-15.1	1,013	8.7	8.1-9.2
2005	1,513	13.5	12.8-14.2	1,043	8.9	8.4-9.5
2006	1,540	13.5	12.9-14.2	1,023	8.7	8.2-9.2
2007	1,544	13.4	12.8-14.1	950	8.0	7.5-8.5
2008	1,553	13.4	12.7-14.1	985	8.1	7.6-8.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Ovary**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	1,538	13.2	12.5-13.9	940	7.6	7.1-8.1
2010	1,540	13.1	12.4-13.7	938	7.6	7.1-8.1
2011	1,580	13.1	12.5-13.8	958	7.5	7.1-8.1
2012	1,483	12.2	11.6-12.9	954	7.5	7.0-8.0
2013	1,589	13.1	12.4-13.8	978	7.5	7.0-8.0
2014	1,474	12.0	11.3-12.6	892	6.8	6.3-7.2
2015	1,504	12.2	11.6-12.9	897	6.8	6.4-7.3
2016	1,484	11.9	11.2-12.5	891	6.6	6.2-7.1
2017	1,483	11.7	11.1-12.3	866	6.3	5.9-6.8
2018	1,372	11.0	10.4-11.6	866	6.2	5.8-6.6
2019	1,426	11.5	10.8-12.1	876	6.2	5.8-6.7
2020	1,283	10.2	9.6-10.8	831	5.8	5.4-6.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Prostate**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	4,712	77.7	75.4-80.1	1,739	31.1	29.6-32.7
1977	4,795	77.5	75.2-79.9	1,721	30.3	28.8-31.9
1978	4,798	77.7	75.4-80.0	1,804	32.0	30.4-33.6
1979	5,046	81.7	79.4-84.1	1,797	31.8	30.2-33.3
1980	5,624	88.7	86.3-91.2	1,824	32.0	30.5-33.6
1981	5,560	87.7	85.3-90.2	1,836	32.1	30.6-33.7
1982	5,631	88.1	85.7-90.5	1,820	31.3	29.8-32.9
1983	5,866	90.2	87.8-92.6	1,982	33.7	32.2-35.3
1984	5,910	90.1	87.7-92.5	2,054	34.9	33.3-36.5
1985	6,196	92.8	90.4-95.3	1,975	32.9	31.4-34.4
1986	6,328	95.1	92.7-97.6	2,068	33.9	32.4-35.5
1987	6,662	98.5	96.1-101.0	2,076	34.0	32.5-35.5
1988	6,759	100.5	98.0-103.0	2,123	34.7	33.2-36.3
1989	7,026	102.8	100.4-105.4	2,128	34.6	33.0-36.1
1990	8,362	119.2	116.6-121.9	2,263	36.5	35.0-38.1
1991	9,918	138.3	135.5-141.1	2,292	36.6	35.1-38.2
1992	12,528	169.9	166.9-173.0	2,373	37.6	36.1-39.2
1993	12,246	163.1	160.2-166.1	2,363	37.3	35.8-38.9
1994	11,861	156.0	153.1-158.8	2,421	37.9	36.4-39.5
1995	11,601	150.2	147.4-153.0	2,355	36.3	34.8-37.8
1996	12,378	158.9	156.1-161.8	2,286	34.6	33.1-36.1
1997	13,044	166.0	163.2-169.0	2,234	33.4	32.0-34.9
1998	13,057	163.7	160.8-166.5	2,114	30.9	29.6-32.3
1999	13,918	172.2	169.3-175.1	2,032	29.2	27.9-30.5
2000	15,411	188.2	185.2-191.2	2,089	29.7	28.4-31.0
2001	15,128	181.7	178.8-184.6	2,070	28.9	27.7-30.2
2002	15,519	183.3	180.4-186.2	2,017	27.7	26.5-29.0
2003	14,520	169.4	166.7-172.2	1,968	26.5	25.3-27.7
2004	14,253	163.2	160.5-165.9	1,881	25.0	23.9-26.2
2005	14,473	162.9	160.3-165.6	1,766	23.0	21.9-24.1
2006	16,125	179.8	177.0-182.6	1,718	22.1	21.1-23.2
2007	16,680	181.2	178.5-184.0	1,809	22.7	21.6-23.8
2008	16,200	171.7	169.0-174.4	1,726	21.2	20.2-22.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Prostate**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	15,964	164.8	162.2-167.4	1,715	20.7	19.8-21.8
2010	15,186	152.9	150.4-155.4	1,807	21.3	20.3-22.3
2011	16,121	158.5	156.0-161.0	1,738	20.0	19.1-21.0
2012	13,979	133.5	131.2-135.8	1,653	18.7	17.8-19.6
2013	13,883	129.7	127.5-131.9	1,600	17.6	16.8-18.5
2014	13,312	121.1	119.0-123.2	1,721	18.3	17.5-19.2
2015	13,890	123.6	121.5-125.7	1,627	17.0	16.2-17.9
2016	14,424	126.1	124.0-128.3	1,741	17.7	16.9-18.6
2017	15,343	131.2	129.1-133.3	1,788	17.8	17.0-18.7
2018	16,034	135.0	132.9-137.2	1,717	16.6	15.8-17.4
2019	17,030	141.0	138.8-143.2	1,661	15.8	15.0-16.6
2020	14,447	118.0	116.1-120.0	1,625	15.2	14.4-15.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Testis**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	263	3.0	2.6-3.4	52	0.6	0.4-0.8
1977	257	3.0	2.6-3.4	47	0.5	0.4-0.7
1978	275	3.3	2.9-3.7	53	0.6	0.5-0.9
1979	315	3.8	3.4-4.3	47	0.6	0.4-0.8
1980	342	4.0	3.6-4.5	47	0.6	0.4-0.8
1981	333	3.8	3.4-4.2	34	0.4	0.3-0.5
1982	344	3.9	3.5-4.4	32	0.4	0.3-0.6
1983	358	4.0	3.6-4.4	38	0.5	0.3-0.7
1984	396	4.4	4.0-4.9	28	0.4	0.2-0.5
1985	380	4.2	3.8-4.7	36	0.4	0.3-0.6
1986	378	4.0	3.6-4.5	36	0.4	0.3-0.6
1987	401	4.3	3.8-4.7	30	0.3	0.2-0.5
1988	378	4.0	3.6-4.4	26	0.3	0.2-0.5
1989	390	4.1	3.7-4.5	34	0.4	0.2-0.5
1990	433	4.6	4.1-5.0	28	0.3	0.2-0.4
1991	410	4.3	3.9-4.7	23	0.2	0.2-0.4
1992	441	4.6	4.2-5.1	31	0.3	0.2-0.5
1993	451	4.7	4.3-5.2	35	0.4	0.3-0.6
1994	442	4.6	4.2-5.1	22	0.3	0.2-0.4
1995	430	4.6	4.1-5.0	23	0.3	0.2-0.4
1996	441	4.7	4.2-5.1	27	0.3	0.2-0.4
1997	447	4.7	4.3-5.2	13	0.2	0.1-0.3
1998	472	5.0	4.6-5.5	27	0.3	0.2-0.4
1999	510	5.4	4.9-5.9	18	0.2	0.1-0.3
2000	470	5.0	4.5-5.5	18	0.2	0.1-0.3
2001	475	5.0	4.6-5.5	20	0.2	0.1-0.3
2002	465	4.9	4.5-5.4	31	0.3	0.2-0.5
2003	520	5.5	5.1-6.0	23	0.2	0.2-0.4
2004	504	5.4	4.9-5.9	20	0.2	0.1-0.3
2005	538	5.8	5.3-6.3	19	0.2	0.1-0.3
2006	512	5.5	5.0-6.0	25	0.3	0.2-0.4
2007	554	5.9	5.5-6.5	11	0.1	0.1-0.2
2008	560	6.0	5.5-6.5	16	0.2	0.1-0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Testis**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	550	5.9	5.4-6.4	16	0.2	0.1-0.3
2010	558	6.0	5.5-6.5	30	0.3	0.2-0.5
2011	570	6.0	5.5-6.5	20	0.2	0.1-0.3
2012	585	6.2	5.7-6.8	21	0.2	0.1-0.3
2013	543	5.7	5.2-6.2	15	0.2	0.1-0.3
2014	564	5.9	5.5-6.5	19	0.2	0.1-0.3
2015	560	5.8	5.3-6.3	19	0.2	0.1-0.3
2016	588	6.2	5.7-6.7	17	0.2	0.1-0.3
2017	583	6.2	5.7-6.7	20	0.2	0.1-0.3
2018	604	6.5	6.0-7.0	18	0.2	0.1-0.3
2019	574	6.1	5.6-6.6	30	0.3	0.2-0.4
2020	547	6.0	5.5-6.5	25	0.3	0.2-0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Urinary bladder (incl. in situ)**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	3,289	20.0	19.3-20.7	2,389	36.2	34.7-37.8	900	9.3	8.7-9.9	1,029	6.7	6.3-7.1	719	12.1	11.2-13.1	310	3.4	3.0-3.8
1977	3,402	20.5	19.8-21.3	2,442	36.5	35.0-38.0	960	10.0	9.4-10.7	1,037	6.7	6.3-7.1	718	12.1	11.2-13.0	319	3.4	3.0-3.8
1978	3,578	21.6	20.9-22.3	2,590	39.1	37.5-40.7	988	10.2	9.5-10.8	1,012	6.4	6.0-6.8	690	11.5	10.6-12.4	322	3.4	3.0-3.8
1979	3,620	21.6	20.9-22.4	2,625	39.6	38.0-41.2	995	10.1	9.4-10.7	1,006	6.4	6.0-6.8	674	11.2	10.3-12.1	332	3.5	3.1-3.9
1980	3,809	22.5	21.7-23.2	2,805	41.6	40.0-43.2	1,004	10.0	9.4-10.7	985	6.0	5.7-6.4	681	11.1	10.3-12.0	304	3.1	2.7-3.5
1981	3,695	21.7	21.0-22.4	2,698	39.9	38.3-41.5	997	9.9	9.3-10.6	996	6.1	5.7-6.5	686	11.3	10.4-12.2	310	3.1	2.8-3.5
1982	3,773	21.9	21.2-22.6	2,703	39.5	37.9-41.0	1,070	10.5	9.9-11.2	962	5.8	5.4-6.2	664	10.7	9.9-11.6	298	2.9	2.6-3.3
1983	3,844	22.2	21.5-22.9	2,780	40.9	39.4-42.6	1,064	10.3	9.7-10.9	922	5.5	5.2-5.9	635	10.5	9.7-11.4	287	2.8	2.5-3.1
1984	3,730	21.3	20.6-22.0	2,689	38.6	37.1-40.2	1,041	10.1	9.5-10.7	935	5.6	5.3-6.0	630	10.3	9.4-11.1	305	2.9	2.6-3.3
1985	3,746	21.4	20.7-22.1	2,681	38.6	37.1-40.2	1,065	10.3	9.6-10.9	939	5.6	5.2-5.9	634	10.3	9.4-11.1	305	2.9	2.6-3.3
1986	3,891	22.1	21.4-22.8	2,817	40.3	38.8-41.9	1,074	10.3	9.7-11.0	914	5.4	5.1-5.8	610	10.1	9.2-10.9	304	2.9	2.5-3.2
1987	3,752	21.1	20.4-21.8	2,680	37.9	36.4-39.4	1,072	10.1	9.5-10.8	859	5.0	4.7-5.4	575	9.2	8.4-10.0	284	2.7	2.4-3.0
1988	3,748	21.0	20.3-21.7	2,669	37.6	36.1-39.1	1,079	10.2	9.6-10.8	919	5.3	4.9-5.6	596	9.2	8.5-10.0	323	3.0	2.7-3.3
1989	3,733	20.9	20.2-21.6	2,686	38.0	36.5-39.5	1,047	9.8	9.2-10.5	906	5.2	4.9-5.6	618	9.8	9.0-10.6	288	2.6	2.3-3.0
1990	3,865	21.6	20.9-22.3	2,739	38.6	37.1-40.1	1,126	10.6	10.0-11.2	890	5.1	4.8-5.5	574	9.0	8.3-9.8	316	2.9	2.6-3.2
1991	3,901	21.6	20.9-22.3	2,797	39.0	37.5-40.5	1,104	10.3	9.7-11.0	857	4.8	4.5-5.2	580	8.9	8.2-9.7	277	2.5	2.2-2.8
1992	3,918	21.6	20.9-22.3	2,773	38.6	37.1-40.1	1,145	10.7	10.1-11.3	885	4.9	4.6-5.3	542	8.4	7.7-9.1	343	3.1	2.7-3.4
1993	3,992	21.8	21.1-22.5	2,786	38.0	36.6-39.5	1,206	11.1	10.5-11.8	940	5.2	4.9-5.6	626	9.5	8.8-10.3	314	2.8	2.5-3.1
1994	3,964	21.6	20.9-22.3	2,795	38.1	36.6-39.6	1,169	10.8	10.2-11.4	918	5.0	4.7-5.4	603	9.1	8.3-9.8	315	2.8	2.5-3.1
1995	4,068	22.0	21.4-22.7	2,868	38.8	37.4-40.3	1,200	11.0	10.4-11.6	912	5.0	4.7-5.3	588	8.7	8.0-9.4	324	2.8	2.5-3.2
1996	4,163	22.4	21.8-23.1	3,007	40.4	38.9-41.9	1,156	10.5	9.9-11.2	932	5.0	4.7-5.4	604	8.9	8.2-9.7	328	2.8	2.5-3.2
1997	4,471	23.9	23.2-24.6	3,185	42.1	40.6-43.6	1,286	11.6	11.0-12.3	942	5.0	4.7-5.4	617	8.9	8.2-9.6	325	2.8	2.5-3.1
1998	4,630	24.5	23.8-25.2	3,316	43.2	41.7-44.7	1,314	11.8	11.2-12.5	929	4.9	4.6-5.2	620	8.8	8.1-9.5	309	2.6	2.3-2.9
1999	4,561	23.8	23.1-24.5	3,264	41.9	40.5-43.4	1,297	11.5	10.9-12.2	913	4.7	4.4-5.1	619	8.7	8.0-9.4	294	2.5	2.2-2.8
2000	4,673	24.1	23.4-24.8	3,332	42.2	40.8-43.7	1,341	11.7	11.1-12.3	925	4.7	4.4-5.0	621	8.4	7.7-9.1	304	2.5	2.2-2.8
2001	4,625	23.6	22.9-24.3	3,345	42.0	40.6-43.5	1,280	11.1	10.5-11.7	946	4.8	4.5-5.1	644	8.7	8.0-9.4	302	2.4	2.2-2.7
2002	4,729	23.8	23.2-24.5	3,463	42.8	41.4-44.3	1,266	10.9	10.3-11.5	904	4.5	4.2-4.8	588	7.8	7.2-8.5	316	2.6	2.3-2.9
2003	4,879	24.4	23.7-25.1	3,568	43.3	41.9-44.7	1,311	11.2	10.6-11.8	847	4.2	3.9-4.5	567	7.4	6.8-8.1	280	2.2	2.0-2.5
2004	5,035	24.9	24.2-25.6	3,668	44.0	42.6-45.5	1,367	11.5	10.9-12.2	922	4.5	4.2-4.8	624	7.9	7.3-8.5	298	2.4	2.1-2.7
2005	5,082	24.9	24.2-25.6	3,680	43.7	42.3-45.1	1,402	11.7	11.1-12.4	888	4.3	4.0-4.6	582	7.4	6.8-8.0	306	2.4	2.1-2.7
2006	4,983	24.2	23.6-24.9	3,677	43.4	42.0-44.8	1,306	11.0	10.4-11.6	891	4.3	4.0-4.6	599	7.5	6.9-8.1	292	2.3	2.0-2.6
2007	5,010	24.1	23.5-24.8	3,665	42.5	41.1-43.9	1,345	11.3	10.7-11.9	981	4.6	4.3-4.9	671	8.2	7.6-8.8	310	2.4	2.1-2.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Urinary bladder (incl. in situ)**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	5,172	24.5	23.8-25.2	3,883	44.2	42.8-45.6	1,289	10.6	10.0-11.2	957	4.5	4.2-4.8	654	7.9	7.3-8.5	303	2.3	2.1-2.6
2009	5,199	24.3	23.7-25.0	3,837	43.0	41.7-44.4	1,362	11.0	10.4-11.6	953	4.4	4.1-4.7	684	8.1	7.5-8.7	269	2.0	1.8-2.3
2010	4,917	22.6	22.0-23.3	3,698	40.7	39.4-42.1	1,219	9.7	9.2-10.3	1,015	4.6	4.3-4.9	693	7.9	7.3-8.5	322	2.4	2.1-2.7
2011	5,285	24.0	23.3-24.7	3,923	42.2	40.9-43.6	1,362	10.9	10.3-11.5	956	4.2	4.0-4.5	653	7.4	6.8-8.0	303	2.2	2.0-2.5
2012	5,366	23.8	23.2-24.5	3,939	41.8	40.5-43.1	1,427	10.9	10.4-11.5	989	4.3	4.0-4.6	674	7.4	6.8-8.0	315	2.3	2.0-2.5
2013	5,458	24.0	23.3-24.6	4,043	42.1	40.8-43.4	1,415	10.8	10.3-11.4	978	4.2	3.9-4.5	692	7.5	6.9-8.1	286	2.0	1.8-2.2
2014	5,358	23.0	22.3-23.6	3,974	40.2	38.9-41.5	1,384	10.3	9.8-10.9	1,058	4.4	4.2-4.7	717	7.6	7.1-8.2	341	2.4	2.1-2.6
2015	5,398	22.8	22.2-23.4	4,032	39.9	38.7-41.2	1,366	10.2	9.7-10.8	986	4.1	3.8-4.3	667	6.8	6.3-7.4	319	2.2	2.0-2.5
2016	5,452	22.5	21.9-23.2	4,096	39.6	38.4-40.9	1,356	9.8	9.3-10.4	1,010	4.1	3.9-4.4	708	7.2	6.7-7.8	302	2.1	1.8-2.3
2017	5,525	22.5	21.9-23.1	4,112	39.1	37.9-40.4	1,413	10.1	9.6-10.7	986	3.9	3.7-4.2	684	6.8	6.3-7.3	302	2.0	1.8-2.3
2018	5,427	21.6	21.1-22.2	4,045	37.4	36.3-38.6	1,382	9.8	9.3-10.3	987	3.9	3.6-4.1	690	6.7	6.2-7.2	297	2.0	1.8-2.2
2019	5,503	21.7	21.1-22.3	4,075	37.0	35.9-38.2	1,428	10.0	9.5-10.5	930	3.6	3.3-3.8	666	6.3	5.8-6.8	264	1.7	1.5-1.9
2020	4,905	19.0	18.5-19.5	3,677	32.8	31.7-33.9	1,228	8.5	8.0-9.0	878	3.3	3.1-3.6	613	5.8	5.3-6.2	265	1.7	1.5-1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Kidney and renal pelvis**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,220	7.2	6.8-7.6	756	10.4	9.7-11.3	464	4.8	4.4-5.3	653	3.9	3.6-4.2	416	5.9	5.3-6.6	237	2.4	2.1-2.7
1977	1,242	7.3	6.9-7.7	782	10.8	10.0-11.6	460	4.8	4.3-5.2	667	4.0	3.7-4.3	415	5.9	5.3-6.5	252	2.6	2.3-2.9
1978	1,275	7.5	7.1-7.9	787	10.9	10.2-11.8	488	5.0	4.6-5.5	634	3.7	3.4-4.0	402	5.6	5.1-6.2	232	2.4	2.1-2.7
1979	1,253	7.3	6.9-7.7	781	10.8	10.0-11.6	472	4.7	4.3-5.1	688	4.1	3.8-4.4	438	6.2	5.6-6.8	250	2.6	2.3-2.9
1980	1,253	7.2	6.8-7.7	761	10.3	9.6-11.1	492	5.0	4.6-5.5	636	3.7	3.4-4.0	387	5.5	4.9-6.1	249	2.4	2.1-2.8
1981	1,331	7.8	7.4-8.3	794	11.2	10.4-12.0	537	5.5	5.0-5.9	635	3.7	3.4-4.0	387	5.4	4.9-6.0	248	2.5	2.2-2.8
1982	1,426	8.2	7.8-8.7	862	11.7	10.9-12.6	564	5.7	5.2-6.2	650	3.7	3.4-4.0	385	5.5	4.9-6.1	265	2.6	2.3-2.9
1983	1,489	8.6	8.1-9.0	908	12.4	11.5-13.2	581	5.9	5.4-6.4	642	3.7	3.4-4.0	384	5.3	4.8-5.9	258	2.6	2.3-2.9
1984	1,530	8.8	8.3-9.2	961	13.1	12.2-14.0	569	5.7	5.2-6.2	658	3.8	3.5-4.1	405	5.8	5.2-6.4	253	2.5	2.2-2.8
1985	1,597	9.1	8.6-9.5	963	13.1	12.2-13.9	634	6.2	5.7-6.7	684	3.9	3.6-4.2	390	5.5	5.0-6.1	294	2.9	2.6-3.2
1986	1,603	9.1	8.6-9.5	975	13.1	12.3-14.0	628	6.2	5.7-6.7	733	4.1	3.8-4.5	442	6.1	5.6-6.8	291	2.8	2.5-3.1
1987	1,746	9.9	9.4-10.3	1,039	14.0	13.1-14.9	707	7.0	6.5-7.5	718	4.1	3.8-4.4	439	6.2	5.6-6.8	279	2.7	2.3-3.0
1988	1,619	9.2	8.7-9.6	967	13.0	12.2-13.9	652	6.4	5.9-7.0	688	3.9	3.6-4.2	397	5.5	5.0-6.1	291	2.8	2.4-3.1
1989	1,635	9.2	8.7-9.7	1,008	13.5	12.7-14.4	627	6.1	5.6-6.6	683	3.9	3.6-4.2	401	5.7	5.1-6.3	282	2.7	2.4-3.0
1990	1,822	10.2	9.7-10.7	1,101	14.7	13.8-15.6	721	7.0	6.5-7.6	680	3.8	3.5-4.1	401	5.5	5.0-6.1	279	2.6	2.3-2.9
1991	1,880	10.5	10.0-11.0	1,125	14.8	13.9-15.7	755	7.4	6.8-7.9	705	3.9	3.6-4.2	416	5.7	5.1-6.2	289	2.7	2.4-3.0
1992	1,895	10.5	10.0-11.0	1,148	14.9	14.1-15.8	747	7.3	6.7-7.8	649	3.6	3.3-3.9	385	5.4	4.9-6.0	264	2.5	2.2-2.8
1993	1,952	10.7	10.2-11.2	1,202	15.5	14.6-16.4	750	7.2	6.7-7.7	665	3.7	3.4-4.0	418	5.5	5.0-6.1	247	2.3	2.0-2.6
1994	2,020	11.1	10.6-11.6	1,196	15.4	14.6-16.3	824	7.9	7.4-8.5	729	4.0	3.7-4.3	451	6.1	5.6-6.7	278	2.6	2.3-2.9
1995	2,017	11.0	10.5-11.5	1,193	15.2	14.3-16.1	824	7.9	7.3-8.4	712	3.9	3.6-4.2	415	5.6	5.1-6.2	297	2.7	2.4-3.1
1996	2,176	11.8	11.3-12.3	1,341	17.0	16.1-18.0	835	7.9	7.4-8.4	722	3.9	3.6-4.2	416	5.5	5.0-6.1	306	2.8	2.5-3.1
1997	2,255	12.1	11.6-12.7	1,379	17.5	16.5-18.4	876	8.3	7.7-8.8	720	3.9	3.6-4.2	441	5.8	5.3-6.4	279	2.5	2.2-2.8
1998	2,386	12.7	12.2-13.2	1,453	17.8	16.9-18.7	933	8.8	8.2-9.3	702	3.7	3.5-4.0	427	5.5	5.0-6.0	275	2.5	2.2-2.8
1999	2,515	13.2	12.7-13.7	1,529	18.5	17.6-19.5	986	9.1	8.6-9.7	613	3.2	3.0-3.5	377	4.8	4.3-5.3	236	2.0	1.8-2.3
2000	2,500	12.9	12.4-13.4	1,462	17.5	16.7-18.5	1,038	9.4	8.9-10.0	680	3.5	3.2-3.8	415	5.2	4.7-5.8	265	2.3	2.0-2.6
2001	2,639	13.5	13.0-14.0	1,627	19.1	18.2-20.1	1,012	9.2	8.6-9.7	719	3.7	3.4-3.9	419	5.2	4.7-5.7	300	2.6	2.3-3.0
2002	2,672	13.5	13.0-14.1	1,661	19.3	18.4-20.3	1,011	9.0	8.5-9.6	676	3.4	3.2-3.7	411	5.0	4.6-5.6	265	2.2	2.0-2.5
2003	2,822	14.1	13.6-14.6	1,767	20.1	19.1-21.0	1,055	9.4	8.8-9.9	700	3.5	3.2-3.8	425	5.1	4.6-5.6	275	2.3	2.0-2.6
2004	3,050	15.1	14.6-15.7	1,890	21.3	20.3-22.3	1,160	10.3	9.7-10.9	724	3.6	3.3-3.8	435	5.2	4.7-5.7	289	2.4	2.2-2.7
2005	3,061	15.1	14.6-15.7	1,845	20.8	19.8-21.8	1,216	10.7	10.1-11.4	692	3.4	3.1-3.6	438	5.1	4.6-5.6	254	2.1	1.9-2.4
2006	3,339	16.3	15.8-16.9	2,113	23.4	22.4-24.4	1,226	10.7	10.1-11.4	724	3.5	3.2-3.8	432	5.1	4.6-5.6	292	2.4	2.1-2.7
2007	3,426	16.6	16.0-17.1	2,173	23.7	22.7-24.7	1,253	11.0	10.4-11.6	720	3.4	3.2-3.7	453	5.2	4.7-5.7	267	2.2	1.9-2.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Kidney and renal pelvis**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	3,428	16.3	15.7-16.8	2,137	22.8	21.8-23.8	1,291	11.1	10.5-11.7	699	3.3	3.0-3.5	447	5.0	4.5-5.5	252	2.0	1.8-2.3
2009	3,363	15.8	15.3-16.4	2,073	21.8	20.8-22.8	1,290	11.0	10.4-11.6	727	3.4	3.1-3.6	480	5.3	4.8-5.8	247	2.0	1.7-2.2
2010	3,482	16.1	15.5-16.6	2,198	22.6	21.7-23.6	1,284	10.8	10.2-11.4	697	3.2	2.9-3.4	437	4.7	4.2-5.2	260	2.0	1.8-2.3
2011	3,576	16.4	15.8-16.9	2,300	23.3	22.4-24.3	1,276	10.7	10.1-11.3	715	3.2	2.9-3.4	444	4.7	4.3-5.2	271	2.1	1.8-2.3
2012	3,666	16.5	16.0-17.0	2,318	23.1	22.2-24.1	1,348	11.1	10.5-11.7	737	3.2	3.0-3.5	480	5.0	4.6-5.5	257	1.9	1.7-2.2
2013	3,745	16.5	16.0-17.1	2,398	23.3	22.4-24.3	1,347	11.0	10.4-11.6	676	2.9	2.7-3.2	429	4.4	4.0-4.8	247	1.8	1.6-2.1
2014	3,854	16.7	16.2-17.3	2,514	23.9	22.9-24.9	1,340	10.7	10.2-11.4	685	2.9	2.7-3.1	436	4.3	3.9-4.7	249	1.8	1.6-2.1
2015	3,963	16.9	16.3-17.4	2,540	23.8	22.9-24.8	1,423	11.0	10.5-11.7	759	3.1	2.9-3.4	488	4.7	4.3-5.2	271	1.9	1.7-2.2
2016	3,967	16.8	16.3-17.4	2,591	24.0	23.1-25.0	1,376	10.8	10.2-11.4	634	2.6	2.4-2.8	419	4.0	3.6-4.4	215	1.6	1.4-1.8
2017	4,054	17.0	16.5-17.6	2,557	23.4	22.5-24.4	1,497	11.6	11.0-12.3	664	2.7	2.5-2.9	428	4.1	3.7-4.5	236	1.7	1.5-1.9
2018	4,183	17.4	16.9-18.0	2,697	24.4	23.5-25.4	1,486	11.6	11.0-12.2	702	2.8	2.6-3.0	453	4.2	3.8-4.6	249	1.7	1.5-1.9
2019	4,297	17.7	17.2-18.3	2,799	25.1	24.1-26.0	1,498	11.6	11.0-12.2	643	2.5	2.3-2.7	394	3.5	3.2-3.9	249	1.7	1.5-1.9
2020	3,720	15.3	14.8-15.8	2,447	21.9	21.0-22.8	1,273	9.6	9.1-10.2	636	2.5	2.3-2.7	410	3.8	3.4-4.2	226	1.5	1.3-1.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Brain and other central nervous system**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	991	5.6	5.2-5.9	559	6.9	6.3-7.5	432	4.5	4.1-5.0	746	4.3	4.0-4.6	407	5.2	4.7-5.8	339	3.5	3.1-3.9
1977	1,016	5.8	5.4-6.1	563	7.1	6.5-7.7	453	4.7	4.3-5.2	745	4.2	3.9-4.5	404	5.1	4.6-5.6	341	3.5	3.1-3.9
1978	959	5.5	5.1-5.9	492	6.2	5.7-6.8	467	4.9	4.5-5.4	703	3.9	3.7-4.3	377	4.8	4.3-5.3	326	3.3	2.9-3.6
1979	1,052	6.0	5.6-6.3	585	7.4	6.8-8.1	467	4.8	4.4-5.3	702	4.0	3.7-4.3	392	5.1	4.6-5.6	310	3.1	2.8-3.5
1980	1,052	5.9	5.6-6.3	585	7.3	6.7-8.0	467	4.9	4.4-5.3	774	4.3	4.0-4.7	450	5.8	5.2-6.3	324	3.3	2.9-3.7
1981	1,034	5.9	5.6-6.3	578	7.5	6.9-8.2	456	4.7	4.2-5.1	711	4.0	3.7-4.3	387	5.1	4.6-5.6	324	3.2	2.9-3.6
1982	1,091	6.2	5.9-6.6	569	7.3	6.7-7.9	522	5.4	4.9-5.9	757	4.3	4.0-4.6	391	5.1	4.6-5.7	366	3.7	3.3-4.1
1983	1,043	5.9	5.6-6.3	536	6.7	6.2-7.3	507	5.2	4.8-5.7	729	4.1	3.8-4.4	382	5.0	4.5-5.5	347	3.4	3.0-3.8
1984	1,157	6.4	6.1-6.8	604	7.6	7.0-8.2	553	5.6	5.1-6.1	738	4.1	3.8-4.4	398	5.2	4.7-5.7	340	3.3	3.0-3.7
1985	1,135	6.4	6.1-6.8	601	7.7	7.1-8.3	534	5.5	5.0-6.0	725	4.1	3.8-4.4	363	4.8	4.3-5.3	362	3.6	3.3-4.0
1986	1,125	6.3	6.0-6.7	625	8.0	7.3-8.6	500	5.1	4.6-5.5	781	4.4	4.1-4.7	429	5.7	5.1-6.2	352	3.4	3.1-3.8
1987	1,225	6.9	6.5-7.3	636	8.0	7.4-8.6	589	5.9	5.4-6.4	756	4.3	3.9-4.6	387	5.0	4.5-5.6	369	3.6	3.2-4.0
1988	1,205	6.7	6.3-7.1	656	8.2	7.6-8.9	549	5.6	5.1-6.1	726	4.0	3.7-4.3	399	5.0	4.5-5.6	327	3.2	2.8-3.5
1989	1,214	6.8	6.4-7.2	671	8.3	7.7-9.0	543	5.6	5.1-6.1	698	3.9	3.6-4.2	380	4.9	4.4-5.4	318	3.1	2.8-3.5
1990	1,277	7.1	6.8-7.5	678	8.6	7.9-9.3	599	6.1	5.6-6.6	678	3.8	3.5-4.1	392	5.1	4.6-5.7	286	2.8	2.5-3.1
1991	1,264	7.0	6.6-7.4	663	8.3	7.7-9.0	601	6.1	5.6-6.6	730	4.1	3.8-4.4	405	5.2	4.7-5.8	325	3.3	2.9-3.6
1992	1,272	7.0	6.6-7.4	670	8.2	7.6-8.9	602	6.0	5.6-6.6	722	4.0	3.7-4.3	368	4.7	4.2-5.2	354	3.4	3.1-3.8
1993	1,255	6.9	6.5-7.3	676	8.3	7.6-8.9	579	5.8	5.3-6.3	710	3.9	3.6-4.2	382	4.8	4.3-5.4	328	3.2	2.9-3.6
1994	1,362	7.4	7.0-7.8	737	8.9	8.3-9.6	625	6.1	5.7-6.6	793	4.3	4.0-4.7	425	5.4	4.9-5.9	368	3.5	3.2-3.9
1995	1,431	7.8	7.4-8.2	723	8.7	8.1-9.4	708	7.0	6.5-7.5	729	4.0	3.7-4.3	389	4.9	4.4-5.4	340	3.3	2.9-3.6
1996	1,446	7.8	7.4-8.2	760	9.2	8.5-9.9	686	6.8	6.3-7.3	767	4.2	3.9-4.5	414	5.1	4.7-5.7	353	3.4	3.0-3.8
1997	1,320	7.1	6.7-7.5	707	8.4	7.8-9.1	613	6.0	5.5-6.5	766	4.1	3.8-4.4	392	4.8	4.3-5.3	374	3.6	3.2-3.9
1998	1,447	7.7	7.3-8.1	792	9.3	8.6-9.9	655	6.4	5.9-6.9	751	4.0	3.7-4.3	406	4.9	4.4-5.4	345	3.2	2.9-3.6
1999	1,311	6.9	6.5-7.3	692	8.1	7.5-8.7	619	5.9	5.5-6.4	754	4.0	3.7-4.3	413	4.9	4.5-5.4	341	3.2	2.8-3.5
2000	1,432	7.5	7.1-7.9	729	8.4	7.8-9.0	703	6.6	6.2-7.2	723	3.7	3.5-4.0	374	4.4	3.9-4.8	349	3.2	2.9-3.5
2001	1,360	7.0	6.7-7.4	703	8.0	7.4-8.6	657	6.2	5.7-6.7	713	3.7	3.4-3.9	377	4.4	4.0-4.9	336	3.1	2.7-3.4
2002	1,364	7.0	6.6-7.4	747	8.5	7.9-9.2	617	5.8	5.4-6.3	759	3.9	3.6-4.1	412	4.8	4.3-5.3	347	3.2	2.8-3.5
2003	1,382	7.0	6.7-7.4	705	7.9	7.3-8.5	677	6.4	5.9-6.9	696	3.5	3.2-3.8	398	4.5	4.0-4.9	298	2.7	2.4-3.0
2004	1,329	6.7	6.4-7.1	736	8.2	7.6-8.8	593	5.6	5.1-6.1	751	3.8	3.5-4.0	398	4.5	4.0-4.9	353	3.2	2.8-3.5
2005	1,382	7.0	6.6-7.4	782	8.7	8.1-9.4	600	5.6	5.2-6.1	781	3.9	3.6-4.2	437	4.9	4.5-5.4	344	3.1	2.8-3.4
2006	1,359	6.8	6.5-7.2	761	8.4	7.8-9.0	598	5.6	5.1-6.0	742	3.7	3.4-3.9	439	4.8	4.4-5.3	303	2.7	2.4-3.0
2007	1,316	6.6	6.3-7.0	698	7.7	7.1-8.3	618	5.8	5.3-6.3	762	3.7	3.4-4.0	435	4.8	4.3-5.3	327	2.8	2.5-3.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Brain and other central nervous system**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	1,361	6.8	6.4-7.1	737	8.0	7.5-8.6	624	5.7	5.3-6.2	728	3.5	3.2-3.8	415	4.5	4.1-5.0	313	2.7	2.4-3.0
2009	1,399	6.9	6.5-7.2	749	8.0	7.5-8.6	650	5.9	5.4-6.4	812	3.9	3.6-4.1	447	4.8	4.4-5.3	365	3.1	2.8-3.5
2010	1,342	6.6	6.2-6.9	732	7.8	7.3-8.4	610	5.5	5.1-6.0	789	3.7	3.5-4.0	422	4.5	4.0-4.9	367	3.1	2.8-3.5
2011	1,352	6.5	6.2-6.9	740	7.7	7.2-8.3	612	5.6	5.1-6.0	771	3.6	3.3-3.8	417	4.3	3.9-4.8	354	2.9	2.6-3.3
2012	1,397	6.6	6.3-7.0	765	7.8	7.3-8.4	632	5.6	5.2-6.1	850	3.8	3.6-4.1	449	4.6	4.1-5.0	401	3.3	3.0-3.6
2013	1,407	6.7	6.3-7.0	783	7.9	7.3-8.4	624	5.6	5.1-6.0	825	3.6	3.4-3.9	446	4.4	4.0-4.9	379	3.0	2.7-3.3
2014	1,481	6.9	6.5-7.3	820	8.2	7.6-8.8	661	5.8	5.4-6.3	871	3.8	3.6-4.1	482	4.7	4.2-5.1	389	3.1	2.8-3.5
2015	1,440	6.7	6.3-7.0	756	7.5	7.0-8.1	684	6.0	5.5-6.4	839	3.6	3.4-3.9	426	4.1	3.7-4.5	413	3.3	3.0-3.6
2016	1,427	6.5	6.1-6.8	823	8.0	7.5-8.6	604	5.1	4.7-5.6	895	3.8	3.6-4.1	509	4.8	4.4-5.3	386	3.0	2.7-3.4
2017	1,512	6.9	6.5-7.3	842	8.3	7.7-8.9	670	5.7	5.3-6.2	907	3.8	3.6-4.1	518	4.8	4.4-5.3	389	3.0	2.7-3.4
2018	1,401	6.3	6.0-6.7	789	7.6	7.0-8.1	612	5.3	4.9-5.7	908	3.9	3.6-4.2	508	4.8	4.3-5.2	400	3.1	2.8-3.5
2019	1,382	6.3	5.9-6.6	760	7.4	6.9-8.0	622	5.3	4.8-5.7	893	3.7	3.5-4.0	505	4.6	4.2-5.1	388	2.9	2.7-3.3
2020	1,362	6.2	5.8-6.5	740	7.0	6.5-7.6	622	5.4	5.0-5.9	899	3.7	3.5-4.0	510	4.6	4.2-5.1	389	2.9	2.6-3.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Thyroid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	561	3.4	3.1-3.7	175	2.4	2.0-2.8	386	4.3	3.9-4.7	114	0.7	0.6-0.9	45	0.7	0.5-0.9	69	0.7	0.6-0.9
1977	631	3.8	3.5-4.1	176	2.3	2.0-2.7	455	5.0	4.5-5.5	120	0.7	0.6-0.9	49	0.7	0.5-1.0	71	0.7	0.6-0.9
1978	613	3.7	3.4-4.0	172	2.3	2.0-2.7	441	4.8	4.4-5.3	107	0.7	0.6-0.8	33	0.5	0.4-0.7	74	0.8	0.6-1.0
1979	585	3.5	3.2-3.8	188	2.6	2.2-3.0	397	4.4	3.9-4.8	86	0.5	0.4-0.6	30	0.4	0.3-0.6	56	0.6	0.4-0.7
1980	604	3.5	3.3-3.8	171	2.2	1.9-2.6	433	4.6	4.2-5.1	119	0.7	0.6-0.9	49	0.7	0.5-1.0	70	0.7	0.6-0.9
1981	579	3.4	3.1-3.7	168	2.2	1.9-2.6	411	4.4	4.0-4.9	100	0.6	0.5-0.7	37	0.5	0.4-0.7	63	0.6	0.5-0.8
1982	576	3.4	3.1-3.7	176	2.3	2.0-2.7	400	4.4	3.9-4.8	109	0.6	0.5-0.8	45	0.6	0.5-0.9	64	0.6	0.5-0.8
1983	674	3.9	3.6-4.2	168	2.1	1.8-2.5	506	5.5	5.0-6.0	86	0.5	0.4-0.6	22	0.3	0.2-0.5	64	0.6	0.5-0.8
1984	729	4.2	3.9-4.5	201	2.7	2.3-3.1	528	5.6	5.1-6.1	113	0.7	0.5-0.8	41	0.6	0.5-0.9	72	0.7	0.5-0.9
1985	752	4.3	4.0-4.7	212	2.7	2.3-3.1	540	5.7	5.3-6.3	101	0.6	0.5-0.7	47	0.7	0.5-1.0	54	0.5	0.4-0.7
1986	708	4.0	3.7-4.3	194	2.5	2.1-2.8	514	5.4	4.9-5.9	93	0.5	0.4-0.7	33	0.5	0.3-0.7	60	0.6	0.4-0.8
1987	713	4.1	3.8-4.4	217	2.8	2.4-3.2	496	5.2	4.8-5.7	96	0.5	0.4-0.7	30	0.4	0.3-0.6	66	0.6	0.5-0.8
1988	691	3.9	3.6-4.2	195	2.5	2.1-2.9	496	5.2	4.8-5.7	86	0.5	0.4-0.6	32	0.5	0.3-0.6	54	0.5	0.4-0.7
1989	683	3.9	3.6-4.2	180	2.3	1.9-2.6	503	5.3	4.9-5.8	95	0.5	0.4-0.7	26	0.4	0.2-0.5	69	0.6	0.5-0.8
1990	752	4.2	3.9-4.5	206	2.6	2.2-3.0	546	5.7	5.2-6.2	90	0.5	0.4-0.6	32	0.4	0.3-0.6	58	0.5	0.4-0.7
1991	845	4.7	4.4-5.0	227	2.8	2.4-3.2	618	6.4	5.9-6.9	89	0.5	0.4-0.6	26	0.4	0.2-0.5	63	0.6	0.4-0.8
1992	902	4.9	4.6-5.3	240	2.9	2.6-3.3	662	6.7	6.2-7.3	99	0.5	0.4-0.7	32	0.4	0.3-0.6	67	0.6	0.5-0.8
1993	913	5.0	4.7-5.3	233	2.8	2.5-3.2	680	7.0	6.5-7.6	97	0.5	0.4-0.7	33	0.4	0.3-0.6	64	0.6	0.4-0.7
1994	1,016	5.5	5.2-5.9	265	3.2	2.8-3.6	751	7.7	7.2-8.3	93	0.5	0.4-0.6	39	0.5	0.4-0.7	54	0.5	0.4-0.6
1995	1,101	5.9	5.6-6.3	286	3.4	3.0-3.9	815	8.2	7.7-8.8	87	0.5	0.4-0.6	32	0.4	0.3-0.6	55	0.5	0.4-0.6
1996	1,177	6.3	6.0-6.7	296	3.5	3.1-3.9	881	9.0	8.4-9.6	131	0.7	0.6-0.8	48	0.7	0.5-0.9	83	0.8	0.6-0.9
1997	1,280	6.8	6.4-7.2	334	3.9	3.5-4.3	946	9.5	8.9-10.1	101	0.5	0.4-0.7	40	0.5	0.4-0.7	61	0.5	0.4-0.7
1998	1,353	7.2	6.8-7.5	356	4.1	3.7-4.6	997	9.9	9.3-10.6	89	0.5	0.4-0.6	29	0.4	0.2-0.5	60	0.5	0.4-0.7
1999	1,537	8.1	7.7-8.5	354	4.0	3.6-4.5	1,183	11.8	11.1-12.5	93	0.5	0.4-0.6	33	0.4	0.3-0.6	60	0.5	0.4-0.7
2000	1,743	9.0	8.6-9.5	426	4.8	4.3-5.2	1,317	13.0	12.3-13.7	98	0.5	0.4-0.6	29	0.4	0.2-0.5	69	0.6	0.5-0.7
2001	1,836	9.4	9.0-9.9	415	4.6	4.2-5.1	1,421	13.9	13.2-14.7	98	0.5	0.4-0.6	36	0.4	0.3-0.6	62	0.5	0.4-0.7
2002	1,881	9.6	9.2-10.1	482	5.3	4.8-5.8	1,399	13.6	12.9-14.4	91	0.5	0.4-0.6	33	0.4	0.3-0.5	58	0.5	0.4-0.6
2003	1,911	9.7	9.3-10.1	475	5.2	4.7-5.6	1,436	13.9	13.2-14.6	89	0.4	0.4-0.5	25	0.3	0.2-0.5	64	0.5	0.4-0.7
2004	2,165	10.9	10.5-11.4	518	5.6	5.1-6.1	1,647	15.9	15.2-16.7	88	0.4	0.3-0.5	39	0.5	0.3-0.7	49	0.4	0.3-0.5
2005	2,583	13.0	12.5-13.5	626	6.7	6.2-7.3	1,957	18.9	18.1-19.8	109	0.5	0.4-0.6	50	0.6	0.4-0.8	59	0.5	0.4-0.6
2006	2,681	13.5	13.0-14.0	647	6.9	6.4-7.5	2,034	19.6	18.7-20.5	111	0.5	0.4-0.6	44	0.5	0.4-0.7	67	0.5	0.4-0.7
2007	2,964	14.8	14.3-15.4	716	7.6	7.0-8.2	2,248	21.5	20.7-22.5	112	0.5	0.4-0.6	43	0.5	0.3-0.6	69	0.6	0.4-0.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Thyroid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	3,413	16.9	16.4-17.5	777	8.2	7.6-8.8	2,636	25.1	24.2-26.1	105	0.5	0.4-0.6	46	0.5	0.4-0.7	59	0.5	0.4-0.6
2009	3,661	18.1	17.5-18.7	896	9.3	8.7-9.9	2,765	26.3	25.3-27.3	119	0.5	0.5-0.7	49	0.6	0.4-0.7	70	0.5	0.4-0.7
2010	3,723	18.2	17.6-18.8	898	9.2	8.6-9.9	2,825	26.7	25.7-27.7	115	0.5	0.4-0.6	51	0.6	0.4-0.7	64	0.5	0.4-0.6
2011	3,914	18.9	18.3-19.5	960	9.7	9.1-10.3	2,954	27.5	26.5-28.5	120	0.5	0.5-0.7	51	0.6	0.4-0.8	69	0.5	0.4-0.7
2012	4,020	19.4	18.8-20.0	969	9.7	9.1-10.4	3,051	28.6	27.5-29.6	122	0.5	0.4-0.6	53	0.6	0.4-0.7	69	0.5	0.4-0.7
2013	4,327	20.8	20.1-21.4	1,085	10.8	10.1-11.5	3,242	30.2	29.2-31.3	128	0.6	0.5-0.7	53	0.5	0.4-0.7	75	0.6	0.4-0.7
2014	4,297	20.6	19.9-21.2	1,090	10.7	10.1-11.4	3,207	29.9	28.9-31.0	113	0.5	0.4-0.6	44	0.4	0.3-0.6	69	0.5	0.4-0.6
2015	4,296	20.7	20.0-21.3	1,079	10.6	9.9-11.2	3,217	30.2	29.1-31.3	125	0.5	0.4-0.6	55	0.5	0.4-0.7	70	0.5	0.4-0.7
2016	4,058	19.4	18.8-20.1	1,031	10.1	9.4-10.7	3,027	28.3	27.3-29.4	127	0.5	0.4-0.6	49	0.5	0.4-0.6	78	0.6	0.4-0.7
2017	3,893	18.6	18.0-19.2	1,007	9.9	9.2-10.5	2,886	26.9	25.9-27.9	124	0.5	0.4-0.6	55	0.5	0.4-0.7	69	0.5	0.4-0.6
2018	3,805	18.3	17.7-18.9	1,033	10.1	9.5-10.8	2,772	26.0	25.0-27.0	122	0.5	0.4-0.6	53	0.5	0.4-0.6	69	0.5	0.4-0.6
2019	4,115	19.8	19.1-20.4	1,134	10.9	10.3-11.6	2,981	28.2	27.2-29.3	136	0.5	0.5-0.7	63	0.6	0.5-0.8	73	0.5	0.4-0.6
2020	3,229	15.5	15.0-16.1	902	8.8	8.2-9.4	2,327	22.0	21.1-23.0	115	0.5	0.4-0.6	43	0.4	0.3-0.6	72	0.5	0.4-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Hodgkin lymphoma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	626	3.4	3.1-3.7	336	4.0	3.5-4.4	290	2.9	2.6-3.3	239	1.4	1.2-1.6	140	1.8	1.5-2.1	99	1.0	0.8-1.3
1977	584	3.2	2.9-3.5	327	3.9	3.4-4.3	257	2.6	2.3-3.0	247	1.5	1.3-1.7	138	1.8	1.5-2.2	109	1.1	0.9-1.4
1978	617	3.3	3.1-3.6	327	3.9	3.4-4.3	290	2.9	2.6-3.3	213	1.2	1.0-1.4	118	1.5	1.2-1.8	95	1.0	0.8-1.2
1979	626	3.5	3.2-3.8	345	4.2	3.8-4.7	281	2.9	2.5-3.2	191	1.1	1.0-1.3	111	1.4	1.2-1.7	80	0.8	0.7-1.1
1980	669	3.6	3.4-3.9	365	4.4	3.9-4.9	304	3.0	2.7-3.4	217	1.3	1.1-1.4	119	1.6	1.3-1.9	98	1.0	0.8-1.2
1981	628	3.4	3.1-3.7	343	4.0	3.6-4.4	285	2.9	2.5-3.2	198	1.1	1.0-1.3	111	1.5	1.2-1.8	87	0.9	0.7-1.1
1982	650	3.5	3.2-3.8	341	4.1	3.7-4.6	309	3.1	2.7-3.4	201	1.1	1.0-1.3	107	1.3	1.1-1.6	94	1.0	0.8-1.2
1983	631	3.5	3.2-3.8	363	4.3	3.9-4.8	268	2.7	2.4-3.0	168	0.9	0.8-1.1	98	1.2	1.0-1.5	70	0.7	0.5-0.9
1984	638	3.4	3.2-3.7	347	4.1	3.7-4.6	291	2.9	2.6-3.2	182	1.0	0.9-1.2	114	1.4	1.2-1.7	68	0.7	0.5-0.9
1985	670	3.6	3.3-3.9	368	4.3	3.8-4.7	302	3.0	2.7-3.4	183	1.0	0.8-1.1	102	1.2	1.0-1.5	81	0.8	0.6-1.0
1986	639	3.4	3.1-3.7	362	4.1	3.7-4.6	277	2.7	2.4-3.1	201	1.1	0.9-1.3	118	1.4	1.2-1.7	83	0.8	0.6-1.0
1987	598	3.2	2.9-3.5	315	3.6	3.2-4.1	283	2.8	2.5-3.2	162	0.9	0.8-1.0	103	1.2	1.0-1.5	59	0.6	0.4-0.8
1988	642	3.4	3.2-3.7	363	4.1	3.7-4.6	279	2.8	2.5-3.1	137	0.8	0.6-0.9	86	1.1	0.8-1.3	51	0.5	0.4-0.7
1989	644	3.4	3.1-3.7	365	4.1	3.7-4.6	279	2.8	2.5-3.1	170	1.0	0.8-1.1	113	1.4	1.2-1.7	57	0.6	0.4-0.8
1990	612	3.3	3.0-3.6	332	3.8	3.4-4.3	280	2.8	2.5-3.2	165	0.9	0.8-1.1	101	1.3	1.0-1.5	64	0.6	0.5-0.8
1991	662	3.5	3.3-3.8	357	4.1	3.7-4.5	305	3.1	2.7-3.5	139	0.8	0.6-0.9	76	1.0	0.7-1.2	63	0.6	0.5-0.8
1992	643	3.4	3.2-3.7	354	4.0	3.6-4.5	289	2.9	2.6-3.2	158	0.8	0.7-1.0	81	1.0	0.8-1.2	77	0.7	0.6-0.9
1993	601	3.2	3.0-3.5	308	3.5	3.1-3.9	293	2.9	2.6-3.3	129	0.7	0.6-0.8	73	0.9	0.7-1.1	56	0.5	0.4-0.7
1994	652	3.5	3.2-3.7	347	3.9	3.5-4.4	305	3.0	2.7-3.4	135	0.7	0.6-0.9	69	0.8	0.7-1.1	66	0.6	0.5-0.8
1995	614	3.3	3.0-3.5	333	3.7	3.3-4.2	281	2.9	2.5-3.2	121	0.6	0.5-0.8	65	0.8	0.6-1.0	56	0.5	0.4-0.7
1996	580	3.1	2.8-3.3	296	3.3	3.0-3.7	284	2.9	2.6-3.2	106	0.6	0.5-0.7	55	0.7	0.5-0.9	51	0.5	0.4-0.7
1997	577	3.1	2.8-3.3	326	3.6	3.2-4.0	251	2.6	2.3-2.9	113	0.6	0.5-0.7	59	0.7	0.5-0.9	54	0.5	0.4-0.7
1998	647	3.4	3.1-3.7	334	3.7	3.3-4.1	313	3.2	2.8-3.5	105	0.6	0.5-0.7	53	0.6	0.5-0.8	52	0.5	0.4-0.7
1999	597	3.1	2.9-3.4	320	3.5	3.1-3.9	277	2.8	2.5-3.1	98	0.5	0.4-0.6	49	0.6	0.4-0.7	49	0.5	0.3-0.6
2000	653	3.4	3.1-3.7	361	4.0	3.6-4.4	292	2.9	2.6-3.3	92	0.5	0.4-0.6	48	0.6	0.4-0.8	44	0.4	0.3-0.5
2001	647	3.3	3.1-3.6	345	3.7	3.4-4.2	302	3.0	2.7-3.4	99	0.5	0.4-0.6	52	0.6	0.4-0.8	47	0.4	0.3-0.6
2002	704	3.6	3.4-3.9	391	4.3	3.9-4.7	313	3.1	2.8-3.5	104	0.5	0.4-0.6	56	0.6	0.5-0.8	48	0.4	0.3-0.6
2003	625	3.2	3.0-3.5	332	3.6	3.2-4.0	293	2.9	2.6-3.3	87	0.4	0.4-0.5	41	0.5	0.3-0.6	46	0.4	0.3-0.6
2004	662	3.4	3.1-3.7	344	3.7	3.3-4.1	318	3.1	2.8-3.5	88	0.4	0.4-0.5	53	0.6	0.4-0.8	35	0.3	0.2-0.4
2005	672	3.5	3.2-3.7	374	4.0	3.6-4.5	298	3.0	2.6-3.3	83	0.4	0.3-0.5	44	0.5	0.4-0.7	39	0.4	0.3-0.5
2006	647	3.3	3.1-3.6	362	3.9	3.5-4.3	285	2.8	2.5-3.1	84	0.4	0.3-0.5	48	0.5	0.4-0.7	36	0.3	0.2-0.5
2007	641	3.3	3.0-3.5	331	3.5	3.2-3.9	310	3.0	2.7-3.4	99	0.5	0.4-0.6	48	0.5	0.4-0.7	51	0.4	0.3-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Hodgkin lymphoma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	669	3.4	3.2-3.7	364	3.8	3.5-4.3	305	3.0	2.7-3.4	94	0.5	0.4-0.6	55	0.6	0.5-0.8	39	0.3	0.2-0.5
2009	668	3.4	3.1-3.7	360	3.8	3.4-4.2	308	3.0	2.7-3.4	75	0.4	0.3-0.5	41	0.4	0.3-0.6	34	0.3	0.2-0.4
2010	658	3.3	3.0-3.5	370	3.8	3.4-4.3	288	2.7	2.4-3.1	75	0.4	0.3-0.5	43	0.4	0.3-0.6	32	0.3	0.2-0.4
2011	666	3.3	3.1-3.6	372	3.9	3.5-4.3	294	2.9	2.5-3.2	87	0.4	0.3-0.5	54	0.6	0.4-0.8	33	0.3	0.2-0.4
2012	649	3.2	3.0-3.5	363	3.8	3.4-4.2	286	2.8	2.5-3.2	73	0.3	0.3-0.4	36	0.4	0.3-0.5	37	0.3	0.2-0.4
2013	655	3.3	3.0-3.5	332	3.4	3.1-3.8	323	3.1	2.8-3.5	68	0.3	0.2-0.4	34	0.4	0.2-0.5	34	0.3	0.2-0.4
2014	699	3.4	3.2-3.7	367	3.7	3.4-4.1	332	3.1	2.8-3.5	67	0.3	0.2-0.4	29	0.3	0.2-0.4	38	0.3	0.2-0.4
2015	633	3.2	2.9-3.4	354	3.6	3.3-4.1	279	2.7	2.4-3.1	65	0.3	0.2-0.4	33	0.4	0.2-0.5	32	0.3	0.2-0.4
2016	667	3.3	3.0-3.5	332	3.4	3.0-3.8	335	3.2	2.8-3.5	69	0.3	0.2-0.4	42	0.4	0.3-0.6	27	0.2	0.1-0.3
2017	643	3.1	2.9-3.4	347	3.5	3.1-3.9	296	2.8	2.5-3.2	58	0.3	0.2-0.3	30	0.3	0.2-0.4	28	0.2	0.1-0.3
2018	573	2.8	2.5-3.0	330	3.3	3.0-3.7	243	2.3	2.0-2.6	69	0.3	0.2-0.4	45	0.4	0.3-0.6	24	0.2	0.1-0.3
2019	661	3.3	3.0-3.5	356	3.6	3.2-4.0	305	3.0	2.6-3.3	61	0.2	0.2-0.3	38	0.3	0.2-0.5	23	0.2	0.1-0.3
2020	557	2.8	2.5-3.0	312	3.2	2.8-3.6	245	2.4	2.1-2.7	54	0.2	0.2-0.3	33	0.3	0.2-0.4	21	0.2	0.1-0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Non-Hodgkin lymphomas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,880	11.1	10.6-11.7	959	13.4	12.5-14.3	921	9.6	8.9-10.2	1,040	6.2	5.8-6.6	558	8.0	7.3-8.8	482	5.0	4.5-5.5
1977	1,938	11.3	10.8-11.9	983	13.5	12.6-14.4	955	9.8	9.2-10.4	1,043	6.2	5.8-6.6	509	7.2	6.5-7.8	534	5.4	5.0-5.9
1978	2,086	12.2	11.7-12.8	1,090	14.7	13.8-15.7	996	10.3	9.6-10.9	1,058	6.3	5.9-6.7	553	7.9	7.2-8.6	505	5.1	4.7-5.6
1979	2,006	11.9	11.4-12.4	987	13.8	13.0-14.8	1,019	10.4	9.8-11.1	1,076	6.4	6.1-6.9	565	8.1	7.4-8.8	511	5.2	4.8-5.7
1980	2,188	12.8	12.3-13.4	1,108	15.1	14.2-16.1	1,080	11.0	10.3-11.7	1,100	6.4	6.1-6.8	550	7.6	7.0-8.3	550	5.5	5.1-6.0
1981	2,187	12.7	12.2-13.3	1,069	14.7	13.8-15.7	1,118	11.2	10.6-11.9	1,102	6.4	6.1-6.8	543	7.7	7.1-8.4	559	5.5	5.1-6.0
1982	2,325	13.5	12.9-14.0	1,155	15.7	14.8-16.7	1,170	11.7	11.1-12.5	1,124	6.6	6.2-7.0	561	8.1	7.4-8.8	563	5.6	5.1-6.0
1983	2,438	14.0	13.5-14.6	1,224	16.7	15.7-17.7	1,214	12.0	11.3-12.7	1,219	7.1	6.7-7.5	625	8.7	8.0-9.5	594	5.9	5.4-6.4
1984	2,513	14.5	13.9-15.1	1,296	17.9	16.9-19.0	1,217	12.0	11.3-12.7	1,178	6.8	6.4-7.2	628	9.0	8.2-9.7	550	5.3	4.9-5.8
1985	2,724	15.6	15.0-16.2	1,449	19.6	18.5-20.6	1,275	12.5	11.8-13.2	1,319	7.6	7.2-8.0	693	9.6	8.9-10.4	626	6.1	5.6-6.6
1986	2,857	16.2	15.6-16.8	1,523	20.3	19.3-21.4	1,334	13.0	12.3-13.8	1,312	7.5	7.1-8.0	719	10.1	9.3-10.9	593	5.7	5.2-6.2
1987	2,843	16.1	15.5-16.7	1,537	20.6	19.5-21.6	1,306	12.6	12.0-13.4	1,309	7.5	7.0-7.9	661	9.1	8.4-9.8	648	6.2	5.7-6.7
1988	2,878	16.3	15.7-16.9	1,525	20.1	19.1-21.2	1,353	13.3	12.6-14.1	1,394	8.0	7.5-8.4	714	10.0	9.2-10.8	680	6.5	6.0-7.0
1989	3,022	17.1	16.5-17.7	1,656	21.6	20.6-22.7	1,366	13.4	12.7-14.2	1,388	7.9	7.4-8.3	763	10.5	9.8-11.3	625	5.9	5.5-6.4
1990	3,281	18.4	17.7-19.0	1,820	23.5	22.4-24.7	1,461	14.1	13.4-14.9	1,441	8.1	7.7-8.6	733	10.1	9.4-10.9	708	6.7	6.2-7.2
1991	3,412	19.0	18.4-19.7	1,860	24.4	23.2-25.5	1,552	14.9	14.2-15.7	1,444	8.1	7.7-8.5	768	10.5	9.7-11.2	676	6.3	5.8-6.8
1992	3,363	18.6	18.0-19.3	1,824	23.3	22.2-24.4	1,539	14.8	14.1-15.6	1,517	8.4	8.0-8.9	777	10.6	9.9-11.4	740	6.9	6.4-7.4
1993	3,462	19.1	18.4-19.7	1,920	24.2	23.1-25.3	1,542	14.8	14.1-15.6	1,429	7.8	7.4-8.3	728	9.7	9.0-10.5	701	6.4	6.0-6.9
1994	3,806	20.8	20.1-21.5	2,124	26.6	25.4-27.8	1,682	16.0	15.2-16.8	1,565	8.6	8.1-9.0	801	10.7	10.0-11.5	764	7.0	6.5-7.5
1995	3,695	20.1	19.5-20.8	2,022	25.4	24.3-26.5	1,673	15.8	15.1-16.6	1,554	8.5	8.1-8.9	811	10.9	10.1-11.7	743	6.8	6.3-7.3
1996	3,877	20.9	20.3-21.6	2,090	26.1	25.0-27.3	1,787	16.8	16.1-17.7	1,581	8.6	8.1-9.0	816	11.0	10.2-11.7	765	6.9	6.5-7.5
1997	3,877	20.8	20.1-21.4	2,031	25.2	24.1-26.4	1,846	17.2	16.5-18.0	1,624	8.7	8.3-9.1	824	10.7	10.0-11.5	800	7.2	6.7-7.7
1998	3,857	20.5	19.8-21.1	2,011	24.6	23.5-25.7	1,846	17.2	16.4-18.0	1,552	8.2	7.8-8.6	747	9.5	8.8-10.2	805	7.2	6.7-7.7
1999	3,943	20.6	20.0-21.3	2,004	24.3	23.3-25.4	1,939	17.9	17.1-18.7	1,489	7.8	7.4-8.2	739	9.5	8.9-10.3	750	6.6	6.2-7.1
2000	3,907	20.2	19.6-20.8	2,002	23.9	22.9-25.0	1,905	17.3	16.5-18.1	1,526	7.8	7.5-8.3	768	9.7	9.0-10.4	758	6.6	6.1-7.1
2001	3,923	20.1	19.4-20.7	2,046	24.4	23.3-25.4	1,877	16.8	16.1-17.6	1,444	7.3	7.0-7.7	785	9.7	9.0-10.4	659	5.6	5.1-6.0
2002	3,983	20.2	19.5-20.8	2,141	25.0	23.9-26.1	1,842	16.4	15.7-17.2	1,406	7.1	6.7-7.5	761	9.3	8.7-10.0	645	5.5	5.1-5.9
2003	4,225	21.2	20.5-21.8	2,142	24.7	23.7-25.8	2,083	18.4	17.6-19.2	1,323	6.6	6.2-6.9	671	8.2	7.6-8.8	652	5.4	5.0-5.9
2004	4,199	20.9	20.3-21.5	2,223	25.5	24.4-26.6	1,976	17.4	16.7-18.2	1,303	6.4	6.1-6.8	649	7.8	7.2-8.5	654	5.4	5.0-5.8
2005	4,391	21.7	21.0-22.3	2,298	26.0	24.9-27.1	2,093	18.3	17.5-19.1	1,272	6.2	5.9-6.5	667	8.0	7.4-8.6	605	5.0	4.6-5.4
2006	4,469	21.9	21.3-22.6	2,351	26.4	25.3-27.5	2,118	18.3	17.6-19.2	1,360	6.6	6.3-7.0	692	8.1	7.5-8.8	668	5.4	5.0-5.9
2007	4,481	21.7	21.1-22.4	2,392	26.6	25.5-27.7	2,089	18.0	17.3-18.8	1,262	6.0	5.7-6.4	677	7.8	7.2-8.4	585	4.8	4.4-5.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Non-Hodgkin lymphomas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	4,610	22.1	21.5-22.8	2,466	27.0	25.9-28.1	2,144	18.4	17.6-19.2	1,281	6.1	5.7-6.4	678	7.8	7.2-8.4	603	4.9	4.5-5.3
2009	4,759	22.6	21.9-23.2	2,503	27.1	26.0-28.2	2,256	19.1	18.3-20.0	1,250	5.8	5.5-6.2	689	7.7	7.1-8.3	561	4.5	4.1-4.9
2010	4,679	21.8	21.2-22.4	2,478	26.3	25.3-27.4	2,201	18.2	17.4-19.0	1,264	5.8	5.5-6.1	661	7.3	6.8-7.9	603	4.7	4.4-5.2
2011	4,775	22.0	21.4-22.6	2,574	26.9	25.8-27.9	2,201	18.2	17.4-19.0	1,270	5.8	5.5-6.1	676	7.3	6.8-7.9	594	4.6	4.3-5.0
2012	4,740	21.5	20.9-22.1	2,553	26.3	25.3-27.3	2,187	17.8	17.1-18.6	1,236	5.5	5.2-5.8	675	7.3	6.7-7.9	561	4.2	3.9-4.6
2013	4,889	21.9	21.3-22.5	2,641	26.8	25.8-27.9	2,248	18.0	17.2-18.8	1,179	5.2	4.9-5.5	635	6.7	6.2-7.3	544	4.1	3.7-4.4
2014	5,006	22.0	21.4-22.6	2,732	27.3	26.2-28.3	2,274	17.9	17.1-18.7	1,236	5.3	5.0-5.6	677	7.0	6.5-7.6	559	4.0	3.7-4.4
2015	5,013	21.8	21.2-22.4	2,727	26.6	25.6-27.7	2,286	18.0	17.3-18.8	1,182	5.1	4.8-5.4	662	6.8	6.3-7.3	520	3.8	3.5-4.2
2016	5,116	21.9	21.3-22.5	2,703	25.9	24.9-26.9	2,413	18.8	18.0-19.6	1,251	5.2	4.9-5.5	675	6.7	6.2-7.2	576	4.1	3.8-4.5
2017	5,118	21.7	21.1-22.3	2,759	26.2	25.2-27.2	2,359	18.2	17.4-19.0	1,274	5.2	4.9-5.5	713	6.9	6.4-7.5	561	3.9	3.6-4.3
2018	5,020	21.1	20.5-21.7	2,654	24.9	23.9-25.9	2,366	18.2	17.4-19.0	1,112	4.5	4.2-4.7	636	6.1	5.6-6.6	476	3.3	3.0-3.6
2019	5,193	21.4	20.8-22.0	2,830	26.0	25.1-27.0	2,363	17.8	17.1-18.6	1,157	4.6	4.3-4.9	647	6.0	5.6-6.5	510	3.5	3.2-3.8
2020	4,583	18.8	18.3-19.4	2,462	22.5	21.6-23.4	2,121	15.9	15.2-16.7	1,064	4.2	3.9-4.4	581	5.4	4.9-5.8	483	3.3	3.0-3.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Myeloma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	769	4.6	4.2-4.9	385	5.6	5.0-6.2	384	3.9	3.5-4.4	488	3.0	2.7-3.3	244	3.7	3.2-4.2	244	2.5	2.2-2.9
1977	757	4.5	4.2-4.9	372	5.5	4.9-6.1	385	3.9	3.5-4.3	499	3.0	2.7-3.2	247	3.6	3.1-4.1	252	2.5	2.2-2.9
1978	773	4.6	4.3-5.0	373	5.5	4.9-6.1	400	4.1	3.7-4.5	464	2.8	2.5-3.0	230	3.4	2.9-3.8	234	2.3	2.0-2.7
1979	831	4.9	4.6-5.3	394	5.8	5.2-6.4	437	4.4	4.0-4.8	505	3.0	2.8-3.3	235	3.5	3.0-4.0	270	2.7	2.4-3.1
1980	813	4.8	4.5-5.1	391	5.8	5.2-6.4	422	4.2	3.8-4.6	580	3.4	3.1-3.7	273	4.1	3.6-4.6	307	3.0	2.7-3.4
1981	914	5.4	5.0-5.7	428	6.2	5.6-6.9	486	4.8	4.4-5.3	542	3.2	2.9-3.5	255	3.9	3.4-4.4	287	2.8	2.5-3.1
1982	886	5.1	4.8-5.5	440	6.3	5.7-7.0	446	4.3	3.9-4.8	564	3.3	3.0-3.6	273	4.0	3.5-4.5	291	2.9	2.5-3.2
1983	866	5.1	4.7-5.4	446	6.6	6.0-7.3	420	4.1	3.7-4.5	562	3.3	3.0-3.6	265	3.9	3.5-4.5	297	2.9	2.5-3.2
1984	907	5.3	4.9-5.6	452	6.5	5.9-7.2	455	4.4	4.0-4.8	608	3.5	3.2-3.8	283	4.1	3.6-4.6	325	3.1	2.8-3.5
1985	924	5.3	5.0-5.7	460	6.6	6.0-7.2	464	4.5	4.1-4.9	607	3.4	3.2-3.7	315	4.6	4.1-5.1	292	2.7	2.4-3.1
1986	895	5.1	4.8-5.5	466	6.6	6.0-7.2	429	4.1	3.7-4.6	616	3.5	3.3-3.8	302	4.4	3.9-5.0	314	3.0	2.6-3.3
1987	924	5.3	4.9-5.6	468	6.7	6.1-7.3	456	4.3	3.9-4.7	607	3.5	3.2-3.7	316	4.6	4.1-5.2	291	2.8	2.4-3.1
1988	891	5.1	4.7-5.4	458	6.5	5.9-7.2	433	4.1	3.7-4.5	600	3.4	3.1-3.7	323	4.7	4.2-5.3	277	2.6	2.3-2.9
1989	984	5.6	5.3-6.0	449	6.4	5.8-7.0	535	5.1	4.7-5.6	578	3.3	3.0-3.5	281	4.0	3.6-4.5	297	2.7	2.4-3.1
1990	1,023	5.7	5.4-6.1	508	7.0	6.4-7.6	515	4.9	4.5-5.4	581	3.3	3.0-3.5	280	4.1	3.6-4.6	301	2.8	2.5-3.1
1991	978	5.4	5.1-5.8	463	6.4	5.8-7.0	515	4.8	4.4-5.3	614	3.4	3.2-3.7	291	4.1	3.6-4.6	323	3.0	2.7-3.4
1992	1,090	6.0	5.6-6.4	545	7.5	6.9-8.2	545	5.0	4.6-5.5	659	3.6	3.4-3.9	325	4.7	4.2-5.2	334	3.1	2.8-3.4
1993	1,037	5.7	5.4-6.1	497	6.7	6.1-7.4	540	5.0	4.6-5.5	706	3.9	3.6-4.2	341	4.7	4.2-5.3	365	3.3	3.0-3.7
1994	1,041	5.7	5.4-6.1	511	6.8	6.2-7.5	530	5.0	4.5-5.4	697	3.8	3.5-4.1	306	4.2	3.7-4.7	391	3.6	3.2-4.0
1995	1,078	5.9	5.5-6.2	539	7.3	6.7-7.9	539	5.0	4.6-5.5	707	3.8	3.6-4.1	353	4.8	4.3-5.3	354	3.2	2.9-3.6
1996	1,120	6.1	5.7-6.4	552	7.4	6.7-8.0	568	5.3	4.8-5.7	656	3.5	3.3-3.8	319	4.5	4.0-5.0	337	3.0	2.7-3.4
1997	1,106	5.9	5.6-6.3	533	6.9	6.3-7.5	573	5.3	4.9-5.7	654	3.5	3.2-3.8	324	4.4	3.9-4.9	330	3.0	2.7-3.3
1998	1,233	6.5	6.2-6.9	610	7.8	7.2-8.5	623	5.7	5.2-6.1	654	3.5	3.2-3.7	310	4.1	3.6-4.6	344	3.0	2.7-3.4
1999	1,248	6.5	6.2-6.9	596	7.6	7.0-8.2	652	5.8	5.4-6.3	677	3.5	3.3-3.8	328	4.3	3.8-4.8	349	3.0	2.7-3.4
2000	1,260	6.5	6.1-6.9	622	7.7	7.1-8.3	638	5.7	5.2-6.1	706	3.6	3.4-3.9	337	4.4	3.9-4.9	369	3.2	2.9-3.5
2001	1,280	6.5	6.2-6.9	677	8.3	7.7-9.0	603	5.3	4.9-5.7	648	3.3	3.1-3.6	328	4.1	3.7-4.6	320	2.7	2.4-3.1
2002	1,348	6.8	6.4-7.2	699	8.5	7.9-9.1	649	5.6	5.2-6.1	693	3.5	3.2-3.7	350	4.4	4.0-4.9	343	2.9	2.6-3.2
2003	1,338	6.7	6.3-7.0	693	8.3	7.7-9.0	645	5.6	5.1-6.0	645	3.2	3.0-3.5	335	4.1	3.7-4.6	310	2.6	2.3-2.9
2004	1,374	6.8	6.4-7.1	698	8.1	7.5-8.8	676	5.8	5.4-6.3	635	3.1	2.9-3.4	326	4.0	3.6-4.4	309	2.6	2.3-2.9
2005	1,425	6.9	6.6-7.3	721	8.3	7.7-8.9	704	6.0	5.5-6.4	650	3.2	2.9-3.4	321	3.8	3.4-4.2	329	2.7	2.4-3.0
2006	1,457	7.1	6.7-7.5	767	8.7	8.1-9.4	690	5.9	5.4-6.3	622	3.0	2.8-3.3	347	4.1	3.7-4.6	275	2.3	2.0-2.6
2007	1,475	7.1	6.7-7.5	774	8.8	8.1-9.4	701	5.9	5.5-6.4	681	3.3	3.0-3.5	345	4.0	3.6-4.4	336	2.7	2.5-3.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Myeloma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	1,600	7.5	7.2-7.9	844	9.3	8.7-10.0	756	6.3	5.9-6.8	631	3.0	2.8-3.2	331	3.9	3.5-4.3	300	2.4	2.2-2.7
2009	1,720	8.0	7.7-8.4	914	9.9	9.3-10.6	806	6.6	6.1-7.1	647	3.0	2.8-3.2	351	3.9	3.5-4.4	296	2.3	2.1-2.6
2010	1,736	8.0	7.6-8.4	920	9.7	9.1-10.4	816	6.7	6.2-7.2	710	3.3	3.0-3.5	355	4.0	3.6-4.4	355	2.8	2.5-3.1
2011	1,832	8.3	7.9-8.7	997	10.4	9.7-11.0	835	6.8	6.3-7.2	649	2.9	2.7-3.1	321	3.5	3.1-3.9	328	2.5	2.3-2.8
2012	1,815	8.0	7.6-8.4	979	9.9	9.3-10.5	836	6.6	6.2-7.1	687	3.0	2.8-3.3	342	3.6	3.2-4.0	345	2.6	2.3-2.9
2013	1,930	8.4	8.0-8.8	1,053	10.5	9.8-11.1	877	6.8	6.4-7.3	662	2.9	2.6-3.1	358	3.7	3.3-4.1	304	2.3	2.0-2.6
2014	2,049	8.8	8.4-9.2	1,115	10.9	10.2-11.6	934	7.3	6.8-7.7	717	3.1	2.8-3.3	375	3.8	3.4-4.2	342	2.6	2.3-2.8
2015	1,953	8.3	7.9-8.7	1,046	10.1	9.5-10.8	907	6.9	6.5-7.4	735	3.1	2.9-3.3	401	4.0	3.6-4.4	334	2.4	2.2-2.7
2016	2,074	8.7	8.3-9.0	1,150	10.8	10.2-11.5	924	7.0	6.5-7.5	727	3.0	2.8-3.2	359	3.4	3.1-3.8	368	2.7	2.4-2.9
2017	2,056	8.4	8.1-8.8	1,113	10.3	9.7-11.0	943	6.9	6.5-7.4	712	2.9	2.7-3.1	369	3.5	3.2-3.9	343	2.4	2.1-2.7
2018	1,997	8.0	7.7-8.4	1,066	9.6	9.0-10.2	931	6.8	6.4-7.3	664	2.6	2.4-2.8	341	3.2	2.8-3.5	323	2.2	2.0-2.4
2019	2,066	8.3	7.9-8.6	1,133	10.1	9.5-10.7	933	6.8	6.4-7.3	670	2.6	2.4-2.8	376	3.4	3.1-3.8	294	1.9	1.7-2.2
2020	1,730	6.8	6.5-7.2	931	8.2	7.7-8.8	799	5.8	5.4-6.2	589	2.3	2.1-2.5	319	2.9	2.6-3.2	270	1.8	1.6-2.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Leukemias**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	2,038	12.3	11.8-12.9	1,147	16.6	15.6-17.7	891	9.4	8.8-10.1	1,337	8.2	7.8-8.7	739	11.2	10.3-12.1	598	6.3	5.8-6.9
1977	1,904	11.6	11.0-12.1	1,052	15.2	14.2-16.2	852	9.0	8.4-9.7	1,355	8.3	7.8-8.7	735	10.8	10.0-11.7	620	6.6	6.1-7.1
1978	2,021	12.3	11.8-12.9	1,127	16.4	15.4-17.5	894	9.5	8.9-10.2	1,330	8.0	7.6-8.5	705	10.3	9.5-11.2	625	6.6	6.1-7.1
1979	2,086	12.5	11.9-13.0	1,140	16.1	15.2-17.2	946	9.9	9.3-10.5	1,357	8.2	7.8-8.7	725	10.7	9.9-11.5	632	6.7	6.1-7.2
1980	2,203	13.1	12.5-13.7	1,165	16.7	15.7-17.8	1,038	10.8	10.1-11.5	1,378	8.2	7.8-8.7	753	11.0	10.2-11.8	625	6.4	5.9-7.0
1981	2,023	12.0	11.5-12.6	1,131	16.2	15.2-17.2	892	9.2	8.6-9.8	1,374	8.3	7.8-8.7	745	11.0	10.2-11.9	629	6.5	6.0-7.1
1982	2,089	12.4	11.8-12.9	1,158	16.6	15.6-17.6	931	9.5	8.9-10.2	1,401	8.3	7.9-8.8	782	11.6	10.7-12.5	619	6.3	5.8-6.9
1983	2,047	12.0	11.5-12.5	1,115	15.7	14.7-16.7	932	9.5	8.9-10.2	1,345	8.0	7.6-8.4	699	10.4	9.6-11.2	646	6.5	6.0-7.0
1984	2,172	12.6	12.1-13.2	1,167	16.5	15.5-17.5	1,005	10.0	9.4-10.7	1,335	7.8	7.4-8.2	667	9.6	8.8-10.4	668	6.6	6.1-7.2
1985	2,229	12.9	12.3-13.4	1,255	17.5	16.5-18.5	974	9.7	9.1-10.4	1,345	7.8	7.4-8.2	755	10.9	10.1-11.7	590	5.8	5.3-6.3
1986	2,151	12.4	11.9-12.9	1,163	16.1	15.2-17.1	988	9.9	9.2-10.5	1,336	7.7	7.3-8.2	732	10.3	9.5-11.1	604	5.9	5.5-6.4
1987	2,011	11.4	10.9-11.9	1,049	14.4	13.5-15.3	962	9.5	8.9-10.1	1,282	7.3	6.9-7.8	673	9.5	8.8-10.3	609	5.9	5.5-6.4
1988	2,046	11.6	11.1-12.1	1,114	15.5	14.5-16.4	932	9.2	8.6-9.8	1,310	7.5	7.1-7.9	730	10.4	9.7-11.2	580	5.6	5.1-6.0
1989	2,157	12.2	11.7-12.7	1,175	16.3	15.3-17.3	982	9.5	8.9-10.1	1,318	7.4	7.0-7.9	714	10.1	9.4-10.9	604	5.8	5.3-6.3
1990	2,350	13.3	12.7-13.8	1,257	16.9	15.9-17.9	1,093	10.7	10.0-11.3	1,328	7.5	7.1-7.9	674	9.5	8.8-10.3	654	6.2	5.8-6.7
1991	2,257	12.7	12.2-13.2	1,228	16.6	15.6-17.5	1,029	10.0	9.4-10.7	1,296	7.3	6.9-7.7	685	9.7	9.0-10.5	611	5.8	5.3-6.3
1992	2,304	12.8	12.3-13.4	1,276	17.2	16.2-18.2	1,028	9.9	9.3-10.5	1,334	7.5	7.1-7.9	720	10.1	9.3-10.9	614	5.8	5.3-6.3
1993	2,264	12.5	12.0-13.0	1,203	15.8	14.9-16.8	1,061	10.3	9.6-10.9	1,267	7.0	6.6-7.4	689	9.5	8.8-10.3	578	5.4	5.0-5.9
1994	2,457	13.5	12.9-14.0	1,326	17.5	16.5-18.5	1,131	10.7	10.1-11.3	1,418	7.8	7.4-8.2	767	10.4	9.7-11.2	651	6.0	5.5-6.5
1995	2,349	12.8	12.2-13.3	1,272	16.5	15.6-17.5	1,077	10.2	9.6-10.8	1,372	7.5	7.1-7.9	755	10.4	9.7-11.2	617	5.7	5.3-6.2
1996	2,446	13.2	12.7-13.7	1,394	17.8	16.8-18.8	1,052	9.9	9.3-10.5	1,403	7.6	7.2-8.0	745	9.9	9.2-10.7	658	6.1	5.6-6.5
1997	2,465	13.2	12.7-13.7	1,309	16.7	15.8-17.6	1,156	10.7	10.1-11.4	1,360	7.3	6.9-7.7	752	10.0	9.3-10.8	608	5.5	5.0-5.9
1998	2,469	13.1	12.6-13.6	1,382	17.2	16.3-18.2	1,087	10.1	9.5-10.7	1,342	7.1	6.7-7.5	731	9.4	8.7-10.1	611	5.5	5.0-5.9
1999	2,686	14.1	13.6-14.6	1,460	18.3	17.3-19.3	1,226	11.2	10.6-11.8	1,424	7.4	7.0-7.8	769	10.0	9.3-10.7	655	5.8	5.3-6.2
2000	2,803	14.5	14.0-15.1	1,532	18.9	17.9-19.9	1,271	11.4	10.8-12.0	1,396	7.2	6.8-7.6	738	9.3	8.7-10.1	658	5.7	5.3-6.2
2001	2,766	14.2	13.7-14.7	1,492	18.1	17.2-19.1	1,274	11.4	10.8-12.0	1,391	7.1	6.7-7.5	738	9.3	8.7-10.0	653	5.6	5.2-6.1
2002	2,648	13.5	12.9-14.0	1,499	17.9	17.0-18.8	1,149	10.2	9.6-10.8	1,371	6.9	6.5-7.3	741	9.2	8.6-9.9	630	5.4	4.9-5.8
2003	2,879	14.5	14.0-15.0	1,579	18.7	17.8-19.7	1,300	11.5	10.9-12.2	1,326	6.6	6.3-7.0	729	9.0	8.3-9.7	597	5.0	4.6-5.5
2004	2,921	14.6	14.1-15.1	1,618	19.0	18.0-19.9	1,303	11.5	10.8-12.1	1,435	7.1	6.7-7.5	789	9.7	9.0-10.4	646	5.4	5.0-5.9
2005	2,948	14.6	14.1-15.2	1,627	18.7	17.8-19.7	1,321	11.5	10.9-12.1	1,404	6.9	6.5-7.3	792	9.5	8.9-10.2	612	5.1	4.7-5.5
2006	2,968	14.7	14.2-15.2	1,699	19.6	18.7-20.6	1,269	11.1	10.4-11.7	1,387	6.8	6.4-7.2	762	9.1	8.4-9.7	625	5.1	4.7-5.6
2007	3,075	15.0	14.5-15.6	1,740	19.6	18.6-20.5	1,335	11.7	11.1-12.4	1,372	6.7	6.3-7.0	776	9.2	8.5-9.8	596	5.0	4.6-5.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State
Leukemias**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	3,020	14.6	14.1-15.2	1,696	18.9	18.0-19.8	1,324	11.5	10.9-12.2	1,399	6.7	6.3-7.0	777	9.0	8.3-9.6	622	5.1	4.7-5.5
2009	3,223	15.4	14.8-15.9	1,843	20.2	19.2-21.1	1,380	11.8	11.1-12.4	1,387	6.6	6.3-7.0	782	8.9	8.3-9.6	605	4.9	4.5-5.3
2010	3,456	16.3	15.8-16.9	1,985	21.5	20.5-22.4	1,471	12.4	11.8-13.1	1,338	6.2	5.9-6.6	751	8.5	7.9-9.1	587	4.7	4.3-5.1
2011	3,649	17.0	16.4-17.5	2,076	22.1	21.1-23.1	1,573	13.1	12.4-13.7	1,440	6.6	6.2-6.9	816	8.9	8.3-9.6	624	4.9	4.5-5.3
2012	3,492	16.0	15.4-16.5	2,011	20.9	20.0-21.8	1,481	12.3	11.7-13.0	1,414	6.4	6.1-6.7	821	9.0	8.4-9.6	593	4.5	4.2-4.9
2013	3,779	17.1	16.5-17.7	2,158	22.2	21.2-23.1	1,621	13.3	12.6-14.0	1,425	6.3	6.0-6.7	826	8.8	8.2-9.4	599	4.6	4.2-5.0
2014	3,866	17.2	16.6-17.7	2,232	22.4	21.5-23.4	1,634	13.2	12.5-13.8	1,374	6.0	5.6-6.3	750	7.9	7.3-8.4	624	4.7	4.3-5.1
2015	3,792	16.6	16.1-17.2	2,180	21.6	20.7-22.6	1,612	12.7	12.1-13.4	1,415	6.0	5.7-6.4	802	8.2	7.6-8.8	613	4.5	4.2-4.9
2016	3,935	16.9	16.4-17.5	2,312	22.4	21.4-23.3	1,623	12.6	12.0-13.2	1,388	5.9	5.6-6.2	759	7.6	7.1-8.2	629	4.7	4.3-5.1
2017	3,684	15.8	15.2-16.3	2,134	20.4	19.6-21.3	1,550	12.2	11.5-12.8	1,358	5.6	5.3-5.9	748	7.4	6.8-7.9	610	4.4	4.0-4.7
2018	3,592	15.3	14.8-15.9	2,089	19.9	19.0-20.8	1,503	11.7	11.1-12.4	1,370	5.6	5.3-5.9	775	7.5	7.0-8.1	595	4.3	3.9-4.6
2019	3,564	14.9	14.4-15.4	2,068	19.3	18.5-20.2	1,496	11.4	10.8-12.0	1,341	5.4	5.1-5.7	775	7.4	6.9-7.9	566	4.0	3.6-4.3
2020	3,286	13.9	13.4-14.4	1,928	18.0	17.2-18.8	1,358	10.5	10.0-11.2	1,260	5.0	4.7-5.3	742	7.0	6.5-7.5	518	3.6	3.3-3.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
All Invasive Malignant Tumors**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	28,475	388.3	383.7-393.0	13,594	456.0	447.9-464.1	14,881	351.9	346.1-357.8	16,436	227.0	223.4-230.6	8,458	290.3	283.8-297.0	7,978	187.9	183.6-192.2
1977	28,244	388.9	384.2-393.6	13,553	460.7	452.6-469.0	14,691	350.1	344.3-356.0	16,059	222.0	218.5-225.6	8,241	285.1	278.6-291.6	7,818	183.6	179.4-187.8
1978	28,355	393.4	388.7-398.2	13,657	468.8	460.6-477.0	14,698	352.5	346.7-358.4	15,602	217.0	213.5-220.5	7,966	277.5	271.1-283.9	7,636	180.4	176.3-184.6
1979	28,909	402.6	397.9-407.4	13,932	482.6	474.2-491.0	14,977	359.8	353.9-365.8	15,498	216.9	213.4-220.4	7,928	279.1	272.7-285.6	7,570	179.4	175.3-183.6
1980	29,773	415.5	410.7-420.4	14,525	505.9	497.4-514.5	15,248	366.8	360.8-372.8	15,904	222.3	218.8-225.9	8,085	288.3	281.8-295.0	7,819	183.5	179.4-187.7
1981	29,231	408.4	403.7-413.3	14,030	490.4	482.0-498.9	15,201	364.0	358.1-370.0	15,478	216.7	213.2-220.2	7,727	275.3	268.9-281.7	7,751	182.1	177.9-186.3
1982	29,670	415.0	410.1-419.8	14,144	496.8	488.4-505.4	15,526	371.4	365.4-377.5	15,175	212.3	208.9-215.8	7,659	274.2	267.9-280.7	7,516	176.4	172.3-180.5
1983	29,871	415.1	410.3-419.9	14,310	498.6	490.2-507.1	15,561	370.3	364.3-376.3	15,281	213.0	209.5-216.4	7,656	273.7	267.4-280.1	7,625	177.2	173.1-181.3
1984	30,489	423.0	418.2-427.9	14,624	508.0	499.5-516.6	15,865	377.6	371.6-383.7	15,358	213.3	209.9-216.8	7,740	275.8	269.5-282.2	7,618	176.9	172.8-181.0
1985	30,897	427.2	422.4-432.1	15,069	520.5	511.9-529.1	15,828	374.7	368.8-380.7	15,177	209.8	206.5-213.3	7,706	272.8	266.6-279.2	7,471	173.1	169.1-177.1
1986	31,033	429.0	424.1-433.8	15,146	524.3	515.7-533.0	15,887	376.2	370.2-382.2	15,051	208.3	204.9-211.7	7,440	264.2	258.0-270.4	7,611	176.0	172.0-180.0
1987	30,684	423.2	418.4-428.0	14,887	513.5	505.0-522.0	15,797	372.2	366.3-378.2	15,019	207.7	204.3-211.1	7,542	268.6	262.4-274.9	7,477	172.3	168.3-176.3
1988	30,118	417.4	412.7-422.2	14,299	497.5	489.2-506.0	15,819	373.8	367.9-379.8	15,088	210.0	206.6-213.4	7,539	271.7	265.5-278.1	7,549	174.2	170.3-178.3
1989	30,270	422.2	417.4-427.0	14,604	507.5	499.1-516.0	15,666	374.0	368.1-380.0	14,589	204.0	200.7-207.4	7,220	261.4	255.3-267.7	7,369	170.9	167.0-174.9
1990	31,593	442.0	437.1-447.0	15,184	530.4	521.8-539.1	16,409	393.4	387.3-399.6	14,867	209.4	206.0-212.8	7,276	266.1	259.9-272.5	7,591	177.4	173.4-181.5
1991	33,381	468.6	463.5-473.7	16,130	564.8	555.9-573.8	17,251	415.1	408.8-421.4	14,646	206.5	203.2-209.9	7,194	262.3	256.2-268.6	7,452	174.7	170.7-178.7
1992	33,944	474.9	469.8-480.0	16,799	584.6	575.5-593.7	17,145	412.6	406.3-418.9	14,557	205.2	201.9-208.6	7,127	260.8	254.6-267.0	7,430	174.1	170.1-178.1
1993	32,978	459.8	454.8-464.8	16,433	567.9	559.0-576.8	16,545	396.4	390.3-402.5	14,466	203.3	200.0-206.7	7,145	260.0	253.9-266.3	7,321	171.3	167.4-175.3
1994	35,360	492.3	487.1-497.5	17,319	593.3	584.3-602.4	18,041	431.1	424.8-437.5	14,586	204.6	201.3-208.0	7,176	260.3	254.2-266.5	7,410	172.8	168.8-176.8
1995	33,313	461.9	456.9-466.9	16,213	553.8	545.2-562.6	17,100	406.5	400.4-412.7	14,436	202.1	198.8-205.4	6,927	250.2	244.3-256.3	7,509	174.4	170.4-178.4
1996	33,855	468.1	463.1-473.1	16,312	555.8	547.1-564.5	17,543	416.5	410.3-422.8	14,067	196.3	193.1-199.6	6,723	241.1	235.3-247.1	7,344	170.4	166.5-174.4
1997	34,053	468.2	463.2-473.2	16,404	556.5	547.9-565.2	17,649	415.2	409.1-421.4	13,608	188.8	185.6-192.0	6,545	233.7	227.9-239.5	7,063	162.9	159.1-166.7
1998	34,725	471.8	466.8-476.8	16,726	556.3	547.7-564.9	17,999	420.7	414.6-426.9	13,521	185.4	182.3-188.6	6,468	225.8	220.3-231.5	7,053	161.3	157.6-165.1
1999	34,711	463.0	458.2-467.9	16,816	550.0	541.6-558.5	17,895	410.6	404.6-416.7	13,481	181.4	178.3-184.5	6,476	222.8	217.4-228.4	7,005	157.5	153.8-161.2
2000	35,836	470.5	465.6-475.4	17,748	571.0	562.5-579.6	18,088	408.9	402.9-414.9	13,300	175.9	172.9-178.9	6,368	215.4	210.0-220.8	6,932	152.4	148.8-156.0
2001	35,216	456.9	452.2-461.7	17,452	553.3	545.0-561.7	17,764	397.1	391.3-403.1	13,261	173.3	170.4-176.3	6,391	213.1	207.8-218.5	6,870	149.6	146.1-153.2
2002	35,108	450.0	445.3-454.7	17,422	542.0	533.9-550.2	17,686	391.6	385.8-397.4	12,870	166.2	163.4-169.1	6,252	205.3	200.1-210.5	6,618	142.7	139.3-146.2
2003	35,149	447.2	442.6-452.0	17,293	535.2	527.1-543.3	17,856	392.8	387.1-398.7	12,899	165.3	162.4-168.2	6,151	199.4	194.4-204.5	6,748	144.6	141.1-148.1
2004	35,601	449.3	444.6-454.0	17,496	533.0	525.0-541.1	18,105	396.7	390.9-402.5	12,640	160.9	158.1-163.7	6,124	197.0	192.0-202.0	6,516	138.9	135.5-142.4
2005	36,158	453.5	448.8-458.2	17,866	538.5	530.5-546.6	18,292	400.0	394.1-405.8	12,477	157.8	155.1-160.6	5,987	190.5	185.6-195.4	6,490	138.2	134.8-141.6
2006	37,131	464.8	460.0-469.6	18,513	555.9	547.8-564.1	18,618	406.7	400.9-412.7	12,283	155.3	152.5-158.1	5,885	187.2	182.4-192.2	6,398	136.1	132.7-139.5
2007	37,865	469.3	464.6-474.1	19,011	561.8	553.7-570.0	18,854	410.0	404.2-416.0	12,450	155.5	152.8-158.3	6,045	189.4	184.5-194.3	6,405	135.1	131.7-138.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
All Invasive Malignant Tumors**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	38,243	467.7	462.9-472.4	18,977	550.9	542.9-558.9	19,266	414.3	408.4-420.3	12,238	151.1	148.4-153.8	5,993	185.0	180.3-189.8	6,245	130.2	126.9-133.5
2009	38,899	469.5	464.8-474.2	19,168	546.5	538.7-554.5	19,731	419.9	414.0-425.9	12,276	149.7	147.1-152.4	6,019	183.4	178.7-188.1	6,257	128.8	125.6-132.1
2010	37,937	450.9	446.3-455.5	18,697	522.2	514.5-529.9	19,240	405.5	399.7-411.4	12,410	148.4	145.7-151.0	6,085	181.5	176.9-186.2	6,325	127.8	124.6-131.1
2011	39,686	462.0	457.4-466.6	19,574	533.5	525.9-541.2	20,112	416.4	410.6-422.3	12,476	146.3	143.8-149.0	6,046	175.8	171.3-180.4	6,430	127.7	124.5-130.9
2012	38,504	439.7	435.3-444.2	18,710	497.5	490.3-504.8	19,794	404.3	398.6-410.1	12,589	144.7	142.2-147.3	6,105	172.7	168.3-177.2	6,484	127.2	124.1-130.4
2013	40,296	451.6	447.1-456.1	19,424	505.1	497.9-512.4	20,872	419.6	413.8-425.4	12,471	140.0	137.5-142.5	6,128	169.2	164.9-173.6	6,343	121.2	118.2-124.3
2014	39,924	439.2	434.8-443.6	18,986	481.1	474.2-488.2	20,938	415.0	409.3-420.7	12,533	137.7	135.3-140.2	6,127	164.6	160.4-168.8	6,406	120.5	117.5-123.5
2015	40,383	436.4	432.1-440.8	19,334	479.5	472.7-486.5	21,049	412.0	406.4-417.8	12,200	131.5	129.1-133.9	5,913	155.0	151.0-159.1	6,287	116.5	113.6-119.5
2016	40,969	435.7	431.4-440.0	19,838	482.0	475.2-488.9	21,131	408.7	403.1-414.4	12,426	131.1	128.8-133.5	6,104	156.1	152.2-160.2	6,322	115.2	112.3-118.1
2017	41,902	440.3	436.0-444.6	20,130	480.1	473.4-486.9	21,772	417.3	411.6-423.0	12,200	126.7	124.4-129.0	5,975	150.2	146.4-154.2	6,225	111.6	108.8-114.5
2018	41,155	428.1	423.9-432.4	19,688	462.5	455.9-469.1	21,467	408.5	402.9-414.2	11,666	118.9	116.7-121.1	5,629	137.7	134.0-141.4	6,037	106.5	103.8-109.3
2019	42,833	441.2	436.9-445.5	20,598	476.3	469.7-483.0	22,235	421.3	415.6-427.0	11,919	119.7	117.6-121.9	5,800	139.5	135.9-143.2	6,119	106.5	103.8-109.3
2020	36,276	372.1	368.2-376.0	17,371	399.5	393.4-405.6	18,905	357.2	351.9-362.5	10,872	108.1	106.1-110.2	5,265	125.1	121.7-128.6	5,607	97.0	94.4-99.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Oral cavity and pharynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	857	11.7	10.9-12.5	582	19.0	17.4-20.7	275	6.4	5.7-7.3	421	5.7	5.2-6.3	299	9.7	8.6-10.9	122	2.9	2.4-3.5
1977	858	11.6	10.8-12.4	574	18.9	17.4-20.6	284	6.4	5.6-7.2	419	5.8	5.2-6.4	301	10.1	8.9-11.3	118	2.7	2.2-3.3
1978	832	11.5	10.7-12.3	561	18.3	16.8-19.9	271	6.5	5.7-7.4	340	4.6	4.1-5.2	233	7.6	6.6-8.7	107	2.5	2.0-3.1
1979	842	11.8	11.0-12.7	585	19.7	18.1-21.4	257	6.2	5.4-7.0	353	5.0	4.4-5.5	252	8.4	7.4-9.6	101	2.5	2.0-3.0
1980	833	11.9	11.0-12.7	556	18.9	17.3-20.6	277	6.9	6.1-7.8	354	5.0	4.5-5.6	258	8.9	7.8-10.1	96	2.2	1.8-2.7
1981	888	12.5	11.7-13.4	636	21.3	19.6-23.1	252	6.1	5.4-6.9	340	4.8	4.3-5.3	229	7.7	6.7-8.8	111	2.6	2.2-3.2
1982	907	12.8	12.0-13.7	617	21.0	19.3-22.7	290	7.1	6.3-8.0	343	4.9	4.4-5.5	262	9.0	7.9-10.2	81	2.0	1.6-2.5
1983	868	12.2	11.4-13.1	602	20.3	18.6-22.0	266	6.3	5.5-7.1	336	4.6	4.1-5.2	255	8.4	7.4-9.6	81	1.9	1.5-2.4
1984	897	12.7	11.9-13.6	604	20.4	18.8-22.2	293	7.1	6.3-8.0	319	4.5	4.0-5.0	234	8.0	7.0-9.1	85	2.1	1.6-2.6
1985	923	13.0	12.2-13.9	599	20.2	18.6-22.0	324	7.8	7.0-8.8	301	4.2	3.7-4.7	212	7.2	6.3-8.3	89	2.1	1.6-2.6
1986	868	12.3	11.4-13.1	590	20.0	18.4-21.7	278	6.7	5.9-7.5	295	4.1	3.6-4.6	205	6.9	6.0-8.0	90	2.1	1.7-2.6
1987	874	12.4	11.5-13.2	592	19.8	18.2-21.5	282	6.9	6.1-7.7	261	3.7	3.3-4.2	187	6.4	5.5-7.4	74	1.7	1.3-2.2
1988	786	11.1	10.3-11.9	517	17.2	15.8-18.8	269	6.4	5.6-7.2	316	4.6	4.1-5.1	213	7.4	6.5-8.5	103	2.5	2.0-3.0
1989	794	11.4	10.6-12.2	518	17.7	16.2-19.3	276	6.7	6.0-7.6	272	3.9	3.5-4.4	185	6.4	5.5-7.4	87	2.1	1.7-2.6
1990	853	12.2	11.3-13.0	567	19.4	17.8-21.1	286	6.9	6.1-7.7	249	3.6	3.2-4.1	163	5.9	5.0-6.9	86	2.0	1.6-2.5
1991	954	13.8	12.9-14.7	627	21.5	19.9-23.3	327	8.0	7.2-8.9	273	3.9	3.4-4.4	192	6.7	5.8-7.8	81	1.9	1.5-2.4
1992	970	13.8	13.0-14.7	643	21.6	20.0-23.4	327	8.1	7.2-9.0	249	3.5	3.1-4.0	164	5.7	4.8-6.6	85	2.0	1.6-2.5
1993	895	12.8	12.0-13.7	591	20.1	18.5-21.8	304	7.5	6.7-8.4	254	3.6	3.2-4.1	191	6.5	5.6-7.5	63	1.5	1.1-1.9
1994	861	12.2	11.4-13.0	565	18.8	17.3-20.5	296	7.2	6.4-8.0	241	3.4	3.0-3.9	177	6.1	5.2-7.0	64	1.5	1.2-2.0
1995	730	10.3	9.5-11.0	502	16.5	15.1-18.0	228	5.5	4.8-6.2	239	3.4	3.0-3.8	161	5.5	4.7-6.4	78	1.8	1.4-2.3
1996	819	11.4	10.6-12.2	541	17.5	16.1-19.1	278	6.7	5.9-7.5	229	3.2	2.8-3.7	148	4.8	4.1-5.7	81	1.9	1.5-2.4
1997	758	10.4	9.7-11.2	487	15.7	14.3-17.2	271	6.4	5.7-7.2	216	3.0	2.6-3.4	160	5.3	4.5-6.2	56	1.3	1.0-1.7
1998	773	10.5	9.8-11.2	493	15.5	14.2-17.0	280	6.6	5.8-7.4	203	2.8	2.4-3.2	131	4.3	3.6-5.1	72	1.6	1.3-2.1
1999	803	10.6	9.9-11.4	556	17.2	15.8-18.7	247	5.7	5.0-6.4	207	2.8	2.4-3.2	144	4.6	3.9-5.5	63	1.4	1.1-1.8
2000	725	9.5	8.8-10.2	490	14.9	13.6-16.3	235	5.4	4.7-6.1	190	2.5	2.1-2.9	128	4.0	3.3-4.8	62	1.4	1.0-1.8
2001	748	9.7	9.0-10.4	495	14.9	13.6-16.3	253	5.7	5.0-6.4	207	2.7	2.3-3.1	147	4.5	3.8-5.3	60	1.3	1.0-1.6
2002	758	9.7	9.0-10.4	491	14.6	13.3-15.9	267	5.9	5.3-6.7	174	2.2	1.9-2.6	117	3.5	2.9-4.2	57	1.2	0.9-1.6
2003	742	9.3	8.7-10.0	493	14.4	13.1-15.7	249	5.5	4.8-6.2	183	2.3	2.0-2.7	127	3.8	3.2-4.6	56	1.2	0.9-1.6
2004	789	9.9	9.2-10.6	507	14.7	13.4-16.1	282	6.2	5.5-7.0	223	2.8	2.4-3.2	148	4.4	3.7-5.2	75	1.6	1.3-2.0
2005	768	9.5	8.8-10.2	537	15.2	14.0-16.6	231	5.1	4.4-5.8	194	2.4	2.1-2.8	125	3.7	3.1-4.4	69	1.5	1.1-1.9
2006	767	9.4	8.8-10.1	501	14.2	13.0-15.6	266	5.7	5.0-6.4	161	2.0	1.7-2.3	120	3.6	2.9-4.3	41	0.9	0.6-1.2
2007	825	10.1	9.5-10.9	536	15.2	14.0-16.6	289	6.3	5.6-7.1	192	2.4	2.0-2.7	137	4.0	3.3-4.7	55	1.1	0.9-1.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Oral cavity and pharynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	818	9.9	9.3-10.7	527	14.7	13.5-16.1	291	6.3	5.6-7.0	175	2.1	1.8-2.5	125	3.6	3.0-4.3	50	1.1	0.8-1.4
2009	832	10.0	9.3-10.7	585	16.0	14.7-17.4	247	5.3	4.6-6.0	196	2.3	2.0-2.7	139	3.9	3.3-4.6	57	1.2	0.9-1.5
2010	812	9.5	8.9-10.2	518	13.9	12.7-15.2	294	6.2	5.5-6.9	203	2.4	2.1-2.8	130	3.7	3.1-4.4	73	1.5	1.2-1.9
2011	840	9.7	9.1-10.4	576	15.2	14.0-16.5	264	5.5	4.8-6.2	195	2.2	1.9-2.6	136	3.7	3.1-4.4	59	1.2	0.9-1.5
2012	860	9.6	9.0-10.3	562	14.3	13.1-15.5	298	6.0	5.3-6.7	193	2.2	1.9-2.5	125	3.3	2.7-3.9	68	1.4	1.1-1.7
2013	847	9.4	8.8-10.1	584	14.6	13.5-15.9	263	5.2	4.6-5.9	203	2.2	1.9-2.6	147	3.7	3.1-4.4	56	1.1	0.8-1.4
2014	904	9.8	9.2-10.5	559	13.6	12.5-14.8	345	6.8	6.1-7.5	186	2.0	1.7-2.3	127	3.2	2.7-3.9	59	1.1	0.9-1.5
2015	928	10.0	9.4-10.7	604	14.7	13.5-16.0	324	6.3	5.6-7.1	206	2.2	1.9-2.5	149	3.7	3.1-4.4	57	1.0	0.8-1.4
2016	943	10.0	9.4-10.7	644	15.5	14.3-16.8	299	5.7	5.1-6.4	197	2.1	1.8-2.4	137	3.4	2.9-4.1	60	1.1	0.8-1.4
2017	986	10.3	9.7-11.0	646	15.3	14.1-16.5	340	6.4	5.7-7.1	201	2.1	1.8-2.4	140	3.4	2.8-4.0	61	1.1	0.8-1.4
2018	919	9.5	8.9-10.1	600	13.9	12.8-15.1	319	5.9	5.3-6.6	191	2.0	1.7-2.3	130	3.1	2.6-3.7	61	1.1	0.8-1.4
2019	961	9.9	9.2-10.5	609	13.9	12.8-15.0	352	6.6	5.9-7.3	192	1.9	1.7-2.2	136	3.2	2.7-3.8	56	0.9	0.7-1.2
2020	849	8.6	8.0-9.2	541	12.3	11.3-13.4	308	5.6	5.0-6.3	210	2.1	1.8-2.4	139	3.2	2.7-3.8	71	1.2	1.0-1.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Esophagus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	445	5.9	5.3-6.5	307	9.7	8.6-10.9	138	3.2	2.7-3.8	432	5.8	5.2-6.4	307	9.7	8.6-10.9	125	3.0	2.5-3.6
1977	434	5.9	5.4-6.5	288	9.3	8.2-10.5	146	3.5	3.0-4.2	404	5.5	4.9-6.0	266	8.8	7.7-10.0	138	3.2	2.7-3.8
1978	452	6.2	5.6-6.8	301	9.8	8.7-11.0	151	3.6	3.0-4.2	386	5.3	4.8-5.9	249	8.3	7.3-9.5	137	3.3	2.7-3.9
1979	454	6.1	5.6-6.7	316	10.4	9.2-11.6	138	3.2	2.7-3.8	414	5.7	5.1-6.3	284	9.3	8.2-10.4	130	3.2	2.6-3.8
1980	499	6.9	6.3-7.5	345	11.7	10.5-13.1	154	3.6	3.0-4.2	395	5.4	4.9-6.0	267	9.2	8.0-10.4	128	2.9	2.4-3.5
1981	467	6.5	5.9-7.2	311	10.5	9.3-11.7	156	3.8	3.2-4.4	421	5.8	5.2-6.4	296	9.9	8.8-11.1	125	2.9	2.4-3.4
1982	462	6.4	5.8-7.0	307	10.4	9.2-11.6	155	3.5	3.0-4.2	377	5.2	4.7-5.8	264	9.1	8.0-10.3	113	2.6	2.1-3.2
1983	445	6.1	5.5-6.7	306	10.1	9.0-11.3	139	3.3	2.7-3.9	374	5.1	4.6-5.7	256	8.6	7.6-9.8	118	2.8	2.3-3.3
1984	457	6.2	5.6-6.8	310	10.3	9.2-11.6	147	3.4	2.8-4.0	384	5.3	4.7-5.8	260	8.5	7.5-9.6	124	2.9	2.4-3.5
1985	436	6.1	5.5-6.7	294	10.0	8.8-11.2	142	3.4	2.9-4.0	378	5.2	4.7-5.7	277	9.2	8.1-10.4	101	2.3	1.9-2.9
1986	447	6.2	5.6-6.8	303	10.3	9.1-11.6	144	3.3	2.8-3.9	382	5.3	4.8-5.9	280	9.7	8.6-11.0	102	2.4	2.0-3.0
1987	412	5.7	5.2-6.3	255	8.6	7.6-9.8	157	3.7	3.1-4.3	342	4.7	4.2-5.3	227	7.7	6.7-8.8	115	2.7	2.2-3.2
1988	399	5.6	5.1-6.2	267	9.2	8.1-10.4	132	3.0	2.5-3.6	362	5.1	4.6-5.6	264	9.3	8.2-10.5	98	2.2	1.8-2.7
1989	447	6.3	5.7-6.9	299	10.4	9.2-11.6	148	3.4	2.9-4.1	354	5.0	4.5-5.5	237	8.1	7.1-9.2	117	2.8	2.3-3.3
1990	418	5.9	5.4-6.5	269	9.3	8.2-10.5	149	3.5	3.0-4.2	356	5.0	4.5-5.6	232	8.2	7.1-9.3	124	2.8	2.4-3.4
1991	395	5.6	5.0-6.2	263	9.2	8.1-10.4	132	3.1	2.6-3.7	322	4.5	4.0-5.0	214	7.4	6.5-8.5	108	2.5	2.0-3.0
1992	416	5.9	5.3-6.5	269	9.3	8.2-10.5	147	3.5	2.9-4.1	327	4.6	4.2-5.2	230	8.0	7.0-9.1	97	2.3	1.9-2.8
1993	405	5.7	5.2-6.3	280	9.6	8.5-10.8	125	3.0	2.5-3.5	336	4.7	4.2-5.3	238	8.3	7.3-9.5	98	2.4	1.9-2.9
1994	440	6.2	5.6-6.8	304	10.5	9.3-11.8	136	3.2	2.7-3.8	336	4.7	4.2-5.2	236	8.2	7.1-9.3	100	2.3	1.9-2.9
1995	379	5.3	4.8-5.9	234	7.9	6.9-9.0	145	3.4	2.9-4.0	317	4.4	4.0-5.0	220	7.6	6.6-8.6	97	2.2	1.8-2.7
1996	379	5.3	4.8-5.8	240	8.2	7.2-9.3	139	3.2	2.7-3.8	327	4.6	4.1-5.1	217	7.4	6.5-8.5	110	2.5	2.1-3.0
1997	376	5.2	4.7-5.8	248	8.4	7.4-9.5	128	2.9	2.5-3.5	298	4.1	3.7-4.6	205	7.0	6.1-8.1	93	2.1	1.7-2.6
1998	384	5.2	4.7-5.8	266	9.0	7.9-10.1	118	2.7	2.2-3.3	305	4.2	3.7-4.7	206	7.0	6.0-8.0	99	2.3	1.8-2.7
1999	364	4.9	4.4-5.4	266	8.6	7.6-9.8	98	2.2	1.8-2.7	293	3.9	3.5-4.4	218	7.2	6.2-8.2	75	1.7	1.3-2.1
2000	351	4.7	4.2-5.2	215	7.0	6.1-8.0	136	3.0	2.5-3.5	322	4.3	3.8-4.7	208	6.8	5.9-7.8	114	2.5	2.0-3.0
2001	351	4.6	4.1-5.1	246	7.8	6.9-8.9	105	2.3	1.9-2.8	267	3.5	3.1-3.9	205	6.6	5.7-7.6	62	1.4	1.0-1.8
2002	356	4.6	4.1-5.1	244	7.6	6.6-8.6	112	2.4	2.0-2.9	269	3.5	3.1-3.9	182	5.9	5.1-6.8	87	1.8	1.5-2.3
2003	368	4.7	4.2-5.2	241	7.5	6.5-8.5	127	2.7	2.3-3.2	284	3.6	3.2-4.1	187	5.7	4.9-6.6	97	2.1	1.7-2.5
2004	362	4.5	4.1-5.0	241	7.2	6.3-8.2	121	2.6	2.1-3.1	276	3.5	3.1-3.9	187	5.8	5.0-6.7	89	1.9	1.5-2.3
2005	358	4.5	4.1-5.0	224	6.7	5.9-7.7	134	2.9	2.4-3.4	296	3.7	3.3-4.1	208	6.3	5.5-7.2	88	1.8	1.5-2.3
2006	340	4.3	3.9-4.8	250	7.7	6.8-8.8	90	1.9	1.6-2.4	285	3.6	3.2-4.0	172	5.2	4.5-6.1	113	2.4	2.0-2.9
2007	327	4.1	3.6-4.5	223	6.6	5.7-7.5	104	2.2	1.8-2.7	276	3.4	3.0-3.9	204	6.2	5.3-7.1	72	1.5	1.2-1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Esophagus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	379	4.6	4.2-5.1	262	7.6	6.7-8.6	117	2.4	2.0-2.9	273	3.3	3.0-3.8	189	5.6	4.8-6.4	84	1.8	1.4-2.2
2009	330	4.0	3.5-4.4	239	6.7	5.9-7.7	91	1.9	1.5-2.3	280	3.4	3.0-3.8	208	6.1	5.3-7.0	72	1.4	1.1-1.8
2010	291	3.4	3.0-3.9	198	5.6	4.8-6.4	93	1.9	1.5-2.3	242	2.8	2.5-3.2	170	4.9	4.2-5.7	72	1.4	1.1-1.8
2011	329	3.8	3.4-4.3	248	6.9	6.0-7.8	81	1.6	1.3-2.1	242	2.8	2.5-3.2	175	4.9	4.2-5.7	67	1.3	1.0-1.7
2012	285	3.2	2.8-3.6	199	5.3	4.6-6.1	86	1.6	1.3-2.0	207	2.4	2.0-2.7	148	4.0	3.4-4.8	59	1.1	0.8-1.4
2013	333	3.8	3.4-4.2	238	6.4	5.6-7.2	95	1.9	1.5-2.3	230	2.6	2.2-2.9	167	4.3	3.7-5.0	63	1.2	0.9-1.6
2014	348	3.8	3.4-4.2	250	6.4	5.6-7.3	98	1.8	1.5-2.3	267	3.0	2.6-3.4	192	5.2	4.4-6.0	75	1.4	1.1-1.8
2015	312	3.3	2.9-3.7	211	5.2	4.5-6.0	101	1.9	1.5-2.3	228	2.5	2.2-2.8	152	3.9	3.3-4.6	76	1.3	1.1-1.7
2016	343	3.5	3.2-3.9	252	6.1	5.3-6.9	91	1.6	1.3-2.0	228	2.4	2.1-2.7	166	4.1	3.5-4.7	62	1.1	0.8-1.4
2017	301	3.1	2.8-3.5	216	5.2	4.5-5.9	85	1.6	1.2-1.9	228	2.3	2.0-2.7	171	4.1	3.5-4.8	57	1.0	0.8-1.3
2018	317	3.2	2.8-3.5	232	5.4	4.7-6.2	85	1.4	1.2-1.8	202	2.0	1.7-2.3	152	3.6	3.0-4.2	50	0.9	0.6-1.2
2019	330	3.3	2.9-3.6	237	5.4	4.7-6.1	93	1.6	1.3-2.0	212	2.1	1.8-2.4	159	3.7	3.2-4.4	53	0.9	0.6-1.1
2020	289	2.8	2.5-3.2	192	4.4	3.8-5.0	97	1.7	1.3-2.1	195	1.9	1.6-2.2	138	3.1	2.6-3.7	57	1.0	0.7-1.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Stomach**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,063	14.9	14.0-15.9	609	21.2	19.4-23.1	454	10.7	9.7-11.8	853	12.1	11.3-13.0	487	17.3	15.7-19.0	366	8.7	7.8-9.7
1977	1,078	15.4	14.5-16.4	611	21.9	20.1-23.8	467	11.2	10.2-12.3	828	12.0	11.1-12.8	458	16.9	15.3-18.6	370	8.8	7.9-9.8
1978	1,028	14.5	13.6-15.5	567	20.2	18.5-22.0	461	10.8	9.9-11.9	774	11.1	10.3-11.9	424	15.4	13.9-17.0	350	8.3	7.5-9.3
1979	1,053	14.9	14.0-15.8	580	20.5	18.8-22.3	473	11.2	10.2-12.3	747	10.6	9.9-11.4	412	14.9	13.4-16.5	335	7.9	7.1-8.8
1980	1,115	15.8	14.9-16.8	640	23.0	21.2-24.9	475	11.3	10.3-12.4	769	11.0	10.2-11.8	438	16.0	14.5-17.6	331	7.8	7.0-8.7
1981	1,056	14.8	13.9-15.8	599	21.5	19.8-23.4	457	10.6	9.7-11.7	778	11.0	10.2-11.8	461	16.9	15.3-18.5	317	7.4	6.6-8.3
1982	1,008	14.1	13.2-15.0	554	19.8	18.2-21.6	454	10.5	9.6-11.6	675	9.6	8.9-10.4	385	14.5	13.1-16.1	290	6.7	5.9-7.5
1983	1,085	15.3	14.4-16.3	563	20.3	18.6-22.1	522	12.1	11.0-13.2	709	10.0	9.2-10.7	390	14.4	12.9-15.9	319	7.3	6.5-8.1
1984	973	13.4	12.6-14.3	531	18.7	17.1-20.4	442	10.1	9.1-11.1	654	9.2	8.5-9.9	340	12.3	11.0-13.8	314	7.2	6.4-8.1
1985	980	13.8	12.9-14.7	545	19.6	18.0-21.4	435	10.0	9.1-11.0	669	9.3	8.6-10.1	375	13.5	12.1-14.9	294	6.7	6.0-7.6
1986	977	13.6	12.8-14.5	540	19.3	17.7-21.1	437	10.0	9.1-11.0	613	8.5	7.9-9.3	345	12.6	11.3-14.1	268	6.0	5.3-6.8
1987	994	13.7	12.9-14.6	572	20.1	18.4-21.8	422	9.4	8.5-10.4	631	8.8	8.1-9.5	356	12.8	11.4-14.2	275	6.2	5.4-6.9
1988	938	13.1	12.2-13.9	526	18.7	17.1-20.5	412	9.3	8.5-10.3	639	8.9	8.2-9.6	348	12.4	11.1-13.8	291	6.6	5.9-7.4
1989	905	12.7	11.9-13.5	488	17.2	15.7-18.9	417	9.5	8.6-10.4	635	8.9	8.2-9.6	357	12.9	11.6-14.4	278	6.3	5.5-7.1
1990	1,017	14.4	13.5-15.3	560	20.4	18.7-22.2	457	10.6	9.6-11.6	609	8.7	8.0-9.4	338	12.5	11.1-13.9	271	6.2	5.5-7.0
1991	1,009	14.4	13.5-15.3	581	21.2	19.5-23.0	428	9.9	9.0-10.9	625	8.9	8.2-9.6	358	13.3	12.0-14.8	267	6.1	5.4-6.9
1992	992	14.0	13.1-14.9	544	19.5	17.9-21.3	448	10.3	9.4-11.3	662	9.4	8.7-10.2	353	13.1	11.8-14.6	309	7.1	6.3-7.9
1993	1,006	14.1	13.3-15.0	566	20.6	18.9-22.4	440	10.1	9.2-11.1	622	8.8	8.1-9.5	337	12.5	11.2-13.9	285	6.6	5.8-7.4
1994	1,067	15.0	14.1-15.9	563	20.0	18.4-21.8	504	11.6	10.6-12.7	666	9.4	8.7-10.1	379	13.9	12.5-15.4	287	6.6	5.8-7.4
1995	959	13.4	12.6-14.3	530	18.9	17.3-20.6	429	9.7	8.8-10.7	623	8.8	8.1-9.5	329	12.1	10.8-13.5	294	6.7	5.9-7.5
1996	1,005	14.0	13.2-14.9	551	19.6	18.0-21.3	454	10.3	9.4-11.3	628	8.8	8.1-9.5	330	12.0	10.7-13.4	298	6.8	6.0-7.6
1997	955	13.2	12.3-14.0	542	19.2	17.6-20.9	413	9.3	8.5-10.3	579	8.0	7.4-8.7	311	11.0	9.8-12.3	268	6.1	5.4-6.9
1998	1,033	14.2	13.3-15.0	565	19.6	18.0-21.3	468	10.7	9.7-11.7	586	8.0	7.4-8.7	316	11.2	9.9-12.5	270	6.0	5.3-6.8
1999	959	12.9	12.1-13.8	552	18.8	17.2-20.4	407	9.0	8.2-9.9	551	7.4	6.8-8.1	294	10.3	9.2-11.6	257	5.7	5.0-6.4
2000	935	12.3	11.6-13.2	496	16.6	15.1-18.1	439	9.6	8.7-10.5	520	6.8	6.3-7.5	276	9.3	8.2-10.5	244	5.2	4.6-5.9
2001	957	12.5	11.7-13.3	528	17.2	15.8-18.8	429	9.3	8.4-10.2	503	6.6	6.0-7.2	271	8.9	7.8-10.0	232	5.0	4.3-5.7
2002	900	11.6	10.8-12.4	495	15.8	14.5-17.3	405	8.7	7.9-9.6	480	6.2	5.6-6.8	266	8.7	7.7-9.8	214	4.5	4.0-5.2
2003	981	12.5	11.7-13.3	541	17.1	15.6-18.6	440	9.4	8.5-10.3	520	6.6	6.1-7.2	274	8.9	7.8-10.0	246	5.2	4.6-5.9
2004	948	12.1	11.3-12.9	531	16.8	15.3-18.3	417	8.9	8.1-9.8	520	6.6	6.0-7.2	263	8.5	7.5-9.6	257	5.5	4.8-6.2
2005	1,001	12.7	11.9-13.5	560	17.7	16.2-19.2	441	9.3	8.5-10.3	513	6.5	5.9-7.1	293	9.4	8.4-10.6	220	4.5	4.0-5.2
2006	941	11.9	11.1-12.7	530	16.4	15.0-17.8	411	8.8	7.9-9.7	470	5.9	5.4-6.5	260	8.3	7.3-9.4	210	4.4	3.8-5.1
2007	964	12.0	11.3-12.8	546	16.6	15.3-18.1	418	8.9	8.0-9.8	435	5.4	4.9-6.0	232	7.2	6.3-8.2	203	4.2	3.6-4.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Stomach**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	894	11.0	10.3-11.8	524	15.8	14.5-17.3	370	7.8	7.0-8.6	438	5.4	4.9-5.9	260	8.0	7.0-9.0	178	3.7	3.2-4.3
2009	937	11.4	10.7-12.2	551	16.2	14.9-17.7	386	8.0	7.2-8.9	449	5.5	5.0-6.0	254	7.8	6.9-8.9	195	3.9	3.4-4.5
2010	939	11.3	10.5-12.0	527	15.1	13.8-16.5	412	8.4	7.6-9.3	412	4.9	4.4-5.4	208	6.3	5.5-7.2	204	4.0	3.5-4.6
2011	892	10.5	9.8-11.2	497	14.1	12.9-15.5	395	7.9	7.1-8.8	439	5.2	4.7-5.7	241	7.0	6.2-8.0	198	3.9	3.3-4.5
2012	967	11.2	10.5-11.9	556	15.4	14.1-16.7	411	8.2	7.4-9.1	455	5.2	4.8-5.8	253	7.2	6.3-8.1	202	3.9	3.4-4.5
2013	1,028	11.6	10.9-12.3	598	16.1	14.8-17.5	430	8.3	7.6-9.2	433	4.8	4.4-5.3	258	7.1	6.2-8.0	175	3.3	2.8-3.8
2014	968	10.8	10.1-11.5	576	15.1	13.9-16.5	392	7.5	6.8-8.4	465	5.2	4.7-5.7	269	7.2	6.3-8.1	196	3.8	3.3-4.4
2015	996	10.7	10.1-11.4	550	13.8	12.7-15.1	446	8.4	7.7-9.3	401	4.4	3.9-4.8	224	6.0	5.2-6.9	177	3.2	2.8-3.8
2016	1,023	10.9	10.2-11.6	610	15.1	13.9-16.4	413	7.8	7.1-8.6	470	5.0	4.5-5.4	307	7.8	6.9-8.7	163	2.9	2.5-3.4
2017	1,013	10.7	10.0-11.4	562	13.8	12.6-15.0	451	8.4	7.6-9.2	428	4.5	4.1-5.0	244	6.1	5.4-7.0	184	3.4	2.9-3.9
2018	921	9.6	8.9-10.2	527	12.6	11.5-13.7	394	7.3	6.6-8.1	386	4.0	3.6-4.4	219	5.4	4.7-6.1	167	3.0	2.5-3.5
2019	1,011	10.5	9.8-11.2	576	13.6	12.5-14.8	435	8.2	7.4-9.0	443	4.6	4.1-5.0	264	6.4	5.6-7.2	179	3.2	2.8-3.8
2020	836	8.6	8.0-9.2	474	11.1	10.1-12.2	362	6.6	5.9-7.3	356	3.6	3.2-4.0	218	5.1	4.5-5.9	138	2.5	2.1-2.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Colorectal**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	4,731	65.1	63.2-67.0	2,276	78.1	74.8-81.6	2,455	56.9	54.6-59.2	2,618	37.1	35.6-38.6	1,316	47.5	44.8-50.3	1,302	30.7	29.0-32.5
1977	4,755	65.9	64.0-67.9	2,302	80.8	77.4-84.4	2,453	56.8	54.5-59.1	2,428	34.3	32.9-35.7	1,192	43.5	41.0-46.2	1,236	28.8	27.2-30.5
1978	4,653	64.5	62.6-66.5	2,246	79.3	75.9-82.8	2,407	55.6	53.4-57.9	2,366	33.7	32.3-35.1	1,186	43.4	40.9-46.1	1,180	27.8	26.2-29.5
1979	4,768	66.4	64.5-68.3	2,300	81.7	78.2-85.3	2,468	57.1	54.8-59.4	2,359	33.4	32.0-34.8	1,119	40.9	38.4-43.4	1,240	28.8	27.2-30.5
1980	4,945	68.7	66.7-70.7	2,435	86.4	82.9-90.1	2,510	57.9	55.6-60.2	2,403	34.3	32.9-35.8	1,195	45.0	42.4-47.8	1,208	28.1	26.5-29.8
1981	4,833	67.6	65.7-69.6	2,312	83.7	80.2-87.3	2,521	58.0	55.7-60.3	2,228	31.7	30.3-33.0	1,019	38.2	35.8-40.8	1,209	27.9	26.3-29.5
1982	4,862	67.8	65.8-69.7	2,326	84.0	80.5-87.6	2,536	58.0	55.8-60.4	2,143	30.3	29.0-31.6	1,056	39.1	36.7-41.6	1,087	25.2	23.7-26.7
1983	4,744	65.6	63.7-67.5	2,280	81.4	78.0-85.0	2,464	56.3	54.0-58.6	2,164	30.5	29.2-31.9	1,036	38.3	35.9-40.8	1,128	25.8	24.3-27.4
1984	4,687	65.0	63.1-66.9	2,226	79.6	76.2-83.0	2,461	56.1	53.9-58.4	2,216	30.9	29.6-32.3	1,098	40.6	38.1-43.1	1,118	25.5	24.0-27.0
1985	4,774	65.5	63.6-67.4	2,350	83.7	80.2-87.2	2,424	54.8	52.6-57.1	2,101	29.2	27.9-30.5	1,009	36.6	34.3-39.0	1,092	24.7	23.3-26.3
1986	4,857	66.9	65.0-68.8	2,394	85.5	82.0-89.1	2,463	55.8	53.6-58.1	2,057	28.6	27.4-29.9	1,008	37.0	34.7-39.4	1,049	23.8	22.4-25.3
1987	4,377	60.1	58.3-61.9	2,136	76.0	72.7-79.4	2,241	50.2	48.2-52.4	2,034	28.2	27.0-29.5	971	35.8	33.5-38.2	1,063	23.8	22.4-25.3
1988	4,276	59.1	57.3-60.9	2,001	71.8	68.6-75.1	2,275	51.2	49.1-53.4	1,940	27.1	25.9-28.4	978	36.6	34.2-39.0	962	21.7	20.3-23.1
1989	4,103	57.3	55.5-59.1	1,990	71.6	68.4-74.9	2,113	48.4	46.3-50.5	1,836	25.8	24.6-27.0	859	32.3	30.1-34.5	977	22.0	20.6-23.4
1990	4,136	57.8	56.0-59.6	1,900	68.3	65.2-71.5	2,236	50.9	48.7-53.0	1,879	26.6	25.4-27.8	886	33.3	31.1-35.6	993	22.6	21.2-24.0
1991	4,236	59.7	57.9-61.5	2,052	75.8	72.4-79.2	2,184	50.1	48.0-52.3	1,814	25.7	24.5-26.9	868	33.3	31.1-35.7	946	21.4	20.0-22.8
1992	4,070	57.3	55.6-59.1	1,859	68.0	64.9-71.2	2,211	51.1	48.9-53.3	1,746	24.7	23.5-25.9	817	30.9	28.7-33.1	929	21.0	19.7-22.5
1993	4,099	57.5	55.8-59.3	1,945	70.2	67.1-73.5	2,154	49.5	47.4-51.6	1,697	23.9	22.7-25.0	805	30.0	27.9-32.1	892	20.2	18.9-21.6
1994	4,491	62.9	61.1-64.8	2,069	74.4	71.2-77.7	2,422	55.6	53.3-57.8	1,794	25.2	24.0-26.4	867	32.5	30.3-34.7	927	20.9	19.6-22.3
1995	4,212	58.8	57.1-60.6	1,958	70.2	67.1-73.5	2,254	51.6	49.5-53.8	1,831	25.7	24.6-26.9	861	32.1	29.9-34.3	970	21.9	20.6-23.4
1996	4,149	57.9	56.2-59.7	1,969	70.0	66.9-73.2	2,180	50.1	48.0-52.3	1,625	22.7	21.6-23.8	786	28.8	26.8-30.9	839	18.9	17.6-20.2
1997	4,358	60.5	58.7-62.3	2,016	71.2	68.1-74.4	2,342	53.3	51.2-55.5	1,740	24.2	23.1-25.4	845	30.9	28.8-33.1	895	20.1	18.8-21.4
1998	4,397	60.3	58.6-62.1	2,053	71.5	68.4-74.7	2,344	53.2	51.0-55.4	1,631	22.5	21.4-23.6	733	26.2	24.3-28.2	898	20.0	18.7-21.4
1999	4,348	58.5	56.8-60.3	2,012	68.3	65.3-71.4	2,336	52.3	50.1-54.4	1,623	21.9	20.9-23.0	744	26.1	24.3-28.1	879	19.4	18.1-20.7
2000	4,362	57.7	56.0-59.4	2,103	69.9	66.9-73.0	2,259	49.5	47.5-51.6	1,607	21.3	20.2-22.3	769	26.4	24.5-28.3	838	18.0	16.7-19.2
2001	4,277	55.9	54.3-57.6	2,047	67.7	64.7-70.7	2,230	48.6	46.6-50.6	1,537	20.1	19.1-21.2	732	24.9	23.1-26.8	805	17.2	16.0-18.4
2002	4,188	54.1	52.5-55.8	1,998	63.8	61.0-66.7	2,190	47.2	45.3-49.2	1,528	19.8	18.8-20.8	750	25.1	23.3-26.9	778	16.4	15.3-17.6
2003	4,229	54.2	52.6-55.8	2,064	65.7	62.9-68.7	2,165	46.5	44.5-48.5	1,579	20.3	19.3-21.3	739	24.3	22.6-26.2	840	17.6	16.4-18.9
2004	4,109	52.2	50.6-53.8	1,935	60.8	58.1-63.6	2,174	46.4	44.4-48.4	1,418	18.1	17.2-19.1	685	22.2	20.6-24.0	733	15.3	14.2-16.4
2005	3,984	50.3	48.7-51.9	1,921	59.7	57.0-62.5	2,063	43.7	41.9-45.7	1,347	17.1	16.2-18.1	624	20.4	18.8-22.1	723	14.9	13.9-16.1
2006	3,985	50.2	48.6-51.8	1,889	58.6	56.0-61.4	2,096	44.6	42.7-46.6	1,400	17.7	16.8-18.7	678	22.0	20.4-23.8	722	15.1	14.0-16.3
2007	3,947	49.1	47.5-50.6	1,931	58.1	55.5-60.8	2,016	42.6	40.8-44.6	1,308	16.4	15.5-17.3	614	19.3	17.8-20.9	694	14.3	13.3-15.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Colorectal**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	3,823	47.1	45.6-48.7	1,866	55.5	52.9-58.1	1,957	41.1	39.3-43.0	1,349	16.7	15.8-17.6	664	20.8	19.2-22.4	685	14.0	13.0-15.1
2009	3,810	46.3	44.8-47.8	1,849	53.8	51.3-56.4	1,961	40.7	38.9-42.6	1,312	16.0	15.1-16.9	616	19.0	17.5-20.6	696	13.9	12.9-15.0
2010	3,634	43.5	42.1-45.0	1,798	51.6	49.2-54.1	1,836	37.9	36.1-39.7	1,314	15.7	14.8-16.6	686	20.7	19.1-22.3	628	12.4	11.4-13.4
2011	3,548	41.7	40.3-43.1	1,767	49.8	47.4-52.2	1,781	35.9	34.3-37.7	1,295	15.1	14.3-16.0	625	18.2	16.8-19.7	670	13.0	12.0-14.1
2012	3,444	39.5	38.2-40.9	1,746	47.6	45.3-49.9	1,698	33.7	32.1-35.4	1,301	14.9	14.1-15.7	623	17.6	16.3-19.1	678	12.9	11.9-13.9
2013	3,630	40.9	39.5-42.2	1,814	48.3	46.1-50.7	1,816	35.3	33.7-37.0	1,236	13.8	13.0-14.6	598	16.3	15.0-17.7	638	11.8	10.9-12.8
2014	3,606	39.9	38.5-41.2	1,820	46.8	44.6-49.0	1,786	34.6	33.0-36.3	1,204	13.2	12.4-14.0	595	16.0	14.8-17.4	609	11.2	10.3-12.1
2015	3,504	38.0	36.8-39.3	1,777	45.0	42.9-47.2	1,727	32.9	31.3-34.5	1,167	12.5	11.8-13.3	572	15.1	13.8-16.4	595	10.7	9.9-11.6
2016	3,564	38.1	36.9-39.4	1,868	46.4	44.3-48.6	1,696	32.1	30.5-33.7	1,211	12.7	12.0-13.4	596	15.2	14.0-16.5	615	10.9	10.0-11.8
2017	3,547	37.3	36.1-38.6	1,824	43.9	41.9-46.0	1,723	32.4	30.9-34.0	1,214	12.6	11.9-13.3	587	14.7	13.5-15.9	627	11.1	10.2-12.0
2018	3,431	35.8	34.6-37.0	1,760	42.2	40.2-44.2	1,671	30.7	29.3-32.3	1,061	10.9	10.3-11.6	530	13.0	11.9-14.2	531	9.4	8.6-10.2
2019	3,565	36.9	35.7-38.1	1,858	43.8	41.8-45.9	1,707	31.5	30.0-33.1	1,067	10.8	10.2-11.5	543	13.2	12.1-14.4	524	8.9	8.2-9.8
2020	3,012	31.0	29.9-32.2	1,550	36.2	34.4-38.1	1,462	26.9	25.5-28.3	1,048	10.4	9.8-11.1	546	13.0	11.9-14.1	502	8.5	7.8-9.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Colon excluding rectum**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	3,341	46.0	44.4-47.6	1,546	53.6	50.8-56.5	1,795	41.4	39.5-43.4	2,082	29.6	28.3-30.9	1,019	37.0	34.6-39.5	1,063	25.1	23.6-26.7
1977	3,380	47.1	45.5-48.8	1,574	56.1	53.2-59.1	1,806	41.8	39.9-43.9	1,986	28.0	26.8-29.3	948	34.8	32.5-37.2	1,038	24.2	22.7-25.7
1978	3,348	46.8	45.2-48.4	1,575	56.2	53.3-59.2	1,773	41.2	39.3-43.2	1,965	27.9	26.7-29.2	969	35.7	33.4-38.1	996	23.3	21.9-24.9
1979	3,478	48.7	47.0-50.4	1,626	58.5	55.5-61.5	1,852	42.9	40.9-44.9	1,970	27.8	26.6-29.1	901	32.9	30.7-35.2	1,069	24.8	23.3-26.4
1980	3,518	49.1	47.5-50.8	1,679	60.9	57.9-64.0	1,839	42.3	40.3-44.3	2,040	29.2	27.9-30.5	995	37.7	35.3-40.3	1,045	24.3	22.8-25.9
1981	3,423	48.0	46.4-49.7	1,586	58.1	55.1-61.1	1,837	42.1	40.2-44.1	1,924	27.3	26.1-28.6	862	32.3	30.1-34.6	1,062	24.5	23.0-26.1
1982	3,462	48.4	46.8-50.1	1,587	58.4	55.4-61.4	1,875	42.8	40.9-44.8	1,850	26.1	24.9-27.3	909	33.8	31.5-36.2	941	21.8	20.4-23.2
1983	3,430	47.6	46.0-49.3	1,594	57.5	54.6-60.5	1,836	41.9	40.0-43.9	1,820	25.6	24.5-26.9	861	31.8	29.6-34.1	959	21.9	20.5-23.3
1984	3,392	47.1	45.5-48.8	1,576	57.1	54.2-60.1	1,816	41.3	39.4-43.2	1,911	26.7	25.5-27.9	939	34.8	32.5-37.2	972	22.2	20.8-23.6
1985	3,546	48.9	47.3-50.6	1,667	60.3	57.4-63.4	1,879	42.5	40.6-44.5	1,841	25.6	24.4-26.8	873	31.7	29.5-33.9	968	22.0	20.6-23.4
1986	3,595	49.7	48.0-51.3	1,697	61.2	58.2-64.3	1,898	43.0	41.1-45.0	1,785	24.8	23.7-26.0	862	31.5	29.4-33.7	923	21.0	19.6-22.4
1987	3,221	44.1	42.6-45.7	1,514	54.3	51.5-57.2	1,707	38.0	36.2-39.8	1,781	24.7	23.6-25.9	861	31.8	29.6-34.0	920	20.6	19.3-22.0
1988	3,146	43.6	42.0-45.1	1,424	51.5	48.8-54.4	1,722	38.7	36.8-40.5	1,705	23.8	22.7-25.0	862	32.2	30.0-34.5	843	19.0	17.7-20.3
1989	3,025	42.3	40.8-43.9	1,427	51.9	49.2-54.8	1,598	36.6	34.8-38.5	1,639	23.0	21.9-24.1	761	28.7	26.7-30.9	878	19.7	18.4-21.1
1990	3,100	43.4	41.9-45.0	1,376	50.1	47.4-52.9	1,724	39.2	37.4-41.1	1,660	23.5	22.4-24.6	771	29.0	27.0-31.2	889	20.2	18.9-21.6
1991	3,166	44.6	43.1-46.2	1,491	55.4	52.5-58.3	1,675	38.4	36.5-40.3	1,585	22.4	21.3-23.6	752	29.0	26.9-31.2	833	18.8	17.5-20.1
1992	3,017	42.5	41.0-44.1	1,350	50.1	47.4-52.9	1,667	38.4	36.5-40.3	1,525	21.5	20.4-22.6	711	26.9	24.9-29.0	814	18.4	17.2-19.7
1993	3,051	42.8	41.3-44.4	1,392	50.6	47.9-53.4	1,659	38.0	36.2-39.9	1,481	20.9	19.8-22.0	689	25.8	23.8-27.8	792	18.0	16.7-19.3
1994	3,368	47.2	45.6-48.8	1,507	54.6	51.8-57.5	1,861	42.5	40.6-44.5	1,587	22.3	21.2-23.4	767	28.7	26.7-30.9	820	18.5	17.3-19.8
1995	3,141	44.0	42.4-45.5	1,421	51.7	49.0-54.5	1,720	39.2	37.3-41.1	1,615	22.7	21.6-23.8	761	28.5	26.5-30.6	854	19.3	18.0-20.6
1996	3,075	42.9	41.4-44.5	1,420	50.8	48.2-53.6	1,655	37.8	36.0-39.7	1,428	20.0	18.9-21.0	695	25.6	23.7-27.6	733	16.5	15.3-17.7
1997	3,275	45.5	44.0-47.1	1,476	52.7	50.0-55.5	1,799	40.8	39.0-42.8	1,511	21.0	20.0-22.1	723	26.5	24.6-28.6	788	17.7	16.5-19.0
1998	3,225	44.3	42.8-45.9	1,460	51.5	48.8-54.2	1,765	39.9	38.0-41.8	1,456	20.1	19.0-21.1	652	23.5	21.7-25.4	804	17.9	16.7-19.2
1999	3,212	43.2	41.8-44.8	1,436	49.4	46.8-52.0	1,776	39.6	37.8-41.5	1,417	19.1	18.2-20.2	641	22.7	20.9-24.5	776	17.1	15.9-18.4
2000	3,218	42.6	41.2-44.1	1,479	49.8	47.3-52.4	1,739	38.0	36.2-39.9	1,416	18.7	17.8-19.7	675	23.3	21.5-25.1	741	15.9	14.7-17.1
2001	3,139	41.1	39.7-42.6	1,468	49.0	46.5-51.6	1,671	36.1	34.4-37.9	1,332	17.5	16.5-18.4	617	21.1	19.5-22.9	715	15.2	14.1-16.4
2002	3,090	40.0	38.6-41.4	1,404	45.5	43.1-48.0	1,686	36.2	34.5-38.0	1,332	17.2	16.3-18.2	652	21.9	20.2-23.7	680	14.4	13.3-15.5
2003	3,077	39.5	38.1-40.9	1,449	46.8	44.4-49.4	1,628	34.8	33.1-36.5	1,376	17.7	16.7-18.6	635	21.0	19.4-22.8	741	15.5	14.4-16.7
2004	2,992	38.1	36.7-39.5	1,381	43.8	41.5-46.2	1,611	34.2	32.5-35.9	1,231	15.7	14.8-16.6	582	18.9	17.4-20.5	649	13.5	12.5-14.6
2005	2,950	37.3	35.9-38.6	1,358	42.6	40.3-45.0	1,592	33.6	32.0-35.3	1,172	14.9	14.1-15.8	533	17.5	16.0-19.1	639	13.2	12.2-14.3
2006	2,850	36.0	34.7-37.4	1,307	41.1	38.9-43.5	1,543	32.8	31.1-34.5	1,222	15.5	14.6-16.4	586	19.1	17.6-20.7	636	13.3	12.3-14.4
2007	2,807	35.0	33.7-36.3	1,333	40.7	38.5-43.0	1,474	31.1	29.5-32.7	1,130	14.1	13.3-15.0	525	16.5	15.1-18.0	605	12.5	11.5-13.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Colon excluding rectum**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	2,759	34.2	32.9-35.5	1,307	39.6	37.4-41.8	1,452	30.4	28.9-32.0	1,135	14.1	13.3-14.9	551	17.4	15.9-18.9	584	12.0	11.0-13.0
2009	2,746	33.5	32.2-34.8	1,263	37.6	35.5-39.7	1,483	30.6	29.0-32.2	1,128	13.8	13.0-14.6	532	16.6	15.2-18.1	596	11.9	10.9-12.9
2010	2,605	31.3	30.1-32.5	1,241	35.8	33.8-37.9	1,364	28.0	26.5-29.5	1,131	13.5	12.7-14.3	577	17.5	16.1-19.0	554	10.9	10.0-11.8
2011	2,533	29.8	28.7-31.0	1,240	35.4	33.4-37.4	1,293	25.9	24.5-27.4	1,087	12.8	12.0-13.5	518	15.2	13.9-16.6	569	11.1	10.2-12.0
2012	2,477	28.6	27.4-29.7	1,189	33.0	31.1-34.9	1,288	25.5	24.1-27.0	1,085	12.4	11.7-13.2	508	14.5	13.2-15.8	577	10.9	10.0-11.9
2013	2,581	29.1	28.0-30.3	1,243	33.6	31.7-35.5	1,338	25.9	24.5-27.3	1,046	11.7	11.0-12.4	487	13.5	12.3-14.8	559	10.3	9.5-11.2
2014	2,574	28.5	27.4-29.7	1,252	32.7	30.9-34.6	1,322	25.5	24.1-26.9	993	10.8	10.2-11.5	479	13.0	11.8-14.2	514	9.4	8.6-10.2
2015	2,414	26.2	25.2-27.3	1,172	30.2	28.4-32.0	1,242	23.5	22.2-24.8	967	10.3	9.7-11.0	469	12.3	11.2-13.5	498	9.0	8.2-9.8
2016	2,519	26.9	25.9-28.0	1,287	32.2	30.5-34.1	1,232	23.1	21.8-24.5	978	10.2	9.6-10.9	469	12.0	10.9-13.2	509	9.0	8.2-9.9
2017	2,462	25.9	24.8-26.9	1,204	29.2	27.5-30.9	1,258	23.4	22.1-24.7	982	10.1	9.5-10.8	455	11.4	10.4-12.5	527	9.3	8.5-10.1
2018	2,405	25.0	23.9-26.0	1,194	28.6	26.9-30.3	1,211	22.1	20.9-23.4	875	9.0	8.4-9.6	428	10.5	9.5-11.6	447	7.8	7.1-8.6
2019	2,543	26.2	25.2-27.2	1,274	30.2	28.5-31.9	1,269	23.1	21.8-24.4	866	8.8	8.2-9.4	422	10.3	9.4-11.4	444	7.6	6.9-8.3
2020	2,122	21.7	20.8-22.7	1,029	24.0	22.6-25.6	1,093	19.8	18.6-21.1	838	8.3	7.7-8.9	427	10.1	9.2-11.1	411	7.0	6.3-7.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Rectum & rectosigmoid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,390	19.1	18.1-20.1	730	24.6	22.7-26.5	660	15.5	14.3-16.7	536	7.5	6.9-8.2	297	10.4	9.2-11.8	239	5.6	4.9-6.4
1977	1,375	18.8	17.8-19.8	728	24.8	22.9-26.7	647	14.9	13.8-16.2	442	6.2	5.6-6.8	244	8.8	7.6-10.0	198	4.6	4.0-5.3
1978	1,305	17.8	16.8-18.8	671	23.1	21.3-25.0	634	14.4	13.3-15.6	401	5.8	5.2-6.4	217	7.7	6.7-8.9	184	4.5	3.8-5.2
1979	1,290	17.7	16.7-18.7	674	23.2	21.4-25.1	616	14.2	13.1-15.4	389	5.5	5.0-6.1	218	8.0	6.9-9.2	171	4.0	3.4-4.7
1980	1,427	19.5	18.5-20.6	756	25.6	23.7-27.5	671	15.6	14.4-16.9	363	5.1	4.6-5.7	200	7.3	6.3-8.4	163	3.8	3.2-4.5
1981	1,410	19.6	18.6-20.7	726	25.6	23.7-27.6	684	15.8	14.6-17.1	304	4.3	3.8-4.8	157	5.9	5.0-7.0	147	3.4	2.8-4.0
1982	1,400	19.3	18.3-20.4	739	25.6	23.8-27.6	661	15.2	14.1-16.4	293	4.2	3.7-4.7	147	5.3	4.4-6.2	146	3.4	2.9-4.0
1983	1,314	18.0	17.0-19.0	686	24.0	22.2-25.9	628	14.3	13.2-15.5	344	4.9	4.4-5.5	175	6.5	5.5-7.6	169	3.9	3.4-4.6
1984	1,295	17.9	16.9-18.9	650	22.5	20.7-24.3	645	14.9	13.7-16.1	305	4.3	3.8-4.8	159	5.8	4.9-6.8	146	3.3	2.8-3.9
1985	1,228	16.6	15.7-17.6	683	23.3	21.6-25.2	545	12.3	11.3-13.4	260	3.6	3.2-4.1	136	4.9	4.1-5.8	124	2.8	2.3-3.3
1986	1,262	17.2	16.2-18.2	697	24.3	22.5-26.2	565	12.8	11.8-14.0	272	3.8	3.3-4.3	146	5.5	4.6-6.5	126	2.8	2.3-3.4
1987	1,156	15.9	15.0-16.9	622	21.8	20.0-23.6	534	12.3	11.2-13.4	253	3.5	3.1-3.9	110	4.0	3.3-4.9	143	3.2	2.7-3.7
1988	1,130	15.6	14.7-16.5	577	20.3	18.6-22.0	553	12.6	11.5-13.7	235	3.3	2.9-3.8	116	4.4	3.6-5.3	119	2.7	2.2-3.2
1989	1,078	14.9	14.0-15.9	563	19.7	18.0-21.4	515	11.8	10.8-12.8	197	2.8	2.4-3.2	98	3.5	2.8-4.3	99	2.2	1.8-2.7
1990	1,036	14.3	13.5-15.2	524	18.2	16.6-19.8	512	11.6	10.6-12.7	219	3.1	2.7-3.5	115	4.3	3.5-5.1	104	2.4	1.9-2.9
1991	1,070	15.1	14.2-16.0	561	20.4	18.7-22.2	509	11.8	10.8-12.9	229	3.2	2.8-3.7	116	4.4	3.6-5.3	113	2.6	2.1-3.1
1992	1,053	14.8	13.9-15.7	509	18.0	16.4-19.6	544	12.7	11.6-13.8	221	3.1	2.7-3.6	106	4.0	3.2-4.8	115	2.6	2.2-3.2
1993	1,048	14.7	13.8-15.6	553	19.6	18.0-21.4	495	11.4	10.5-12.5	216	3.0	2.6-3.4	116	4.2	3.4-5.0	100	2.2	1.8-2.7
1994	1,123	15.7	14.8-16.7	562	19.8	18.1-21.5	561	13.0	12.0-14.2	207	2.9	2.5-3.4	100	3.7	3.0-4.6	107	2.4	2.0-2.9
1995	1,071	14.9	14.0-15.8	537	18.6	17.0-20.3	534	12.4	11.4-13.5	216	3.0	2.6-3.4	100	3.6	2.9-4.4	116	2.7	2.2-3.2
1996	1,074	15.0	14.1-15.9	549	19.2	17.6-20.9	525	12.3	11.3-13.4	197	2.7	2.4-3.1	91	3.2	2.6-3.9	106	2.4	2.0-3.0
1997	1,083	15.0	14.1-15.9	540	18.5	16.9-20.1	543	12.5	11.4-13.6	229	3.2	2.8-3.6	122	4.4	3.6-5.2	107	2.4	2.0-2.9
1998	1,172	16.0	15.1-17.0	593	20.0	18.4-21.7	579	13.3	12.2-14.4	175	2.4	2.1-2.8	81	2.7	2.2-3.4	94	2.1	1.7-2.6
1999	1,136	15.2	14.4-16.2	576	18.9	17.4-20.5	560	12.7	11.6-13.8	206	2.8	2.4-3.2	103	3.5	2.8-4.2	103	2.3	1.9-2.8
2000	1,144	15.1	14.2-16.0	624	20.1	18.5-21.7	520	11.5	10.6-12.6	191	2.5	2.2-2.9	94	3.1	2.5-3.8	97	2.1	1.7-2.5
2001	1,138	14.8	14.0-15.7	579	18.6	17.1-20.2	559	12.4	11.4-13.5	205	2.7	2.3-3.1	115	3.8	3.1-4.6	90	1.9	1.6-2.4
2002	1,098	14.1	13.3-15.0	594	18.3	16.8-19.8	504	11.0	10.0-12.0	196	2.5	2.2-2.9	98	3.1	2.5-3.8	98	2.1	1.7-2.5
2003	1,152	14.7	13.8-15.6	615	18.9	17.4-20.5	537	11.7	10.7-12.7	203	2.6	2.3-3.0	104	3.3	2.7-4.0	99	2.1	1.7-2.6
2004	1,117	14.1	13.3-15.0	554	17.0	15.6-18.5	563	12.2	11.2-13.3	187	2.4	2.1-2.8	103	3.3	2.7-4.1	84	1.8	1.4-2.2
2005	1,034	13.0	12.2-13.8	563	17.1	15.7-18.6	471	10.1	9.2-11.1	175	2.2	1.9-2.5	91	2.9	2.3-3.6	84	1.7	1.4-2.2
2006	1,135	14.2	13.3-15.0	582	17.5	16.1-19.0	553	11.8	10.8-12.9	178	2.2	1.9-2.6	92	2.9	2.3-3.6	86	1.8	1.4-2.2
2007	1,140	14.1	13.3-14.9	598	17.4	16.0-18.9	542	11.6	10.6-12.6	178	2.2	1.9-2.6	89	2.8	2.2-3.4	89	1.8	1.5-2.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Rectum & rectosigmoid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	1,064	12.9	12.2-13.8	559	15.9	14.5-17.2	505	10.7	9.8-11.7	214	2.6	2.3-3.0	113	3.4	2.8-4.1	101	2.0	1.6-2.4
2009	1,064	12.8	12.0-13.6	586	16.2	14.9-17.6	478	10.2	9.3-11.1	184	2.2	1.9-2.6	84	2.5	2.0-3.1	100	2.0	1.6-2.5
2010	1,029	12.3	11.5-13.0	557	15.8	14.5-17.2	472	9.9	9.0-10.8	183	2.2	1.9-2.5	109	3.2	2.6-3.8	74	1.5	1.2-1.9
2011	1,015	11.9	11.1-12.6	527	14.4	13.2-15.7	488	10.0	9.1-10.9	208	2.4	2.1-2.7	107	3.0	2.4-3.6	101	1.9	1.6-2.4
2012	967	11.0	10.3-11.7	557	14.6	13.4-15.9	410	8.2	7.4-9.1	216	2.5	2.1-2.8	115	3.1	2.6-3.8	101	2.0	1.6-2.4
2013	1,049	11.8	11.0-12.5	571	14.7	13.5-16.0	478	9.4	8.6-10.3	190	2.1	1.8-2.4	111	2.8	2.3-3.4	79	1.5	1.2-1.9
2014	1,032	11.3	10.7-12.1	568	14.1	12.9-15.3	464	9.1	8.3-10.0	211	2.4	2.0-2.7	116	3.1	2.5-3.7	95	1.8	1.4-2.2
2015	1,090	11.8	11.1-12.5	605	14.8	13.6-16.1	485	9.4	8.6-10.3	200	2.2	1.9-2.5	103	2.8	2.2-3.3	97	1.7	1.4-2.1
2016	1,045	11.2	10.5-11.9	581	14.2	13.0-15.4	464	8.9	8.1-9.8	233	2.4	2.1-2.8	127	3.2	2.7-3.8	106	1.9	1.5-2.3
2017	1,085	11.5	10.8-12.2	620	14.7	13.5-15.9	465	9.0	8.2-9.9	232	2.4	2.1-2.8	132	3.3	2.7-3.9	100	1.8	1.5-2.2
2018	1,026	10.8	10.2-11.5	566	13.6	12.5-14.8	460	8.6	7.8-9.5	186	2.0	1.7-2.3	102	2.5	2.0-3.0	84	1.5	1.2-1.9
2019	1,022	10.7	10.0-11.4	584	13.7	12.6-14.8	438	8.4	7.6-9.3	201	2.1	1.8-2.4	121	2.9	2.4-3.5	80	1.4	1.1-1.7
2020	890	9.3	8.7-10.0	521	12.2	11.1-13.3	369	7.1	6.3-7.8	210	2.1	1.8-2.4	119	2.9	2.4-3.4	91	1.5	1.2-1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Liver / intrahepatic bile duct**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	304	4.2	3.7-4.7	178	6.0	5.1-7.0	126	2.9	2.4-3.5	316	4.3	3.9-4.9	181	5.9	5.0-6.9	135	3.2	2.7-3.9
1977	296	4.0	3.6-4.5	192	6.3	5.4-7.3	104	2.4	2.0-3.0	352	4.8	4.3-5.3	204	6.7	5.8-7.7	148	3.5	2.9-4.1
1978	300	4.2	3.7-4.7	181	6.1	5.2-7.1	119	2.8	2.3-3.3	327	4.6	4.1-5.1	191	6.5	5.6-7.5	136	3.2	2.7-3.8
1979	272	3.8	3.4-4.3	166	5.7	4.9-6.7	106	2.5	2.1-3.1	348	4.9	4.4-5.5	188	6.6	5.7-7.7	160	3.8	3.2-4.5
1980	292	4.1	3.6-4.6	174	6.0	5.1-6.9	118	2.9	2.4-3.4	303	4.2	3.7-4.7	167	5.8	4.9-6.7	136	3.2	2.7-3.8
1981	308	4.3	3.8-4.8	187	6.4	5.5-7.5	121	2.8	2.3-3.3	350	4.9	4.4-5.5	202	7.2	6.2-8.3	148	3.4	2.9-4.1
1982	274	3.8	3.3-4.3	194	6.7	5.8-7.8	80	1.8	1.4-2.3	337	4.6	4.1-5.2	204	7.0	6.0-8.1	133	3.1	2.6-3.6
1983	304	4.3	3.8-4.8	188	6.5	5.6-7.6	116	2.7	2.2-3.3	379	5.3	4.8-5.9	216	7.4	6.4-8.4	163	3.8	3.2-4.5
1984	325	4.5	4.0-5.0	208	7.1	6.1-8.2	117	2.7	2.2-3.3	374	5.1	4.6-5.7	212	7.4	6.4-8.5	162	3.7	3.1-4.3
1985	330	4.6	4.1-5.1	208	7.2	6.2-8.3	122	2.8	2.3-3.4	397	5.6	5.0-6.1	235	8.4	7.3-9.6	162	3.7	3.1-4.3
1986	354	4.9	4.4-5.5	233	8.1	7.1-9.3	121	2.8	2.4-3.4	400	5.6	5.1-6.2	230	8.1	7.1-9.2	170	3.9	3.4-4.6
1987	320	4.4	3.9-4.9	198	6.8	5.9-7.9	122	2.7	2.3-3.3	400	5.6	5.0-6.1	226	8.0	7.0-9.2	174	4.0	3.4-4.6
1988	348	4.9	4.4-5.5	220	7.6	6.6-8.7	128	3.0	2.5-3.6	400	5.6	5.1-6.2	233	8.1	7.1-9.3	167	3.8	3.3-4.5
1989	316	4.4	3.9-4.9	198	6.8	5.9-7.9	118	2.7	2.2-3.3	362	5.1	4.6-5.6	215	7.5	6.5-8.5	147	3.4	2.9-4.0
1990	398	5.6	5.0-6.2	250	8.7	7.6-9.8	148	3.4	2.9-4.0	423	5.9	5.3-6.5	243	8.7	7.6-9.9	180	4.1	3.5-4.8
1991	441	6.2	5.6-6.8	295	10.2	9.0-11.4	146	3.4	2.8-4.0	458	6.4	5.8-7.0	276	9.7	8.6-10.9	182	4.1	3.5-4.8
1992	450	6.4	5.8-7.0	283	9.7	8.6-11.0	167	3.9	3.4-4.6	477	6.7	6.1-7.4	289	10.2	9.0-11.4	188	4.4	3.8-5.1
1993	417	5.8	5.3-6.4	268	9.1	8.1-10.3	149	3.5	2.9-4.1	443	6.2	5.6-6.8	255	8.8	7.7-9.9	188	4.3	3.7-5.0
1994	474	6.7	6.1-7.3	302	10.2	9.0-11.4	172	4.0	3.4-4.7	388	5.5	4.9-6.1	253	8.8	7.8-10.0	135	3.1	2.6-3.7
1995	489	6.8	6.3-7.5	316	10.7	9.5-11.9	173	4.0	3.5-4.7	421	5.9	5.4-6.5	251	8.6	7.6-9.8	170	3.9	3.3-4.5
1996	549	7.6	7.0-8.3	389	13.0	11.7-14.4	160	3.7	3.2-4.3	432	6.0	5.4-6.6	275	9.4	8.3-10.6	157	3.6	3.0-4.2
1997	565	7.8	7.2-8.5	368	12.2	10.9-13.5	197	4.5	3.9-5.2	414	5.7	5.2-6.3	259	8.8	7.7-9.9	155	3.5	3.0-4.1
1998	645	8.8	8.1-9.5	432	14.0	12.7-15.4	213	4.9	4.3-5.6	464	6.3	5.8-6.9	304	10.0	8.9-11.2	160	3.6	3.1-4.2
1999	644	8.6	8.0-9.3	419	13.2	11.9-14.5	225	5.1	4.4-5.8	432	5.8	5.3-6.4	272	8.9	7.8-10.0	160	3.6	3.0-4.2
2000	662	8.7	8.1-9.4	459	14.2	12.9-15.6	203	4.5	3.9-5.2	484	6.4	5.8-7.0	314	10.0	8.9-11.1	170	3.8	3.2-4.4
2001	660	8.6	7.9-9.2	471	14.3	13.0-15.6	189	4.2	3.6-4.8	468	6.1	5.6-6.7	303	9.4	8.3-10.5	165	3.6	3.0-4.2
2002	658	8.4	7.8-9.1	467	13.9	12.7-15.2	191	4.2	3.6-4.9	481	6.2	5.6-6.8	316	9.7	8.6-10.8	165	3.5	3.0-4.1
2003	738	9.4	8.7-10.1	508	15.0	13.7-16.4	230	5.1	4.4-5.8	499	6.4	5.8-6.9	319	9.8	8.8-11.0	180	3.8	3.3-4.4
2004	791	9.9	9.2-10.6	565	16.5	15.1-17.9	226	4.9	4.3-5.6	542	6.9	6.3-7.5	344	10.2	9.1-11.3	198	4.3	3.7-4.9
2005	830	10.3	9.6-11.0	620	17.7	16.4-19.2	210	4.5	3.9-5.2	547	6.9	6.3-7.5	365	10.8	9.7-11.9	182	3.8	3.3-4.4
2006	848	10.5	9.8-11.2	600	17.0	15.6-18.4	248	5.4	4.7-6.1	567	7.1	6.5-7.7	389	11.5	10.4-12.7	178	3.8	3.3-4.4
2007	911	11.1	10.3-11.8	662	18.4	17.0-19.9	249	5.3	4.7-6.0	596	7.3	6.8-8.0	405	11.7	10.6-12.9	191	4.1	3.5-4.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Liver / intrahepatic bile duct**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	908	10.9	10.2-11.6	671	18.2	16.9-19.7	237	5.1	4.4-5.8	530	6.4	5.8-6.9	373	10.5	9.4-11.6	157	3.2	2.7-3.8
2009	949	11.2	10.5-11.9	676	18.2	16.8-19.7	273	5.7	5.0-6.4	599	7.2	6.7-7.9	408	11.5	10.4-12.7	191	4.0	3.5-4.7
2010	994	11.5	10.8-12.3	708	18.7	17.3-20.2	286	5.9	5.2-6.7	618	7.2	6.7-7.8	421	11.6	10.5-12.8	197	3.9	3.4-4.5
2011	998	11.3	10.6-12.0	696	17.9	16.6-19.3	302	6.1	5.4-6.8	588	6.8	6.2-7.4	379	10.1	9.1-11.2	209	4.2	3.6-4.8
2012	1,093	12.1	11.4-12.8	772	19.4	18.1-20.9	321	6.4	5.7-7.1	655	7.4	6.8-8.0	436	11.5	10.4-12.6	219	4.3	3.7-4.9
2013	1,062	11.5	10.8-12.2	744	18.2	16.9-19.6	318	6.1	5.4-6.8	655	7.2	6.6-7.8	438	11.2	10.1-12.3	217	4.1	3.6-4.7
2014	1,071	11.3	10.6-12.0	736	17.4	16.1-18.7	335	6.3	5.6-7.0	669	7.2	6.6-7.7	442	10.8	9.8-11.9	227	4.3	3.7-4.9
2015	1,118	11.7	11.0-12.4	778	18.5	17.2-19.9	340	6.3	5.6-7.0	666	7.0	6.4-7.5	441	10.7	9.7-11.8	225	4.1	3.6-4.7
2016	1,126	11.4	10.8-12.1	804	18.5	17.2-19.8	322	5.9	5.2-6.6	693	7.2	6.6-7.7	475	11.3	10.3-12.4	218	3.9	3.4-4.5
2017	1,112	11.3	10.6-12.0	789	18.1	16.9-19.5	323	5.8	5.2-6.5	632	6.5	6.0-7.0	425	10.1	9.1-11.1	207	3.7	3.2-4.3
2018	1,047	10.5	9.8-11.1	728	16.6	15.3-17.8	319	5.7	5.0-6.3	609	6.1	5.6-6.6	400	9.3	8.4-10.2	209	3.6	3.1-4.2
2019	1,018	10.1	9.5-10.7	699	15.7	14.6-17.0	319	5.6	5.0-6.3	640	6.3	5.8-6.8	405	9.4	8.4-10.3	235	4.0	3.5-4.6
2020	858	8.4	7.9-9.0	572	12.9	11.8-14.0	286	4.9	4.3-5.5	567	5.5	5.1-6.0	364	8.3	7.5-9.3	203	3.4	2.9-3.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Pancreas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	991	13.6	12.7-14.5	488	16.5	15.0-18.2	503	11.6	10.6-12.7	913	12.6	11.8-13.5	450	15.4	13.9-16.9	463	10.8	9.8-11.8
1977	974	13.5	12.7-14.4	470	16.4	14.9-18.0	504	11.7	10.7-12.8	905	12.6	11.7-13.4	436	15.2	13.7-16.7	469	10.8	9.9-11.9
1978	947	12.9	12.0-13.7	460	15.4	13.9-16.9	487	11.1	10.1-12.2	862	11.7	10.9-12.5	415	14.0	12.6-15.5	447	10.2	9.2-11.2
1979	924	12.8	11.9-13.6	444	15.1	13.6-16.6	480	11.1	10.1-12.1	877	12.2	11.4-13.0	414	14.3	12.9-15.8	463	10.7	9.8-11.8
1980	971	13.4	12.6-14.3	485	17.1	15.5-18.7	486	11.0	10.0-12.1	869	12.0	11.2-12.8	443	15.4	14.0-17.0	426	9.7	8.8-10.7
1981	938	13.0	12.2-13.9	479	16.8	15.3-18.4	459	10.5	9.6-11.6	859	11.9	11.1-12.8	419	14.7	13.3-16.3	440	9.9	9.0-10.9
1982	975	13.6	12.8-14.5	428	15.7	14.2-17.3	547	12.4	11.3-13.5	838	11.7	10.9-12.5	380	13.8	12.4-15.4	458	10.3	9.4-11.3
1983	946	13.0	12.1-13.8	446	15.3	13.9-16.9	500	11.3	10.3-12.3	847	11.6	10.8-12.4	419	14.4	13.0-15.9	428	9.7	8.8-10.7
1984	976	13.4	12.6-14.3	440	15.6	14.1-17.2	536	12.0	11.0-13.1	856	11.8	11.0-12.7	397	14.1	12.7-15.5	459	10.3	9.4-11.3
1985	903	12.4	11.6-13.3	422	14.7	13.3-16.2	481	10.8	9.8-11.8	768	10.6	9.8-11.4	372	13.2	11.8-14.6	396	8.8	8.0-9.8
1986	942	13.1	12.3-14.0	417	14.9	13.5-16.5	525	11.8	10.8-12.9	801	11.0	10.2-11.8	349	12.3	11.0-13.7	452	10.0	9.1-11.0
1987	925	12.8	12.0-13.6	419	14.8	13.3-16.3	506	11.3	10.3-12.3	783	10.8	10.0-11.6	365	12.8	11.5-14.2	418	9.4	8.5-10.3
1988	917	12.7	11.9-13.6	396	14.2	12.8-15.7	521	11.6	10.6-12.6	745	10.3	9.6-11.1	351	12.6	11.3-14.0	394	8.7	7.9-9.6
1989	893	12.5	11.7-13.4	397	14.3	12.9-15.8	496	11.3	10.3-12.3	814	11.4	10.6-12.2	362	13.1	11.8-14.5	452	10.2	9.3-11.2
1990	892	12.5	11.7-13.4	395	14.2	12.8-15.7	497	11.3	10.4-12.4	776	10.9	10.1-11.7	346	12.7	11.4-14.2	430	9.7	8.8-10.7
1991	887	12.4	11.6-13.3	397	14.3	12.9-15.8	490	11.1	10.1-12.1	766	10.8	10.0-11.6	356	12.8	11.4-14.2	410	9.3	8.5-10.3
1992	919	12.9	12.1-13.8	414	15.1	13.7-16.7	505	11.5	10.5-12.5	742	10.4	9.7-11.2	319	11.7	10.4-13.1	423	9.5	8.6-10.5
1993	913	12.8	12.0-13.7	396	14.1	12.7-15.6	517	11.8	10.8-12.8	799	11.2	10.4-12.0	369	13.5	12.1-14.9	430	9.7	8.8-10.7
1994	960	13.5	12.7-14.4	413	14.8	13.4-16.3	547	12.5	11.4-13.6	825	11.6	10.8-12.4	334	12.1	10.8-13.5	491	11.2	10.2-12.2
1995	854	12.0	11.2-12.8	365	13.2	11.9-14.7	489	11.1	10.2-12.2	748	10.5	9.7-11.3	332	12.0	10.7-13.4	416	9.4	8.5-10.3
1996	917	12.8	12.0-13.6	399	14.2	12.8-15.7	518	11.9	10.9-12.9	823	11.5	10.7-12.3	351	12.7	11.4-14.1	472	10.8	9.8-11.8
1997	909	12.6	11.8-13.5	411	14.3	12.9-15.8	498	11.3	10.3-12.4	729	10.2	9.4-10.9	320	11.4	10.2-12.8	409	9.2	8.3-10.2
1998	926	12.7	11.9-13.6	427	14.9	13.5-16.4	499	11.3	10.3-12.3	748	10.3	9.6-11.0	335	11.9	10.6-13.2	413	9.3	8.4-10.2
1999	964	13.0	12.2-13.9	446	15.2	13.8-16.7	518	11.5	10.6-12.6	807	10.9	10.2-11.7	383	13.1	11.8-14.4	424	9.3	8.5-10.3
2000	1,023	13.6	12.8-14.4	438	14.7	13.3-16.1	585	12.8	11.8-13.9	765	10.2	9.5-10.9	328	11.1	9.9-12.3	437	9.4	8.6-10.4
2001	950	12.5	11.7-13.3	401	13.2	11.9-14.6	549	11.9	10.9-12.9	830	10.9	10.2-11.6	353	12.0	10.7-13.3	477	10.2	9.3-11.2
2002	927	12.0	11.2-12.8	421	13.4	12.1-14.8	506	10.7	9.8-11.7	771	10.0	9.3-10.7	333	10.8	9.7-12.0	438	9.3	8.4-10.2
2003	996	12.8	12.0-13.6	463	15.0	13.7-16.5	533	11.2	10.3-12.2	776	10.0	9.3-10.7	367	11.8	10.6-13.1	409	8.5	7.7-9.4
2004	995	12.7	11.9-13.5	438	13.8	12.5-15.2	557	11.8	10.8-12.8	829	10.6	9.8-11.3	379	12.1	10.9-13.4	450	9.4	8.5-10.3
2005	949	12.0	11.2-12.7	439	13.7	12.4-15.0	510	10.8	9.9-11.8	793	10.0	9.3-10.7	376	11.8	10.6-13.1	417	8.8	7.9-9.7
2006	1,052	13.2	12.4-14.1	476	14.7	13.3-16.1	576	12.1	11.1-13.2	804	10.2	9.5-10.9	361	11.4	10.3-12.7	443	9.3	8.4-10.2
2007	1,071	13.3	12.5-14.1	514	15.7	14.3-17.1	557	11.6	10.6-12.6	852	10.6	9.9-11.3	394	12.3	11.1-13.6	458	9.5	8.6-10.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Pancreas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	1,099	13.6	12.8-14.4	504	15.2	13.9-16.6	595	12.3	11.3-13.3	886	11.0	10.2-11.7	396	12.1	10.9-13.4	490	10.0	9.1-10.9
2009	1,125	13.6	12.8-14.5	545	16.3	14.9-17.8	580	11.8	10.9-12.8	893	10.9	10.2-11.6	447	13.5	12.2-14.8	446	9.0	8.2-9.9
2010	1,156	13.9	13.1-14.7	565	16.3	15.0-17.7	591	12.0	11.1-13.1	877	10.5	9.8-11.3	417	12.2	11.0-13.5	460	9.2	8.3-10.1
2011	1,176	13.8	13.0-14.6	551	15.5	14.2-16.9	625	12.5	11.5-13.5	943	11.1	10.4-11.9	453	13.1	11.9-14.4	490	9.7	8.8-10.6
2012	1,196	13.7	13.0-14.6	556	15.4	14.1-16.7	640	12.7	11.7-13.7	948	10.9	10.2-11.6	432	12.0	10.9-13.2	516	10.2	9.3-11.1
2013	1,231	13.7	12.9-14.4	591	15.6	14.3-16.9	640	12.2	11.2-13.2	896	10.1	9.4-10.7	434	11.9	10.8-13.1	462	8.7	8.0-9.6
2014	1,198	13.2	12.4-14.0	582	15.2	13.9-16.5	616	11.6	10.7-12.6	952	10.4	9.7-11.1	454	12.0	10.9-13.2	498	9.2	8.4-10.1
2015	1,261	13.5	12.8-14.3	618	15.6	14.4-16.9	643	11.9	11.0-12.8	926	9.9	9.3-10.6	439	11.3	10.2-12.4	487	8.9	8.1-9.7
2016	1,250	13.1	12.4-13.9	602	14.9	13.7-16.2	648	11.7	10.8-12.7	980	10.3	9.6-11.0	492	12.3	11.2-13.4	488	8.7	8.0-9.6
2017	1,363	14.2	13.4-14.9	663	16.3	15.1-17.6	700	12.6	11.6-13.6	1,004	10.4	9.7-11.0	474	11.9	10.8-13.0	530	9.3	8.5-10.1
2018	1,345	13.7	12.9-14.4	626	14.9	13.7-16.2	719	12.6	11.7-13.6	948	9.6	9.0-10.2	441	10.7	9.7-11.8	507	8.7	7.9-9.5
2019	1,313	13.2	12.5-13.9	628	14.7	13.5-15.9	685	11.9	11.0-12.9	984	9.8	9.2-10.4	460	10.7	9.8-11.8	524	8.9	8.2-9.8
2020	1,289	12.7	12.0-13.4	627	14.3	13.1-15.4	662	11.5	10.6-12.4	954	9.3	8.8-10.0	472	11.0	10.0-12.0	482	8.1	7.4-8.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York City

Larynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	437	5.7	5.1-6.2	356	11.0	9.8-12.3	81	1.9	1.5-2.3	194	2.6	2.2-3.0	154	5.0	4.2-5.9	40	0.9	0.7-1.3
1977	432	5.8	5.2-6.3	363	11.5	10.3-12.8	69	1.7	1.3-2.1	186	2.5	2.1-2.9	158	5.1	4.3-6.0	28	0.7	0.4-1.0
1978	419	5.7	5.2-6.3	336	11.2	10.0-12.5	83	1.9	1.5-2.4	191	2.6	2.2-3.0	157	5.1	4.3-6.0	34	0.8	0.5-1.1
1979	468	6.3	5.7-6.9	380	12.1	10.9-13.4	88	2.1	1.7-2.7	189	2.5	2.2-2.9	153	5.0	4.3-5.9	36	0.8	0.6-1.1
1980	470	6.5	5.9-7.1	379	12.5	11.2-13.9	91	2.2	1.8-2.8	176	2.4	2.1-2.8	140	4.7	4.0-5.6	36	0.9	0.6-1.3
1981	436	6.1	5.5-6.7	348	11.6	10.4-12.9	88	2.1	1.7-2.7	172	2.3	2.0-2.7	130	4.3	3.6-5.1	42	1.0	0.7-1.4
1982	423	5.8	5.2-6.4	341	11.3	10.1-12.7	82	1.9	1.5-2.4	163	2.2	1.8-2.5	131	4.3	3.6-5.1	32	0.8	0.5-1.1
1983	436	6.0	5.4-6.6	341	11.0	9.9-12.3	95	2.3	1.9-2.9	157	2.2	1.8-2.6	123	4.2	3.4-5.0	34	0.8	0.5-1.1
1984	478	6.6	6.0-7.2	369	12.2	11.0-13.5	109	2.7	2.2-3.2	173	2.4	2.0-2.8	138	4.5	3.8-5.4	35	0.8	0.6-1.2
1985	497	6.9	6.3-7.5	389	12.8	11.6-14.2	108	2.6	2.1-3.2	174	2.3	2.0-2.7	140	4.7	3.9-5.5	34	0.8	0.5-1.1
1986	455	6.3	5.7-6.9	352	11.8	10.5-13.1	103	2.6	2.1-3.1	155	2.1	1.8-2.5	126	4.3	3.6-5.1	29	0.7	0.5-1.0
1987	422	6.0	5.4-6.6	333	11.2	10.0-12.5	89	2.3	1.8-2.8	171	2.4	2.0-2.8	131	4.4	3.7-5.3	40	0.9	0.7-1.3
1988	477	6.6	6.0-7.3	364	11.9	10.7-13.2	113	2.8	2.3-3.4	176	2.5	2.1-2.9	142	4.9	4.1-5.8	34	0.8	0.6-1.2
1989	495	7.0	6.4-7.7	337	11.3	10.1-12.5	158	4.0	3.4-4.7	164	2.3	2.0-2.7	119	4.1	3.4-4.9	45	1.1	0.8-1.5
1990	476	6.7	6.1-7.4	378	12.9	11.6-14.3	98	2.5	2.0-3.0	141	2.0	1.7-2.3	119	4.1	3.4-4.9	22	0.5	0.3-0.8
1991	428	6.1	5.5-6.7	334	11.4	10.2-12.7	94	2.4	2.0-3.0	157	2.2	1.9-2.6	118	4.1	3.4-4.9	39	1.0	0.7-1.3
1992	468	6.6	6.0-7.3	360	12.1	10.9-13.4	108	2.7	2.2-3.2	178	2.5	2.2-2.9	135	4.8	4.0-5.7	43	1.0	0.8-1.4
1993	404	5.7	5.1-6.3	313	10.6	9.4-11.8	91	2.3	1.8-2.8	180	2.5	2.1-2.9	139	4.8	4.0-5.6	41	1.0	0.7-1.3
1994	390	5.5	5.0-6.1	303	10.4	9.2-11.7	87	2.1	1.7-2.6	161	2.3	1.9-2.6	123	4.2	3.5-5.0	38	0.9	0.6-1.2
1995	348	4.9	4.4-5.4	266	8.8	7.8-9.9	82	2.0	1.6-2.5	166	2.3	2.0-2.7	127	4.3	3.6-5.1	39	0.9	0.7-1.3
1996	412	5.8	5.2-6.4	329	11.0	9.8-12.2	83	2.0	1.6-2.5	166	2.3	2.0-2.7	141	4.8	4.1-5.7	25	0.6	0.4-0.9
1997	363	5.0	4.5-5.6	293	9.7	8.6-10.9	70	1.7	1.3-2.1	136	1.9	1.6-2.2	107	3.6	3.0-4.4	29	0.7	0.5-1.0
1998	362	4.9	4.4-5.5	270	8.8	7.8-9.9	92	2.2	1.8-2.7	140	1.9	1.6-2.3	113	3.7	3.1-4.5	27	0.6	0.4-0.9
1999	326	4.4	3.9-4.9	260	8.3	7.3-9.4	66	1.6	1.2-2.0	127	1.7	1.4-2.0	96	3.1	2.5-3.8	31	0.7	0.5-1.0
2000	344	4.5	4.1-5.0	260	8.2	7.2-9.3	84	1.9	1.5-2.4	147	1.9	1.6-2.3	118	3.8	3.2-4.6	29	0.7	0.4-0.9
2001	325	4.2	3.8-4.7	244	7.6	6.7-8.7	81	1.8	1.4-2.3	100	1.3	1.1-1.6	79	2.5	2.0-3.1	21	0.4	0.3-0.7
2002	317	4.1	3.6-4.6	254	7.8	6.9-8.8	63	1.4	1.1-1.8	125	1.6	1.3-1.9	97	3.0	2.5-3.7	28	0.6	0.4-0.9
2003	349	4.4	4.0-4.9	268	8.2	7.2-9.2	81	1.8	1.4-2.2	120	1.5	1.3-1.8	96	3.0	2.4-3.7	24	0.5	0.3-0.8
2004	338	4.2	3.8-4.7	281	8.2	7.3-9.3	57	1.3	1.0-1.6	123	1.5	1.3-1.8	108	3.2	2.6-3.9	15	0.3	0.2-0.5
2005	317	4.0	3.5-4.4	245	7.2	6.3-8.2	72	1.6	1.2-2.0	106	1.3	1.1-1.6	86	2.6	2.1-3.2	20	0.4	0.3-0.7
2006	279	3.4	3.0-3.9	223	6.5	5.7-7.4	56	1.2	0.9-1.6	111	1.4	1.2-1.7	93	2.9	2.3-3.6	18	0.4	0.2-0.6
2007	310	3.8	3.4-4.3	242	6.9	6.1-7.9	68	1.4	1.1-1.8	111	1.4	1.1-1.6	88	2.6	2.1-3.2	23	0.5	0.3-0.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York City

Larynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	354	4.3	3.9-4.8	262	7.6	6.7-8.6	92	2.0	1.6-2.4	104	1.3	1.1-1.6	81	2.4	1.9-3.0	23	0.5	0.3-0.7
2009	308	3.6	3.2-4.1	238	6.6	5.8-7.5	70	1.5	1.2-1.9	114	1.4	1.1-1.6	87	2.5	2.0-3.1	27	0.5	0.4-0.8
2010	299	3.5	3.1-3.9	241	6.7	5.8-7.6	58	1.2	0.9-1.6	98	1.2	0.9-1.4	77	2.2	1.8-2.8	21	0.4	0.3-0.6
2011	267	3.0	2.6-3.4	206	5.4	4.7-6.2	61	1.2	0.9-1.5	104	1.2	1.0-1.5	79	2.3	1.8-2.8	25	0.5	0.3-0.7
2012	291	3.2	2.9-3.6	237	6.1	5.4-7.0	54	1.1	0.8-1.4	96	1.1	0.9-1.3	76	2.1	1.6-2.6	20	0.4	0.2-0.6
2013	289	3.2	2.8-3.6	242	6.3	5.5-7.2	47	0.9	0.7-1.2	100	1.1	0.9-1.4	79	2.2	1.7-2.7	21	0.4	0.2-0.6
2014	289	3.2	2.8-3.6	224	5.6	4.9-6.4	65	1.3	1.0-1.6	93	1.0	0.8-1.2	77	2.0	1.5-2.5	16	0.3	0.2-0.5
2015	295	3.1	2.7-3.5	236	5.8	5.0-6.6	59	1.1	0.8-1.4	81	0.9	0.7-1.1	60	1.5	1.2-2.0	21	0.4	0.3-0.6
2016	252	2.6	2.3-3.0	205	5.0	4.3-5.8	47	0.9	0.6-1.2	96	1.0	0.8-1.2	81	2.0	1.6-2.5	15	0.3	0.1-0.4
2017	277	2.8	2.5-3.2	213	5.0	4.3-5.7	64	1.2	0.9-1.6	71	0.7	0.5-0.9	56	1.3	1.0-1.8	15	0.2	0.1-0.4
2018	237	2.4	2.1-2.7	187	4.2	3.7-4.9	50	0.9	0.7-1.2	73	0.7	0.6-0.9	57	1.4	1.0-1.8	16	0.3	0.2-0.5
2019	258	2.6	2.3-3.0	200	4.7	4.1-5.4	58	1.1	0.8-1.4	74	0.7	0.6-0.9	63	1.5	1.1-1.9	11	0.2	0.1-0.3
2020	193	1.9	1.7-2.2	157	3.6	3.0-4.2	36	0.6	0.4-0.9	49	0.5	0.3-0.6	37	0.9	0.6-1.2	12	0.2	0.1-0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Lung and bronchus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	3,928	51.5	49.9-53.2	2,777	88.6	85.1-92.1	1,151	26.3	24.8-27.9	3,188	42.0	40.5-43.5	2,302	74.4	71.3-77.7	886	20.2	18.8-21.6
1977	3,904	51.6	50.0-53.3	2,706	88.4	84.9-91.9	1,198	27.2	25.6-28.8	3,240	42.9	41.4-44.4	2,342	76.4	73.2-79.7	898	20.5	19.2-22.0
1978	3,968	53.3	51.6-55.0	2,809	92.5	89.0-96.1	1,159	27.1	25.5-28.7	3,236	43.6	42.1-45.2	2,303	76.6	73.4-79.9	933	21.7	20.3-23.2
1979	4,087	55.1	53.4-56.9	2,784	92.9	89.4-96.6	1,303	30.1	28.5-31.9	3,255	44.1	42.6-45.7	2,289	77.2	73.9-80.5	966	22.4	20.9-23.9
1980	4,047	55.0	53.3-56.8	2,744	92.3	88.7-95.9	1,303	30.4	28.7-32.1	3,376	45.8	44.3-47.4	2,305	78.1	74.8-81.5	1,071	24.9	23.4-26.4
1981	3,962	53.8	52.1-55.5	2,602	88.5	85.0-92.0	1,360	31.0	29.4-32.8	3,250	44.3	42.8-45.9	2,215	75.4	72.2-78.8	1,035	23.9	22.4-25.4
1982	4,168	56.9	55.2-58.7	2,678	91.5	88.0-95.2	1,490	34.4	32.6-36.2	3,291	44.8	43.3-46.4	2,187	74.7	71.5-78.0	1,104	25.6	24.1-27.2
1983	4,171	56.6	54.9-58.4	2,695	91.7	88.1-95.3	1,476	33.8	32.1-35.6	3,348	45.7	44.1-47.3	2,118	73.4	70.2-76.7	1,230	27.9	26.4-29.6
1984	4,237	57.8	56.1-59.6	2,760	93.8	90.3-97.5	1,477	34.4	32.7-36.3	3,287	44.9	43.4-46.5	2,192	75.9	72.7-79.3	1,095	25.0	23.5-26.6
1985	4,230	57.4	55.6-59.2	2,709	91.9	88.4-95.5	1,521	35.0	33.2-36.8	3,375	45.8	44.3-47.4	2,145	73.7	70.5-76.9	1,230	28.1	26.6-29.8
1986	4,202	57.3	55.5-59.1	2,591	88.2	84.8-91.8	1,611	37.1	35.3-39.0	3,324	45.3	43.7-46.9	2,052	70.6	67.5-73.8	1,272	29.2	27.6-30.9
1987	4,336	58.8	57.1-60.6	2,702	91.8	88.3-95.5	1,634	37.3	35.5-39.2	3,442	46.9	45.3-48.5	2,169	74.9	71.7-78.2	1,273	29.1	27.5-30.8
1988	4,173	57.3	55.5-59.0	2,516	87.2	83.8-90.8	1,657	38.2	36.3-40.1	3,401	46.6	45.0-48.2	2,155	75.1	71.9-78.4	1,246	28.6	27.0-30.3
1989	4,257	59.0	57.2-60.8	2,549	88.1	84.7-91.7	1,708	40.1	38.2-42.1	3,353	46.4	44.8-48.0	2,077	73.3	70.2-76.6	1,276	29.4	27.8-31.1
1990	4,292	59.7	57.9-61.6	2,541	88.8	85.3-92.4	1,751	41.0	39.1-43.0	3,442	47.9	46.3-49.6	2,093	73.8	70.7-77.1	1,349	31.4	29.7-33.1
1991	4,336	61.0	59.2-62.9	2,507	88.0	84.5-91.5	1,829	43.4	41.4-45.4	3,451	48.4	46.8-50.1	2,042	72.7	69.5-76.0	1,409	32.9	31.2-34.7
1992	4,484	62.8	61.0-64.7	2,649	92.7	89.1-96.4	1,835	43.5	41.5-45.6	3,378	47.4	45.8-49.0	1,998	70.9	67.8-74.2	1,380	32.4	30.7-34.1
1993	4,330	60.7	58.9-62.6	2,487	87.3	83.8-90.8	1,843	43.7	41.7-45.8	3,454	48.4	46.8-50.1	2,020	72.0	68.9-75.3	1,434	33.7	32.0-35.5
1994	4,353	61.0	59.2-62.8	2,438	84.7	81.3-88.2	1,915	45.4	43.4-47.5	3,361	47.1	45.5-48.7	2,020	71.4	68.3-74.6	1,341	31.5	29.8-33.2
1995	4,184	58.3	56.5-60.1	2,376	82.2	78.8-85.6	1,808	42.3	40.4-44.3	3,311	46.2	44.6-47.8	1,890	66.2	63.2-69.3	1,421	33.0	31.3-34.8
1996	4,285	59.7	57.9-61.5	2,388	82.8	79.5-86.3	1,897	44.7	42.7-46.7	3,265	45.5	44.0-47.1	1,883	65.6	62.6-68.7	1,382	32.1	30.4-33.8
1997	4,278	59.3	57.6-61.1	2,358	81.1	77.8-84.4	1,920	44.9	43.0-47.0	3,097	43.0	41.5-44.6	1,798	62.8	59.9-65.8	1,299	30.1	28.5-31.8
1998	4,274	58.6	56.9-60.4	2,332	78.2	75.0-81.5	1,942	45.0	43.0-47.1	3,149	43.2	41.7-44.8	1,738	59.4	56.6-62.2	1,411	32.5	30.8-34.3
1999	4,058	54.6	52.9-56.3	2,169	72.0	69.0-75.2	1,889	42.9	41.0-44.9	3,102	41.8	40.4-43.3	1,707	57.5	54.8-60.3	1,395	31.6	30.0-33.3
2000	4,251	56.3	54.6-58.0	2,285	74.8	71.8-78.0	1,966	44.1	42.2-46.1	3,037	40.3	38.9-41.8	1,693	56.2	53.6-59.0	1,344	29.7	28.1-31.3
2001	4,184	54.8	53.2-56.5	2,237	72.4	69.4-75.5	1,947	43.0	41.1-45.0	3,049	40.0	38.6-41.4	1,675	55.0	52.4-57.8	1,374	30.1	28.5-31.7
2002	4,222	54.6	52.9-56.3	2,252	71.5	68.5-74.5	1,970	43.2	41.3-45.2	3,026	39.2	37.8-40.6	1,655	53.2	50.7-55.9	1,371	29.7	28.1-31.3
2003	4,095	52.7	51.1-54.3	2,199	69.6	66.7-72.6	1,896	41.3	39.4-43.2	3,021	38.9	37.5-40.3	1,637	52.4	49.8-55.0	1,384	29.9	28.3-31.5
2004	4,156	53.0	51.4-54.7	2,244	70.1	67.2-73.1	1,912	41.4	39.6-43.3	2,877	36.7	35.4-38.1	1,612	51.2	48.7-53.8	1,265	27.1	25.6-28.6
2005	4,194	53.1	51.5-54.8	2,262	70.2	67.3-73.2	1,932	41.8	40.0-43.7	2,852	36.3	35.0-37.7	1,579	49.7	47.3-52.3	1,273	27.5	26.0-29.1
2006	4,125	52.3	50.7-53.9	2,167	67.2	64.3-70.1	1,958	42.2	40.4-44.2	2,765	35.2	33.9-36.5	1,511	47.5	45.1-49.9	1,254	26.8	25.3-28.3
2007	4,339	54.5	52.9-56.2	2,269	69.4	66.5-72.4	2,070	44.6	42.6-46.5	2,850	35.8	34.5-37.2	1,532	47.9	45.5-50.4	1,318	27.9	26.4-29.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Lung and bronchus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	4,253	52.8	51.2-54.4	2,179	65.8	63.0-68.7	2,074	44.2	42.3-46.1	2,798	34.8	33.5-36.1	1,520	46.6	44.2-49.0	1,278	26.8	25.3-28.3
2009	4,315	52.9	51.3-54.5	2,232	66.6	63.8-69.5	2,083	43.6	41.8-45.6	2,681	32.9	31.6-34.2	1,423	43.1	40.9-45.5	1,258	26.0	24.5-27.5
2010	4,190	50.6	49.0-52.1	2,179	63.4	60.7-66.1	2,011	41.7	39.9-43.6	2,805	33.8	32.5-35.1	1,478	43.9	41.6-46.2	1,327	27.1	25.7-28.7
2011	4,370	51.6	50.0-53.2	2,233	63.7	61.0-66.5	2,137	43.3	41.5-45.2	2,722	32.0	30.8-33.3	1,444	41.7	39.6-44.0	1,278	25.2	23.9-26.7
2012	4,301	49.7	48.2-51.2	2,136	59.5	57.0-62.1	2,165	43.4	41.5-45.3	2,762	32.0	30.8-33.2	1,518	42.8	40.6-45.0	1,244	24.7	23.3-26.1
2013	4,308	48.7	47.3-50.2	2,173	59.0	56.4-61.5	2,135	41.8	40.1-43.7	2,792	31.5	30.4-32.7	1,492	41.3	39.2-43.5	1,300	25.0	23.7-26.4
2014	4,326	47.4	46.0-48.9	2,190	57.2	54.8-59.7	2,136	40.7	39.0-42.5	2,534	27.9	26.8-29.0	1,359	36.4	34.4-38.4	1,175	22.0	20.7-23.3
2015	4,320	46.5	45.1-47.9	2,247	57.5	55.1-59.9	2,073	38.8	37.2-40.6	2,529	27.2	26.1-28.3	1,339	34.9	33.0-36.8	1,190	21.9	20.6-23.2
2016	4,374	46.2	44.8-47.6	2,162	54.1	51.8-56.5	2,212	40.7	39.0-42.5	2,375	25.0	24.0-26.1	1,270	32.5	30.7-34.4	1,105	19.9	18.7-21.1
2017	4,578	47.6	46.2-49.0	2,326	57.3	55.0-59.7	2,252	40.7	39.0-42.5	2,323	24.0	23.0-25.0	1,226	30.5	28.8-32.3	1,097	19.4	18.2-20.6
2018	4,454	45.3	44.0-46.7	2,188	52.3	50.0-54.5	2,266	40.5	38.8-42.2	2,234	22.6	21.6-23.5	1,181	28.7	27.0-30.4	1,053	18.2	17.1-19.4
2019	4,492	44.8	43.5-46.1	2,181	50.9	48.8-53.1	2,311	40.5	38.8-42.2	2,127	21.1	20.2-22.0	1,110	26.4	24.9-28.0	1,017	17.3	16.2-18.4
2020	3,653	36.1	34.9-37.3	1,861	43.5	41.5-45.6	1,792	30.7	29.3-32.2	1,930	19.0	18.2-19.9	1,033	24.4	22.9-25.9	897	15.2	14.2-16.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Melanoma of the skin**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	395	5.4	4.8-6.0	200	6.6	5.6-7.6	195	4.6	4.0-5.3	152	2.1	1.8-2.5	85	2.8	2.2-3.5	67	1.7	1.3-2.1
1977	397	5.4	4.8-5.9	175	5.6	4.8-6.5	222	5.2	4.5-6.0	148	2.0	1.7-2.4	75	2.5	2.0-3.2	73	1.7	1.3-2.2
1978	368	5.3	4.7-5.8	175	5.9	5.0-6.8	193	4.8	4.2-5.6	121	1.6	1.4-2.0	71	2.5	1.9-3.1	50	1.1	0.8-1.5
1979	412	5.8	5.3-6.4	203	6.5	5.6-7.5	209	5.3	4.6-6.1	143	2.0	1.7-2.4	77	2.6	2.0-3.2	66	1.6	1.2-2.1
1980	429	6.0	5.4-6.6	210	6.8	5.9-7.9	219	5.5	4.8-6.3	152	2.1	1.8-2.5	79	2.7	2.1-3.4	73	1.8	1.4-2.2
1981	398	5.6	5.1-6.2	191	6.3	5.4-7.2	207	5.2	4.5-6.0	170	2.5	2.1-2.9	93	3.2	2.5-3.9	77	1.9	1.5-2.4
1982	397	5.7	5.1-6.3	211	7.2	6.3-8.3	186	4.6	3.9-5.3	136	1.9	1.6-2.2	77	2.6	2.0-3.3	59	1.4	1.0-1.8
1983	444	6.2	5.7-6.9	220	7.5	6.5-8.5	224	5.5	4.7-6.2	154	2.2	1.9-2.6	93	3.3	2.6-4.1	61	1.5	1.1-1.9
1984	495	6.9	6.3-7.5	246	8.1	7.1-9.2	249	6.1	5.3-6.9	170	2.3	2.0-2.7	103	3.4	2.8-4.1	67	1.6	1.2-2.0
1985	444	6.2	5.6-6.8	251	8.3	7.3-9.4	193	4.8	4.1-5.5	181	2.6	2.2-3.0	100	3.5	2.8-4.3	81	2.0	1.6-2.5
1986	451	6.2	5.6-6.8	221	7.4	6.4-8.5	230	5.5	4.8-6.3	161	2.2	1.9-2.6	87	2.9	2.3-3.6	74	1.8	1.4-2.2
1987	440	6.1	5.6-6.8	234	8.0	7.0-9.1	206	5.0	4.3-5.7	152	2.1	1.8-2.5	87	3.0	2.4-3.7	65	1.5	1.2-2.0
1988	412	5.7	5.2-6.3	218	7.4	6.4-8.5	194	4.6	3.9-5.3	139	2.0	1.7-2.3	81	2.9	2.3-3.6	58	1.4	1.1-1.8
1989	416	5.8	5.3-6.4	211	7.2	6.2-8.2	205	4.8	4.2-5.6	123	1.7	1.5-2.1	74	2.6	2.0-3.3	49	1.2	0.9-1.6
1990	421	5.9	5.3-6.5	227	7.7	6.7-8.8	194	4.7	4.1-5.4	125	1.8	1.5-2.1	71	2.5	1.9-3.1	54	1.3	1.0-1.7
1991	447	6.1	5.6-6.8	244	8.1	7.1-9.2	203	4.8	4.1-5.5	126	1.7	1.5-2.1	65	2.3	1.8-2.9	61	1.4	1.1-1.8
1992	367	5.1	4.6-5.7	188	6.3	5.4-7.3	179	4.2	3.6-4.9	104	1.4	1.2-1.7	61	2.0	1.5-2.6	43	1.0	0.7-1.4
1993	398	5.5	5.0-6.1	222	7.6	6.6-8.7	176	4.1	3.5-4.8	103	1.4	1.2-1.8	57	2.0	1.5-2.7	46	1.1	0.8-1.4
1994	463	6.4	5.8-7.0	254	8.6	7.5-9.7	209	5.0	4.3-5.7	118	1.6	1.4-2.0	76	2.6	2.1-3.3	42	1.0	0.7-1.3
1995	470	6.5	5.9-7.1	264	8.7	7.6-9.8	206	4.9	4.2-5.6	119	1.7	1.4-2.0	65	2.2	1.7-2.8	54	1.3	1.0-1.7
1996	491	6.7	6.1-7.3	254	8.5	7.5-9.7	237	5.6	4.9-6.3	123	1.7	1.4-2.0	60	2.0	1.5-2.6	63	1.5	1.1-1.9
1997	547	7.5	6.8-8.1	290	9.9	8.8-11.1	257	6.0	5.3-6.8	116	1.6	1.3-1.9	63	2.1	1.6-2.7	53	1.2	0.9-1.6
1998	565	7.5	6.9-8.2	297	9.6	8.5-10.8	268	6.2	5.4-6.9	135	1.8	1.5-2.2	75	2.6	2.0-3.2	60	1.4	1.0-1.7
1999	527	6.9	6.3-7.5	290	9.2	8.2-10.4	237	5.4	4.7-6.1	116	1.6	1.3-1.9	70	2.2	1.7-2.8	46	1.0	0.8-1.4
2000	625	8.1	7.4-8.7	329	10.5	9.4-11.7	296	6.6	5.9-7.5	138	1.8	1.5-2.2	79	2.7	2.1-3.4	59	1.3	1.0-1.7
2001	668	8.6	7.9-9.2	350	10.8	9.7-12.0	318	7.1	6.3-7.9	118	1.5	1.3-1.8	64	2.2	1.7-2.8	54	1.2	0.9-1.5
2002	673	8.5	7.9-9.2	353	10.9	9.8-12.1	320	7.1	6.3-7.9	110	1.4	1.2-1.7	53	1.7	1.2-2.2	57	1.3	0.9-1.6
2003	585	7.3	6.8-8.0	299	9.1	8.1-10.2	286	6.3	5.6-7.1	119	1.5	1.3-1.8	58	1.9	1.4-2.5	61	1.3	1.0-1.7
2004	804	10.0	9.3-10.7	426	12.8	11.6-14.1	378	8.3	7.5-9.2	105	1.3	1.1-1.6	55	1.7	1.3-2.3	50	1.1	0.8-1.4
2005	890	11.0	10.3-11.8	469	14.0	12.7-15.3	421	9.2	8.4-10.2	105	1.3	1.1-1.6	55	1.8	1.3-2.3	50	1.1	0.8-1.4
2006	1,077	13.3	12.5-14.2	582	17.0	15.7-18.5	495	10.8	9.9-11.8	121	1.6	1.3-1.9	72	2.3	1.8-2.9	49	1.1	0.8-1.4
2007	871	10.7	10.0-11.4	449	13.1	11.9-14.4	422	9.2	8.3-10.1	125	1.5	1.3-1.8	72	2.3	1.8-2.9	53	1.1	0.8-1.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Melanoma of the skin**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	895	10.9	10.2-11.7	481	14.0	12.8-15.3	414	9.0	8.1-9.9	111	1.4	1.1-1.7	63	1.9	1.5-2.5	48	1.0	0.7-1.3
2009	970	11.6	10.9-12.4	520	14.9	13.6-16.2	450	9.5	8.6-10.4	103	1.2	1.0-1.5	66	1.9	1.5-2.5	37	0.8	0.5-1.1
2010	839	10.0	9.3-10.7	447	12.7	11.5-14.0	392	8.2	7.4-9.1	113	1.3	1.1-1.6	62	1.8	1.4-2.3	51	1.0	0.8-1.4
2011	1,028	11.9	11.2-12.7	549	15.1	13.8-16.4	479	10.0	9.1-10.9	96	1.1	0.9-1.4	49	1.4	1.1-1.9	47	0.9	0.7-1.2
2012	886	10.1	9.5-10.8	481	13.0	11.8-14.2	405	8.2	7.5-9.1	137	1.6	1.3-1.8	73	2.0	1.6-2.6	64	1.2	0.9-1.6
2013	994	11.2	10.5-11.9	554	14.7	13.5-16.1	440	8.8	8.0-9.7	116	1.3	1.1-1.5	72	1.9	1.5-2.4	44	0.8	0.6-1.1
2014	925	10.3	9.6-11.0	488	13.0	11.8-14.2	437	8.6	7.8-9.4	110	1.2	1.0-1.5	71	1.9	1.5-2.4	39	0.7	0.5-1.0
2015	916	10.0	9.4-10.7	505	13.2	12.0-14.4	411	7.9	7.2-8.7	81	0.9	0.7-1.1	46	1.2	0.9-1.6	35	0.7	0.5-1.0
2016	940	10.1	9.5-10.8	515	13.0	11.9-14.2	425	8.2	7.4-9.0	91	1.0	0.8-1.2	52	1.3	1.0-1.8	39	0.7	0.5-1.0
2017	919	9.9	9.2-10.5	530	13.2	12.1-14.4	389	7.6	6.8-8.4	99	1.0	0.8-1.3	59	1.5	1.1-2.0	40	0.7	0.5-0.9
2018	920	9.6	9.0-10.2	517	12.4	11.3-13.5	403	7.6	6.8-8.4	78	0.8	0.6-1.0	48	1.2	0.9-1.6	30	0.5	0.3-0.8
2019	1,003	10.4	9.8-11.1	543	13.1	12.0-14.3	460	8.7	7.9-9.6	79	0.8	0.6-1.0	51	1.3	0.9-1.7	28	0.5	0.3-0.7
2020	775	8.0	7.4-8.6	426	10.2	9.2-11.2	349	6.5	5.8-7.2	75	0.8	0.6-1.0	37	0.9	0.7-1.3	38	0.7	0.5-1.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Female breast**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	4,107	99.5	96.4-102.7	1,647	38.9	37.0-40.9
1977	3,943	96.9	93.8-100.1	1,591	38.1	36.2-40.0
1978	4,014	99.3	96.1-102.5	1,597	38.5	36.6-40.5
1979	3,954	98.2	95.0-101.4	1,438	35.1	33.2-37.0
1980	3,956	99.1	95.9-102.3	1,609	38.8	36.9-40.8
1981	4,067	102.1	98.9-105.4	1,564	37.9	36.0-39.9
1982	4,049	101.2	98.0-104.5	1,481	35.7	33.9-37.7
1983	4,060	100.5	97.4-103.8	1,436	34.8	33.0-36.7
1984	4,372	108.3	105.1-111.7	1,527	36.8	34.9-38.7
1985	4,371	107.8	104.5-111.1	1,446	34.6	32.8-36.4
1986	4,271	105.3	102.1-108.6	1,458	34.9	33.1-36.8
1987	4,422	108.2	105.0-111.5	1,439	34.4	32.6-36.3
1988	4,515	110.8	107.5-114.1	1,559	37.0	35.2-39.0
1989	4,445	109.8	106.6-113.2	1,501	36.0	34.2-37.9
1990	4,746	118.2	114.8-121.6	1,470	35.6	33.7-37.5
1991	4,955	123.2	119.7-126.7	1,414	34.2	32.4-36.1
1992	4,894	121.8	118.4-125.3	1,444	35.0	33.2-36.9
1993	4,675	115.6	112.3-119.0	1,429	34.3	32.5-36.1
1994	5,130	126.3	122.8-129.8	1,411	33.8	32.0-35.6
1995	4,995	122.1	118.8-125.6	1,489	35.5	33.7-37.3
1996	5,323	129.6	126.1-133.2	1,475	35.0	33.2-36.9
1997	5,188	125.1	121.7-128.6	1,332	31.4	29.7-33.1
1998	5,403	128.7	125.3-132.2	1,261	29.4	27.8-31.0
1999	5,280	123.6	120.3-127.0	1,229	27.9	26.4-29.6
2000	5,234	120.7	117.4-124.0	1,192	26.5	25.0-28.1
2001	5,059	115.0	111.8-118.2	1,284	28.4	26.8-30.0
2002	5,031	113.2	110.0-116.3	1,151	25.1	23.6-26.6
2003	5,050	113.0	109.9-116.2	1,165	25.2	23.8-26.7
2004	5,034	112.3	109.2-115.4	1,192	25.7	24.2-27.2
2005	4,995	111.0	107.9-114.1	1,180	25.4	24.0-26.9
2006	5,210	115.8	112.6-119.0	1,126	24.1	22.7-25.6
2007	5,389	118.9	115.8-122.2	1,049	22.2	20.9-23.6
2008	5,410	118.2	115.1-121.4	1,037	21.7	20.4-23.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Female breast**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	5,572	120.3	117.2-123.6	1,046	22.0	20.6-23.3
2010	5,445	116.4	113.3-119.6	1,013	20.6	19.3-21.9
2011	5,787	121.8	118.6-125.0	1,027	20.8	19.5-22.1
2012	5,587	115.9	112.9-119.1	1,070	21.2	20.0-22.6
2013	6,000	123.1	120.0-126.4	1,010	19.5	18.3-20.8
2014	6,080	123.0	119.9-126.2	1,038	19.8	18.6-21.1
2015	6,190	123.8	120.7-127.0	990	18.9	17.7-20.1
2016	6,145	121.4	118.4-124.6	1,021	19.0	17.9-20.3
2017	6,458	127.5	124.3-130.7	958	17.6	16.5-18.8
2018	6,647	130.5	127.3-133.8	1,028	18.5	17.4-19.7
2019	6,959	135.2	132.0-138.5	1,032	18.6	17.5-19.8
2020	5,903	116.1	113.1-119.2	899	16.2	15.1-17.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Cervix uteri**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	626	16.2	14.9-17.5	238	5.9	5.1-6.7
1977	610	15.7	14.4-17.1	237	5.9	5.1-6.7
1978	589	16.0	14.7-17.4	218	5.5	4.8-6.3
1979	620	16.6	15.2-18.0	186	4.9	4.2-5.7
1980	566	15.3	14.0-16.7	203	5.1	4.4-5.9
1981	530	13.9	12.7-15.2	195	5.0	4.3-5.8
1982	554	14.9	13.6-16.2	199	5.2	4.4-6.0
1983	546	14.4	13.2-15.7	172	4.2	3.6-4.9
1984	549	14.4	13.2-15.7	147	3.7	3.1-4.4
1985	519	13.3	12.2-14.6	159	4.0	3.4-4.7
1986	516	13.6	12.4-14.8	142	3.6	3.0-4.3
1987	558	14.3	13.1-15.6	135	3.3	2.8-4.0
1988	508	13.1	11.9-14.3	177	4.5	3.8-5.2
1989	500	12.7	11.6-13.9	127	3.1	2.6-3.8
1990	597	15.3	14.1-16.6	163	4.2	3.5-4.9
1991	581	15.1	13.8-16.4	145	3.7	3.1-4.3
1992	593	14.8	13.6-16.1	134	3.3	2.8-3.9
1993	545	13.8	12.6-15.0	160	4.0	3.4-4.7
1994	618	15.5	14.3-16.7	167	4.1	3.5-4.8
1995	604	14.9	13.7-16.1	180	4.4	3.8-5.1
1996	624	15.3	14.2-16.6	163	3.9	3.4-4.6
1997	575	13.9	12.7-15.0	164	4.0	3.4-4.6
1998	547	13.1	12.0-14.2	148	3.5	3.0-4.1
1999	527	12.4	11.4-13.5	178	4.2	3.6-4.9
2000	558	13.0	12.0-14.2	159	3.6	3.1-4.2
2001	454	10.5	9.6-11.5	139	3.2	2.7-3.8
2002	504	11.6	10.6-12.6	142	3.2	2.7-3.8
2003	458	10.4	9.4-11.4	157	3.5	3.0-4.1
2004	450	10.1	9.2-11.1	130	2.9	2.4-3.5
2005	498	11.4	10.4-12.4	145	3.2	2.7-3.7
2006	460	10.3	9.4-11.3	145	3.2	2.7-3.8
2007	461	10.4	9.5-11.4	152	3.4	2.8-3.9
2008	459	10.2	9.3-11.2	133	2.8	2.4-3.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Cervix uteri**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	434	9.7	8.8-10.6	134	2.8	2.4-3.4
2010	412	9.2	8.3-10.1	120	2.5	2.1-3.0
2011	418	9.1	8.3-10.1	148	3.1	2.6-3.7
2012	436	9.5	8.6-10.4	131	2.7	2.3-3.3
2013	443	9.3	8.5-10.3	146	2.9	2.5-3.5
2014	421	9.0	8.1-9.9	152	3.0	2.5-3.5
2015	423	8.9	8.0-9.8	119	2.3	1.9-2.8
2016	434	9.1	8.3-10.1	132	2.6	2.2-3.1
2017	459	9.5	8.7-10.5	117	2.3	1.9-2.8
2018	420	8.6	7.8-9.5	115	2.2	1.8-2.7
2019	428	8.9	8.1-9.9	86	1.7	1.3-2.1
2020	335	6.8	6.1-7.6	104	2.0	1.6-2.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York City

Corpus uteris and uterus, not otherwise specified

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	984	22.6	21.2-24.1	308	6.9	6.2-7.8
1977	866	20.2	18.8-21.6	251	5.7	5.0-6.4
1978	814	18.7	17.4-20.1	247	5.6	4.9-6.4
1979	872	20.3	19.0-21.8	222	5.0	4.4-5.7
1980	941	21.9	20.5-23.4	261	5.9	5.2-6.7
1981	861	20.1	18.7-21.5	225	5.1	4.5-5.8
1982	871	20.5	19.1-21.9	216	4.9	4.2-5.6
1983	898	20.9	19.5-22.4	227	5.1	4.4-5.8
1984	852	19.9	18.5-21.3	225	5.0	4.4-5.8
1985	877	20.4	19.0-21.8	199	4.5	3.9-5.2
1986	793	18.6	17.3-20.0	197	4.4	3.8-5.1
1987	845	19.9	18.5-21.3	202	4.6	4.0-5.3
1988	847	19.8	18.5-21.2	197	4.4	3.8-5.1
1989	801	18.9	17.6-20.2	194	4.4	3.8-5.1
1990	782	18.8	17.5-20.2	208	4.7	4.1-5.4
1991	938	22.7	21.3-24.3	213	5.0	4.3-5.7
1992	902	22.0	20.6-23.5	218	5.0	4.4-5.8
1993	904	21.8	20.4-23.3	233	5.4	4.7-6.1
1994	996	23.9	22.5-25.5	234	5.5	4.8-6.2
1995	979	23.6	22.1-25.1	220	5.1	4.4-5.8
1996	998	23.9	22.4-25.4	237	5.5	4.8-6.2
1997	1,066	25.4	23.9-27.0	256	5.9	5.2-6.7
1998	1,110	26.3	24.8-27.9	250	5.8	5.1-6.6
1999	1,165	27.0	25.4-28.6	248	5.6	4.9-6.4
2000	1,101	25.2	23.7-26.8	226	5.0	4.3-5.7
2001	1,175	26.7	25.2-28.2	239	5.3	4.6-6.0
2002	1,172	26.2	24.8-27.8	225	4.9	4.3-5.6
2003	1,202	26.6	25.1-28.1	255	5.6	4.9-6.3
2004	1,283	28.3	26.7-29.9	244	5.2	4.6-6.0
2005	1,280	27.9	26.4-29.5	268	5.7	5.1-6.5
2006	1,309	28.4	26.9-30.0	273	5.8	5.2-6.6
2007	1,239	26.6	25.2-28.2	265	5.6	5.0-6.4
2008	1,509	32.1	30.5-33.7	268	5.6	4.9-6.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York City

Corpus uteris and uterus, not otherwise specified

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	1,413	29.8	28.2-31.4	265	5.5	4.9-6.2
2010	1,416	29.4	27.8-30.9	310	6.3	5.6-7.1
2011	1,505	30.6	29.0-32.2	324	6.5	5.8-7.2
2012	1,463	29.1	27.6-30.6	309	6.1	5.4-6.8
2013	1,611	31.5	29.9-33.1	313	6.0	5.3-6.7
2014	1,573	30.2	28.7-31.8	348	6.6	5.9-7.3
2015	1,682	32.1	30.5-33.7	382	7.0	6.3-7.7
2016	1,724	32.5	30.9-34.1	403	7.4	6.6-8.1
2017	1,678	31.2	29.7-32.8	368	6.5	5.9-7.3
2018	1,688	31.1	29.6-32.6	362	6.3	5.7-7.0
2019	1,695	31.2	29.7-32.8	413	7.3	6.6-8.0
2020	1,485	26.9	25.5-28.3	387	6.6	6.0-7.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York City

Ovary

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	650	15.1	13.9-16.3	443	10.2	9.3-11.3
1977	610	14.7	13.5-16.0	458	10.5	9.6-11.6
1978	617	14.9	13.7-16.2	399	9.3	8.4-10.3
1979	637	15.8	14.5-17.1	390	9.1	8.2-10.1
1980	635	15.3	14.1-16.6	434	10.2	9.2-11.2
1981	650	15.9	14.6-17.2	387	9.0	8.1-10.0
1982	653	16.0	14.8-17.3	373	8.8	7.9-9.7
1983	677	16.5	15.2-17.8	395	9.3	8.3-10.2
1984	659	15.9	14.7-17.2	403	9.4	8.5-10.4
1985	688	16.6	15.3-17.9	384	9.1	8.2-10.1
1986	605	14.6	13.5-15.9	395	9.2	8.3-10.1
1987	629	15.4	14.2-16.7	376	8.8	7.9-9.7
1988	609	14.8	13.6-16.1	417	9.7	8.7-10.7
1989	603	14.5	13.4-15.8	346	8.1	7.2-9.0
1990	639	15.6	14.4-16.8	381	8.9	8.0-9.9
1991	673	16.5	15.3-17.8	410	9.8	8.9-10.8
1992	682	16.8	15.6-18.2	434	10.4	9.5-11.5
1993	624	15.3	14.1-16.6	390	9.3	8.4-10.3
1994	690	16.6	15.4-17.9	378	8.9	8.0-9.8
1995	719	17.5	16.2-18.8	394	9.3	8.4-10.3
1996	639	15.4	14.2-16.6	356	8.5	7.6-9.4
1997	673	16.1	14.9-17.4	381	8.9	8.1-9.9
1998	603	14.4	13.3-15.6	344	8.0	7.2-8.9
1999	612	14.3	13.2-15.4	366	8.4	7.5-9.3
2000	579	13.2	12.2-14.4	359	8.0	7.2-8.9
2001	643	14.6	13.5-15.8	335	7.4	6.7-8.3
2002	647	14.4	13.3-15.6	368	8.0	7.2-8.9
2003	571	12.6	11.6-13.7	364	7.9	7.1-8.8
2004	618	13.7	12.6-14.8	363	7.8	7.0-8.7
2005	563	12.5	11.5-13.6	355	7.7	6.9-8.6
2006	549	12.1	11.1-13.1	351	7.5	6.7-8.4
2007	588	12.8	11.7-13.9	337	7.2	6.4-8.0
2008	580	12.4	11.4-13.5	340	7.1	6.3-7.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Ovary**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	618	13.3	12.2-14.4	330	6.8	6.0-7.5
2010	578	12.3	11.3-13.4	341	6.9	6.2-7.7
2011	622	12.9	11.9-13.9	350	7.0	6.3-7.8
2012	573	11.7	10.7-12.7	367	7.3	6.6-8.1
2013	591	12.0	11.0-13.0	346	6.7	6.0-7.4
2014	598	11.9	10.9-12.9	319	6.1	5.4-6.8
2015	607	12.1	11.1-13.1	320	6.1	5.5-6.9
2016	575	11.3	10.4-12.3	341	6.2	5.6-7.0
2017	615	11.9	10.9-12.9	348	6.3	5.7-7.1
2018	542	10.6	9.7-11.5	330	5.9	5.3-6.6
2019	557	11.0	10.1-12.0	351	6.3	5.6-7.0
2020	551	10.8	9.8-11.7	288	5.0	4.5-5.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Prostate**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	2,001	73.5	70.1-77.0	736	29.5	27.3-31.9
1977	2,018	73.7	70.3-77.1	727	28.4	26.3-30.6
1978	2,037	75.1	71.7-78.6	766	30.4	28.2-32.7
1979	2,209	82.0	78.5-85.7	728	29.6	27.4-31.9
1980	2,435	89.9	86.2-93.7	772	31.3	29.0-33.7
1981	2,320	85.5	81.9-89.2	800	32.5	30.2-34.9
1982	2,251	83.8	80.3-87.5	731	29.5	27.4-31.8
1983	2,278	83.5	79.9-87.1	814	32.7	30.4-35.1
1984	2,401	88.0	84.4-91.7	850	34.3	31.9-36.7
1985	2,491	89.8	86.2-93.5	798	31.1	28.9-33.4
1986	2,534	93.1	89.4-96.9	784	30.9	28.7-33.2
1987	2,536	91.8	88.1-95.5	821	32.2	30.0-34.6
1988	2,490	92.6	88.9-96.4	835	33.4	31.1-35.8
1989	2,540	93.5	89.8-97.3	823	32.8	30.6-35.2
1990	3,049	112.3	108.2-116.5	923	37.4	35.0-39.9
1991	3,458	125.8	121.5-130.1	830	33.6	31.3-36.0
1992	4,305	154.2	149.6-159.0	976	39.2	36.7-41.8
1993	4,248	150.2	145.7-154.9	925	37.4	35.0-40.0
1994	4,557	159.3	154.6-164.1	931	37.8	35.4-40.4
1995	4,327	150.1	145.6-154.7	948	37.9	35.4-40.4
1996	4,358	150.7	146.2-155.3	870	34.7	32.4-37.1
1997	4,680	160.2	155.6-164.9	820	32.5	30.3-34.9
1998	4,750	159.9	155.3-164.5	849	32.9	30.7-35.2
1999	5,125	169.6	164.9-174.3	770	29.3	27.2-31.4
2000	5,855	189.6	184.8-194.6	806	30.2	28.1-32.3
2001	5,496	175.1	170.4-179.8	807	29.8	27.8-32.0
2002	5,481	171.6	167.0-176.2	825	29.9	27.9-32.0
2003	5,083	157.9	153.5-162.3	761	27.3	25.3-29.3
2004	5,135	156.1	151.8-160.4	730	26.1	24.2-28.1
2005	5,199	156.0	151.7-160.4	691	24.2	22.4-26.1
2006	5,716	171.0	166.5-175.6	671	23.6	21.8-25.5
2007	6,050	177.2	172.7-181.8	707	24.3	22.6-26.2
2008	5,887	167.9	163.6-172.4	666	22.8	21.1-24.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Prostate**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	5,856	163.6	159.4-167.9	717	24.1	22.3-25.9
2010	5,517	149.9	145.9-154.0	736	24.0	22.3-25.8
2011	6,090	160.9	156.8-165.1	720	22.8	21.2-24.6
2012	5,386	138.1	134.4-141.9	663	20.5	18.9-22.1
2013	5,425	135.9	132.2-139.6	653	19.7	18.2-21.3
2014	5,033	122.3	118.9-125.8	706	20.3	18.8-21.9
2015	5,143	121.8	118.4-125.3	658	18.3	17.0-19.8
2016	5,385	125.2	121.8-128.6	696	19.1	17.7-20.5
2017	5,595	127.4	124.0-130.8	714	18.9	17.5-20.3
2018	5,687	127.6	124.3-131.1	674	17.3	16.0-18.7
2019	6,129	135.4	131.9-138.9	686	17.2	15.9-18.5
2020	5,006	109.1	106.0-112.2	573	14.0	12.9-15.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Testis**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	84	2.3	1.8-2.9	19	0.5	0.3-0.9
1977	79	2.3	1.8-2.8	18	0.5	0.3-0.8
1978	74	2.1	1.7-2.7	18	0.5	0.3-0.9
1979	88	2.7	2.1-3.3	16	0.5	0.3-0.8
1980	106	3.2	2.6-3.9	17	0.5	0.3-0.9
1981	104	2.9	2.3-3.5	6	0.2	0.1-0.4
1982	116	3.2	2.6-3.9	7	0.2	0.1-0.4
1983	108	3.0	2.4-3.6	14	0.4	0.2-0.7
1984	101	2.8	2.2-3.4	11	0.4	0.2-0.7
1985	113	3.1	2.5-3.8	15	0.4	0.2-0.7
1986	124	3.2	2.6-3.8	11	0.3	0.2-0.6
1987	119	3.0	2.5-3.7	8	0.2	0.1-0.4
1988	108	2.8	2.3-3.4	10	0.3	0.1-0.5
1989	122	3.2	2.6-3.8	12	0.3	0.2-0.6
1990	137	3.6	3.0-4.3	12	0.3	0.2-0.6
1991	135	3.4	2.8-4.1	9	0.3	0.1-0.5
1992	143	3.7	3.1-4.4	11	0.3	0.2-0.6
1993	139	3.5	2.9-4.2	11	0.3	0.2-0.6
1994	138	3.5	2.9-4.2	10	0.3	0.1-0.6
1995	117	2.9	2.4-3.5	7	0.2	0.1-0.4
1996	131	3.3	2.7-3.9	6	0.2	0.1-0.4
1997	114	2.8	2.3-3.4	7	0.2	0.1-0.4
1998	130	3.2	2.7-3.8	9	0.2	0.1-0.4
1999	176	4.2	3.6-4.9	4	0.1	0.0-0.3
2000	164	4.0	3.4-4.7	5	0.2	0.0-0.4
2001	145	3.4	2.9-4.1	7	0.2	0.1-0.4
2002	153	3.7	3.1-4.3	7	0.2	0.1-0.4
2003	167	4.0	3.4-4.7	11	0.3	0.1-0.5
2004	173	4.2	3.6-4.8	9	0.2	0.1-0.4
2005	189	4.5	3.9-5.3	11	0.3	0.1-0.5
2006	186	4.5	3.9-5.2	12	0.3	0.2-0.6
2007	202	4.9	4.2-5.6	3	0.1	0.0-0.2
2008	177	4.3	3.7-5.0	5	0.1	0.0-0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Testis**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	208	5.0	4.3-5.8	9	0.2	0.1-0.4
2010	199	4.8	4.2-5.5	15	0.4	0.2-0.6
2011	208	4.9	4.2-5.6	11	0.3	0.1-0.5
2012	207	4.9	4.2-5.6	9	0.2	0.1-0.4
2013	201	4.6	4.0-5.3	7	0.2	0.1-0.3
2014	201	4.7	4.1-5.4	4	0.1	0.0-0.2
2015	208	4.7	4.1-5.4	6	0.1	0.0-0.3
2016	215	4.9	4.3-5.7	6	0.2	0.1-0.4
2017	229	5.2	4.5-5.9	11	0.3	0.1-0.5
2018	219	5.3	4.6-6.0	8	0.2	0.1-0.4
2019	236	5.5	4.8-6.3	2	0.1	0.0-0.2
2020	228	5.6	4.8-6.3	8	0.2	0.1-0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Urinary bladder (incl. in situ)**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,305	17.7	16.7-18.7	928	31.9	29.7-34.1	377	8.6	7.7-9.6	428	6.4	5.8-7.0	288	11.0	9.7-12.5	140	3.5	3.0-4.2
1977	1,379	18.8	17.8-19.9	966	33.4	31.2-35.7	413	9.5	8.6-10.5	438	6.4	5.8-7.0	299	11.5	10.2-13.0	139	3.3	2.8-3.9
1978	1,437	19.9	18.8-20.9	1,031	36.4	34.1-38.8	406	9.3	8.4-10.2	398	5.8	5.2-6.4	262	10.0	8.8-11.4	136	3.3	2.8-4.0
1979	1,454	20.3	19.2-21.4	1,003	36.1	33.8-38.5	451	10.5	9.5-11.5	404	5.9	5.3-6.5	271	10.5	9.3-11.9	133	3.1	2.6-3.7
1980	1,552	21.4	20.3-22.5	1,146	40.7	38.3-43.2	406	9.3	8.4-10.3	409	5.9	5.3-6.5	276	10.6	9.3-12.0	133	3.1	2.6-3.7
1981	1,488	20.6	19.6-21.7	1,065	38.1	35.8-40.6	423	9.7	8.8-10.7	364	5.2	4.7-5.8	236	9.0	7.9-10.3	128	3.0	2.5-3.6
1982	1,394	19.1	18.1-20.2	967	34.3	32.1-36.7	427	9.7	8.8-10.7	378	5.4	4.8-5.9	254	9.8	8.6-11.1	124	2.8	2.3-3.4
1983	1,537	21.2	20.1-22.3	1,065	38.7	36.3-41.2	472	10.5	9.6-11.5	340	4.8	4.3-5.4	216	8.6	7.4-9.8	124	2.8	2.3-3.3
1984	1,401	19.0	18.0-20.0	1,005	35.2	33.0-37.5	396	8.9	8.0-9.8	343	4.8	4.3-5.4	225	8.5	7.4-9.7	118	2.7	2.2-3.2
1985	1,299	17.9	16.9-18.9	900	32.3	30.1-34.5	399	9.0	8.1-9.9	358	5.1	4.6-5.6	238	9.4	8.2-10.8	120	2.7	2.2-3.2
1986	1,368	18.9	17.9-20.0	961	35.0	32.7-37.3	407	9.3	8.4-10.3	323	4.5	4.0-5.1	199	7.9	6.8-9.1	124	2.7	2.3-3.3
1987	1,378	18.8	17.8-19.8	977	34.9	32.7-37.3	401	9.0	8.1-10.0	315	4.4	3.9-4.9	212	8.3	7.2-9.5	103	2.3	1.9-2.8
1988	1,345	18.4	17.4-19.4	926	32.9	30.8-35.2	419	9.4	8.5-10.3	342	4.7	4.3-5.3	210	8.1	7.0-9.3	132	2.9	2.4-3.4
1989	1,265	17.6	16.7-18.6	883	32.3	30.1-34.5	382	8.7	7.8-9.6	343	4.9	4.4-5.4	215	8.5	7.4-9.8	128	2.9	2.4-3.4
1990	1,310	18.3	17.3-19.3	907	33.3	31.1-35.6	403	9.1	8.2-10.0	315	4.5	4.0-5.0	196	7.8	6.7-9.0	119	2.7	2.2-3.3
1991	1,384	19.4	18.4-20.5	970	35.7	33.4-38.0	414	9.6	8.7-10.6	289	4.1	3.6-4.6	188	7.4	6.4-8.6	101	2.3	1.8-2.7
1992	1,394	19.5	18.5-20.6	950	35.0	32.8-37.3	444	10.3	9.3-11.3	302	4.3	3.8-4.8	192	7.7	6.7-8.9	110	2.4	2.0-2.9
1993	1,304	18.3	17.3-19.3	907	33.1	30.9-35.3	397	9.2	8.3-10.1	354	5.0	4.5-5.6	236	9.2	8.0-10.5	118	2.7	2.2-3.2
1994	1,414	19.7	18.7-20.8	956	34.8	32.5-37.1	458	10.5	9.6-11.5	325	4.6	4.1-5.1	209	8.3	7.2-9.6	116	2.6	2.1-3.1
1995	1,354	18.9	17.9-19.9	957	34.9	32.6-37.2	397	9.1	8.2-10.0	349	4.9	4.4-5.5	216	8.4	7.3-9.6	133	2.9	2.4-3.5
1996	1,225	17.1	16.2-18.1	860	31.3	29.2-33.5	365	8.3	7.5-9.2	305	4.3	3.8-4.8	187	7.3	6.3-8.5	118	2.6	2.1-3.1
1997	1,403	19.5	18.5-20.5	977	34.6	32.5-36.9	426	9.7	8.8-10.7	301	4.2	3.7-4.7	190	7.4	6.4-8.6	111	2.4	2.0-3.0
1998	1,440	19.8	18.8-20.9	986	34.5	32.4-36.8	454	10.2	9.3-11.2	326	4.5	4.0-5.0	197	7.4	6.4-8.5	129	2.8	2.3-3.3
1999	1,343	18.1	17.1-19.1	937	32.4	30.3-34.6	406	9.0	8.2-10.0	303	4.1	3.7-4.6	200	7.5	6.5-8.6	103	2.2	1.8-2.7
2000	1,427	19.0	18.0-20.0	984	33.8	31.7-36.0	443	9.7	8.8-10.7	308	4.1	3.6-4.6	198	7.1	6.1-8.2	110	2.3	1.9-2.8
2001	1,378	18.1	17.1-19.1	966	32.4	30.4-34.6	412	9.0	8.1-9.9	314	4.1	3.7-4.6	202	7.2	6.2-8.2	112	2.3	1.9-2.8
2002	1,327	17.3	16.3-18.2	934	31.3	29.3-33.4	393	8.4	7.6-9.3	273	3.5	3.1-4.0	168	6.0	5.1-7.0	105	2.2	1.8-2.6
2003	1,404	18.2	17.2-19.1	1,004	33.2	31.1-35.3	400	8.5	7.7-9.4	258	3.3	2.9-3.8	171	6.1	5.2-7.0	87	1.8	1.4-2.2
2004	1,438	18.4	17.4-19.3	1,004	32.4	30.4-34.5	434	9.1	8.3-10.0	295	3.8	3.4-4.2	194	6.6	5.7-7.6	101	2.0	1.7-2.5
2005	1,451	18.5	17.5-19.4	1,004	32.0	30.0-34.1	447	9.4	8.6-10.4	296	3.8	3.4-4.3	184	6.4	5.5-7.4	112	2.3	1.9-2.7
2006	1,427	18.1	17.2-19.1	1,052	33.8	31.7-35.9	375	7.9	7.1-8.8	300	3.8	3.4-4.3	194	6.7	5.7-7.7	106	2.1	1.7-2.6
2007	1,464	18.5	17.6-19.5	1,047	33.5	31.4-35.6	417	8.8	8.0-9.7	304	3.8	3.4-4.3	195	6.6	5.7-7.6	109	2.2	1.8-2.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Urinary bladder (incl. in situ)**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	1,473	18.3	17.4-19.3	1,102	34.0	32.0-36.1	371	7.8	7.0-8.6	289	3.6	3.2-4.1	196	6.5	5.6-7.5	93	1.9	1.5-2.3
2009	1,452	17.9	16.9-18.8	1,034	31.8	29.9-33.9	418	8.6	7.8-9.5	279	3.4	3.0-3.8	176	5.8	5.0-6.8	103	2.0	1.6-2.4
2010	1,416	17.1	16.2-18.0	1,055	31.8	29.9-33.8	361	7.4	6.6-8.2	323	3.9	3.5-4.3	214	6.7	5.8-7.7	109	2.1	1.7-2.6
2011	1,468	17.4	16.5-18.3	1,069	31.5	29.6-33.5	399	8.0	7.2-8.8	308	3.6	3.2-4.1	197	6.1	5.3-7.0	111	2.2	1.8-2.6
2012	1,448	16.7	15.9-17.6	1,044	30.1	28.3-32.0	404	7.8	7.0-8.6	321	3.7	3.3-4.1	214	6.3	5.5-7.2	107	2.0	1.6-2.4
2013	1,506	17.1	16.2-18.0	1,093	30.4	28.6-32.3	413	8.0	7.2-8.8	298	3.3	2.9-3.7	203	6.0	5.2-6.9	95	1.7	1.3-2.0
2014	1,493	16.5	15.7-17.4	1,087	29.5	27.8-31.4	406	7.6	6.8-8.4	342	3.8	3.4-4.2	229	6.7	5.9-7.7	113	1.9	1.6-2.4
2015	1,627	17.6	16.8-18.5	1,226	32.4	30.6-34.3	401	7.5	6.7-8.2	317	3.4	3.0-3.8	207	5.8	5.0-6.6	110	1.9	1.6-2.3
2016	1,532	16.2	15.4-17.1	1,153	29.9	28.1-31.7	379	6.8	6.2-7.6	317	3.4	3.0-3.8	216	6.0	5.2-6.8	101	1.8	1.4-2.1
2017	1,647	17.2	16.3-18.0	1,197	30.1	28.4-31.9	450	8.1	7.4-8.9	313	3.3	2.9-3.6	221	5.9	5.2-6.8	92	1.6	1.3-2.0
2018	1,560	15.9	15.1-16.7	1,153	28.3	26.6-30.0	407	7.2	6.5-7.9	300	3.0	2.7-3.4	195	5.0	4.3-5.8	105	1.8	1.4-2.1
2019	1,621	16.3	15.5-17.2	1,165	28.0	26.4-29.7	456	8.0	7.2-8.7	306	3.0	2.6-3.3	215	5.3	4.6-6.1	91	1.5	1.2-1.8
2020	1,351	13.4	12.7-14.2	989	23.3	21.9-24.9	362	6.3	5.7-7.0	237	2.3	2.0-2.6	146	3.6	3.0-4.2	91	1.5	1.2-1.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Kidney and renal pelvis**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	536	7.2	6.6-7.9	330	10.6	9.4-11.8	206	4.8	4.2-5.5	286	3.9	3.4-4.4	187	6.3	5.4-7.3	99	2.3	1.8-2.8
1977	510	6.9	6.3-7.5	333	10.8	9.7-12.1	177	4.1	3.5-4.8	259	3.5	3.1-4.0	168	5.7	4.8-6.6	91	2.1	1.7-2.6
1978	533	7.2	6.6-7.9	331	11.0	9.8-12.2	202	4.6	4.0-5.3	264	3.6	3.2-4.1	176	5.8	5.0-6.8	88	2.1	1.7-2.6
1979	496	6.7	6.1-7.3	316	10.5	9.3-11.7	180	4.0	3.5-4.7	264	3.7	3.3-4.2	170	5.9	5.0-6.9	94	2.2	1.8-2.7
1980	492	6.6	6.1-7.3	302	9.7	8.6-10.9	190	4.6	3.9-5.3	219	3.0	2.6-3.4	139	4.7	3.9-5.6	80	1.8	1.4-2.3
1981	490	6.7	6.1-7.4	292	10.0	8.8-11.2	198	4.6	3.9-5.3	236	3.2	2.8-3.7	139	4.7	3.9-5.5	97	2.2	1.8-2.7
1982	548	7.7	7.0-8.4	329	11.3	10.1-12.6	219	5.2	4.5-5.9	240	3.3	2.9-3.8	142	5.1	4.3-6.0	98	2.2	1.8-2.7
1983	547	7.6	6.9-8.2	329	11.1	9.9-12.5	218	5.2	4.6-6.0	234	3.2	2.8-3.6	143	4.9	4.1-5.8	91	2.1	1.7-2.6
1984	588	8.1	7.4-8.8	367	12.3	11.1-13.7	221	5.2	4.5-5.9	246	3.4	3.0-3.9	149	5.2	4.4-6.1	97	2.3	1.8-2.8
1985	606	8.3	7.6-9.0	365	12.2	11.0-13.6	241	5.6	4.9-6.4	234	3.2	2.8-3.7	132	4.6	3.8-5.5	102	2.4	1.9-2.9
1986	604	8.2	7.5-8.9	361	12.0	10.8-13.3	243	5.5	4.8-6.3	261	3.5	3.1-4.0	160	5.6	4.7-6.5	101	2.3	1.8-2.8
1987	676	9.3	8.6-10.0	406	13.7	12.3-15.1	270	6.4	5.6-7.2	236	3.2	2.8-3.7	146	5.1	4.3-6.0	90	2.1	1.7-2.6
1988	575	8.1	7.4-8.8	350	12.0	10.8-13.4	225	5.4	4.7-6.2	224	3.1	2.7-3.6	134	4.8	4.0-5.7	90	2.1	1.7-2.6
1989	613	8.5	7.8-9.2	371	12.6	11.4-14.0	242	5.7	5.0-6.4	233	3.3	2.9-3.7	130	4.7	3.9-5.6	103	2.4	1.9-2.9
1990	652	9.1	8.4-9.8	370	12.7	11.4-14.1	282	6.7	5.9-7.5	242	3.4	3.0-3.8	134	4.9	4.1-5.8	108	2.5	2.0-3.0
1991	690	9.6	8.9-10.4	403	13.7	12.4-15.2	287	6.8	6.0-7.6	219	3.0	2.7-3.5	123	4.2	3.5-5.1	96	2.2	1.8-2.7
1992	679	9.5	8.8-10.3	411	14.1	12.7-15.5	268	6.4	5.6-7.2	201	2.8	2.5-3.3	112	4.2	3.4-5.0	89	2.0	1.6-2.5
1993	699	9.8	9.1-10.5	451	15.3	13.9-16.8	248	5.9	5.2-6.7	229	3.2	2.8-3.6	156	5.4	4.6-6.3	73	1.7	1.3-2.1
1994	761	10.7	9.9-11.5	454	15.5	14.1-17.0	307	7.3	6.5-8.1	237	3.3	2.9-3.7	145	5.1	4.3-6.0	92	2.1	1.7-2.6
1995	718	10.0	9.3-10.7	429	14.4	13.1-15.9	289	6.9	6.1-7.8	243	3.4	3.0-3.9	150	5.3	4.5-6.2	93	2.2	1.7-2.6
1996	810	11.2	10.4-12.0	492	16.6	15.2-18.2	318	7.5	6.7-8.3	226	3.1	2.7-3.6	130	4.5	3.8-5.4	96	2.2	1.8-2.7
1997	799	11.0	10.3-11.8	461	15.6	14.2-17.1	338	8.0	7.1-8.9	240	3.3	2.9-3.8	141	5.0	4.2-5.9	99	2.3	1.8-2.8
1998	833	11.4	10.6-12.2	503	16.4	15.0-17.9	330	7.7	6.9-8.6	225	3.1	2.7-3.5	129	4.5	3.7-5.3	96	2.2	1.8-2.7
1999	911	12.2	11.4-13.0	548	17.5	16.0-19.0	363	8.3	7.5-9.3	202	2.7	2.4-3.1	124	4.1	3.4-4.9	78	1.7	1.4-2.1
2000	893	11.8	11.0-12.6	513	16.3	14.9-17.7	380	8.6	7.7-9.5	210	2.8	2.4-3.2	133	4.4	3.7-5.2	77	1.7	1.3-2.1
2001	932	12.1	11.4-12.9	568	17.5	16.1-19.0	364	8.2	7.3-9.0	220	2.9	2.5-3.3	126	4.1	3.4-4.9	94	2.1	1.7-2.6
2002	929	11.9	11.2-12.7	576	17.3	15.9-18.8	353	7.8	7.0-8.7	227	2.9	2.6-3.4	144	4.6	3.9-5.4	83	1.8	1.4-2.2
2003	1,032	13.2	12.4-14.0	653	19.6	18.1-21.1	379	8.4	7.6-9.3	213	2.7	2.4-3.1	132	4.2	3.5-5.0	81	1.7	1.4-2.1
2004	1,053	13.3	12.5-14.1	652	19.4	17.9-20.9	401	8.8	8.0-9.7	228	2.9	2.6-3.3	139	4.4	3.7-5.2	89	1.9	1.5-2.3
2005	1,075	13.5	12.7-14.4	638	18.9	17.4-20.4	437	9.6	8.7-10.5	186	2.4	2.0-2.7	109	3.4	2.7-4.1	77	1.6	1.3-2.0
2006	1,158	14.6	13.8-15.5	723	21.4	19.8-23.0	435	9.6	8.7-10.5	226	2.9	2.5-3.3	128	4.1	3.4-4.8	98	2.1	1.7-2.6
2007	1,135	14.2	13.4-15.0	727	21.2	19.7-22.8	408	8.9	8.1-9.9	229	2.9	2.5-3.3	145	4.5	3.8-5.3	84	1.7	1.4-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Kidney and renal pelvis**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	1,191	14.6	13.8-15.4	739	21.1	19.6-22.7	452	9.7	8.8-10.6	202	2.5	2.1-2.8	120	3.6	3.0-4.4	82	1.7	1.3-2.1
2009	1,124	13.6	12.8-14.4	687	19.3	17.8-20.8	437	9.3	8.4-10.2	247	3.0	2.6-3.4	172	5.2	4.4-6.0	75	1.5	1.2-1.9
2010	1,189	14.1	13.3-14.9	762	20.8	19.3-22.3	427	9.1	8.2-10.0	231	2.7	2.4-3.1	144	4.3	3.6-5.0	87	1.7	1.4-2.1
2011	1,204	14.1	13.3-14.9	749	20.0	18.5-21.5	455	9.5	8.6-10.4	205	2.3	2.0-2.7	132	3.7	3.1-4.4	73	1.4	1.1-1.7
2012	1,220	14.0	13.3-14.9	769	20.1	18.7-21.6	451	9.3	8.5-10.2	206	2.4	2.1-2.7	130	3.6	3.0-4.3	76	1.5	1.1-1.8
2013	1,303	14.6	13.8-15.4	820	20.9	19.5-22.4	483	9.7	8.8-10.6	205	2.3	2.0-2.6	120	3.3	2.7-3.9	85	1.6	1.2-1.9
2014	1,304	14.3	13.5-15.1	864	21.4	19.9-22.9	440	8.7	7.9-9.6	222	2.4	2.1-2.8	132	3.5	2.9-4.2	90	1.7	1.4-2.1
2015	1,344	14.4	13.6-15.2	842	20.3	18.9-21.7	502	9.6	8.8-10.5	261	2.8	2.5-3.2	176	4.6	3.9-5.3	85	1.5	1.2-1.9
2016	1,376	14.7	13.9-15.5	864	20.8	19.5-22.3	512	9.9	9.0-10.8	198	2.1	1.8-2.4	131	3.4	2.8-4.0	67	1.2	0.9-1.6
2017	1,433	15.0	14.2-15.8	903	21.2	19.9-22.7	530	10.0	9.1-10.9	227	2.4	2.1-2.7	154	3.9	3.3-4.5	73	1.3	1.0-1.7
2018	1,383	14.4	13.7-15.2	911	21.5	20.1-23.0	472	8.8	8.0-9.7	227	2.3	2.0-2.6	156	3.8	3.2-4.4	71	1.2	0.9-1.5
2019	1,519	15.8	15.0-16.6	970	22.5	21.0-24.0	549	10.5	9.6-11.4	207	2.0	1.8-2.3	126	2.9	2.4-3.5	81	1.4	1.1-1.7
2020	1,232	12.8	12.1-13.5	784	18.2	16.9-19.5	448	8.3	7.5-9.1	208	2.1	1.9-2.5	132	3.3	2.7-3.9	76	1.3	1.0-1.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Brain and other central nervous system**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	405	5.2	4.7-5.8	219	6.4	5.6-7.4	186	4.3	3.7-5.0	314	4.2	3.7-4.7	175	5.5	4.7-6.4	139	3.3	2.8-3.9
1977	412	5.6	5.0-6.1	223	6.8	5.9-7.8	189	4.6	4.0-5.3	307	4.1	3.6-4.6	161	4.9	4.1-5.7	146	3.5	2.9-4.2
1978	392	5.4	4.8-5.9	199	6.2	5.3-7.2	193	4.7	4.0-5.4	294	3.9	3.4-4.4	152	4.7	4.0-5.6	142	3.2	2.7-3.8
1979	394	5.3	4.8-5.9	202	6.3	5.5-7.3	192	4.7	4.0-5.4	285	3.8	3.4-4.3	156	5.1	4.3-6.0	129	2.9	2.4-3.5
1980	401	5.5	5.0-6.1	205	6.6	5.7-7.6	196	4.7	4.0-5.4	309	4.1	3.6-4.6	168	5.3	4.5-6.2	141	3.2	2.7-3.8
1981	398	5.6	5.0-6.2	224	7.4	6.4-8.4	174	4.2	3.6-4.9	284	3.9	3.4-4.4	144	4.7	3.9-5.6	140	3.3	2.7-3.9
1982	414	5.7	5.1-6.3	222	7.1	6.1-8.1	192	4.6	4.0-5.4	313	4.3	3.8-4.8	163	5.4	4.6-6.3	150	3.5	3.0-4.2
1983	389	5.3	4.8-5.9	208	6.5	5.6-7.4	181	4.4	3.8-5.1	292	4.0	3.5-4.5	151	5.0	4.2-5.9	141	3.3	2.7-3.9
1984	431	5.8	5.3-6.4	207	6.5	5.6-7.5	224	5.3	4.6-6.1	290	3.9	3.4-4.3	152	4.9	4.1-5.8	138	3.1	2.6-3.7
1985	450	6.3	5.7-6.9	236	7.7	6.7-8.8	214	5.2	4.5-6.0	298	4.1	3.6-4.6	151	5.0	4.2-5.9	147	3.4	2.8-4.0
1986	413	5.7	5.2-6.3	239	7.7	6.8-8.8	174	4.3	3.6-5.0	312	4.3	3.8-4.8	174	5.9	5.0-6.9	138	3.2	2.7-3.8
1987	461	6.3	5.7-6.9	235	7.4	6.5-8.4	226	5.4	4.7-6.2	287	4.0	3.5-4.5	149	4.9	4.2-5.8	138	3.3	2.7-3.9
1988	454	6.2	5.7-6.8	236	7.5	6.6-8.6	218	5.3	4.6-6.1	224	3.1	2.7-3.5	118	3.9	3.2-4.6	106	2.5	2.0-3.0
1989	476	6.7	6.1-7.3	253	8.2	7.2-9.4	223	5.5	4.8-6.3	230	3.2	2.8-3.6	120	4.0	3.3-4.8	110	2.5	2.1-3.0
1990	475	6.7	6.1-7.3	237	7.7	6.7-8.8	238	6.0	5.2-6.8	246	3.5	3.1-4.0	136	4.8	4.0-5.7	110	2.7	2.2-3.2
1991	488	6.8	6.2-7.5	238	7.7	6.7-8.7	250	6.1	5.4-6.9	267	3.7	3.3-4.2	134	4.4	3.7-5.2	133	3.2	2.7-3.8
1992	485	6.7	6.1-7.3	259	8.0	7.1-9.1	226	5.6	4.9-6.4	226	3.2	2.8-3.6	118	4.0	3.3-4.9	108	2.5	2.1-3.1
1993	469	6.4	5.8-7.0	238	7.6	6.6-8.6	231	5.6	4.9-6.4	226	3.1	2.7-3.6	121	4.0	3.3-4.8	105	2.6	2.1-3.1
1994	516	7.0	6.4-7.7	263	8.1	7.1-9.2	253	6.1	5.4-6.9	248	3.4	3.0-3.9	123	4.0	3.3-4.8	125	3.0	2.5-3.6
1995	531	7.2	6.6-7.9	278	8.7	7.6-9.8	253	6.1	5.4-7.0	245	3.4	3.0-3.9	127	4.2	3.5-5.0	118	2.8	2.3-3.4
1996	561	7.6	7.0-8.3	276	8.5	7.5-9.5	285	6.9	6.1-7.7	230	3.2	2.8-3.6	114	3.7	3.1-4.5	116	2.8	2.3-3.3
1997	489	6.6	6.0-7.2	238	7.2	6.3-8.3	251	6.0	5.3-6.8	262	3.6	3.2-4.0	112	3.6	3.0-4.4	150	3.5	3.0-4.2
1998	573	7.5	6.9-8.2	299	8.8	7.9-9.9	274	6.6	5.8-7.4	258	3.5	3.1-3.9	123	3.9	3.2-4.7	135	3.1	2.6-3.7
1999	478	6.2	5.7-6.8	243	7.1	6.2-8.1	235	5.5	4.8-6.2	254	3.4	3.0-3.8	134	4.2	3.5-5.0	120	2.8	2.3-3.3
2000	503	6.5	5.9-7.1	246	7.2	6.3-8.2	257	5.9	5.2-6.7	234	3.1	2.7-3.5	117	3.5	2.9-4.3	117	2.7	2.2-3.2
2001	479	6.1	5.6-6.7	231	6.7	5.9-7.6	248	5.7	5.0-6.5	227	3.0	2.6-3.4	125	3.8	3.2-4.5	102	2.3	1.9-2.8
2002	480	6.1	5.6-6.7	251	7.3	6.4-8.3	229	5.2	4.6-6.0	228	2.9	2.6-3.3	113	3.4	2.8-4.1	115	2.6	2.2-3.1
2003	482	6.1	5.5-6.6	240	6.7	5.9-7.7	242	5.4	4.8-6.2	210	2.6	2.3-3.0	124	3.6	3.0-4.3	86	1.9	1.5-2.3
2004	465	5.9	5.4-6.4	258	7.4	6.6-8.4	207	4.7	4.1-5.4	218	2.8	2.4-3.2	116	3.4	2.8-4.1	102	2.3	1.8-2.8
2005	501	6.3	5.7-6.9	272	7.7	6.8-8.7	229	5.2	4.6-6.0	246	3.1	2.7-3.5	142	4.2	3.5-5.0	104	2.3	1.9-2.8
2006	497	6.2	5.7-6.8	264	7.5	6.6-8.4	233	5.3	4.6-6.0	250	3.1	2.8-3.6	139	4.1	3.4-4.8	111	2.4	2.0-3.0
2007	465	5.8	5.3-6.4	238	6.6	5.8-7.5	227	5.2	4.5-5.9	256	3.2	2.8-3.6	132	3.8	3.2-4.5	124	2.7	2.2-3.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Brain and other central nervous system**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	490	6.1	5.6-6.7	256	7.1	6.3-8.1	234	5.3	4.6-6.0	218	2.7	2.3-3.0	120	3.4	2.8-4.1	98	2.1	1.7-2.6
2009	500	6.1	5.6-6.6	267	7.3	6.4-8.2	233	5.1	4.5-5.8	253	3.1	2.7-3.5	138	4.0	3.3-4.7	115	2.5	2.0-3.0
2010	464	5.7	5.1-6.2	245	6.7	5.9-7.6	219	4.9	4.2-5.6	239	2.8	2.5-3.2	119	3.3	2.7-3.9	120	2.5	2.1-3.0
2011	482	5.7	5.2-6.3	254	6.8	6.0-7.7	228	4.9	4.3-5.6	267	3.2	2.8-3.6	139	3.8	3.2-4.5	128	2.7	2.3-3.2
2012	486	5.7	5.2-6.2	261	6.8	6.0-7.7	225	4.8	4.2-5.4	247	2.8	2.5-3.2	111	2.9	2.4-3.5	136	2.8	2.3-3.3
2013	512	6.0	5.5-6.5	271	6.8	6.0-7.7	241	5.2	4.6-5.9	258	2.9	2.6-3.3	132	3.5	2.9-4.2	126	2.4	2.0-2.9
2014	516	5.9	5.4-6.4	294	7.4	6.5-8.3	222	4.7	4.1-5.4	286	3.2	2.8-3.6	154	4.0	3.3-4.6	132	2.6	2.2-3.1
2015	527	6.0	5.5-6.5	269	6.7	5.9-7.6	258	5.4	4.7-6.1	257	2.8	2.5-3.2	126	3.1	2.6-3.7	131	2.6	2.1-3.1
2016	541	6.0	5.5-6.5	307	7.5	6.7-8.4	234	4.7	4.1-5.4	287	3.1	2.7-3.5	156	3.8	3.2-4.5	131	2.5	2.1-3.0
2017	545	6.1	5.6-6.7	299	7.4	6.6-8.3	246	5.0	4.4-5.7	293	3.2	2.8-3.5	163	4.0	3.4-4.7	130	2.5	2.1-3.0
2018	513	5.7	5.2-6.2	291	7.0	6.2-7.8	222	4.6	4.0-5.3	283	3.1	2.7-3.5	160	3.9	3.3-4.6	123	2.4	2.0-2.8
2019	536	5.9	5.4-6.4	294	7.0	6.2-7.9	242	4.9	4.3-5.6	299	3.2	2.8-3.6	162	3.9	3.3-4.5	137	2.6	2.2-3.1
2020	489	5.5	5.0-6.0	263	6.3	5.6-7.1	226	4.7	4.1-5.4	290	3.0	2.7-3.4	161	3.7	3.2-4.4	129	2.4	2.0-2.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Thyroid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	232	3.2	2.8-3.7	63	2.0	1.5-2.6	169	4.1	3.5-4.9	59	0.9	0.6-1.1	23	0.8	0.5-1.1	36	0.9	0.6-1.3
1977	265	3.6	3.2-4.1	64	2.0	1.5-2.5	201	4.9	4.3-5.7	51	0.7	0.5-0.9	27	0.9	0.6-1.3	24	0.6	0.4-0.9
1978	264	3.7	3.3-4.2	72	2.3	1.8-3.0	192	4.8	4.1-5.6	45	0.7	0.5-0.9	11	0.4	0.2-0.8	34	0.8	0.6-1.2
1979	251	3.6	3.2-4.1	92	3.0	2.4-3.7	159	4.2	3.5-4.9	40	0.6	0.4-0.8	17	0.6	0.3-1.0	23	0.5	0.3-0.8
1980	248	3.5	3.1-4.0	62	2.1	1.6-2.7	186	4.7	4.0-5.4	49	0.7	0.5-0.9	16	0.6	0.3-0.9	33	0.8	0.5-1.1
1981	229	3.3	2.8-3.7	63	2.0	1.5-2.6	166	4.3	3.6-5.0	44	0.6	0.4-0.8	12	0.4	0.2-0.7	32	0.7	0.5-1.1
1982	238	3.4	3.0-3.9	73	2.5	1.9-3.1	165	4.2	3.6-4.9	56	0.8	0.6-1.0	18	0.6	0.4-1.0	38	0.9	0.6-1.2
1983	271	3.9	3.4-4.4	64	2.0	1.5-2.6	207	5.4	4.7-6.3	33	0.5	0.3-0.7	8	0.3	0.1-0.5	25	0.6	0.4-0.9
1984	320	4.5	4.0-5.0	84	2.8	2.2-3.4	236	6.0	5.2-6.8	49	0.7	0.5-0.9	17	0.6	0.4-1.0	32	0.7	0.5-1.0
1985	334	4.6	4.1-5.2	99	3.1	2.5-3.8	235	5.8	5.1-6.6	55	0.8	0.6-1.0	28	1.0	0.7-1.5	27	0.6	0.4-0.9
1986	286	3.9	3.4-4.4	71	2.3	1.8-2.9	215	5.3	4.6-6.0	38	0.5	0.4-0.7	8	0.3	0.1-0.6	30	0.7	0.5-1.0
1987	290	4.0	3.6-4.5	80	2.7	2.1-3.3	210	5.2	4.5-6.0	42	0.6	0.4-0.8	14	0.5	0.3-0.9	28	0.6	0.4-0.9
1988	285	3.9	3.5-4.4	90	2.9	2.3-3.6	195	4.8	4.1-5.6	35	0.5	0.3-0.7	12	0.4	0.2-0.8	23	0.5	0.3-0.8
1989	249	3.5	3.0-3.9	62	1.9	1.5-2.5	187	4.8	4.1-5.5	38	0.5	0.4-0.7	9	0.3	0.1-0.6	29	0.6	0.4-0.9
1990	301	4.1	3.7-4.6	72	2.3	1.8-2.9	229	5.7	5.0-6.5	41	0.6	0.4-0.8	11	0.4	0.2-0.7	30	0.7	0.5-1.0
1991	331	4.5	4.0-5.0	77	2.4	1.9-3.0	254	6.2	5.4-7.0	43	0.6	0.4-0.8	10	0.4	0.2-0.7	33	0.7	0.5-1.1
1992	378	5.1	4.6-5.7	92	2.9	2.3-3.6	286	7.0	6.2-7.9	42	0.6	0.4-0.8	15	0.5	0.3-0.9	27	0.6	0.4-0.9
1993	380	5.2	4.7-5.8	90	2.9	2.3-3.6	290	7.2	6.4-8.1	37	0.5	0.4-0.7	10	0.3	0.2-0.6	27	0.6	0.4-0.9
1994	426	5.8	5.2-6.4	113	3.5	2.8-4.2	313	7.7	6.9-8.6	41	0.6	0.4-0.8	16	0.6	0.3-0.9	25	0.6	0.4-0.9
1995	439	5.8	5.2-6.3	110	3.4	2.8-4.1	329	7.9	7.0-8.8	29	0.4	0.3-0.6	7	0.2	0.1-0.4	22	0.5	0.3-0.7
1996	447	5.9	5.3-6.4	103	3.2	2.6-3.9	344	8.2	7.4-9.2	47	0.6	0.5-0.9	19	0.6	0.4-1.0	28	0.6	0.4-0.9
1997	484	6.3	5.7-6.9	122	3.6	3.0-4.3	362	8.6	7.7-9.5	40	0.6	0.4-0.8	13	0.4	0.2-0.8	27	0.6	0.4-0.9
1998	543	7.0	6.4-7.6	132	3.9	3.3-4.6	411	9.6	8.7-10.6	32	0.4	0.3-0.6	11	0.4	0.2-0.6	21	0.5	0.3-0.7
1999	583	7.4	6.8-8.0	127	3.6	3.0-4.3	456	10.7	9.7-11.7	33	0.4	0.3-0.6	12	0.4	0.2-0.7	21	0.5	0.3-0.7
2000	620	7.8	7.2-8.4	142	3.9	3.3-4.7	478	11.1	10.1-12.1	47	0.6	0.5-0.8	12	0.4	0.2-0.7	35	0.8	0.5-1.1
2001	691	8.6	7.9-9.2	167	4.7	4.0-5.5	524	12.0	11.0-13.0	41	0.5	0.4-0.7	13	0.4	0.2-0.7	28	0.6	0.4-0.9
2002	714	8.8	8.2-9.5	186	5.1	4.4-6.0	528	12.1	11.1-13.2	40	0.5	0.4-0.7	13	0.4	0.2-0.7	27	0.6	0.4-0.8
2003	679	8.3	7.7-9.0	145	4.0	3.3-4.7	534	12.0	11.0-13.1	30	0.4	0.3-0.6	10	0.4	0.2-0.6	20	0.4	0.3-0.6
2004	803	9.8	9.1-10.5	194	5.4	4.7-6.2	609	13.7	12.6-14.9	35	0.4	0.3-0.6	17	0.6	0.3-0.9	18	0.4	0.2-0.6
2005	1,068	13.1	12.3-13.9	252	6.9	6.1-7.8	816	18.6	17.3-19.9	56	0.7	0.5-0.9	21	0.7	0.4-1.0	35	0.7	0.5-1.0
2006	1,041	12.7	11.9-13.5	233	6.2	5.4-7.1	808	18.3	17.1-19.6	51	0.6	0.5-0.9	16	0.5	0.3-0.9	35	0.7	0.5-1.0
2007	1,160	14.1	13.3-14.9	249	6.6	5.8-7.5	911	20.6	19.3-22.0	54	0.7	0.5-0.9	15	0.5	0.3-0.8	39	0.8	0.6-1.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York City

Thyroid

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	1,414	16.8	15.9-17.7	308	8.1	7.2-9.1	1,106	24.5	23.1-26.0	42	0.5	0.4-0.7	18	0.6	0.4-1.0	24	0.5	0.3-0.8
2009	1,582	18.7	17.8-19.7	371	9.5	8.6-10.6	1,211	26.8	25.3-28.4	46	0.6	0.4-0.8	22	0.7	0.4-1.1	24	0.5	0.3-0.7
2010	1,590	18.7	17.8-19.7	358	9.3	8.4-10.4	1,232	27.1	25.6-28.7	49	0.6	0.4-0.8	22	0.6	0.4-1.0	27	0.5	0.4-0.8
2011	1,622	18.8	17.9-19.8	340	8.7	7.8-9.7	1,282	27.8	26.3-29.4	40	0.5	0.3-0.7	12	0.4	0.2-0.7	28	0.5	0.4-0.8
2012	1,719	19.8	18.8-20.7	367	9.1	8.2-10.2	1,352	29.2	27.6-30.8	42	0.5	0.4-0.7	19	0.6	0.3-0.9	23	0.5	0.3-0.7
2013	1,952	22.3	21.3-23.3	449	11.2	10.1-12.3	1,503	32.3	30.7-34.1	54	0.6	0.5-0.8	24	0.7	0.4-1.0	30	0.6	0.4-0.8
2014	1,927	21.7	20.7-22.7	459	11.2	10.2-12.3	1,468	31.2	29.6-32.9	58	0.6	0.5-0.8	17	0.4	0.2-0.7	41	0.8	0.5-1.0
2015	1,887	21.3	20.3-22.3	439	10.5	9.5-11.6	1,448	31.0	29.4-32.7	49	0.5	0.4-0.7	19	0.5	0.3-0.7	30	0.5	0.4-0.8
2016	1,871	21.0	20.0-22.0	427	10.2	9.2-11.2	1,444	30.6	29.0-32.3	47	0.5	0.4-0.7	17	0.4	0.3-0.7	30	0.5	0.3-0.7
2017	1,770	19.6	18.7-20.5	441	10.4	9.4-11.5	1,329	27.9	26.4-29.5	51	0.6	0.4-0.7	17	0.5	0.3-0.7	34	0.6	0.4-0.9
2018	1,726	19.5	18.6-20.5	410	9.9	8.9-10.9	1,316	28.2	26.7-29.9	46	0.4	0.3-0.6	14	0.3	0.2-0.6	32	0.5	0.4-0.8
2019	1,852	20.8	19.9-21.8	476	11.3	10.3-12.4	1,376	29.6	28.0-31.3	52	0.5	0.4-0.7	25	0.6	0.4-0.9	27	0.5	0.3-0.7
2020	1,431	16.3	15.4-17.2	381	9.3	8.3-10.3	1,050	22.7	21.3-24.2	43	0.4	0.3-0.6	18	0.4	0.3-0.7	25	0.4	0.3-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Hodgkin lymphoma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	255	3.3	2.9-3.7	136	4.0	3.3-4.8	119	2.7	2.3-3.3	97	1.3	1.1-1.6	64	1.9	1.5-2.5	33	0.8	0.6-1.2
1977	243	3.2	2.8-3.6	139	4.0	3.4-4.8	104	2.5	2.1-3.1	106	1.5	1.2-1.8	56	1.8	1.3-2.3	50	1.3	0.9-1.7
1978	243	3.2	2.8-3.6	127	3.7	3.0-4.4	116	2.8	2.3-3.3	92	1.2	1.0-1.5	50	1.5	1.1-2.0	42	1.0	0.7-1.4
1979	238	3.3	2.9-3.8	125	3.9	3.2-4.7	113	2.8	2.3-3.4	79	1.1	0.9-1.4	48	1.5	1.1-2.0	31	0.8	0.5-1.2
1980	250	3.3	2.9-3.8	129	3.9	3.3-4.7	121	2.8	2.3-3.4	82	1.1	0.9-1.4	40	1.4	1.0-1.9	42	1.0	0.7-1.3
1981	231	3.0	2.6-3.5	131	3.8	3.2-4.6	100	2.3	1.9-2.9	81	1.2	0.9-1.4	42	1.4	1.0-1.9	39	0.9	0.7-1.3
1982	254	3.4	3.0-3.8	148	4.4	3.7-5.3	106	2.5	2.1-3.1	67	0.8	0.7-1.1	34	1.0	0.7-1.4	33	0.8	0.5-1.1
1983	253	3.5	3.0-3.9	149	4.6	3.8-5.4	104	2.5	2.0-3.0	75	1.0	0.8-1.2	43	1.3	0.9-1.8	32	0.7	0.5-1.0
1984	237	3.1	2.7-3.6	135	3.9	3.3-4.7	102	2.4	2.0-3.0	85	1.2	0.9-1.5	52	1.6	1.2-2.1	33	0.8	0.5-1.1
1985	253	3.3	2.9-3.7	155	4.5	3.8-5.3	98	2.3	1.8-2.8	80	1.0	0.8-1.3	46	1.3	0.9-1.7	34	0.8	0.6-1.1
1986	261	3.4	3.0-3.9	167	4.8	4.1-5.7	94	2.2	1.8-2.7	74	1.0	0.7-1.2	49	1.4	1.0-1.9	25	0.6	0.4-0.9
1987	223	2.9	2.5-3.3	124	3.6	3.0-4.4	99	2.4	1.9-2.9	57	0.7	0.6-1.0	38	1.1	0.8-1.6	19	0.4	0.3-0.7
1988	261	3.4	3.0-3.8	149	4.2	3.5-5.0	112	2.7	2.2-3.2	50	0.7	0.5-0.9	27	0.9	0.6-1.3	23	0.5	0.3-0.8
1989	244	3.2	2.8-3.6	145	4.1	3.4-4.9	99	2.4	2.0-3.0	74	1.0	0.8-1.3	49	1.5	1.1-2.0	25	0.6	0.4-0.9
1990	219	2.9	2.5-3.3	128	3.8	3.1-4.5	91	2.2	1.7-2.7	75	1.0	0.8-1.3	51	1.6	1.2-2.1	24	0.6	0.4-0.8
1991	280	3.7	3.3-4.2	150	4.4	3.7-5.2	130	3.2	2.7-3.8	63	0.9	0.7-1.1	36	1.1	0.8-1.6	27	0.6	0.4-0.9
1992	248	3.2	2.8-3.6	143	4.0	3.3-4.7	105	2.5	2.0-3.0	63	0.8	0.6-1.0	35	1.0	0.7-1.4	28	0.7	0.4-1.0
1993	221	2.8	2.5-3.3	118	3.4	2.8-4.1	103	2.3	1.9-2.9	56	0.7	0.6-1.0	33	0.9	0.6-1.3	23	0.5	0.3-0.8
1994	238	3.1	2.7-3.5	142	4.1	3.4-4.8	96	2.3	1.8-2.8	58	0.8	0.6-1.0	28	0.9	0.6-1.3	30	0.7	0.5-1.0
1995	251	3.2	2.8-3.7	132	3.7	3.1-4.4	119	2.8	2.4-3.4	43	0.6	0.4-0.8	24	0.7	0.5-1.1	19	0.5	0.3-0.7
1996	204	2.6	2.3-3.0	109	3.0	2.5-3.7	95	2.2	1.8-2.7	45	0.6	0.4-0.8	24	0.7	0.5-1.1	21	0.5	0.3-0.7
1997	236	3.0	2.6-3.4	136	3.8	3.2-4.5	100	2.4	1.9-2.9	50	0.7	0.5-0.9	22	0.7	0.4-1.0	28	0.7	0.4-1.0
1998	259	3.2	2.8-3.6	136	3.6	3.0-4.3	123	2.9	2.4-3.4	45	0.6	0.4-0.8	22	0.6	0.4-0.9	23	0.5	0.3-0.8
1999	225	2.8	2.4-3.1	113	2.9	2.4-3.6	112	2.6	2.1-3.1	32	0.4	0.3-0.6	18	0.5	0.3-0.8	14	0.3	0.2-0.5
2000	281	3.4	3.1-3.9	155	4.1	3.5-4.8	126	2.9	2.4-3.5	42	0.5	0.4-0.7	23	0.7	0.4-1.1	19	0.4	0.3-0.7
2001	247	3.0	2.6-3.4	135	3.5	2.9-4.1	112	2.5	2.0-3.0	34	0.4	0.3-0.6	18	0.5	0.3-0.8	16	0.3	0.2-0.6
2002	283	3.4	3.0-3.9	147	3.9	3.3-4.6	136	3.1	2.6-3.6	36	0.5	0.3-0.6	26	0.8	0.5-1.2	10	0.2	0.1-0.4
2003	266	3.3	2.9-3.7	131	3.5	2.9-4.1	135	3.2	2.7-3.8	40	0.5	0.4-0.7	19	0.6	0.3-0.9	21	0.5	0.3-0.7
2004	258	3.1	2.8-3.5	138	3.6	3.0-4.2	120	2.7	2.2-3.3	33	0.4	0.3-0.6	19	0.5	0.3-0.8	14	0.3	0.2-0.5
2005	269	3.3	2.9-3.7	157	4.2	3.6-4.9	112	2.5	2.1-3.0	30	0.4	0.3-0.5	19	0.5	0.3-0.8	11	0.2	0.1-0.4
2006	271	3.3	2.9-3.7	154	4.0	3.4-4.7	117	2.6	2.2-3.2	23	0.3	0.2-0.4	15	0.4	0.2-0.7	8	0.2	0.1-0.4
2007	260	3.1	2.7-3.5	124	3.2	2.6-3.8	136	3.0	2.5-3.6	41	0.5	0.4-0.7	20	0.6	0.4-0.9	21	0.5	0.3-0.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Hodgkin lymphoma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	288	3.5	3.1-3.9	149	3.8	3.2-4.5	139	3.1	2.6-3.7	39	0.5	0.3-0.6	25	0.7	0.4-1.0	14	0.3	0.2-0.5
2009	277	3.3	2.9-3.7	146	3.8	3.2-4.5	131	2.9	2.4-3.5	31	0.4	0.3-0.5	19	0.5	0.3-0.8	12	0.3	0.1-0.5
2010	278	3.2	2.8-3.6	152	3.8	3.2-4.4	126	2.7	2.3-3.3	30	0.3	0.2-0.5	19	0.5	0.3-0.7	11	0.3	0.1-0.5
2011	285	3.3	2.9-3.7	157	3.8	3.2-4.4	128	2.8	2.3-3.3	42	0.5	0.4-0.7	23	0.7	0.4-1.0	19	0.4	0.2-0.6
2012	246	2.8	2.5-3.2	121	3.0	2.5-3.6	125	2.8	2.3-3.3	24	0.3	0.2-0.4	14	0.4	0.2-0.6	10	0.2	0.1-0.4
2013	292	3.3	3.0-3.7	142	3.5	2.9-4.1	150	3.2	2.7-3.8	32	0.4	0.2-0.5	19	0.5	0.3-0.8	13	0.3	0.1-0.5
2014	290	3.3	2.9-3.7	154	3.8	3.2-4.4	136	2.9	2.4-3.5	32	0.4	0.2-0.5	13	0.3	0.2-0.6	19	0.4	0.2-0.6
2015	284	3.3	2.9-3.7	160	3.9	3.3-4.6	124	2.7	2.2-3.2	31	0.3	0.2-0.5	16	0.4	0.2-0.7	15	0.3	0.2-0.5
2016	281	3.2	2.8-3.6	141	3.5	2.9-4.1	140	3.0	2.5-3.5	27	0.3	0.2-0.4	17	0.4	0.2-0.7	10	0.2	0.1-0.3
2017	262	2.9	2.6-3.3	140	3.3	2.7-3.9	122	2.6	2.2-3.2	19	0.2	0.1-0.3	8	0.2	0.1-0.4	11	0.2	0.1-0.4
2018	217	2.4	2.1-2.8	124	3.0	2.5-3.6	93	1.9	1.6-2.4	22	0.2	0.1-0.4	16	0.4	0.2-0.6	6	0.1	0.0-0.3
2019	267	3.1	2.8-3.5	147	3.6	3.1-4.3	120	2.6	2.2-3.2	19	0.2	0.1-0.3	10	0.2	0.1-0.4	9	0.2	0.1-0.3
2020	228	2.6	2.3-3.0	128	3.1	2.6-3.7	100	2.3	1.8-2.8	26	0.3	0.2-0.4	14	0.3	0.2-0.5	12	0.2	0.1-0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Non-Hodgkin lymphomas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	818	11.0	10.3-11.8	407	13.4	12.1-14.8	411	9.5	8.6-10.5	453	6.2	5.6-6.8	249	8.5	7.5-9.7	204	4.7	4.1-5.4
1977	783	10.6	9.9-11.4	386	12.4	11.1-13.7	397	9.3	8.4-10.3	401	5.5	5.0-6.1	189	6.3	5.4-7.3	212	4.9	4.3-5.7
1978	813	11.2	10.4-12.0	415	13.4	12.1-14.8	398	9.5	8.6-10.5	427	5.8	5.3-6.4	233	7.7	6.7-8.9	194	4.5	3.9-5.2
1979	842	11.8	11.0-12.6	400	13.8	12.4-15.3	442	10.3	9.4-11.3	417	6.0	5.4-6.6	207	7.3	6.3-8.5	210	5.0	4.4-5.8
1980	941	13.1	12.2-14.0	461	15.3	13.9-16.8	480	11.3	10.3-12.4	436	6.0	5.4-6.6	228	7.5	6.5-8.6	208	4.9	4.2-5.6
1981	913	12.6	11.8-13.5	435	14.5	13.2-16.0	478	11.2	10.2-12.3	442	6.2	5.6-6.8	201	7.0	6.0-8.1	241	5.6	4.9-6.4
1982	1,020	14.2	13.3-15.1	470	15.6	14.2-17.1	550	12.9	11.8-14.0	463	6.5	5.9-7.2	217	7.6	6.6-8.8	246	5.7	5.0-6.5
1983	1,019	14.3	13.5-15.3	521	17.9	16.3-19.6	498	11.7	10.7-12.8	515	7.2	6.6-7.9	255	8.8	7.7-10.0	260	6.1	5.3-6.9
1984	1,051	14.5	13.7-15.5	530	17.9	16.4-19.6	521	12.1	11.0-13.2	451	6.3	5.7-6.9	233	8.0	7.0-9.2	218	5.0	4.3-5.7
1985	1,184	16.4	15.5-17.4	633	21.4	19.7-23.1	551	12.7	11.7-13.9	552	7.6	7.0-8.3	290	9.9	8.8-11.2	262	6.0	5.3-6.8
1986	1,253	17.2	16.2-18.2	661	21.5	19.9-23.3	592	13.7	12.6-14.8	501	7.0	6.4-7.6	274	9.4	8.3-10.6	227	5.2	4.5-5.9
1987	1,265	17.6	16.6-18.6	696	23.3	21.5-25.1	569	13.1	12.0-14.2	507	7.0	6.4-7.7	263	8.9	7.8-10.1	244	5.6	4.9-6.3
1988	1,172	16.2	15.3-17.2	645	21.1	19.5-22.9	527	12.4	11.3-13.5	567	7.9	7.3-8.6	295	10.2	9.0-11.5	272	6.2	5.4-6.9
1989	1,231	17.3	16.4-18.4	716	23.6	21.8-25.4	515	12.2	11.2-13.3	532	7.5	6.9-8.2	304	10.6	9.4-11.8	228	5.3	4.6-6.1
1990	1,395	19.4	18.3-20.4	811	26.4	24.5-28.3	584	13.7	12.6-14.9	544	7.6	7.0-8.3	273	9.4	8.3-10.6	271	6.3	5.5-7.1
1991	1,543	21.4	20.3-22.5	879	28.9	26.9-30.9	664	15.6	14.4-16.8	539	7.6	6.9-8.2	295	10.0	8.8-11.2	244	5.6	4.9-6.3
1992	1,497	20.7	19.7-21.8	832	26.8	25.0-28.8	665	15.8	14.6-17.1	562	7.9	7.2-8.5	288	10.1	8.9-11.4	274	6.3	5.6-7.2
1993	1,472	20.3	19.2-21.4	867	27.7	25.8-29.6	605	14.3	13.2-15.5	536	7.4	6.8-8.1	285	9.8	8.7-11.0	251	5.8	5.1-6.6
1994	1,724	23.7	22.6-24.9	1,000	31.5	29.5-33.6	724	17.2	16.0-18.5	571	8.0	7.3-8.7	288	9.9	8.7-11.1	283	6.5	5.8-7.4
1995	1,591	21.8	20.7-22.9	914	29.0	27.1-31.0	677	15.9	14.7-17.1	533	7.4	6.8-8.1	279	9.6	8.5-10.8	254	5.9	5.2-6.6
1996	1,589	21.6	20.6-22.7	881	28.2	26.3-30.1	708	16.6	15.4-17.9	554	7.7	7.1-8.4	285	10.1	8.9-11.3	269	6.2	5.5-7.0
1997	1,554	21.1	20.1-22.2	829	26.5	24.7-28.5	725	16.9	15.6-18.1	558	7.7	7.0-8.3	286	9.8	8.7-11.0	272	6.2	5.5-7.0
1998	1,470	19.7	18.7-20.7	799	25.0	23.2-26.8	671	15.6	14.5-16.9	562	7.7	7.1-8.3	268	8.9	7.9-10.1	294	6.7	5.9-7.5
1999	1,505	19.8	18.9-20.9	774	24.1	22.4-25.8	731	16.7	15.5-18.0	537	7.2	6.6-7.8	274	9.2	8.1-10.4	263	5.9	5.2-6.7
2000	1,465	19.1	18.1-20.1	751	22.8	21.2-24.5	714	16.1	14.9-17.3	512	6.7	6.2-7.3	258	8.5	7.5-9.6	254	5.6	4.9-6.3
2001	1,473	19.0	18.0-20.0	795	24.3	22.6-26.1	678	15.1	14.0-16.3	517	6.7	6.1-7.3	286	9.3	8.2-10.4	231	4.9	4.3-5.6
2002	1,460	18.6	17.7-19.6	804	23.9	22.3-25.7	656	14.5	13.4-15.7	469	6.0	5.5-6.6	256	8.1	7.1-9.2	213	4.6	4.0-5.2
2003	1,599	20.2	19.2-21.2	806	23.9	22.3-25.7	793	17.4	16.2-18.7	481	6.1	5.6-6.7	245	7.6	6.6-8.6	236	5.0	4.4-5.7
2004	1,537	19.4	18.5-20.4	798	23.7	22.1-25.5	739	16.2	15.1-17.4	464	5.9	5.4-6.5	239	7.6	6.6-8.6	225	4.8	4.2-5.4
2005	1,647	20.7	19.7-21.7	822	24.1	22.4-25.8	825	18.1	16.9-19.4	465	5.9	5.3-6.4	237	7.4	6.5-8.4	228	4.8	4.2-5.5
2006	1,595	19.9	19.0-20.9	869	25.3	23.6-27.1	726	15.8	14.7-17.0	467	5.9	5.4-6.5	238	7.4	6.5-8.4	229	4.9	4.3-5.6
2007	1,593	19.8	18.8-20.8	839	24.3	22.6-26.0	754	16.4	15.2-17.6	468	5.9	5.3-6.4	240	7.4	6.4-8.4	228	4.9	4.2-5.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Non-Hodgkin lymphomas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	1,674	20.5	19.5-21.5	900	25.7	24.0-27.4	774	16.6	15.4-17.8	420	5.2	4.7-5.7	228	6.9	6.0-7.9	192	4.0	3.4-4.6
2009	1,771	21.6	20.6-22.6	887	25.3	23.6-27.1	884	18.7	17.5-20.0	432	5.2	4.7-5.7	230	6.7	5.9-7.7	202	4.1	3.6-4.8
2010	1,672	20.0	19.1-21.0	882	24.6	22.9-26.3	790	16.5	15.4-17.7	409	5.0	4.5-5.5	211	6.3	5.5-7.3	198	4.0	3.5-4.6
2011	1,774	20.8	19.8-21.8	915	24.8	23.2-26.5	859	17.6	16.4-18.9	429	5.1	4.6-5.6	225	6.5	5.7-7.4	204	4.1	3.5-4.7
2012	1,744	20.1	19.2-21.1	938	25.1	23.5-26.8	806	16.5	15.4-17.7	433	5.0	4.5-5.5	235	6.7	5.8-7.6	198	3.8	3.3-4.4
2013	1,820	20.6	19.7-21.6	971	25.5	23.9-27.2	849	16.8	15.7-18.0	436	5.0	4.5-5.5	234	6.5	5.7-7.4	202	3.9	3.3-4.5
2014	1,813	20.1	19.2-21.1	965	25.0	23.4-26.6	848	16.5	15.4-17.7	426	4.7	4.3-5.2	227	6.1	5.3-7.0	199	3.7	3.2-4.3
2015	1,801	19.7	18.8-20.7	965	24.3	22.7-25.9	836	16.2	15.1-17.4	389	4.3	3.9-4.8	220	6.0	5.2-6.8	169	3.2	2.8-3.8
2016	1,863	20.0	19.1-20.9	985	24.2	22.7-25.8	878	16.8	15.6-17.9	442	4.7	4.3-5.2	237	6.1	5.3-6.9	205	3.7	3.2-4.3
2017	1,907	20.4	19.5-21.4	990	24.3	22.7-25.9	917	17.5	16.4-18.8	429	4.5	4.1-4.9	226	5.8	5.0-6.6	203	3.6	3.1-4.1
2018	1,882	20.0	19.1-21.0	990	24.2	22.6-25.7	892	16.8	15.7-18.0	384	3.9	3.5-4.3	218	5.4	4.7-6.2	166	2.8	2.4-3.3
2019	1,940	20.1	19.2-21.1	1,039	24.6	23.1-26.2	901	16.7	15.6-17.9	415	4.2	3.8-4.6	243	5.9	5.2-6.7	172	2.9	2.5-3.4
2020	1,629	16.9	16.1-17.8	847	20.0	18.7-21.4	782	14.5	13.4-15.5	359	3.6	3.2-4.0	209	5.0	4.4-5.8	150	2.5	2.1-3.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Myeloma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	341	4.6	4.1-5.2	172	5.8	4.9-6.8	169	3.9	3.3-4.6	207	2.9	2.5-3.3	97	3.5	2.8-4.3	110	2.6	2.1-3.1
1977	355	4.9	4.4-5.4	167	5.8	4.9-6.8	188	4.3	3.7-5.0	231	3.1	2.7-3.6	112	3.8	3.1-4.6	119	2.7	2.2-3.3
1978	349	4.8	4.3-5.3	179	6.2	5.3-7.2	170	3.9	3.3-4.5	196	2.6	2.3-3.1	93	3.1	2.5-3.8	103	2.3	1.9-2.8
1979	377	5.2	4.7-5.8	191	6.6	5.6-7.6	186	4.3	3.7-5.0	230	3.2	2.8-3.6	95	3.2	2.6-4.0	135	3.1	2.6-3.7
1980	365	5.1	4.6-5.7	164	5.9	5.0-6.9	201	4.7	4.0-5.4	246	3.4	3.0-3.8	114	4.0	3.3-4.8	132	3.0	2.5-3.6
1981	419	5.7	5.2-6.3	200	6.9	6.0-8.0	219	5.0	4.4-5.7	226	3.1	2.7-3.6	108	4.0	3.2-4.8	118	2.7	2.2-3.2
1982	390	5.4	4.9-6.0	175	6.2	5.3-7.2	215	4.9	4.3-5.6	249	3.5	3.0-3.9	114	4.0	3.3-4.8	135	3.2	2.6-3.7
1983	402	5.6	5.1-6.2	192	6.9	5.9-7.9	210	4.9	4.2-5.6	238	3.3	2.9-3.7	97	3.4	2.7-4.2	141	3.2	2.7-3.8
1984	375	5.3	4.7-5.8	169	6.2	5.3-7.3	206	4.7	4.1-5.4	249	3.4	3.0-3.9	105	3.7	3.0-4.5	144	3.2	2.7-3.8
1985	399	5.5	5.0-6.1	194	6.9	5.9-8.0	205	4.7	4.1-5.4	233	3.2	2.8-3.6	123	4.4	3.6-5.3	110	2.4	2.0-3.0
1986	396	5.5	4.9-6.1	194	6.8	5.8-7.8	202	4.6	4.0-5.3	237	3.3	2.9-3.7	110	3.9	3.2-4.7	127	2.9	2.4-3.4
1987	374	5.1	4.6-5.7	197	7.0	6.0-8.1	177	4.0	3.4-4.6	243	3.4	3.0-3.9	133	4.8	4.0-5.7	110	2.5	2.0-3.0
1988	368	5.1	4.6-5.7	185	6.7	5.8-7.8	183	4.2	3.6-4.8	215	3.0	2.6-3.4	111	4.0	3.3-4.9	104	2.3	1.9-2.8
1989	406	5.7	5.2-6.3	182	6.5	5.5-7.5	224	5.3	4.6-6.0	227	3.2	2.8-3.6	105	3.8	3.1-4.6	122	2.8	2.3-3.3
1990	419	5.9	5.3-6.5	193	6.8	5.8-7.8	226	5.4	4.7-6.2	222	3.1	2.7-3.5	99	3.7	3.0-4.5	123	2.8	2.3-3.4
1991	404	5.7	5.1-6.2	169	5.9	5.0-6.9	235	5.5	4.8-6.3	233	3.3	2.9-3.7	105	3.9	3.1-4.7	128	3.0	2.5-3.6
1992	468	6.5	5.9-7.1	234	8.4	7.3-9.5	234	5.4	4.7-6.2	254	3.6	3.1-4.0	116	4.3	3.5-5.2	138	3.2	2.7-3.8
1993	407	5.7	5.2-6.3	179	6.5	5.5-7.5	228	5.3	4.6-6.0	277	3.9	3.4-4.4	135	4.9	4.1-5.8	142	3.3	2.7-3.8
1994	411	5.7	5.2-6.3	188	6.5	5.6-7.5	223	5.3	4.7-6.1	240	3.4	2.9-3.8	104	3.6	3.0-4.4	136	3.2	2.6-3.7
1995	428	6.0	5.4-6.6	194	6.8	5.8-7.8	234	5.5	4.8-6.2	270	3.8	3.3-4.2	126	4.5	3.7-5.3	144	3.3	2.8-3.9
1996	472	6.6	6.0-7.2	218	7.6	6.6-8.7	254	6.0	5.2-6.7	242	3.4	3.0-3.8	103	3.7	3.0-4.5	139	3.2	2.7-3.8
1997	432	6.0	5.4-6.6	205	7.1	6.1-8.1	227	5.3	4.6-6.0	231	3.2	2.8-3.6	110	3.9	3.2-4.7	121	2.8	2.3-3.3
1998	501	6.9	6.3-7.5	241	8.2	7.2-9.3	260	6.0	5.3-6.8	226	3.1	2.7-3.5	101	3.4	2.8-4.2	125	2.9	2.4-3.4
1999	488	6.5	6.0-7.2	195	6.4	5.6-7.4	293	6.6	5.9-7.4	231	3.1	2.7-3.5	96	3.3	2.6-4.0	135	3.0	2.5-3.6
2000	489	6.4	5.9-7.0	215	6.9	6.0-7.9	274	6.1	5.4-6.9	250	3.3	2.9-3.8	93	3.1	2.5-3.9	157	3.4	2.9-4.0
2001	541	7.1	6.5-7.7	257	8.3	7.3-9.4	284	6.3	5.6-7.1	244	3.2	2.8-3.6	118	3.9	3.3-4.7	126	2.8	2.3-3.3
2002	550	7.1	6.5-7.7	260	8.3	7.3-9.4	290	6.4	5.7-7.2	259	3.4	3.0-3.8	116	3.9	3.2-4.6	143	3.1	2.6-3.6
2003	572	7.3	6.7-7.9	272	8.7	7.7-9.8	300	6.5	5.8-7.3	233	3.0	2.6-3.4	102	3.3	2.7-4.0	131	2.8	2.3-3.3
2004	551	7.0	6.4-7.6	253	7.8	6.8-8.8	298	6.4	5.7-7.2	228	2.9	2.6-3.3	101	3.3	2.7-4.0	127	2.7	2.3-3.2
2005	616	7.7	7.1-8.4	292	8.9	7.9-10.0	324	6.9	6.2-7.7	232	2.9	2.6-3.3	106	3.3	2.7-4.0	126	2.6	2.2-3.1
2006	614	7.8	7.2-8.4	295	9.1	8.0-10.2	319	6.9	6.2-7.7	214	2.7	2.4-3.1	120	3.8	3.1-4.5	94	2.0	1.6-2.5
2007	581	7.3	6.7-7.9	296	9.0	8.0-10.1	285	6.1	5.4-6.8	248	3.1	2.8-3.6	122	3.8	3.1-4.5	126	2.7	2.2-3.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Myeloma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	670	8.2	7.6-8.9	351	10.4	9.4-11.6	319	6.7	6.0-7.5	225	2.8	2.4-3.2	106	3.3	2.7-4.0	119	2.5	2.1-3.0
2009	736	9.0	8.3-9.6	388	11.2	10.1-12.4	348	7.3	6.5-8.1	223	2.7	2.4-3.1	125	3.8	3.1-4.5	98	2.0	1.6-2.4
2010	739	8.8	8.2-9.5	389	11.1	10.0-12.3	350	7.3	6.6-8.1	260	3.2	2.8-3.6	135	4.1	3.4-4.9	125	2.5	2.1-3.0
2011	813	9.5	8.9-10.2	440	12.2	11.0-13.4	373	7.8	7.0-8.6	234	2.7	2.4-3.1	108	3.1	2.6-3.8	126	2.5	2.1-3.0
2012	785	9.0	8.4-9.6	407	11.1	10.0-12.2	378	7.5	6.8-8.3	238	2.8	2.4-3.1	117	3.3	2.7-4.0	121	2.4	1.9-2.8
2013	802	8.9	8.3-9.5	409	10.8	9.7-11.9	393	7.6	6.8-8.4	264	2.9	2.6-3.3	150	4.0	3.4-4.7	114	2.2	1.8-2.6
2014	871	9.7	9.0-10.4	469	12.2	11.1-13.3	402	7.9	7.1-8.7	269	3.0	2.7-3.4	133	3.6	3.0-4.3	136	2.6	2.2-3.1
2015	838	9.0	8.4-9.7	425	10.8	9.8-11.9	413	7.8	7.1-8.6	284	3.1	2.7-3.5	146	3.9	3.2-4.5	138	2.5	2.1-3.0
2016	882	9.3	8.7-9.9	481	11.8	10.7-12.9	401	7.4	6.7-8.2	307	3.3	2.9-3.6	162	4.1	3.5-4.8	145	2.7	2.2-3.2
2017	919	9.6	9.0-10.2	458	11.2	10.1-12.3	461	8.4	7.6-9.2	277	2.9	2.5-3.2	140	3.5	2.9-4.2	137	2.4	2.0-2.8
2018	844	8.5	8.0-9.1	403	9.4	8.5-10.4	441	7.9	7.2-8.7	235	2.3	2.0-2.6	118	2.8	2.3-3.4	117	2.0	1.6-2.4
2019	866	8.8	8.2-9.4	468	10.9	10.0-12.0	398	7.2	6.5-8.0	267	2.6	2.3-2.9	135	3.2	2.7-3.8	132	2.2	1.8-2.6
2020	720	7.3	6.7-7.8	366	8.5	7.6-9.4	354	6.4	5.7-7.1	224	2.2	1.9-2.5	128	3.0	2.5-3.6	96	1.6	1.3-2.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Leukemias**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	808	11.3	10.5-12.1	446	14.9	13.5-16.4	362	8.8	7.8-9.7	589	8.5	7.8-9.2	312	10.8	9.6-12.2	277	6.8	6.0-7.7
1977	740	10.4	9.7-11.2	402	13.6	12.2-15.0	338	8.2	7.3-9.2	548	7.7	7.1-8.4	299	10.4	9.2-11.7	249	6.0	5.3-6.8
1978	815	11.6	10.8-12.4	463	16.0	14.5-17.6	352	8.6	7.7-9.6	555	7.9	7.2-8.6	302	10.3	9.2-11.6	253	6.2	5.4-7.0
1979	768	10.9	10.1-11.7	394	13.3	12.0-14.8	374	9.2	8.3-10.2	551	7.9	7.2-8.6	288	10.0	8.9-11.3	263	6.6	5.8-7.4
1980	875	12.3	11.5-13.2	443	15.2	13.8-16.8	432	10.5	9.5-11.5	522	7.4	6.8-8.1	291	10.3	9.1-11.6	231	5.6	4.8-6.4
1981	794	11.2	10.4-12.0	438	15.1	13.7-16.6	356	8.6	7.7-9.5	545	7.7	7.1-8.4	286	10.0	8.8-11.3	259	6.3	5.5-7.1
1982	833	11.8	11.0-12.6	453	15.7	14.2-17.3	380	9.1	8.2-10.1	557	7.9	7.3-8.6	308	10.9	9.7-12.2	249	6.0	5.3-6.8
1983	798	11.3	10.5-12.1	405	14.0	12.6-15.5	393	9.7	8.7-10.7	512	7.3	6.7-8.0	270	9.7	8.5-10.9	242	5.8	5.1-6.6
1984	846	12.0	11.1-12.8	458	16.3	14.8-17.9	388	9.2	8.3-10.2	556	7.8	7.1-8.4	289	10.2	9.0-11.5	267	6.2	5.5-7.0
1985	822	11.4	10.7-12.3	455	15.7	14.2-17.2	367	8.7	7.8-9.7	514	7.1	6.5-7.8	267	9.3	8.2-10.6	247	5.7	5.0-6.5
1986	859	11.9	11.1-12.8	439	14.8	13.4-16.3	420	10.1	9.1-11.1	505	7.0	6.4-7.7	256	8.7	7.6-9.9	249	5.8	5.1-6.6
1987	725	10.0	9.3-10.8	346	11.9	10.6-13.2	379	8.8	7.9-9.8	459	6.3	5.7-6.9	222	7.7	6.7-8.8	237	5.5	4.8-6.3
1988	742	10.3	9.6-11.1	388	13.6	12.3-15.1	354	8.3	7.5-9.2	461	6.4	5.8-7.0	246	8.7	7.6-9.8	215	5.0	4.3-5.7
1989	875	12.2	11.4-13.1	460	16.1	14.6-17.7	415	9.8	8.8-10.8	487	6.7	6.2-7.4	258	9.2	8.1-10.4	229	5.3	4.6-6.0
1990	896	12.5	11.7-13.4	464	15.8	14.3-17.3	432	10.3	9.4-11.4	503	7.1	6.5-7.7	253	9.2	8.1-10.4	250	5.8	5.1-6.6
1991	856	12.0	11.2-12.9	467	15.9	14.5-17.5	389	9.3	8.4-10.3	461	6.5	5.9-7.1	260	9.5	8.3-10.7	201	4.8	4.1-5.5
1992	874	12.2	11.4-13.1	464	16.0	14.5-17.6	410	9.8	8.8-10.8	459	6.5	5.9-7.1	235	8.5	7.4-9.7	224	5.2	4.6-6.0
1993	824	11.4	10.6-12.2	398	13.4	12.1-14.8	426	10.2	9.2-11.2	483	6.7	6.1-7.4	263	9.2	8.1-10.4	220	5.2	4.5-5.9
1994	908	12.6	11.8-13.5	464	15.9	14.4-17.4	444	10.4	9.4-11.4	529	7.3	6.7-8.0	263	9.1	8.0-10.3	266	6.1	5.4-6.9
1995	820	11.4	10.6-12.2	416	14.0	12.7-15.4	404	9.6	8.7-10.6	489	6.8	6.2-7.4	255	9.0	7.9-10.2	234	5.5	4.8-6.2
1996	847	11.6	10.8-12.4	467	15.5	14.1-17.0	380	8.9	8.0-9.8	459	6.3	5.8-6.9	224	7.7	6.7-8.8	235	5.4	4.8-6.2
1997	871	11.9	11.1-12.7	442	14.7	13.4-16.2	429	10.0	9.1-11.0	456	6.3	5.7-6.9	255	8.9	7.8-10.1	201	4.6	4.0-5.3
1998	854	11.5	10.8-12.3	455	14.8	13.4-16.2	399	9.2	8.4-10.2	458	6.2	5.7-6.8	249	8.3	7.3-9.5	209	4.8	4.1-5.5
1999	936	12.4	11.6-13.2	488	15.8	14.4-17.2	448	10.2	9.3-11.2	486	6.5	5.9-7.1	264	8.9	7.8-10.1	222	5.0	4.3-5.7
2000	940	12.3	11.6-13.2	476	15.2	13.8-16.6	464	10.4	9.5-11.4	458	6.0	5.5-6.6	208	6.8	5.9-7.8	250	5.5	4.8-6.2
2001	925	12.0	11.3-12.8	466	14.7	13.4-16.1	459	10.2	9.3-11.2	469	6.1	5.5-6.7	239	7.7	6.7-8.8	230	4.9	4.3-5.6
2002	890	11.4	10.7-12.2	492	15.1	13.8-16.6	398	8.8	7.9-9.7	469	6.0	5.5-6.6	242	7.9	6.9-9.0	227	4.9	4.3-5.6
2003	978	12.5	11.8-13.3	502	15.3	14.0-16.8	476	10.5	9.6-11.5	433	5.5	5.0-6.1	220	6.9	6.0-7.9	213	4.6	4.0-5.3
2004	937	11.9	11.2-12.7	502	15.5	14.1-16.9	435	9.5	8.6-10.5	466	5.9	5.4-6.5	246	8.0	7.0-9.1	220	4.7	4.1-5.3
2005	959	12.1	11.4-12.9	534	16.1	14.8-17.6	425	9.3	8.5-10.3	474	6.0	5.5-6.6	258	8.1	7.1-9.1	216	4.7	4.0-5.3
2006	967	12.3	11.6-13.1	540	16.4	15.0-17.8	427	9.4	8.5-10.4	452	5.7	5.2-6.3	243	7.6	6.6-8.6	209	4.4	3.8-5.0
2007	1,015	12.8	12.0-13.6	546	16.1	14.7-17.5	469	10.4	9.4-11.4	464	5.9	5.3-6.4	257	8.1	7.2-9.2	207	4.5	3.9-5.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York City
Leukemias**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	1,001	12.5	11.7-13.3	536	16.0	14.7-17.4	465	10.2	9.2-11.1	474	5.9	5.4-6.5	239	7.3	6.4-8.3	235	5.0	4.3-5.6
2009	1,085	13.3	12.5-14.1	582	16.9	15.5-18.3	503	10.8	9.8-11.8	434	5.4	4.9-5.9	206	6.2	5.3-7.1	228	4.8	4.2-5.4
2010	1,184	14.3	13.5-15.2	659	18.7	17.3-20.3	525	11.2	10.2-12.2	459	5.5	5.0-6.1	247	7.4	6.5-8.4	212	4.4	3.8-5.0
2011	1,226	14.5	13.7-15.4	670	18.6	17.2-20.1	556	11.6	10.6-12.6	492	5.8	5.3-6.3	274	8.0	7.0-9.0	218	4.3	3.7-4.9
2012	1,135	13.2	12.4-14.0	647	17.5	16.2-18.9	488	10.1	9.2-11.1	460	5.4	4.9-5.9	257	7.5	6.6-8.4	203	4.0	3.4-4.6
2013	1,238	14.2	13.4-15.0	685	18.4	17.0-19.8	553	11.3	10.3-12.3	497	5.7	5.2-6.2	291	8.2	7.2-9.2	206	4.1	3.5-4.7
2014	1,333	15.1	14.3-15.9	693	18.3	16.9-19.7	640	12.9	11.9-13.9	449	5.1	4.6-5.6	231	6.4	5.6-7.3	218	4.2	3.6-4.8
2015	1,304	14.4	13.6-15.2	726	18.7	17.4-20.2	578	11.3	10.3-12.2	495	5.4	5.0-6.0	268	7.2	6.3-8.1	227	4.3	3.7-4.9
2016	1,337	14.5	13.8-15.4	762	19.2	17.8-20.6	575	11.1	10.2-12.1	486	5.2	4.8-5.7	239	6.3	5.5-7.2	247	4.6	4.0-5.2
2017	1,230	13.2	12.5-14.0	676	16.7	15.4-18.0	554	10.7	9.8-11.6	481	5.1	4.6-5.5	250	6.4	5.6-7.3	231	4.2	3.6-4.8
2018	1,232	13.3	12.5-14.1	692	17.1	15.8-18.4	540	10.4	9.5-11.3	458	4.8	4.4-5.3	243	6.0	5.3-6.9	215	4.0	3.5-4.6
2019	1,158	12.4	11.7-13.1	649	15.8	14.6-17.1	509	9.7	8.8-10.6	462	4.8	4.3-5.3	259	6.5	5.7-7.3	203	3.6	3.1-4.1
2020	1,096	11.8	11.1-12.5	616	15.0	13.8-16.3	480	9.3	8.5-10.2	409	4.2	3.8-4.6	238	5.9	5.1-6.7	171	3.0	2.6-3.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State Excl New York City
All Invasive Malignant Tumors**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	37,817	405.5	401.4-409.7	18,574	480.1	472.8-487.5	19,243	366.5	361.3-371.8	20,094	218.3	215.2-221.4	10,509	279.1	273.5-284.9	9,585	181.3	177.7-185.0
1977	38,176	403.8	399.7-408.0	18,797	479.9	472.6-487.2	19,379	363.5	358.3-368.7	20,642	221.4	218.3-224.5	10,877	285.2	279.5-290.9	9,765	182.0	178.3-185.7
1978	38,807	406.8	402.7-410.9	19,095	484.5	477.2-491.8	19,712	365.7	360.5-370.9	20,868	220.4	217.4-223.5	10,909	284.6	279.0-290.3	9,959	182.1	178.5-185.8
1979	39,627	410.2	406.1-414.3	19,447	487.2	480.0-494.5	20,180	369.4	364.3-374.7	21,013	220.0	217.0-223.1	10,887	281.3	275.8-287.0	10,126	183.2	179.6-186.8
1980	41,104	419.4	415.3-423.5	20,597	507.1	499.8-514.5	20,507	370.9	365.7-376.1	21,571	222.4	219.3-225.4	11,281	286.6	281.1-292.3	10,290	183.4	179.8-187.0
1981	41,601	421.5	417.4-425.7	20,684	509.1	501.8-516.5	20,917	375.3	370.1-380.5	21,691	221.5	218.5-224.5	11,357	287.8	282.2-293.5	10,334	182.1	178.6-185.7
1982	42,460	424.9	420.8-429.0	21,024	510.4	503.1-517.7	21,436	380.2	375.0-385.4	21,934	221.7	218.7-224.7	11,407	286.9	281.4-292.6	10,527	183.3	179.8-186.9
1983	43,688	433.8	429.6-437.9	21,580	518.7	511.5-526.1	22,108	390.0	384.8-395.3	22,206	221.9	218.9-224.9	11,543	288.0	282.4-293.6	10,663	183.5	180.0-187.1
1984	44,105	434.4	430.3-438.6	21,573	513.0	505.9-520.3	22,532	393.6	388.4-398.9	22,439	222.4	219.4-225.3	11,461	283.0	277.5-288.5	10,978	187.4	183.8-191.0
1985	45,642	446.8	442.7-451.0	22,372	527.0	519.8-534.3	23,270	405.2	399.9-410.5	22,800	224.5	221.6-227.5	11,668	285.9	280.5-291.4	11,132	188.9	185.4-192.5
1986	45,693	444.3	440.1-448.4	22,339	524.1	517.0-531.4	23,354	403.8	398.5-409.1	23,260	227.3	224.3-230.2	11,997	291.8	286.3-297.3	11,263	189.6	186.1-193.2
1987	46,505	447.6	443.5-451.7	22,528	523.8	516.7-531.0	23,977	409.9	404.7-415.3	23,170	224.3	221.4-227.3	11,858	286.6	281.2-292.0	11,312	188.0	184.5-191.5
1988	46,843	446.6	442.5-450.7	22,692	522.8	515.7-530.0	24,151	409.9	404.7-415.2	23,509	225.0	222.1-227.9	12,076	288.3	282.9-293.7	11,433	188.3	184.8-191.8
1989	47,356	446.9	442.8-451.0	23,298	529.8	522.7-537.0	24,058	404.6	399.4-409.8	23,240	220.6	217.8-223.5	11,874	280.1	274.8-285.4	11,366	186.0	182.5-189.4
1990	49,819	464.5	460.4-468.6	24,622	549.3	542.1-556.4	25,197	419.4	414.2-424.7	23,242	217.6	214.8-220.5	11,812	275.3	270.2-280.6	11,430	183.9	180.5-187.3
1991	50,739	468.0	463.9-472.1	25,530	562.4	555.3-569.6	25,209	416.2	411.0-421.4	23,702	219.7	216.9-222.6	12,073	278.8	273.7-284.0	11,629	185.3	181.9-188.8
1992	52,914	482.2	478.0-486.3	27,408	595.2	588.0-602.6	25,506	416.6	411.4-421.8	23,942	219.3	216.5-222.1	11,953	273.8	268.7-279.0	11,989	188.9	185.5-192.4
1993	53,334	481.4	477.3-485.5	27,253	581.1	574.0-588.3	26,081	421.8	416.7-427.1	23,777	215.3	212.6-218.1	11,925	269.3	264.4-274.4	11,852	184.4	181.0-187.8
1994	53,629	481.2	477.2-485.4	27,126	575.0	568.0-582.1	26,503	426.1	421.0-431.3	24,543	220.3	217.5-223.1	12,515	279.7	274.7-284.8	12,028	185.3	182.0-188.7
1995	54,198	483.3	479.3-487.4	26,981	563.2	556.4-570.1	27,217	436.3	431.1-441.6	24,218	215.5	212.8-218.2	12,203	270.5	265.5-275.4	12,015	183.9	180.6-187.3
1996	56,147	497.9	493.8-502.0	28,374	587.2	580.2-594.2	27,773	442.8	437.5-448.1	24,045	212.4	209.7-215.1	12,121	264.6	259.8-269.4	11,924	181.4	178.1-184.7
1997	57,326	504.1	499.9-508.2	28,932	592.2	585.3-599.1	28,394	448.6	443.3-453.9	23,681	206.9	204.3-209.6	11,974	257.8	253.1-262.6	11,707	176.0	172.8-179.2
1998	58,906	512.7	508.6-516.9	29,535	594.8	587.9-601.7	29,371	461.2	455.9-466.6	23,300	201.2	198.6-203.8	11,686	245.4	240.9-249.9	11,614	172.9	169.8-176.2
1999	59,817	515.1	511.0-519.2	30,165	601.7	594.8-608.6	29,652	459.4	454.2-464.7	23,854	203.4	200.8-206.0	11,935	249.1	244.5-253.6	11,919	175.3	172.1-178.5
2000	61,378	522.1	517.9-526.2	31,138	614.8	608.0-621.8	30,240	462.4	457.1-467.7	23,922	201.1	198.5-203.7	11,937	246.7	242.2-251.2	11,985	173.6	170.5-176.8
2001	61,334	517.1	513.0-521.2	31,143	606.2	599.4-613.0	30,191	459.3	454.1-464.5	23,543	195.9	193.4-198.4	11,722	239.1	234.7-243.5	11,821	170.1	167.0-173.2
2002	62,503	520.7	516.6-524.8	32,311	619.4	612.6-626.3	30,192	454.8	449.7-460.1	23,618	194.1	191.6-196.6	11,752	235.6	231.3-240.0	11,866	169.2	166.1-172.3
2003	62,435	513.7	509.7-517.8	31,811	599.8	593.2-606.5	30,624	456.6	451.5-461.8	23,152	187.7	185.3-190.2	11,408	224.7	220.5-228.9	11,744	165.3	162.3-168.3
2004	63,003	513.3	509.3-517.3	32,022	595.5	588.9-602.1	30,981	459.0	453.9-464.2	23,186	186.0	183.6-188.4	11,510	222.4	218.3-226.6	11,676	162.8	159.8-165.8
2005	63,838	515.5	511.5-519.6	32,478	595.4	588.9-602.0	31,360	461.0	455.9-466.3	22,795	181.0	178.6-183.3	11,335	216.3	212.3-220.4	11,460	158.3	155.4-161.3
2006	65,752	526.3	522.2-530.4	33,906	612.9	606.4-619.6	31,846	466.1	460.9-471.4	22,758	179.0	176.6-181.3	11,289	211.9	207.9-215.8	11,469	157.6	154.7-160.6
2007	66,664	526.1	522.0-530.1	34,613	613.1	606.6-619.7	32,051	465.5	460.3-470.7	22,742	176.0	173.7-178.4	11,408	209.4	205.5-213.3	11,334	154.3	151.4-157.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

All Invasive Malignant Tumors

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	68,027	529.9	525.9-534.0	34,880	607.7	601.2-614.2	33,147	476.8	471.6-482.1	22,861	174.1	171.9-176.4	11,497	207.4	203.6-211.2	11,364	152.3	149.5-155.2
2009	68,027	522.0	518.0-525.9	34,923	595.3	589.0-601.7	33,104	470.3	465.1-475.5	22,540	169.6	167.4-171.9	11,475	203.6	199.9-207.4	11,065	147.1	144.3-149.9
2010	66,878	506.0	502.1-509.9	34,053	569.7	563.5-575.9	32,825	462.0	456.9-467.2	22,614	167.4	165.2-169.6	11,253	195.7	192.0-199.4	11,361	148.7	145.9-151.5
2011	69,148	516.5	512.6-520.4	35,414	581.6	575.5-587.9	33,734	470.5	465.3-475.7	22,554	164.4	162.2-166.6	11,269	191.8	188.2-195.5	11,285	145.8	143.1-148.6
2012	67,528	497.6	493.8-501.5	34,013	549.5	543.6-555.5	33,515	463.6	458.5-468.7	23,002	165.7	163.5-167.9	11,691	196.6	193.0-200.3	11,311	144.5	141.7-147.2
2013	69,334	504.9	501.0-508.7	34,639	550.0	544.1-556.0	34,695	476.4	471.2-481.6	22,611	159.8	157.7-162.0	11,477	187.9	184.4-191.5	11,134	140.2	137.5-142.9
2014	69,578	499.2	495.4-503.0	34,819	542.3	536.4-548.1	34,759	472.1	466.9-477.3	22,146	154.1	152.1-156.2	11,353	182.6	179.2-186.1	10,793	134.5	131.9-137.1
2015	70,653	501.4	497.6-505.2	35,231	539.6	533.8-545.4	35,422	478.2	473.0-483.4	22,264	152.7	150.7-154.8	11,242	178.1	174.7-181.5	11,022	135.1	132.5-137.7
2016	71,381	500.4	496.6-504.2	36,209	545.5	539.8-551.4	35,172	470.7	465.6-475.9	22,163	150.5	148.5-152.5	11,223	174.6	171.3-177.9	10,940	133.7	131.2-136.4
2017	72,406	501.3	497.5-505.1	36,747	545.1	539.4-550.9	35,659	473.0	467.8-478.1	22,029	146.8	144.8-148.8	11,239	171.5	168.3-174.8	10,790	129.3	126.8-131.9
2018	72,475	495.2	491.4-498.9	37,053	539.3	533.7-545.0	35,422	464.9	459.8-470.0	22,048	144.8	142.8-146.7	11,168	167.6	164.5-170.8	10,880	129.1	126.6-131.7
2019	74,614	503.8	500.1-507.6	38,134	545.7	540.1-551.4	36,480	475.7	470.6-480.9	21,578	139.6	137.7-141.5	10,918	160.8	157.7-163.9	10,660	124.4	122.0-126.9
2020	67,152	448.9	445.4-452.4	34,337	486.1	480.8-491.4	32,815	423.8	419.0-428.7	21,273	136.3	134.4-138.2	10,753	156.8	153.7-159.8	10,520	122.0	119.6-124.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Oral cavity and pharynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,076	11.1	10.5-11.8	731	17.4	16.1-18.8	345	6.4	5.8-7.2	417	4.3	3.9-4.8	300	7.3	6.5-8.2	117	2.1	1.7-2.5
1977	1,043	10.6	10.0-11.3	707	16.5	15.2-17.8	336	6.2	5.6-6.9	424	4.3	3.9-4.8	290	6.9	6.1-7.8	134	2.4	2.0-2.9
1978	1,092	11.0	10.3-11.7	715	16.7	15.5-18.1	377	6.8	6.2-7.6	497	5.1	4.6-5.6	348	8.5	7.6-9.5	149	2.7	2.3-3.2
1979	1,127	11.2	10.5-11.8	729	16.6	15.4-18.0	398	7.1	6.4-7.9	455	4.6	4.1-5.0	290	6.7	5.9-7.5	165	3.0	2.5-3.5
1980	1,099	10.9	10.2-11.6	719	16.5	15.2-17.8	380	6.7	6.0-7.4	439	4.4	3.9-4.8	288	6.8	6.0-7.6	151	2.6	2.2-3.1
1981	1,155	11.3	10.7-12.0	784	17.7	16.4-19.0	371	6.5	5.9-7.2	428	4.2	3.8-4.6	294	6.7	6.0-7.6	134	2.3	1.9-2.7
1982	1,196	11.8	11.1-12.5	798	18.3	16.9-19.6	398	7.1	6.4-7.8	453	4.5	4.0-4.9	304	7.2	6.4-8.1	149	2.5	2.1-3.0
1983	1,140	11.0	10.4-11.7	755	16.8	15.6-18.1	385	6.8	6.1-7.5	438	4.2	3.8-4.7	290	6.6	5.8-7.4	148	2.5	2.1-3.0
1984	1,149	11.1	10.5-11.8	778	17.6	16.3-18.9	371	6.4	5.8-7.1	420	4.1	3.7-4.5	274	6.3	5.6-7.2	146	2.5	2.1-2.9
1985	1,152	11.2	10.5-11.9	766	17.2	16.0-18.6	386	6.6	6.0-7.3	414	4.0	3.6-4.4	266	6.1	5.4-6.9	148	2.5	2.1-3.0
1986	1,116	10.8	10.2-11.5	740	16.7	15.4-18.0	376	6.4	5.8-7.1	447	4.3	3.9-4.8	293	6.6	5.8-7.5	154	2.6	2.2-3.1
1987	1,128	10.9	10.3-11.6	710	15.8	14.6-17.1	418	7.1	6.4-7.8	418	4.1	3.7-4.5	266	6.2	5.4-7.0	152	2.5	2.1-3.0
1988	1,124	10.7	10.1-11.4	712	15.6	14.5-16.9	412	7.0	6.3-7.7	439	4.2	3.8-4.6	288	6.8	6.0-7.7	151	2.5	2.1-2.9
1989	1,117	10.5	9.9-11.1	746	16.0	14.8-17.2	371	6.3	5.6-6.9	380	3.6	3.3-4.0	235	5.3	4.6-6.0	145	2.4	2.0-2.9
1990	1,067	10.0	9.4-10.7	686	14.6	13.5-15.8	381	6.4	5.8-7.1	377	3.5	3.1-3.9	240	5.2	4.5-5.9	137	2.2	1.8-2.6
1991	1,119	10.5	9.9-11.1	727	15.4	14.3-16.6	392	6.5	5.9-7.2	381	3.6	3.2-3.9	236	5.2	4.5-5.9	145	2.3	2.0-2.7
1992	1,114	10.2	9.6-10.9	712	14.9	13.8-16.1	402	6.6	6.0-7.3	345	3.1	2.8-3.5	216	4.5	3.9-5.2	129	2.0	1.7-2.4
1993	1,104	10.1	9.5-10.7	729	15.3	14.2-16.5	375	6.0	5.4-6.7	319	2.9	2.6-3.2	191	4.0	3.5-4.7	128	2.0	1.7-2.4
1994	1,096	9.9	9.3-10.5	725	15.0	13.9-16.2	371	6.0	5.4-6.7	336	3.0	2.7-3.4	236	4.9	4.3-5.6	100	1.6	1.3-1.9
1995	1,040	9.4	8.8-10.0	643	13.1	12.1-14.2	397	6.3	5.7-7.0	365	3.3	3.0-3.6	246	5.2	4.6-5.9	119	1.9	1.5-2.2
1996	1,136	10.2	9.6-10.8	744	14.9	13.8-16.0	392	6.3	5.7-7.0	314	2.8	2.5-3.1	204	4.3	3.7-4.9	110	1.7	1.4-2.0
1997	1,205	10.7	10.1-11.3	781	15.5	14.4-16.6	424	6.7	6.1-7.4	301	2.6	2.4-3.0	198	4.1	3.5-4.7	103	1.6	1.3-1.9
1998	1,160	10.2	9.6-10.8	733	14.5	13.4-15.5	427	6.7	6.1-7.4	330	2.9	2.6-3.2	215	4.4	3.8-5.0	115	1.7	1.4-2.0
1999	1,125	9.7	9.2-10.3	769	14.9	13.8-16.0	356	5.5	4.9-6.1	295	2.5	2.3-2.8	191	3.8	3.2-4.3	104	1.5	1.3-1.9
2000	1,122	9.6	9.0-10.2	725	14.0	13.0-15.0	397	6.1	5.5-6.8	278	2.4	2.1-2.6	175	3.5	3.0-4.1	103	1.5	1.2-1.8
2001	1,161	9.8	9.3-10.4	755	14.0	13.0-15.0	406	6.1	5.5-6.8	296	2.5	2.2-2.8	187	3.6	3.1-4.2	109	1.5	1.2-1.8
2002	1,222	10.2	9.6-10.8	799	14.7	13.6-15.7	423	6.3	5.7-6.9	280	2.3	2.0-2.6	180	3.4	3.0-4.0	100	1.4	1.1-1.7
2003	1,210	9.9	9.4-10.5	794	14.4	13.4-15.5	416	6.2	5.6-6.8	272	2.2	2.0-2.5	174	3.2	2.8-3.7	98	1.4	1.1-1.7
2004	1,267	10.3	9.7-10.9	862	15.4	14.4-16.5	405	6.0	5.4-6.6	306	2.4	2.2-2.7	207	3.7	3.2-4.3	99	1.4	1.1-1.7
2005	1,210	9.7	9.2-10.3	777	13.6	12.6-14.6	433	6.4	5.8-7.1	265	2.1	1.8-2.4	180	3.2	2.8-3.8	85	1.2	0.9-1.4
2006	1,308	10.4	9.8-11.0	913	15.8	14.8-16.9	395	5.8	5.3-6.5	278	2.2	1.9-2.5	186	3.3	2.8-3.8	92	1.2	1.0-1.5
2007	1,410	11.0	10.4-11.6	985	16.6	15.6-17.7	425	6.2	5.6-6.8	261	2.0	1.8-2.3	176	3.0	2.6-3.5	85	1.2	0.9-1.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Oral cavity and pharynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	1,392	10.7	10.1-11.3	937	15.6	14.6-16.7	455	6.5	5.9-7.1	273	2.0	1.8-2.3	195	3.2	2.8-3.7	78	1.0	0.8-1.3
2009	1,466	11.0	10.4-11.6	989	15.9	14.9-17.0	477	6.7	6.1-7.3	273	2.0	1.8-2.3	181	3.1	2.7-3.6	92	1.1	0.9-1.4
2010	1,514	11.2	10.6-11.8	1,025	16.4	15.4-17.4	489	6.8	6.2-7.5	285	2.1	1.9-2.4	187	3.1	2.6-3.6	98	1.3	1.0-1.6
2011	1,519	11.0	10.5-11.6	1,062	16.4	15.4-17.5	457	6.2	5.7-6.9	291	2.1	1.9-2.4	191	3.1	2.6-3.5	100	1.3	1.0-1.6
2012	1,537	11.1	10.5-11.7	1,047	16.2	15.2-17.3	490	6.6	6.0-7.2	308	2.2	1.9-2.4	220	3.5	3.0-4.0	88	1.0	0.8-1.3
2013	1,597	11.4	10.9-12.0	1,120	17.3	16.3-18.3	477	6.3	5.8-7.0	302	2.1	1.8-2.3	207	3.2	2.8-3.7	95	1.1	0.9-1.4
2014	1,646	11.6	11.0-12.2	1,137	17.1	16.1-18.1	509	6.8	6.2-7.4	253	1.7	1.5-2.0	176	2.7	2.3-3.2	77	1.0	0.8-1.2
2015	1,747	12.2	11.6-12.8	1,253	18.6	17.6-19.7	494	6.6	6.0-7.3	319	2.2	2.0-2.5	214	3.3	2.9-3.8	105	1.3	1.1-1.6
2016	1,855	12.6	12.1-13.3	1,282	18.7	17.6-19.8	573	7.4	6.8-8.1	339	2.2	2.0-2.5	236	3.5	3.0-3.9	103	1.2	1.0-1.5
2017	1,763	12.0	11.4-12.6	1,227	17.9	16.9-19.0	536	6.8	6.3-7.5	338	2.2	2.0-2.5	238	3.5	3.0-4.0	100	1.2	1.0-1.5
2018	1,842	12.4	11.8-13.0	1,280	18.3	17.3-19.3	562	7.2	6.5-7.8	340	2.2	2.0-2.4	230	3.3	2.9-3.8	110	1.3	1.0-1.6
2019	1,934	13.0	12.4-13.6	1,331	18.8	17.7-19.8	603	7.8	7.1-8.5	358	2.3	2.1-2.5	256	3.6	3.2-4.1	102	1.2	0.9-1.4
2020	1,807	11.8	11.2-12.4	1,281	17.7	16.7-18.8	526	6.5	5.9-7.1	340	2.2	1.9-2.4	228	3.1	2.7-3.6	112	1.3	1.1-1.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Esophagus

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	411	4.3	3.9-4.8	309	7.7	6.8-8.7	102	1.9	1.6-2.3	381	4.0	3.6-4.5	292	7.3	6.4-8.2	89	1.7	1.3-2.1
1977	420	4.4	4.0-4.9	297	7.4	6.5-8.3	123	2.3	1.9-2.7	374	3.9	3.5-4.3	273	6.7	5.9-7.6	101	1.8	1.5-2.2
1978	427	4.4	4.0-4.9	306	7.6	6.7-8.5	121	2.2	1.8-2.6	425	4.4	4.0-4.9	312	7.8	6.9-8.7	113	2.0	1.7-2.5
1979	430	4.4	4.0-4.8	302	7.2	6.4-8.1	128	2.3	1.9-2.7	369	3.8	3.4-4.2	263	6.5	5.7-7.3	106	1.9	1.5-2.3
1980	439	4.4	4.0-4.8	309	7.4	6.6-8.4	130	2.3	1.9-2.7	390	3.9	3.5-4.3	271	6.5	5.7-7.3	119	2.1	1.7-2.5
1981	434	4.3	3.9-4.7	303	7.1	6.3-8.0	131	2.3	1.9-2.7	402	4.0	3.6-4.4	297	7.1	6.3-8.0	105	1.9	1.5-2.2
1982	420	4.1	3.7-4.5	284	6.6	5.8-7.4	136	2.3	1.9-2.7	400	3.9	3.6-4.4	276	6.6	5.8-7.4	124	2.1	1.7-2.5
1983	451	4.3	3.9-4.8	322	7.4	6.6-8.3	129	2.2	1.8-2.6	419	4.2	3.8-4.6	315	7.7	6.8-8.6	104	1.8	1.5-2.2
1984	457	4.4	4.0-4.8	316	7.3	6.5-8.2	141	2.4	2.0-2.8	418	4.0	3.6-4.4	289	6.9	6.1-7.8	129	2.1	1.8-2.6
1985	427	4.2	3.8-4.6	323	7.4	6.6-8.3	104	1.7	1.4-2.1	408	3.9	3.6-4.3	305	7.0	6.2-7.9	103	1.7	1.4-2.0
1986	490	4.8	4.4-5.2	348	7.9	7.1-8.8	142	2.4	2.0-2.8	424	4.1	3.7-4.5	303	6.9	6.1-7.8	121	2.0	1.7-2.4
1987	443	4.2	3.8-4.6	307	6.9	6.1-7.8	136	2.2	1.9-2.7	435	4.2	3.8-4.6	318	7.3	6.4-8.1	117	1.9	1.6-2.3
1988	471	4.4	4.0-4.9	337	7.5	6.6-8.3	134	2.2	1.8-2.6	479	4.5	4.1-5.0	355	8.0	7.2-8.9	124	2.0	1.7-2.4
1989	483	4.5	4.1-4.9	351	7.6	6.8-8.5	132	2.2	1.8-2.6	449	4.2	3.8-4.6	338	7.5	6.7-8.4	111	1.8	1.5-2.1
1990	527	4.9	4.5-5.3	361	7.7	6.9-8.6	166	2.6	2.2-3.1	456	4.2	3.8-4.6	328	7.0	6.3-7.8	128	2.0	1.7-2.4
1991	507	4.6	4.2-5.1	352	7.4	6.6-8.2	155	2.5	2.1-2.9	482	4.4	4.0-4.8	346	7.4	6.6-8.3	136	2.1	1.8-2.5
1992	569	5.1	4.7-5.6	393	8.2	7.4-9.1	176	2.7	2.3-3.1	501	4.5	4.1-4.9	364	7.7	6.9-8.6	137	2.1	1.8-2.5
1993	561	5.0	4.6-5.5	398	8.2	7.4-9.1	163	2.5	2.1-2.9	506	4.6	4.2-5.0	379	8.0	7.2-8.8	127	1.9	1.6-2.3
1994	574	5.1	4.7-5.5	408	8.5	7.6-9.3	166	2.5	2.1-2.9	507	4.5	4.1-4.9	361	7.5	6.7-8.3	146	2.2	1.9-2.6
1995	547	4.9	4.5-5.3	392	8.2	7.4-9.1	155	2.3	2.0-2.7	554	4.9	4.5-5.4	417	8.8	7.9-9.7	137	2.1	1.7-2.4
1996	594	5.3	4.9-5.7	422	8.6	7.8-9.5	172	2.6	2.2-3.0	522	4.6	4.2-5.0	371	7.7	6.9-8.5	151	2.2	1.9-2.6
1997	533	4.7	4.3-5.1	375	7.7	6.9-8.5	158	2.4	2.0-2.8	522	4.6	4.2-5.0	397	8.2	7.4-9.0	125	1.8	1.5-2.2
1998	603	5.2	4.8-5.7	452	9.0	8.2-9.9	151	2.3	1.9-2.7	538	4.7	4.3-5.1	403	8.1	7.3-8.9	135	2.0	1.7-2.4
1999	600	5.1	4.7-5.6	436	8.6	7.8-9.5	164	2.4	2.0-2.8	536	4.6	4.2-5.0	392	7.8	7.1-8.7	144	2.1	1.8-2.5
2000	643	5.4	5.0-5.9	465	9.1	8.3-10.0	178	2.6	2.2-3.0	564	4.8	4.4-5.2	408	8.1	7.3-9.0	156	2.2	1.9-2.6
2001	594	5.0	4.6-5.4	441	8.4	7.7-9.3	153	2.2	1.9-2.6	579	4.8	4.5-5.3	440	8.5	7.8-9.4	139	1.9	1.6-2.3
2002	629	5.2	4.8-5.6	474	9.0	8.2-9.9	155	2.2	1.8-2.6	559	4.6	4.2-5.0	424	8.1	7.3-8.9	135	1.9	1.6-2.2
2003	668	5.5	5.1-5.9	507	9.4	8.6-10.3	161	2.2	1.9-2.6	607	4.9	4.5-5.3	450	8.6	7.8-9.4	157	2.1	1.8-2.5
2004	682	5.5	5.1-5.9	515	9.6	8.8-10.4	167	2.3	2.0-2.7	637	5.1	4.7-5.6	491	9.1	8.3-9.9	146	2.0	1.7-2.3
2005	734	5.9	5.5-6.3	566	10.3	9.5-11.2	168	2.3	2.0-2.7	645	5.1	4.7-5.6	499	9.3	8.5-10.1	146	2.0	1.7-2.3
2006	749	5.9	5.5-6.4	583	10.5	9.6-11.3	166	2.3	2.0-2.7	584	4.6	4.3-5.0	461	8.4	7.6-9.2	123	1.7	1.4-2.1
2007	721	5.6	5.2-6.0	555	9.7	8.9-10.6	166	2.3	2.0-2.7	624	4.8	4.5-5.2	463	8.3	7.5-9.1	161	2.2	1.8-2.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Esophagus

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	748	5.8	5.3-6.2	582	10.1	9.3-11.0	166	2.2	1.9-2.6	616	4.7	4.3-5.1	485	8.4	7.6-9.2	131	1.8	1.5-2.1
2009	786	5.9	5.5-6.3	615	10.4	9.5-11.2	171	2.3	1.9-2.6	603	4.5	4.1-4.9	484	8.2	7.5-9.0	119	1.5	1.3-1.8
2010	729	5.4	5.0-5.8	552	9.1	8.4-10.0	177	2.3	2.0-2.7	638	4.7	4.4-5.1	510	8.5	7.8-9.3	128	1.6	1.3-1.9
2011	713	5.2	4.8-5.6	526	8.4	7.7-9.2	187	2.4	2.1-2.8	654	4.8	4.4-5.1	499	8.2	7.5-9.0	155	2.0	1.7-2.3
2012	690	4.9	4.6-5.3	549	8.7	8.0-9.5	141	1.8	1.5-2.1	622	4.5	4.1-4.8	486	7.9	7.2-8.7	136	1.7	1.4-2.0
2013	742	5.2	4.8-5.6	577	8.9	8.2-9.7	165	2.0	1.7-2.4	641	4.5	4.1-4.9	494	7.8	7.1-8.5	147	1.8	1.6-2.2
2014	732	5.0	4.6-5.3	568	8.5	7.8-9.3	164	2.0	1.7-2.4	628	4.4	4.0-4.8	511	8.0	7.3-8.8	117	1.4	1.2-1.7
2015	794	5.4	5.1-5.8	604	9.0	8.3-9.8	190	2.4	2.1-2.8	589	4.0	3.7-4.3	457	6.9	6.3-7.6	132	1.6	1.3-1.9
2016	790	5.3	5.0-5.7	603	9.0	8.2-9.7	187	2.3	2.0-2.7	644	4.3	4.0-4.7	509	7.6	6.9-8.3	135	1.6	1.4-2.0
2017	739	4.9	4.6-5.3	568	8.2	7.5-9.0	171	2.1	1.8-2.5	624	4.1	3.8-4.4	505	7.3	6.6-8.0	119	1.4	1.2-1.7
2018	776	5.1	4.8-5.5	601	8.7	8.0-9.5	175	2.1	1.8-2.4	606	4.0	3.6-4.3	467	6.7	6.1-7.4	139	1.7	1.4-2.0
2019	797	5.1	4.7-5.5	615	8.6	7.9-9.4	182	2.1	1.8-2.4	604	3.8	3.5-4.1	479	6.7	6.1-7.4	125	1.4	1.2-1.7
2020	714	4.5	4.2-4.9	566	8.0	7.3-8.7	148	1.7	1.4-2.0	601	3.8	3.5-4.2	477	6.9	6.2-7.5	124	1.4	1.2-1.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State Excl New York City
Stomach**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,044	11.6	10.9-12.3	602	16.0	14.6-17.4	442	8.4	7.7-9.3	805	9.1	8.4-9.7	448	12.3	11.1-13.5	357	6.9	6.2-7.6
1977	1,028	11.3	10.6-12.0	607	16.2	14.9-17.6	421	7.9	7.1-8.7	804	8.9	8.3-9.5	477	13.1	11.9-14.4	327	6.1	5.5-6.8
1978	1,038	11.2	10.5-11.9	593	15.7	14.4-17.1	445	8.2	7.4-9.0	756	8.3	7.7-9.0	419	11.3	10.2-12.5	337	6.2	5.6-7.0
1979	1,055	11.2	10.5-11.9	617	16.0	14.7-17.4	438	7.9	7.2-8.7	775	8.3	7.7-8.9	447	11.8	10.6-13.0	328	5.9	5.3-6.6
1980	1,013	10.6	10.0-11.3	606	15.4	14.1-16.8	407	7.4	6.7-8.1	782	8.3	7.8-9.0	443	11.8	10.6-13.0	339	6.1	5.5-6.8
1981	1,089	11.3	10.6-12.0	660	16.6	15.3-18.0	429	7.6	6.9-8.3	769	8.0	7.5-8.6	454	11.6	10.5-12.8	315	5.6	5.0-6.2
1982	1,050	10.7	10.0-11.3	632	16.0	14.7-17.3	418	7.2	6.5-7.9	748	7.7	7.2-8.3	458	11.8	10.7-13.0	290	5.0	4.5-5.6
1983	1,040	10.5	9.9-11.2	626	15.5	14.2-16.8	414	7.1	6.4-7.8	734	7.5	6.9-8.0	440	11.1	10.1-12.3	294	5.0	4.5-5.7
1984	1,035	10.3	9.7-11.0	635	15.3	14.1-16.6	400	6.7	6.0-7.4	775	7.7	7.2-8.3	459	11.1	10.1-12.2	316	5.3	4.7-5.9
1985	1,063	10.5	9.9-11.2	665	16.1	14.9-17.5	398	6.6	6.0-7.3	705	7.0	6.5-7.6	408	10.3	9.3-11.4	297	5.0	4.4-5.6
1986	1,021	10.0	9.4-10.6	641	15.6	14.3-16.9	380	6.2	5.6-6.9	751	7.4	6.9-8.0	459	11.2	10.1-12.3	292	4.8	4.2-5.3
1987	1,052	10.2	9.6-10.9	631	15.1	13.9-16.4	421	6.9	6.2-7.6	728	7.1	6.6-7.6	435	10.7	9.6-11.8	293	4.8	4.2-5.4
1988	1,016	9.9	9.3-10.5	583	13.8	12.7-15.1	433	7.1	6.4-7.8	698	6.8	6.3-7.3	390	9.5	8.5-10.5	308	5.0	4.5-5.6
1989	984	9.4	8.8-10.0	648	15.2	14.0-16.5	336	5.3	4.8-5.9	746	7.1	6.6-7.7	442	10.7	9.7-11.8	304	4.8	4.3-5.4
1990	1,083	10.2	9.6-10.8	688	15.7	14.5-16.9	395	6.2	5.6-6.9	673	6.4	5.9-6.9	424	10.0	9.0-11.0	249	3.9	3.4-4.4
1991	979	9.1	8.5-9.7	609	13.8	12.7-15.0	370	5.8	5.2-6.4	663	6.2	5.7-6.7	430	10.1	9.1-11.1	233	3.6	3.2-4.1
1992	1,017	9.3	8.8-9.9	629	14.2	13.1-15.4	388	5.9	5.4-6.6	688	6.3	5.9-6.8	419	9.8	8.8-10.8	269	4.1	3.6-4.6
1993	1,074	9.8	9.2-10.4	658	14.6	13.5-15.8	416	6.2	5.6-6.9	652	6.0	5.5-6.4	400	9.3	8.3-10.3	252	3.8	3.3-4.3
1994	1,024	9.2	8.7-9.8	633	13.9	12.8-15.1	391	5.9	5.3-6.5	683	6.2	5.7-6.7	408	9.2	8.3-10.2	275	4.1	3.6-4.6
1995	1,004	8.9	8.4-9.5	596	12.9	11.9-14.0	408	6.1	5.6-6.8	644	5.8	5.3-6.2	365	8.2	7.4-9.2	279	4.1	3.6-4.6
1996	995	8.8	8.3-9.4	615	13.2	12.2-14.4	380	5.6	5.0-6.2	592	5.2	4.8-5.7	349	7.7	6.9-8.6	243	3.5	3.1-4.0
1997	1,022	8.9	8.4-9.5	623	13.1	12.1-14.2	399	5.9	5.3-6.5	569	5.0	4.6-5.4	325	6.9	6.2-7.8	244	3.5	3.1-4.0
1998	1,035	8.9	8.4-9.5	661	13.5	12.5-14.6	374	5.4	4.9-6.0	569	4.9	4.5-5.3	329	6.9	6.2-7.7	240	3.4	3.0-3.9
1999	983	8.4	7.9-9.0	625	12.8	11.8-13.9	358	5.2	4.7-5.8	558	4.8	4.4-5.2	342	7.1	6.3-7.9	216	3.1	2.7-3.5
2000	1,004	8.4	7.9-9.0	615	12.5	11.5-13.5	389	5.4	4.9-6.0	556	4.7	4.3-5.1	312	6.6	5.8-7.3	244	3.4	3.0-3.9
2001	961	8.0	7.5-8.6	602	12.0	11.0-13.0	359	5.2	4.6-5.7	518	4.3	3.9-4.7	301	6.1	5.4-6.9	217	3.0	2.6-3.4
2002	972	8.0	7.5-8.5	591	11.6	10.7-12.6	381	5.4	4.9-6.0	509	4.2	3.8-4.5	293	5.9	5.3-6.7	216	2.9	2.5-3.4
2003	1,012	8.3	7.8-8.8	631	12.1	11.2-13.1	381	5.4	4.8-6.0	483	3.9	3.6-4.3	284	5.6	5.0-6.3	199	2.7	2.3-3.1
2004	959	7.7	7.3-8.3	575	11.0	10.2-12.0	384	5.4	4.8-5.9	533	4.3	3.9-4.7	310	6.1	5.4-6.8	223	3.0	2.6-3.5
2005	928	7.4	6.9-7.9	557	10.2	9.4-11.1	371	5.2	4.7-5.8	498	3.9	3.6-4.3	287	5.5	4.9-6.2	211	2.8	2.4-3.2
2006	920	7.3	6.8-7.8	564	10.3	9.5-11.2	356	5.0	4.5-5.5	435	3.4	3.1-3.7	252	4.6	4.1-5.2	183	2.5	2.1-2.9
2007	963	7.5	7.1-8.0	613	10.9	10.1-11.9	350	4.9	4.4-5.4	437	3.4	3.1-3.7	258	4.6	4.1-5.3	179	2.5	2.1-2.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Stomach

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	881	6.8	6.3-7.2	567	10.1	9.3-11.0	314	4.3	3.8-4.8	466	3.5	3.2-3.8	284	5.1	4.5-5.7	182	2.3	2.0-2.7
2009	871	6.7	6.2-7.1	550	9.6	8.8-10.5	321	4.3	3.8-4.8	410	3.1	2.8-3.4	251	4.5	3.9-5.1	159	2.1	1.8-2.4
2010	933	7.0	6.6-7.5	594	10.0	9.2-10.9	339	4.5	4.1-5.1	399	3.0	2.7-3.3	245	4.3	3.8-4.9	154	2.0	1.7-2.3
2011	1,032	7.6	7.1-8.1	642	10.7	9.9-11.6	390	5.2	4.7-5.8	378	2.7	2.5-3.0	229	3.9	3.4-4.4	149	1.9	1.6-2.2
2012	988	7.2	6.8-7.7	612	10.0	9.2-10.9	376	5.0	4.5-5.5	418	3.0	2.7-3.3	247	4.1	3.6-4.7	171	2.2	1.9-2.5
2013	1,073	7.8	7.3-8.3	672	10.9	10.1-11.8	401	5.3	4.8-5.9	420	3.0	2.7-3.3	254	4.1	3.6-4.7	166	2.1	1.8-2.4
2014	1,010	7.2	6.7-7.6	662	10.5	9.7-11.3	348	4.6	4.1-5.1	410	2.9	2.6-3.2	260	4.2	3.7-4.7	150	1.9	1.6-2.3
2015	1,010	7.1	6.7-7.6	632	9.8	9.0-10.6	378	5.0	4.5-5.5	377	2.6	2.4-2.9	233	3.7	3.2-4.2	144	1.8	1.5-2.2
2016	1,059	7.4	7.0-7.9	645	9.8	9.1-10.7	414	5.5	4.9-6.1	379	2.6	2.3-2.9	235	3.5	3.1-4.0	144	1.8	1.5-2.2
2017	1,082	7.5	7.0-8.0	654	9.9	9.1-10.7	428	5.6	5.0-6.2	363	2.5	2.2-2.7	203	3.0	2.6-3.5	160	2.0	1.7-2.4
2018	1,055	7.3	6.8-7.8	633	9.5	8.7-10.3	422	5.5	5.0-6.1	379	2.5	2.3-2.8	228	3.5	3.0-4.0	151	1.8	1.5-2.1
2019	1,068	7.1	6.6-7.5	660	9.6	8.8-10.4	408	5.0	4.5-5.5	379	2.6	2.3-2.8	227	3.5	3.0-4.0	152	1.8	1.5-2.1
2020	974	6.6	6.1-7.0	613	8.9	8.2-9.7	361	4.7	4.2-5.2	354	2.3	2.1-2.6	199	2.9	2.5-3.4	155	1.9	1.6-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State Excl New York City
Colorectal**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	6,202	68.0	66.3-69.8	3,029	80.5	77.5-83.6	3,173	59.9	57.8-62.0	3,102	34.8	33.6-36.1	1,473	40.8	38.6-43.1	1,629	31.1	29.5-32.6
1977	6,377	68.7	67.0-70.4	3,089	81.0	78.0-84.1	3,288	60.8	58.7-62.9	3,120	34.7	33.4-35.9	1,497	41.4	39.2-43.6	1,623	30.3	28.9-31.8
1978	6,467	69.2	67.5-71.0	3,150	82.1	79.1-85.1	3,317	60.7	58.6-62.8	3,304	36.0	34.8-37.3	1,609	43.8	41.5-46.1	1,695	31.1	29.7-32.7
1979	6,524	68.7	67.0-70.4	3,126	80.2	77.3-83.2	3,398	61.0	58.9-63.1	3,280	35.2	34.0-36.5	1,557	42.4	40.2-44.7	1,723	31.0	29.5-32.5
1980	6,741	69.7	68.0-71.4	3,373	84.9	81.9-87.9	3,368	59.4	57.4-61.5	3,226	34.1	32.9-35.3	1,563	41.3	39.1-43.5	1,663	29.5	28.1-30.9
1981	6,860	70.4	68.7-72.1	3,437	87.3	84.2-90.4	3,423	59.6	57.6-61.6	3,298	34.5	33.3-35.7	1,610	42.3	40.1-44.5	1,688	29.5	28.1-30.9
1982	6,845	69.3	67.6-71.0	3,382	83.6	80.6-86.6	3,463	59.5	57.5-61.5	3,303	34.1	33.0-35.3	1,621	42.7	40.5-44.9	1,682	28.9	27.5-30.3
1983	7,008	70.3	68.7-72.0	3,491	86.2	83.2-89.3	3,517	59.8	57.8-61.8	3,232	32.9	31.8-34.1	1,579	41.0	38.9-43.2	1,653	27.9	26.5-29.3
1984	7,076	70.4	68.7-72.1	3,459	84.3	81.4-87.3	3,617	60.9	58.9-63.0	3,372	34.1	32.9-35.2	1,658	42.5	40.3-44.7	1,714	28.8	27.4-30.2
1985	7,312	71.9	70.3-73.6	3,671	88.5	85.5-91.5	3,641	60.8	58.8-62.8	3,301	33.1	32.0-34.3	1,593	40.5	38.4-42.6	1,708	28.3	27.0-29.7
1986	7,051	68.9	67.3-70.5	3,474	83.5	80.7-86.5	3,577	59.2	57.2-61.2	3,296	32.7	31.5-33.8	1,627	41.2	39.1-43.3	1,669	27.4	26.1-28.8
1987	7,014	67.8	66.2-69.4	3,486	83.5	80.6-86.4	3,528	57.8	55.8-59.7	3,292	32.3	31.2-33.4	1,659	41.5	39.4-43.6	1,633	26.5	25.2-27.8
1988	6,782	64.8	63.2-66.4	3,342	78.4	75.7-81.3	3,440	55.6	53.8-57.5	3,271	31.7	30.6-32.8	1,638	40.1	38.1-42.2	1,633	26.3	25.0-27.6
1989	6,748	64.0	62.5-65.6	3,382	79.0	76.2-81.8	3,366	54.1	52.3-56.0	3,054	29.2	28.2-30.3	1,494	36.3	34.4-38.3	1,560	24.8	23.5-26.0
1990	6,722	62.7	61.2-64.2	3,323	75.9	73.3-78.7	3,399	53.5	51.7-55.4	2,958	27.9	26.9-28.9	1,508	36.2	34.3-38.1	1,450	22.6	21.5-23.8
1991	6,676	61.7	60.2-63.2	3,320	75.0	72.4-77.7	3,356	52.6	50.8-54.4	3,005	28.1	27.1-29.1	1,455	34.3	32.5-36.2	1,550	23.9	22.7-25.2
1992	6,604	60.3	58.9-61.8	3,336	74.6	72.0-77.3	3,268	50.5	48.8-52.3	2,855	26.3	25.3-27.3	1,381	32.2	30.5-34.1	1,474	22.5	21.3-23.7
1993	6,707	60.5	59.0-62.0	3,256	71.4	68.9-74.0	3,451	52.7	50.9-54.5	2,947	26.7	25.8-27.7	1,437	33.1	31.4-35.0	1,510	22.6	21.4-23.8
1994	6,901	61.8	60.4-63.3	3,422	75.2	72.6-77.8	3,479	52.9	51.2-54.7	3,005	27.1	26.1-28.1	1,433	33.0	31.2-34.8	1,572	23.3	22.1-24.5
1995	6,807	60.4	59.0-61.9	3,367	73.0	70.5-75.6	3,440	51.7	50.0-53.5	2,929	26.0	25.1-27.0	1,402	32.1	30.4-33.9	1,527	22.4	21.2-23.5
1996	6,681	58.9	57.5-60.3	3,378	72.5	70.0-75.0	3,303	49.4	47.7-51.1	2,792	24.6	23.7-25.5	1,382	30.8	29.2-32.5	1,410	20.5	19.4-21.6
1997	7,005	61.1	59.7-62.5	3,487	73.2	70.7-75.7	3,518	52.3	50.5-54.0	2,675	23.3	22.4-24.2	1,333	29.2	27.6-30.8	1,342	19.3	18.3-20.4
1998	7,364	63.5	62.1-65.0	3,639	75.1	72.7-77.7	3,725	55.0	53.3-56.9	2,660	22.9	22.0-23.8	1,290	27.4	25.9-29.0	1,370	19.5	18.5-20.6
1999	7,477	63.7	62.3-65.2	3,719	76.1	73.7-78.6	3,758	54.8	53.0-56.6	2,646	22.4	21.6-23.3	1,292	27.5	26.0-29.1	1,354	19.0	18.0-20.0
2000	7,436	62.6	61.2-64.0	3,685	74.2	71.8-76.7	3,751	54.1	52.3-55.9	2,673	22.3	21.5-23.2	1,346	28.2	26.7-29.8	1,327	18.2	17.2-19.2
2001	7,122	59.4	58.0-60.8	3,534	70.1	67.7-72.4	3,588	51.3	49.6-53.1	2,560	21.2	20.3-22.0	1,214	25.1	23.7-26.6	1,346	18.5	17.5-19.5
2002	7,121	58.7	57.3-60.1	3,545	68.9	66.6-71.2	3,576	50.7	49.1-52.4	2,578	21.0	20.2-21.9	1,250	25.3	23.9-26.8	1,328	18.0	17.1-19.1
2003	6,802	55.2	53.9-56.6	3,317	63.5	61.4-65.8	3,485	49.0	47.3-50.6	2,359	19.0	18.2-19.7	1,165	23.2	21.8-24.5	1,194	16.1	15.2-17.0
2004	6,719	54.0	52.7-55.3	3,254	61.4	59.3-63.5	3,465	48.4	46.8-50.1	2,344	18.6	17.9-19.4	1,130	21.9	20.6-23.2	1,214	16.0	15.1-16.9
2005	6,378	50.7	49.4-51.9	3,206	59.7	57.6-61.8	3,172	44.0	42.5-45.6	2,145	16.8	16.1-17.5	1,049	20.3	19.1-21.6	1,096	14.4	13.5-15.3
2006	6,078	47.9	46.7-49.2	3,008	55.2	53.2-57.2	3,070	42.2	40.7-43.8	2,104	16.4	15.7-17.1	1,008	19.3	18.1-20.5	1,096	14.3	13.4-15.2
2007	6,224	48.5	47.3-49.8	3,073	55.1	53.2-57.1	3,151	43.3	41.8-44.9	2,128	16.3	15.6-17.0	1,046	19.4	18.2-20.6	1,082	14.0	13.2-14.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State Excl New York City
Colorectal**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	6,126	47.0	45.8-48.2	3,023	53.5	51.5-55.4	3,103	42.2	40.7-43.7	2,049	15.5	14.8-16.2	994	18.1	17.0-19.3	1,055	13.4	12.6-14.3
2009	5,736	43.5	42.3-44.6	2,861	49.6	47.8-51.5	2,875	38.5	37.1-40.0	2,006	15.0	14.4-15.7	998	17.9	16.8-19.1	1,008	12.9	12.1-13.8
2010	5,807	43.4	42.3-44.6	2,866	48.7	46.9-50.6	2,941	39.1	37.6-40.6	1,963	14.4	13.7-15.0	962	16.7	15.7-17.9	1,001	12.4	11.7-13.3
2011	5,648	42.0	40.9-43.2	2,791	47.0	45.2-48.8	2,857	38.0	36.6-39.5	1,948	14.0	13.3-14.6	965	16.3	15.3-17.4	983	12.0	11.3-12.8
2012	5,373	39.4	38.3-40.5	2,760	45.4	43.7-47.2	2,613	34.5	33.2-35.9	1,889	13.4	12.8-14.1	911	15.4	14.4-16.5	978	12.0	11.2-12.8
2013	5,459	39.5	38.4-40.6	2,806	45.3	43.6-47.0	2,653	34.9	33.5-36.3	1,882	13.3	12.7-13.9	992	16.4	15.4-17.5	890	10.7	10.0-11.5
2014	5,464	39.0	38.0-40.1	2,807	44.7	43.1-46.5	2,657	34.4	33.1-35.8	1,852	12.8	12.2-13.4	955	15.4	14.4-16.4	897	10.8	10.1-11.6
2015	5,492	39.1	38.0-40.1	2,793	43.8	42.1-45.5	2,699	35.1	33.7-36.5	1,889	12.9	12.3-13.5	964	15.3	14.3-16.3	925	11.0	10.3-11.8
2016	5,512	38.9	37.8-40.0	2,824	43.8	42.2-45.5	2,688	34.7	33.3-36.1	1,727	11.7	11.2-12.3	858	13.3	12.4-14.3	869	10.4	9.6-11.1
2017	5,529	38.8	37.7-39.8	2,834	43.6	42.0-45.3	2,695	34.7	33.4-36.1	1,879	12.6	12.0-13.2	943	14.5	13.6-15.5	936	11.1	10.3-11.8
2018	5,395	37.4	36.4-38.4	2,790	42.3	40.7-44.0	2,605	33.1	31.8-34.5	1,739	11.6	11.1-12.2	878	13.5	12.6-14.4	861	10.2	9.5-10.9
2019	5,437	37.4	36.4-38.5	2,808	42.0	40.4-43.7	2,629	33.5	32.2-34.9	1,744	11.3	10.8-11.9	908	13.6	12.7-14.5	836	9.4	8.8-10.1
2020	4,959	34.0	33.0-35.0	2,638	39.3	37.8-40.9	2,321	29.5	28.3-30.8	1,753	11.4	10.8-12.0	916	13.5	12.6-14.4	837	9.7	9.0-10.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Colon excluding rectum

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	4,317	47.8	46.3-49.3	1,984	53.7	51.2-56.3	2,333	44.1	42.3-45.9	2,435	27.3	26.2-28.5	1,092	30.2	28.4-32.2	1,343	25.6	24.2-27.0
1977	4,459	48.5	47.1-50.0	2,047	55.1	52.6-57.7	2,412	44.7	42.9-46.6	2,522	28.1	27.0-29.2	1,177	32.6	30.6-34.6	1,345	25.2	23.8-26.6
1978	4,551	49.0	47.6-50.5	2,101	55.7	53.2-58.3	2,450	44.8	43.1-46.7	2,648	28.9	27.8-30.0	1,233	33.6	31.6-35.6	1,415	26.0	24.6-27.4
1979	4,634	49.3	47.9-50.8	2,106	55.2	52.7-57.8	2,528	45.5	43.7-47.3	2,711	29.1	28.0-30.2	1,254	34.1	32.1-36.1	1,457	26.1	24.8-27.5
1980	4,789	49.9	48.5-51.4	2,305	58.8	56.3-61.4	2,484	43.9	42.2-45.7	2,629	27.7	26.7-28.8	1,254	33.2	31.3-35.2	1,375	24.3	23.0-25.6
1981	4,917	50.8	49.4-52.3	2,364	60.8	58.2-63.4	2,553	44.6	42.8-46.4	2,774	29.1	28.0-30.3	1,335	35.3	33.3-37.4	1,439	25.2	23.9-26.6
1982	4,833	49.3	47.9-50.8	2,265	56.9	54.5-59.4	2,568	44.1	42.4-45.9	2,725	28.2	27.1-29.3	1,311	34.5	32.6-36.6	1,414	24.3	23.0-25.6
1983	4,948	50.0	48.6-51.5	2,321	58.7	56.2-61.2	2,627	44.8	43.1-46.5	2,711	27.6	26.5-28.7	1,295	33.7	31.8-35.7	1,416	23.9	22.6-25.2
1984	5,095	51.0	49.6-52.4	2,333	57.8	55.3-60.3	2,762	46.5	44.7-48.2	2,831	28.6	27.5-29.7	1,362	34.8	32.9-36.8	1,469	24.6	23.4-25.9
1985	5,231	51.7	50.3-53.1	2,513	61.2	58.7-63.8	2,718	45.3	43.6-47.1	2,781	27.9	26.8-28.9	1,314	33.3	31.4-35.2	1,467	24.4	23.1-25.7
1986	5,092	50.0	48.6-51.4	2,386	58.2	55.8-60.7	2,706	44.7	43.0-46.5	2,795	27.7	26.7-28.7	1,354	34.4	32.5-36.4	1,441	23.6	22.4-24.9
1987	4,987	48.4	47.0-49.8	2,381	57.9	55.5-60.4	2,606	42.5	40.8-44.2	2,787	27.3	26.3-28.3	1,396	34.9	33.0-36.8	1,391	22.6	21.4-23.8
1988	4,872	46.7	45.4-48.0	2,263	53.7	51.4-56.1	2,609	42.1	40.5-43.7	2,810	27.2	26.2-28.2	1,389	33.9	32.1-35.9	1,421	22.8	21.6-24.1
1989	4,774	45.5	44.2-46.8	2,307	55.0	52.7-57.4	2,467	39.5	37.9-41.1	2,597	24.8	23.9-25.8	1,239	30.1	28.4-31.9	1,358	21.5	20.4-22.7
1990	4,753	44.5	43.2-45.7	2,246	52.3	50.0-54.6	2,507	39.4	37.8-41.0	2,566	24.2	23.2-25.1	1,313	31.6	29.9-33.5	1,253	19.5	18.5-20.7
1991	4,823	44.6	43.4-45.9	2,310	52.9	50.7-55.2	2,513	39.2	37.6-40.7	2,541	23.8	22.9-24.7	1,215	28.7	27.1-30.5	1,326	20.4	19.3-21.6
1992	4,737	43.4	42.2-44.7	2,259	51.2	49.0-53.4	2,478	38.2	36.7-39.8	2,419	22.3	21.4-23.2	1,137	26.7	25.1-28.4	1,282	19.6	18.5-20.7
1993	4,858	43.9	42.7-45.2	2,234	49.6	47.5-51.8	2,624	39.9	38.4-41.5	2,532	23.0	22.1-23.9	1,227	28.4	26.8-30.1	1,305	19.5	18.5-20.6
1994	5,010	44.9	43.7-46.2	2,396	53.3	51.1-55.5	2,614	39.6	38.1-41.1	2,582	23.2	22.3-24.1	1,204	27.9	26.3-29.6	1,378	20.4	19.3-21.5
1995	4,945	43.8	42.6-45.1	2,353	51.5	49.3-53.6	2,592	38.7	37.2-40.2	2,472	22.0	21.1-22.8	1,156	26.6	25.1-28.3	1,316	19.2	18.2-20.3
1996	4,769	42.0	40.8-43.2	2,284	49.8	47.7-51.9	2,485	36.9	35.4-38.4	2,359	20.8	19.9-21.6	1,150	25.8	24.3-27.4	1,209	17.5	16.5-18.5
1997	5,164	45.0	43.8-46.2	2,436	51.6	49.5-53.7	2,728	40.3	38.8-41.8	2,312	20.1	19.3-21.0	1,124	24.6	23.1-26.1	1,188	17.0	16.1-18.1
1998	5,378	46.3	45.1-47.6	2,570	53.7	51.6-55.8	2,808	41.3	39.8-42.9	2,254	19.4	18.6-20.2	1,076	23.0	21.6-24.4	1,178	16.7	15.8-17.7
1999	5,398	45.9	44.7-47.2	2,560	52.8	50.8-54.9	2,838	41.0	39.5-42.5	2,220	18.8	18.0-19.6	1,061	22.7	21.3-24.1	1,159	16.2	15.3-17.2
2000	5,421	45.5	44.3-46.8	2,558	52.1	50.1-54.2	2,863	40.9	39.4-42.5	2,255	18.8	18.0-19.6	1,117	23.6	22.2-25.0	1,138	15.6	14.7-16.5
2001	5,170	43.0	41.8-44.2	2,456	49.1	47.2-51.1	2,714	38.5	37.0-40.0	2,171	17.9	17.2-18.7	1,011	21.0	19.7-22.4	1,160	15.9	15.0-16.9
2002	5,185	42.6	41.5-43.8	2,476	48.6	46.7-50.6	2,709	38.2	36.8-39.7	2,134	17.4	16.7-18.1	1,017	20.7	19.4-22.0	1,117	15.1	14.2-16.0
2003	4,947	40.1	39.0-41.2	2,308	44.7	42.8-46.5	2,639	36.8	35.4-38.3	1,990	16.0	15.3-16.7	954	19.1	17.9-20.3	1,036	13.9	13.1-14.8
2004	4,842	38.8	37.8-40.0	2,231	42.6	40.8-44.4	2,611	36.1	34.7-37.5	1,946	15.4	14.8-16.1	917	17.8	16.7-19.0	1,029	13.5	12.7-14.4
2005	4,650	36.9	35.9-38.0	2,253	42.4	40.6-44.2	2,397	33.0	31.7-34.4	1,768	13.8	13.2-14.5	841	16.4	15.3-17.6	927	12.1	11.4-13.0
2006	4,466	35.1	34.1-36.2	2,129	39.5	37.8-41.3	2,337	31.7	30.5-33.1	1,750	13.6	13.0-14.2	826	15.8	14.8-17.0	924	12.0	11.2-12.8
2007	4,555	35.5	34.5-36.6	2,161	39.2	37.5-40.9	2,394	32.5	31.2-33.9	1,781	13.6	13.0-14.3	852	15.9	14.8-17.0	929	12.0	11.2-12.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Colon excluding rectum

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	4,449	34.1	33.1-35.1	2,078	37.0	35.4-38.6	2,371	32.1	30.8-33.4	1,675	12.7	12.0-13.3	796	14.6	13.6-15.6	879	11.2	10.4-11.9
2009	4,102	31.0	30.0-32.0	1,943	33.9	32.4-35.5	2,159	28.6	27.4-29.8	1,653	12.4	11.8-13.0	800	14.5	13.5-15.5	853	10.9	10.2-11.7
2010	4,157	31.1	30.1-32.1	1,948	33.7	32.2-35.2	2,209	29.0	27.8-30.3	1,598	11.6	11.1-12.2	759	13.3	12.3-14.3	839	10.3	9.6-11.1
2011	4,030	29.9	29.0-30.9	1,873	31.9	30.5-33.5	2,157	28.3	27.1-29.5	1,600	11.4	10.9-12.0	764	13.0	12.1-14.0	836	10.2	9.5-10.9
2012	3,861	28.2	27.3-29.1	1,874	31.1	29.7-32.5	1,987	25.9	24.7-27.1	1,564	11.1	10.6-11.7	720	12.2	11.3-13.1	844	10.3	9.6-11.1
2013	3,819	27.5	26.6-28.4	1,872	30.5	29.1-31.9	1,947	25.1	24.0-26.3	1,532	10.8	10.2-11.3	786	13.0	12.1-14.0	746	8.9	8.3-9.6
2014	3,869	27.5	26.7-28.5	1,875	30.2	28.8-31.6	1,994	25.5	24.4-26.7	1,489	10.3	9.8-10.8	747	12.1	11.3-13.1	742	8.9	8.2-9.6
2015	3,900	27.8	26.9-28.7	1,875	29.8	28.5-31.3	2,025	26.0	24.9-27.2	1,538	10.5	9.9-11.0	759	12.1	11.2-13.0	779	9.2	8.6-9.9
2016	3,948	27.8	26.9-28.7	1,898	29.6	28.3-31.0	2,050	26.2	25.1-27.4	1,353	9.1	8.7-9.7	646	10.0	9.2-10.9	707	8.4	7.7-9.0
2017	3,938	27.4	26.6-28.4	1,914	29.6	28.3-31.0	2,024	25.7	24.6-26.9	1,493	10.0	9.5-10.6	732	11.3	10.5-12.2	761	8.9	8.3-9.6
2018	3,803	26.3	25.4-27.1	1,864	28.6	27.3-29.9	1,939	24.2	23.1-25.4	1,356	9.0	8.6-9.6	657	10.2	9.4-11.0	699	8.2	7.6-8.9
2019	3,826	26.0	25.2-26.9	1,869	27.9	26.6-29.2	1,957	24.5	23.4-25.7	1,363	8.9	8.4-9.4	688	10.4	9.6-11.2	675	7.5	7.0-8.2
2020	3,491	23.8	23.0-24.6	1,769	26.5	25.2-27.8	1,722	21.6	20.5-22.7	1,334	8.6	8.2-9.1	675	10.0	9.2-10.8	659	7.6	7.0-8.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Rectum & rectosigmoid

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,885	20.2	19.3-21.2	1,045	26.8	25.1-28.6	840	15.8	14.7-16.9	667	7.5	6.9-8.1	381	10.6	9.5-11.8	286	5.5	4.8-6.1
1977	1,918	20.1	19.2-21.1	1,042	25.9	24.2-27.6	876	16.1	15.0-17.2	598	6.6	6.1-7.1	320	8.8	7.8-9.8	278	5.1	4.6-5.8
1978	1,916	20.2	19.3-21.1	1,049	26.3	24.7-28.1	867	15.9	14.8-17.0	656	7.1	6.6-7.7	376	10.2	9.1-11.3	280	5.2	4.6-5.8
1979	1,890	19.4	18.6-20.3	1,020	25.0	23.4-26.7	870	15.5	14.5-16.6	569	6.2	5.6-6.7	303	8.4	7.4-9.4	266	4.8	4.2-5.4
1980	1,952	19.8	18.9-20.7	1,068	26.1	24.5-27.8	884	15.5	14.5-16.6	597	6.3	5.8-6.9	309	8.1	7.2-9.1	288	5.1	4.6-5.8
1981	1,943	19.6	18.7-20.5	1,073	26.5	24.9-28.2	870	15.0	14.0-16.0	524	5.4	4.9-5.9	275	7.0	6.1-7.9	249	4.3	3.8-4.8
1982	2,012	20.0	19.1-20.9	1,117	26.7	25.1-28.4	895	15.4	14.4-16.4	578	6.0	5.5-6.5	310	8.1	7.2-9.1	268	4.6	4.1-5.2
1983	2,060	20.3	19.4-21.2	1,170	27.5	25.9-29.2	890	15.0	14.1-16.1	521	5.3	4.9-5.8	284	7.2	6.4-8.2	237	4.0	3.5-4.6
1984	1,981	19.4	18.5-20.3	1,126	26.5	24.9-28.2	855	14.5	13.5-15.5	541	5.5	5.0-6.0	296	7.6	6.7-8.6	245	4.1	3.6-4.7
1985	2,081	20.2	19.4-21.1	1,158	27.2	25.6-28.9	923	15.5	14.5-16.5	520	5.2	4.8-5.7	279	7.2	6.3-8.2	241	4.0	3.5-4.5
1986	1,959	18.9	18.1-19.8	1,088	25.3	23.8-27.0	871	14.5	13.5-15.5	501	5.0	4.5-5.4	273	6.8	6.0-7.7	228	3.8	3.3-4.3
1987	2,027	19.4	18.6-20.3	1,105	25.6	24.0-27.2	922	15.3	14.3-16.3	505	5.0	4.5-5.4	263	6.6	5.8-7.5	242	3.9	3.4-4.4
1988	1,910	18.1	17.3-18.9	1,079	24.7	23.2-26.3	831	13.6	12.6-14.5	461	4.5	4.1-4.9	249	6.1	5.3-7.0	212	3.4	3.0-3.9
1989	1,974	18.5	17.7-19.3	1,075	23.9	22.5-25.5	899	14.6	13.6-15.6	457	4.4	4.0-4.8	255	6.2	5.4-7.0	202	3.2	2.8-3.7
1990	1,969	18.2	17.4-19.1	1,077	23.7	22.2-25.2	892	14.1	13.2-15.1	392	3.7	3.3-4.1	195	4.5	3.9-5.2	197	3.1	2.7-3.6
1991	1,853	17.1	16.3-17.9	1,010	22.1	20.7-23.5	843	13.4	12.5-14.4	464	4.3	3.9-4.7	240	5.6	4.9-6.4	224	3.5	3.0-4.0
1992	1,867	16.9	16.2-17.7	1,077	23.5	22.0-25.0	790	12.3	11.5-13.2	436	4.0	3.6-4.4	244	5.5	4.8-6.3	192	2.9	2.5-3.4
1993	1,849	16.6	15.8-17.3	1,022	21.8	20.4-23.2	827	12.8	11.9-13.7	415	3.7	3.4-4.1	210	4.7	4.1-5.4	205	3.0	2.6-3.5
1994	1,891	16.9	16.1-17.7	1,026	21.9	20.5-23.3	865	13.4	12.5-14.3	423	3.8	3.5-4.2	229	5.1	4.4-5.8	194	2.9	2.5-3.3
1995	1,862	16.6	15.8-17.4	1,014	21.5	20.2-22.9	848	13.1	12.2-14.0	457	4.1	3.7-4.5	246	5.4	4.8-6.2	211	3.1	2.7-3.6
1996	1,912	16.9	16.2-17.7	1,094	22.7	21.3-24.1	818	12.5	11.6-13.4	433	3.8	3.5-4.2	232	5.1	4.4-5.8	201	3.0	2.6-3.4
1997	1,841	16.1	15.4-16.8	1,051	21.6	20.3-23.0	790	12.0	11.1-12.8	363	3.2	2.8-3.5	209	4.6	3.9-5.2	154	2.2	1.9-2.6
1998	1,986	17.2	16.4-17.9	1,069	21.5	20.2-22.8	917	13.8	12.9-14.7	406	3.5	3.2-3.9	214	4.5	3.9-5.1	192	2.8	2.4-3.2
1999	2,079	17.8	17.0-18.6	1,159	23.3	21.9-24.7	920	13.8	12.9-14.7	426	3.6	3.3-4.0	231	4.9	4.2-5.5	195	2.8	2.4-3.2
2000	2,015	17.1	16.3-17.8	1,127	22.1	20.8-23.5	888	13.2	12.3-14.1	418	3.5	3.2-3.9	229	4.6	4.0-5.3	189	2.7	2.3-3.1
2001	1,952	16.3	15.6-17.1	1,078	21.0	19.7-22.3	874	12.8	12.0-13.7	389	3.2	2.9-3.6	203	4.1	3.6-4.8	186	2.6	2.2-3.0
2002	1,936	16.0	15.3-16.8	1,069	20.3	19.1-21.6	867	12.5	11.7-13.4	444	3.6	3.3-4.0	233	4.6	4.0-5.3	211	2.9	2.5-3.4
2003	1,855	15.1	14.5-15.9	1,009	18.9	17.7-20.1	846	12.2	11.3-13.0	369	3.0	2.7-3.3	211	4.1	3.6-4.7	158	2.2	1.8-2.5
2004	1,877	15.1	14.5-15.9	1,023	18.8	17.6-20.0	854	12.3	11.5-13.2	398	3.2	2.9-3.5	213	4.0	3.5-4.6	185	2.5	2.1-2.9
2005	1,728	13.7	13.1-14.4	953	17.4	16.3-18.5	775	11.0	10.2-11.8	377	3.0	2.7-3.3	208	3.9	3.4-4.5	169	2.2	1.9-2.6
2006	1,612	12.8	12.2-13.5	879	15.7	14.6-16.7	733	10.5	9.7-11.3	354	2.8	2.5-3.1	182	3.4	2.9-4.0	172	2.3	2.0-2.7
2007	1,669	13.0	12.4-13.7	912	15.9	14.9-17.0	757	10.8	10.0-11.6	347	2.7	2.4-3.0	194	3.5	3.0-4.0	153	2.0	1.7-2.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Rectum & rectosigmoid

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	1,677	12.9	12.3-13.6	945	16.5	15.4-17.6	732	10.1	9.4-10.9	374	2.8	2.5-3.1	198	3.5	3.0-4.1	176	2.2	1.9-2.6
2009	1,634	12.5	11.9-13.1	918	15.7	14.7-16.8	716	9.9	9.2-10.7	353	2.6	2.4-2.9	198	3.5	3.0-4.0	155	2.0	1.7-2.4
2010	1,650	12.3	11.7-12.9	918	15.0	14.1-16.1	732	10.1	9.3-10.9	365	2.7	2.4-3.0	203	3.5	3.0-4.0	162	2.1	1.8-2.5
2011	1,618	12.1	11.5-12.8	918	15.0	14.0-16.0	700	9.7	9.0-10.5	348	2.5	2.2-2.8	201	3.3	2.9-3.8	147	1.9	1.6-2.2
2012	1,512	11.2	10.6-11.8	886	14.3	13.4-15.3	626	8.6	8.0-9.4	325	2.3	2.1-2.6	191	3.2	2.8-3.7	134	1.6	1.4-2.0
2013	1,640	12.0	11.4-12.6	934	14.8	13.8-15.8	706	9.7	9.0-10.5	350	2.5	2.2-2.8	206	3.4	2.9-3.9	144	1.8	1.5-2.1
2014	1,595	11.5	10.9-12.1	932	14.5	13.6-15.5	663	8.9	8.2-9.6	363	2.5	2.3-2.8	208	3.3	2.8-3.7	155	1.9	1.6-2.3
2015	1,592	11.3	10.7-11.9	918	13.9	13.0-14.9	674	9.1	8.4-9.8	351	2.4	2.2-2.7	205	3.2	2.8-3.7	146	1.8	1.5-2.2
2016	1,564	11.1	10.5-11.7	926	14.2	13.3-15.2	638	8.5	7.8-9.2	374	2.6	2.3-2.9	212	3.3	2.9-3.8	162	2.0	1.7-2.3
2017	1,591	11.3	10.7-11.9	920	14.0	13.1-15.0	671	9.0	8.3-9.7	386	2.6	2.3-2.9	211	3.2	2.8-3.7	175	2.1	1.8-2.5
2018	1,592	11.1	10.6-11.7	926	13.8	12.9-14.7	666	8.9	8.2-9.6	383	2.6	2.3-2.9	221	3.3	2.9-3.8	162	2.0	1.7-2.3
2019	1,611	11.4	10.8-12.0	939	14.1	13.2-15.1	672	9.0	8.3-9.8	381	2.5	2.2-2.8	220	3.2	2.8-3.7	161	1.9	1.6-2.2
2020	1,468	10.2	9.7-10.8	869	12.8	12.0-13.8	599	8.0	7.3-8.7	419	2.8	2.5-3.1	241	3.5	3.1-4.0	178	2.1	1.8-2.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Liver / intrahepatic bile duct

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	257	2.7	2.4-3.1	157	3.9	3.3-4.6	100	1.9	1.5-2.3	199	2.1	1.8-2.4	127	3.2	2.7-3.9	72	1.4	1.1-1.7
1977	296	3.0	2.7-3.4	198	4.7	4.0-5.4	98	1.8	1.4-2.1	253	2.6	2.3-2.9	154	3.6	3.0-4.2	99	1.8	1.5-2.2
1978	293	3.0	2.7-3.4	173	4.2	3.6-4.9	120	2.2	1.8-2.6	250	2.6	2.3-2.9	145	3.6	3.0-4.2	105	1.9	1.6-2.3
1979	273	2.8	2.5-3.2	145	3.4	2.9-4.1	128	2.3	2.0-2.8	212	2.2	1.9-2.5	117	2.8	2.3-3.4	95	1.7	1.4-2.1
1980	275	2.8	2.5-3.2	166	4.2	3.6-5.0	109	1.9	1.5-2.3	235	2.4	2.1-2.7	128	3.2	2.7-3.9	107	1.9	1.5-2.3
1981	287	2.9	2.6-3.3	169	4.2	3.5-4.9	118	2.1	1.7-2.5	264	2.7	2.4-3.0	147	3.7	3.1-4.4	117	2.0	1.7-2.4
1982	291	2.9	2.6-3.3	185	4.5	3.8-5.2	106	1.8	1.5-2.2	245	2.5	2.2-2.8	160	4.1	3.4-4.8	85	1.5	1.2-1.8
1983	300	3.0	2.6-3.3	206	4.8	4.1-5.5	94	1.7	1.3-2.0	258	2.5	2.2-2.9	175	4.2	3.6-4.9	83	1.4	1.1-1.7
1984	288	2.8	2.5-3.2	167	4.0	3.3-4.6	121	2.1	1.7-2.5	257	2.5	2.2-2.9	153	3.7	3.1-4.3	104	1.8	1.4-2.2
1985	275	2.7	2.4-3.1	171	4.2	3.5-4.9	104	1.8	1.4-2.2	250	2.5	2.2-2.8	154	3.7	3.1-4.4	96	1.7	1.3-2.0
1986	291	2.8	2.5-3.2	187	4.3	3.7-5.0	104	1.8	1.4-2.1	271	2.6	2.3-2.9	168	3.9	3.3-4.6	103	1.7	1.4-2.1
1987	306	2.9	2.6-3.3	191	4.3	3.7-5.0	115	1.9	1.6-2.3	306	3.0	2.7-3.3	185	4.4	3.8-5.1	121	2.0	1.7-2.4
1988	364	3.5	3.1-3.8	236	5.4	4.7-6.1	128	2.1	1.7-2.5	318	3.0	2.7-3.4	209	4.8	4.1-5.5	109	1.8	1.5-2.1
1989	353	3.3	3.0-3.7	228	5.0	4.3-5.7	125	2.0	1.7-2.4	285	2.7	2.4-3.0	175	4.0	3.4-4.6	110	1.8	1.4-2.1
1990	393	3.7	3.3-4.1	254	5.7	5.0-6.4	139	2.2	1.8-2.6	346	3.3	2.9-3.6	219	5.1	4.4-5.8	127	2.0	1.7-2.4
1991	383	3.5	3.2-3.9	239	5.0	4.4-5.7	144	2.3	1.9-2.7	346	3.1	2.8-3.5	214	4.6	4.0-5.2	132	2.1	1.7-2.4
1992	422	3.9	3.5-4.3	248	5.6	4.9-6.4	174	2.7	2.3-3.2	358	3.3	2.9-3.6	215	4.8	4.2-5.5	143	2.2	1.9-2.6
1993	456	4.1	3.7-4.5	280	6.1	5.4-6.9	176	2.7	2.3-3.1	434	3.9	3.6-4.3	256	5.7	5.0-6.5	178	2.7	2.3-3.1
1994	425	3.8	3.4-4.2	282	6.1	5.4-6.9	143	2.2	1.8-2.6	401	3.6	3.2-4.0	237	5.3	4.6-6.0	164	2.5	2.1-2.9
1995	504	4.5	4.1-4.9	327	6.9	6.2-7.7	177	2.7	2.3-3.1	443	3.9	3.6-4.3	284	6.2	5.5-6.9	159	2.4	2.0-2.8
1996	454	4.0	3.7-4.4	281	5.9	5.2-6.7	173	2.6	2.2-3.0	429	3.8	3.4-4.2	257	5.5	4.8-6.2	172	2.5	2.2-2.9
1997	535	4.7	4.3-5.1	349	7.1	6.4-7.9	186	2.8	2.4-3.3	431	3.8	3.4-4.2	272	5.7	5.0-6.4	159	2.3	2.0-2.7
1998	570	4.9	4.5-5.4	386	7.7	7.0-8.6	184	2.7	2.3-3.1	428	3.7	3.4-4.1	258	5.2	4.6-5.9	170	2.5	2.1-2.9
1999	536	4.6	4.2-5.0	355	7.1	6.4-7.9	181	2.7	2.3-3.1	447	3.8	3.5-4.2	269	5.4	4.8-6.1	178	2.5	2.2-3.0
2000	540	4.6	4.2-5.0	346	6.8	6.1-7.5	194	2.9	2.5-3.3	457	3.8	3.5-4.2	278	5.6	4.9-6.2	179	2.5	2.2-3.0
2001	579	4.9	4.5-5.3	410	7.9	7.2-8.7	169	2.5	2.1-2.9	461	3.8	3.5-4.2	297	5.9	5.2-6.6	164	2.3	2.0-2.7
2002	609	5.0	4.6-5.5	434	8.2	7.5-9.0	175	2.5	2.1-2.9	517	4.2	3.9-4.6	338	6.6	5.9-7.3	179	2.5	2.1-2.9
2003	622	5.1	4.7-5.5	402	7.4	6.6-8.1	220	3.2	2.8-3.6	467	3.8	3.5-4.2	306	5.8	5.1-6.5	161	2.3	1.9-2.6
2004	675	5.5	5.0-5.9	485	8.8	8.0-9.6	190	2.7	2.3-3.2	540	4.3	4.0-4.7	337	6.2	5.6-6.9	203	2.8	2.4-3.2
2005	723	5.8	5.4-6.2	508	9.0	8.2-9.8	215	3.1	2.7-3.6	558	4.4	4.1-4.8	370	6.7	6.1-7.5	188	2.6	2.2-3.0
2006	725	5.7	5.3-6.1	530	9.2	8.5-10.1	195	2.8	2.4-3.2	578	4.5	4.2-4.9	349	6.3	5.7-7.0	229	3.2	2.8-3.6
2007	824	6.4	6.0-6.8	587	10.0	9.2-10.9	237	3.3	2.9-3.8	571	4.4	4.0-4.8	382	6.7	6.1-7.5	189	2.5	2.2-2.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Liver / intrahepatic bile duct

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	869	6.6	6.2-7.1	620	10.3	9.5-11.2	249	3.5	3.0-3.9	670	5.1	4.7-5.5	441	7.5	6.8-8.3	229	3.0	2.6-3.5
2009	879	6.5	6.1-7.0	628	10.2	9.4-11.0	251	3.4	3.0-3.9	652	4.9	4.5-5.3	435	7.3	6.6-8.1	217	2.8	2.5-3.2
2010	827	6.0	5.6-6.5	630	10.0	9.3-10.9	197	2.7	2.3-3.1	667	4.9	4.5-5.3	434	7.1	6.5-7.8	233	3.1	2.7-3.5
2011	914	6.5	6.1-7.0	661	10.1	9.3-10.9	253	3.4	3.0-3.9	720	5.2	4.8-5.6	460	7.3	6.6-8.0	260	3.3	2.9-3.8
2012	966	6.8	6.3-7.2	715	10.8	10.0-11.6	251	3.3	2.9-3.8	776	5.5	5.1-5.9	515	8.0	7.3-8.8	261	3.4	3.0-3.8
2013	1,066	7.4	6.9-7.8	748	11.2	10.4-12.0	318	4.1	3.7-4.6	795	5.5	5.1-5.9	543	8.4	7.7-9.2	252	3.1	2.8-3.6
2014	1,150	7.8	7.4-8.3	844	12.3	11.5-13.2	306	3.9	3.5-4.4	742	5.1	4.7-5.5	511	7.6	7.0-8.3	231	3.0	2.6-3.4
2015	1,096	7.3	6.9-7.8	783	11.2	10.4-12.1	313	4.0	3.5-4.5	796	5.4	5.0-5.8	525	7.8	7.2-8.6	271	3.3	2.9-3.7
2016	1,234	8.1	7.7-8.6	905	12.7	11.9-13.6	329	4.1	3.7-4.6	812	5.3	5.0-5.7	545	7.9	7.2-8.6	267	3.2	2.8-3.6
2017	1,212	7.8	7.4-8.3	872	12.2	11.4-13.1	340	4.1	3.7-4.6	772	5.1	4.8-5.5	537	7.8	7.1-8.5	235	2.9	2.6-3.4
2018	1,092	7.0	6.6-7.5	774	10.7	10.0-11.5	318	3.9	3.5-4.4	808	5.2	4.8-5.6	537	7.6	6.9-8.3	271	3.2	2.8-3.6
2019	1,190	7.6	7.1-8.0	826	11.3	10.5-12.1	364	4.4	4.0-4.9	760	4.8	4.5-5.2	516	7.2	6.6-7.8	244	2.9	2.5-3.3
2020	1,127	7.1	6.7-7.5	765	10.4	9.7-11.2	362	4.3	3.9-4.8	776	4.9	4.5-5.2	501	6.9	6.3-7.5	275	3.2	2.8-3.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Pancreas

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,114	12.0	11.3-12.8	592	15.2	13.9-16.5	522	9.8	9.0-10.7	1,066	11.6	10.9-12.3	544	14.1	12.9-15.4	522	9.8	9.0-10.7
1977	1,174	12.7	12.0-13.5	623	16.5	15.2-18.0	551	10.2	9.3-11.0	1,143	12.4	11.7-13.1	602	15.7	14.4-17.1	541	10.1	9.2-11.0
1978	1,191	12.6	11.9-13.4	618	15.6	14.4-17.0	573	10.4	9.6-11.3	1,138	12.0	11.3-12.7	593	15.1	13.9-16.5	545	9.9	9.0-10.7
1979	1,297	13.6	12.9-14.4	689	17.6	16.2-19.0	608	10.9	10.0-11.8	1,232	12.9	12.2-13.7	647	16.6	15.3-18.0	585	10.4	9.5-11.2
1980	1,225	12.6	11.9-13.4	605	15.0	13.7-16.3	620	10.9	10.1-11.8	1,147	11.9	11.2-12.6	581	14.7	13.4-16.0	566	9.9	9.1-10.8
1981	1,201	12.3	11.6-13.1	604	15.0	13.8-16.3	597	10.5	9.6-11.4	1,160	11.9	11.2-12.6	583	14.6	13.3-15.9	577	10.1	9.3-11.0
1982	1,207	12.1	11.4-12.8	593	14.6	13.4-15.9	614	10.5	9.6-11.3	1,105	11.1	10.5-11.8	567	14.0	12.8-15.3	538	9.2	8.4-10.0
1983	1,167	11.7	11.0-12.4	582	14.1	12.9-15.4	585	10.0	9.2-10.8	1,200	12.0	11.3-12.7	590	14.1	13.0-15.4	610	10.3	9.5-11.2
1984	1,205	11.9	11.3-12.6	591	14.6	13.4-15.9	614	10.3	9.5-11.1	1,160	11.5	10.8-12.2	562	13.9	12.7-15.1	598	10.0	9.2-10.8
1985	1,303	12.7	12.0-13.4	615	14.5	13.3-15.7	688	11.4	10.6-12.3	1,187	11.7	11.0-12.4	558	13.6	12.4-14.8	629	10.5	9.6-11.3
1986	1,239	12.1	11.5-12.8	608	14.3	13.1-15.5	631	10.4	9.6-11.3	1,204	11.7	11.1-12.4	597	13.9	12.7-15.1	607	10.0	9.2-10.9
1987	1,275	12.2	11.6-12.9	593	14.2	13.1-15.5	682	11.0	10.2-11.9	1,205	11.7	11.0-12.4	581	14.0	12.8-15.2	624	10.1	9.3-10.9
1988	1,322	12.7	12.0-13.4	648	15.3	14.1-16.6	674	10.8	10.0-11.7	1,251	12.0	11.3-12.6	609	14.5	13.3-15.8	642	10.2	9.4-11.0
1989	1,228	11.5	10.9-12.2	602	13.9	12.8-15.1	626	10.0	9.2-10.8	1,147	10.8	10.2-11.5	574	13.5	12.3-14.7	573	9.1	8.4-9.9
1990	1,334	12.5	11.9-13.2	634	14.6	13.4-15.8	700	11.1	10.2-11.9	1,222	11.5	10.8-12.1	569	13.1	12.0-14.3	653	10.3	9.5-11.1
1991	1,369	12.7	12.0-13.4	662	14.8	13.6-16.0	707	11.0	10.2-11.9	1,209	11.2	10.6-11.9	572	13.1	12.0-14.2	637	9.9	9.1-10.7
1992	1,323	12.2	11.5-12.8	630	14.2	13.1-15.4	693	10.7	9.9-11.6	1,224	11.2	10.6-11.9	568	13.0	11.9-14.1	656	10.0	9.3-10.8
1993	1,338	12.1	11.5-12.8	653	14.3	13.2-15.4	685	10.4	9.7-11.3	1,282	11.6	11.0-12.2	611	13.4	12.4-14.6	671	10.2	9.4-11.0
1994	1,443	12.9	12.2-13.6	719	15.5	14.4-16.7	724	10.8	10.1-11.7	1,334	12.0	11.3-12.6	665	14.6	13.5-15.8	669	9.9	9.2-10.7
1995	1,425	12.6	12.0-13.3	655	13.8	12.7-14.9	770	11.5	10.7-12.4	1,324	11.7	11.1-12.4	615	13.0	11.9-14.0	709	10.5	9.7-11.3
1996	1,371	12.1	11.5-12.8	694	14.8	13.7-16.0	677	10.2	9.4-11.0	1,278	11.3	10.7-11.9	614	13.3	12.3-14.4	664	9.9	9.1-10.7
1997	1,424	12.4	11.8-13.1	684	14.2	13.2-15.4	740	10.9	10.1-11.8	1,358	11.8	11.2-12.5	675	14.4	13.3-15.5	683	9.9	9.2-10.7
1998	1,512	13.0	12.4-13.7	720	14.8	13.7-15.9	792	11.7	10.9-12.5	1,345	11.6	11.0-12.2	653	13.7	12.6-14.8	692	10.0	9.3-10.8
1999	1,489	12.7	12.1-13.4	698	14.1	13.1-15.2	791	11.5	10.7-12.3	1,395	11.9	11.2-12.5	673	13.7	12.7-14.8	722	10.4	9.7-11.2
2000	1,492	12.5	11.9-13.2	697	14.0	12.9-15.1	795	11.4	10.6-12.3	1,369	11.5	10.9-12.1	639	13.1	12.1-14.1	730	10.3	9.6-11.1
2001	1,552	12.9	12.3-13.6	711	14.0	13.0-15.1	841	11.9	11.1-12.8	1,376	11.4	10.8-12.0	628	12.6	11.6-13.6	748	10.5	9.7-11.3
2002	1,606	13.2	12.6-13.9	757	14.7	13.6-15.8	849	12.0	11.2-12.9	1,411	11.6	11.0-12.2	680	13.2	12.2-14.3	731	10.2	9.4-10.9
2003	1,614	13.1	12.4-13.7	786	15.0	14.0-16.1	828	11.6	10.8-12.4	1,392	11.2	10.7-11.8	640	12.4	11.5-13.4	752	10.3	9.6-11.1
2004	1,602	12.8	12.2-13.5	772	14.5	13.5-15.6	830	11.5	10.7-12.4	1,418	11.3	10.7-11.9	707	13.4	12.4-14.4	711	9.8	9.1-10.6
2005	1,659	13.2	12.5-13.8	811	14.9	13.9-16.0	848	11.5	10.8-12.4	1,456	11.5	10.9-12.1	687	12.8	11.9-13.8	769	10.3	9.6-11.1
2006	1,680	13.2	12.6-13.9	802	14.7	13.7-15.8	878	12.0	11.2-12.9	1,442	11.3	10.7-11.9	703	12.9	11.9-13.9	739	9.9	9.2-10.6
2007	1,735	13.4	12.8-14.1	840	15.1	14.0-16.1	895	12.0	11.2-12.9	1,437	11.1	10.5-11.6	703	12.7	11.8-13.7	734	9.8	9.1-10.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Pancreas

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	1,827	14.0	13.3-14.6	895	15.8	14.8-16.9	932	12.4	11.6-13.3	1,506	11.5	10.9-12.1	718	12.8	11.9-13.8	788	10.3	9.6-11.1
2009	1,825	13.7	13.1-14.4	955	16.4	15.4-17.5	870	11.5	10.7-12.3	1,534	11.4	10.8-12.0	764	13.3	12.3-14.3	770	10.0	9.3-10.8
2010	1,860	13.8	13.2-14.5	889	15.1	14.1-16.2	971	12.7	11.9-13.6	1,552	11.3	10.8-11.9	759	12.8	11.9-13.8	793	10.1	9.4-10.9
2011	1,938	14.1	13.5-14.8	981	16.2	15.2-17.3	957	12.3	11.5-13.2	1,531	11.1	10.5-11.7	739	12.4	11.5-13.4	792	10.0	9.3-10.7
2012	1,951	14.0	13.4-14.7	995	16.3	15.3-17.4	956	12.2	11.4-13.0	1,620	11.6	11.1-12.2	839	13.8	12.9-14.8	781	9.8	9.1-10.6
2013	1,996	14.1	13.4-14.7	1,001	16.0	15.0-17.1	995	12.4	11.6-13.2	1,646	11.6	11.0-12.2	819	13.3	12.3-14.2	827	10.2	9.5-10.9
2014	2,122	14.8	14.2-15.5	1,084	16.8	15.8-17.9	1,038	13.1	12.3-13.9	1,645	11.4	10.8-11.9	837	13.2	12.3-14.2	808	9.8	9.1-10.5
2015	2,097	14.3	13.7-15.0	1,033	15.7	14.7-16.7	1,064	13.1	12.3-14.0	1,595	10.9	10.3-11.4	805	12.4	11.6-13.4	790	9.4	8.8-10.1
2016	2,214	15.2	14.6-15.9	1,136	17.3	16.3-18.4	1,078	13.5	12.7-14.4	1,695	11.4	10.9-12.0	836	12.7	11.8-13.6	859	10.4	9.7-11.1
2017	2,288	15.3	14.6-15.9	1,193	17.7	16.7-18.8	1,095	13.3	12.5-14.1	1,723	11.4	10.9-12.0	872	13.1	12.2-14.0	851	10.1	9.4-10.8
2018	2,271	15.0	14.3-15.6	1,159	16.9	15.9-17.9	1,112	13.3	12.5-14.1	1,740	11.3	10.7-11.8	886	12.9	12.1-13.8	854	9.9	9.2-10.6
2019	2,389	15.5	14.9-16.2	1,208	17.5	16.5-18.6	1,181	13.8	13.0-14.7	1,788	11.5	10.9-12.0	930	13.4	12.5-14.3	858	9.8	9.2-10.5
2020	2,291	14.6	14.0-15.3	1,204	17.0	16.0-18.0	1,087	12.6	11.8-13.4	1,767	11.2	10.7-11.8	889	12.7	11.9-13.6	878	10.0	9.3-10.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Larynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	547	5.5	5.0-6.0	468	10.8	9.8-11.9	79	1.4	1.1-1.8	191	2.0	1.7-2.3	158	4.0	3.3-4.7	33	0.6	0.4-0.9
1977	538	5.4	4.9-5.8	441	10.2	9.2-11.3	97	1.7	1.4-2.1	181	1.8	1.6-2.1	151	3.7	3.1-4.3	30	0.5	0.4-0.8
1978	579	5.7	5.2-6.2	482	11.0	10.0-12.1	97	1.7	1.4-2.1	204	2.1	1.8-2.4	161	4.0	3.3-4.7	43	0.8	0.6-1.0
1979	619	6.0	5.5-6.5	486	10.8	9.8-11.9	133	2.3	1.9-2.8	219	2.2	1.9-2.5	180	4.3	3.7-5.0	39	0.7	0.5-0.9
1980	622	6.0	5.5-6.5	512	11.5	10.5-12.6	110	2.0	1.6-2.4	198	1.9	1.7-2.2	159	3.8	3.2-4.4	39	0.7	0.5-1.0
1981	672	6.5	6.0-7.0	550	12.2	11.2-13.3	122	2.1	1.7-2.5	203	2.0	1.7-2.3	171	4.1	3.5-4.8	32	0.5	0.4-0.8
1982	606	5.8	5.4-6.3	494	11.0	10.0-12.0	112	2.0	1.6-2.4	211	2.0	1.7-2.3	168	3.8	3.3-4.5	43	0.7	0.5-1.0
1983	596	5.7	5.3-6.2	480	10.6	9.6-11.6	116	2.1	1.7-2.5	201	1.9	1.7-2.2	154	3.6	3.0-4.2	47	0.8	0.6-1.1
1984	610	5.8	5.3-6.3	477	10.4	9.5-11.4	133	2.3	1.9-2.8	175	1.7	1.4-1.9	137	3.2	2.7-3.8	38	0.6	0.4-0.9
1985	633	6.0	5.5-6.5	497	10.7	9.7-11.7	136	2.4	2.0-2.8	194	1.8	1.5-2.1	151	3.3	2.8-4.0	43	0.7	0.5-1.0
1986	613	5.8	5.3-6.3	476	10.5	9.5-11.5	137	2.4	2.0-2.8	184	1.7	1.5-2.0	144	3.3	2.7-3.9	40	0.6	0.5-0.9
1987	642	6.0	5.6-6.5	514	11.2	10.2-12.3	128	2.1	1.8-2.6	190	1.8	1.5-2.1	148	3.4	2.8-4.0	42	0.7	0.5-0.9
1988	573	5.3	4.9-5.8	434	9.2	8.4-10.2	139	2.4	2.0-2.8	192	1.8	1.6-2.1	156	3.5	3.0-4.2	36	0.6	0.4-0.8
1989	649	6.1	5.6-6.5	498	10.7	9.8-11.8	151	2.6	2.2-3.0	183	1.7	1.5-2.0	128	2.8	2.3-3.3	55	0.9	0.7-1.2
1990	643	5.9	5.4-6.3	485	10.0	9.1-11.0	158	2.7	2.3-3.1	171	1.6	1.4-1.9	137	3.1	2.6-3.7	34	0.5	0.4-0.8
1991	583	5.3	4.9-5.8	467	9.7	8.9-10.7	116	1.9	1.5-2.3	178	1.6	1.4-1.9	141	3.0	2.5-3.5	37	0.6	0.4-0.8
1992	609	5.5	5.1-6.0	461	9.5	8.7-10.5	148	2.5	2.1-2.9	223	2.0	1.7-2.3	170	3.7	3.2-4.4	53	0.8	0.6-1.1
1993	587	5.3	4.9-5.7	433	9.0	8.2-9.9	154	2.5	2.1-2.9	184	1.7	1.4-1.9	145	3.1	2.6-3.7	39	0.6	0.4-0.8
1994	569	5.1	4.7-5.5	441	9.1	8.2-10.0	128	2.1	1.7-2.5	188	1.7	1.4-1.9	142	3.1	2.6-3.7	46	0.7	0.5-0.9
1995	583	5.2	4.8-5.7	444	9.0	8.2-9.9	139	2.3	1.9-2.7	185	1.6	1.4-1.9	143	3.0	2.5-3.6	42	0.6	0.5-0.9
1996	524	4.7	4.3-5.1	407	8.2	7.4-9.0	117	1.9	1.6-2.3	183	1.6	1.4-1.9	138	2.9	2.4-3.4	45	0.7	0.5-0.9
1997	540	4.8	4.4-5.2	419	8.4	7.6-9.2	121	2.0	1.6-2.4	154	1.3	1.1-1.6	113	2.4	1.9-2.9	41	0.6	0.5-0.9
1998	524	4.6	4.2-5.0	380	7.5	6.8-8.3	144	2.3	1.9-2.7	169	1.5	1.3-1.7	129	2.6	2.2-3.1	40	0.6	0.4-0.8
1999	563	4.9	4.5-5.3	447	8.6	7.8-9.4	116	1.8	1.5-2.2	175	1.5	1.3-1.7	135	2.7	2.2-3.2	40	0.6	0.4-0.8
2000	537	4.6	4.2-5.0	419	8.1	7.3-8.9	118	1.8	1.5-2.2	153	1.3	1.1-1.5	120	2.4	2.0-2.9	33	0.5	0.3-0.7
2001	514	4.3	4.0-4.7	402	7.5	6.8-8.3	112	1.7	1.4-2.1	149	1.2	1.0-1.5	125	2.5	2.1-3.0	24	0.3	0.2-0.5
2002	505	4.2	3.8-4.6	393	7.4	6.7-8.2	112	1.7	1.4-2.1	171	1.4	1.2-1.6	134	2.6	2.2-3.1	37	0.5	0.4-0.8
2003	495	4.1	3.7-4.4	383	7.0	6.3-7.7	112	1.7	1.4-2.0	150	1.2	1.0-1.4	111	2.1	1.7-2.5	39	0.6	0.4-0.8
2004	495	4.0	3.6-4.4	401	7.1	6.4-7.9	94	1.4	1.1-1.7	144	1.2	1.0-1.4	108	2.1	1.7-2.5	36	0.5	0.4-0.7
2005	546	4.3	4.0-4.7	428	7.6	6.9-8.3	118	1.7	1.4-2.1	164	1.3	1.1-1.6	122	2.3	1.9-2.7	42	0.6	0.4-0.8
2006	540	4.3	3.9-4.7	409	7.2	6.6-8.0	131	1.9	1.6-2.3	150	1.2	1.0-1.4	113	2.1	1.7-2.5	37	0.5	0.4-0.7
2007	491	3.8	3.5-4.1	387	6.7	6.1-7.4	104	1.4	1.2-1.8	138	1.1	0.9-1.2	114	2.0	1.6-2.4	24	0.3	0.2-0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Larynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	543	4.2	3.8-4.5	429	7.3	6.6-8.0	114	1.6	1.3-2.0	132	1.0	0.8-1.2	102	1.8	1.4-2.1	30	0.4	0.3-0.6
2009	493	3.7	3.4-4.1	381	6.4	5.8-7.1	112	1.6	1.3-1.9	149	1.1	0.9-1.3	109	1.9	1.6-2.3	40	0.5	0.4-0.7
2010	486	3.6	3.3-4.0	381	6.2	5.6-6.9	105	1.5	1.2-1.8	159	1.2	1.0-1.4	121	2.1	1.7-2.5	38	0.5	0.3-0.7
2011	490	3.5	3.2-3.8	401	6.2	5.6-6.9	89	1.2	1.0-1.5	144	1.0	0.9-1.2	106	1.7	1.4-2.1	38	0.5	0.3-0.7
2012	495	3.5	3.2-3.9	384	6.1	5.5-6.8	111	1.4	1.2-1.8	140	1.0	0.8-1.2	112	1.8	1.5-2.2	28	0.4	0.2-0.5
2013	511	3.5	3.2-3.9	401	6.1	5.5-6.7	110	1.5	1.2-1.8	169	1.1	1.0-1.3	139	2.1	1.8-2.5	30	0.4	0.3-0.6
2014	491	3.4	3.1-3.7	388	5.8	5.2-6.4	103	1.3	1.1-1.6	156	1.1	0.9-1.3	125	1.9	1.6-2.3	31	0.4	0.3-0.6
2015	513	3.5	3.2-3.8	411	6.1	5.5-6.7	102	1.3	1.1-1.6	132	0.9	0.7-1.0	103	1.6	1.3-1.9	29	0.3	0.2-0.5
2016	473	3.2	2.9-3.5	377	5.6	5.0-6.2	96	1.2	1.0-1.5	141	1.0	0.8-1.2	112	1.8	1.4-2.1	29	0.4	0.2-0.5
2017	488	3.2	3.0-3.6	372	5.4	4.8-6.0	116	1.5	1.2-1.8	123	0.8	0.6-0.9	96	1.4	1.1-1.7	27	0.3	0.2-0.5
2018	465	3.0	2.7-3.3	364	5.1	4.6-5.6	101	1.3	1.0-1.5	138	0.9	0.7-1.0	104	1.5	1.2-1.8	34	0.4	0.3-0.6
2019	463	2.9	2.7-3.2	374	5.1	4.6-5.6	89	1.1	0.9-1.4	117	0.7	0.6-0.9	92	1.2	1.0-1.5	25	0.3	0.2-0.4
2020	423	2.7	2.4-3.0	336	4.6	4.1-5.1	87	1.1	0.9-1.3	110	0.7	0.6-0.8	90	1.3	1.0-1.6	20	0.2	0.1-0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Lung and bronchus

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	5,605	58.0	56.5-59.6	4,064	99.9	96.7-103.2	1,541	28.5	27.1-30.0	4,460	46.5	45.2-48.0	3,295	82.7	79.8-85.8	1,165	21.6	20.4-22.9
1977	5,681	58.0	56.4-59.5	4,073	99.0	95.8-102.2	1,608	29.4	28.0-30.9	4,748	48.9	47.5-50.3	3,476	86.1	83.1-89.1	1,272	23.2	22.0-24.6
1978	5,806	58.7	57.2-60.2	4,171	100.6	97.4-103.8	1,635	29.6	28.1-31.1	4,778	48.4	47.0-49.8	3,460	84.7	81.7-87.7	1,318	23.6	22.4-25.0
1979	5,962	59.8	58.3-61.4	4,148	99.7	96.5-102.9	1,814	32.5	31.0-34.1	4,750	47.8	46.4-49.2	3,366	82.0	79.1-84.9	1,384	24.6	23.3-25.9
1980	6,356	62.8	61.2-64.4	4,469	105.2	102.0-108.5	1,887	33.2	31.7-34.8	5,161	51.3	49.8-52.7	3,683	88.2	85.2-91.2	1,478	25.9	24.6-27.3
1981	6,394	62.9	61.3-64.5	4,387	103.5	100.3-106.8	2,007	35.4	33.8-37.0	5,213	51.4	50.0-52.8	3,685	88.2	85.3-91.3	1,528	26.7	25.4-28.1
1982	6,629	64.5	62.9-66.1	4,476	104.5	101.3-107.7	2,153	37.4	35.8-39.0	5,360	52.6	51.2-54.1	3,637	86.7	83.7-89.7	1,723	29.9	28.5-31.3
1983	6,914	66.7	65.1-68.3	4,548	104.8	101.6-108.0	2,366	40.6	39.0-42.3	5,566	54.0	52.6-55.4	3,764	88.9	85.9-91.9	1,802	30.8	29.3-32.3
1984	6,845	65.7	64.2-67.3	4,457	101.7	98.6-104.9	2,388	41.1	39.4-42.8	5,541	53.3	51.8-54.7	3,671	84.8	82.0-87.8	1,870	31.8	30.4-33.3
1985	7,194	68.7	67.1-70.3	4,615	105.0	101.9-108.2	2,579	44.1	42.4-45.9	5,885	56.3	54.9-57.8	3,859	89.4	86.5-92.4	2,026	34.1	32.7-35.7
1986	7,161	68.2	66.6-69.8	4,547	103.7	100.6-106.9	2,614	44.4	42.7-46.2	5,939	56.5	55.1-58.0	3,823	88.5	85.6-91.4	2,116	35.7	34.1-37.3
1987	7,316	69.0	67.4-70.7	4,571	102.6	99.6-105.8	2,745	46.2	44.5-48.0	5,954	56.4	54.9-57.9	3,837	88.0	85.1-91.0	2,117	35.2	33.7-36.8
1988	7,573	70.8	69.2-72.4	4,656	104.4	101.3-107.5	2,917	48.5	46.7-50.3	6,102	57.2	55.7-58.6	3,836	87.1	84.2-90.0	2,266	37.5	35.9-39.1
1989	7,711	71.5	69.9-73.1	4,669	102.8	99.8-105.9	3,042	50.3	48.5-52.1	6,260	58.2	56.7-59.7	3,891	86.9	84.0-89.7	2,369	38.9	37.3-40.5
1990	7,738	71.0	69.4-72.6	4,709	103.2	100.1-106.3	3,029	49.2	47.5-51.0	6,352	58.4	56.9-59.9	3,862	85.7	82.9-88.5	2,490	40.1	38.6-41.8
1991	7,858	71.7	70.1-73.3	4,625	100.6	97.7-103.7	3,233	52.3	50.4-54.1	6,555	59.8	58.4-61.3	3,957	87.3	84.5-90.2	2,598	41.6	40.0-43.2
1992	7,945	71.7	70.2-73.4	4,623	99.3	96.4-102.3	3,322	53.1	51.3-55.0	6,588	59.6	58.2-61.1	3,862	84.8	82.0-87.6	2,726	42.9	41.3-44.6
1993	8,002	71.7	70.1-73.3	4,535	96.3	93.5-99.3	3,467	55.0	53.2-56.9	6,445	57.7	56.3-59.2	3,767	81.4	78.7-84.1	2,678	42.1	40.5-43.7
1994	8,278	73.6	72.0-75.2	4,651	98.3	95.4-101.2	3,627	57.4	55.5-59.3	6,628	58.9	57.5-60.3	3,931	84.4	81.7-87.2	2,697	41.8	40.3-43.5
1995	8,252	72.9	71.3-74.5	4,657	96.8	94.0-99.7	3,595	56.3	54.5-58.2	6,625	58.5	57.1-60.0	3,827	81.4	78.8-84.1	2,798	43.3	41.7-45.0
1996	8,370	73.6	72.1-75.2	4,708	97.4	94.6-100.3	3,662	57.2	55.3-59.1	6,534	57.4	56.0-58.8	3,745	78.7	76.2-81.3	2,789	43.1	41.5-44.7
1997	8,422	73.5	71.9-75.1	4,624	94.9	92.1-97.7	3,798	58.9	57.1-60.9	6,439	56.1	54.7-57.5	3,614	75.4	72.9-77.9	2,825	43.1	41.5-44.7
1998	8,554	73.9	72.4-75.5	4,623	93.1	90.4-95.8	3,931	60.7	58.8-62.6	6,464	55.7	54.4-57.1	3,663	74.6	72.2-77.1	2,801	42.3	40.7-43.9
1999	8,616	73.7	72.1-75.3	4,647	93.0	90.3-95.8	3,969	60.4	58.6-62.4	6,403	54.6	53.3-56.0	3,525	71.6	69.2-74.0	2,878	43.1	41.5-44.7
2000	8,800	74.4	72.8-75.9	4,694	92.7	90.1-95.5	4,106	61.6	59.7-63.6	6,602	55.6	54.3-57.0	3,621	72.4	70.1-74.9	2,981	44.0	42.4-45.6
2001	8,622	72.3	70.7-73.8	4,488	88.1	85.5-90.7	4,134	61.8	59.9-63.8	6,463	53.9	52.6-55.3	3,531	70.6	68.2-73.0	2,932	43.0	41.5-44.6
2002	8,800	72.8	71.3-74.4	4,652	90.0	87.4-92.7	4,148	61.3	59.4-63.2	6,599	54.4	53.1-55.8	3,538	69.3	67.0-71.7	3,061	44.6	43.0-46.2
2003	8,914	72.9	71.4-74.4	4,691	89.1	86.5-91.7	4,223	61.8	60.0-63.7	6,441	52.5	51.2-53.8	3,469	66.8	64.5-69.1	2,972	42.7	41.1-44.2
2004	8,962	72.5	71.0-74.0	4,564	85.4	83.0-88.0	4,398	63.6	61.7-65.5	6,483	52.3	51.0-53.6	3,357	63.7	61.6-65.9	3,126	44.4	42.9-46.0
2005	8,992	72.2	70.7-73.7	4,609	85.4	82.9-87.9	4,383	62.9	61.0-64.8	6,421	51.4	50.1-52.7	3,364	63.3	61.2-65.5	3,057	43.0	41.5-44.6
2006	9,224	73.6	72.1-75.1	4,561	83.5	81.1-86.0	4,663	67.0	65.1-69.0	6,416	50.8	49.6-52.1	3,401	63.0	60.9-65.2	3,015	42.4	40.9-43.9
2007	9,204	72.2	70.7-73.7	4,660	83.9	81.5-86.4	4,544	64.1	62.2-66.0	6,467	50.5	49.2-51.7	3,351	60.7	58.6-62.8	3,116	43.3	41.8-44.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Lung and bronchus

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	9,497	73.4	71.9-74.9	4,752	84.3	81.9-86.7	4,745	66.2	64.3-68.1	6,382	48.9	47.7-50.1	3,291	58.8	56.8-60.9	3,091	42.1	40.6-43.7
2009	9,393	71.5	70.0-72.9	4,805	83.4	81.0-85.9	4,588	62.9	61.1-64.8	6,238	47.4	46.2-48.6	3,274	57.4	55.4-59.4	2,964	40.2	38.8-41.7
2010	9,305	69.7	68.3-71.2	4,598	78.0	75.8-80.4	4,707	64.3	62.4-66.2	6,236	46.7	45.5-47.9	3,197	55.0	53.1-57.0	3,039	40.8	39.3-42.3
2011	9,451	69.6	68.2-71.1	4,694	78.4	76.1-80.7	4,757	63.7	61.8-65.5	6,218	45.8	44.6-46.9	3,186	53.9	52.0-55.9	3,032	40.0	38.5-41.4
2012	9,435	68.3	66.9-69.8	4,706	77.2	74.9-79.4	4,729	62.3	60.5-64.1	6,369	46.2	45.0-47.3	3,320	55.3	53.4-57.2	3,049	39.6	38.2-41.1
2013	9,497	67.8	66.4-69.2	4,651	74.6	72.4-76.8	4,846	63.0	61.2-64.8	6,108	43.3	42.2-44.4	3,113	50.2	48.4-52.1	2,995	38.1	36.7-39.5
2014	9,269	64.8	63.4-66.1	4,546	71.5	69.4-73.7	4,723	60.0	58.3-61.8	5,877	41.1	40.1-42.2	3,037	48.4	46.7-50.2	2,840	35.8	34.5-37.2
2015	9,822	67.7	66.4-69.1	4,727	73.3	71.2-75.5	5,095	64.1	62.3-65.9	5,891	40.5	39.4-41.5	3,012	47.2	45.5-48.9	2,879	35.6	34.3-36.9
2016	9,690	65.5	64.2-66.9	4,718	71.4	69.3-73.5	4,972	61.7	59.9-63.4	5,564	37.6	36.6-38.7	2,827	43.5	41.9-45.2	2,737	33.3	32.0-34.6
2017	9,986	66.4	65.0-67.7	4,744	70.2	68.2-72.3	5,242	64.0	62.2-65.8	5,347	35.4	34.4-36.4	2,690	40.4	38.8-42.0	2,657	31.7	30.4-32.9
2018	9,855	64.2	62.9-65.5	4,685	67.9	65.9-69.9	5,170	61.9	60.2-63.6	5,344	34.8	33.8-35.7	2,708	39.9	38.4-41.5	2,636	31.0	29.8-32.2
2019	9,811	62.6	61.3-63.9	4,632	65.7	63.8-67.7	5,179	60.8	59.1-62.5	5,172	32.8	31.9-33.8	2,584	37.2	35.8-38.7	2,588	29.6	28.5-30.8
2020	8,817	55.3	54.1-56.5	4,116	57.1	55.3-58.9	4,701	54.3	52.8-55.9	4,825	30.2	29.3-31.1	2,438	34.4	33.0-35.8	2,387	27.1	26.0-28.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State Excl New York City
Melanoma of the skin**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	559	5.9	5.4-6.4	286	6.6	5.8-7.4	273	5.4	4.7-6.1	237	2.5	2.2-2.9	131	3.1	2.6-3.7	106	2.1	1.7-2.5
1977	708	7.4	6.9-8.0	388	9.0	8.1-9.9	320	6.3	5.6-7.0	250	2.7	2.4-3.1	161	3.9	3.3-4.6	89	1.8	1.4-2.2
1978	675	7.1	6.6-7.7	343	8.1	7.2-9.0	332	6.6	5.9-7.3	220	2.4	2.0-2.7	139	3.3	2.8-3.9	81	1.6	1.2-2.0
1979	694	7.2	6.7-7.8	368	8.6	7.7-9.5	326	6.3	5.6-7.1	238	2.5	2.2-2.9	134	3.2	2.6-3.8	104	2.0	1.6-2.4
1980	857	8.8	8.2-9.5	470	10.9	9.9-12.0	387	7.4	6.7-8.2	251	2.6	2.3-2.9	152	3.5	2.9-4.1	99	1.9	1.5-2.3
1981	841	8.6	8.0-9.2	435	9.9	9.0-11.0	406	7.7	6.9-8.5	224	2.3	2.0-2.6	133	3.1	2.5-3.6	91	1.7	1.3-2.1
1982	883	8.9	8.3-9.5	429	9.6	8.7-10.6	454	8.5	7.7-9.3	256	2.6	2.3-3.0	159	3.7	3.1-4.3	97	1.8	1.4-2.2
1983	952	9.6	9.0-10.2	521	11.8	10.8-12.9	431	8.2	7.5-9.1	246	2.5	2.2-2.8	137	3.1	2.6-3.7	109	2.0	1.7-2.5
1984	840	8.4	7.8-9.0	456	10.1	9.1-11.1	384	7.3	6.5-8.0	255	2.5	2.2-2.9	145	3.3	2.8-3.9	110	2.0	1.6-2.4
1985	939	9.3	8.7-10.0	501	11.0	10.1-12.1	438	8.1	7.3-8.9	254	2.5	2.2-2.8	152	3.5	2.9-4.1	102	1.8	1.4-2.2
1986	989	9.7	9.1-10.4	532	11.8	10.8-12.9	457	8.5	7.7-9.4	281	2.8	2.5-3.1	163	3.7	3.1-4.3	118	2.2	1.8-2.6
1987	926	9.0	8.4-9.6	507	11.0	10.0-12.0	419	7.6	6.9-8.4	298	2.9	2.6-3.3	180	4.0	3.4-4.7	118	2.2	1.8-2.6
1988	893	8.6	8.0-9.2	507	11.1	10.1-12.2	386	6.9	6.2-7.6	295	2.8	2.5-3.2	191	4.2	3.6-4.9	104	1.8	1.5-2.2
1989	941	9.0	8.5-9.6	530	11.4	10.4-12.4	411	7.3	6.6-8.0	321	3.1	2.8-3.5	204	4.7	4.0-5.4	117	2.0	1.6-2.4
1990	990	9.3	8.7-9.9	527	11.1	10.1-12.1	463	8.1	7.3-8.9	297	2.8	2.5-3.2	181	3.9	3.4-4.6	116	2.0	1.7-2.4
1991	931	8.7	8.1-9.2	539	11.3	10.4-12.4	392	6.7	6.1-7.4	300	2.8	2.5-3.1	187	4.1	3.5-4.8	113	1.9	1.6-2.3
1992	944	8.7	8.2-9.3	551	11.5	10.5-12.5	393	6.7	6.0-7.4	298	2.8	2.5-3.1	174	3.7	3.2-4.4	124	2.0	1.7-2.4
1993	959	8.7	8.1-9.3	544	11.1	10.2-12.1	415	6.9	6.3-7.6	297	2.7	2.4-3.0	185	4.0	3.4-4.6	112	1.8	1.5-2.2
1994	1,029	9.3	8.8-9.9	602	12.3	11.3-13.3	427	7.2	6.5-7.9	298	2.7	2.4-3.0	185	3.9	3.4-4.5	113	1.8	1.5-2.2
1995	1,136	10.3	9.7-10.9	674	13.6	12.6-14.7	462	7.7	7.0-8.5	315	2.8	2.5-3.2	184	3.9	3.3-4.5	131	2.1	1.7-2.5
1996	1,206	10.8	10.2-11.5	691	13.9	12.9-15.0	515	8.6	7.8-9.3	325	2.9	2.6-3.2	200	4.0	3.5-4.6	125	2.0	1.7-2.4
1997	1,297	11.5	10.9-12.2	757	15.1	14.0-16.2	540	8.9	8.2-9.7	305	2.7	2.4-3.0	205	4.3	3.7-4.9	100	1.5	1.2-1.9
1998	1,381	12.2	11.6-12.9	787	15.5	14.4-16.6	594	9.9	9.1-10.7	296	2.6	2.3-2.9	189	3.8	3.3-4.4	107	1.7	1.4-2.0
1999	1,411	12.3	11.7-13.0	815	15.9	14.8-17.1	596	9.7	8.9-10.5	285	2.5	2.2-2.8	168	3.4	2.9-4.0	117	1.8	1.5-2.2
2000	1,571	13.6	12.9-14.3	896	17.3	16.2-18.5	675	10.9	10.1-11.8	308	2.6	2.3-2.9	199	3.9	3.3-4.4	109	1.7	1.4-2.0
2001	1,754	15.0	14.3-15.7	987	18.8	17.7-20.1	767	12.3	11.5-13.2	361	3.0	2.7-3.4	232	4.6	4.0-5.2	129	1.9	1.6-2.3
2002	1,739	14.7	14.0-15.4	1,007	19.0	17.9-20.2	732	11.6	10.8-12.5	315	2.6	2.3-2.9	169	3.2	2.8-3.8	146	2.2	1.8-2.6
2003	1,803	15.0	14.4-15.8	1,050	19.6	18.5-20.9	753	11.8	11.0-12.7	318	2.6	2.3-2.9	193	3.7	3.2-4.2	125	1.8	1.5-2.2
2004	2,145	17.8	17.0-18.5	1,231	22.6	21.3-23.9	914	14.4	13.4-15.3	325	2.6	2.4-2.9	198	3.7	3.2-4.3	127	1.9	1.6-2.2
2005	2,345	19.4	18.6-20.2	1,368	25.0	23.7-26.4	977	15.3	14.3-16.3	338	2.7	2.4-3.0	210	3.9	3.4-4.4	128	1.9	1.6-2.2
2006	2,522	20.6	19.8-21.4	1,451	26.1	24.8-27.5	1,071	16.6	15.6-17.7	345	2.7	2.5-3.0	224	4.1	3.6-4.7	121	1.7	1.4-2.0
2007	2,436	19.7	18.9-20.5	1,402	25.0	23.7-26.4	1,034	16.1	15.1-17.1	329	2.6	2.3-2.9	236	4.3	3.8-4.9	93	1.3	1.0-1.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Melanoma of the skin

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	2,689	21.5	20.7-22.4	1,552	27.5	26.1-28.9	1,137	17.5	16.4-18.6	348	2.7	2.4-3.0	229	4.0	3.5-4.6	119	1.6	1.4-2.0
2009	2,847	22.5	21.6-23.3	1,670	29.0	27.6-30.5	1,177	17.9	16.9-19.0	357	2.7	2.5-3.0	211	3.8	3.3-4.3	146	2.0	1.7-2.4
2010	2,727	21.3	20.5-22.2	1,563	26.8	25.4-28.2	1,164	17.4	16.4-18.5	354	2.7	2.4-3.0	215	3.7	3.2-4.2	139	1.9	1.6-2.3
2011	3,041	23.2	22.3-24.0	1,771	29.7	28.3-31.1	1,270	18.5	17.5-19.6	374	2.8	2.5-3.1	244	4.1	3.6-4.7	130	1.7	1.4-2.1
2012	2,961	22.4	21.6-23.3	1,680	27.7	26.3-29.0	1,281	18.9	17.8-20.0	399	2.9	2.6-3.2	251	4.3	3.8-4.9	148	1.9	1.6-2.2
2013	3,296	24.7	23.9-25.6	1,889	30.5	29.1-31.9	1,407	20.7	19.6-21.9	387	2.8	2.6-3.2	255	4.3	3.8-4.9	132	1.7	1.4-2.0
2014	3,267	24.4	23.5-25.2	1,911	30.7	29.3-32.1	1,356	19.8	18.7-20.9	340	2.4	2.1-2.7	219	3.5	3.0-4.0	121	1.5	1.2-1.8
2015	3,227	23.5	22.7-24.4	1,903	30.0	28.6-31.4	1,324	18.9	17.8-20.0	297	2.1	1.8-2.3	195	3.1	2.7-3.6	102	1.3	1.1-1.6
2016	3,248	23.4	22.5-24.2	1,927	29.8	28.5-31.3	1,321	18.7	17.7-19.8	326	2.2	2.0-2.5	208	3.3	2.8-3.8	118	1.4	1.2-1.7
2017	3,325	23.8	23.0-24.7	1,968	30.5	29.1-31.9	1,357	19.1	18.1-20.2	295	2.0	1.8-2.3	191	3.0	2.6-3.5	104	1.2	1.0-1.5
2018	3,397	23.8	22.9-24.6	1,955	29.3	28.0-30.7	1,442	19.8	18.7-20.9	313	2.0	1.8-2.3	191	2.8	2.4-3.3	122	1.4	1.2-1.7
2019	3,484	24.3	23.4-25.1	2,040	30.2	28.9-31.6	1,444	20.1	19.0-21.2	277	1.9	1.6-2.1	173	2.6	2.3-3.1	104	1.2	1.0-1.5
2020	3,058	21.1	20.3-21.9	1,722	25.3	24.1-26.6	1,336	18.1	17.1-19.2	306	2.0	1.8-2.3	206	3.2	2.7-3.6	100	1.1	0.9-1.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Female breast

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	5,329	103.8	101.0-106.7	2,030	38.8	37.1-40.5
1977	5,338	102.8	100.0-105.7	2,083	39.4	37.7-41.2
1978	5,443	103.6	100.8-106.5	2,091	38.8	37.1-40.5
1979	5,399	101.8	99.0-104.6	2,065	38.2	36.5-39.9
1980	5,595	104.3	101.5-107.1	2,065	37.5	35.9-39.2
1981	5,630	104.5	101.7-107.3	2,084	37.5	35.8-39.1
1982	5,795	106.1	103.3-109.0	2,115	37.7	36.1-39.4
1983	6,175	112.6	109.8-115.5	2,140	37.9	36.2-39.5
1984	6,309	113.7	110.8-116.6	2,229	39.0	37.4-40.7
1985	6,687	120.7	117.8-123.7	2,261	39.6	37.9-41.3
1986	6,851	122.4	119.4-125.4	2,295	39.6	38.0-41.3
1987	7,086	125.4	122.4-128.4	2,292	39.0	37.4-40.6
1988	7,236	126.7	123.7-129.7	2,283	38.9	37.3-40.5
1989	7,094	123.4	120.5-126.4	2,275	38.5	36.9-40.2
1990	7,571	130.5	127.6-133.6	2,231	37.0	35.5-38.6
1991	7,477	127.1	124.2-130.1	2,178	36.0	34.4-37.5
1992	7,559	127.8	124.8-130.7	2,170	35.3	33.8-36.9
1993	7,589	127.1	124.3-130.1	2,153	34.6	33.1-36.1
1994	7,693	128.2	125.3-131.1	2,153	34.3	32.9-35.8
1995	8,045	133.6	130.7-136.6	2,055	32.7	31.2-34.1
1996	8,517	140.3	137.3-143.3	2,078	32.4	31.0-33.9
1997	8,723	142.2	139.2-145.2	1,977	30.6	29.2-32.0
1998	9,004	145.2	142.2-148.2	2,000	30.5	29.2-31.9
1999	8,928	142.2	139.3-145.2	1,853	27.9	26.6-29.2
2000	8,971	140.7	137.8-143.6	1,896	28.1	26.8-29.4
2001	8,857	137.8	134.9-140.7	1,840	27.1	25.9-28.4
2002	8,897	137.2	134.3-140.1	1,824	26.4	25.2-27.7
2003	8,857	134.8	132.0-137.7	1,823	26.2	25.0-27.4
2004	8,781	132.6	129.8-135.5	1,688	23.9	22.7-25.1
2005	8,926	133.3	130.5-136.2	1,662	23.4	22.2-24.5
2006	9,098	135.8	133.0-138.7	1,588	22.2	21.1-23.4
2007	9,077	133.6	130.9-136.5	1,668	23.0	21.9-24.2
2008	9,450	138.1	135.3-141.0	1,602	21.8	20.8-23.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Female breast

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	9,675	139.1	136.3-142.0	1,608	21.6	20.5-22.7
2010	9,120	130.2	127.5-133.0	1,653	21.7	20.6-22.8
2011	9,721	137.9	135.1-140.8	1,628	21.3	20.2-22.4
2012	9,557	134.9	132.1-137.7	1,537	19.8	18.8-20.9
2013	9,901	138.4	135.6-141.3	1,566	20.3	19.2-21.3
2014	10,169	141.1	138.3-144.0	1,410	17.9	16.9-18.9
2015	10,266	142.0	139.1-144.9	1,493	18.7	17.7-19.7
2016	10,007	137.4	134.6-140.3	1,540	19.5	18.5-20.5
2017	10,334	141.2	138.4-144.1	1,456	17.9	16.9-18.9
2018	10,377	140.5	137.7-143.4	1,548	19.0	18.0-20.0
2019	10,884	147.0	144.1-149.9	1,460	17.9	16.9-18.8
2020	9,688	130.3	127.5-133.1	1,488	18.1	17.1-19.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Cervix uteri

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	690	13.7	12.6-14.7	264	5.0	4.4-5.7
1977	593	11.6	10.7-12.6	219	4.1	3.6-4.7
1978	603	11.7	10.8-12.7	246	4.7	4.1-5.3
1979	595	11.6	10.6-12.5	241	4.6	4.0-5.2
1980	626	11.7	10.8-12.7	236	4.5	3.9-5.1
1981	505	9.4	8.6-10.3	248	4.5	3.9-5.1
1982	517	9.8	8.9-10.7	208	3.7	3.2-4.3
1983	544	10.2	9.4-11.2	191	3.5	3.0-4.0
1984	581	11.0	10.1-11.9	178	3.2	2.8-3.8
1985	526	9.8	9.0-10.7	185	3.4	2.9-3.9
1986	497	9.3	8.5-10.2	203	3.7	3.2-4.3
1987	562	10.2	9.4-11.1	201	3.6	3.1-4.1
1988	567	10.4	9.6-11.4	189	3.3	2.8-3.8
1989	545	9.9	9.1-10.8	170	2.9	2.5-3.4
1990	594	10.8	9.9-11.7	187	3.2	2.8-3.7
1991	573	10.1	9.3-11.0	161	2.8	2.4-3.3
1992	513	9.0	8.2-9.8	182	3.1	2.7-3.6
1993	560	9.7	8.9-10.6	172	2.9	2.5-3.4
1994	539	9.3	8.5-10.2	173	2.9	2.5-3.4
1995	634	11.0	10.1-11.8	135	2.3	1.9-2.7
1996	622	10.7	9.8-11.6	168	2.9	2.4-3.3
1997	540	9.2	8.5-10.0	183	3.0	2.6-3.5
1998	507	8.7	7.9-9.5	142	2.3	2.0-2.8
1999	508	8.6	7.8-9.4	161	2.6	2.2-3.0
2000	463	7.8	7.1-8.5	144	2.3	1.9-2.7
2001	492	8.1	7.4-8.9	141	2.2	1.8-2.6
2002	428	7.1	6.4-7.8	150	2.4	2.0-2.8
2003	484	8.1	7.3-8.8	140	2.1	1.8-2.5
2004	411	6.9	6.2-7.6	137	2.1	1.8-2.5
2005	463	7.7	7.0-8.5	142	2.2	1.8-2.6
2006	444	7.4	6.7-8.1	130	2.0	1.7-2.4
2007	454	7.4	6.8-8.2	143	2.1	1.8-2.5
2008	442	7.3	6.6-8.0	117	1.7	1.4-2.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Cervix uteri

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	462	7.6	6.9-8.3	118	1.8	1.4-2.1
2010	454	7.6	6.9-8.4	129	1.8	1.5-2.2
2011	375	6.1	5.5-6.8	128	1.9	1.6-2.3
2012	412	6.6	6.0-7.3	163	2.4	2.0-2.8
2013	446	7.2	6.5-8.0	140	2.0	1.7-2.4
2014	416	6.7	6.1-7.4	136	2.0	1.6-2.4
2015	442	7.3	6.6-8.0	121	1.7	1.4-2.0
2016	393	6.6	5.9-7.3	128	1.8	1.5-2.2
2017	444	7.3	6.6-8.0	138	1.9	1.6-2.3
2018	378	6.3	5.7-7.0	111	1.5	1.2-1.9
2019	388	6.7	6.0-7.4	130	1.9	1.5-2.3
2020	363	6.0	5.3-6.7	105	1.4	1.1-1.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State Excl New York City
Corpus uteris and uterus, not otherwise specified**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,584	29.5	28.0-31.0	298	5.6	5.0-6.3
1977	1,454	26.4	25.1-27.8	301	5.5	4.9-6.2
1978	1,281	23.0	21.8-24.3	267	4.8	4.2-5.4
1979	1,420	25.5	24.1-26.9	304	5.4	4.8-6.0
1980	1,418	25.5	24.2-26.9	313	5.4	4.8-6.0
1981	1,511	26.8	25.4-28.2	304	5.3	4.7-5.9
1982	1,475	25.8	24.5-27.2	311	5.3	4.8-6.0
1983	1,403	24.6	23.3-25.9	295	4.9	4.4-5.5
1984	1,449	25.2	23.9-26.6	307	5.1	4.5-5.7
1985	1,391	24.0	22.7-25.3	301	5.0	4.4-5.6
1986	1,399	24.1	22.9-25.5	287	4.6	4.1-5.2
1987	1,495	25.5	24.2-26.9	284	4.6	4.1-5.2
1988	1,372	23.3	22.1-24.6	294	4.7	4.2-5.3
1989	1,376	23.1	21.9-24.4	302	4.8	4.3-5.4
1990	1,453	24.4	23.1-25.7	253	4.0	3.5-4.5
1991	1,463	24.9	23.6-26.3	289	4.5	4.0-5.0
1992	1,559	25.9	24.6-27.2	304	4.7	4.2-5.3
1993	1,494	24.7	23.4-26.0	313	4.7	4.2-5.3
1994	1,611	26.3	25.0-27.6	302	4.6	4.1-5.1
1995	1,593	26.1	24.8-27.5	319	4.8	4.3-5.3
1996	1,719	27.9	26.6-29.3	301	4.5	4.0-5.1
1997	1,660	26.9	25.6-28.2	285	4.3	3.8-4.8
1998	1,824	29.3	28.0-30.7	300	4.5	4.0-5.0
1999	1,756	27.8	26.5-29.1	273	4.0	3.5-4.5
2000	1,750	27.3	26.0-28.6	308	4.4	4.0-5.0
2001	1,914	29.7	28.4-31.1	299	4.2	3.8-4.8
2002	1,925	29.4	28.1-30.8	319	4.6	4.1-5.1
2003	1,933	29.2	27.9-30.5	344	4.9	4.4-5.4
2004	1,978	29.5	28.2-30.8	314	4.4	3.9-4.9
2005	2,030	29.9	28.6-31.3	327	4.6	4.1-5.1
2006	1,997	29.0	27.7-30.3	353	4.9	4.4-5.4
2007	2,110	30.5	29.2-31.8	344	4.6	4.2-5.2
2008	2,214	31.6	30.3-33.0	358	4.9	4.4-5.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State Excl New York City
Corpus uteris and uterus, not otherwise specified**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	2,134	30.0	28.7-31.3	340	4.6	4.1-5.1
2010	2,184	30.2	29.0-31.6	355	4.6	4.1-5.1
2011	2,263	30.8	29.5-32.1	372	4.9	4.4-5.5
2012	2,388	31.8	30.5-33.1	394	5.1	4.6-5.6
2013	2,496	32.9	31.6-34.3	427	5.3	4.8-5.8
2014	2,528	33.0	31.7-34.4	425	5.2	4.7-5.7
2015	2,469	31.9	30.6-33.2	372	4.6	4.1-5.1
2016	2,568	33.2	31.9-34.6	427	5.2	4.7-5.7
2017	2,526	32.0	30.7-33.3	440	5.3	4.8-5.9
2018	2,459	30.9	29.6-32.2	473	5.6	5.1-6.2
2019	2,545	31.3	30.1-32.6	480	5.7	5.2-6.2
2020	2,471	30.2	29.0-31.5	460	5.3	4.8-5.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Ovary

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	922	17.6	16.4-18.8	671	12.5	11.6-13.5
1977	897	16.8	15.7-17.9	573	10.6	9.7-11.5
1978	875	16.1	15.1-17.3	610	11.0	10.1-11.9
1979	921	17.0	15.9-18.1	595	10.8	9.9-11.7
1980	906	16.5	15.4-17.7	604	10.8	10.0-11.7
1981	984	17.9	16.8-19.1	642	11.3	10.4-12.2
1982	934	16.9	15.8-18.1	637	11.1	10.2-12.0
1983	998	17.9	16.7-19.0	628	10.7	9.9-11.6
1984	938	16.7	15.7-17.9	626	10.8	9.9-11.7
1985	975	17.5	16.4-18.7	603	10.4	9.5-11.3
1986	991	17.6	16.5-18.8	589	10.1	9.3-10.9
1987	970	17.0	15.9-18.1	645	10.9	10.0-11.8
1988	993	17.3	16.2-18.4	641	10.7	9.8-11.5
1989	999	17.2	16.1-18.3	648	10.7	9.9-11.6
1990	1,047	17.9	16.8-19.0	661	10.7	9.9-11.6
1991	1,043	17.7	16.6-18.8	685	11.1	10.3-12.0
1992	1,047	17.6	16.6-18.8	671	10.9	10.1-11.8
1993	1,054	17.4	16.3-18.5	648	10.2	9.4-11.0
1994	1,070	17.6	16.6-18.7	644	10.2	9.4-11.0
1995	1,087	17.8	16.8-18.9	667	10.4	9.6-11.2
1996	1,097	17.8	16.8-18.9	627	9.7	9.0-10.5
1997	1,068	17.3	16.3-18.4	680	10.5	9.7-11.3
1998	981	15.8	14.8-16.8	600	9.1	8.4-9.9
1999	1,021	16.1	15.1-17.2	611	9.2	8.5-10.0
2000	1,056	16.4	15.4-17.4	668	9.9	9.2-10.7
2001	1,042	16.0	15.1-17.1	667	9.7	9.0-10.5
2002	1,011	15.5	14.5-16.5	644	9.4	8.7-10.2
2003	1,024	15.4	14.5-16.4	674	9.7	8.9-10.4
2004	997	14.9	14.0-15.8	650	9.2	8.5-9.9
2005	948	14.1	13.2-15.0	688	9.7	9.0-10.5
2006	991	14.5	13.6-15.5	672	9.5	8.8-10.2
2007	956	13.8	13.0-14.8	613	8.6	7.9-9.3
2008	973	14.0	13.1-14.9	645	8.8	8.1-9.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Ovary

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	920	13.1	12.2-14.0	610	8.2	7.5-8.9
2010	962	13.5	12.7-14.5	597	8.0	7.4-8.7
2011	958	13.2	12.4-14.1	608	7.9	7.3-8.6
2012	910	12.6	11.8-13.5	587	7.6	7.0-8.2
2013	998	13.8	12.9-14.7	632	8.1	7.4-8.7
2014	876	12.0	11.2-12.8	573	7.2	6.6-7.8
2015	897	12.3	11.5-13.2	577	7.2	6.6-7.9
2016	909	12.2	11.4-13.1	550	6.9	6.3-7.5
2017	868	11.5	10.7-12.3	518	6.3	5.8-6.9
2018	830	11.3	10.5-12.2	536	6.4	5.8-7.0
2019	869	11.7	10.9-12.5	525	6.2	5.7-6.8
2020	731	9.7	8.9-10.4	543	6.4	5.8-6.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Prostate

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	2,711	81.2	78.0-84.5	1,003	32.5	30.4-34.6
1977	2,777	80.7	77.5-83.9	994	31.9	29.8-34.0
1978	2,761	79.9	76.8-83.1	1,038	33.3	31.2-35.5
1979	2,837	81.6	78.5-84.9	1,069	33.5	31.4-35.7
1980	3,188	87.9	84.7-91.2	1,052	32.7	30.6-34.8
1981	3,239	89.5	86.3-92.8	1,036	31.9	29.9-34.0
1982	3,378	91.3	88.1-94.6	1,089	32.7	30.6-34.7
1983	3,588	95.3	92.0-98.6	1,168	34.5	32.4-36.6
1984	3,507	91.6	88.5-94.9	1,204	35.3	33.2-37.4
1985	3,703	95.2	92.0-98.5	1,177	34.3	32.2-36.4
1986	3,793	96.5	93.3-99.8	1,284	36.1	34.1-38.2
1987	4,125	103.4	100.2-106.8	1,255	35.4	33.3-37.5
1988	4,269	105.8	102.5-109.2	1,288	35.7	33.7-37.8
1989	4,486	109.3	105.9-112.7	1,305	35.9	33.8-38.0
1990	5,312	123.5	120.1-127.1	1,340	36.0	34.0-38.1
1991	6,460	146.2	142.5-149.9	1,462	38.7	36.6-40.8
1992	8,222	179.6	175.6-183.6	1,397	36.7	34.8-38.8
1993	7,995	170.9	167.1-174.9	1,438	37.3	35.4-39.4
1994	7,302	154.0	150.4-157.7	1,490	38.0	36.0-40.0
1995	7,272	150.2	146.7-153.8	1,407	35.5	33.6-37.5
1996	8,020	163.9	160.3-167.6	1,416	34.6	32.8-36.5
1997	8,363	169.5	165.8-173.2	1,414	34.0	32.2-35.9
1998	8,306	165.8	162.2-169.4	1,265	29.7	28.1-31.5
1999	8,790	173.5	169.9-177.2	1,262	29.1	27.5-30.8
2000	9,530	186.8	183.0-190.6	1,283	29.6	28.0-31.3
2001	9,608	185.1	181.4-188.9	1,263	28.5	27.0-30.2
2002	10,012	189.7	186.0-193.5	1,192	26.4	24.9-28.0
2003	9,418	175.9	172.3-179.5	1,207	26.0	24.6-27.6
2004	9,113	167.2	163.7-170.7	1,151	24.4	23.0-25.9
2005	9,268	166.9	163.4-170.3	1,075	22.2	20.9-23.6
2006	10,405	184.8	181.2-188.4	1,047	21.3	20.0-22.6
2007	10,628	183.6	180.0-187.1	1,102	21.7	20.4-23.0
2008	10,310	173.7	170.3-177.1	1,060	20.3	19.1-21.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Prostate

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	10,107	165.4	162.1-168.7	998	18.9	17.7-20.1
2010	9,669	154.6	151.4-157.7	1,071	19.7	18.5-20.9
2011	10,031	157.0	153.9-160.2	1,018	18.4	17.3-19.6
2012	8,593	130.7	127.9-133.6	990	17.7	16.6-18.8
2013	8,457	126.0	123.3-128.8	947	16.4	15.3-17.5
2014	8,278	120.3	117.7-123.0	1,015	17.2	16.1-18.3
2015	8,746	124.5	121.9-127.2	969	16.3	15.2-17.3
2016	9,037	126.7	124.0-129.4	1,045	16.9	15.9-18.0
2017	9,744	133.4	130.7-136.1	1,074	17.2	16.2-18.3
2018	10,340	139.3	136.6-142.1	1,043	16.2	15.2-17.2
2019	10,892	144.1	141.4-146.9	975	15.0	14.1-16.0
2020	9,412	123.0	120.5-125.6	1,052	15.9	14.9-16.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Testis

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	179	3.6	3.0-4.1	33	0.6	0.4-0.9
1977	177	3.5	3.0-4.1	29	0.5	0.4-0.8
1978	201	4.1	3.5-4.7	35	0.7	0.5-1.0
1979	227	4.6	4.0-5.2	31	0.7	0.4-1.0
1980	236	4.6	4.0-5.2	30	0.6	0.4-0.8
1981	228	4.3	3.8-5.0	28	0.5	0.3-0.7
1982	228	4.4	3.8-5.0	25	0.5	0.3-0.8
1983	250	4.7	4.1-5.3	24	0.5	0.3-0.8
1984	295	5.6	4.9-6.3	17	0.3	0.2-0.5
1985	265	5.0	4.4-5.6	21	0.4	0.2-0.6
1986	254	4.6	4.1-5.3	25	0.5	0.3-0.7
1987	281	5.1	4.5-5.7	22	0.4	0.2-0.6
1988	270	4.8	4.2-5.4	16	0.3	0.2-0.5
1989	268	4.8	4.2-5.4	22	0.4	0.2-0.6
1990	296	5.3	4.7-5.9	16	0.3	0.2-0.5
1991	275	4.9	4.3-5.5	14	0.2	0.1-0.4
1992	298	5.3	4.7-6.0	20	0.4	0.2-0.6
1993	312	5.6	5.0-6.3	24	0.5	0.3-0.7
1994	304	5.5	4.9-6.1	12	0.2	0.1-0.4
1995	313	5.7	5.1-6.4	16	0.3	0.2-0.5
1996	310	5.7	5.1-6.3	21	0.4	0.2-0.6
1997	333	6.2	5.5-6.9	6	0.1	0.0-0.3
1998	342	6.4	5.7-7.1	18	0.3	0.2-0.5
1999	334	6.2	5.6-7.0	14	0.3	0.1-0.4
2000	306	5.7	5.1-6.4	13	0.2	0.1-0.4
2001	329	6.2	5.5-6.9	13	0.3	0.1-0.4
2002	312	5.9	5.3-6.6	24	0.5	0.3-0.7
2003	353	6.7	6.0-7.4	12	0.2	0.1-0.4
2004	331	6.3	5.7-7.0	11	0.2	0.1-0.4
2005	349	6.7	6.0-7.4	8	0.1	0.1-0.3
2006	326	6.2	5.5-6.9	13	0.2	0.1-0.4
2007	352	6.8	6.1-7.5	8	0.1	0.1-0.3
2008	383	7.4	6.7-8.2	11	0.2	0.1-0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Testis

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	342	6.6	5.9-7.3	7	0.1	0.1-0.3
2010	359	6.9	6.2-7.7	15	0.3	0.2-0.5
2011	362	6.9	6.2-7.7	9	0.2	0.1-0.3
2012	378	7.3	6.5-8.1	12	0.2	0.1-0.4
2013	342	6.6	5.9-7.4	8	0.1	0.1-0.3
2014	363	7.0	6.3-7.8	15	0.3	0.2-0.5
2015	352	6.7	6.0-7.5	13	0.2	0.1-0.4
2016	373	7.2	6.5-8.0	11	0.2	0.1-0.3
2017	354	6.9	6.2-7.7	9	0.1	0.1-0.3
2018	385	7.5	6.7-8.3	10	0.2	0.1-0.3
2019	338	6.6	5.9-7.4	28	0.5	0.3-0.7
2020	319	6.3	5.6-7.0	17	0.3	0.2-0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Urinary bladder (incl. in situ)

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,984	21.7	20.8-22.7	1,461	39.6	37.4-41.8	523	9.8	9.0-10.7	601	7.0	6.4-7.6	431	13.0	11.7-14.4	170	3.3	2.8-3.9
1977	2,023	21.8	20.9-22.8	1,476	38.8	36.7-40.9	547	10.3	9.5-11.2	599	6.9	6.4-7.5	419	12.6	11.3-13.9	180	3.5	3.0-4.0
1978	2,141	22.8	21.9-23.8	1,559	41.0	38.9-43.3	582	10.8	9.9-11.7	614	6.9	6.3-7.5	428	12.7	11.4-14.0	186	3.4	2.9-4.0
1979	2,166	22.7	21.7-23.6	1,622	42.1	40.0-44.3	544	9.8	8.9-10.6	602	6.7	6.2-7.3	403	11.7	10.6-13.0	199	3.7	3.2-4.2
1980	2,255	23.2	22.3-24.2	1,657	42.1	40.0-44.3	598	10.6	9.8-11.5	576	6.2	5.7-6.7	405	11.6	10.4-12.8	171	3.1	2.6-3.6
1981	2,207	22.4	21.5-23.4	1,633	41.1	39.0-43.2	574	10.1	9.3-10.9	632	6.8	6.2-7.3	450	12.9	11.7-14.2	182	3.2	2.8-3.7
1982	2,378	23.9	23.0-24.9	1,735	43.0	40.9-45.2	643	11.1	10.2-12.0	584	6.1	5.7-6.7	410	11.4	10.3-12.6	174	3.0	2.6-3.5
1983	2,307	22.9	22.0-23.9	1,715	42.4	40.3-44.6	592	10.1	9.3-10.9	582	6.1	5.6-6.6	419	12.0	10.8-13.2	163	2.8	2.3-3.2
1984	2,329	22.9	21.9-23.8	1,684	41.0	38.9-43.1	645	10.9	10.1-11.8	592	6.2	5.7-6.7	405	11.6	10.4-12.9	187	3.1	2.7-3.6
1985	2,446	23.9	22.9-24.8	1,780	42.9	40.9-45.1	666	11.2	10.4-12.1	581	5.9	5.5-6.4	396	10.8	9.7-12.0	185	3.1	2.7-3.6
1986	2,522	24.3	23.3-25.3	1,856	43.9	41.8-46.0	666	11.1	10.3-12.0	591	6.0	5.5-6.5	411	11.6	10.5-12.9	180	3.0	2.5-3.4
1987	2,374	22.6	21.7-23.6	1,703	39.8	37.9-41.8	671	11.0	10.1-11.8	544	5.4	5.0-5.9	363	9.8	8.8-10.9	181	2.9	2.5-3.4
1988	2,402	22.8	21.9-23.8	1,742	40.7	38.7-42.8	660	10.8	10.0-11.6	577	5.6	5.2-6.1	386	10.0	9.0-11.1	191	3.0	2.6-3.5
1989	2,468	23.1	22.2-24.1	1,803	41.7	39.7-43.7	665	10.7	9.9-11.5	563	5.5	5.0-5.9	403	10.7	9.6-11.8	160	2.5	2.1-2.9
1990	2,555	23.8	22.8-24.7	1,832	41.9	39.9-44.0	723	11.6	10.8-12.5	575	5.5	5.1-6.0	378	9.8	8.8-10.9	197	3.0	2.6-3.5
1991	2,517	23.0	22.1-24.0	1,827	41.0	39.1-43.0	690	10.8	10.0-11.7	568	5.3	4.9-5.8	392	9.9	8.9-11.0	176	2.7	2.3-3.1
1992	2,524	22.9	22.0-23.9	1,823	40.8	38.9-42.8	701	11.0	10.2-11.9	583	5.4	4.9-5.8	350	8.8	7.9-9.8	233	3.5	3.0-3.9
1993	2,688	24.1	23.2-25.0	1,879	41.0	39.2-43.0	809	12.5	11.6-13.3	586	5.3	4.9-5.8	390	9.8	8.8-10.8	196	2.9	2.5-3.3
1994	2,549	22.8	21.9-23.7	1,838	40.0	38.2-42.0	711	11.0	10.2-11.8	593	5.3	4.9-5.8	394	9.5	8.6-10.5	199	2.9	2.5-3.3
1995	2,714	24.1	23.2-25.0	1,911	41.1	39.3-43.1	803	12.3	11.4-13.2	563	5.0	4.6-5.4	372	8.8	7.9-9.8	191	2.8	2.4-3.2
1996	2,938	25.8	24.9-26.8	2,147	45.9	43.9-47.9	791	12.0	11.2-12.9	627	5.5	5.1-6.0	417	9.9	8.9-10.9	210	3.0	2.6-3.5
1997	3,068	26.7	25.8-27.7	2,208	46.6	44.6-48.6	860	12.9	12.0-13.8	641	5.6	5.1-6.0	427	9.7	8.8-10.7	214	3.0	2.6-3.5
1998	3,189	27.5	26.6-28.5	2,330	48.4	46.4-50.4	859	13.0	12.1-13.9	603	5.2	4.8-5.6	423	9.7	8.8-10.7	180	2.5	2.1-2.9
1999	3,217	27.5	26.5-28.4	2,326	47.6	45.7-49.6	891	13.2	12.3-14.1	610	5.2	4.8-5.6	419	9.4	8.5-10.4	191	2.6	2.2-3.0
2000	3,245	27.3	26.4-28.3	2,348	47.4	45.5-49.4	897	13.0	12.1-13.9	617	5.1	4.7-5.6	423	9.1	8.3-10.1	194	2.6	2.3-3.0
2001	3,244	27.0	26.1-28.0	2,377	47.7	45.7-49.6	867	12.5	11.7-13.4	632	5.2	4.8-5.6	442	9.6	8.7-10.5	190	2.5	2.2-2.9
2002	3,400	28.0	27.1-29.0	2,527	49.7	47.7-51.6	873	12.6	11.7-13.4	631	5.1	4.7-5.5	420	8.9	8.0-9.8	211	2.9	2.5-3.3
2003	3,468	28.3	27.3-29.2	2,560	49.2	47.3-51.2	908	12.9	12.1-13.8	589	4.7	4.3-5.1	396	8.2	7.4-9.1	193	2.5	2.2-2.9
2004	3,597	29.0	28.1-30.0	2,664	51.0	49.1-53.0	933	13.2	12.4-14.1	627	5.0	4.6-5.4	430	8.6	7.8-9.5	197	2.6	2.2-3.0
2005	3,630	29.0	28.1-30.0	2,675	50.5	48.6-52.5	955	13.3	12.5-14.2	592	4.6	4.2-5.0	398	8.0	7.2-8.8	194	2.5	2.1-2.9
2006	3,555	28.1	27.1-29.0	2,624	48.9	47.1-50.9	931	13.0	12.1-13.8	591	4.6	4.2-5.0	405	7.9	7.2-8.8	186	2.4	2.1-2.8
2007	3,546	27.7	26.8-28.6	2,618	47.8	46.0-49.7	928	12.9	12.1-13.8	677	5.1	4.8-5.6	476	9.1	8.3-10.0	201	2.5	2.2-2.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Urinary bladder (incl. in situ)

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	3,699	28.3	27.4-29.3	2,781	50.0	48.1-51.9	918	12.4	11.6-13.2	668	5.0	4.6-5.4	458	8.7	7.9-9.5	210	2.6	2.2-3.0
2009	3,747	28.3	27.4-29.2	2,803	49.5	47.6-51.4	944	12.5	11.7-13.4	674	5.0	4.6-5.4	508	9.4	8.6-10.3	166	2.0	1.7-2.4
2010	3,501	26.0	25.2-26.9	2,643	45.8	44.1-47.6	858	11.3	10.5-12.1	692	5.0	4.6-5.4	479	8.6	7.8-9.4	213	2.6	2.2-2.9
2011	3,817	28.2	27.3-29.1	2,854	48.5	46.7-50.3	963	12.8	12.0-13.6	648	4.6	4.2-5.0	456	8.1	7.4-8.9	192	2.3	2.0-2.6
2012	3,918	28.3	27.4-29.2	2,895	48.6	46.8-50.5	1,023	13.0	12.2-13.9	668	4.7	4.4-5.1	460	8.0	7.3-8.8	208	2.5	2.1-2.8
2013	3,951	28.3	27.4-29.2	2,949	49.0	47.2-50.8	1,002	12.7	11.9-13.6	680	4.7	4.4-5.1	489	8.4	7.6-9.1	191	2.2	1.9-2.6
2014	3,865	27.0	26.2-27.9	2,887	46.5	44.8-48.3	978	12.1	11.3-12.9	716	4.9	4.5-5.2	488	8.2	7.4-8.9	228	2.7	2.3-3.1
2015	3,771	26.1	25.3-27.0	2,806	44.4	42.8-46.1	965	12.1	11.3-12.9	669	4.5	4.2-4.9	460	7.5	6.8-8.2	209	2.5	2.1-2.8
2016	3,920	26.6	25.8-27.5	2,943	45.5	43.8-47.2	977	11.9	11.1-12.7	693	4.6	4.3-5.0	492	8.0	7.3-8.8	201	2.3	1.9-2.6
2017	3,878	26.0	25.1-26.8	2,915	44.6	43.0-46.3	963	11.5	10.8-12.3	673	4.4	4.0-4.7	463	7.4	6.7-8.1	210	2.3	2.0-2.6
2018	3,866	25.4	24.5-26.2	2,891	43.0	41.4-44.7	975	11.5	10.8-12.3	687	4.4	4.1-4.7	495	7.7	7.0-8.4	192	2.2	1.9-2.5
2019	3,880	25.1	24.3-25.9	2,909	42.5	40.9-44.1	971	11.4	10.6-12.1	624	3.9	3.6-4.3	451	6.9	6.3-7.6	173	1.9	1.6-2.2
2020	3,551	22.6	21.8-23.3	2,685	38.6	37.1-40.1	866	9.9	9.3-10.6	641	4.0	3.7-4.3	467	7.1	6.5-7.8	174	1.9	1.6-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State Excl New York City
Kidney and renal pelvis**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	684	7.2	6.6-7.7	426	10.3	9.3-11.4	258	4.8	4.2-5.4	367	3.9	3.5-4.3	229	5.6	4.9-6.4	138	2.5	2.1-3.0
1977	732	7.6	7.1-8.2	449	10.8	9.8-11.9	283	5.2	4.6-5.9	408	4.3	3.9-4.8	247	6.1	5.3-7.0	161	3.0	2.5-3.5
1978	742	7.7	7.1-8.3	456	11.0	9.9-12.1	286	5.3	4.7-5.9	370	3.8	3.4-4.2	226	5.5	4.8-6.3	144	2.6	2.2-3.0
1979	757	7.7	7.2-8.3	465	10.9	9.9-12.0	292	5.2	4.6-5.8	424	4.4	4.0-4.8	268	6.4	5.7-7.3	156	2.8	2.4-3.3
1980	761	7.7	7.1-8.3	459	10.7	9.7-11.8	302	5.4	4.8-6.1	417	4.2	3.8-4.6	248	6.1	5.3-6.9	169	2.9	2.5-3.4
1981	838	8.5	8.0-9.2	501	12.0	10.9-13.2	337	6.0	5.4-6.7	399	4.0	3.6-4.4	248	6.0	5.2-6.8	151	2.6	2.2-3.1
1982	877	8.6	8.1-9.2	532	12.0	10.9-13.1	345	6.1	5.4-6.7	410	4.0	3.6-4.5	243	5.7	5.0-6.5	167	2.9	2.4-3.3
1983	942	9.3	8.7-9.9	579	13.1	12.1-14.3	363	6.3	5.7-7.0	408	4.1	3.7-4.5	241	5.6	4.9-6.4	167	2.9	2.5-3.4
1984	942	9.3	8.7-9.9	594	13.6	12.5-14.8	348	6.1	5.4-6.7	412	4.1	3.7-4.5	256	6.3	5.5-7.1	156	2.6	2.2-3.1
1985	991	9.6	9.0-10.3	598	13.6	12.5-14.8	393	6.7	6.0-7.4	450	4.4	4.0-4.9	258	6.2	5.4-7.0	192	3.3	2.8-3.8
1986	999	9.7	9.1-10.4	614	13.8	12.7-15.0	385	6.7	6.1-7.4	472	4.6	4.2-5.0	282	6.5	5.8-7.4	190	3.2	2.7-3.7
1987	1,070	10.3	9.7-10.9	633	14.2	13.1-15.4	437	7.4	6.8-8.2	482	4.7	4.2-5.1	293	6.9	6.1-7.8	189	3.1	2.7-3.6
1988	1,044	9.9	9.3-10.5	617	13.7	12.6-14.9	427	7.2	6.5-7.9	464	4.4	4.0-4.8	263	6.0	5.3-6.8	201	3.2	2.8-3.7
1989	1,022	9.7	9.1-10.3	637	14.2	13.1-15.4	385	6.4	5.8-7.1	450	4.3	3.9-4.7	271	6.3	5.5-7.1	179	2.9	2.5-3.4
1990	1,170	10.9	10.3-11.6	731	16.0	14.8-17.2	439	7.3	6.6-8.0	438	4.1	3.7-4.5	267	5.9	5.2-6.7	171	2.7	2.3-3.2
1991	1,190	11.0	10.4-11.7	722	15.5	14.3-16.7	468	7.7	7.1-8.5	486	4.5	4.1-4.9	293	6.6	5.8-7.4	193	3.0	2.6-3.5
1992	1,216	11.1	10.5-11.8	737	15.5	14.4-16.7	479	7.9	7.2-8.6	448	4.1	3.7-4.5	273	6.1	5.4-6.9	175	2.7	2.3-3.2
1993	1,253	11.3	10.7-12.0	751	15.7	14.6-16.9	502	8.1	7.4-8.8	436	4.0	3.6-4.4	262	5.6	4.9-6.3	174	2.7	2.3-3.1
1994	1,259	11.3	10.7-12.0	742	15.4	14.3-16.6	517	8.4	7.7-9.1	492	4.4	4.0-4.8	306	6.8	6.0-7.6	186	2.9	2.5-3.3
1995	1,299	11.6	11.0-12.3	764	15.6	14.5-16.8	535	8.5	7.8-9.3	469	4.2	3.8-4.6	265	5.8	5.1-6.6	204	3.1	2.7-3.6
1996	1,366	12.2	11.5-12.8	849	17.3	16.1-18.5	517	8.2	7.5-8.9	496	4.4	4.0-4.8	286	6.1	5.4-6.9	210	3.2	2.8-3.7
1997	1,456	12.8	12.2-13.5	918	18.6	17.4-19.9	538	8.4	7.7-9.2	480	4.2	3.8-4.6	300	6.4	5.6-7.1	180	2.7	2.3-3.1
1998	1,553	13.6	12.9-14.3	950	18.7	17.5-19.9	603	9.5	8.7-10.3	477	4.1	3.8-4.5	298	6.0	5.4-6.8	179	2.6	2.2-3.0
1999	1,604	13.9	13.2-14.6	981	19.2	18.0-20.4	623	9.7	8.9-10.5	411	3.5	3.2-3.9	253	5.2	4.6-5.9	158	2.3	1.9-2.7
2000	1,605	13.7	13.0-14.4	948	18.3	17.2-19.5	657	10.1	9.3-10.9	470	4.0	3.6-4.3	282	5.7	5.1-6.5	188	2.7	2.3-3.1
2001	1,706	14.4	13.7-15.1	1,059	20.2	19.0-21.5	647	9.8	9.1-10.6	499	4.2	3.8-4.5	293	5.9	5.2-6.6	206	3.0	2.6-3.5
2002	1,743	14.5	13.9-15.2	1,085	20.5	19.3-21.8	658	9.9	9.1-10.7	449	3.7	3.4-4.1	267	5.3	4.7-6.0	182	2.6	2.2-3.0
2003	1,789	14.7	14.1-15.4	1,113	20.3	19.2-21.6	676	10.0	9.3-10.8	487	4.0	3.6-4.3	293	5.6	5.0-6.3	194	2.6	2.3-3.0
2004	1,997	16.3	15.6-17.0	1,238	22.4	21.2-23.7	759	11.3	10.5-12.2	496	4.0	3.6-4.4	296	5.6	5.0-6.3	200	2.8	2.4-3.2
2005	1,986	16.1	15.4-16.9	1,207	21.9	20.7-23.2	779	11.6	10.8-12.4	506	4.0	3.7-4.4	329	6.1	5.5-6.8	177	2.4	2.1-2.8
2006	2,181	17.4	16.7-18.2	1,390	24.6	23.3-26.0	791	11.5	10.7-12.4	498	3.9	3.5-4.2	304	5.7	5.1-6.4	194	2.6	2.3-3.1
2007	2,291	18.1	17.4-18.9	1,446	25.2	23.9-26.5	845	12.4	11.6-13.3	491	3.8	3.5-4.2	308	5.6	5.0-6.2	183	2.5	2.1-2.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Kidney and renal pelvis

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	2,237	17.4	16.7-18.2	1,398	23.9	22.7-25.2	839	12.0	11.2-12.9	497	3.8	3.4-4.1	327	5.8	5.2-6.5	170	2.2	1.9-2.6
2009	2,239	17.3	16.6-18.0	1,386	23.3	22.1-24.6	853	12.3	11.4-13.1	480	3.6	3.3-3.9	308	5.4	4.8-6.0	172	2.2	1.9-2.6
2010	2,293	17.3	16.6-18.1	1,436	23.8	22.6-25.1	857	12.0	11.2-12.8	466	3.4	3.1-3.8	293	4.9	4.4-5.5	173	2.2	1.9-2.6
2011	2,372	17.8	17.1-18.6	1,551	25.3	24.1-26.7	821	11.5	10.7-12.4	510	3.7	3.4-4.1	312	5.3	4.7-6.0	198	2.5	2.2-2.9
2012	2,446	18.1	17.3-18.8	1,549	24.9	23.6-26.2	897	12.3	11.5-13.1	531	3.8	3.5-4.1	350	5.8	5.2-6.4	181	2.3	1.9-2.6
2013	2,442	17.9	17.2-18.6	1,578	24.8	23.6-26.1	864	11.9	11.1-12.8	471	3.3	3.0-3.7	309	5.0	4.5-5.7	162	2.0	1.7-2.3
2014	2,550	18.3	17.6-19.1	1,650	25.4	24.2-26.7	900	12.2	11.4-13.1	463	3.2	2.9-3.5	304	4.7	4.2-5.3	159	1.9	1.6-2.3
2015	2,619	18.5	17.8-19.2	1,698	25.9	24.7-27.2	921	12.1	11.3-12.9	498	3.4	3.1-3.7	312	4.8	4.3-5.4	186	2.2	1.9-2.6
2016	2,591	18.4	17.6-19.1	1,727	26.2	24.9-27.5	864	11.5	10.7-12.4	436	2.9	2.6-3.2	288	4.3	3.8-4.9	148	1.8	1.5-2.1
2017	2,621	18.4	17.7-19.1	1,654	24.8	23.6-26.1	967	12.8	12.0-13.7	437	2.9	2.7-3.2	274	4.2	3.7-4.7	163	1.9	1.6-2.3
2018	2,799	19.5	18.7-20.2	1,785	26.3	25.0-27.6	1,014	13.6	12.7-14.5	475	3.1	2.8-3.4	297	4.4	3.9-5.0	178	2.1	1.8-2.4
2019	2,778	19.0	18.3-19.7	1,829	26.6	25.4-27.9	949	12.3	11.5-13.2	436	2.8	2.5-3.1	268	3.8	3.4-4.3	168	1.9	1.6-2.3
2020	2,486	16.9	16.2-17.6	1,661	24.1	22.9-25.4	825	10.5	9.8-11.3	428	2.8	2.5-3.1	278	4.0	3.6-4.6	150	1.7	1.4-2.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State Excl New York City
Brain and other central nervous system**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	586	5.8	5.3-6.3	340	7.2	6.4-8.1	246	4.6	4.0-5.2	432	4.3	3.9-4.8	232	5.0	4.3-5.7	200	3.7	3.2-4.3
1977	604	5.9	5.5-6.4	340	7.3	6.5-8.1	264	4.8	4.2-5.4	438	4.3	3.9-4.7	243	5.2	4.6-6.0	195	3.5	3.0-4.0
1978	567	5.6	5.1-6.1	293	6.3	5.6-7.1	274	5.1	4.5-5.7	409	4.0	3.6-4.4	225	4.8	4.2-5.5	184	3.3	2.8-3.8
1979	657	6.4	5.9-6.9	382	8.1	7.3-9.0	275	5.0	4.4-5.6	417	4.1	3.7-4.5	236	5.0	4.4-5.7	181	3.2	2.8-3.8
1980	651	6.3	5.8-6.8	380	7.8	7.0-8.7	271	5.0	4.4-5.6	465	4.5	4.1-5.0	282	6.1	5.4-6.9	183	3.3	2.8-3.8
1981	636	6.2	5.7-6.7	354	7.6	6.8-8.5	282	5.0	4.4-5.7	427	4.1	3.8-4.6	243	5.4	4.7-6.1	184	3.2	2.8-3.8
1982	677	6.6	6.1-7.1	347	7.5	6.7-8.3	330	5.9	5.3-6.6	444	4.3	3.9-4.8	228	5.0	4.3-5.7	216	3.8	3.3-4.4
1983	654	6.3	5.9-6.9	328	6.9	6.2-7.8	326	5.9	5.2-6.5	437	4.2	3.8-4.6	231	4.9	4.3-5.6	206	3.5	3.0-4.0
1984	726	6.9	6.4-7.4	397	8.4	7.6-9.3	329	5.8	5.2-6.4	448	4.3	3.9-4.7	246	5.4	4.7-6.1	202	3.5	3.0-4.0
1985	685	6.6	6.1-7.1	365	7.7	6.9-8.5	320	5.7	5.1-6.3	427	4.1	3.8-4.6	212	4.7	4.1-5.4	215	3.8	3.3-4.4
1986	712	6.8	6.3-7.3	386	8.1	7.3-9.0	326	5.7	5.1-6.3	469	4.5	4.1-4.9	255	5.5	4.8-6.2	214	3.6	3.1-4.2
1987	763	7.2	6.7-7.8	401	8.3	7.5-9.2	362	6.3	5.6-7.0	469	4.4	4.0-4.9	238	5.1	4.4-5.8	231	3.9	3.4-4.4
1988	751	7.0	6.5-7.6	420	8.6	7.8-9.5	331	5.8	5.1-6.4	502	4.6	4.2-5.0	281	5.8	5.1-6.5	221	3.6	3.2-4.2
1989	738	6.9	6.4-7.4	418	8.4	7.6-9.3	320	5.6	5.0-6.3	468	4.4	4.0-4.8	260	5.4	4.8-6.2	208	3.5	3.0-4.0
1990	802	7.5	6.9-8.0	441	9.1	8.3-10.1	361	6.2	5.5-6.9	432	4.0	3.6-4.4	256	5.4	4.7-6.1	176	2.9	2.4-3.3
1991	776	7.2	6.7-7.7	425	8.8	7.9-9.7	351	6.0	5.4-6.7	463	4.3	3.9-4.7	271	5.8	5.1-6.6	192	3.3	2.8-3.8
1992	787	7.2	6.7-7.7	411	8.4	7.6-9.2	376	6.4	5.8-7.1	496	4.5	4.1-4.9	250	5.1	4.5-5.8	246	4.1	3.6-4.6
1993	786	7.2	6.7-7.7	438	8.7	7.9-9.6	348	5.9	5.3-6.5	484	4.4	4.0-4.8	261	5.3	4.7-6.0	223	3.7	3.2-4.2
1994	846	7.7	7.1-8.2	474	9.5	8.6-10.4	372	6.1	5.5-6.8	545	4.9	4.5-5.3	302	6.2	5.5-6.9	243	3.9	3.4-4.4
1995	900	8.1	7.6-8.6	445	8.7	7.9-9.6	455	7.5	6.9-8.3	484	4.4	4.0-4.8	262	5.3	4.6-6.0	222	3.6	3.1-4.1
1996	885	8.0	7.5-8.5	484	9.6	8.8-10.5	401	6.7	6.1-7.4	537	4.8	4.4-5.2	300	6.0	5.4-6.8	237	3.8	3.3-4.3
1997	831	7.4	6.9-8.0	469	9.1	8.3-10.0	362	6.0	5.4-6.6	504	4.5	4.1-4.9	280	5.5	4.9-6.2	224	3.6	3.1-4.1
1998	874	7.8	7.3-8.3	493	9.4	8.6-10.3	381	6.3	5.7-7.0	493	4.3	4.0-4.7	283	5.5	4.9-6.2	210	3.3	2.9-3.8
1999	832	7.3	6.9-7.9	448	8.6	7.8-9.4	384	6.3	5.7-6.9	500	4.4	4.0-4.8	279	5.4	4.8-6.1	221	3.5	3.0-4.0
2000	923	8.1	7.5-8.6	479	9.1	8.3-9.9	444	7.1	6.5-7.8	489	4.2	3.8-4.6	257	4.9	4.3-5.5	232	3.5	3.1-4.0
2001	880	7.7	7.2-8.2	472	8.9	8.1-9.8	408	6.5	5.9-7.2	486	4.1	3.8-4.5	252	4.8	4.2-5.4	234	3.6	3.1-4.1
2002	884	7.6	7.1-8.2	496	9.3	8.5-10.2	388	6.2	5.6-6.9	531	4.5	4.1-4.9	299	5.6	5.0-6.3	232	3.6	3.1-4.1
2003	899	7.7	7.2-8.3	465	8.6	7.8-9.4	434	7.1	6.4-7.8	486	4.1	3.7-4.5	274	5.0	4.4-5.7	212	3.3	2.9-3.8
2004	863	7.4	6.9-7.9	478	8.8	8.0-9.7	385	6.1	5.5-6.8	533	4.4	4.0-4.8	282	5.1	4.5-5.8	251	3.8	3.3-4.3
2005	881	7.5	7.0-8.0	510	9.4	8.6-10.2	371	5.9	5.3-6.5	535	4.4	4.1-4.8	295	5.4	4.8-6.0	240	3.6	3.2-4.1
2006	862	7.2	6.8-7.8	497	8.9	8.1-9.7	365	5.8	5.2-6.4	492	4.0	3.6-4.4	300	5.3	4.8-6.0	192	2.8	2.4-3.2
2007	850	7.1	6.7-7.7	459	8.3	7.6-9.1	391	6.2	5.6-6.9	506	4.0	3.7-4.4	303	5.4	4.8-6.0	203	3.0	2.6-3.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State Excl New York City
Brain and other central nervous system**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	871	7.2	6.8-7.8	481	8.6	7.9-9.5	390	6.1	5.5-6.7	510	4.0	3.7-4.4	295	5.2	4.6-5.8	215	3.1	2.7-3.5
2009	898	7.4	6.9-7.9	481	8.5	7.7-9.3	417	6.4	5.8-7.1	559	4.3	4.0-4.7	309	5.3	4.7-5.9	250	3.6	3.1-4.1
2010	878	7.2	6.7-7.7	487	8.5	7.8-9.3	391	6.0	5.4-6.7	550	4.3	3.9-4.7	303	5.2	4.6-5.9	247	3.5	3.1-4.0
2011	870	7.0	6.6-7.5	486	8.3	7.6-9.1	384	5.9	5.3-6.6	504	3.8	3.4-4.1	278	4.6	4.1-5.2	226	3.1	2.7-3.5
2012	911	7.2	6.8-7.7	504	8.5	7.7-9.3	407	6.2	5.6-6.9	603	4.5	4.2-4.9	338	5.6	5.0-6.3	265	3.7	3.2-4.2
2013	895	7.1	6.6-7.6	512	8.5	7.7-9.3	383	5.8	5.2-6.4	567	4.1	3.8-4.5	314	5.0	4.4-5.6	253	3.4	3.0-3.9
2014	965	7.6	7.1-8.1	526	8.7	8.0-9.5	439	6.7	6.0-7.3	585	4.3	3.9-4.6	328	5.1	4.6-5.7	257	3.5	3.1-4.0
2015	913	7.2	6.7-7.7	487	8.0	7.3-8.8	426	6.4	5.8-7.1	582	4.2	3.9-4.6	300	4.7	4.2-5.3	282	3.8	3.3-4.3
2016	886	6.8	6.3-7.3	516	8.3	7.6-9.1	370	5.4	4.9-6.1	608	4.4	4.0-4.7	353	5.4	4.9-6.1	255	3.4	3.0-3.9
2017	967	7.5	7.0-8.0	543	8.8	8.1-9.7	424	6.3	5.7-7.0	614	4.3	4.0-4.7	355	5.3	4.8-5.9	259	3.4	3.0-3.9
2018	888	6.8	6.4-7.3	498	8.0	7.2-8.7	390	5.8	5.2-6.5	625	4.4	4.1-4.8	348	5.3	4.7-5.9	277	3.7	3.2-4.2
2019	844	6.6	6.1-7.0	465	7.7	7.0-8.5	379	5.5	5.0-6.2	594	4.1	3.7-4.4	343	5.1	4.5-5.7	251	3.2	2.8-3.6
2020	871	6.6	6.2-7.1	477	7.5	6.8-8.2	394	5.8	5.2-6.5	609	4.2	3.9-4.6	349	5.2	4.7-5.8	260	3.3	2.9-3.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Thyroid

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	329	3.5	3.1-3.9	112	2.6	2.1-3.2	217	4.3	3.8-5.0	55	0.6	0.5-0.8	22	0.6	0.4-0.9	33	0.6	0.4-0.9
1977	366	3.8	3.5-4.3	112	2.5	2.1-3.1	254	5.0	4.4-5.7	69	0.8	0.6-1.0	22	0.6	0.4-0.9	47	0.9	0.6-1.2
1978	349	3.6	3.2-4.0	100	2.3	1.9-2.8	249	4.8	4.2-5.5	62	0.7	0.5-0.9	22	0.6	0.4-0.9	40	0.7	0.5-1.0
1979	333	3.4	3.0-3.8	96	2.2	1.8-2.7	237	4.5	3.9-5.1	46	0.5	0.3-0.6	13	0.3	0.2-0.6	33	0.6	0.4-0.8
1980	356	3.5	3.2-3.9	109	2.4	1.9-2.9	247	4.6	4.0-5.2	70	0.7	0.6-0.9	33	0.8	0.6-1.2	37	0.7	0.5-0.9
1981	350	3.5	3.1-3.9	105	2.4	2.0-2.9	245	4.6	4.0-5.2	56	0.6	0.4-0.8	25	0.6	0.4-0.9	31	0.6	0.4-0.8
1982	338	3.4	3.0-3.8	103	2.2	1.8-2.7	235	4.4	3.9-5.1	53	0.5	0.4-0.7	27	0.7	0.4-1.0	26	0.5	0.3-0.7
1983	403	4.0	3.6-4.4	104	2.2	1.8-2.7	299	5.5	4.9-6.2	53	0.6	0.4-0.7	14	0.4	0.2-0.6	39	0.7	0.5-0.9
1984	409	4.0	3.6-4.4	117	2.6	2.1-3.1	292	5.4	4.7-6.0	64	0.7	0.5-0.8	24	0.6	0.4-1.0	40	0.7	0.5-0.9
1985	418	4.1	3.7-4.5	113	2.4	1.9-2.9	305	5.7	5.1-6.4	46	0.5	0.3-0.6	19	0.5	0.3-0.8	27	0.5	0.3-0.7
1986	422	4.1	3.7-4.5	123	2.6	2.1-3.1	299	5.5	4.9-6.2	55	0.6	0.4-0.7	25	0.6	0.4-1.0	30	0.5	0.3-0.7
1987	423	4.1	3.7-4.5	137	2.8	2.4-3.4	286	5.3	4.7-5.9	54	0.5	0.4-0.7	16	0.3	0.2-0.5	38	0.6	0.4-0.9
1988	406	3.9	3.6-4.3	105	2.2	1.8-2.7	301	5.5	4.9-6.2	51	0.5	0.4-0.7	20	0.5	0.3-0.7	31	0.5	0.3-0.7
1989	433	4.1	3.7-4.5	118	2.4	2.0-2.9	315	5.7	5.1-6.4	57	0.5	0.4-0.7	17	0.4	0.2-0.6	40	0.6	0.5-0.9
1990	451	4.3	3.9-4.7	134	2.8	2.3-3.3	317	5.7	5.0-6.3	49	0.4	0.3-0.6	21	0.4	0.3-0.7	28	0.4	0.3-0.6
1991	514	4.8	4.4-5.2	150	3.0	2.6-3.6	364	6.5	5.8-7.2	46	0.4	0.3-0.6	16	0.4	0.2-0.6	30	0.5	0.3-0.7
1992	523	4.8	4.4-5.2	148	3.0	2.5-3.5	375	6.5	5.9-7.2	57	0.5	0.4-0.7	17	0.4	0.2-0.6	40	0.6	0.4-0.8
1993	533	4.9	4.5-5.3	143	2.8	2.3-3.3	390	6.9	6.2-7.7	60	0.5	0.4-0.7	23	0.5	0.3-0.8	37	0.6	0.4-0.8
1994	590	5.4	5.0-5.9	152	3.0	2.6-3.6	438	7.7	7.0-8.5	52	0.5	0.4-0.6	23	0.5	0.3-0.8	29	0.4	0.3-0.6
1995	662	6.0	5.6-6.5	176	3.5	3.0-4.1	486	8.5	7.8-9.3	58	0.5	0.4-0.7	25	0.5	0.3-0.8	33	0.5	0.3-0.7
1996	730	6.6	6.2-7.1	193	3.7	3.2-4.3	537	9.5	8.7-10.3	84	0.8	0.6-0.9	29	0.7	0.4-0.9	55	0.8	0.6-1.1
1997	794	7.2	6.7-7.7	212	4.1	3.5-4.6	582	10.2	9.4-11.0	61	0.5	0.4-0.7	27	0.6	0.4-0.9	34	0.5	0.3-0.7
1998	810	7.3	6.8-7.8	224	4.2	3.7-4.8	586	10.2	9.4-11.1	57	0.5	0.4-0.6	18	0.4	0.2-0.6	39	0.5	0.4-0.8
1999	953	8.6	8.0-9.1	227	4.3	3.7-4.8	726	12.7	11.8-13.7	60	0.5	0.4-0.7	21	0.4	0.3-0.7	39	0.5	0.4-0.7
2000	1,120	10.0	9.4-10.6	283	5.3	4.7-6.0	837	14.5	13.6-15.6	51	0.4	0.3-0.6	17	0.3	0.2-0.6	34	0.5	0.3-0.7
2001	1,139	10.1	9.5-10.7	246	4.5	4.0-5.1	893	15.5	14.5-16.5	57	0.5	0.4-0.6	23	0.4	0.3-0.7	34	0.5	0.3-0.7
2002	1,167	10.2	9.7-10.9	296	5.4	4.8-6.0	871	15.0	14.0-16.0	51	0.4	0.3-0.6	20	0.4	0.2-0.6	31	0.4	0.3-0.6
2003	1,231	10.7	10.1-11.3	330	5.9	5.3-6.6	901	15.3	14.3-16.4	59	0.5	0.4-0.6	15	0.3	0.2-0.5	44	0.6	0.4-0.8
2004	1,362	11.8	11.2-12.4	324	5.7	5.1-6.4	1,038	17.6	16.6-18.8	53	0.4	0.3-0.6	22	0.4	0.3-0.7	31	0.4	0.3-0.6
2005	1,514	13.1	12.4-13.8	374	6.6	5.9-7.3	1,140	19.4	18.3-20.6	53	0.4	0.3-0.5	29	0.5	0.4-0.8	24	0.3	0.2-0.5
2006	1,640	14.2	13.5-14.9	414	7.4	6.7-8.1	1,226	20.7	19.6-22.0	60	0.5	0.4-0.6	28	0.5	0.3-0.7	32	0.4	0.3-0.6
2007	1,803	15.4	14.7-16.1	466	8.3	7.5-9.1	1,337	22.3	21.1-23.6	58	0.4	0.3-0.6	28	0.5	0.3-0.7	30	0.4	0.3-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Thyroid

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	1,999	17.1	16.3-17.8	469	8.2	7.4-9.0	1,530	25.7	24.4-27.0	63	0.5	0.4-0.6	28	0.5	0.3-0.7	35	0.5	0.3-0.7
2009	2,079	17.7	17.0-18.5	525	9.2	8.4-10.0	1,554	26.0	24.7-27.4	73	0.5	0.4-0.7	27	0.5	0.3-0.7	46	0.6	0.4-0.8
2010	2,130	17.9	17.2-18.7	538	9.1	8.4-10.0	1,592	26.5	25.2-27.9	66	0.5	0.4-0.6	29	0.5	0.3-0.7	37	0.4	0.3-0.6
2011	2,291	19.2	18.4-20.1	620	10.4	9.6-11.3	1,671	27.7	26.4-29.1	80	0.6	0.5-0.7	39	0.7	0.5-0.9	41	0.5	0.4-0.7
2012	2,301	19.3	18.5-20.1	602	10.0	9.2-10.9	1,699	28.4	27.0-29.9	80	0.6	0.4-0.7	34	0.6	0.4-0.8	46	0.6	0.4-0.8
2013	2,375	19.8	19.0-20.6	636	10.5	9.7-11.4	1,739	28.9	27.5-30.3	74	0.5	0.4-0.7	29	0.5	0.3-0.7	45	0.5	0.4-0.7
2014	2,370	19.9	19.0-20.7	631	10.4	9.6-11.3	1,739	29.2	27.8-30.7	55	0.4	0.3-0.5	27	0.4	0.3-0.6	28	0.3	0.2-0.5
2015	2,409	20.3	19.4-21.1	640	10.6	9.8-11.5	1,769	29.8	28.4-31.3	76	0.5	0.4-0.7	36	0.6	0.4-0.8	40	0.5	0.3-0.7
2016	2,187	18.5	17.7-19.3	604	10.0	9.2-10.9	1,583	26.8	25.4-28.2	80	0.5	0.4-0.7	32	0.5	0.3-0.7	48	0.6	0.4-0.8
2017	2,122	17.9	17.1-18.7	566	9.4	8.6-10.3	1,556	26.3	24.9-27.7	73	0.5	0.4-0.6	38	0.6	0.4-0.8	35	0.4	0.3-0.6
2018	2,078	17.4	16.6-18.2	623	10.3	9.5-11.2	1,455	24.5	23.2-25.8	76	0.5	0.4-0.6	39	0.6	0.4-0.8	37	0.4	0.3-0.6
2019	2,259	19.0	18.2-19.8	658	10.6	9.8-11.5	1,601	27.3	25.9-28.7	84	0.6	0.4-0.7	38	0.6	0.4-0.8	46	0.5	0.4-0.7
2020	1,790	14.9	14.2-15.7	519	8.3	7.6-9.1	1,271	21.5	20.3-22.8	72	0.5	0.4-0.6	25	0.4	0.3-0.6	47	0.6	0.4-0.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State Excl New York City
Hodgkin lymphoma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	371	3.5	3.1-3.8	200	3.9	3.4-4.5	171	3.0	2.6-3.5	142	1.4	1.2-1.7	76	1.7	1.3-2.1	66	1.2	0.9-1.5
1977	341	3.2	2.9-3.6	188	3.7	3.2-4.3	153	2.7	2.3-3.2	141	1.4	1.2-1.7	82	1.9	1.5-2.4	59	1.1	0.8-1.4
1978	373	3.4	3.1-3.8	200	4.0	3.4-4.6	173	3.0	2.5-3.5	121	1.2	1.0-1.4	68	1.5	1.2-2.0	53	1.0	0.7-1.3
1979	388	3.6	3.2-4.0	220	4.4	3.8-5.1	168	2.9	2.5-3.4	112	1.1	0.9-1.3	63	1.4	1.0-1.8	49	0.9	0.6-1.2
1980	418	3.8	3.5-4.2	235	4.6	4.0-5.3	183	3.1	2.7-3.6	135	1.3	1.1-1.6	79	1.7	1.4-2.2	56	1.0	0.7-1.3
1981	397	3.6	3.3-4.0	212	4.1	3.5-4.7	185	3.2	2.8-3.8	117	1.1	0.9-1.4	69	1.5	1.1-1.9	48	0.8	0.6-1.1
1982	396	3.6	3.3-4.0	193	3.9	3.3-4.5	203	3.4	3.0-3.9	134	1.3	1.1-1.6	73	1.6	1.2-2.0	61	1.1	0.8-1.4
1983	378	3.5	3.1-3.9	214	4.2	3.6-4.8	164	2.8	2.4-3.3	93	0.9	0.7-1.1	55	1.1	0.8-1.4	38	0.6	0.5-0.9
1984	401	3.7	3.3-4.1	212	4.3	3.7-4.9	189	3.2	2.8-3.7	97	0.9	0.8-1.2	62	1.3	1.0-1.7	35	0.6	0.4-0.8
1985	416	3.8	3.4-4.1	213	4.1	3.5-4.7	203	3.5	3.0-4.0	103	1.0	0.8-1.2	56	1.1	0.9-1.5	47	0.8	0.6-1.1
1986	378	3.4	3.0-3.7	195	3.6	3.1-4.2	183	3.1	2.7-3.6	127	1.2	1.0-1.4	69	1.4	1.1-1.8	58	1.0	0.7-1.2
1987	374	3.4	3.0-3.7	190	3.6	3.1-4.2	184	3.2	2.7-3.7	105	1.0	0.8-1.2	65	1.3	1.0-1.7	40	0.7	0.5-1.0
1988	381	3.5	3.1-3.8	214	4.1	3.5-4.7	167	2.9	2.5-3.4	87	0.8	0.7-1.0	59	1.2	0.9-1.5	28	0.5	0.3-0.7
1989	400	3.6	3.2-3.9	220	4.2	3.6-4.8	180	3.0	2.6-3.5	96	0.9	0.7-1.1	64	1.4	1.0-1.7	32	0.6	0.4-0.8
1990	392	3.5	3.2-3.9	204	3.9	3.4-4.5	188	3.2	2.8-3.8	90	0.8	0.7-1.0	50	1.0	0.8-1.4	40	0.7	0.5-0.9
1991	382	3.4	3.1-3.8	207	3.8	3.3-4.4	175	3.0	2.6-3.5	76	0.7	0.5-0.9	40	0.8	0.6-1.1	36	0.6	0.4-0.9
1992	395	3.6	3.2-4.0	211	4.1	3.5-4.7	184	3.2	2.7-3.7	95	0.9	0.7-1.1	46	0.9	0.7-1.2	49	0.8	0.6-1.1
1993	380	3.5	3.1-3.8	190	3.6	3.1-4.1	190	3.3	2.9-3.9	73	0.7	0.5-0.8	40	0.8	0.6-1.1	33	0.6	0.4-0.8
1994	414	3.7	3.4-4.1	205	3.9	3.4-4.5	209	3.6	3.1-4.2	77	0.7	0.6-0.9	41	0.8	0.6-1.1	36	0.6	0.4-0.8
1995	363	3.3	3.0-3.7	201	3.8	3.3-4.3	162	2.9	2.4-3.3	78	0.7	0.6-0.9	41	0.8	0.6-1.1	37	0.6	0.4-0.8
1996	375	3.4	3.1-3.8	187	3.6	3.1-4.1	188	3.4	2.9-3.9	61	0.6	0.4-0.7	31	0.6	0.4-0.9	30	0.5	0.4-0.8
1997	341	3.1	2.8-3.5	190	3.6	3.1-4.1	151	2.7	2.3-3.2	63	0.6	0.4-0.7	37	0.7	0.5-1.0	26	0.4	0.3-0.6
1998	388	3.6	3.2-3.9	198	3.8	3.2-4.3	190	3.4	2.9-3.9	60	0.5	0.4-0.7	31	0.6	0.4-0.8	29	0.5	0.3-0.7
1999	372	3.4	3.1-3.8	207	3.9	3.4-4.5	165	2.9	2.5-3.4	66	0.6	0.5-0.7	31	0.6	0.4-0.8	35	0.6	0.4-0.8
2000	372	3.4	3.1-3.8	206	3.9	3.4-4.5	166	2.9	2.5-3.4	50	0.4	0.3-0.6	25	0.5	0.3-0.7	25	0.4	0.3-0.6
2001	400	3.7	3.3-4.0	210	3.9	3.4-4.5	190	3.5	3.0-4.0	65	0.6	0.4-0.7	34	0.7	0.5-0.9	31	0.5	0.3-0.7
2002	420	3.8	3.5-4.2	243	4.6	4.0-5.2	177	3.2	2.7-3.7	68	0.6	0.5-0.7	30	0.6	0.4-0.8	38	0.6	0.4-0.9
2003	359	3.2	2.9-3.6	201	3.7	3.2-4.3	158	2.8	2.4-3.2	47	0.4	0.3-0.6	22	0.4	0.3-0.6	25	0.4	0.3-0.6
2004	404	3.6	3.3-4.0	206	3.8	3.3-4.3	198	3.5	3.0-4.1	55	0.5	0.3-0.6	34	0.6	0.4-0.9	21	0.3	0.2-0.5
2005	403	3.6	3.3-4.0	217	4.0	3.5-4.5	186	3.3	2.9-3.9	53	0.5	0.3-0.6	25	0.5	0.3-0.7	28	0.4	0.3-0.7
2006	376	3.3	3.0-3.7	208	3.8	3.3-4.3	168	2.9	2.5-3.4	61	0.5	0.4-0.7	33	0.6	0.4-0.8	28	0.4	0.3-0.6
2007	381	3.4	3.1-3.7	207	3.8	3.3-4.3	174	3.1	2.7-3.6	58	0.5	0.4-0.6	28	0.5	0.3-0.7	30	0.4	0.3-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Hodgkin lymphoma

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	381	3.4	3.0-3.7	215	3.9	3.4-4.5	166	2.9	2.5-3.4	55	0.5	0.3-0.6	30	0.6	0.4-0.8	25	0.4	0.2-0.6
2009	391	3.5	3.1-3.9	214	3.9	3.4-4.4	177	3.2	2.7-3.7	44	0.4	0.3-0.5	22	0.4	0.2-0.6	22	0.3	0.2-0.5
2010	380	3.3	3.0-3.6	218	3.9	3.4-4.4	162	2.8	2.3-3.2	45	0.4	0.3-0.5	24	0.4	0.3-0.6	21	0.3	0.2-0.5
2011	381	3.4	3.0-3.7	215	3.9	3.3-4.4	166	2.9	2.5-3.4	45	0.3	0.2-0.5	31	0.5	0.4-0.8	14	0.2	0.1-0.3
2012	403	3.6	3.3-4.0	242	4.4	3.9-5.0	161	2.9	2.4-3.4	49	0.4	0.3-0.5	22	0.4	0.2-0.6	27	0.4	0.3-0.6
2013	363	3.2	2.8-3.5	190	3.3	2.9-3.9	173	3.0	2.6-3.5	36	0.3	0.2-0.4	15	0.3	0.1-0.5	21	0.3	0.2-0.5
2014	409	3.5	3.2-3.9	213	3.8	3.3-4.3	196	3.3	2.8-3.8	35	0.3	0.2-0.4	16	0.3	0.1-0.4	19	0.2	0.1-0.4
2015	349	3.1	2.8-3.4	194	3.4	3.0-4.0	155	2.8	2.3-3.2	34	0.3	0.2-0.4	17	0.3	0.2-0.5	17	0.2	0.1-0.4
2016	386	3.3	3.0-3.7	191	3.4	2.9-3.9	195	3.3	2.9-3.8	42	0.3	0.2-0.4	25	0.4	0.3-0.6	17	0.2	0.1-0.4
2017	381	3.3	2.9-3.6	207	3.6	3.1-4.2	174	3.0	2.5-3.4	39	0.3	0.2-0.4	22	0.3	0.2-0.5	17	0.2	0.1-0.3
2018	354	3.0	2.6-3.3	205	3.5	3.0-4.0	149	2.5	2.1-2.9	47	0.3	0.2-0.5	29	0.5	0.3-0.7	18	0.2	0.1-0.4
2019	393	3.4	3.0-3.7	209	3.6	3.1-4.1	184	3.2	2.8-3.8	42	0.3	0.2-0.4	28	0.4	0.3-0.6	14	0.2	0.1-0.3
2020	329	2.9	2.6-3.2	184	3.3	2.8-3.8	145	2.5	2.1-3.0	28	0.2	0.1-0.3	19	0.3	0.2-0.4	9	0.1	0.1-0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State Excl New York City
Non-Hodgkin lymphomas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,062	11.2	10.5-11.9	552	13.3	12.1-14.5	510	9.6	8.8-10.5	587	6.2	5.7-6.8	309	7.6	6.8-8.6	278	5.2	4.6-5.9
1977	1,155	11.9	11.2-12.6	597	14.3	13.1-15.5	558	10.2	9.4-11.1	642	6.7	6.2-7.2	320	7.8	6.9-8.7	322	5.9	5.2-6.6
1978	1,273	13.0	12.3-13.8	675	15.7	14.5-17.0	598	10.9	10.0-11.8	631	6.6	6.1-7.2	320	8.1	7.1-9.0	311	5.7	5.0-6.3
1979	1,163	11.9	11.2-12.6	586	13.8	12.7-15.1	577	10.4	9.6-11.3	659	6.8	6.3-7.4	358	8.6	7.7-9.6	301	5.4	4.8-6.0
1980	1,246	12.6	11.9-13.4	647	15.0	13.8-16.2	599	10.7	9.9-11.6	664	6.8	6.3-7.3	322	7.7	6.8-8.6	342	6.1	5.5-6.8
1981	1,274	12.8	12.1-13.5	634	14.8	13.7-16.1	640	11.2	10.4-12.2	660	6.7	6.2-7.2	342	8.3	7.4-9.2	318	5.5	4.9-6.2
1982	1,304	13.0	12.3-13.7	684	15.8	14.6-17.1	620	10.9	10.1-11.8	661	6.7	6.2-7.2	344	8.4	7.5-9.4	317	5.4	4.9-6.1
1983	1,417	13.8	13.1-14.6	703	15.9	14.7-17.2	714	12.2	11.3-13.1	704	7.0	6.5-7.5	370	8.7	7.8-9.6	334	5.8	5.2-6.5
1984	1,461	14.4	13.7-15.2	766	17.9	16.6-19.3	695	11.9	11.0-12.8	727	7.2	6.6-7.7	395	9.7	8.7-10.7	332	5.6	5.0-6.2
1985	1,539	15.0	14.2-15.8	816	18.3	17.0-19.6	723	12.3	11.4-13.2	767	7.5	7.0-8.1	403	9.4	8.5-10.4	364	6.1	5.5-6.8
1986	1,604	15.5	14.7-16.3	862	19.5	18.1-20.9	742	12.6	11.7-13.5	811	8.0	7.4-8.5	445	10.6	9.6-11.6	366	6.0	5.4-6.7
1987	1,578	15.1	14.4-15.9	841	18.7	17.4-20.0	737	12.3	11.4-13.3	802	7.8	7.2-8.3	398	9.2	8.3-10.2	404	6.7	6.0-7.4
1988	1,706	16.3	15.6-17.2	880	19.5	18.1-20.9	826	14.0	13.1-15.0	827	8.0	7.4-8.5	419	9.8	8.9-10.8	408	6.7	6.1-7.4
1989	1,789	16.9	16.1-17.7	939	20.3	18.9-21.6	850	14.3	13.3-15.3	856	8.1	7.6-8.7	459	10.4	9.5-11.5	397	6.4	5.8-7.1
1990	1,886	17.6	16.8-18.4	1,009	21.6	20.3-23.1	877	14.3	13.4-15.3	897	8.5	7.9-9.1	460	10.5	9.5-11.6	437	7.0	6.3-7.7
1991	1,869	17.3	16.6-18.1	981	21.4	20.0-22.8	888	14.5	13.6-15.5	905	8.4	7.8-8.9	473	10.8	9.8-11.8	432	6.7	6.1-7.4
1992	1,866	17.1	16.4-18.0	992	20.9	19.6-22.3	874	14.1	13.2-15.1	955	8.8	8.2-9.4	489	11.0	10.0-12.1	466	7.3	6.6-8.0
1993	1,989	18.2	17.4-19.0	1,053	21.8	20.5-23.2	936	15.1	14.1-16.1	893	8.1	7.6-8.6	443	9.7	8.8-10.7	450	6.9	6.2-7.5
1994	2,082	18.8	18.0-19.6	1,124	23.3	21.9-24.7	958	15.1	14.2-16.1	994	8.9	8.4-9.5	513	11.3	10.3-12.3	481	7.3	6.7-8.0
1995	2,103	18.9	18.1-19.7	1,107	22.9	21.6-24.3	996	15.8	14.8-16.8	1,021	9.1	8.6-9.7	532	11.7	10.7-12.7	489	7.4	6.7-8.1
1996	2,288	20.4	19.5-21.2	1,209	24.7	23.3-26.2	1,079	17.0	16.0-18.0	1,027	9.1	8.5-9.7	531	11.5	10.5-12.5	496	7.4	6.8-8.1
1997	2,323	20.5	19.7-21.3	1,202	24.3	22.9-25.7	1,121	17.5	16.5-18.6	1,066	9.3	8.8-9.9	538	11.3	10.3-12.3	528	7.9	7.2-8.6
1998	2,387	20.9	20.0-21.7	1,212	24.2	22.9-25.6	1,175	18.2	17.2-19.3	990	8.5	8.0-9.1	479	9.9	9.0-10.8	511	7.5	6.9-8.2
1999	2,437	21.1	20.2-21.9	1,229	24.3	22.9-25.7	1,208	18.6	17.5-19.7	952	8.1	7.6-8.7	465	9.7	8.8-10.7	487	7.1	6.5-7.8
2000	2,441	20.8	20.0-21.7	1,250	24.5	23.1-25.9	1,191	18.0	17.0-19.1	1,014	8.5	8.0-9.1	510	10.4	9.5-11.4	504	7.2	6.6-7.9
2001	2,448	20.7	19.9-21.5	1,251	24.3	22.9-25.7	1,197	18.0	16.9-19.0	927	7.7	7.2-8.2	499	9.9	9.1-10.8	428	6.0	5.4-6.6
2002	2,520	21.1	20.3-21.9	1,336	25.5	24.2-27.0	1,184	17.7	16.7-18.8	937	7.7	7.2-8.2	505	10.1	9.2-11.0	432	6.1	5.5-6.7
2003	2,616	21.6	20.8-22.5	1,332	25.0	23.7-26.4	1,284	18.9	17.9-20.0	842	6.8	6.4-7.3	426	8.5	7.7-9.4	416	5.7	5.1-6.3
2004	2,661	21.8	21.0-22.7	1,424	26.5	25.1-27.9	1,237	18.3	17.3-19.3	839	6.7	6.3-7.2	410	7.9	7.2-8.7	429	5.8	5.2-6.3
2005	2,743	22.3	21.4-23.1	1,475	27.1	25.7-28.5	1,268	18.4	17.4-19.5	807	6.4	5.9-6.8	430	8.3	7.5-9.1	377	5.0	4.5-5.6
2006	2,874	23.1	22.3-24.0	1,482	26.9	25.6-28.4	1,392	20.0	18.9-21.1	893	7.0	6.6-7.5	454	8.5	7.7-9.4	439	5.8	5.2-6.3
2007	2,888	22.9	22.0-23.7	1,553	27.8	26.4-29.2	1,335	19.1	18.0-20.2	794	6.1	5.7-6.5	437	8.1	7.3-8.9	357	4.7	4.2-5.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Non-Hodgkin lymphomas

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	2,934	23.1	22.3-24.0	1,564	27.7	26.3-29.1	1,370	19.5	18.5-20.6	861	6.6	6.1-7.0	450	8.2	7.5-9.0	411	5.4	4.9-6.0
2009	2,987	23.1	22.3-24.0	1,615	28.0	26.6-29.4	1,372	19.3	18.3-20.4	818	6.2	5.7-6.6	459	8.2	7.5-9.0	359	4.7	4.2-5.2
2010	3,007	22.9	22.1-23.7	1,596	27.3	26.0-28.7	1,411	19.3	18.2-20.3	855	6.3	5.9-6.8	450	7.9	7.2-8.7	405	5.2	4.7-5.8
2011	3,001	22.7	21.9-23.6	1,659	28.0	26.6-29.4	1,342	18.6	17.6-19.6	841	6.2	5.8-6.6	451	7.8	7.1-8.6	390	5.0	4.5-5.5
2012	2,996	22.2	21.4-23.1	1,615	26.9	25.6-28.3	1,381	18.6	17.6-19.6	803	5.9	5.5-6.3	440	7.6	6.9-8.4	363	4.5	4.0-5.0
2013	3,069	22.6	21.8-23.4	1,670	27.4	26.0-28.7	1,399	18.8	17.8-19.8	743	5.3	4.9-5.7	401	6.7	6.1-7.5	342	4.1	3.7-4.6
2014	3,193	23.1	22.2-23.9	1,767	28.5	27.2-29.9	1,426	18.6	17.7-19.7	810	5.7	5.3-6.1	450	7.5	6.8-8.2	360	4.3	3.8-4.8
2015	3,212	23.1	22.2-23.9	1,762	27.9	26.5-29.2	1,450	19.2	18.2-20.3	793	5.5	5.1-5.9	442	7.2	6.5-7.9	351	4.2	3.8-4.7
2016	3,253	23.1	22.3-24.0	1,718	26.9	25.6-28.2	1,535	20.2	19.1-21.2	809	5.5	5.1-5.9	438	7.0	6.3-7.7	371	4.4	4.0-4.9
2017	3,209	22.4	21.6-23.2	1,768	27.2	25.9-28.5	1,441	18.6	17.6-19.6	845	5.7	5.3-6.1	487	7.6	7.0-8.4	358	4.1	3.7-4.6
2018	3,137	21.8	21.0-22.6	1,664	25.2	24.0-26.5	1,473	19.1	18.1-20.1	728	4.8	4.5-5.2	418	6.5	5.9-7.2	310	3.7	3.2-4.1
2019	3,252	22.2	21.4-23.0	1,790	26.7	25.5-28.1	1,462	18.5	17.5-19.5	742	4.8	4.5-5.2	404	6.1	5.5-6.8	338	3.9	3.5-4.3
2020	2,949	20.0	19.3-20.8	1,613	23.9	22.7-25.2	1,336	16.9	16.0-17.9	705	4.6	4.2-4.9	372	5.6	5.0-6.2	333	3.8	3.4-4.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Myeloma

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	428	4.6	4.1-5.0	213	5.5	4.7-6.3	215	4.0	3.5-4.6	281	3.1	2.7-3.5	147	3.9	3.2-4.6	134	2.6	2.1-3.0
1977	402	4.3	3.9-4.7	205	5.3	4.5-6.1	197	3.6	3.1-4.1	268	2.8	2.5-3.2	135	3.4	2.8-4.1	133	2.4	2.0-2.9
1978	424	4.5	4.1-5.0	194	5.0	4.3-5.8	230	4.2	3.7-4.8	268	2.8	2.5-3.2	137	3.6	3.0-4.3	131	2.3	2.0-2.8
1979	454	4.8	4.3-5.2	203	5.2	4.5-6.0	251	4.5	3.9-5.1	275	2.9	2.6-3.3	140	3.7	3.1-4.5	135	2.4	2.0-2.9
1980	448	4.6	4.2-5.0	227	5.7	5.0-6.6	221	3.9	3.4-4.4	334	3.5	3.1-3.9	159	4.1	3.5-4.8	175	3.0	2.6-3.5
1981	495	5.1	4.6-5.5	228	5.8	5.0-6.6	267	4.7	4.1-5.3	316	3.2	2.9-3.6	147	3.8	3.2-4.5	169	2.9	2.5-3.4
1982	496	4.9	4.5-5.4	265	6.4	5.6-7.3	231	3.9	3.4-4.4	315	3.2	2.8-3.6	159	4.0	3.4-4.7	156	2.6	2.2-3.1
1983	464	4.6	4.2-5.1	254	6.5	5.7-7.4	210	3.6	3.1-4.1	324	3.3	2.9-3.7	168	4.4	3.7-5.1	156	2.6	2.2-3.1
1984	532	5.3	4.8-5.7	283	6.7	5.9-7.6	249	4.2	3.7-4.7	359	3.6	3.2-3.9	178	4.3	3.7-5.0	181	3.0	2.6-3.5
1985	525	5.2	4.7-5.6	266	6.4	5.6-7.2	259	4.3	3.8-4.9	374	3.7	3.3-4.1	192	4.7	4.0-5.4	182	2.9	2.5-3.4
1986	498	4.9	4.5-5.3	271	6.5	5.7-7.3	227	3.8	3.3-4.3	379	3.7	3.4-4.1	192	4.8	4.1-5.6	187	3.0	2.6-3.5
1987	550	5.4	4.9-5.8	271	6.5	5.7-7.3	279	4.6	4.1-5.2	364	3.5	3.1-3.9	183	4.5	3.8-5.2	181	3.0	2.5-3.4
1988	523	5.0	4.6-5.5	273	6.4	5.6-7.2	250	4.1	3.6-4.6	385	3.7	3.4-4.1	212	5.2	4.5-6.0	173	2.8	2.4-3.2
1989	578	5.5	5.1-6.0	267	6.3	5.6-7.2	311	5.1	4.5-5.7	351	3.3	3.0-3.7	176	4.2	3.6-4.9	175	2.7	2.3-3.2
1990	604	5.6	5.2-6.1	315	7.1	6.3-8.0	289	4.6	4.1-5.2	359	3.4	3.0-3.7	181	4.3	3.7-5.0	178	2.8	2.4-3.2
1991	574	5.3	4.9-5.7	294	6.8	6.0-7.6	280	4.4	3.9-4.9	381	3.5	3.2-3.9	186	4.3	3.7-4.9	195	3.0	2.6-3.5
1992	622	5.6	5.2-6.1	311	7.0	6.2-7.8	311	4.8	4.3-5.4	405	3.7	3.3-4.1	209	4.9	4.3-5.7	196	3.0	2.6-3.4
1993	630	5.7	5.2-6.1	318	6.9	6.1-7.7	312	4.9	4.3-5.4	429	3.9	3.5-4.3	206	4.6	4.0-5.3	223	3.3	2.9-3.8
1994	630	5.7	5.2-6.1	323	7.1	6.3-7.9	307	4.7	4.2-5.2	457	4.1	3.7-4.5	202	4.6	3.9-5.2	255	3.9	3.4-4.4
1995	650	5.8	5.4-6.3	345	7.6	6.8-8.5	305	4.7	4.2-5.2	437	3.9	3.5-4.3	227	5.0	4.4-5.7	210	3.1	2.7-3.6
1996	648	5.7	5.3-6.2	334	7.2	6.4-8.0	314	4.8	4.3-5.3	414	3.6	3.3-4.0	216	4.9	4.3-5.7	198	2.9	2.5-3.3
1997	674	5.9	5.5-6.4	328	6.8	6.1-7.6	346	5.3	4.7-5.9	423	3.7	3.3-4.0	214	4.7	4.1-5.4	209	3.1	2.7-3.6
1998	732	6.3	5.9-6.8	369	7.5	6.8-8.4	363	5.4	4.9-6.0	428	3.7	3.3-4.0	209	4.5	3.9-5.1	219	3.1	2.7-3.6
1999	760	6.5	6.0-7.0	401	8.2	7.4-9.1	359	5.3	4.8-5.9	446	3.8	3.4-4.2	232	4.8	4.2-5.5	214	3.0	2.6-3.5
2000	770	6.5	6.0-7.0	406	8.1	7.3-9.0	364	5.3	4.8-5.9	456	3.8	3.5-4.2	244	5.1	4.5-5.8	212	3.0	2.6-3.5
2001	738	6.2	5.7-6.6	419	8.3	7.5-9.2	319	4.6	4.1-5.1	404	3.4	3.0-3.7	210	4.3	3.7-4.9	194	2.7	2.3-3.1
2002	798	6.6	6.1-7.1	439	8.6	7.8-9.4	359	5.1	4.6-5.7	434	3.5	3.2-3.9	234	4.8	4.2-5.4	200	2.8	2.4-3.2
2003	764	6.2	5.8-6.7	420	8.0	7.3-8.9	344	4.9	4.4-5.5	412	3.3	3.0-3.7	233	4.6	4.0-5.2	179	2.5	2.1-2.9
2004	822	6.6	6.2-7.1	445	8.3	7.6-9.2	377	5.3	4.8-5.9	407	3.3	3.0-3.6	225	4.4	3.8-5.0	182	2.5	2.1-2.9
2005	809	6.4	6.0-6.9	429	7.9	7.2-8.7	380	5.3	4.8-5.9	418	3.3	3.0-3.6	215	4.1	3.5-4.7	203	2.8	2.4-3.2
2006	843	6.6	6.2-7.1	472	8.5	7.8-9.3	371	5.2	4.6-5.7	408	3.2	2.9-3.5	227	4.3	3.7-4.9	181	2.5	2.1-2.9
2007	894	7.0	6.5-7.4	478	8.6	7.8-9.4	416	5.8	5.2-6.4	433	3.3	3.0-3.7	223	4.1	3.5-4.6	210	2.8	2.4-3.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Myeloma

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	930	7.1	6.6-7.6	493	8.6	7.9-9.4	437	6.0	5.4-6.6	406	3.1	2.8-3.4	225	4.2	3.6-4.7	181	2.4	2.1-2.8
2009	984	7.4	7.0-7.9	526	9.1	8.3-9.9	458	6.1	5.6-6.7	424	3.1	2.8-3.5	226	4.0	3.5-4.5	198	2.5	2.2-2.9
2010	997	7.4	7.0-7.9	531	8.9	8.2-9.8	466	6.3	5.7-6.9	450	3.3	3.0-3.6	220	3.9	3.4-4.5	230	2.9	2.5-3.3
2011	1,018	7.5	7.1-8.0	557	9.3	8.5-10.1	461	6.1	5.5-6.7	415	3.0	2.7-3.3	213	3.6	3.1-4.2	202	2.5	2.2-2.9
2012	1,030	7.3	6.9-7.8	572	9.1	8.4-10.0	458	6.0	5.4-6.6	449	3.2	2.9-3.5	225	3.8	3.3-4.3	224	2.8	2.4-3.2
2013	1,128	8.1	7.6-8.6	644	10.3	9.5-11.1	484	6.3	5.8-6.9	398	2.8	2.5-3.1	208	3.5	3.0-4.0	190	2.4	2.0-2.7
2014	1,178	8.3	7.8-8.8	646	10.1	9.3-10.9	532	6.8	6.2-7.5	448	3.1	2.8-3.4	242	3.8	3.4-4.4	206	2.5	2.2-2.9
2015	1,115	7.8	7.3-8.3	621	9.7	8.9-10.5	494	6.3	5.7-6.9	451	3.1	2.8-3.4	255	4.1	3.6-4.6	196	2.3	2.0-2.7
2016	1,192	8.3	7.8-8.8	669	10.3	9.5-11.1	523	6.7	6.2-7.4	420	2.8	2.5-3.1	197	3.0	2.6-3.5	223	2.6	2.3-3.0
2017	1,137	7.7	7.2-8.2	655	9.8	9.0-10.6	482	6.0	5.4-6.5	435	2.9	2.6-3.2	229	3.5	3.1-4.0	206	2.4	2.1-2.7
2018	1,153	7.7	7.3-8.2	663	9.8	9.0-10.6	490	6.0	5.5-6.6	429	2.8	2.5-3.1	223	3.4	3.0-3.9	206	2.3	2.0-2.7
2019	1,200	7.9	7.5-8.4	665	9.6	8.8-10.4	535	6.5	6.0-7.2	403	2.6	2.3-2.8	241	3.6	3.1-4.1	162	1.8	1.5-2.1
2020	1,009	6.5	6.1-7.0	564	8.1	7.4-8.8	445	5.3	4.8-5.8	365	2.3	2.1-2.6	191	2.8	2.4-3.3	174	2.0	1.7-2.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York State Excl New York City
Leukemias**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,230	13.1	12.4-13.9	701	18.0	16.6-19.5	529	9.9	9.1-10.8	748	8.1	7.5-8.7	427	11.5	10.4-12.8	321	6.0	5.4-6.7
1977	1,164	12.4	11.7-13.2	650	16.5	15.1-17.9	514	9.6	8.8-10.5	807	8.7	8.1-9.3	436	11.2	10.1-12.3	371	7.0	6.3-7.8
1978	1,206	12.9	12.2-13.7	664	16.8	15.5-18.3	542	10.2	9.3-11.1	775	8.2	7.6-8.8	403	10.3	9.3-11.5	372	6.9	6.2-7.6
1979	1,317	13.6	12.9-14.4	746	18.2	16.8-19.6	571	10.4	9.5-11.3	806	8.5	7.9-9.1	437	11.1	10.1-12.3	369	6.8	6.1-7.5
1980	1,328	13.7	12.9-14.4	722	17.8	16.5-19.3	606	11.0	10.1-11.9	856	8.9	8.3-9.5	462	11.5	10.4-12.7	394	7.0	6.4-7.8
1981	1,229	12.6	11.9-13.4	693	17.0	15.7-18.4	536	9.6	8.8-10.5	829	8.7	8.1-9.3	459	11.8	10.7-13.0	370	6.7	6.0-7.4
1982	1,256	12.8	12.1-13.6	705	17.3	16.0-18.7	551	9.9	9.0-10.7	844	8.7	8.1-9.3	474	12.1	11.0-13.3	370	6.6	5.9-7.3
1983	1,249	12.6	11.9-13.3	710	16.9	15.6-18.3	539	9.5	8.7-10.3	833	8.5	7.9-9.1	429	10.9	9.9-12.1	404	7.0	6.3-7.7
1984	1,326	13.1	12.4-13.9	709	16.6	15.3-17.9	617	10.6	9.8-11.5	779	7.8	7.3-8.4	378	9.1	8.2-10.1	401	6.9	6.3-7.7
1985	1,406	13.8	13.1-14.6	799	18.7	17.3-20.1	607	10.5	9.6-11.4	831	8.2	7.7-8.8	488	11.9	10.8-13.1	343	5.8	5.2-6.5
1986	1,291	12.7	12.0-13.5	724	17.1	15.8-18.5	567	9.7	8.9-10.5	831	8.3	7.7-8.8	476	11.4	10.4-12.5	355	6.0	5.4-6.6
1987	1,286	12.4	11.7-13.1	703	16.1	14.8-17.4	583	9.9	9.1-10.8	823	8.1	7.5-8.7	451	10.8	9.8-11.9	372	6.3	5.6-6.9
1988	1,303	12.5	11.8-13.2	726	16.7	15.5-18.0	577	9.8	9.0-10.6	849	8.2	7.6-8.8	484	11.7	10.6-12.8	365	6.0	5.4-6.7
1989	1,282	12.2	11.5-12.9	715	16.4	15.2-17.7	567	9.3	8.6-10.1	831	7.9	7.4-8.5	456	10.7	9.7-11.8	375	6.1	5.5-6.8
1990	1,454	13.7	13.0-14.5	793	17.6	16.3-18.9	661	10.9	10.1-11.8	825	7.8	7.3-8.4	421	9.8	8.8-10.8	404	6.5	5.9-7.2
1991	1,401	13.1	12.4-13.8	761	17.0	15.7-18.3	640	10.5	9.7-11.4	835	7.8	7.3-8.4	425	9.8	8.9-10.9	410	6.5	5.9-7.2
1992	1,430	13.2	12.6-13.9	812	17.9	16.7-19.3	618	10.0	9.2-10.8	875	8.1	7.5-8.6	485	11.1	10.1-12.2	390	6.1	5.5-6.8
1993	1,440	13.2	12.5-13.9	805	17.3	16.1-18.6	635	10.3	9.5-11.1	784	7.2	6.7-7.7	426	9.8	8.8-10.8	358	5.6	5.0-6.2
1994	1,548	14.0	13.3-14.7	862	18.4	17.2-19.8	686	10.8	10.0-11.7	889	8.0	7.5-8.6	504	11.2	10.2-12.3	385	5.9	5.3-6.5
1995	1,529	13.6	13.0-14.4	856	18.1	16.9-19.4	673	10.6	9.8-11.5	883	7.9	7.4-8.4	500	11.3	10.3-12.3	383	5.9	5.3-6.5
1996	1,598	14.2	13.5-14.9	927	19.1	17.9-20.4	671	10.5	9.7-11.3	944	8.4	7.8-8.9	521	11.2	10.3-12.3	423	6.5	5.9-7.1
1997	1,594	14.1	13.4-14.8	867	17.9	16.7-19.1	727	11.2	10.4-12.0	904	7.9	7.4-8.4	497	10.6	9.7-11.6	407	6.0	5.4-6.6
1998	1,614	14.1	13.4-14.8	927	18.7	17.5-20.0	687	10.6	9.8-11.4	884	7.7	7.2-8.2	482	10.1	9.2-11.0	402	5.9	5.4-6.6
1999	1,746	15.1	14.4-15.8	970	19.7	18.5-21.0	776	11.8	11.0-12.7	938	8.0	7.5-8.5	505	10.5	9.6-11.5	433	6.3	5.7-6.9
2000	1,860	15.9	15.2-16.6	1,055	21.2	19.9-22.5	805	12.0	11.2-12.8	938	7.9	7.4-8.5	530	10.9	10.0-11.9	408	5.9	5.3-6.5
2001	1,840	15.6	14.9-16.3	1,025	20.2	19.0-21.5	815	12.2	11.4-13.1	922	7.7	7.2-8.2	499	10.3	9.4-11.2	423	6.1	5.5-6.7
2002	1,758	14.8	14.1-15.5	1,007	19.5	18.3-20.8	751	11.2	10.4-12.0	902	7.4	7.0-8.0	499	10.1	9.2-11.0	403	5.6	5.1-6.2
2003	1,900	15.8	15.1-16.5	1,076	20.7	19.5-22.0	824	12.2	11.4-13.1	893	7.3	6.8-7.8	509	10.2	9.3-11.1	384	5.3	4.8-5.9
2004	1,983	16.3	15.6-17.0	1,116	21.1	19.8-22.3	867	12.7	11.9-13.6	969	7.8	7.3-8.3	543	10.6	9.7-11.6	426	5.9	5.4-6.5
2005	1,988	16.2	15.5-17.0	1,092	20.3	19.1-21.6	896	13.0	12.1-13.9	930	7.4	6.9-7.9	534	10.4	9.5-11.3	396	5.4	4.8-5.9
2006	2,001	16.2	15.5-17.0	1,159	21.5	20.3-22.8	842	12.2	11.3-13.0	935	7.5	7.0-8.0	519	9.9	9.1-10.8	416	5.6	5.1-6.2
2007	2,058	16.4	15.7-17.2	1,194	21.6	20.4-22.8	864	12.6	11.8-13.5	908	7.1	6.7-7.6	519	9.7	8.9-10.6	389	5.3	4.8-5.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York State Excl New York City

Leukemias

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	2,017	16.0	15.3-16.7	1,158	20.5	19.4-21.8	859	12.5	11.6-13.4	925	7.1	6.6-7.6	538	9.8	9.0-10.7	387	5.1	4.6-5.7
2009	2,137	16.7	16.0-17.4	1,260	22.1	20.9-23.4	877	12.4	11.6-13.3	953	7.4	6.9-7.9	576	10.5	9.7-11.4	377	5.0	4.5-5.6
2010	2,271	17.5	16.8-18.3	1,325	23.0	21.7-24.3	946	13.2	12.3-14.1	879	6.6	6.2-7.1	504	9.1	8.3-9.9	375	4.9	4.4-5.4
2011	2,423	18.5	17.7-19.3	1,406	24.1	22.9-25.4	1,017	14.0	13.1-14.9	948	7.0	6.6-7.5	542	9.4	8.6-10.3	406	5.3	4.8-5.9
2012	2,357	17.7	17.0-18.5	1,364	22.9	21.7-24.2	993	13.8	12.9-14.7	954	7.0	6.5-7.5	564	9.8	9.0-10.7	390	4.9	4.4-5.4
2013	2,541	19.0	18.3-19.8	1,473	24.5	23.2-25.8	1,068	14.7	13.8-15.7	928	6.7	6.3-7.2	535	9.2	8.4-10.0	393	4.9	4.4-5.5
2014	2,533	18.5	17.8-19.3	1,539	25.0	23.7-26.3	994	13.3	12.5-14.2	925	6.5	6.1-6.9	519	8.7	7.9-9.5	406	5.0	4.5-5.5
2015	2,488	18.1	17.3-18.8	1,454	23.4	22.2-24.7	1,034	13.8	12.9-14.7	920	6.4	6.0-6.8	534	8.8	8.0-9.5	386	4.6	4.2-5.2
2016	2,598	18.4	17.7-19.1	1,550	24.3	23.0-25.6	1,048	13.5	12.7-14.4	902	6.3	5.9-6.7	520	8.5	7.8-9.3	382	4.6	4.2-5.2
2017	2,453	17.4	16.7-18.1	1,457	22.7	21.5-23.9	996	13.2	12.3-14.1	877	5.9	5.5-6.3	498	8.0	7.3-8.7	379	4.4	4.0-4.9
2018	2,359	16.6	15.9-17.3	1,396	21.6	20.4-22.8	963	12.6	11.8-13.5	912	6.1	5.7-6.6	532	8.5	7.8-9.2	380	4.5	4.0-5.0
2019	2,405	16.5	15.8-17.2	1,418	21.3	20.2-22.5	987	12.5	11.7-13.4	879	5.8	5.4-6.2	516	7.9	7.2-8.6	363	4.2	3.8-4.7
2020	2,186	15.2	14.6-15.9	1,311	19.8	18.8-21.0	875	11.3	10.5-12.2	851	5.6	5.2-6.0	504	7.7	7.0-8.4	347	4.0	3.5-4.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
All Invasive Malignant Tumors**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	4,319	355.0	344.2-366.0	2,060	424.8	405.9-444.3	2,259	316.9	303.7-330.6	3,030	250.4	241.3-259.7	1,548	325.6	308.9-342.9	1,482	204.9	194.4-215.8
1977	4,696	395.8	384.3-407.5	2,260	481.7	461.3-502.6	2,436	347.7	333.7-362.2	2,672	225.6	216.9-234.5	1,362	291.3	275.5-307.7	1,310	185.1	175.0-195.6
1978	4,686	406.2	394.4-418.2	2,251	494.8	474.0-516.3	2,435	356.1	341.8-370.9	2,506	216.1	207.6-224.9	1,238	274.8	259.2-291.0	1,268	181.9	171.9-192.4
1979	4,545	399.9	388.2-411.9	2,219	501.6	480.5-523.5	2,326	343.7	329.6-358.3	2,529	223.4	214.6-232.4	1,281	293.4	277.2-310.4	1,248	181.7	171.6-192.3
1980	4,601	410.9	398.9-423.1	2,207	505.0	483.6-527.0	2,394	360.0	345.4-375.1	2,563	228.8	219.9-237.9	1,307	305.2	288.5-322.6	1,256	185.0	174.7-195.8
1981	4,245	382.3	370.7-394.2	2,027	470.5	449.7-491.9	2,218	332.4	318.4-346.8	2,468	222.5	213.7-231.6	1,244	293.3	276.9-310.4	1,224	181.3	171.1-192.0
1982	4,538	411.8	399.7-424.1	2,127	498.5	477.0-520.6	2,411	364.3	349.6-379.5	2,340	211.9	203.3-220.8	1,158	272.3	256.5-288.8	1,182	176.8	166.7-187.4
1983	4,436	402.2	390.3-414.4	2,016	472.9	452.0-494.4	2,420	364.8	350.1-380.0	2,341	211.7	203.1-220.5	1,168	279.2	263.1-296.0	1,173	171.5	161.7-181.8
1984	4,386	399.8	387.9-412.0	2,055	487.2	465.9-509.1	2,331	353.7	339.2-368.7	2,351	213.8	205.2-222.8	1,189	286.7	270.3-303.8	1,162	172.8	162.9-183.3
1985	4,475	409.1	397.1-421.5	2,168	511.8	490.1-534.3	2,307	350.6	336.2-365.5	2,300	209.4	200.9-218.3	1,160	276.0	260.1-292.7	1,140	171.2	161.2-181.6
1986	4,485	409.3	397.3-421.6	2,141	506.4	484.7-528.8	2,344	357.2	342.6-372.3	2,310	210.2	201.6-219.0	1,118	269.4	253.5-285.9	1,192	176.8	166.8-187.4
1987	4,406	400.1	388.3-412.3	2,071	492.0	470.6-514.0	2,335	350.5	336.1-365.2	2,291	207.8	199.3-216.6	1,135	272.4	256.5-289.0	1,156	170.4	160.6-180.7
1988	4,425	403.9	392.0-416.1	2,081	499.5	477.8-521.8	2,344	353.3	338.9-368.1	2,375	218.0	209.2-227.0	1,212	301.0	284.0-318.7	1,163	171.7	161.9-182.0
1989	4,227	389.9	378.1-401.9	1,966	473.6	452.5-495.4	2,261	346.2	331.9-361.0	2,271	209.4	200.8-218.3	1,120	279.4	263.0-296.5	1,151	171.5	161.6-181.8
1990	4,551	425.2	412.8-437.8	2,119	516.0	493.9-538.9	2,432	376.7	361.7-392.2	2,273	214.1	205.3-223.1	1,098	280.1	263.6-297.4	1,175	178.1	168.0-188.8
1991	4,957	466.9	453.9-480.2	2,310	568.5	545.1-592.6	2,647	415.0	399.2-431.3	2,272	215.2	206.3-224.2	1,114	283.5	266.8-300.9	1,158	177.7	167.5-188.3
1992	4,757	449.1	436.3-462.1	2,285	560.3	537.1-584.2	2,472	388.2	372.9-404.0	2,292	218.3	209.4-227.4	1,142	293.9	276.9-311.7	1,150	177.4	167.2-188.1
1993	4,664	438.8	426.2-451.6	2,285	555.8	532.8-579.5	2,379	373.2	358.2-388.7	2,327	221.8	212.9-231.0	1,125	288.3	271.4-305.9	1,202	186.2	175.7-197.1
1994	5,008	473.5	460.4-486.9	2,389	581.0	557.5-605.2	2,619	410.9	395.2-427.1	2,375	225.9	216.9-235.2	1,147	293.0	276.0-310.7	1,228	188.7	178.2-199.6
1995	4,611	434.7	422.1-447.4	2,194	531.2	508.7-554.3	2,417	378.7	363.6-394.2	2,287	218.0	209.2-227.2	1,097	279.8	263.3-297.1	1,190	183.4	173.0-194.2
1996	4,776	447.6	434.9-460.5	2,291	550.7	527.9-574.1	2,485	388.6	373.4-404.3	2,153	205.5	196.9-214.4	1,004	255.8	239.9-272.3	1,149	177.6	167.4-188.2
1997	4,839	453.2	440.4-466.2	2,275	546.8	524.1-570.1	2,564	397.6	382.3-413.4	2,077	197.0	188.6-205.7	953	240.2	224.9-256.1	1,124	173.0	163.0-183.5
1998	4,962	460.1	447.4-473.2	2,360	558.0	535.3-581.4	2,602	402.2	386.8-418.0	2,075	195.4	187.1-204.0	974	241.3	226.1-257.1	1,101	168.2	158.3-178.5
1999	4,919	448.4	435.9-461.2	2,376	556.1	533.5-579.3	2,543	386.4	371.4-401.8	2,122	195.7	187.4-204.2	1,000	245.4	230.1-261.3	1,122	169.4	159.6-179.7
2000	5,384	482.4	469.6-495.6	2,695	621.1	597.5-645.4	2,689	401.0	385.9-416.5	2,025	184.0	176.1-192.2	948	226.4	211.9-241.5	1,077	159.1	149.7-168.9
2001	5,247	464.2	451.7-477.0	2,647	593.2	570.4-616.7	2,600	384.7	370.0-399.8	2,116	189.9	181.9-198.2	1,023	241.7	226.8-257.3	1,093	159.8	150.4-169.6
2002	5,295	460.7	448.3-473.3	2,590	569.3	547.0-592.1	2,705	395.3	380.5-410.6	1,970	174.6	167.0-182.5	944	219.7	205.6-234.4	1,026	148.1	139.2-157.5
2003	5,491	472.1	459.7-484.9	2,666	583.3	560.8-606.3	2,825	406.6	391.7-422.0	2,156	189.3	181.3-197.5	1,010	234.1	219.5-249.3	1,146	163.9	154.5-173.7
2004	5,363	457.2	444.9-469.6	2,631	564.5	542.6-587.0	2,732	391.7	377.1-406.7	2,021	176.2	168.6-184.1	956	218.4	204.4-233.0	1,065	151.8	142.8-161.3
2005	5,405	457.2	445.0-469.6	2,717	576.6	554.6-599.3	2,688	384.0	369.6-398.9	2,015	173.7	166.1-181.5	967	218.4	204.5-233.0	1,048	147.7	138.8-156.9
2006	5,605	468.6	456.3-481.1	2,840	596.4	574.0-619.4	2,765	391.0	376.5-406.0	1,953	168.1	160.7-175.8	925	208.4	194.9-222.7	1,028	145.0	136.2-154.2
2007	5,751	473.5	461.2-486.0	2,935	598.2	576.1-620.9	2,816	394.1	379.5-409.0	2,024	171.4	163.9-179.0	991	218.7	205.0-233.2	1,033	143.3	134.7-152.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
All Invasive Malignant Tumors**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	5,596	454.1	442.1-466.3	2,766	552.1	531.1-573.6	2,830	392.0	377.6-406.9	2,011	167.1	159.8-174.7	952	208.0	194.7-222.1	1,059	144.7	136.0-153.7
2009	5,901	469.6	457.5-481.8	2,890	566.3	545.2-587.9	3,011	409.0	394.4-424.0	2,060	168.8	161.5-176.3	1,022	217.2	203.7-231.3	1,038	140.6	132.1-149.5
2010	5,781	451.6	439.9-463.6	2,934	560.7	540.0-582.1	2,847	382.7	368.6-397.1	1,973	158.3	151.3-165.5	954	200.4	187.5-213.9	1,019	135.1	126.8-143.8
2011	6,109	465.7	453.9-477.6	3,017	561.2	540.7-582.2	3,092	405.0	390.7-419.7	2,018	159.2	152.2-166.4	1,026	208.8	195.8-222.3	992	129.8	121.7-138.2
2012	6,075	454.8	443.3-466.5	2,999	544.3	524.4-564.7	3,076	399.6	385.5-414.1	2,089	160.0	153.1-167.1	1,018	198.9	186.5-211.9	1,071	136.1	128.0-144.6
2013	6,060	443.1	431.9-454.6	2,961	519.7	500.6-539.3	3,099	394.4	380.4-408.7	1,960	146.5	140.0-153.2	966	185.0	173.2-197.4	994	123.4	115.8-131.4
2014	6,248	445.4	434.3-456.7	2,998	512.2	493.5-531.4	3,250	404.9	390.9-419.2	2,118	154.4	147.8-161.2	1,016	188.7	177.0-201.0	1,102	134.0	126.1-142.3
2015	6,310	442.9	431.9-454.1	3,033	506.9	488.5-525.8	3,277	403.5	389.6-417.7	2,019	144.4	138.1-151.0	973	177.3	166.0-189.1	1,046	125.1	117.5-133.0
2016	6,160	422.8	412.2-433.7	3,023	492.1	474.2-510.4	3,137	379.3	365.9-393.0	2,166	151.5	145.1-158.1	1,063	187.1	175.7-199.0	1,103	129.8	122.2-137.8
2017	6,409	432.5	421.8-443.4	3,125	495.8	478.1-514.0	3,284	391.8	378.3-405.7	2,011	137.1	131.1-143.3	985	168.1	157.5-179.2	1,026	118.2	110.9-125.8
2018	6,471	430.7	420.1-441.5	3,102	483.9	466.6-501.7	3,369	396.3	382.8-410.2	2,010	134.7	128.8-140.8	1,019	169.7	159.2-180.7	991	111.7	104.8-119.0
2019	6,539	433.2	422.6-444.0	3,235	499.9	482.4-517.9	3,304	390.7	377.2-404.6	1,997	131.6	125.8-137.6	971	158.2	148.2-168.7	1,026	114.2	107.2-121.6
2020	5,443	356.6	347.0-366.4	2,707	415.6	399.7-431.9	2,736	319.4	307.3-331.9	1,863	121.0	115.5-126.7	913	146.9	137.3-156.9	950	105.1	98.4-112.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Oral cavity and pharynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	135	11.4	9.5-13.5	92	18.7	14.9-23.0	43	6.6	4.7-8.9	101	8.3	6.7-10.1	73	14.5	11.3-18.4	28	3.8	2.5-5.6
1977	146	12.2	10.2-14.3	95	19.9	16.0-24.4	51	6.9	5.1-9.1	74	6.6	5.1-8.3	52	11.3	8.3-14.8	22	3.2	2.0-4.9
1978	147	13.1	11.0-15.4	102	22.1	18.0-27.0	45	6.6	4.8-8.9	57	5.2	3.9-6.7	38	8.4	5.9-11.5	19	3.1	1.8-4.8
1979	143	12.6	10.6-14.9	96	20.8	16.8-25.4	47	6.9	5.0-9.2	64	5.8	4.5-7.5	42	9.3	6.7-12.7	22	3.4	2.1-5.2
1980	137	12.6	10.5-14.9	101	22.8	18.5-27.8	36	5.7	4.0-8.0	75	6.8	5.3-8.6	57	13.2	9.9-17.2	18	2.4	1.4-3.9
1981	149	13.6	11.5-16.0	111	24.7	20.2-29.9	38	5.8	4.1-8.1	72	6.6	5.1-8.3	54	12.3	9.2-16.1	18	2.9	1.7-4.6
1982	159	14.8	12.5-17.3	105	23.8	19.4-28.9	54	8.4	6.3-11.0	59	5.3	4.0-6.8	43	9.6	6.9-13.0	16	2.4	1.3-3.9
1983	128	11.9	9.9-14.2	82	19.0	15.0-23.7	46	6.9	5.1-9.3	65	5.7	4.4-7.3	49	10.7	7.9-14.2	16	2.3	1.3-3.8
1984	142	13.9	11.7-16.4	96	23.3	18.8-28.5	46	7.4	5.4-10.0	45	4.4	3.2-5.9	34	8.1	5.6-11.3	11	2.0	1.0-3.6
1985	161	15.0	12.8-17.6	111	26.0	21.3-31.3	50	7.7	5.7-10.2	48	4.4	3.3-5.9	41	9.7	6.9-13.2	7	0.9	0.4-2.0
1986	153	14.7	12.5-17.3	100	24.4	19.8-29.7	53	8.6	6.4-11.3	57	5.2	3.9-6.7	41	9.5	6.8-12.9	16	2.6	1.4-4.2
1987	140	13.1	11.0-15.5	93	21.3	17.1-26.1	47	7.4	5.4-9.9	48	4.5	3.3-5.9	36	8.4	5.8-11.6	12	1.8	0.9-3.1
1988	139	13.0	10.9-15.3	99	22.7	18.4-27.7	40	6.1	4.3-8.3	57	5.4	4.1-7.0	43	10.1	7.3-13.6	14	2.2	1.2-3.7
1989	128	12.2	10.1-14.5	75	18.0	14.1-22.6	53	8.0	6.0-10.5	54	5.0	3.7-6.5	36	8.5	5.9-11.8	18	2.6	1.5-4.1
1990	135	13.0	10.9-15.4	88	21.1	16.9-26.1	47	7.5	5.5-9.9	44	4.2	3.0-5.6	25	6.5	4.2-9.5	19	2.8	1.7-4.4
1991	159	15.6	13.3-18.3	107	25.9	21.2-31.3	52	8.5	6.4-11.2	54	5.2	3.9-6.8	41	10.1	7.2-13.7	13	2.0	1.0-3.4
1992	159	15.5	13.2-18.1	100	23.8	19.3-29.0	59	9.5	7.2-12.3	35	3.3	2.3-4.6	22	5.3	3.3-8.0	13	2.0	1.1-3.5
1993	166	16.0	13.7-18.7	109	26.1	21.4-31.5	57	9.2	6.9-11.9	44	4.3	3.1-5.7	36	8.5	5.9-11.7	8	1.3	0.6-2.6
1994	151	14.6	12.3-17.1	102	24.0	19.5-29.2	49	7.8	5.7-10.3	46	4.5	3.3-6.0	31	7.5	5.1-10.6	15	2.4	1.3-3.9
1995	92	8.8	7.1-10.8	65	15.4	11.9-19.7	27	4.4	2.9-6.4	45	4.4	3.2-5.8	34	8.0	5.5-11.2	11	1.8	0.9-3.1
1996	122	11.4	9.5-13.7	81	18.2	14.4-22.6	41	6.6	4.7-8.9	40	3.8	2.7-5.2	25	5.7	3.6-8.4	15	2.5	1.4-4.1
1997	124	11.4	9.5-13.6	79	17.8	14.0-22.2	45	7.0	5.1-9.4	45	4.2	3.1-5.6	31	7.1	4.8-10.1	14	2.1	1.1-3.5
1998	113	10.4	8.5-12.5	70	15.6	12.1-19.8	43	6.6	4.8-8.9	37	3.5	2.5-4.8	26	6.3	4.1-9.2	11	1.7	0.8-3.0
1999	128	11.6	9.7-13.8	94	20.5	16.5-25.1	34	5.2	3.6-7.3	39	3.6	2.6-4.9	27	6.7	4.4-9.8	12	1.9	1.0-3.2
2000	104	9.3	7.6-11.3	69	15.4	11.9-19.5	35	5.2	3.6-7.3	24	2.1	1.3-3.1	18	4.0	2.3-6.3	6	0.9	0.3-2.0
2001	126	10.9	9.1-13.0	97	20.0	16.2-24.4	29	4.1	2.8-5.9	36	3.1	2.2-4.3	32	6.7	4.6-9.5	4	0.5	0.1-1.4
2002	135	11.4	9.6-13.5	88	17.7	14.1-21.8	47	6.9	5.1-9.2	31	2.7	1.8-3.8	17	3.7	2.1-5.9	14	2.0	1.1-3.3
2003	111	9.4	7.7-11.3	76	15.7	12.3-19.7	35	5.2	3.6-7.2	36	3.1	2.2-4.3	25	5.5	3.5-8.1	11	1.6	0.8-2.8
2004	130	10.8	9.1-12.9	88	17.9	14.3-22.1	42	5.9	4.2-7.9	45	3.7	2.7-5.0	37	7.3	5.1-10.1	8	1.1	0.5-2.2
2005	143	11.7	9.8-13.8	102	19.8	16.1-24.1	41	5.9	4.2-8.0	37	3.1	2.1-4.2	26	5.3	3.4-7.8	11	1.5	0.7-2.6
2006	128	10.4	8.6-12.4	76	15.5	12.1-19.4	52	7.2	5.3-9.4	25	2.0	1.3-3.0	22	4.4	2.7-6.7	3	0.5	0.1-1.3
2007	152	12.5	10.6-14.7	106	21.3	17.3-25.9	46	6.5	4.8-8.7	37	3.2	2.2-4.4	28	6.3	4.1-9.1	9	1.3	0.6-2.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Oral cavity and pharynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	125	9.8	8.1-11.7	80	14.7	11.6-18.4	45	6.1	4.4-8.2	30	2.4	1.6-3.5	24	4.9	3.1-7.4	6	0.9	0.3-1.9
2009	154	12.3	10.4-14.4	107	20.5	16.7-24.9	47	6.4	4.7-8.5	38	3.1	2.2-4.2	29	5.6	3.7-8.1	9	1.2	0.6-2.4
2010	110	8.4	6.9-10.1	79	14.5	11.4-18.2	31	3.8	2.6-5.5	27	2.1	1.4-3.1	22	4.4	2.7-6.6	5	0.7	0.2-1.6
2011	134	10.1	8.4-12.0	88	15.6	12.4-19.3	46	6.1	4.4-8.1	38	2.8	1.9-3.8	30	5.0	3.4-7.3	8	1.0	0.4-1.9
2012	150	11.1	9.3-13.0	94	16.2	13.0-19.9	56	7.2	5.4-9.4	27	2.1	1.3-3.0	13	2.3	1.2-4.0	14	1.8	1.0-3.1
2013	131	9.4	7.9-11.2	99	17.0	13.7-20.7	32	4.0	2.7-5.6	30	2.2	1.5-3.2	25	4.5	2.9-6.7	5	0.6	0.2-1.5
2014	144	9.8	8.3-11.6	102	16.4	13.3-20.0	42	5.1	3.7-6.9	32	2.3	1.6-3.3	24	4.3	2.7-6.4	8	1.1	0.5-2.1
2015	148	10.2	8.6-12.1	93	15.0	12.0-18.4	55	6.7	5.0-8.7	41	2.8	2.0-3.8	26	4.4	2.8-6.4	15	1.6	0.9-2.7
2016	138	9.5	7.9-11.2	102	16.3	13.2-19.9	36	4.4	3.0-6.1	43	2.9	2.1-3.9	34	5.6	3.8-7.9	9	1.0	0.5-2.0
2017	170	11.4	9.7-13.2	112	17.5	14.3-21.2	58	6.8	5.2-8.9	39	2.7	1.9-3.7	25	4.1	2.6-6.1	14	1.8	0.9-3.0
2018	132	8.9	7.4-10.5	80	12.9	10.2-16.1	52	5.9	4.4-7.8	32	2.2	1.5-3.1	22	3.6	2.2-5.5	10	1.2	0.6-2.2
2019	160	10.4	8.8-12.2	114	16.9	13.9-20.4	46	5.4	3.9-7.2	41	2.8	2.0-3.8	28	4.5	3.0-6.6	13	1.5	0.8-2.6
2020	152	9.7	8.2-11.4	98	14.8	11.9-18.1	54	6.0	4.5-7.8	43	2.7	1.9-3.6	32	4.9	3.3-6.9	11	1.1	0.6-2.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Esophagus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	65	5.2	3.9-6.6	47	8.9	6.5-11.9	18	2.5	1.5-4.0	89	7.2	5.8-9.0	59	11.9	9.0-15.5	30	4.1	2.8-5.9
1977	72	5.8	4.5-7.3	46	8.8	6.4-11.8	26	3.8	2.4-5.5	78	6.3	5.0-7.9	46	8.9	6.5-12.0	32	4.5	3.0-6.3
1978	81	7.3	5.7-9.1	51	11.1	8.2-14.6	30	4.5	3.0-6.4	57	4.9	3.7-6.4	33	7.1	4.8-10.0	24	3.5	2.2-5.3
1979	74	6.6	5.1-8.3	48	10.8	7.9-14.4	26	3.9	2.5-5.7	71	6.4	5.0-8.1	41	8.8	6.3-11.9	30	4.8	3.2-6.8
1980	85	7.5	5.9-9.2	52	11.1	8.3-14.6	33	5.0	3.4-7.0	56	4.9	3.7-6.4	40	8.9	6.3-12.2	16	2.4	1.3-3.9
1981	79	7.2	5.7-9.0	53	12.2	9.1-16.0	26	3.8	2.5-5.6	77	7.0	5.5-8.8	50	11.6	8.6-15.3	27	3.9	2.6-5.8
1982	101	9.6	7.8-11.7	63	15.6	12.0-20.0	38	5.5	3.8-7.5	73	6.8	5.3-8.6	51	12.3	9.1-16.1	22	3.2	2.0-4.9
1983	83	7.4	5.9-9.3	60	13.4	10.2-17.3	23	3.4	2.1-5.2	73	6.9	5.4-8.7	45	10.7	7.8-14.4	28	4.4	2.9-6.4
1984	96	8.8	7.1-10.8	66	15.5	12.0-19.8	30	4.6	3.1-6.6	74	6.7	5.2-8.4	57	12.9	9.7-16.7	17	2.5	1.5-4.1
1985	79	7.3	5.8-9.1	44	10.4	7.5-14.0	35	5.1	3.6-7.2	72	6.5	5.0-8.2	50	11.5	8.5-15.2	22	3.0	1.9-4.6
1986	79	7.1	5.6-8.9	50	11.4	8.4-15.1	29	4.2	2.8-6.1	58	5.5	4.1-7.1	39	9.4	6.6-12.8	19	3.2	1.9-5.0
1987	73	6.9	5.4-8.7	51	12.1	8.9-15.9	22	3.5	2.1-5.3	59	5.4	4.1-7.0	36	8.3	5.8-11.5	23	3.4	2.2-5.2
1988	81	7.4	5.9-9.2	58	13.6	10.3-17.6	23	3.4	2.1-5.1	62	5.8	4.4-7.4	49	11.8	8.7-15.6	13	1.8	1.0-3.1
1989	71	6.6	5.1-8.3	44	10.6	7.6-14.2	27	3.9	2.6-5.7	63	5.8	4.5-7.4	46	10.8	7.9-14.4	17	2.4	1.4-3.9
1990	89	8.4	6.7-10.3	57	14.1	10.6-18.3	32	4.9	3.4-7.0	72	6.7	5.2-8.4	46	11.0	8.1-14.7	26	3.8	2.5-5.6
1991	75	7.3	5.7-9.1	45	11.1	8.1-14.8	30	4.7	3.1-6.7	65	6.2	4.8-8.0	44	11.2	8.1-15.0	21	3.2	2.0-4.9
1992	88	8.5	6.8-10.4	60	14.5	11.0-18.7	28	4.3	2.9-6.2	75	7.3	5.7-9.1	50	12.2	9.0-16.1	25	4.0	2.6-6.0
1993	79	7.5	6.0-9.4	55	13.4	10.1-17.5	24	3.7	2.4-5.6	74	7.1	5.6-9.0	51	12.7	9.4-16.6	23	3.8	2.4-5.7
1994	80	7.9	6.2-9.8	55	13.6	10.2-17.7	25	4.1	2.6-6.0	64	6.3	4.8-8.0	49	12.2	9.0-16.2	15	2.4	1.3-3.9
1995	71	6.7	5.2-8.5	44	10.4	7.5-14.0	27	4.0	2.6-5.9	55	5.3	4.0-6.8	38	9.2	6.5-12.6	17	2.5	1.5-4.1
1996	75	7.1	5.6-8.9	42	9.9	7.1-13.4	33	5.1	3.5-7.1	68	6.4	5.0-8.1	42	9.8	7.0-13.2	26	3.7	2.4-5.5
1997	83	8.0	6.3-9.9	51	12.1	8.9-15.8	32	5.0	3.4-7.1	57	5.5	4.1-7.1	38	9.4	6.6-12.8	19	3.0	1.8-4.6
1998	79	7.5	5.9-9.3	55	13.2	9.9-17.2	24	3.8	2.4-5.6	61	5.8	4.4-7.4	40	9.5	6.8-12.9	21	3.3	2.0-5.1
1999	67	6.2	4.8-7.8	50	11.7	8.6-15.4	17	2.6	1.5-4.2	60	5.5	4.2-7.1	45	10.8	7.8-14.4	15	2.4	1.3-3.9
2000	73	6.7	5.3-8.5	44	10.7	7.7-14.3	29	4.4	2.9-6.3	56	5.1	3.9-6.6	36	8.5	5.9-11.8	20	2.9	1.8-4.5
2001	59	5.3	4.0-6.9	37	8.3	5.8-11.5	22	3.2	2.0-4.8	56	5.0	3.8-6.5	38	8.7	6.1-11.9	18	2.7	1.6-4.2
2002	64	5.6	4.3-7.1	44	9.4	6.8-12.7	20	2.9	1.8-4.5	43	3.8	2.8-5.1	25	5.8	3.7-8.6	18	2.5	1.5-4.0
2003	61	5.3	4.0-6.8	45	10.0	7.3-13.5	16	2.2	1.3-3.6	47	4.1	3.0-5.4	27	5.7	3.7-8.3	20	2.8	1.7-4.4
2004	64	5.5	4.3-7.1	44	9.5	6.8-12.8	20	2.9	1.7-4.4	45	3.9	2.8-5.2	31	6.9	4.6-9.8	14	2.0	1.1-3.4
2005	50	4.3	3.2-5.6	40	8.8	6.2-12.0	10	1.5	0.7-2.8	56	4.9	3.7-6.4	40	9.1	6.5-12.4	16	2.2	1.3-3.6
2006	57	4.9	3.7-6.3	43	9.0	6.4-12.1	14	2.0	1.1-3.4	35	3.0	2.1-4.1	23	4.6	2.9-6.9	12	1.7	0.9-3.0
2007	59	4.9	3.7-6.3	42	8.6	6.1-11.7	17	2.3	1.3-3.7	47	4.0	2.9-5.3	37	8.3	5.8-11.4	10	1.3	0.6-2.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Esophagus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	68	5.5	4.3-7.0	46	9.1	6.6-12.2	22	2.9	1.8-4.4	53	4.2	3.1-5.5	36	6.6	4.6-9.2	17	2.4	1.4-3.9
2009	62	5.1	3.9-6.6	46	9.4	6.8-12.6	16	2.2	1.2-3.6	52	4.4	3.3-5.7	37	8.0	5.5-11.0	15	2.0	1.1-3.3
2010	48	3.7	2.7-4.9	31	6.0	4.0-8.5	17	2.2	1.3-3.5	36	3.0	2.1-4.1	25	5.4	3.4-7.9	11	1.5	0.7-2.7
2011	63	4.8	3.7-6.1	43	8.2	5.8-11.1	20	2.6	1.6-4.0	45	3.4	2.5-4.6	36	7.0	4.9-9.8	9	1.1	0.5-2.2
2012	43	3.2	2.3-4.3	29	5.4	3.5-7.7	14	1.7	0.9-2.9	41	3.0	2.2-4.1	27	5.0	3.2-7.3	14	1.6	0.9-2.8
2013	48	3.6	2.6-4.7	35	6.5	4.5-9.1	13	1.6	0.9-2.8	37	2.8	1.9-3.8	28	5.1	3.4-7.5	9	1.1	0.5-2.1
2014	64	4.6	3.5-5.9	46	8.1	5.9-10.9	18	2.2	1.3-3.5	56	4.0	3.0-5.2	44	7.9	5.7-10.6	12	1.6	0.8-2.7
2015	49	3.4	2.5-4.5	31	5.3	3.5-7.5	18	2.2	1.3-3.4	49	3.5	2.6-4.7	28	5.1	3.3-7.4	21	2.4	1.5-3.8
2016	43	2.9	2.1-3.9	32	5.3	3.6-7.5	11	1.3	0.6-2.4	36	2.4	1.6-3.3	26	4.2	2.7-6.2	10	1.2	0.6-2.2
2017	42	2.9	2.0-3.9	27	4.3	2.8-6.3	15	1.8	1.0-3.0	23	1.5	0.9-2.3	17	2.8	1.6-4.5	6	0.7	0.2-1.5
2018	55	3.6	2.7-4.7	43	6.9	5.0-9.4	12	1.3	0.7-2.4	35	2.3	1.6-3.2	21	3.3	2.0-5.2	14	1.6	0.9-2.7
2019	62	4.0	3.0-5.1	51	7.9	5.8-10.4	11	1.2	0.6-2.2	37	2.4	1.7-3.3	31	4.9	3.3-7.0	6	0.6	0.2-1.4
2020	38	2.5	1.7-3.4	23	3.9	2.4-5.8	15	1.5	0.8-2.5	30	1.8	1.2-2.6	18	2.7	1.6-4.4	12	1.2	0.6-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Stomach**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	169	14.2	12.1-16.5	89	19.2	15.3-23.8	80	11.0	8.7-13.7	162	13.4	11.4-15.7	92	19.3	15.4-23.9	70	9.7	7.5-12.3
1977	194	16.7	14.4-19.2	116	24.9	20.4-30.0	78	11.3	8.9-14.1	158	13.7	11.6-16.1	82	18.4	14.5-22.9	76	10.8	8.5-13.5
1978	182	15.6	13.4-18.1	99	22.0	17.7-26.8	83	11.6	9.2-14.4	125	11.1	9.2-13.2	59	13.9	10.5-18.0	66	9.4	7.2-12.0
1979	159	13.9	11.8-16.2	83	19.0	15.0-23.6	76	11.0	8.6-13.8	132	11.5	9.6-13.7	69	15.8	12.2-20.1	63	8.9	6.8-11.4
1980	184	16.4	14.1-19.0	97	22.7	18.3-27.8	87	12.6	10.1-15.6	121	10.9	9.0-13.0	66	15.9	12.2-20.3	55	8.0	6.0-10.4
1981	174	15.5	13.3-18.0	109	25.5	20.8-30.8	65	9.2	7.1-11.7	135	12.1	10.1-14.3	89	21.2	17.0-26.2	46	6.6	4.8-8.8
1982	178	16.2	13.9-18.8	99	22.9	18.5-28.0	79	11.6	9.2-14.6	116	10.7	8.8-12.8	64	15.8	12.1-20.2	52	7.7	5.7-10.1
1983	157	14.4	12.2-16.8	75	18.0	14.1-22.7	82	11.9	9.4-14.8	110	10.1	8.3-12.2	62	15.4	11.8-19.8	48	7.0	5.1-9.3
1984	145	13.1	11.0-15.4	76	18.1	14.2-22.7	69	9.8	7.6-12.5	105	9.6	7.8-11.6	47	11.5	8.4-15.3	58	8.2	6.2-10.7
1985	178	16.4	14.0-19.0	105	25.5	20.8-31.0	73	10.5	8.2-13.3	98	9.0	7.3-11.0	59	14.3	10.9-18.5	39	5.7	4.1-7.9
1986	143	13.1	11.0-15.5	71	17.4	13.5-21.9	72	10.4	8.1-13.1	100	9.1	7.4-11.1	51	12.6	9.3-16.6	49	7.1	5.2-9.4
1987	135	12.3	10.3-14.6	71	17.1	13.3-21.6	64	9.3	7.2-12.0	102	9.3	7.6-11.3	49	12.1	8.9-15.9	53	7.6	5.7-10.0
1988	142	13.2	11.1-15.5	75	18.3	14.4-22.9	67	9.7	7.5-12.4	101	9.5	7.7-11.5	50	12.6	9.3-16.6	51	7.4	5.5-9.7
1989	140	12.8	10.7-15.1	73	18.2	14.2-22.9	67	9.5	7.4-12.1	113	10.4	8.5-12.5	62	15.7	12.0-20.2	51	7.2	5.4-9.5
1990	151	14.2	12.0-16.7	68	17.2	13.3-21.8	83	12.4	9.9-15.4	91	8.5	6.8-10.4	47	11.8	8.6-15.6	44	6.5	4.7-8.8
1991	140	13.3	11.2-15.7	77	19.9	15.7-24.9	63	9.2	7.1-11.8	108	10.1	8.3-12.2	60	15.7	11.9-20.2	48	7.0	5.1-9.3
1992	153	14.7	12.4-17.2	80	20.3	16.0-25.3	73	10.9	8.5-13.8	100	9.5	7.7-11.6	53	14.0	10.5-18.3	47	6.8	5.0-9.1
1993	136	13.0	10.9-15.4	68	17.4	13.5-22.1	68	10.3	8.0-13.0	93	8.9	7.2-10.9	41	10.5	7.5-14.3	52	7.8	5.8-10.3
1994	166	15.9	13.6-18.5	81	20.0	15.8-24.9	85	13.1	10.4-16.2	105	10.0	8.2-12.2	60	15.4	11.7-19.9	45	6.8	4.9-9.1
1995	134	12.8	10.7-15.1	73	19.3	15.1-24.3	61	9.2	7.1-11.9	105	10.1	8.2-12.2	48	12.3	9.0-16.3	57	8.7	6.6-11.3
1996	145	13.7	11.6-16.1	67	17.1	13.2-21.7	78	11.8	9.3-14.8	105	10.0	8.2-12.1	47	12.8	9.4-17.0	58	8.8	6.6-11.3
1997	147	13.9	11.8-16.4	65	16.4	12.6-20.9	82	12.3	9.8-15.3	99	9.4	7.6-11.4	44	10.9	7.9-14.6	55	8.3	6.3-10.9
1998	154	14.5	12.3-16.9	90	22.2	17.8-27.3	64	9.9	7.6-12.6	100	9.3	7.6-11.3	54	13.1	9.8-17.1	46	6.9	5.0-9.2
1999	132	12.2	10.2-14.4	71	17.4	13.5-22.0	61	9.1	6.9-11.7	102	9.3	7.6-11.3	57	14.3	10.8-18.6	45	6.6	4.8-8.8
2000	142	13.1	11.0-15.4	84	20.1	16.0-24.9	58	8.6	6.5-11.1	76	7.0	5.5-8.8	47	11.5	8.4-15.2	29	4.2	2.8-6.0
2001	153	13.7	11.6-16.1	85	19.6	15.6-24.3	68	9.8	7.6-12.4	78	7.1	5.6-8.9	38	8.7	6.1-11.9	40	5.8	4.1-7.9
2002	134	11.8	9.9-14.0	59	13.6	10.3-17.6	75	10.7	8.4-13.4	74	6.5	5.1-8.2	44	10.4	7.5-14.0	30	4.2	2.8-6.0
2003	155	13.4	11.4-15.7	83	18.6	14.8-23.2	72	10.2	8.0-12.9	91	8.0	6.5-9.9	47	10.9	8.0-14.5	44	6.3	4.5-8.4
2004	129	11.3	9.5-13.5	67	15.7	12.1-20.0	62	8.9	6.8-11.4	93	8.1	6.6-10.0	38	8.7	6.1-12.0	55	7.8	5.9-10.1
2005	139	12.0	10.1-14.2	70	16.4	12.7-20.7	69	9.6	7.5-12.2	77	6.6	5.2-8.2	44	10.2	7.3-13.7	33	4.6	3.1-6.4
2006	132	11.5	9.6-13.7	68	15.1	11.7-19.3	64	9.1	7.0-11.7	67	5.7	4.4-7.3	35	8.1	5.6-11.3	32	4.3	3.0-6.1
2007	141	11.7	9.9-13.8	73	15.2	11.8-19.2	68	9.2	7.1-11.7	74	6.3	5.0-8.0	34	7.3	5.0-10.2	40	5.4	3.9-7.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Stomach**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	138	11.3	9.5-13.4	79	16.4	12.9-20.5	59	8.2	6.2-10.6	67	5.7	4.4-7.2	33	7.1	4.8-10.0	34	4.8	3.3-6.7
2009	154	12.8	10.8-15.0	75	15.6	12.2-19.6	79	10.7	8.5-13.4	58	4.7	3.5-6.0	33	7.1	4.8-10.0	25	3.2	2.1-4.7
2010	142	11.4	9.6-13.4	75	15.1	11.7-19.0	67	9.0	6.9-11.4	78	6.3	4.9-7.8	31	6.5	4.4-9.3	47	6.2	4.5-8.3
2011	140	11.1	9.4-13.2	65	13.2	10.1-16.8	75	9.9	7.8-12.5	74	5.8	4.6-7.3	38	8.0	5.6-11.1	36	4.7	3.3-6.5
2012	158	12.2	10.3-14.2	85	16.4	13.0-20.3	73	9.5	7.4-11.9	81	6.4	5.0-7.9	37	7.5	5.2-10.3	44	5.5	4.0-7.5
2013	167	12.6	10.7-14.7	98	18.6	15.0-22.7	69	8.6	6.7-10.9	71	5.4	4.2-6.8	36	7.4	5.1-10.2	35	4.3	3.0-6.0
2014	148	10.9	9.2-12.8	78	14.3	11.2-17.9	70	9.0	7.0-11.4	59	4.5	3.4-5.8	39	7.2	5.1-9.9	20	2.7	1.6-4.1
2015	137	9.7	8.1-11.5	60	10.5	7.9-13.6	77	9.2	7.2-11.5	58	4.3	3.2-5.5	29	5.7	3.8-8.2	29	3.4	2.3-4.9
2016	147	10.5	8.8-12.3	82	14.3	11.3-17.8	65	7.9	6.1-10.1	56	3.9	2.9-5.1	31	5.3	3.5-7.5	25	2.9	1.9-4.4
2017	143	9.8	8.3-11.6	82	13.4	10.6-16.7	61	7.3	5.5-9.4	61	4.4	3.3-5.6	35	6.1	4.2-8.5	26	3.3	2.1-4.8
2018	134	9.2	7.7-10.9	74	12.3	9.6-15.5	60	7.2	5.5-9.3	59	4.0	3.1-5.2	37	6.1	4.3-8.5	22	2.6	1.6-4.0
2019	161	11.0	9.4-12.9	82	13.2	10.5-16.5	79	9.5	7.5-11.9	59	4.1	3.1-5.3	41	7.1	5.1-9.7	18	2.1	1.2-3.4
2020	139	9.4	7.9-11.1	83	13.3	10.5-16.5	56	6.5	4.9-8.5	53	3.5	2.6-4.7	34	5.4	3.7-7.6	19	2.1	1.2-3.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Colorectal**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	708	58.6	54.2-63.2	341	71.9	64.1-80.2	367	50.4	45.3-55.9	508	41.9	38.3-45.8	243	50.8	44.4-57.9	265	36.4	32.1-41.2
1977	803	67.7	63.0-72.7	372	82.3	73.9-91.4	431	59.0	53.5-65.0	417	36.5	33.0-40.2	195	44.9	38.6-51.9	222	31.5	27.4-36.0
1978	768	65.6	61.0-70.5	379	84.3	75.8-93.5	389	54.5	49.1-60.3	367	31.8	28.5-35.2	184	41.7	35.7-48.3	183	25.7	22.1-29.8
1979	817	71.5	66.6-76.7	410	93.7	84.6-103.4	407	58.3	52.7-64.4	383	33.5	30.1-37.0	189	43.6	37.5-50.5	194	27.5	23.7-31.7
1980	821	72.1	67.2-77.2	369	86.5	77.7-96.0	452	64.5	58.7-70.9	412	37.1	33.6-40.9	204	50.2	43.4-57.8	208	30.0	26.0-34.4
1981	689	61.9	57.3-66.8	326	78.3	69.8-87.4	363	52.1	46.8-57.8	347	31.4	28.1-34.9	163	40.6	34.4-47.4	184	26.5	22.8-30.7
1982	740	66.3	61.6-71.4	356	85.9	77.0-95.4	384	55.2	49.8-61.1	329	29.7	26.5-33.1	148	35.3	29.7-41.5	181	26.3	22.5-30.4
1983	670	60.2	55.7-65.0	310	74.4	66.2-83.3	360	52.3	47.0-58.1	356	32.2	28.9-35.8	150	37.0	31.2-43.6	206	29.2	25.3-33.5
1984	665	59.8	55.3-64.5	288	71.3	63.1-80.2	377	54.2	48.8-60.0	318	28.4	25.4-31.7	164	40.3	34.2-47.0	154	22.3	18.9-26.2
1985	676	61.2	56.6-66.0	326	79.3	70.8-88.6	350	50.9	45.6-56.6	336	30.4	27.2-33.9	165	40.5	34.5-47.3	171	24.8	21.1-28.8
1986	680	61.6	57.1-66.5	337	81.8	73.1-91.1	343	50.1	44.9-55.8	312	28.3	25.2-31.7	141	35.6	29.8-42.0	171	24.5	20.9-28.5
1987	673	60.6	56.1-65.4	309	74.9	66.7-83.9	364	52.6	47.3-58.4	324	29.0	25.9-32.3	162	39.6	33.6-46.2	162	22.9	19.5-26.8
1988	622	56.3	51.9-60.9	291	72.1	64.0-81.0	331	47.0	42.1-52.4	316	28.7	25.6-32.1	157	39.1	33.1-45.8	159	22.8	19.3-26.6
1989	591	54.7	50.4-59.4	267	67.7	59.7-76.4	324	47.9	42.8-53.5	281	25.9	22.9-29.1	135	34.8	29.1-41.2	146	21.3	17.9-25.1
1990	609	56.8	52.4-61.5	267	67.7	59.7-76.4	342	49.8	44.7-55.5	280	26.4	23.4-29.7	135	36.5	30.5-43.2	145	21.2	17.9-25.0
1991	655	62.3	57.6-67.3	306	80.0	71.2-89.5	349	52.6	47.1-58.4	279	26.2	23.2-29.5	116	30.6	25.2-36.8	163	24.0	20.5-28.1
1992	575	54.7	50.3-59.4	259	66.8	58.8-75.5	316	48.0	42.8-53.6	292	27.7	24.6-31.1	136	36.3	30.4-43.0	156	22.9	19.4-26.8
1993	530	50.7	46.4-55.2	240	61.3	53.6-69.6	290	44.3	39.3-49.7	260	24.8	21.8-28.0	122	31.8	26.3-38.0	138	20.6	17.3-24.4
1994	634	60.7	56.1-65.7	258	66.7	58.7-75.4	376	57.0	51.4-63.1	300	28.5	25.3-31.9	134	34.8	29.0-41.3	166	24.8	21.1-28.9
1995	575	55.0	50.5-59.6	242	60.8	53.3-69.1	333	50.9	45.6-56.7	281	26.9	23.8-30.2	125	33.4	27.8-39.9	156	23.3	19.8-27.3
1996	564	53.7	49.3-58.3	252	64.3	56.5-72.8	312	47.4	42.2-53.0	234	22.3	19.6-25.4	105	26.9	21.9-32.6	129	19.5	16.2-23.1
1997	650	61.8	57.2-66.8	301	75.9	67.4-85.1	349	53.0	47.6-58.9	269	25.5	22.5-28.7	126	32.1	26.6-38.2	143	21.6	18.2-25.5
1998	619	58.6	54.1-63.4	272	67.5	59.6-76.1	347	53.0	47.5-58.9	244	23.2	20.4-26.3	105	27.7	22.6-33.6	139	20.6	17.3-24.4
1999	607	56.1	51.7-60.8	243	60.2	52.7-68.3	364	54.1	48.6-60.0	246	22.9	20.1-26.0	91	22.6	18.1-27.7	155	23.2	19.7-27.2
2000	676	61.4	56.9-66.3	308	74.7	66.4-83.6	368	54.2	48.8-60.0	255	23.1	20.3-26.1	120	29.1	24.0-34.8	135	19.4	16.2-23.0
2001	630	56.5	52.2-61.1	297	69.8	61.9-78.3	333	48.9	43.8-54.5	242	21.7	19.0-24.6	104	25.2	20.5-30.6	138	19.8	16.6-23.4
2002	596	52.8	48.6-57.2	278	61.5	54.4-69.3	318	46.8	41.8-52.2	243	21.6	19.0-24.5	115	26.9	22.1-32.3	128	18.3	15.3-21.8
2003	674	59.7	55.2-64.4	310	72.7	64.7-81.4	364	52.4	47.1-58.1	271	24.3	21.4-27.3	122	29.5	24.4-35.3	149	21.0	17.8-24.7
2004	614	53.2	49.0-57.6	266	59.5	52.4-67.3	348	49.5	44.4-55.0	239	21.2	18.6-24.0	107	25.1	20.5-30.4	132	18.7	15.6-22.2
2005	600	51.9	47.8-56.2	297	66.1	58.6-74.2	303	43.2	38.5-48.4	198	17.3	14.9-19.8	80	19.1	15.0-23.8	118	16.2	13.4-19.4
2006	619	52.8	48.6-57.1	296	65.1	57.7-73.2	323	45.3	40.5-50.6	218	19.1	16.7-21.9	105	25.2	20.5-30.6	113	16.0	13.2-19.2
2007	617	51.3	47.3-55.6	296	61.5	54.5-69.1	321	44.2	39.4-49.3	213	18.2	15.8-20.9	100	22.6	18.3-27.6	113	15.5	12.8-18.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Colorectal**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	604	50.0	46.1-54.2	301	62.7	55.6-70.4	303	41.7	37.1-46.7	229	19.3	16.9-22.0	108	24.3	19.8-29.4	121	16.4	13.6-19.6
2009	576	46.5	42.7-50.5	263	52.6	46.2-59.5	313	42.3	37.7-47.3	222	18.4	16.1-21.0	105	23.0	18.7-27.9	117	15.6	12.9-18.7
2010	572	45.5	41.8-49.4	266	52.9	46.6-59.8	306	41.1	36.6-46.0	213	17.1	14.8-19.6	115	24.3	19.9-29.2	98	12.9	10.4-15.7
2011	500	38.7	35.3-42.3	235	45.3	39.5-51.6	265	34.2	30.1-38.6	199	15.8	13.7-18.2	94	19.0	15.2-23.3	105	13.6	11.1-16.5
2012	536	40.8	37.4-44.5	273	52.3	46.1-59.0	263	33.5	29.5-37.8	236	18.1	15.8-20.6	112	22.0	18.0-26.6	124	15.4	12.8-18.4
2013	575	42.8	39.3-46.5	279	50.5	44.5-56.9	296	37.2	33.0-41.7	209	15.5	13.4-17.8	102	19.2	15.6-23.4	107	12.8	10.4-15.5
2014	580	41.8	38.4-45.4	290	50.9	45.1-57.3	290	35.5	31.5-39.9	212	15.6	13.5-17.8	106	20.1	16.3-24.4	106	12.9	10.5-15.6
2015	527	37.6	34.4-41.0	276	47.1	41.5-53.1	251	30.6	26.9-34.7	197	14.2	12.3-16.4	94	16.8	13.5-20.6	103	12.3	10.0-15.0
2016	545	37.6	34.4-40.9	270	44.3	39.1-50.1	275	33.3	29.4-37.5	197	13.7	11.9-15.8	89	15.6	12.5-19.3	108	12.4	10.1-15.0
2017	515	35.3	32.3-38.5	274	44.5	39.3-50.2	241	28.5	24.9-32.4	196	13.5	11.7-15.6	108	19.0	15.5-23.0	88	10.2	8.1-12.6
2018	528	35.8	32.8-39.1	254	40.7	35.7-46.1	274	32.1	28.4-36.3	167	11.3	9.7-13.2	81	13.7	10.8-17.1	86	9.8	7.8-12.1
2019	561	37.7	34.6-41.0	295	46.9	41.6-52.7	266	30.8	27.2-34.8	187	12.4	10.7-14.4	93	15.1	12.1-18.5	94	10.1	8.1-12.5
2020	457	31.0	28.2-34.0	243	39.0	34.2-44.4	214	25.2	21.8-28.9	155	10.3	8.8-12.1	80	13.3	10.5-16.6	75	8.4	6.6-10.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Colon excluding rectum**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	502	41.5	37.8-45.3	235	49.9	43.5-57.0	267	36.5	32.2-41.3	385	31.6	28.5-35.0	177	36.9	31.4-42.9	208	28.4	24.6-32.7
1977	590	49.7	45.7-54.0	261	58.7	51.5-66.5	329	44.7	39.9-49.9	344	30.3	27.1-33.7	156	36.5	30.8-42.9	188	26.8	23.1-31.0
1978	579	50.0	45.9-54.3	271	61.5	54.2-69.5	308	43.4	38.6-48.6	311	26.8	23.9-30.0	156	35.5	30.0-41.7	155	21.8	18.4-25.5
1979	630	55.3	51.0-59.9	311	71.8	63.8-80.4	319	45.7	40.7-51.1	327	28.6	25.5-31.9	159	37.1	31.4-43.5	168	23.7	20.2-27.7
1980	591	52.0	47.8-56.4	259	61.4	54.0-69.5	332	47.2	42.2-52.7	358	32.3	29.0-35.9	171	42.1	35.9-49.1	187	27.0	23.2-31.3
1981	480	43.0	39.2-47.1	212	51.7	44.8-59.3	268	38.2	33.7-43.1	301	27.1	24.1-30.4	136	33.9	28.3-40.2	165	23.6	20.1-27.6
1982	537	48.1	44.1-52.4	252	61.3	53.8-69.5	285	41.1	36.4-46.3	267	24.1	21.3-27.2	118	28.5	23.5-34.3	149	21.6	18.2-25.4
1983	484	43.3	39.5-47.4	221	53.2	46.2-60.8	263	37.8	33.4-42.8	292	26.3	23.4-29.6	117	28.5	23.5-34.3	175	24.8	21.2-28.8
1984	468	42.0	38.3-46.0	202	50.0	43.2-57.5	266	38.2	33.7-43.2	285	25.4	22.5-28.6	149	36.6	30.9-43.1	136	19.7	16.5-23.3
1985	517	46.9	42.9-51.1	229	56.0	48.9-63.9	288	41.8	37.1-47.0	289	26.2	23.2-29.4	144	35.4	29.7-41.7	145	21.1	17.7-24.8
1986	490	44.6	40.7-48.7	226	55.3	48.2-63.1	264	38.4	33.9-43.4	270	24.5	21.6-27.6	123	30.9	25.6-36.9	147	21.0	17.7-24.7
1987	504	44.9	41.1-49.1	240	57.9	50.7-65.8	264	37.9	33.4-42.8	290	25.9	23.0-29.1	147	35.9	30.2-42.2	143	20.3	17.1-23.9
1988	466	42.0	38.3-46.0	221	55.5	48.3-63.4	245	34.4	30.2-39.0	263	23.9	21.1-27.0	134	33.4	27.9-39.6	129	18.5	15.4-22.1
1989	456	42.2	38.4-46.3	207	53.0	45.9-60.8	249	36.6	32.2-41.5	255	23.4	20.6-26.5	122	31.7	26.2-37.9	133	19.4	16.2-23.0
1990	470	43.8	39.9-48.0	199	51.1	44.1-58.7	271	39.3	34.7-44.3	246	23.1	20.3-26.2	122	33.1	27.4-39.5	124	17.9	14.9-21.4
1991	484	46.0	41.9-50.2	220	58.0	50.5-66.2	264	39.7	35.0-44.8	232	21.7	19.0-24.7	99	26.3	21.3-32.0	133	19.6	16.4-23.2
1992	416	39.5	35.8-43.5	188	49.5	42.5-57.1	228	34.4	30.0-39.2	241	22.9	20.1-26.0	109	29.2	23.9-35.2	132	19.4	16.2-23.1
1993	396	37.9	34.2-41.8	170	43.9	37.5-51.1	226	34.5	30.1-39.3	225	21.5	18.7-24.5	104	27.4	22.3-33.2	121	18.1	15.0-21.7
1994	485	46.4	42.3-50.7	194	50.6	43.6-58.2	291	43.7	38.8-49.0	263	24.8	21.9-28.0	122	31.8	26.3-38.0	141	20.9	17.6-24.7
1995	430	41.1	37.3-45.2	184	46.4	39.9-53.7	246	37.4	32.8-42.4	250	23.9	21.0-27.1	110	29.3	24.0-35.3	140	20.9	17.6-24.7
1996	425	40.4	36.7-44.5	188	48.3	41.5-55.7	237	36.0	31.5-40.9	202	19.3	16.7-22.1	92	23.9	19.2-29.3	110	16.5	13.5-19.9
1997	481	45.8	41.8-50.1	219	55.5	48.2-63.4	262	39.9	35.2-45.1	233	22.0	19.3-25.1	110	28.1	23.0-34.0	123	18.8	15.6-22.4
1998	470	44.6	40.7-48.9	199	50.3	43.5-57.9	271	41.2	36.4-46.4	217	20.7	18.0-23.6	93	24.9	20.0-30.5	124	18.5	15.4-22.1
1999	450	41.6	37.9-45.7	169	42.4	36.1-49.3	281	41.5	36.8-46.7	215	20.0	17.4-22.8	76	18.9	14.8-23.7	139	20.8	17.4-24.6
2000	508	46.3	42.3-50.5	219	54.0	47.0-61.7	289	42.5	37.7-47.7	230	20.8	18.2-23.7	109	26.6	21.8-32.1	121	17.4	14.5-20.9
2001	465	41.9	38.1-45.9	219	52.2	45.3-59.6	246	35.8	31.5-40.6	203	18.2	15.8-20.9	85	20.7	16.5-25.6	118	16.9	14.0-20.3
2002	435	38.7	35.2-42.5	190	42.7	36.7-49.3	245	36.0	31.6-40.8	216	19.2	16.7-21.9	104	24.3	19.8-29.5	112	16.0	13.1-19.2
2003	502	44.6	40.8-48.7	218	51.9	45.1-59.4	284	40.9	36.3-46.0	232	20.9	18.3-23.7	102	25.3	20.5-30.7	130	18.4	15.4-21.9
2004	450	39.2	35.6-43.0	197	44.9	38.7-51.7	253	35.9	31.6-40.6	208	18.4	16.0-21.1	93	21.8	17.5-26.7	115	16.3	13.4-19.6
2005	469	40.7	37.0-44.5	224	50.5	43.9-57.7	245	35.0	30.7-39.7	172	15.0	12.9-17.5	64	15.3	11.7-19.6	108	14.8	12.2-17.9
2006	451	38.7	35.2-42.5	204	45.8	39.5-52.6	247	34.8	30.6-39.5	192	16.9	14.6-19.5	91	22.4	17.9-27.5	101	14.4	11.7-17.5
2007	448	37.4	34.0-41.1	210	44.2	38.2-50.7	238	32.8	28.7-37.3	186	15.9	13.7-18.4	84	19.0	15.1-23.6	102	14.0	11.4-17.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Colon excluding rectum**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	425	35.6	32.3-39.2	207	44.4	38.4-51.0	218	30.2	26.3-34.5	200	16.9	14.6-19.4	94	21.1	17.0-25.9	106	14.4	11.7-17.4
2009	403	32.8	29.7-36.2	182	37.4	32.0-43.4	221	29.6	25.8-33.9	186	15.5	13.3-17.9	87	19.4	15.4-24.0	99	13.2	10.7-16.1
2010	412	33.0	29.9-36.4	192	38.5	33.1-44.5	220	29.8	26.0-34.0	176	14.3	12.2-16.5	88	19.0	15.1-23.5	88	11.6	9.3-14.4
2011	361	28.1	25.2-31.2	168	32.8	27.9-38.3	193	24.9	21.5-28.7	170	13.6	11.6-15.8	78	16.0	12.5-20.0	92	11.8	9.5-14.6
2012	392	30.2	27.2-33.4	195	38.3	32.9-44.2	197	25.2	21.7-29.0	197	15.2	13.2-17.5	90	18.0	14.4-22.2	107	13.2	10.8-16.1
2013	406	30.5	27.5-33.6	197	36.4	31.4-42.0	209	26.3	22.8-30.1	180	13.5	11.6-15.6	85	16.4	13.0-20.4	95	11.4	9.2-14.0
2014	421	30.6	27.7-33.7	203	36.7	31.7-42.3	218	26.5	23.1-30.3	174	12.6	10.7-14.6	84	15.7	12.4-19.5	90	10.8	8.6-13.3
2015	367	26.4	23.8-29.3	186	32.6	27.9-37.7	181	22.1	18.9-25.6	165	11.9	10.1-13.9	77	13.9	10.9-17.4	88	10.5	8.4-13.0
2016	370	25.7	23.1-28.5	185	30.7	26.4-35.6	185	22.4	19.3-26.0	160	11.2	9.5-13.1	65	11.3	8.7-14.5	95	11.1	9.0-13.6
2017	367	25.2	22.6-27.9	187	30.5	26.2-35.3	180	21.3	18.2-24.7	150	10.3	8.7-12.1	77	13.7	10.7-17.1	73	8.4	6.5-10.6
2018	370	25.0	22.5-27.7	168	26.9	22.9-31.4	202	23.4	20.2-26.9	141	9.5	8.0-11.3	65	10.9	8.4-14.0	76	8.5	6.7-10.7
2019	395	26.6	24.0-29.4	205	32.9	28.5-37.8	190	22.0	19.0-25.5	159	10.5	8.9-12.2	75	12.1	9.5-15.2	84	8.9	7.1-11.1
2020	325	22.1	19.7-24.6	163	26.3	22.4-30.8	162	19.0	16.1-22.2	133	8.9	7.4-10.6	68	11.3	8.7-14.4	65	7.4	5.7-9.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Rectum & rectosigmoid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	206	17.1	14.8-19.7	106	21.9	17.8-26.7	100	13.8	11.2-16.9	123	10.3	8.5-12.3	66	14.0	10.7-17.9	57	8.0	6.0-10.4
1977	213	18.0	15.6-20.6	111	23.6	19.3-28.6	102	14.4	11.7-17.5	73	6.2	4.8-7.8	39	8.4	5.9-11.6	34	4.7	3.2-6.6
1978	189	15.7	13.5-18.1	108	22.8	18.6-27.7	81	11.1	8.8-13.8	56	4.9	3.7-6.4	28	6.2	4.0-9.0	28	4.0	2.6-5.8
1979	187	16.2	13.9-18.7	99	21.9	17.7-26.7	88	12.6	10.1-15.6	56	4.9	3.7-6.3	30	6.5	4.3-9.4	26	3.7	2.4-5.5
1980	230	20.1	17.5-22.9	110	25.1	20.5-30.4	120	17.3	14.3-20.8	54	4.8	3.6-6.3	33	8.1	5.5-11.4	21	3.0	1.8-4.6
1981	209	18.9	16.4-21.7	114	26.6	21.8-32.0	95	13.9	11.2-17.1	46	4.2	3.1-5.7	27	6.7	4.4-9.8	19	2.9	1.7-4.5
1982	203	18.2	15.8-20.9	104	24.6	20.0-29.8	99	14.1	11.5-17.2	62	5.6	4.3-7.2	30	6.7	4.5-9.6	32	4.7	3.2-6.7
1983	186	17.0	14.6-19.6	89	21.2	17.0-26.2	97	14.5	11.7-17.7	64	5.9	4.5-7.5	33	8.5	5.8-11.9	31	4.4	3.0-6.3
1984	197	17.8	15.3-20.4	86	21.3	17.0-26.3	111	15.9	13.1-19.2	33	3.0	2.0-4.2	15	3.7	2.0-6.0	18	2.7	1.6-4.2
1985	159	14.3	12.2-16.8	97	23.3	18.8-28.5	62	9.1	6.9-11.7	47	4.3	3.1-5.7	21	5.1	3.2-7.9	26	3.7	2.4-5.4
1986	190	17.1	14.7-19.7	111	26.4	21.6-31.9	79	11.7	9.2-14.6	42	3.9	2.8-5.2	18	4.7	2.7-7.3	24	3.5	2.2-5.2
1987	169	15.7	13.4-18.3	69	17.0	13.2-21.6	100	14.7	11.9-18.0	34	3.0	2.1-4.3	15	3.7	2.1-6.1	19	2.6	1.6-4.1
1988	156	14.3	12.1-16.7	70	16.6	12.9-21.0	86	12.6	10.1-15.6	53	4.8	3.6-6.3	23	5.7	3.6-8.6	30	4.2	2.8-6.0
1989	135	12.5	10.5-14.8	60	14.7	11.1-18.9	75	11.3	8.9-14.2	26	2.4	1.6-3.6	13	3.1	1.6-5.3	13	1.9	1.0-3.3
1990	139	13.0	10.9-15.4	68	16.6	12.8-21.1	71	10.6	8.2-13.4	34	3.3	2.3-4.6	13	3.4	1.8-5.8	21	3.3	2.0-5.1
1991	171	16.3	14.0-19.0	86	22.0	17.5-27.2	85	12.9	10.3-16.0	47	4.4	3.3-5.9	17	4.3	2.5-6.9	30	4.5	3.0-6.4
1992	159	15.2	12.9-17.8	71	17.3	13.5-21.9	88	13.6	10.9-16.8	51	4.8	3.6-6.3	27	7.1	4.7-10.3	24	3.5	2.2-5.2
1993	134	12.8	10.7-15.1	70	17.3	13.5-21.9	64	9.8	7.5-12.6	35	3.3	2.3-4.6	18	4.4	2.6-7.0	17	2.5	1.4-4.0
1994	149	14.4	12.2-16.9	64	16.1	12.4-20.6	85	13.3	10.6-16.5	37	3.6	2.6-5.0	12	3.0	1.5-5.2	25	3.9	2.5-5.7
1995	145	13.8	11.7-16.3	58	14.4	10.9-18.6	87	13.5	10.8-16.7	31	3.0	2.0-4.2	15	4.1	2.3-6.7	16	2.4	1.4-3.9
1996	139	13.2	11.1-15.6	64	16.0	12.3-20.5	75	11.4	8.9-14.3	32	3.1	2.1-4.3	13	3.1	1.6-5.2	19	3.0	1.8-4.7
1997	169	16.0	13.7-18.6	82	20.4	16.2-25.4	87	13.1	10.5-16.2	36	3.5	2.4-4.8	16	3.9	2.2-6.3	20	2.8	1.7-4.4
1998	149	14.0	11.8-16.4	73	17.2	13.4-21.6	76	11.8	9.3-14.7	27	2.6	1.7-3.7	12	2.8	1.4-4.9	15	2.2	1.2-3.6
1999	157	14.5	12.3-16.9	74	17.8	13.9-22.4	83	12.6	10.0-15.6	31	2.9	2.0-4.1	15	3.7	2.0-6.0	16	2.5	1.4-4.0
2000	168	15.2	12.9-17.6	89	20.7	16.5-25.5	79	11.7	9.3-14.6	25	2.3	1.5-3.3	11	2.5	1.2-4.4	14	2.0	1.1-3.3
2001	165	14.6	12.5-17.0	78	17.6	13.9-22.0	87	13.1	10.5-16.1	39	3.4	2.4-4.7	19	4.5	2.7-7.0	20	2.9	1.8-4.5
2002	161	14.1	12.0-16.4	88	18.9	15.1-23.3	73	10.8	8.5-13.6	27	2.4	1.6-3.5	11	2.6	1.3-4.6	16	2.4	1.4-3.9
2003	172	15.1	12.9-17.5	92	20.8	16.7-25.6	80	11.5	9.1-14.3	39	3.4	2.4-4.6	20	4.3	2.6-6.6	19	2.6	1.6-4.1
2004	164	14.0	12.0-16.4	69	14.6	11.3-18.6	95	13.6	11.0-16.7	31	2.8	1.9-3.9	14	3.3	1.8-5.5	17	2.4	1.4-3.9
2005	131	11.2	9.3-13.3	73	15.6	12.1-19.6	58	8.3	6.3-10.7	26	2.2	1.4-3.2	16	3.8	2.1-6.1	10	1.3	0.6-2.4
2006	168	14.1	12.0-16.4	92	19.3	15.5-23.8	76	10.5	8.2-13.1	26	2.2	1.4-3.2	14	2.9	1.5-4.8	12	1.6	0.8-2.8
2007	169	13.9	11.9-16.2	86	17.3	13.8-21.5	83	11.4	9.0-14.1	27	2.3	1.5-3.4	16	3.6	2.0-5.8	11	1.5	0.8-2.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Rectum & rectosigmoid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	179	14.4	12.4-16.7	94	18.3	14.7-22.5	85	11.5	9.2-14.3	29	2.5	1.6-3.5	14	3.1	1.7-5.2	15	2.0	1.1-3.3
2009	173	13.7	11.7-15.9	81	15.1	11.9-18.9	92	12.6	10.2-15.5	36	2.9	2.0-4.0	18	3.6	2.1-5.8	18	2.5	1.4-3.9
2010	160	12.5	10.6-14.6	74	14.4	11.2-18.1	86	11.3	9.0-14.0	37	2.8	2.0-3.9	27	5.3	3.4-7.7	10	1.2	0.6-2.3
2011	139	10.6	8.9-12.5	67	12.4	9.6-15.8	72	9.2	7.2-11.7	29	2.3	1.5-3.3	16	3.0	1.7-4.9	13	1.8	0.9-3.0
2012	144	10.7	9.0-12.6	78	14.0	11.0-17.5	66	8.3	6.4-10.6	39	2.9	2.0-3.9	22	4.0	2.5-6.2	17	2.1	1.2-3.4
2013	169	12.3	10.5-14.3	82	14.0	11.1-17.5	87	10.9	8.7-13.5	29	2.0	1.3-2.9	17	2.8	1.6-4.5	12	1.4	0.7-2.5
2014	159	11.2	9.5-13.1	87	14.2	11.3-17.6	72	9.0	7.0-11.3	38	3.0	2.1-4.1	22	4.4	2.7-6.6	16	2.1	1.2-3.4
2015	160	11.1	9.5-13.0	90	14.5	11.6-17.9	70	8.6	6.6-10.9	32	2.3	1.6-3.3	17	2.9	1.7-4.7	15	1.8	1.0-2.9
2016	175	11.9	10.2-13.8	85	13.6	10.8-16.9	90	10.9	8.7-13.4	37	2.5	1.8-3.5	24	4.3	2.7-6.4	13	1.3	0.7-2.3
2017	148	10.1	8.5-11.9	87	14.0	11.2-17.3	61	7.2	5.5-9.3	46	3.2	2.3-4.3	31	5.3	3.5-7.5	15	1.8	1.0-3.0
2018	158	10.8	9.2-12.7	86	13.8	11.0-17.1	72	8.7	6.8-11.0	26	1.8	1.2-2.7	16	2.8	1.6-4.5	10	1.2	0.6-2.3
2019	166	11.1	9.4-12.9	90	14.0	11.2-17.3	76	8.8	6.9-11.0	28	2.0	1.3-2.8	18	3.0	1.7-4.7	10	1.2	0.6-2.2
2020	132	8.9	7.4-10.6	80	12.7	10.0-15.9	52	6.2	4.6-8.2	22	1.5	0.9-2.2	12	2.0	1.0-3.5	10	1.0	0.5-2.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Liver / intrahepatic bile duct**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	32	2.7	1.8-3.8	17	4.0	2.3-6.5	15	2.0	1.1-3.3	33	2.7	1.9-3.8	14	3.3	1.8-5.6	19	2.5	1.5-4.0
1977	46	3.7	2.7-5.0	31	6.2	4.2-8.9	15	2.0	1.1-3.4	61	5.1	3.9-6.6	39	8.1	5.7-11.2	22	3.3	2.0-5.0
1978	52	4.5	3.4-6.0	32	7.1	4.8-10.1	20	2.7	1.7-4.3	71	6.2	4.8-7.9	36	7.7	5.3-10.7	35	4.9	3.4-6.9
1979	53	4.7	3.5-6.2	31	6.6	4.5-9.5	22	3.3	2.1-5.0	52	4.6	3.4-6.0	28	6.2	4.1-8.9	24	3.6	2.3-5.4
1980	47	4.1	3.0-5.5	28	6.3	4.1-9.1	19	2.8	1.7-4.5	46	4.1	3.0-5.5	24	5.6	3.5-8.3	22	3.3	2.0-5.0
1981	52	4.7	3.5-6.2	34	7.9	5.4-11.1	18	2.5	1.5-4.0	53	4.8	3.6-6.3	32	7.3	5.0-10.4	21	3.1	1.9-4.7
1982	47	4.4	3.2-5.9	36	8.5	5.9-11.8	11	1.7	0.8-3.1	50	4.5	3.4-6.0	28	6.6	4.3-9.5	22	3.2	2.0-4.9
1983	52	4.6	3.4-6.0	33	7.4	5.1-10.5	19	2.7	1.6-4.3	46	4.2	3.0-5.6	24	5.5	3.5-8.2	22	3.2	2.0-4.8
1984	44	3.9	2.8-5.3	29	6.4	4.2-9.2	15	2.3	1.3-3.8	57	4.9	3.7-6.4	33	7.5	5.1-10.6	24	3.4	2.2-5.1
1985	63	5.9	4.5-7.5	33	7.8	5.3-11.0	30	4.5	3.0-6.4	52	4.7	3.5-6.2	28	6.1	4.0-8.9	24	3.6	2.3-5.4
1986	57	5.3	4.0-6.8	33	8.0	5.5-11.3	24	3.5	2.2-5.2	56	5.2	3.9-6.7	31	7.9	5.3-11.2	25	3.5	2.3-5.2
1987	48	4.3	3.1-5.7	31	7.0	4.7-9.9	17	2.5	1.4-4.0	70	6.4	4.9-8.0	38	9.0	6.3-12.4	32	4.8	3.3-6.8
1988	50	4.6	3.4-6.1	31	7.6	5.1-10.8	19	2.8	1.7-4.4	46	4.3	3.1-5.7	22	5.4	3.3-8.1	24	3.6	2.3-5.4
1989	53	4.8	3.6-6.3	30	6.9	4.6-9.8	23	3.3	2.1-5.0	55	5.1	3.8-6.7	30	7.1	4.7-10.1	25	3.9	2.5-5.8
1990	51	4.9	3.6-6.4	37	9.0	6.3-12.3	14	2.2	1.2-3.7	50	4.7	3.5-6.2	30	7.5	5.0-10.7	20	3.0	1.8-4.7
1991	62	5.9	4.5-7.5	44	10.4	7.5-13.9	18	2.7	1.6-4.3	70	6.5	5.1-8.3	48	11.8	8.6-15.6	22	3.3	2.1-5.0
1992	74	7.1	5.6-8.9	48	11.7	8.6-15.6	26	4.1	2.6-6.0	70	6.8	5.3-8.5	45	11.2	8.1-15.0	25	3.9	2.5-5.7
1993	57	5.3	4.0-6.9	33	7.6	5.2-10.6	24	3.6	2.3-5.4	70	6.5	5.1-8.3	41	9.6	6.8-13.0	29	4.3	2.9-6.2
1994	74	7.2	5.6-9.0	50	11.9	8.8-15.6	24	3.7	2.4-5.5	62	6.1	4.6-7.8	37	9.3	6.5-12.8	25	3.9	2.5-5.8
1995	100	9.5	7.7-11.6	74	18.4	14.3-23.1	26	4.0	2.6-5.8	75	7.2	5.7-9.1	49	12.0	8.8-15.8	26	4.0	2.6-5.9
1996	90	8.6	6.9-10.5	64	14.8	11.3-18.9	26	4.0	2.6-5.9	72	6.9	5.4-8.7	46	11.3	8.2-15.0	26	4.0	2.6-5.9
1997	85	8.1	6.4-10.0	60	13.8	10.5-17.7	25	3.9	2.5-5.7	46	4.4	3.2-5.9	34	8.2	5.6-11.4	12	1.9	1.0-3.2
1998	93	8.7	7.0-10.7	60	13.4	10.2-17.2	33	5.0	3.4-7.1	68	6.3	4.9-8.0	39	8.6	6.1-11.8	29	4.3	2.9-6.2
1999	97	8.8	7.2-10.8	64	13.7	10.5-17.4	33	5.0	3.4-7.0	56	5.2	3.9-6.7	33	7.7	5.3-10.8	23	3.3	2.1-5.0
2000	101	9.0	7.3-10.9	73	15.6	12.2-19.6	28	4.1	2.7-6.0	71	6.4	5.0-8.1	45	9.6	7.0-12.8	26	4.0	2.6-5.8
2001	124	10.9	9.1-13.0	82	17.5	13.8-21.7	42	6.3	4.5-8.4	79	7.0	5.6-8.8	44	9.4	6.8-12.7	35	5.0	3.5-7.0
2002	125	10.9	9.0-13.0	85	17.4	13.8-21.5	40	6.0	4.3-8.2	85	7.4	5.9-9.1	53	10.8	8.1-14.2	32	4.7	3.2-6.7
2003	133	11.2	9.4-13.3	89	17.9	14.3-22.0	44	6.3	4.6-8.5	103	8.9	7.3-10.8	63	13.6	10.4-17.4	40	5.7	4.1-7.8
2004	175	14.7	12.6-17.0	130	26.0	21.7-31.0	45	6.4	4.6-8.5	94	8.0	6.5-9.8	64	13.2	10.1-16.9	30	4.3	2.9-6.1
2005	163	13.7	11.6-15.9	129	25.4	21.1-30.2	34	5.0	3.5-7.0	109	9.4	7.7-11.3	81	17.0	13.4-21.2	28	4.0	2.6-5.8
2006	172	14.0	12.0-16.3	123	23.7	19.6-28.3	49	6.9	5.1-9.2	113	9.4	7.8-11.4	78	15.4	12.1-19.3	35	5.0	3.5-7.0
2007	177	14.1	12.1-16.4	131	24.5	20.4-29.2	46	6.5	4.7-8.7	115	9.4	7.7-11.3	79	15.0	11.8-18.8	36	5.2	3.6-7.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Liver / intrahepatic bile duct**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	187	14.9	12.8-17.2	138	25.7	21.5-30.5	49	7.0	5.1-9.2	104	8.2	6.7-9.9	78	14.5	11.4-18.2	26	3.6	2.3-5.2
2009	224	17.4	15.1-19.8	161	29.9	25.3-35.0	63	8.5	6.6-10.9	129	10.3	8.6-12.2	92	18.1	14.4-22.3	37	5.0	3.5-6.9
2010	222	16.5	14.4-18.9	170	29.9	25.4-34.9	52	6.8	5.1-9.0	118	9.2	7.6-11.1	83	15.8	12.5-19.7	35	4.6	3.2-6.4
2011	230	16.9	14.8-19.3	157	26.9	22.8-31.6	73	9.5	7.4-11.9	116	8.7	7.2-10.5	81	14.5	11.4-18.1	35	4.6	3.2-6.4
2012	236	17.1	14.9-19.4	174	29.5	25.1-34.3	62	8.0	6.1-10.3	144	10.5	8.8-12.3	98	17.1	13.8-21.0	46	5.6	4.1-7.5
2013	242	16.9	14.8-19.2	165	26.8	22.8-31.3	77	9.4	7.4-11.8	131	9.4	7.8-11.2	94	16.3	13.1-20.1	37	4.6	3.2-6.4
2014	248	16.8	14.7-19.0	174	27.5	23.5-32.0	74	8.8	6.9-11.1	144	9.9	8.3-11.7	98	15.5	12.5-19.0	46	5.6	4.1-7.5
2015	260	17.4	15.3-19.7	195	30.9	26.6-35.7	65	7.6	5.9-9.7	140	9.5	8.0-11.3	96	15.8	12.7-19.3	44	5.0	3.6-6.8
2016	229	14.9	13.0-17.0	157	23.6	20.0-27.7	72	8.2	6.4-10.4	170	11.6	9.9-13.5	118	19.3	15.9-23.2	52	6.0	4.5-8.0
2017	235	15.6	13.6-17.8	170	26.4	22.5-30.8	65	7.5	5.8-9.6	116	7.7	6.4-9.3	82	13.2	10.4-16.5	34	3.7	2.5-5.2
2018	221	14.3	12.5-16.4	143	21.5	18.1-25.5	78	8.7	6.9-10.9	125	8.3	6.8-9.9	86	13.5	10.7-16.7	39	4.4	3.1-6.0
2019	223	13.9	12.1-15.9	160	23.7	20.1-27.8	63	6.8	5.2-8.8	118	7.5	6.2-9.0	73	10.9	8.5-13.8	45	4.9	3.6-6.6
2020	183	11.5	9.8-13.3	126	18.8	15.6-22.4	57	6.2	4.7-8.1	147	9.2	7.8-10.8	90	13.7	11.0-16.9	57	6.0	4.5-7.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Pancreas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	161	13.2	11.2-15.5	80	16.3	12.9-20.4	81	11.1	8.8-13.9	163	13.5	11.4-15.7	86	18.0	14.3-22.4	77	10.5	8.2-13.2
1977	153	13.0	11.0-15.2	76	16.6	13.0-20.9	77	10.8	8.5-13.5	140	11.7	9.8-13.8	67	14.0	10.8-18.0	73	10.1	7.9-12.8
1978	135	11.5	9.6-13.6	78	16.8	13.2-21.1	57	8.0	6.1-10.5	124	10.6	8.8-12.7	61	13.5	10.3-17.5	63	8.9	6.9-11.5
1979	149	13.1	11.0-15.4	71	16.6	12.9-21.0	78	10.9	8.6-13.7	141	12.4	10.4-14.6	65	15.0	11.5-19.2	76	10.7	8.4-13.5
1980	136	12.0	10.1-14.3	73	17.1	13.3-21.5	63	8.9	6.8-11.4	126	10.9	9.1-13.0	67	15.0	11.6-19.2	59	8.3	6.3-10.8
1981	133	11.8	9.9-14.0	61	13.8	10.5-17.8	72	10.3	8.0-13.0	134	11.9	10.0-14.2	68	15.5	12.0-19.8	66	9.6	7.4-12.2
1982	136	12.1	10.2-14.4	58	14.1	10.6-18.2	78	11.2	8.8-14.0	125	11.0	9.1-13.1	56	12.8	9.6-16.7	69	9.8	7.6-12.5
1983	133	12.0	10.0-14.2	67	15.2	11.7-19.4	66	9.8	7.5-12.5	117	10.4	8.6-12.5	60	13.6	10.4-17.6	57	8.3	6.3-10.8
1984	152	13.9	11.7-16.3	64	15.6	11.9-19.9	88	12.7	10.2-15.7	117	10.6	8.8-12.8	54	12.4	9.3-16.2	63	9.2	7.1-11.8
1985	112	10.2	8.4-12.2	54	12.9	9.6-16.9	58	8.3	6.3-10.8	108	9.8	8.0-11.8	61	15.2	11.5-19.5	47	6.8	5.0-9.1
1986	131	11.8	9.8-14.0	53	12.5	9.3-16.4	78	11.0	8.7-13.7	97	8.6	7.0-10.5	44	10.3	7.5-13.9	53	7.4	5.5-9.7
1987	144	13.0	10.9-15.3	68	16.4	12.6-20.8	76	10.7	8.4-13.4	126	11.4	9.5-13.6	56	13.8	10.4-18.0	70	9.8	7.6-12.4
1988	137	12.5	10.5-14.8	66	15.7	12.1-19.9	71	10.3	8.0-13.0	114	10.5	8.6-12.6	53	13.1	9.8-17.1	61	8.6	6.6-11.1
1989	128	11.9	9.9-14.1	50	12.5	9.2-16.5	78	11.4	9.0-14.3	126	11.7	9.7-13.9	61	14.9	11.4-19.2	65	9.6	7.4-12.3
1990	114	10.7	8.8-12.8	45	11.6	8.4-15.5	69	9.9	7.7-12.5	108	10.1	8.3-12.2	37	9.5	6.7-13.1	71	10.2	7.9-12.8
1991	126	11.9	9.9-14.1	56	13.8	10.3-17.9	70	10.4	8.1-13.1	110	10.4	8.5-12.5	55	13.7	10.3-17.9	55	8.2	6.1-10.7
1992	120	11.5	9.5-13.8	52	13.6	10.1-17.8	68	10.1	7.8-12.8	119	11.5	9.5-13.7	43	11.5	8.3-15.5	76	11.3	8.9-14.2
1993	152	14.6	12.4-17.1	50	12.9	9.5-17.0	102	15.4	12.5-18.7	120	11.5	9.5-13.7	44	11.8	8.6-15.9	76	11.1	8.8-14.0
1994	125	12.1	10.1-14.4	52	13.2	9.8-17.2	73	11.1	8.6-13.9	121	11.7	9.7-14.0	47	11.9	8.7-15.8	74	11.3	8.9-14.2
1995	125	12.0	10.0-14.3	48	12.3	9.0-16.3	77	11.8	9.3-14.7	119	11.4	9.5-13.7	50	12.4	9.2-16.4	69	10.5	8.2-13.3
1996	138	13.2	11.1-15.6	58	14.7	11.1-19.0	80	12.2	9.7-15.2	116	11.1	9.2-13.3	44	11.7	8.4-15.7	72	10.9	8.5-13.7
1997	121	11.6	9.6-13.8	46	11.6	8.4-15.4	75	11.4	9.0-14.3	103	9.8	8.0-11.9	40	10.0	7.1-13.6	63	9.5	7.3-12.1
1998	139	13.2	11.1-15.5	49	12.4	9.1-16.4	90	13.6	11.0-16.8	113	10.8	8.9-13.0	45	11.6	8.4-15.5	68	10.2	7.9-13.0
1999	153	14.2	12.0-16.6	67	15.9	12.2-20.2	86	13.0	10.4-16.0	126	11.7	9.7-13.9	49	12.2	9.0-16.1	77	11.5	9.0-14.3
2000	129	11.8	9.9-14.0	50	12.1	9.0-16.0	79	11.4	9.0-14.3	94	8.6	7.0-10.5	37	8.8	6.1-12.1	57	8.3	6.3-10.7
2001	143	13.0	11.0-15.3	61	13.8	10.5-17.7	82	12.1	9.6-15.0	110	9.9	8.2-12.0	50	11.8	8.7-15.6	60	8.7	6.7-11.3
2002	130	11.4	9.6-13.6	49	10.7	7.9-14.2	81	11.4	9.0-14.2	117	10.4	8.6-12.4	48	10.6	7.8-14.1	69	9.9	7.7-12.5
2003	160	14.2	12.0-16.5	71	16.4	12.7-20.7	89	12.6	10.1-15.5	110	9.8	8.0-11.8	44	10.2	7.4-13.7	66	9.1	7.0-11.6
2004	152	13.3	11.3-15.6	68	15.0	11.6-19.1	84	11.9	9.5-14.8	114	10.0	8.3-12.0	49	11.1	8.2-14.8	65	9.2	7.1-11.7
2005	157	13.6	11.5-15.9	71	15.6	12.1-19.7	86	12.3	9.8-15.2	132	11.4	9.6-13.6	63	14.1	10.7-18.1	69	9.8	7.6-12.4
2006	120	10.4	8.6-12.4	56	12.5	9.4-16.3	64	9.1	7.0-11.6	118	10.2	8.5-12.3	45	10.4	7.5-13.9	73	10.4	8.2-13.1
2007	174	14.6	12.5-16.9	78	15.3	12.0-19.2	96	13.6	11.0-16.6	124	10.5	8.7-12.5	52	10.6	7.9-14.0	72	9.9	7.8-12.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Pancreas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	164	13.7	11.7-16.0	57	12.1	9.1-15.8	107	14.7	12.0-17.8	124	10.4	8.7-12.4	49	9.9	7.3-13.1	75	10.2	8.0-12.8
2009	149	12.2	10.3-14.4	65	13.7	10.5-17.5	84	11.2	8.9-13.9	126	10.3	8.6-12.3	62	13.4	10.2-17.2	64	8.6	6.6-11.0
2010	145	11.5	9.7-13.6	67	13.3	10.2-17.0	78	10.3	8.1-12.9	111	9.1	7.4-11.0	48	9.8	7.2-13.1	63	8.5	6.5-10.9
2011	204	16.0	13.9-18.4	89	17.5	13.9-21.6	115	14.9	12.3-18.0	151	12.0	10.1-14.0	73	14.5	11.3-18.4	78	10.1	7.9-12.6
2012	203	15.7	13.6-18.0	97	19.0	15.3-23.3	106	13.8	11.3-16.7	143	11.0	9.2-12.9	65	12.4	9.5-15.9	78	10.0	7.9-12.6
2013	190	13.8	11.9-16.0	85	14.5	11.5-18.1	105	12.9	10.6-15.7	122	9.1	7.5-10.8	59	11.2	8.4-14.5	63	7.9	6.1-10.2
2014	172	12.5	10.6-14.5	73	12.9	10.0-16.3	99	12.1	9.8-14.8	167	12.1	10.3-14.1	67	12.3	9.5-15.7	100	12.0	9.7-14.6
2015	202	14.4	12.4-16.5	97	16.5	13.3-20.2	105	12.8	10.4-15.5	130	9.2	7.7-11.0	60	10.3	7.8-13.4	70	8.6	6.7-10.8
2016	198	13.9	12.0-16.1	107	18.0	14.7-21.9	91	10.9	8.7-13.4	167	11.8	10.0-13.7	87	15.0	11.9-18.6	80	9.5	7.5-11.9
2017	181	12.3	10.5-14.2	84	14.4	11.4-17.9	97	10.9	8.8-13.3	144	10.0	8.4-11.8	69	11.9	9.2-15.2	75	8.6	6.7-10.8
2018	207	13.8	12.0-15.9	112	18.1	14.8-21.8	95	10.7	8.6-13.1	145	9.6	8.1-11.3	81	13.8	10.9-17.2	64	6.9	5.3-8.9
2019	224	14.3	12.5-16.4	110	17.0	13.9-20.5	114	12.4	10.2-15.0	146	9.5	8.0-11.2	67	10.7	8.2-13.6	79	8.5	6.7-10.7
2020	194	12.5	10.8-14.4	106	16.2	13.2-19.6	88	10.0	8.0-12.4	135	8.5	7.1-10.1	75	11.3	8.9-14.3	60	6.7	5.1-8.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Larynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	64	4.9	3.8-6.4	58	11.1	8.3-14.4	6	0.8	0.3-1.8	33	2.9	2.0-4.1	27	6.0	3.9-8.8	6	0.8	0.3-1.8
1977	72	6.2	4.8-7.8	56	11.7	8.8-15.3	16	2.4	1.4-4.0	27	2.2	1.4-3.2	21	4.0	2.4-6.2	6	0.9	0.3-1.9
1978	59	5.4	4.1-7.0	43	9.9	7.1-13.3	16	2.4	1.4-4.0	26	2.2	1.4-3.2	20	4.3	2.6-6.7	6	0.9	0.3-1.9
1979	77	6.7	5.2-8.4	63	13.3	10.2-17.1	14	2.3	1.2-3.9	35	3.0	2.1-4.2	28	6.1	4.0-8.9	7	1.1	0.4-2.3
1980	66	5.8	4.5-7.4	56	12.0	9.0-15.6	10	1.7	0.8-3.1	27	2.5	1.6-3.7	23	5.3	3.3-7.9	4	0.7	0.2-1.8
1981	88	8.1	6.5-10.0	69	16.2	12.6-20.6	19	2.9	1.7-4.6	38	3.4	2.4-4.7	28	6.0	4.0-8.7	10	1.5	0.7-2.8
1982	75	6.7	5.2-8.4	60	13.4	10.2-17.3	15	2.3	1.3-3.8	26	2.3	1.5-3.4	21	4.7	2.9-7.2	5	0.9	0.3-2.1
1983	78	7.3	5.7-9.1	62	14.1	10.8-18.2	16	2.5	1.4-4.2	31	2.8	1.9-4.0	26	5.7	3.7-8.4	5	0.8	0.2-1.9
1984	67	6.2	4.8-8.0	51	11.9	8.8-15.7	16	2.5	1.4-4.2	21	1.9	1.2-3.0	16	3.6	2.0-5.8	5	0.8	0.3-2.0
1985	81	7.9	6.2-9.8	64	15.5	11.8-19.8	17	2.9	1.7-4.6	29	2.7	1.8-3.8	22	5.2	3.2-7.9	7	1.1	0.4-2.2
1986	64	6.0	4.6-7.7	52	12.3	9.2-16.2	12	1.9	1.0-3.3	25	2.4	1.5-3.5	21	4.8	2.9-7.3	4	0.6	0.2-1.6
1987	67	6.4	5.0-8.2	54	13.0	9.7-16.9	13	2.2	1.1-3.7	38	3.6	2.5-4.9	27	6.4	4.2-9.3	11	1.7	0.8-3.1
1988	80	7.4	5.8-9.2	52	11.8	8.8-15.5	28	4.5	3.0-6.5	37	3.4	2.4-4.7	28	6.7	4.4-9.7	9	1.4	0.6-2.7
1989	94	8.8	7.1-10.8	56	12.6	9.5-16.5	38	6.5	4.6-8.9	25	2.5	1.6-3.7	18	4.4	2.6-6.9	7	1.2	0.5-2.5
1990	78	7.6	6.0-9.5	61	14.8	11.3-19.0	17	2.8	1.6-4.4	18	1.7	1.0-2.7	15	3.7	2.1-6.1	3	0.5	0.1-1.4
1991	73	7.0	5.5-8.8	57	13.8	10.4-17.9	16	2.6	1.5-4.3	21	2.1	1.3-3.1	16	3.9	2.2-6.3	5	0.8	0.2-1.8
1992	81	7.9	6.2-9.8	63	14.9	11.4-19.1	18	3.0	1.7-4.7	30	2.9	2.0-4.2	25	6.2	4.0-9.1	5	0.9	0.3-2.0
1993	71	6.8	5.3-8.5	51	12.1	9.0-15.9	20	3.3	2.0-5.1	36	3.4	2.4-4.7	28	7.0	4.6-10.0	8	1.3	0.5-2.5
1994	69	6.8	5.3-8.6	54	13.0	9.8-17.0	15	2.4	1.4-4.0	40	3.8	2.7-5.1	32	7.3	5.0-10.3	8	1.2	0.5-2.4
1995	65	6.3	4.8-8.0	43	10.1	7.3-13.6	22	3.6	2.3-5.5	29	2.8	1.9-4.0	21	4.8	3.0-7.3	8	1.3	0.6-2.6
1996	76	7.3	5.8-9.2	64	15.0	11.5-19.2	12	1.9	1.0-3.3	46	4.5	3.3-6.0	39	9.3	6.6-12.6	7	1.2	0.5-2.4
1997	56	5.3	4.0-6.8	47	10.9	8.0-14.5	9	1.4	0.6-2.7	28	2.6	1.8-3.8	25	6.2	4.0-9.1	3	0.5	0.1-1.4
1998	69	6.5	5.1-8.2	49	11.4	8.4-15.0	20	3.2	1.9-4.9	30	2.8	1.9-4.0	24	5.6	3.5-8.2	6	1.0	0.3-2.1
1999	55	5.1	3.8-6.6	40	9.0	6.4-12.2	15	2.4	1.3-3.9	29	2.6	1.8-3.8	23	5.2	3.3-7.8	6	1.0	0.4-2.1
2000	73	6.6	5.1-8.2	49	11.0	8.1-14.5	24	3.6	2.3-5.3	32	2.9	2.0-4.1	24	5.2	3.3-7.7	8	1.2	0.5-2.3
2001	69	6.1	4.7-7.7	50	11.1	8.2-14.7	19	2.9	1.7-4.5	24	2.1	1.3-3.1	18	3.8	2.2-6.0	6	0.9	0.3-1.9
2002	63	5.5	4.2-7.1	46	9.9	7.2-13.2	17	2.5	1.5-4.0	30	2.7	1.8-3.8	22	5.0	3.1-7.6	8	1.2	0.5-2.3
2003	65	5.5	4.3-7.0	48	10.0	7.4-13.4	17	2.5	1.4-3.9	24	2.1	1.3-3.1	19	4.5	2.7-7.0	5	0.7	0.2-1.7
2004	50	4.2	3.1-5.5	39	7.8	5.5-10.7	11	1.6	0.8-2.8	30	2.5	1.7-3.6	29	6.0	4.0-8.6	1	0.1	0.0-0.8
2005	50	4.3	3.2-5.6	38	7.7	5.4-10.6	12	1.7	0.9-3.0	24	2.0	1.3-3.0	21	4.4	2.7-6.7	3	0.5	0.1-1.3
2006	42	3.3	2.4-4.5	33	6.1	4.2-8.7	9	1.3	0.6-2.4	26	2.2	1.4-3.2	22	4.8	2.9-7.2	4	0.6	0.2-1.4
2007	61	5.1	3.9-6.5	48	9.7	7.1-12.9	13	1.9	1.0-3.2	18	1.5	0.9-2.3	14	2.6	1.4-4.3	4	0.6	0.2-1.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Larynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	83	6.8	5.4-8.4	61	12.1	9.2-15.7	22	3.0	1.9-4.6	25	2.0	1.3-3.0	18	3.7	2.2-5.9	7	0.9	0.4-1.9
2009	56	4.3	3.2-5.6	43	8.0	5.7-10.8	13	1.7	0.9-3.0	23	1.7	1.1-2.6	18	3.4	2.0-5.4	5	0.6	0.2-1.5
2010	53	4.0	3.0-5.2	42	7.8	5.5-10.6	11	1.4	0.7-2.5	18	1.4	0.8-2.2	16	3.1	1.8-5.1	2	0.3	0.0-1.0
2011	54	4.0	3.0-5.2	39	7.0	4.9-9.7	15	1.9	1.0-3.1	21	1.6	1.0-2.5	18	3.5	2.0-5.5	3	0.4	0.1-1.2
2012	59	4.3	3.3-5.6	45	8.2	5.9-11.0	14	1.7	0.9-2.9	34	2.5	1.7-3.5	25	4.5	2.9-6.8	9	1.1	0.5-2.1
2013	48	3.5	2.6-4.7	41	7.3	5.2-10.0	7	0.9	0.4-1.9	22	1.6	1.0-2.4	14	2.5	1.4-4.3	8	1.0	0.4-1.9
2014	61	4.4	3.3-5.6	46	7.8	5.6-10.4	15	1.9	1.1-3.2	23	1.7	1.0-2.5	20	3.6	2.2-5.6	3	0.3	0.1-1.0
2015	66	4.5	3.4-5.7	43	7.1	5.1-9.6	23	2.7	1.7-4.1	20	1.4	0.8-2.2	14	2.6	1.4-4.4	6	0.7	0.3-1.6
2016	54	3.7	2.8-4.9	44	7.1	5.1-9.6	10	1.3	0.6-2.3	16	1.1	0.6-1.8	14	2.3	1.3-4.0	2	0.2	0.0-0.9
2017	55	3.7	2.8-4.8	41	6.6	4.7-9.0	14	1.7	0.9-2.9	21	1.4	0.8-2.1	16	2.5	1.4-4.1	5	0.6	0.2-1.4
2018	58	3.7	2.8-4.8	42	6.5	4.6-8.8	16	1.8	1.0-3.0	19	1.3	0.8-2.0	13	2.3	1.2-3.9	6	0.7	0.3-1.5
2019	66	4.3	3.3-5.5	45	7.0	5.1-9.5	21	2.4	1.5-3.8	15	1.0	0.5-1.6	14	2.3	1.2-3.9	1	0.1	0.0-0.7
2020	44	2.9	2.1-3.9	36	5.7	4.0-8.0	8	0.9	0.4-1.7	12	0.8	0.4-1.4	8	1.3	0.6-2.7	4	0.4	0.1-1.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Lung and bronchus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	608	49.0	45.1-53.1	443	88.8	80.4-97.8	165	22.8	19.4-26.7	557	44.7	41.0-48.7	420	85.0	76.8-93.9	137	18.6	15.6-22.1
1977	636	51.8	47.8-56.1	451	93.3	84.7-102.5	185	25.3	21.7-29.3	524	42.3	38.7-46.2	394	80.2	72.3-88.8	130	17.7	14.7-21.1
1978	663	56.9	52.6-61.5	486	104.6	95.3-114.5	177	25.9	22.1-30.0	497	42.3	38.6-46.2	369	79.6	71.5-88.3	128	18.1	15.1-21.6
1979	612	52.9	48.7-57.4	432	95.4	86.4-105.0	180	26.4	22.6-30.6	522	46.0	42.1-50.2	371	84.0	75.5-93.2	151	22.0	18.6-25.9
1980	630	55.6	51.3-60.2	433	97.8	88.6-107.6	197	29.1	25.1-33.6	514	45.2	41.3-49.4	366	82.5	74.1-91.6	148	22.2	18.7-26.1
1981	582	51.5	47.3-55.9	395	90.0	81.1-99.4	187	27.1	23.3-31.4	498	44.3	40.4-48.4	338	77.6	69.4-86.5	160	23.3	19.8-27.3
1982	603	54.2	49.9-58.8	392	91.1	82.1-100.7	211	31.0	26.9-35.6	480	42.9	39.1-46.9	339	77.9	69.6-86.8	141	21.0	17.6-24.9
1983	667	60.1	55.6-64.9	401	94.1	85.0-103.9	266	39.2	34.6-44.3	497	44.8	40.9-48.9	310	73.5	65.4-82.3	187	27.2	23.4-31.5
1984	642	58.3	53.8-63.0	429	100.3	90.9-110.4	213	32.3	28.0-37.0	504	45.8	41.8-50.0	345	83.0	74.3-92.4	159	23.6	20.0-27.7
1985	626	56.6	52.2-61.3	396	91.6	82.7-101.2	230	34.1	29.8-38.9	534	48.8	44.7-53.1	339	79.9	71.5-89.0	195	29.7	25.6-34.2
1986	677	61.2	56.7-66.1	412	97.1	87.8-107.0	265	38.7	34.1-43.7	537	48.7	44.6-53.0	325	77.2	68.9-86.2	212	31.2	27.1-35.8
1987	631	57.3	52.9-62.0	392	92.5	83.5-102.3	239	35.6	31.2-40.5	521	47.1	43.1-51.4	305	71.6	63.6-80.2	216	32.1	27.9-36.7
1988	608	55.8	51.4-60.4	380	92.7	83.5-102.6	228	34.0	29.7-38.8	503	46.1	42.1-50.3	335	81.8	73.2-91.1	168	24.6	21.0-28.7
1989	651	60.7	56.1-65.6	387	93.9	84.7-103.9	264	40.8	35.9-46.0	505	46.8	42.8-51.1	311	76.9	68.4-86.0	194	29.2	25.2-33.7
1990	626	58.9	54.4-63.7	373	91.9	82.7-101.8	253	38.6	34.0-43.8	523	49.5	45.3-53.9	312	77.6	69.2-86.8	211	32.3	28.1-37.1
1991	622	59.5	54.9-64.3	362	90.5	81.3-100.4	260	41.2	36.3-46.6	498	47.6	43.5-51.9	301	75.6	67.2-84.7	197	30.3	26.2-34.9
1992	583	55.8	51.3-60.5	318	78.9	70.3-88.1	265	42.5	37.5-48.0	470	44.8	40.9-49.1	290	72.6	64.3-81.5	180	28.4	24.4-32.9
1993	641	61.5	56.8-66.4	358	88.9	79.8-98.7	283	45.0	39.9-50.6	523	50.3	46.1-54.8	293	73.4	65.2-82.4	230	36.9	32.2-42.0
1994	637	61.1	56.5-66.1	370	91.7	82.5-101.6	267	42.0	37.1-47.4	516	49.4	45.2-53.8	307	76.8	68.4-86.0	209	32.7	28.4-37.4
1995	598	57.3	52.8-62.1	323	78.7	70.3-87.9	275	43.1	38.2-48.6	521	50.0	45.8-54.5	296	73.0	64.8-81.9	225	35.1	30.6-40.0
1996	614	59.0	54.4-63.8	356	89.4	80.2-99.2	258	40.7	35.8-46.0	507	48.7	44.5-53.1	300	75.2	66.8-84.2	207	32.2	27.9-36.9
1997	576	55.2	50.8-59.9	295	72.9	64.7-81.8	281	44.3	39.2-49.8	435	41.7	37.9-45.8	231	57.9	50.6-65.9	204	31.9	27.6-36.6
1998	647	61.2	56.6-66.1	366	88.1	79.2-97.7	281	44.0	39.0-49.5	491	46.5	42.5-50.8	279	68.7	60.7-77.3	212	32.7	28.4-37.4
1999	536	49.6	45.5-54.0	304	72.5	64.4-81.2	232	35.7	31.2-40.6	476	44.6	40.6-48.7	271	65.5	57.9-73.9	205	31.4	27.2-36.0
2000	624	56.7	52.3-61.3	333	77.8	69.6-86.7	291	43.4	38.5-48.7	433	39.7	36.0-43.6	233	55.0	48.0-62.5	200	29.7	25.7-34.2
2001	592	53.4	49.2-57.9	334	76.2	68.1-84.9	258	38.6	34.0-43.6	482	43.5	39.7-47.6	275	63.7	56.3-71.8	207	30.7	26.7-35.2
2002	599	53.2	49.0-57.6	305	68.5	60.9-76.8	294	43.4	38.6-48.7	441	39.5	35.9-43.4	235	54.6	47.7-62.1	206	30.1	26.1-34.5
2003	646	56.8	52.5-61.4	319	71.3	63.5-79.7	327	47.3	42.3-52.8	493	43.6	39.8-47.6	256	58.3	51.2-66.0	237	34.4	30.2-39.1
2004	604	52.9	48.8-57.4	329	73.7	65.8-82.3	275	40.2	35.6-45.3	438	38.3	34.8-42.1	236	53.3	46.6-60.7	202	29.1	25.2-33.4
2005	603	52.2	48.0-56.5	329	73.3	65.4-81.8	274	39.3	34.8-44.2	451	39.8	36.2-43.6	236	54.3	47.5-61.8	215	31.1	27.1-35.6
2006	631	54.1	49.9-58.5	352	76.8	68.8-85.4	279	39.7	35.2-44.7	419	36.4	33.0-40.1	225	49.9	43.4-57.0	194	27.8	24.0-32.0
2007	706	59.9	55.5-64.5	361	77.4	69.5-86.1	345	48.9	43.8-54.4	450	38.2	34.7-41.9	255	55.1	48.4-62.5	195	27.3	23.6-31.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Lung and bronchus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	638	53.4	49.3-57.7	336	71.0	63.4-79.2	302	42.4	37.7-47.5	447	37.8	34.4-41.6	235	52.4	45.7-59.7	212	29.3	25.4-33.5
2009	661	53.9	49.8-58.2	330	67.9	60.6-75.9	331	45.2	40.5-50.4	437	36.1	32.7-39.6	231	48.4	42.2-55.2	206	28.2	24.4-32.3
2010	619	50.0	46.1-54.1	337	67.9	60.6-75.7	282	38.2	33.8-42.9	442	35.3	32.0-38.8	237	48.4	42.2-55.1	205	27.5	23.8-31.6
2011	697	54.2	50.2-58.4	367	72.4	64.9-80.4	330	43.0	38.4-47.9	426	33.9	30.7-37.3	243	49.2	43.1-56.0	183	24.4	21.0-28.2
2012	643	49.7	45.9-53.8	321	62.3	55.5-69.6	322	41.9	37.4-46.8	452	34.8	31.6-38.2	248	48.6	42.6-55.2	204	26.4	22.8-30.3
2013	605	45.4	41.8-49.2	298	55.5	49.2-62.4	307	38.8	34.5-43.4	400	29.7	26.8-32.8	221	40.8	35.5-46.7	179	22.1	18.9-25.6
2014	679	48.9	45.2-52.7	323	57.4	51.2-64.2	356	43.3	38.9-48.1	395	29.0	26.1-32.0	195	35.5	30.6-41.0	200	24.5	21.2-28.2
2015	630	44.8	41.3-48.5	341	60.2	53.8-67.1	289	34.7	30.8-39.0	408	29.5	26.7-32.5	219	40.3	35.0-46.1	189	22.7	19.5-26.2
2016	662	45.9	42.4-49.6	319	54.3	48.4-60.8	343	40.3	36.1-44.8	410	28.6	25.9-31.5	218	38.5	33.4-44.1	192	22.0	18.9-25.4
2017	687	46.1	42.7-49.8	353	58.1	52.1-64.7	334	38.1	34.0-42.4	399	26.7	24.1-29.5	210	34.6	30.0-39.8	189	21.3	18.4-24.7
2018	728	48.3	44.9-52.0	365	58.9	52.9-65.4	363	40.8	36.6-45.2	398	26.3	23.7-29.0	224	36.2	31.5-41.4	174	19.3	16.5-22.4
2019	633	41.5	38.3-44.9	309	48.9	43.4-54.8	324	36.4	32.5-40.7	335	22.1	19.8-24.6	179	28.6	24.5-33.3	156	17.5	14.8-20.6
2020	511	32.9	30.1-35.9	280	44.8	39.6-50.5	231	25.0	21.8-28.5	330	21.3	19.0-23.8	167	26.5	22.5-30.9	163	17.7	15.0-20.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Melanoma of the skin**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	62	5.0	3.8-6.4	40	8.2	5.8-11.2	22	2.9	1.8-4.4	27	2.2	1.4-3.2	18	3.7	2.1-5.9	9	1.2	0.6-2.4
1977	64	5.4	4.1-6.9	32	6.5	4.4-9.3	32	4.7	3.2-6.6	22	1.9	1.2-2.9	14	3.1	1.7-5.3	8	1.1	0.5-2.2
1978	45	4.1	3.0-5.5	21	4.5	2.7-6.9	24	3.9	2.5-5.8	15	1.2	0.7-2.0	9	2.0	0.9-3.8	6	0.7	0.3-1.7
1979	41	3.9	2.8-5.4	19	4.3	2.5-6.7	22	3.7	2.3-5.7	15	1.5	0.8-2.4	7	1.5	0.6-3.2	8	1.4	0.6-2.8
1980	45	4.0	2.9-5.4	22	4.8	3.0-7.3	23	3.6	2.3-5.5	23	2.2	1.4-3.3	13	3.2	1.6-5.5	10	1.6	0.7-2.9
1981	42	3.7	2.7-5.1	23	4.8	3.0-7.3	19	3.0	1.8-4.7	19	1.8	1.1-2.9	11	2.6	1.3-4.6	8	1.3	0.6-2.7
1982	49	4.5	3.3-6.0	23	5.3	3.3-8.0	26	4.1	2.7-6.1	13	1.2	0.6-2.0	8	1.8	0.8-3.5	5	0.8	0.3-1.9
1983	51	4.7	3.5-6.2	25	5.8	3.7-8.6	26	3.9	2.6-5.8	20	2.0	1.2-3.1	13	3.5	1.8-5.9	7	1.0	0.4-2.2
1984	68	6.2	4.8-7.8	32	7.4	5.0-10.6	36	5.4	3.7-7.5	24	2.2	1.4-3.3	16	3.8	2.1-6.2	8	1.3	0.5-2.6
1985	46	4.3	3.2-5.8	24	6.0	3.8-9.0	22	3.4	2.1-5.3	20	1.9	1.1-2.9	13	3.1	1.6-5.3	7	1.2	0.5-2.4
1986	51	4.6	3.4-6.0	28	6.9	4.6-10.0	23	3.4	2.2-5.2	25	2.1	1.4-3.2	16	3.8	2.1-6.2	9	1.3	0.6-2.4
1987	29	2.7	1.8-3.9	15	3.4	1.9-5.7	14	2.2	1.2-3.8	24	2.2	1.4-3.3	16	3.9	2.2-6.3	8	1.4	0.6-2.7
1988	37	3.4	2.4-4.7	19	4.6	2.7-7.1	18	2.7	1.6-4.3	11	1.0	0.5-1.8	8	2.2	1.0-4.3	3	0.4	0.1-1.2
1989	29	2.6	1.8-3.8	14	3.3	1.8-5.6	15	2.2	1.2-3.6	15	1.4	0.8-2.3	5	1.4	0.4-3.1	10	1.5	0.7-2.8
1990	41	3.8	2.7-5.2	26	6.2	4.0-9.1	15	2.3	1.3-3.8	16	1.6	0.9-2.7	13	3.3	1.8-5.6	3	0.5	0.1-1.4
1991	42	3.9	2.8-5.3	18	4.2	2.4-6.6	24	3.5	2.3-5.3	17	1.6	0.9-2.5	7	1.9	0.8-3.9	10	1.4	0.7-2.7
1992	36	3.4	2.4-4.7	17	4.3	2.5-6.8	19	2.9	1.7-4.6	13	1.3	0.7-2.2	6	1.5	0.5-3.1	7	1.1	0.4-2.3
1993	53	5.0	3.7-6.5	30	7.0	4.7-10.0	23	3.5	2.2-5.2	18	1.7	1.0-2.7	8	1.9	0.8-3.7	10	1.4	0.7-2.7
1994	37	3.5	2.5-4.8	20	5.0	3.0-7.7	17	2.7	1.6-4.4	12	1.1	0.6-2.0	8	2.2	0.9-4.2	4	0.6	0.2-1.5
1995	42	3.9	2.8-5.3	25	6.0	3.8-8.8	17	2.7	1.6-4.4	22	2.1	1.3-3.2	14	3.2	1.8-5.4	8	1.2	0.5-2.5
1996	33	3.0	2.0-4.2	17	3.8	2.2-6.1	16	2.4	1.4-3.9	16	1.5	0.8-2.4	6	1.6	0.6-3.4	10	1.5	0.7-2.8
1997	47	4.5	3.3-5.9	25	6.6	4.2-9.7	22	3.3	2.1-5.0	9	0.9	0.4-1.6	4	1.0	0.3-2.5	5	0.8	0.3-1.9
1998	45	4.2	3.1-5.6	26	6.3	4.1-9.2	19	2.8	1.7-4.4	15	1.4	0.8-2.3	8	1.8	0.8-3.5	7	1.1	0.4-2.2
1999	40	3.6	2.6-4.9	15	3.7	2.0-6.0	25	3.6	2.3-5.3	11	1.0	0.5-1.8	8	1.7	0.7-3.3	3	0.4	0.1-1.3
2000	46	4.1	3.0-5.4	25	5.8	3.8-8.6	21	3.0	1.8-4.5	20	1.8	1.1-2.8	10	2.6	1.2-4.7	10	1.5	0.7-2.7
2001	43	3.9	2.8-5.2	23	5.2	3.3-7.9	20	2.9	1.8-4.5	17	1.5	0.9-2.4	9	2.3	1.0-4.3	8	1.1	0.5-2.2
2002	39	3.5	2.5-4.8	21	5.0	3.0-7.6	18	2.6	1.5-4.1	15	1.3	0.7-2.1	10	2.0	1.0-3.8	5	0.7	0.2-1.6
2003	51	4.4	3.3-5.8	28	6.4	4.2-9.3	23	3.2	2.0-4.8	12	1.0	0.5-1.8	8	1.6	0.7-3.1	4	0.6	0.2-1.5
2004	74	6.4	5.0-8.0	35	7.8	5.4-10.9	39	5.6	4.0-7.7	11	1.0	0.5-1.7	6	1.5	0.5-3.1	5	0.7	0.2-1.6
2005	58	4.9	3.7-6.3	31	6.9	4.7-9.8	27	3.7	2.5-5.5	10	0.8	0.4-1.5	2	0.5	0.0-1.6	8	1.1	0.5-2.2
2006	65	5.6	4.3-7.1	35	8.0	5.5-11.2	30	4.3	2.9-6.1	18	1.6	1.0-2.6	13	2.9	1.5-5.0	5	0.8	0.2-1.7
2007	47	3.9	2.9-5.2	28	6.2	4.1-9.0	19	2.8	1.7-4.4	16	1.4	0.8-2.2	9	2.2	1.0-4.1	7	1.0	0.4-2.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Melanoma of the skin**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	45	3.7	2.7-4.9	19	3.9	2.3-6.2	26	3.4	2.2-5.1	12	1.0	0.5-1.7	8	1.8	0.8-3.5	4	0.5	0.1-1.4
2009	55	4.4	3.3-5.8	26	5.7	3.6-8.3	29	4.0	2.7-5.7	8	0.6	0.3-1.2	4	0.7	0.2-1.8	4	0.5	0.1-1.3
2010	60	4.8	3.7-6.2	30	6.4	4.2-9.1	30	3.9	2.6-5.6	9	0.7	0.3-1.4	6	1.3	0.5-2.9	3	0.4	0.1-1.1
2011	69	5.4	4.2-6.9	36	7.3	5.1-10.2	33	4.4	3.0-6.2	6	0.5	0.2-1.1	3	0.8	0.2-2.0	3	0.3	0.1-1.0
2012	54	4.1	3.0-5.3	33	6.0	4.1-8.5	21	2.6	1.6-4.1	19	1.4	0.8-2.2	11	2.1	1.0-3.7	8	0.9	0.4-1.8
2013	62	4.5	3.4-5.8	35	6.6	4.5-9.2	27	3.4	2.2-4.9	10	0.8	0.4-1.4	5	1.0	0.3-2.3	5	0.7	0.2-1.6
2014	65	4.9	3.8-6.3	37	7.0	4.9-9.7	28	3.4	2.3-5.0	15	1.1	0.6-1.8	9	1.7	0.8-3.3	6	0.6	0.2-1.5
2015	52	3.7	2.8-4.9	21	3.7	2.2-5.7	31	3.8	2.6-5.4	10	0.7	0.4-1.4	6	1.2	0.4-2.5	4	0.5	0.1-1.3
2016	65	4.7	3.6-6.0	34	6.2	4.2-8.7	31	3.9	2.6-5.6	6	0.4	0.1-0.8	4	0.6	0.2-1.5	2	0.2	0.0-0.8
2017	66	4.7	3.6-6.0	31	5.8	3.9-8.2	35	4.1	2.9-5.8	7	0.4	0.2-0.9	4	0.6	0.2-1.6	3	0.3	0.1-1.0
2018	50	3.5	2.6-4.6	32	5.3	3.6-7.6	18	2.1	1.2-3.3	10	0.7	0.3-1.3	6	1.1	0.4-2.4	4	0.5	0.1-1.2
2019	61	4.0	3.1-5.2	37	6.1	4.3-8.5	24	2.8	1.8-4.2	5	0.3	0.1-0.7	5	0.8	0.3-1.9	0	0.0	0.0-0.5
2020	50	3.3	2.4-4.3	27	4.3	2.8-6.4	23	2.6	1.6-3.9	5	0.4	0.1-0.8	1	0.2	0.0-1.0	4	0.5	0.1-1.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Female breast**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	595	85.1	78.2-92.4	313	44.0	39.2-49.3
1977	643	94.2	86.9-102.0	253	36.4	32.0-41.3
1978	639	96.3	88.7-104.2	259	37.9	33.3-43.0
1979	572	87.5	80.3-95.2	217	32.4	28.1-37.1
1980	616	96.2	88.6-104.3	269	40.7	35.9-46.0
1981	569	89.9	82.5-97.8	244	37.4	32.8-42.6
1982	607	96.5	88.8-104.7	229	35.5	31.0-40.6
1983	619	96.3	88.7-104.4	196	29.7	25.6-34.2
1984	585	91.4	84.0-99.3	247	37.3	32.7-42.4
1985	582	92.5	85.0-100.5	229	35.4	30.9-40.5
1986	603	97.3	89.5-105.6	209	32.1	27.8-36.8
1987	617	95.7	88.1-103.7	192	29.2	25.2-33.7
1988	680	106.4	98.4-114.8	239	36.7	32.1-41.8
1989	582	92.7	85.2-100.7	221	34.0	29.6-38.8
1990	659	107.1	99.0-115.7	243	38.3	33.5-43.4
1991	742	120.2	111.6-129.2	227	36.7	32.0-41.9
1992	664	107.8	99.6-116.4	218	34.9	30.4-39.9
1993	610	98.7	90.9-106.9	207	32.7	28.3-37.5
1994	708	114.3	106.0-123.1	220	34.4	30.0-39.3
1995	677	109.1	101.0-117.7	231	36.4	31.9-41.5
1996	694	111.2	103.0-119.8	237	37.4	32.8-42.5
1997	719	113.8	105.6-122.4	215	33.9	29.5-38.7
1998	715	112.6	104.5-121.2	177	27.4	23.4-31.7
1999	718	111.2	103.2-119.7	182	27.6	23.8-32.0
2000	738	111.9	103.9-120.3	177	26.7	22.9-30.9
2001	746	111.7	103.8-120.0	191	28.3	24.4-32.6
2002	725	106.5	98.8-114.5	175	25.4	21.8-29.5
2003	734	106.6	99.0-114.6	190	27.2	23.5-31.4
2004	734	106.1	98.6-114.1	190	27.1	23.4-31.2
2005	724	104.0	96.5-111.9	168	23.6	20.1-27.4
2006	763	108.1	100.5-116.1	170	23.7	20.3-27.6
2007	739	103.8	96.4-111.6	157	21.4	18.1-25.0
2008	804	111.3	103.7-119.3	170	22.9	19.5-26.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Female breast**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	808	110.5	103.0-118.5	182	25.1	21.5-29.0
2010	787	106.5	99.2-114.3	180	23.6	20.3-27.4
2011	853	112.5	105.0-120.4	161	21.2	18.0-24.7
2012	860	112.2	104.8-120.1	183	23.4	20.1-27.1
2013	827	106.1	98.9-113.7	155	19.6	16.6-23.0
2014	918	115.6	108.2-123.4	190	23.2	20.0-26.8
2015	952	118.3	110.8-126.1	162	19.6	16.7-23.0
2016	863	105.1	98.1-112.5	172	20.6	17.6-24.0
2017	970	117.7	110.3-125.5	185	21.8	18.7-25.2
2018	984	118.2	110.8-126.0	162	18.7	15.8-21.9
2019	975	117.4	110.0-125.2	170	19.3	16.4-22.5
2020	833	99.8	93.0-107.0	143	16.4	13.7-19.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Cervix uteri**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	118	18.6	15.3-22.4	58	8.2	6.2-10.6
1977	103	16.1	13.0-19.6	43	6.9	4.9-9.3
1978	107	18.5	15.1-22.5	51	8.0	5.9-10.6
1979	116	19.7	16.2-23.8	38	6.5	4.5-8.9
1980	101	17.6	14.3-21.5	36	5.7	3.9-7.9
1981	101	16.6	13.4-20.2	43	7.1	5.1-9.6
1982	97	16.6	13.4-20.4	47	7.8	5.7-10.4
1983	111	18.4	15.0-22.3	34	5.2	3.6-7.3
1984	125	21.6	17.9-25.8	25	4.1	2.6-6.0
1985	109	18.7	15.3-22.6	32	5.1	3.4-7.2
1986	96	16.1	13.0-19.7	35	6.0	4.2-8.4
1987	105	16.9	13.8-20.6	25	4.0	2.5-5.9
1988	96	15.7	12.7-19.2	31	5.2	3.5-7.4
1989	89	14.4	11.5-17.8	23	3.6	2.2-5.4
1990	117	19.2	15.8-23.0	32	5.3	3.6-7.5
1991	113	18.9	15.6-22.8	35	5.4	3.8-7.6
1992	103	16.7	13.6-20.2	21	3.4	2.1-5.2
1993	89	14.4	11.6-17.8	34	5.7	3.9-8.0
1994	105	17.3	14.1-20.9	33	5.3	3.6-7.5
1995	110	17.4	14.3-21.0	31	4.7	3.2-6.7
1996	130	20.6	17.2-24.5	35	5.7	3.9-7.9
1997	107	16.6	13.6-20.0	44	6.9	5.0-9.2
1998	119	18.5	15.3-22.1	28	4.3	2.9-6.3
1999	103	15.7	12.8-19.0	52	8.0	5.9-10.5
2000	109	16.5	13.6-19.9	34	5.1	3.5-7.1
2001	88	13.2	10.6-16.3	35	5.2	3.6-7.3
2002	106	15.6	12.8-18.9	23	3.5	2.2-5.2
2003	90	13.1	10.5-16.1	32	4.7	3.2-6.6
2004	97	13.8	11.1-16.8	28	4.1	2.7-6.0
2005	81	11.6	9.2-14.5	36	5.1	3.5-7.0
2006	94	13.4	10.8-16.4	28	3.9	2.6-5.7
2007	93	13.4	10.8-16.4	31	4.4	3.0-6.2
2008	81	11.4	9.0-14.1	35	4.7	3.3-6.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Cervix uteri**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	72	9.8	7.6-12.3	22	3.0	1.9-4.6
2010	67	9.1	7.1-11.6	25	3.5	2.2-5.2
2011	69	9.4	7.3-11.9	23	3.1	2.0-4.7
2012	78	10.6	8.3-13.2	20	2.7	1.6-4.1
2013	67	8.5	6.6-10.9	28	3.6	2.4-5.2
2014	77	9.9	7.8-12.4	38	4.6	3.2-6.3
2015	81	10.6	8.4-13.2	17	2.1	1.2-3.4
2016	71	9.3	7.2-11.7	25	3.1	2.0-4.5
2017	77	9.7	7.6-12.2	20	2.6	1.6-4.0
2018	72	9.3	7.3-11.8	21	2.5	1.5-3.9
2019	80	10.6	8.3-13.3	23	2.9	1.8-4.4
2020	60	7.3	5.5-9.5	23	2.6	1.7-4.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Bronx

Corpus uteris and uterus, not otherwise specified

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	139	19.1	16.0-22.6	68	9.0	6.9-11.4
1977	135	19.2	16.0-22.9	42	5.8	4.1-7.9
1978	133	19.0	15.8-22.6	37	5.5	3.8-7.6
1979	144	20.7	17.4-24.4	36	4.9	3.4-6.8
1980	124	18.1	15.0-21.6	45	7.0	5.0-9.4
1981	133	20.2	16.8-24.1	39	5.6	4.0-7.8
1982	136	19.9	16.6-23.6	36	5.2	3.6-7.2
1983	131	19.5	16.2-23.2	29	4.1	2.7-5.9
1984	99	14.7	11.9-18.0	38	5.7	4.0-7.9
1985	111	16.3	13.4-19.7	31	4.6	3.1-6.5
1986	101	15.1	12.3-18.5	27	3.8	2.5-5.6
1987	131	19.8	16.5-23.6	25	3.8	2.4-5.6
1988	126	19.3	16.0-23.0	37	5.4	3.8-7.5
1989	96	14.9	12.0-18.3	26	4.0	2.6-5.9
1990	115	18.0	14.8-21.7	35	5.2	3.6-7.3
1991	140	22.1	18.6-26.1	29	4.5	3.0-6.5
1992	129	20.2	16.8-24.0	40	6.2	4.4-8.4
1993	117	18.6	15.4-22.3	42	6.4	4.6-8.6
1994	120	19.0	15.7-22.7	37	5.8	4.1-8.0
1995	131	20.9	17.4-24.8	39	5.9	4.2-8.1
1996	123	19.8	16.4-23.6	30	4.7	3.2-6.7
1997	140	22.0	18.5-26.0	49	7.5	5.6-10.0
1998	144	22.8	19.2-26.8	39	6.1	4.3-8.3
1999	170	25.7	22.0-29.9	32	4.8	3.3-6.8
2000	162	24.6	21.0-28.7	40	6.0	4.3-8.1
2001	166	25.0	21.3-29.1	37	5.5	3.9-7.6
2002	172	25.6	21.9-29.7	30	4.5	3.0-6.4
2003	171	24.7	21.2-28.7	40	5.8	4.1-7.9
2004	193	27.9	24.1-32.2	36	5.2	3.7-7.2
2005	187	26.7	23.0-30.9	42	5.9	4.2-8.0
2006	206	29.2	25.3-33.5	45	6.4	4.6-8.5
2007	187	26.0	22.4-30.0	48	6.8	5.0-9.1
2008	206	28.8	25.0-33.0	58	8.1	6.2-10.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Bronx

Corpus uteris and uterus, not otherwise specified

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	243	33.2	29.1-37.7	51	7.0	5.2-9.2
2010	233	30.8	26.9-35.0	54	7.3	5.5-9.6
2011	231	29.6	25.9-33.7	59	7.7	5.9-10.0
2012	241	30.7	26.9-34.9	62	8.2	6.3-10.5
2013	261	32.6	28.7-36.8	68	8.7	6.7-11.0
2014	257	31.4	27.6-35.5	54	6.6	4.9-8.6
2015	257	30.7	27.1-34.8	57	6.7	5.0-8.7
2016	256	30.4	26.7-34.4	88	10.3	8.2-12.7
2017	254	29.5	25.9-33.4	56	6.5	4.9-8.4
2018	289	33.2	29.4-37.3	79	9.0	7.1-11.2
2019	280	32.2	28.5-36.2	84	9.4	7.5-11.7
2020	250	27.8	24.4-31.5	77	8.3	6.5-10.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Ovary**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	103	13.8	11.2-16.8	73	10.1	7.8-12.7
1977	95	13.7	11.0-16.8	66	9.1	7.0-11.7
1978	92	13.6	10.9-16.8	66	9.3	7.1-11.8
1979	98	14.5	11.7-17.8	56	8.0	6.0-10.4
1980	100	15.1	12.2-18.4	64	9.5	7.3-12.2
1981	92	14.0	11.2-17.3	61	8.9	6.8-11.5
1982	88	13.9	11.0-17.2	47	6.7	4.9-9.0
1983	96	15.1	12.2-18.5	55	8.4	6.3-11.0
1984	81	12.7	10.0-15.8	69	10.9	8.4-13.8
1985	87	13.6	10.8-16.8	55	8.3	6.2-10.9
1986	77	11.7	9.2-14.7	47	7.1	5.2-9.5
1987	66	10.6	8.2-13.6	46	6.8	5.0-9.1
1988	71	11.0	8.6-14.0	51	7.6	5.6-10.0
1989	63	9.6	7.3-12.3	53	8.1	6.1-10.7
1990	86	13.3	10.6-16.4	40	6.0	4.3-8.2
1991	81	12.7	10.1-15.9	52	8.0	5.9-10.5
1992	90	14.2	11.4-17.5	62	9.6	7.3-12.3
1993	72	11.3	8.8-14.2	58	9.2	7.0-11.9
1994	100	15.8	12.8-19.2	51	7.8	5.8-10.3
1995	79	12.7	10.1-15.9	45	7.1	5.2-9.5
1996	85	13.6	10.8-16.8	46	7.3	5.3-9.7
1997	97	15.2	12.3-18.6	51	7.9	5.9-10.5
1998	64	10.1	7.8-12.9	46	7.3	5.4-9.8
1999	76	11.8	9.3-14.7	50	7.7	5.7-10.1
2000	75	11.3	8.9-14.1	50	7.6	5.6-10.0
2001	92	13.7	11.0-16.8	56	8.4	6.4-10.9
2002	94	13.5	10.9-16.6	59	8.5	6.4-11.0
2003	81	11.6	9.2-14.4	45	6.5	4.8-8.7
2004	83	11.8	9.4-14.6	46	6.5	4.8-8.7
2005	82	11.8	9.4-14.6	47	6.8	5.0-9.1
2006	82	11.4	9.0-14.1	51	7.1	5.2-9.3
2007	86	11.9	9.5-14.7	57	7.9	6.0-10.2
2008	78	10.7	8.4-13.4	40	5.5	3.9-7.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Ovary**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	102	13.9	11.3-16.9	48	6.4	4.7-8.5
2010	66	8.9	6.9-11.4	47	6.0	4.4-8.1
2011	83	10.8	8.6-13.5	39	5.1	3.6-7.0
2012	86	11.0	8.8-13.6	27	3.5	2.3-5.1
2013	81	10.1	8.0-12.6	48	6.0	4.4-7.9
2014	85	10.7	8.5-13.2	53	6.5	4.8-8.5
2015	80	9.8	7.8-12.3	37	4.6	3.2-6.4
2016	91	10.9	8.8-13.5	58	6.9	5.2-9.0
2017	91	11.0	8.8-13.5	40	4.7	3.3-6.4
2018	84	10.0	7.9-12.4	40	4.5	3.2-6.2
2019	88	10.7	8.5-13.2	48	5.4	4.0-7.2
2020	91	10.7	8.6-13.3	45	5.0	3.6-6.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Prostate**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	319	71.2	63.3-79.9	155	37.1	31.3-43.7
1977	340	78.4	70.1-87.5	111	25.8	21.1-31.2
1978	330	75.0	66.9-83.8	130	31.7	26.3-37.8
1979	357	84.9	76.2-94.4	118	30.0	24.7-36.0
1980	381	89.3	80.4-99.0	133	33.6	28.0-40.0
1981	304	73.2	65.1-82.1	118	30.3	25.0-36.4
1982	335	80.5	71.9-89.7	116	29.3	24.1-35.2
1983	326	78.2	69.8-87.3	123	31.6	26.1-37.7
1984	336	83.1	74.3-92.6	138	35.8	30.0-42.4
1985	394	96.2	86.8-106.3	111	28.4	23.3-34.2
1986	360	89.6	80.4-99.4	123	31.4	26.0-37.5
1987	361	89.7	80.5-99.5	125	32.7	27.1-38.9
1988	389	98.1	88.5-108.5	156	41.7	35.3-48.8
1989	343	87.9	78.7-97.8	141	37.6	31.6-44.3
1990	428	109.6	99.4-120.6	143	39.8	33.5-46.9
1991	516	132.7	121.3-144.8	133	36.9	30.8-43.7
1992	556	142.8	131.0-155.3	183	51.2	43.9-59.1
1993	595	151.8	139.7-164.6	157	44.7	37.9-52.3
1994	631	158.9	146.6-171.9	169	48.3	41.2-56.1
1995	573	143.9	132.2-156.3	162	46.7	39.7-54.5
1996	603	150.0	138.1-162.5	127	36.3	30.2-43.2
1997	646	160.1	147.8-173.0	133	36.9	30.8-43.7
1998	676	164.3	152.0-177.3	124	34.4	28.5-41.0
1999	823	199.0	185.5-213.2	134	36.8	30.7-43.5
2000	963	226.5	212.2-241.5	139	37.5	31.5-44.3
2001	894	205.3	191.9-219.4	146	39.9	33.6-46.9
2002	894	203.4	190.0-217.4	146	38.1	32.1-44.8
2003	831	185.5	172.8-198.8	147	38.6	32.6-45.4
2004	871	189.3	176.7-202.5	128	33.2	27.6-39.5
2005	864	184.3	171.9-197.3	120	31.0	25.6-37.1
2006	910	193.0	180.3-206.2	121	31.4	26.0-37.5
2007	1,027	209.3	196.3-222.8	134	34.5	28.8-40.9
2008	890	176.4	164.7-188.7	134	33.8	28.2-40.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Prostate**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	952	184.2	172.4-196.6	125	30.8	25.5-36.7
2010	937	175.8	164.4-187.7	125	30.3	25.2-36.2
2011	1,043	188.6	177.0-200.7	126	29.2	24.2-34.8
2012	961	168.5	157.7-179.8	106	24.4	20.0-29.6
2013	910	155.2	145.0-165.9	107	23.4	19.1-28.2
2014	864	143.0	133.3-153.1	131	27.9	23.2-33.1
2015	945	152.1	142.3-162.4	126	25.7	21.3-30.6
2016	866	134.6	125.6-144.2	130	25.7	21.4-30.5
2017	986	149.3	140.0-159.2	124	22.9	19.0-27.3
2018	998	150.0	140.6-159.8	118	21.6	17.9-25.9
2019	1,015	151.4	142.1-161.3	133	23.1	19.3-27.4
2020	820	119.2	111.0-127.9	116	20.1	16.5-24.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Testis**

Year of diagnosis	Incidence		
	Cases	Rate per 100,000	95% CI
1976	7	1.0	0.4-2.1
1977	10	1.7	0.8-3.3
1978	8	1.5	0.6-3.0
1979	9	1.8	0.8-3.5
1980	14	3.0	1.6-5.0
1981	9	1.6	0.7-3.2
1982	11	2.0	1.0-3.8
1983	9	1.6	0.7-3.2
1984	8	1.4	0.6-2.9
1985	12	2.2	1.1-4.0
1986	14	2.4	1.3-4.2
1987	11	1.6	0.8-3.1
1988	8	1.4	0.6-2.8
1989	12	1.9	1.0-3.5
1990	14	2.3	1.2-4.0
1991	16	2.7	1.5-4.5
1992	14	2.4	1.3-4.1
1993	20	3.1	1.9-5.0
1994	17	2.8	1.6-4.5
1995	8	1.3	0.5-2.6
1996	17	2.6	1.5-4.2
1997	10	1.6	0.8-3.1
1998	10	1.8	0.8-3.4
1999	18	2.9	1.7-4.7
2000	19	3.1	1.9-4.9
2001	15	2.3	1.3-3.9
2002	18	2.7	1.6-4.3
2003	16	2.5	1.4-4.1
2004	16	2.5	1.4-4.1
2005	16	2.6	1.5-4.3
2006	27	4.5	2.9-6.5
2007	21	3.4	2.1-5.2
2008	30	4.6	3.1-6.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

Mortality data not provided due to small number of deaths per year.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Testis**

Year of diagnosis	Incidence		
	Cases	Rate per 100,000	95% CI
2009	23	3.8	2.4-5.8
2010	26	4.0	2.6-6.0
2011	26	3.9	2.5-5.8
2012	21	3.2	1.9-5.0
2013	24	3.5	2.2-5.3
2014	30	4.5	3.0-6.5
2015	23	3.5	2.2-5.3
2016	22	3.3	2.0-5.0
2017	24	3.4	2.1-5.1
2018	22	3.2	2.0-4.9
2019	23	3.4	2.2-5.2
2020	16	2.5	1.4-4.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

Mortality data not provided due to small number of deaths per year.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Urinary bladder (incl. in situ)**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	166	13.2	11.2-15.4	108	22.0	17.9-26.7	58	7.7	5.8-10.0	77	6.7	5.2-8.4	45	10.6	7.6-14.2	32	4.5	3.0-6.3
1977	219	18.4	16.0-21.1	154	33.1	27.9-39.0	65	9.1	7.0-11.7	68	5.8	4.5-7.4	45	10.3	7.4-13.9	23	3.2	2.0-4.8
1978	232	19.7	17.2-22.5	161	36.4	30.8-42.6	71	9.6	7.5-12.1	59	5.2	4.0-6.8	42	10.4	7.4-14.1	17	2.4	1.4-3.9
1979	221	19.5	17.0-22.3	148	34.8	29.3-41.1	73	10.4	8.1-13.2	63	5.6	4.3-7.2	44	10.9	7.8-14.6	19	2.6	1.6-4.1
1980	200	17.3	15.0-19.9	145	34.1	28.7-40.3	55	7.5	5.7-9.8	71	6.3	4.9-7.9	45	11.1	8.0-14.9	26	3.6	2.4-5.4
1981	183	16.2	13.9-18.7	126	30.3	25.1-36.2	57	7.9	6.0-10.3	57	5.1	3.9-6.6	42	10.3	7.4-14.0	15	2.1	1.2-3.5
1982	197	17.4	15.1-20.1	143	34.0	28.5-40.1	54	7.7	5.7-10.0	50	4.4	3.3-5.8	34	8.1	5.5-11.3	16	2.2	1.3-3.7
1983	209	18.6	16.1-21.3	139	34.0	28.5-40.3	70	9.8	7.6-12.4	48	4.3	3.2-5.7	29	7.5	5.0-10.7	19	2.7	1.6-4.2
1984	166	14.8	12.6-17.3	115	27.5	22.6-33.1	51	7.3	5.4-9.6	50	4.5	3.4-6.0	33	8.2	5.6-11.6	17	2.5	1.4-4.0
1985	178	16.2	13.9-18.8	114	27.8	22.9-33.5	64	9.2	7.1-11.8	47	4.3	3.1-5.7	29	7.6	5.1-11.0	18	2.5	1.5-4.1
1986	175	15.9	13.6-18.4	122	29.8	24.7-35.6	53	7.8	5.8-10.2	48	4.3	3.2-5.7	23	6.3	3.9-9.4	25	3.4	2.2-5.1
1987	182	16.3	14.0-18.8	122	30.4	25.1-36.3	60	8.4	6.4-10.8	44	4.0	2.9-5.4	29	7.5	5.0-10.7	15	2.2	1.2-3.7
1988	186	16.8	14.5-19.4	119	29.5	24.3-35.3	67	9.5	7.4-12.1	51	4.6	3.4-6.0	34	9.0	6.2-12.5	17	2.4	1.4-3.8
1989	163	15.0	12.7-17.4	117	29.9	24.6-35.8	46	6.5	4.7-8.7	49	4.4	3.3-5.8	31	8.3	5.6-11.8	18	2.5	1.5-4.0
1990	167	15.7	13.4-18.3	113	29.4	24.2-35.4	54	8.1	6.1-10.6	55	5.2	3.9-6.8	33	8.8	6.1-12.4	22	3.2	2.0-4.9
1991	181	17.3	14.9-20.0	126	33.1	27.5-39.5	55	8.6	6.5-11.2	52	4.9	3.7-6.5	35	9.6	6.7-13.4	17	2.5	1.5-4.0
1992	164	15.6	13.3-18.2	105	28.1	23.0-34.1	59	8.9	6.8-11.5	55	5.2	3.9-6.8	34	9.5	6.5-13.2	21	3.1	1.9-4.7
1993	147	13.9	11.8-16.4	90	23.4	18.8-28.8	57	8.6	6.5-11.2	54	5.2	3.9-6.8	36	10.1	7.1-14.0	18	2.7	1.6-4.3
1994	150	14.5	12.2-17.0	99	26.7	21.6-32.5	51	7.7	5.7-10.2	54	5.2	3.9-6.8	30	8.7	5.8-12.3	24	3.6	2.3-5.4
1995	136	13.0	10.9-15.4	108	28.7	23.5-34.7	28	4.2	2.8-6.0	44	4.2	3.1-5.7	30	8.2	5.5-11.6	14	2.1	1.1-3.5
1996	147	14.1	11.9-16.5	99	25.6	20.8-31.2	48	7.4	5.4-9.8	44	4.2	3.1-5.7	19	5.2	3.1-8.0	25	3.7	2.4-5.5
1997	155	14.9	12.6-17.4	105	27.1	22.1-32.8	50	7.6	5.7-10.1	46	4.4	3.2-5.9	27	7.3	4.8-10.5	19	2.8	1.7-4.4
1998	161	15.3	13.1-17.9	101	26.0	21.2-31.7	60	8.7	6.6-11.2	49	4.7	3.5-6.2	25	6.7	4.3-9.8	24	3.5	2.3-5.3
1999	143	13.3	11.2-15.7	91	22.7	18.2-27.9	52	7.7	5.7-10.1	34	3.2	2.2-4.5	23	6.3	4.0-9.4	11	1.6	0.8-2.9
2000	190	17.5	15.1-20.1	115	28.6	23.5-34.3	75	11.0	8.6-13.8	55	4.9	3.7-6.4	27	6.8	4.4-9.9	28	4.0	2.6-5.8
2001	160	14.7	12.5-17.1	107	26.4	21.6-31.9	53	7.7	5.8-10.1	44	4.0	2.9-5.4	26	6.8	4.4-9.9	18	2.5	1.5-4.0
2002	169	15.3	13.1-17.8	103	25.9	21.1-31.4	66	9.4	7.3-12.0	36	3.2	2.3-4.5	19	5.3	3.2-8.1	17	2.4	1.4-3.8
2003	172	15.3	13.1-17.8	116	27.9	23.0-33.5	56	7.9	5.9-10.2	44	3.9	2.8-5.3	26	6.8	4.4-9.9	18	2.4	1.4-3.9
2004	165	14.6	12.5-17.1	106	25.1	20.5-30.5	59	8.2	6.3-10.6	49	4.4	3.2-5.8	25	6.8	4.3-9.9	24	3.3	2.1-4.9
2005	194	17.0	14.7-19.6	133	30.2	25.2-35.9	61	8.5	6.5-11.0	46	4.2	3.1-5.6	28	7.2	4.8-10.4	18	2.5	1.5-4.0
2006	160	14.2	12.0-16.5	124	29.6	24.5-35.4	36	5.0	3.5-7.0	47	4.2	3.1-5.6	29	7.0	4.7-10.1	18	2.4	1.4-3.8
2007	155	13.5	11.4-15.8	109	25.4	20.7-30.7	46	6.3	4.6-8.4	43	3.8	2.7-5.1	23	5.7	3.6-8.5	20	2.7	1.6-4.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Urinary bladder (incl. in situ)**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	137	11.7	9.8-13.9	104	23.8	19.3-28.9	33	4.6	3.2-6.5	36	3.1	2.2-4.3	24	6.0	3.8-8.9	12	1.6	0.8-2.8
2009	168	13.8	11.8-16.1	113	24.5	20.1-29.6	55	7.3	5.5-9.5	40	3.3	2.4-4.6	23	5.7	3.6-8.5	17	2.2	1.3-3.5
2010	170	13.9	11.9-16.2	125	26.7	22.1-31.9	45	6.1	4.5-8.2	41	3.3	2.4-4.5	25	5.9	3.8-8.7	16	2.0	1.1-3.3
2011	175	14.1	12.1-16.4	122	26.1	21.5-31.2	53	6.8	5.0-8.9	41	3.4	2.4-4.6	26	5.8	3.7-8.5	15	2.0	1.1-3.3
2012	174	13.4	11.5-15.6	127	25.9	21.5-30.9	47	5.8	4.3-7.8	42	3.2	2.3-4.3	24	5.1	3.2-7.5	18	2.2	1.3-3.6
2013	189	14.3	12.3-16.5	134	25.9	21.6-30.7	55	6.9	5.2-9.0	42	3.4	2.4-4.6	32	7.2	4.9-10.2	10	1.2	0.6-2.2
2014	179	13.2	11.3-15.3	119	22.8	18.8-27.4	60	7.3	5.6-9.5	40	3.0	2.1-4.1	32	6.7	4.6-9.5	8	0.8	0.3-1.7
2015	197	14.2	12.3-16.4	132	24.0	19.9-28.5	65	7.9	6.1-10.1	46	3.3	2.4-4.5	23	4.8	3.0-7.1	23	2.7	1.7-4.1
2016	187	13.2	11.3-15.2	137	24.6	20.6-29.2	50	5.7	4.2-7.5	49	3.5	2.6-4.6	30	5.8	3.9-8.3	19	2.0	1.2-3.2
2017	199	13.9	12.1-16.1	137	23.8	19.9-28.2	62	7.4	5.6-9.5	44	3.2	2.3-4.3	21	4.0	2.5-6.1	23	2.7	1.7-4.0
2018	202	13.3	11.5-15.3	134	22.2	18.5-26.3	68	7.6	5.9-9.7	48	3.2	2.4-4.3	33	5.6	3.9-8.0	15	1.6	0.9-2.6
2019	214	14.2	12.3-16.3	155	25.9	21.9-30.4	59	6.7	5.1-8.7	43	2.9	2.1-3.9	35	6.1	4.2-8.5	8	0.9	0.4-1.8
2020	165	10.7	9.1-12.5	126	20.2	16.8-24.1	39	4.4	3.1-6.0	37	2.4	1.7-3.3	21	3.7	2.2-5.6	16	1.7	1.0-2.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Kidney and renal pelvis**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	83	6.9	5.5-8.6	46	9.8	7.1-13.1	37	5.0	3.5-7.0	59	4.6	3.5-6.0	35	7.0	4.8-9.9	24	3.1	2.0-4.6
1977	86	7.0	5.6-8.7	55	10.8	8.0-14.1	31	4.4	3.0-6.3	49	4.2	3.1-5.6	32	7.0	4.7-10.0	17	2.3	1.3-3.7
1978	96	8.2	6.6-10.0	53	11.7	8.7-15.3	43	5.9	4.2-8.0	38	3.2	2.2-4.4	23	4.8	3.0-7.3	15	2.2	1.2-3.7
1979	65	5.6	4.3-7.2	44	9.8	7.1-13.2	21	2.8	1.7-4.3	46	4.0	2.9-5.4	27	6.0	4.0-8.8	19	2.6	1.6-4.1
1980	72	6.3	4.9-8.0	42	9.0	6.4-12.2	30	4.7	3.2-6.8	33	2.9	2.0-4.2	20	4.7	2.8-7.3	13	1.9	1.0-3.2
1981	62	5.6	4.3-7.2	36	8.3	5.8-11.5	26	3.7	2.4-5.5	36	3.3	2.3-4.5	21	4.8	2.9-7.3	15	2.2	1.2-3.6
1982	60	5.4	4.1-7.0	32	7.6	5.1-10.7	28	4.0	2.6-5.8	27	2.4	1.6-3.5	19	4.7	2.8-7.3	8	1.1	0.5-2.1
1983	73	6.8	5.3-8.5	33	7.8	5.3-11.0	40	6.3	4.5-8.7	32	2.9	2.0-4.1	22	5.2	3.2-7.9	10	1.4	0.7-2.7
1984	76	6.9	5.4-8.6	47	10.7	7.8-14.2	29	4.4	2.9-6.3	32	3.0	2.0-4.2	16	4.0	2.2-6.4	16	2.3	1.3-3.8
1985	101	8.9	7.3-10.9	53	11.8	8.8-15.5	48	7.1	5.2-9.4	35	3.2	2.2-4.4	15	3.4	1.9-5.6	20	3.1	1.9-4.8
1986	82	7.4	5.8-9.2	52	11.8	8.7-15.5	30	4.6	3.0-6.6	40	3.6	2.5-4.9	24	5.6	3.6-8.3	16	2.4	1.3-3.9
1987	92	8.4	6.8-10.4	54	12.3	9.2-16.1	38	5.9	4.2-8.2	43	4.0	2.9-5.5	25	6.0	3.8-8.8	18	2.7	1.6-4.4
1988	78	7.3	5.7-9.1	44	10.3	7.5-13.9	34	5.2	3.5-7.2	37	3.4	2.4-4.7	19	4.8	2.9-7.4	18	2.5	1.5-4.1
1989	73	6.6	5.2-8.3	49	11.1	8.2-14.7	24	3.6	2.3-5.4	33	3.0	2.1-4.2	20	4.9	3.0-7.5	13	1.8	1.0-3.2
1990	86	8.0	6.4-9.9	45	10.7	7.8-14.3	41	6.2	4.4-8.4	33	3.1	2.1-4.3	21	5.1	3.1-7.8	12	1.8	0.9-3.1
1991	83	7.8	6.2-9.7	45	10.5	7.6-14.1	38	6.0	4.2-8.2	35	3.3	2.3-4.6	19	4.8	2.9-7.5	16	2.5	1.4-4.0
1992	94	8.9	7.2-10.9	62	14.7	11.2-18.9	32	5.0	3.4-7.0	35	3.4	2.3-4.7	18	4.5	2.6-7.1	17	2.6	1.5-4.2
1993	81	7.6	6.1-9.5	58	13.6	10.3-17.6	23	3.6	2.3-5.4	36	3.4	2.4-4.7	23	5.9	3.7-8.8	13	2.0	1.0-3.4
1994	93	8.9	7.1-10.9	49	11.8	8.7-15.6	44	6.8	4.9-9.2	40	3.8	2.7-5.2	23	5.6	3.5-8.4	17	2.6	1.5-4.1
1995	98	9.3	7.6-11.3	57	13.9	10.5-18.0	41	6.5	4.6-8.8	35	3.3	2.3-4.5	20	4.9	3.0-7.5	15	2.3	1.3-3.8
1996	93	8.8	7.1-10.8	54	13.3	10.0-17.4	39	6.2	4.4-8.4	30	2.9	1.9-4.1	19	4.7	2.8-7.4	11	1.7	0.9-3.1
1997	112	10.6	8.7-12.7	61	14.7	11.2-18.9	51	8.0	6.0-10.5	27	2.5	1.6-3.6	15	3.6	2.0-6.0	12	1.8	0.9-3.1
1998	103	9.5	7.8-11.6	61	13.9	10.6-17.9	42	6.5	4.7-8.8	37	3.5	2.4-4.8	16	3.9	2.2-6.3	21	3.2	2.0-4.9
1999	110	10.2	8.4-12.3	60	13.6	10.4-17.6	50	7.7	5.7-10.2	24	2.2	1.4-3.3	16	3.9	2.2-6.2	8	1.2	0.5-2.3
2000	120	10.7	8.9-12.8	66	14.8	11.4-18.8	54	8.2	6.1-10.6	32	2.8	1.9-4.0	22	5.0	3.1-7.6	10	1.4	0.7-2.6
2001	119	10.3	8.5-12.3	73	15.3	12.0-19.3	46	6.7	4.9-9.0	38	3.4	2.4-4.6	24	5.4	3.4-8.0	14	2.0	1.1-3.4
2002	143	12.5	10.5-14.7	79	16.9	13.3-21.1	64	9.4	7.3-12.0	33	2.9	2.0-4.1	22	4.9	3.0-7.4	11	1.6	0.8-2.9
2003	163	13.9	11.9-16.3	92	19.0	15.3-23.4	71	10.3	8.0-13.0	40	3.6	2.5-4.8	24	5.5	3.5-8.2	16	2.3	1.3-3.7
2004	142	11.9	10.0-14.0	72	14.5	11.3-18.3	70	9.9	7.7-12.6	24	2.1	1.4-3.2	14	3.3	1.8-5.5	10	1.3	0.6-2.5
2005	161	13.6	11.6-15.9	83	16.8	13.3-20.9	78	11.3	8.9-14.0	34	2.8	1.9-3.9	20	4.0	2.4-6.3	14	1.8	1.0-3.1
2006	173	14.4	12.4-16.8	103	21.1	17.1-25.7	70	10.0	7.8-12.6	36	3.2	2.2-4.4	17	4.1	2.3-6.5	19	2.8	1.7-4.3
2007	174	14.4	12.4-16.8	106	21.4	17.4-25.9	68	9.7	7.5-12.2	38	3.2	2.3-4.4	26	5.6	3.6-8.3	12	1.6	0.8-2.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Kidney and renal pelvis**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	167	13.8	11.8-16.1	93	18.7	15.0-23.0	74	10.3	8.1-12.9	37	3.1	2.2-4.2	20	4.4	2.7-6.9	17	2.3	1.3-3.7
2009	180	14.3	12.2-16.5	106	20.1	16.3-24.4	74	10.0	7.8-12.6	37	3.0	2.1-4.1	28	5.7	3.7-8.3	9	1.2	0.5-2.3
2010	196	15.1	13.0-17.4	117	21.4	17.6-25.7	79	10.6	8.4-13.2	35	2.9	2.0-4.0	20	4.3	2.6-6.7	15	1.9	1.1-3.2
2011	202	15.6	13.5-17.9	121	22.5	18.6-27.0	81	10.6	8.4-13.2	38	3.0	2.1-4.1	27	5.7	3.7-8.3	11	1.3	0.6-2.4
2012	222	16.8	14.7-19.2	135	23.9	20.0-28.4	87	11.6	9.2-14.3	45	3.5	2.5-4.7	30	6.0	4.0-8.6	15	1.9	1.0-3.1
2013	212	15.7	13.6-18.0	130	22.7	18.9-27.1	82	10.5	8.3-13.0	36	2.8	1.9-3.8	22	4.2	2.6-6.4	14	1.7	0.9-2.9
2014	249	17.7	15.5-20.0	162	27.2	23.1-31.9	87	10.6	8.5-13.1	38	2.7	1.9-3.7	25	4.5	2.9-6.7	13	1.5	0.8-2.6
2015	247	17.2	15.1-19.6	147	23.8	20.0-28.1	100	12.2	9.9-14.9	42	3.0	2.2-4.1	26	4.7	3.0-6.9	16	1.8	1.0-3.0
2016	253	17.5	15.4-19.8	151	25.7	21.6-30.2	102	12.1	9.8-14.7	41	3.0	2.1-4.1	27	4.8	3.1-7.0	14	1.7	0.9-2.9
2017	227	14.9	13.0-17.0	136	21.0	17.6-24.9	91	10.5	8.4-12.9	38	2.6	1.8-3.6	24	4.2	2.7-6.3	14	1.6	0.9-2.7
2018	205	13.5	11.7-15.5	132	20.6	17.2-24.5	73	8.3	6.5-10.5	47	3.1	2.3-4.2	34	5.5	3.8-7.7	13	1.4	0.8-2.5
2019	260	17.1	15.0-19.3	155	23.3	19.7-27.4	105	12.5	10.2-15.2	44	2.8	2.1-3.8	25	4.0	2.5-5.9	19	2.0	1.2-3.2
2020	189	12.3	10.6-14.2	121	18.3	15.2-22.0	68	7.9	6.1-10.1	38	2.6	1.8-3.6	20	3.5	2.1-5.4	18	2.0	1.2-3.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Brain and other central nervous system**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	52	3.9	2.9-5.1	23	4.0	2.5-6.0	29	3.8	2.5-5.5	43	3.3	2.3-4.4	19	3.3	1.9-5.2	24	3.4	2.1-5.1
1977	64	5.4	4.1-6.9	34	6.6	4.5-9.3	30	4.7	3.1-6.7	42	3.4	2.4-4.6	21	4.2	2.5-6.4	21	2.9	1.8-4.5
1978	76	6.3	4.9-7.9	35	6.9	4.7-9.6	41	5.9	4.2-8.1	40	3.2	2.3-4.4	17	3.4	1.9-5.5	23	3.2	2.0-4.9
1979	50	4.2	3.1-5.6	19	3.8	2.3-6.0	31	4.6	3.1-6.6	39	3.4	2.4-4.7	16	3.5	2.0-5.7	23	3.2	2.0-4.9
1980	65	5.8	4.5-7.4	26	5.5	3.6-8.2	39	6.0	4.2-8.3	50	4.3	3.1-5.6	22	4.4	2.7-6.7	28	4.1	2.7-6.0
1981	53	4.5	3.3-5.9	24	5.1	3.2-7.6	29	4.1	2.7-6.0	41	3.5	2.5-4.8	13	2.8	1.5-4.9	28	4.0	2.6-5.8
1982	62	5.3	4.1-6.8	33	6.7	4.6-9.5	29	4.4	2.9-6.3	37	3.4	2.3-4.7	20	4.7	2.8-7.2	17	2.6	1.5-4.1
1983	52	4.7	3.5-6.1	30	6.4	4.2-9.2	22	3.3	2.1-5.1	34	2.9	2.0-4.1	16	3.2	1.8-5.2	18	2.7	1.6-4.3
1984	67	5.9	4.6-7.5	33	7.0	4.7-9.8	34	4.9	3.4-6.9	44	4.0	2.9-5.3	20	4.9	2.9-7.5	24	3.4	2.2-5.1
1985	60	5.4	4.1-7.0	36	7.9	5.4-10.9	24	3.6	2.3-5.4	44	3.9	2.8-5.3	23	4.8	3.0-7.3	21	3.1	1.9-4.8
1986	60	5.4	4.1-6.9	33	6.8	4.6-9.7	27	4.3	2.8-6.3	58	5.3	4.0-6.8	28	6.2	4.0-9.0	30	4.4	3.0-6.3
1987	73	6.2	4.9-7.9	35	7.5	5.2-10.5	38	5.3	3.7-7.3	42	3.8	2.7-5.1	20	4.5	2.7-6.9	22	3.2	2.0-4.9
1988	61	5.3	4.0-6.8	32	6.9	4.7-9.9	29	4.4	2.9-6.4	39	3.5	2.5-4.8	20	4.6	2.8-7.1	19	2.7	1.6-4.3
1989	71	6.3	4.9-8.0	34	7.1	4.8-10.0	37	5.8	4.1-8.1	21	1.9	1.2-2.9	10	2.2	1.0-4.1	11	1.6	0.8-2.8
1990	77	7.1	5.5-8.8	38	8.5	5.9-11.7	39	6.1	4.3-8.4	41	3.7	2.7-5.0	24	5.6	3.5-8.3	17	2.6	1.5-4.2
1991	80	7.2	5.7-9.0	31	6.7	4.5-9.6	49	7.5	5.5-10.0	43	4.0	2.9-5.4	21	4.6	2.8-7.0	22	3.5	2.2-5.4
1992	76	6.9	5.5-8.7	48	10.3	7.5-13.8	28	4.5	3.0-6.6	29	2.7	1.8-3.8	19	4.4	2.6-6.9	10	1.6	0.7-2.9
1993	64	5.5	4.3-7.1	31	6.1	4.0-8.8	33	5.2	3.6-7.3	44	4.1	3.0-5.5	22	5.4	3.4-8.2	22	3.5	2.2-5.2
1994	72	6.3	4.9-8.0	38	7.9	5.5-11.0	34	5.1	3.5-7.1	42	3.8	2.8-5.2	19	4.2	2.5-6.5	23	3.5	2.2-5.3
1995	70	6.2	4.8-7.8	42	8.7	6.2-11.9	28	4.2	2.8-6.1	32	2.9	2.0-4.1	16	3.7	2.1-6.0	16	2.4	1.4-3.9
1996	75	6.6	5.1-8.3	36	7.3	5.0-10.2	39	6.1	4.3-8.3	26	2.4	1.5-3.5	8	1.9	0.8-3.6	18	2.7	1.6-4.2
1997	66	5.6	4.3-7.1	31	6.0	4.0-8.6	35	5.2	3.6-7.3	33	3.0	2.0-4.2	16	3.4	1.9-5.5	17	2.7	1.6-4.3
1998	86	7.2	5.8-9.0	38	7.8	5.4-10.8	48	6.9	5.1-9.1	44	4.0	2.9-5.4	19	3.9	2.3-6.1	25	3.8	2.5-5.7
1999	67	5.7	4.4-7.3	35	7.0	4.8-9.9	32	4.7	3.2-6.6	26	2.3	1.5-3.3	11	2.0	1.0-3.5	15	2.3	1.3-3.7
2000	60	5.0	3.8-6.4	31	6.0	4.0-8.7	29	4.2	2.8-6.0	48	4.1	3.0-5.5	22	4.5	2.8-6.9	26	3.8	2.5-5.6
2001	54	4.5	3.4-5.9	22	4.6	2.8-7.0	32	4.6	3.1-6.5	29	2.5	1.7-3.6	17	3.7	2.1-6.0	12	1.8	0.9-3.1
2002	61	5.1	3.9-6.5	29	5.4	3.6-7.9	32	4.6	3.2-6.5	21	1.8	1.1-2.7	10	2.1	1.0-3.9	11	1.6	0.8-2.8
2003	77	6.2	4.9-7.8	41	7.6	5.4-10.4	36	5.1	3.6-7.1	38	3.2	2.2-4.4	21	4.4	2.7-6.8	17	2.4	1.4-3.9
2004	63	5.1	3.9-6.5	34	6.2	4.3-8.8	29	4.2	2.8-6.0	32	2.7	1.8-3.8	16	2.9	1.6-4.8	16	2.4	1.4-3.9
2005	65	5.1	3.9-6.5	39	6.9	4.9-9.5	26	3.7	2.4-5.4	32	2.6	1.7-3.6	24	4.5	2.8-6.7	8	1.1	0.5-2.2
2006	74	5.9	4.6-7.4	38	7.1	4.9-9.8	36	4.9	3.4-6.8	46	3.8	2.8-5.0	25	4.8	3.1-7.2	21	3.0	1.8-4.5
2007	65	5.0	3.8-6.4	34	6.3	4.3-8.9	31	4.2	2.9-6.0	30	2.4	1.6-3.5	15	2.8	1.5-4.6	15	2.1	1.2-3.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Brain and other central nervous system**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	56	4.4	3.3-5.8	31	5.8	3.9-8.4	25	3.5	2.2-5.1	25	1.9	1.3-2.9	11	2.4	1.2-4.4	14	1.8	1.0-3.0
2009	79	6.1	4.8-7.6	42	7.5	5.4-10.2	37	5.0	3.5-6.9	40	3.2	2.3-4.3	21	4.0	2.4-6.1	19	2.6	1.6-4.1
2010	52	3.9	2.9-5.1	28	4.8	3.2-7.0	24	3.2	2.0-4.7	36	2.7	1.9-3.7	17	3.2	1.8-5.1	19	2.4	1.5-3.8
2011	67	4.9	3.8-6.3	35	5.9	4.1-8.3	32	4.3	2.9-6.1	32	2.5	1.7-3.5	14	2.6	1.4-4.4	18	2.4	1.4-3.8
2012	72	5.3	4.1-6.7	38	6.3	4.4-8.7	34	4.5	3.1-6.2	35	2.6	1.8-3.6	21	3.6	2.2-5.6	14	1.9	1.0-3.1
2013	71	5.1	4.0-6.5	34	5.4	3.7-7.6	37	4.9	3.5-6.8	30	2.3	1.5-3.2	13	2.5	1.3-4.3	17	2.1	1.2-3.4
2014	64	4.7	3.6-6.0	35	5.8	4.0-8.0	29	3.9	2.6-5.6	39	2.8	2.0-3.9	16	3.0	1.7-4.8	23	2.8	1.8-4.2
2015	81	5.7	4.6-7.2	33	5.4	3.6-7.6	48	6.0	4.4-8.0	39	2.8	2.0-3.8	16	2.7	1.5-4.4	23	2.9	1.8-4.3
2016	70	4.8	3.8-6.1	36	5.6	3.9-7.8	34	4.2	2.9-5.9	44	3.1	2.2-4.1	23	3.9	2.4-5.9	21	2.6	1.6-4.0
2017	81	5.8	4.6-7.2	47	7.6	5.5-10.2	34	4.4	3.0-6.1	34	2.3	1.6-3.3	20	3.3	2.0-5.1	14	1.7	0.9-2.9
2018	79	5.4	4.2-6.7	40	6.2	4.4-8.4	39	4.8	3.4-6.5	50	3.4	2.5-4.5	28	4.7	3.1-6.8	22	2.5	1.5-3.8
2019	76	5.2	4.1-6.5	46	7.2	5.2-9.6	30	3.6	2.4-5.2	50	3.4	2.5-4.5	29	4.5	3.0-6.5	21	2.5	1.5-3.9
2020	71	4.7	3.7-5.9	43	6.4	4.6-8.7	28	3.2	2.1-4.7	49	3.3	2.4-4.4	25	4.0	2.5-5.9	24	2.9	1.9-4.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Thyroid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	29	2.4	1.6-3.4	6	1.1	0.4-2.4	23	3.4	2.1-5.1	13	1.2	0.6-2.1	4	0.7	0.2-1.9	9	1.6	0.7-3.0
1977	42	3.6	2.6-4.9	16	3.2	1.8-5.3	26	4.0	2.6-5.9	4	0.4	0.1-0.9	3	0.7	0.1-1.9	1	0.2	0.0-0.9
1978	36	3.3	2.3-4.6	5	1.1	0.4-2.6	31	4.9	3.3-7.0	6	0.5	0.2-1.2	1	0.2	0.0-1.0	5	0.8	0.2-1.8
1979	46	4.1	3.0-5.5	11	2.5	1.2-4.5	35	5.3	3.6-7.4	4	0.4	0.1-1.0	0	0.0	0.0-0.8	4	0.6	0.2-1.7
1980	25	2.2	1.4-3.2	4	0.9	0.2-2.3	21	3.0	1.9-4.7	6	0.6	0.2-1.3	1	0.3	0.0-1.4	5	0.8	0.2-1.8
1981	24	2.4	1.5-3.5	9	2.2	1.0-4.1	15	2.4	1.3-4.1	9	0.8	0.4-1.5	2	0.5	0.1-1.8	7	1.0	0.4-2.0
1982	37	3.6	2.5-5.0	10	2.4	1.1-4.3	27	4.5	2.9-6.6	6	0.5	0.2-1.1	1	0.2	0.0-1.2	5	0.7	0.2-1.7
1983	38	3.6	2.5-5.0	10	2.2	1.0-4.0	28	4.7	3.1-6.8	3	0.2	0.1-0.7	1	0.2	0.0-1.2	2	0.3	0.0-1.0
1984	46	4.5	3.3-6.0	7	1.6	0.6-3.4	39	6.5	4.6-9.0	6	0.5	0.2-1.1	2	0.4	0.0-1.4	4	0.6	0.2-1.5
1985	42	3.9	2.8-5.3	13	3.1	1.6-5.3	29	4.5	3.0-6.5	3	0.3	0.1-0.8	1	0.3	0.0-1.3	2	0.3	0.0-1.1
1986	37	3.5	2.4-4.8	6	1.6	0.6-3.4	31	5.0	3.3-7.1	6	0.6	0.2-1.3	0	0.0	0.0-0.8	6	1.0	0.4-2.2
1987	32	2.9	2.0-4.1	8	1.9	0.8-3.8	24	3.7	2.4-5.6	8	0.7	0.3-1.4	2	0.5	0.1-1.8	6	0.8	0.3-1.8
1988	34	3.0	2.1-4.2	11	2.3	1.1-4.2	23	3.5	2.2-5.3	6	0.6	0.2-1.3	3	0.8	0.2-2.1	3	0.5	0.1-1.5
1989	23	2.1	1.3-3.2	7	1.3	0.5-2.8	16	2.8	1.6-4.5	9	0.8	0.4-1.5	1	0.2	0.0-1.2	8	1.1	0.5-2.2
1990	29	2.6	1.7-3.8	2	0.4	0.0-1.4	27	4.3	2.8-6.3	6	0.6	0.2-1.2	3	0.8	0.2-2.2	3	0.4	0.1-1.2
1991	35	3.2	2.2-4.4	5	1.0	0.3-2.4	30	4.7	3.2-6.8	5	0.5	0.2-1.1	1	0.2	0.0-1.1	4	0.6	0.2-1.5
1992	48	4.1	3.0-5.4	11	2.2	1.0-4.0	37	5.6	3.9-7.8	11	1.0	0.5-1.8	2	0.5	0.1-1.7	9	1.3	0.6-2.5
1993	39	3.6	2.5-4.9	5	1.3	0.4-2.9	34	5.5	3.8-7.7	7	0.7	0.3-1.3	0	0.0	0.0-0.8	7	1.0	0.4-2.1
1994	52	4.7	3.5-6.2	11	2.6	1.3-4.6	41	6.5	4.6-8.8	6	0.6	0.2-1.3	3	0.7	0.1-2.1	3	0.5	0.1-1.5
1995	44	3.8	2.7-5.1	6	1.5	0.5-3.1	38	5.6	4.0-7.8	5	0.5	0.2-1.1	1	0.3	0.0-1.3	4	0.6	0.2-1.5
1996	49	4.2	3.1-5.6	9	2.0	0.9-3.7	40	6.0	4.3-8.2	4	0.4	0.1-1.0	0	0.0	0.0-0.8	4	0.6	0.2-1.6
1997	44	4.0	2.9-5.4	11	2.3	1.1-4.1	33	5.2	3.6-7.3	0	0.0	0.0-0.3	0	0.0	0.0-0.8	0	0.0	0.0-0.5
1998	71	6.0	4.7-7.6	15	3.3	1.8-5.5	56	8.3	6.2-10.8	3	0.3	0.1-0.8	3	0.7	0.1-2.0	0	0.0	0.0-0.5
1999	64	5.4	4.2-7.0	10	2.2	1.0-4.1	54	8.1	6.1-10.6	4	0.4	0.1-1.0	0	0.0	0.0-0.7	4	0.6	0.2-1.6
2000	74	6.2	4.8-7.7	15	3.1	1.7-5.1	59	8.7	6.6-11.3	6	0.6	0.2-1.2	2	0.6	0.1-2.0	4	0.6	0.2-1.5
2001	85	7.0	5.6-8.6	23	4.5	2.8-6.7	62	9.0	6.9-11.5	6	0.5	0.2-1.2	3	0.7	0.1-2.0	3	0.5	0.1-1.3
2002	91	7.4	6.0-9.1	18	3.4	2.0-5.4	73	10.6	8.3-13.4	8	0.7	0.3-1.4	1	0.2	0.0-1.1	7	1.0	0.4-2.1
2003	90	7.1	5.7-8.7	15	2.8	1.6-4.7	75	10.6	8.3-13.2	4	0.4	0.1-0.9	2	0.5	0.1-1.7	2	0.3	0.0-1.1
2004	91	7.3	5.8-8.9	22	4.5	2.8-6.8	69	9.7	7.5-12.3	4	0.4	0.1-0.9	2	0.5	0.1-1.8	2	0.3	0.0-1.0
2005	121	9.6	8.0-11.5	28	5.3	3.4-7.6	93	13.2	10.6-16.2	6	0.5	0.2-1.1	3	0.6	0.1-1.8	3	0.4	0.1-1.3
2006	100	8.0	6.5-9.7	19	3.6	2.1-5.6	81	11.6	9.2-14.4	4	0.4	0.1-0.9	0	0.0	0.0-0.7	4	0.6	0.2-1.5
2007	117	9.1	7.5-11.0	20	3.6	2.2-5.7	97	13.6	11.0-16.6	5	0.4	0.1-1.0	0	0.0	0.0-0.7	5	0.7	0.2-1.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Thyroid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	155	12.0	10.1-14.0	29	5.0	3.3-7.3	126	17.5	14.6-20.9	7	0.6	0.2-1.3	3	0.9	0.2-2.3	4	0.5	0.1-1.3
2009	157	11.9	10.1-14.0	38	6.9	4.8-9.5	119	16.4	13.6-19.6	5	0.4	0.1-1.0	4	0.8	0.2-2.0	1	0.2	0.0-0.9
2010	171	12.8	11.0-14.9	34	6.3	4.3-8.8	137	18.5	15.5-21.9	4	0.3	0.1-0.9	1	0.3	0.0-1.3	3	0.4	0.1-1.3
2011	195	14.3	12.3-16.5	39	6.9	4.9-9.5	156	20.6	17.5-24.1	5	0.4	0.1-1.0	3	0.7	0.2-2.0	2	0.2	0.0-0.9
2012	221	16.1	14.0-18.4	29	4.7	3.2-6.8	192	25.2	21.7-29.1	5	0.4	0.1-1.0	1	0.3	0.0-1.2	4	0.5	0.1-1.4
2013	234	17.4	15.2-19.8	34	5.8	4.0-8.2	200	26.9	23.2-30.9	6	0.5	0.2-1.0	1	0.2	0.0-1.0	5	0.6	0.2-1.5
2014	260	18.6	16.3-21.0	42	6.9	4.9-9.4	218	28.3	24.6-32.3	13	0.9	0.5-1.6	3	0.6	0.1-1.7	10	1.2	0.6-2.2
2015	235	17.0	14.9-19.4	36	5.9	4.1-8.2	199	26.2	22.6-30.2	7	0.5	0.2-1.0	4	0.6	0.2-1.5	3	0.3	0.1-1.0
2016	232	16.2	14.1-18.4	43	6.7	4.8-9.1	189	23.9	20.5-27.6	8	0.6	0.2-1.1	3	0.6	0.1-1.7	5	0.5	0.2-1.3
2017	212	15.1	13.1-17.3	43	7.0	5.0-9.5	169	22.0	18.7-25.6	7	0.5	0.2-1.1	0	0.0	0.0-0.6	7	0.9	0.4-1.8
2018	200	13.9	12.0-16.0	39	5.9	4.2-8.2	161	20.7	17.6-24.2	8	0.6	0.2-1.1	5	0.8	0.2-1.9	3	0.4	0.1-1.1
2019	203	14.4	12.4-16.6	43	6.6	4.7-8.9	160	21.1	17.9-24.7	8	0.6	0.2-1.2	5	0.9	0.3-2.1	3	0.4	0.1-1.1
2020	172	12.3	10.5-14.3	41	6.8	4.8-9.2	131	17.2	14.3-20.5	4	0.3	0.1-0.7	2	0.3	0.0-1.1	2	0.2	0.0-0.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Hodgkin lymphoma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	41	3.2	2.3-4.4	16	3.1	1.7-5.1	25	3.4	2.2-5.1	17	1.4	0.8-2.2	12	2.1	1.1-3.7	5	0.7	0.2-1.8
1977	37	2.9	2.0-4.0	23	3.8	2.4-5.8	14	2.2	1.2-3.7	13	1.1	0.6-1.9	6	1.0	0.4-2.4	7	1.1	0.4-2.3
1978	40	3.2	2.3-4.4	18	3.4	1.9-5.4	22	3.2	2.0-4.9	18	1.6	0.9-2.5	9	1.8	0.8-3.5	9	1.4	0.6-2.7
1979	36	2.9	2.0-4.1	19	3.6	2.1-5.7	17	2.5	1.5-4.1	14	1.1	0.6-1.9	9	1.7	0.7-3.2	5	0.6	0.2-1.5
1980	35	2.9	2.0-4.1	22	4.2	2.6-6.5	13	1.9	1.0-3.2	8	0.8	0.3-1.5	3	0.8	0.2-2.2	5	0.8	0.2-1.8
1981	23	2.0	1.2-3.0	13	2.5	1.3-4.3	10	1.5	0.7-2.9	14	1.2	0.7-2.1	7	1.6	0.6-3.3	7	0.9	0.4-2.0
1982	39	3.2	2.3-4.5	20	3.8	2.3-6.1	19	2.7	1.6-4.2	10	0.9	0.4-1.6	5	1.1	0.3-2.5	5	0.7	0.2-1.7
1983	43	3.9	2.8-5.3	25	5.2	3.3-7.8	18	2.7	1.6-4.3	8	0.6	0.3-1.2	4	0.6	0.2-1.7	4	0.6	0.2-1.6
1984	35	3.0	2.1-4.2	19	3.6	2.2-5.8	16	2.4	1.3-3.9	17	1.6	0.9-2.6	8	1.7	0.7-3.4	9	1.4	0.6-2.7
1985	24	2.0	1.3-3.0	13	2.6	1.3-4.6	11	1.4	0.7-2.6	8	0.7	0.3-1.4	4	0.6	0.2-1.7	4	0.7	0.2-1.8
1986	33	2.9	1.9-4.1	19	3.9	2.3-6.2	14	2.1	1.1-3.6	5	0.4	0.1-0.9	4	0.7	0.2-1.8	1	0.1	0.0-0.8
1987	39	3.3	2.3-4.5	21	3.9	2.4-6.1	18	2.9	1.7-4.6	8	0.6	0.3-1.2	6	0.9	0.3-2.2	2	0.3	0.0-1.1
1988	38	3.0	2.1-4.2	29	5.3	3.4-7.7	9	1.2	0.5-2.3	10	0.9	0.4-1.7	7	1.6	0.6-3.3	3	0.4	0.1-1.3
1989	29	2.4	1.6-3.5	18	3.2	1.8-5.1	11	1.8	0.9-3.2	10	0.9	0.4-1.6	8	1.6	0.7-3.3	2	0.3	0.0-1.1
1990	44	3.6	2.6-4.9	27	5.1	3.3-7.5	17	2.5	1.4-4.0	9	0.8	0.4-1.6	6	1.4	0.5-3.0	3	0.4	0.1-1.3
1991	57	4.7	3.5-6.1	33	6.3	4.3-8.9	24	3.5	2.2-5.2	9	0.9	0.4-1.7	5	1.2	0.4-2.8	4	0.7	0.2-1.7
1992	53	4.4	3.3-5.8	28	5.2	3.4-7.5	25	3.6	2.3-5.4	19	1.6	0.9-2.5	12	2.4	1.2-4.2	7	1.0	0.4-2.1
1993	28	2.5	1.6-3.6	17	3.4	2.0-5.6	11	1.7	0.8-3.1	13	1.2	0.6-2.1	6	1.2	0.4-2.7	7	1.1	0.4-2.2
1994	38	3.0	2.1-4.2	23	3.9	2.4-6.0	15	2.2	1.2-3.7	13	1.1	0.6-2.0	6	1.4	0.5-3.0	7	1.1	0.4-2.2
1995	32	2.8	1.9-4.0	20	4.0	2.4-6.3	12	1.9	0.9-3.2	12	1.1	0.5-1.9	7	1.5	0.6-3.1	5	0.8	0.2-1.8
1996	40	3.2	2.3-4.4	25	4.7	2.9-7.0	15	2.3	1.3-3.7	11	1.0	0.5-1.7	5	1.1	0.3-2.5	6	0.9	0.3-1.9
1997	40	3.2	2.3-4.4	24	4.6	2.9-6.9	16	2.3	1.3-3.8	9	0.8	0.4-1.5	5	0.9	0.3-2.2	4	0.7	0.2-1.7
1998	42	3.3	2.4-4.5	28	4.9	3.2-7.2	14	2.1	1.1-3.5	9	0.8	0.4-1.5	4	0.8	0.2-2.0	5	0.8	0.3-1.8
1999	31	2.5	1.7-3.5	16	2.9	1.6-4.8	15	2.2	1.2-3.6	10	0.8	0.4-1.5	5	1.1	0.3-2.6	5	0.7	0.2-1.6
2000	44	3.6	2.6-4.8	32	5.9	4.0-8.4	12	1.7	0.9-3.0	11	1.0	0.5-1.7	8	1.7	0.7-3.4	3	0.5	0.1-1.3
2001	40	3.0	2.1-4.1	22	3.6	2.3-5.5	18	2.5	1.5-3.9	8	0.7	0.3-1.3	6	1.1	0.4-2.4	2	0.3	0.0-1.0
2002	49	3.8	2.8-5.1	31	5.6	3.8-8.1	18	2.5	1.5-4.0	5	0.4	0.1-1.0	4	0.9	0.2-2.2	1	0.1	0.0-0.7
2003	50	3.8	2.8-5.1	29	4.7	3.1-6.8	21	3.0	1.9-4.6	6	0.5	0.2-1.1	4	0.8	0.2-2.1	2	0.3	0.0-1.0
2004	46	3.5	2.6-4.7	22	3.7	2.3-5.6	24	3.4	2.2-5.0	7	0.6	0.2-1.2	5	0.9	0.3-2.1	2	0.3	0.0-1.0
2005	42	3.2	2.3-4.4	18	3.0	1.8-4.8	24	3.4	2.2-5.1	5	0.4	0.1-0.9	3	0.5	0.1-1.5	2	0.3	0.0-1.0
2006	42	3.3	2.4-4.5	22	3.8	2.4-5.8	20	2.8	1.7-4.3	7	0.6	0.2-1.2	4	0.7	0.2-1.8	3	0.5	0.1-1.4
2007	38	2.8	1.9-3.8	15	2.3	1.3-3.8	23	3.1	1.9-4.6	6	0.5	0.2-1.0	4	0.8	0.2-2.1	2	0.3	0.0-1.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Hodgkin lymphoma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	45	3.3	2.4-4.4	22	3.6	2.2-5.5	23	3.1	1.9-4.6	10	0.8	0.4-1.5	5	0.9	0.3-2.0	5	0.6	0.2-1.5
2009	44	3.2	2.3-4.2	21	3.6	2.2-5.6	23	2.9	1.8-4.4	7	0.6	0.2-1.1	5	1.0	0.3-2.4	2	0.3	0.0-0.9
2010	48	3.5	2.5-4.6	26	4.0	2.6-5.9	22	3.1	1.9-4.6	1	0.1	0.0-0.5	0	0.0	0.0-0.7	1	0.2	0.0-0.8
2011	50	3.4	2.5-4.5	25	3.6	2.3-5.4	25	3.2	2.0-4.7	6	0.5	0.2-1.0	5	1.0	0.3-2.3	1	0.2	0.0-0.8
2012	44	3.0	2.2-4.1	18	2.5	1.5-4.0	26	3.5	2.3-5.1	3	0.2	0.0-0.7	2	0.3	0.0-1.2	1	0.1	0.0-0.8
2013	61	4.3	3.3-5.5	33	5.1	3.5-7.3	28	3.6	2.4-5.2	7	0.5	0.2-1.1	3	0.5	0.1-1.4	4	0.5	0.1-1.4
2014	52	3.6	2.7-4.8	22	3.7	2.3-5.7	30	3.7	2.5-5.3	7	0.5	0.2-0.9	2	0.3	0.0-1.1	5	0.5	0.2-1.3
2015	55	3.8	2.8-5.0	32	4.7	3.2-6.7	23	3.0	1.9-4.5	6	0.4	0.1-0.9	2	0.4	0.1-1.4	4	0.4	0.1-1.2
2016	37	2.6	1.9-3.7	20	3.3	2.0-5.1	17	2.2	1.2-3.5	6	0.5	0.2-1.0	5	0.9	0.3-2.1	1	0.1	0.0-0.8
2017	41	2.8	2.0-3.8	23	3.4	2.2-5.2	18	2.2	1.3-3.6	2	0.2	0.0-0.6	0	0.0	0.0-0.6	2	0.3	0.0-1.0
2018	39	2.7	1.9-3.7	22	3.3	2.1-5.1	17	2.1	1.2-3.4	2	0.1	0.0-0.5	2	0.3	0.0-1.0	0	0.0	0.0-0.5
2019	56	3.9	3.0-5.1	26	3.9	2.5-5.7	30	4.0	2.7-5.7	7	0.5	0.2-1.0	5	0.7	0.2-1.6	2	0.3	0.0-0.9
2020	33	2.3	1.6-3.2	20	2.9	1.8-4.6	13	1.8	0.9-3.1	6	0.4	0.2-1.0	3	0.5	0.1-1.4	3	0.4	0.1-1.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Non-Hodgkin lymphomas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	133	10.8	9.0-12.8	70	14.4	11.1-18.3	63	8.5	6.5-10.9	82	6.8	5.4-8.5	48	10.4	7.6-14.0	34	4.5	3.1-6.4
1977	140	11.2	9.4-13.3	63	11.7	9.0-15.1	77	10.7	8.4-13.4	71	5.8	4.5-7.3	28	5.6	3.6-8.1	43	5.8	4.2-7.8
1978	131	10.9	9.1-13.0	61	12.4	9.4-16.0	70	9.9	7.7-12.6	81	6.7	5.3-8.3	41	8.3	5.9-11.3	40	5.6	4.0-7.7
1979	106	9.1	7.4-11.1	53	11.6	8.6-15.2	53	7.2	5.4-9.4	50	4.4	3.3-5.9	19	4.6	2.7-7.1	31	4.4	3.0-6.3
1980	142	12.9	10.8-15.2	74	16.1	12.6-20.3	68	10.5	8.1-13.4	78	6.6	5.2-8.3	43	8.6	6.2-11.7	35	5.0	3.5-7.0
1981	128	11.5	9.5-13.7	68	15.5	12.0-19.8	60	8.7	6.6-11.3	67	6.3	4.8-8.0	36	8.4	5.8-11.6	31	4.9	3.3-7.0
1982	146	12.9	10.8-15.2	57	12.6	9.4-16.4	89	13.0	10.4-16.0	63	5.8	4.4-7.4	27	6.5	4.2-9.4	36	5.5	3.8-7.6
1983	148	13.1	11.1-15.5	67	15.2	11.7-19.5	81	11.9	9.4-14.9	80	7.3	5.7-9.1	42	10.2	7.3-13.8	38	5.5	3.9-7.7
1984	150	13.2	11.1-15.5	65	14.8	11.4-18.9	85	12.2	9.7-15.2	67	6.1	4.7-7.7	33	8.0	5.5-11.3	34	4.8	3.3-6.7
1985	168	15.4	13.1-17.9	81	18.8	14.8-23.5	87	13.1	10.5-16.3	69	6.1	4.7-7.7	30	6.8	4.5-9.7	39	5.7	4.0-7.8
1986	174	15.4	13.1-17.9	90	19.2	15.3-23.7	84	12.5	10.0-15.6	84	7.4	5.9-9.2	40	8.9	6.3-12.2	44	6.5	4.7-8.8
1987	193	17.4	15.0-20.0	102	23.0	18.6-28.0	91	13.1	10.5-16.2	85	7.7	6.1-9.5	42	9.7	6.9-13.2	43	6.2	4.4-8.3
1988	160	14.5	12.3-17.0	81	17.9	14.1-22.4	79	12.0	9.5-15.0	89	8.2	6.6-10.1	46	11.4	8.3-15.2	43	6.1	4.4-8.2
1989	166	15.4	13.1-17.9	89	20.2	16.1-25.0	77	11.5	9.1-14.4	78	7.2	5.7-9.0	44	10.5	7.5-14.1	34	5.0	3.4-7.0
1990	183	16.7	14.3-19.3	96	21.1	17.0-25.9	87	13.2	10.5-16.3	77	7.3	5.8-9.2	36	8.6	6.0-11.9	41	6.4	4.6-8.8
1991	218	20.1	17.5-23.0	113	25.5	20.9-30.8	105	16.2	13.2-19.7	89	8.2	6.5-10.1	47	11.1	8.0-14.8	42	6.1	4.4-8.3
1992	187	17.2	14.8-19.9	97	21.4	17.2-26.2	90	13.7	11.0-16.9	80	7.6	6.0-9.4	38	9.4	6.6-12.9	42	6.5	4.7-8.9
1993	216	19.7	17.1-22.5	120	26.1	21.5-31.3	96	14.7	11.9-18.0	87	8.1	6.5-10.0	52	12.6	9.3-16.6	35	5.2	3.6-7.3
1994	266	24.3	21.4-27.4	147	31.9	26.8-37.6	119	18.6	15.4-22.3	85	8.0	6.4-9.9	34	7.8	5.3-10.9	51	8.1	6.0-10.6
1995	235	21.4	18.7-24.3	126	27.1	22.5-32.4	109	16.7	13.7-20.2	78	7.4	5.8-9.2	39	9.4	6.6-12.8	39	6.0	4.2-8.2
1996	252	22.9	20.1-25.9	137	29.7	24.8-35.2	115	17.7	14.6-21.3	86	8.1	6.5-10.0	45	11.1	8.0-14.8	41	6.3	4.5-8.5
1997	244	22.0	19.3-25.0	129	27.5	22.9-32.8	115	17.7	14.6-21.2	74	6.7	5.3-8.5	37	8.2	5.7-11.4	37	5.2	3.7-7.3
1998	247	22.1	19.4-25.0	116	24.1	19.8-29.0	131	19.9	16.7-23.7	81	7.6	6.0-9.4	33	7.8	5.3-10.9	48	7.3	5.4-9.7
1999	199	17.4	15.0-20.0	107	21.9	17.9-26.6	92	14.1	11.3-17.3	97	8.8	7.1-10.7	44	9.5	6.9-12.8	53	8.0	6.0-10.5
2000	231	20.0	17.5-22.7	110	22.5	18.4-27.2	121	17.9	14.8-21.4	77	6.8	5.3-8.5	33	7.5	5.1-10.5	44	6.4	4.7-8.6
2001	198	17.1	14.8-19.7	105	22.1	18.0-26.9	93	13.6	11.0-16.6	77	6.8	5.3-8.5	34	7.4	5.0-10.3	43	6.1	4.4-8.2
2002	237	20.0	17.6-22.8	132	27.3	22.8-32.5	105	15.2	12.4-18.4	61	5.4	4.1-6.9	29	6.2	4.1-8.9	32	4.5	3.1-6.4
2003	269	22.4	19.8-25.3	131	26.0	21.6-30.9	138	19.7	16.6-23.3	75	6.3	5.0-8.0	39	8.0	5.6-11.0	36	5.1	3.6-7.1
2004	234	19.6	17.2-22.3	115	22.5	18.5-27.1	119	17.1	14.2-20.5	70	6.0	4.7-7.6	33	6.8	4.6-9.6	37	5.3	3.7-7.3
2005	237	19.8	17.3-22.5	112	21.8	17.8-26.3	125	18.1	15.0-21.5	64	5.6	4.3-7.1	29	5.9	3.9-8.6	35	5.1	3.5-7.1
2006	283	22.8	20.2-25.7	155	30.0	25.3-35.3	128	17.8	14.9-21.2	71	6.0	4.7-7.6	34	6.7	4.6-9.5	37	5.1	3.5-7.0
2007	254	20.4	17.9-23.0	132	24.4	20.3-29.1	122	17.0	14.1-20.3	80	6.7	5.3-8.3	36	7.7	5.3-10.7	44	6.1	4.4-8.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Non-Hodgkin lymphomas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	233	18.7	16.4-21.3	131	24.2	20.1-28.8	102	14.4	11.7-17.5	68	5.5	4.3-7.0	38	7.6	5.3-10.6	30	4.1	2.7-5.8
2009	279	21.9	19.4-24.6	137	25.0	20.9-29.7	142	19.3	16.2-22.7	72	5.7	4.5-7.2	39	7.2	5.0-9.8	33	4.5	3.1-6.4
2010	263	20.6	18.2-23.3	144	26.6	22.3-31.4	119	16.1	13.3-19.3	61	4.9	3.8-6.3	26	5.4	3.5-8.0	35	4.7	3.2-6.5
2011	273	20.9	18.5-23.6	142	25.7	21.6-30.5	131	17.1	14.3-20.3	66	5.2	4.0-6.7	40	8.3	5.8-11.3	26	3.4	2.2-5.0
2012	262	19.5	17.2-22.0	141	24.8	20.8-29.4	121	15.8	13.1-18.9	74	5.6	4.4-7.0	34	6.4	4.4-9.0	40	5.0	3.5-6.8
2013	256	18.6	16.4-21.1	129	22.5	18.7-26.9	127	16.2	13.5-19.3	74	5.5	4.3-7.0	41	7.8	5.5-10.6	33	4.1	2.8-5.7
2014	263	18.5	16.3-21.0	140	24.0	20.1-28.4	123	15.1	12.5-18.1	66	4.9	3.8-6.3	39	7.4	5.2-10.1	27	3.4	2.2-5.0
2015	265	18.9	16.7-21.4	138	23.6	19.8-28.0	127	15.8	13.1-18.8	74	5.6	4.4-7.0	40	7.8	5.5-10.6	34	4.3	2.9-6.0
2016	270	18.8	16.6-21.2	158	26.1	22.1-30.7	112	13.5	11.1-16.3	73	5.1	4.0-6.4	43	7.4	5.3-10.1	30	3.6	2.4-5.2
2017	282	19.6	17.3-22.0	135	21.9	18.2-26.0	147	17.7	14.9-20.9	70	4.8	3.7-6.1	36	6.3	4.4-8.8	34	3.8	2.6-5.3
2018	317	21.5	19.1-24.0	145	22.9	19.2-27.0	172	20.1	17.2-23.4	75	5.0	3.9-6.3	41	6.5	4.6-8.9	34	3.9	2.7-5.5
2019	278	18.8	16.6-21.2	142	22.6	19.0-26.7	136	16.6	13.8-19.7	63	4.2	3.2-5.4	37	5.9	4.1-8.2	26	3.0	1.9-4.4
2020	239	15.8	13.8-18.0	139	21.5	18.0-25.5	100	11.7	9.5-14.3	63	4.2	3.2-5.4	39	6.2	4.4-8.6	24	2.8	1.8-4.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Myeloma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	57	4.6	3.4-6.0	32	6.2	4.2-8.8	25	3.4	2.2-5.1	39	3.2	2.3-4.4	21	4.3	2.6-6.7	18	2.4	1.4-3.9
1977	57	4.9	3.7-6.4	29	6.8	4.5-9.9	28	3.8	2.5-5.6	45	3.8	2.7-5.1	26	5.5	3.5-8.1	19	2.7	1.6-4.3
1978	67	5.6	4.3-7.1	32	7.0	4.8-10.0	35	4.7	3.3-6.6	37	3.1	2.1-4.2	15	3.2	1.8-5.3	22	2.9	1.8-4.5
1979	59	5.0	3.8-6.5	31	6.9	4.6-9.8	28	3.8	2.5-5.6	43	3.7	2.7-5.1	15	3.6	2.0-5.9	28	3.9	2.6-5.7
1980	64	5.8	4.5-7.4	32	7.4	5.0-10.4	32	4.8	3.3-6.9	35	3.1	2.1-4.3	19	4.4	2.6-6.8	16	2.3	1.3-3.8
1981	64	5.8	4.4-7.4	27	6.5	4.2-9.5	37	5.4	3.8-7.4	37	3.3	2.3-4.5	17	4.3	2.4-6.8	20	2.8	1.7-4.4
1982	66	6.0	4.6-7.7	33	7.5	5.1-10.6	33	5.1	3.5-7.2	41	3.8	2.7-5.1	14	3.1	1.7-5.3	27	4.1	2.7-6.1
1983	65	5.9	4.6-7.6	29	6.8	4.5-9.8	36	5.4	3.7-7.5	37	3.2	2.3-4.5	15	3.4	1.9-5.6	22	3.0	1.9-4.6
1984	55	5.2	3.9-6.8	21	5.4	3.3-8.3	34	5.3	3.6-7.5	42	3.8	2.7-5.1	23	5.5	3.5-8.3	19	2.8	1.7-4.5
1985	82	7.6	6.0-9.4	35	8.5	5.9-11.8	47	6.7	4.9-9.0	32	2.9	1.9-4.0	22	5.1	3.2-7.7	10	1.5	0.7-2.7
1986	70	6.4	5.0-8.1	33	8.0	5.5-11.3	37	5.4	3.8-7.5	32	2.9	2.0-4.1	14	3.3	1.8-5.5	18	2.7	1.6-4.3
1987	55	4.8	3.6-6.3	27	6.3	4.2-9.2	28	3.9	2.6-5.7	45	4.3	3.1-5.7	17	4.3	2.5-6.8	28	4.0	2.7-5.8
1988	50	4.7	3.5-6.2	25	6.5	4.2-9.5	25	3.8	2.4-5.6	37	3.5	2.4-4.8	16	4.2	2.4-6.8	21	3.0	1.9-4.7
1989	54	5.1	3.8-6.6	23	5.4	3.4-8.2	31	4.7	3.2-6.7	37	3.4	2.4-4.7	19	4.8	2.9-7.5	18	2.5	1.5-4.0
1990	71	6.6	5.2-8.4	27	6.4	4.2-9.3	44	7.0	5.1-9.4	31	2.9	2.0-4.1	16	4.1	2.3-6.7	15	2.3	1.3-3.8
1991	60	5.6	4.3-7.3	25	6.1	3.9-8.9	35	5.3	3.7-7.4	29	2.7	1.8-3.9	7	1.8	0.7-3.6	22	3.3	2.0-5.0
1992	93	8.9	7.2-10.9	43	10.7	7.7-14.4	50	7.9	5.8-10.4	54	5.2	3.9-6.7	23	5.6	3.6-8.4	31	4.8	3.2-6.8
1993	64	6.1	4.7-7.8	36	9.3	6.5-12.9	28	4.4	2.9-6.3	57	5.4	4.1-7.1	28	7.0	4.6-10.2	29	4.6	3.1-6.6
1994	72	6.9	5.4-8.6	30	7.3	4.9-10.5	42	6.6	4.7-8.9	47	4.5	3.3-5.9	17	4.5	2.6-7.1	30	4.8	3.2-6.8
1995	70	6.7	5.2-8.5	26	6.2	4.0-9.0	44	7.0	5.1-9.4	36	3.4	2.4-4.8	16	3.8	2.2-6.2	20	3.1	1.9-4.8
1996	82	7.9	6.3-9.8	36	8.9	6.2-12.3	46	7.3	5.3-9.7	47	4.5	3.3-6.0	26	6.4	4.1-9.3	21	3.3	2.0-5.1
1997	80	7.6	6.1-9.5	38	9.2	6.5-12.7	42	6.7	4.8-9.0	49	4.7	3.5-6.2	25	6.4	4.1-9.4	24	3.7	2.4-5.5
1998	76	7.2	5.7-9.0	31	7.6	5.1-10.8	45	7.1	5.2-9.5	35	3.3	2.3-4.6	17	4.0	2.3-6.4	18	2.7	1.6-4.3
1999	82	7.6	6.0-9.4	30	7.3	4.9-10.4	52	7.9	5.9-10.4	42	3.8	2.8-5.2	19	4.5	2.7-7.0	23	3.4	2.2-5.2
2000	96	8.6	7.0-10.6	51	11.9	8.8-15.7	45	6.7	4.9-9.0	54	5.0	3.7-6.5	17	4.0	2.3-6.3	37	5.5	3.9-7.6
2001	103	9.3	7.6-11.2	49	11.0	8.1-14.6	54	8.1	6.1-10.5	47	4.3	3.1-5.6	21	4.9	3.0-7.5	26	3.9	2.5-5.7
2002	120	10.6	8.7-12.6	49	10.6	7.8-14.0	71	10.5	8.2-13.2	46	4.1	3.0-5.5	21	5.0	3.0-7.6	25	3.6	2.3-5.3
2003	114	10.0	8.2-12.0	47	10.6	7.8-14.2	67	9.8	7.6-12.4	43	3.8	2.7-5.1	20	4.8	2.9-7.3	23	3.2	2.0-4.9
2004	93	8.0	6.4-9.7	35	7.2	5.0-10.1	58	8.4	6.4-10.9	39	3.5	2.5-4.7	14	2.9	1.6-4.9	25	3.7	2.4-5.4
2005	117	10.0	8.2-12.0	53	11.5	8.6-15.1	64	8.8	6.8-11.3	51	4.3	3.2-5.7	16	3.5	1.9-5.6	35	4.8	3.3-6.7
2006	127	11.0	9.1-13.1	56	11.4	8.6-14.9	71	10.4	8.1-13.1	44	3.9	2.8-5.2	23	5.0	3.1-7.5	21	3.0	1.9-4.7
2007	111	9.3	7.7-11.2	54	11.9	8.9-15.6	57	8.0	6.1-10.4	51	4.4	3.3-5.8	30	6.6	4.4-9.5	21	3.0	1.9-4.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Myeloma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	134	10.9	9.2-13.0	62	12.9	9.8-16.7	72	9.8	7.6-12.3	42	3.6	2.6-4.8	22	5.0	3.1-7.6	20	2.8	1.7-4.4
2009	127	10.3	8.6-12.3	66	13.4	10.3-17.1	61	8.3	6.4-10.7	49	4.0	3.0-5.3	24	5.0	3.2-7.5	25	3.4	2.2-5.0
2010	145	11.7	9.9-13.8	80	16.3	12.8-20.4	65	8.9	6.9-11.4	42	3.5	2.5-4.7	20	4.5	2.7-6.9	22	2.9	1.8-4.4
2011	144	11.1	9.3-13.1	68	12.8	9.9-16.4	76	10.1	7.9-12.6	49	3.9	2.8-5.1	19	4.1	2.4-6.4	30	3.8	2.6-5.5
2012	163	12.7	10.8-14.9	82	15.7	12.4-19.6	81	10.7	8.4-13.3	50	3.9	2.9-5.2	27	5.7	3.8-8.4	23	2.9	1.8-4.4
2013	140	10.4	8.7-12.3	76	13.5	10.6-17.0	64	8.2	6.3-10.5	46	3.5	2.5-4.7	23	4.4	2.7-6.6	23	2.9	1.9-4.4
2014	153	10.9	9.3-12.9	91	15.7	12.5-19.3	62	7.8	5.9-10.0	40	2.9	2.1-4.0	21	3.8	2.3-5.8	19	2.3	1.4-3.7
2015	148	10.5	8.9-12.4	73	12.6	9.8-15.9	75	9.2	7.2-11.6	43	3.0	2.1-4.0	19	3.2	1.9-5.0	24	2.8	1.8-4.2
2016	190	13.2	11.3-15.2	106	17.9	14.6-21.7	84	10.1	8.0-12.5	63	4.5	3.5-5.8	31	5.3	3.6-7.6	32	4.0	2.7-5.7
2017	189	12.7	10.9-14.6	95	14.7	11.9-18.1	94	11.1	8.9-13.6	51	3.4	2.5-4.5	32	5.2	3.5-7.4	19	2.1	1.3-3.4
2018	180	11.6	9.9-13.4	80	12.4	9.7-15.5	100	11.3	9.2-13.8	46	3.0	2.2-4.0	28	4.7	3.1-6.8	18	1.8	1.1-2.9
2019	160	10.6	9.0-12.4	85	13.2	10.5-16.4	75	8.6	6.8-10.9	56	3.6	2.7-4.7	29	4.7	3.1-6.8	27	2.9	1.9-4.3
2020	138	9.0	7.5-10.6	59	9.3	7.0-12.1	79	9.0	7.1-11.3	48	3.0	2.2-4.1	26	3.9	2.5-5.8	22	2.5	1.6-3.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Leukemias**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	134	10.8	9.0-12.8	75	14.6	11.4-18.5	59	8.2	6.2-10.6	94	7.8	6.3-9.6	49	9.5	6.9-12.7	45	6.5	4.7-8.8
1977	135	11.4	9.6-13.6	73	15.1	11.7-19.1	62	9.0	6.8-11.5	93	7.8	6.3-9.6	51	10.7	7.9-14.2	42	5.9	4.2-8.1
1978	122	10.6	8.8-12.7	63	14.3	10.9-18.5	59	8.5	6.5-11.1	96	8.2	6.6-10.0	49	10.6	7.7-14.1	47	6.7	4.9-9.0
1979	129	11.0	9.2-13.1	65	13.9	10.6-17.9	64	9.0	6.9-11.5	98	8.7	7.0-10.6	45	9.8	7.1-13.3	53	8.0	5.9-10.5
1980	115	10.0	8.3-12.1	56	12.4	9.2-16.2	59	8.7	6.6-11.2	94	8.4	6.7-10.3	53	12.2	9.1-16.1	41	6.0	4.3-8.2
1981	123	10.5	8.7-12.5	62	13.6	10.3-17.6	61	8.7	6.6-11.2	76	6.6	5.2-8.3	39	9.2	6.5-12.7	37	5.3	3.7-7.4
1982	134	12.1	10.1-14.3	73	16.2	12.6-20.5	61	9.2	7.0-11.9	96	8.6	7.0-10.6	47	10.2	7.4-13.7	49	7.7	5.6-10.3
1983	133	11.7	9.8-13.9	70	15.9	12.3-20.2	63	9.1	7.0-11.7	79	7.1	5.6-8.9	45	10.3	7.4-13.8	34	4.9	3.4-6.8
1984	143	13.1	11.0-15.5	80	18.9	14.9-23.6	63	9.5	7.3-12.2	84	7.6	6.0-9.4	51	12.3	9.1-16.2	33	4.7	3.3-6.7
1985	109	9.8	8.0-11.8	72	15.9	12.4-20.1	37	5.6	3.9-7.7	93	8.2	6.6-10.1	50	11.0	8.1-14.6	43	6.2	4.5-8.4
1986	124	11.0	9.1-13.1	53	11.9	8.9-15.7	71	10.8	8.4-13.7	67	6.1	4.7-7.8	32	7.5	5.1-10.7	35	5.3	3.6-7.4
1987	112	9.9	8.1-11.9	45	10.6	7.7-14.3	67	9.7	7.5-12.3	71	6.1	4.7-7.7	29	6.4	4.2-9.3	42	6.1	4.3-8.2
1988	122	10.9	9.0-13.1	66	15.3	11.8-19.6	56	8.2	6.2-10.7	66	5.8	4.5-7.4	28	6.9	4.5-10.0	38	5.4	3.8-7.5
1989	129	11.5	9.6-13.7	65	15.0	11.5-19.2	64	9.3	7.1-11.9	75	6.7	5.2-8.4	35	8.5	5.9-11.9	40	5.7	4.0-7.8
1990	143	12.7	10.7-15.0	68	14.9	11.4-19.0	75	11.1	8.7-13.9	80	7.4	5.8-9.2	41	9.8	7.0-13.4	39	5.7	4.1-7.9
1991	149	13.5	11.4-15.9	69	16.3	12.6-20.8	80	11.9	9.4-14.8	84	7.6	6.1-9.4	49	12.2	8.9-16.2	35	5.2	3.6-7.2
1992	133	12.1	10.1-14.4	73	17.4	13.5-22.0	60	9.0	6.9-11.6	64	5.9	4.6-7.6	39	9.5	6.7-13.1	25	3.6	2.3-5.3
1993	129	11.4	9.5-13.6	57	13.1	9.8-17.0	72	10.9	8.5-13.7	76	7.2	5.6-9.0	40	10.2	7.2-13.8	36	5.5	3.9-7.7
1994	123	11.2	9.3-13.4	66	15.5	11.8-19.8	57	8.6	6.5-11.1	85	7.6	6.0-9.4	41	9.4	6.7-12.9	44	6.3	4.6-8.5
1995	100	9.1	7.4-11.1	47	10.5	7.6-14.0	53	8.2	6.2-10.8	69	6.4	5.0-8.1	33	7.9	5.3-11.1	36	5.6	3.9-7.7
1996	122	10.8	8.9-12.9	69	15.0	11.5-19.0	53	7.6	5.7-10.0	55	5.1	3.9-6.7	25	6.0	3.8-8.9	30	4.5	3.0-6.4
1997	144	12.7	10.7-14.9	75	16.6	12.9-20.9	69	10.4	8.1-13.1	66	6.2	4.8-7.9	39	9.7	6.8-13.3	27	4.1	2.7-6.0
1998	122	11.1	9.2-13.2	69	16.1	12.4-20.4	53	7.9	5.9-10.3	80	7.3	5.8-9.1	46	10.7	7.8-14.4	34	5.2	3.6-7.2
1999	141	12.3	10.3-14.5	73	16.6	12.9-21.0	68	10.0	7.7-12.6	80	7.2	5.7-9.0	43	10.2	7.3-13.8	37	5.6	3.9-7.7
2000	141	12.2	10.3-14.4	68	14.8	11.4-18.9	73	10.5	8.2-13.2	62	5.4	4.1-7.0	32	7.2	4.9-10.3	30	4.4	3.0-6.3
2001	148	12.9	10.9-15.2	73	16.1	12.5-20.3	75	10.9	8.6-13.7	73	6.4	5.0-8.1	39	8.8	6.2-12.1	34	4.9	3.4-6.8
2002	142	12.0	10.1-14.1	79	16.7	13.1-20.9	63	8.9	6.8-11.4	58	4.9	3.7-6.4	30	6.6	4.4-9.4	28	3.9	2.6-5.6
2003	171	14.2	12.1-16.5	83	17.3	13.6-21.5	88	12.5	10.0-15.4	70	5.9	4.6-7.5	30	6.8	4.5-9.7	40	5.6	4.0-7.6
2004	157	13.1	11.2-15.4	89	18.7	14.9-23.1	68	9.6	7.4-12.2	78	6.7	5.3-8.3	37	8.6	6.0-11.9	41	5.7	4.1-7.8
2005	139	11.4	9.6-13.5	70	14.2	11.0-18.1	69	9.8	7.6-12.4	75	6.3	4.9-7.9	40	8.3	5.9-11.4	35	5.0	3.5-7.0
2006	161	13.5	11.5-15.8	91	18.9	15.1-23.3	70	10.0	7.8-12.6	60	5.1	3.9-6.6	30	6.7	4.5-9.6	30	4.2	2.8-6.0
2007	146	11.9	10.0-14.0	72	13.8	10.7-17.5	74	10.3	8.1-13.0	72	6.1	4.8-7.7	38	8.2	5.7-11.3	34	4.8	3.3-6.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Bronx
Leukemias**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	139	11.1	9.3-13.1	69	13.0	10.0-16.6	70	9.8	7.6-12.4	60	4.9	3.7-6.3	23	4.4	2.8-6.7	37	5.1	3.6-7.1
2009	153	12.1	10.2-14.2	78	15.5	12.1-19.4	75	10.1	7.9-12.7	67	5.5	4.2-6.9	33	6.5	4.4-9.1	34	4.6	3.2-6.5
2010	181	14.2	12.2-16.4	107	20.5	16.7-24.9	74	9.8	7.7-12.3	58	4.6	3.5-5.9	30	6.0	4.0-8.7	28	3.7	2.5-5.4
2011	186	14.4	12.3-16.6	96	17.9	14.4-22.0	90	12.0	9.6-14.8	81	6.4	5.1-8.0	49	9.3	6.8-12.4	32	4.3	2.9-6.1
2012	169	12.9	11.0-15.0	93	17.1	13.7-21.1	76	10.1	7.9-12.7	55	4.3	3.2-5.6	24	4.6	2.9-6.8	31	3.9	2.7-5.6
2013	191	13.8	11.9-16.0	96	16.7	13.5-20.6	95	12.0	9.7-14.7	66	5.0	3.9-6.4	43	8.2	5.8-11.1	23	3.0	1.9-4.5
2014	175	12.5	10.7-14.5	95	16.2	13.0-19.9	80	10.3	8.1-12.8	75	5.3	4.2-6.7	42	7.7	5.5-10.5	33	3.9	2.7-5.5
2015	191	13.6	11.7-15.7	89	15.8	12.6-19.5	102	12.2	10.0-14.9	83	5.9	4.7-7.3	41	7.2	5.1-9.9	42	5.0	3.6-6.8
2016	175	12.1	10.4-14.1	100	16.7	13.5-20.4	75	9.0	7.1-11.4	69	4.8	3.7-6.1	32	5.8	3.9-8.2	37	4.3	3.0-5.9
2017	183	12.5	10.7-14.4	92	15.0	12.0-18.4	91	10.9	8.8-13.5	72	4.9	3.8-6.2	39	6.7	4.7-9.2	33	3.8	2.6-5.3
2018	190	12.9	11.1-14.9	102	15.8	12.8-19.3	88	10.6	8.5-13.1	64	4.4	3.4-5.7	33	5.4	3.7-7.7	31	3.7	2.5-5.2
2019	176	11.7	10.0-13.6	90	13.8	11.1-17.1	86	10.0	8.0-12.4	90	5.9	4.8-7.3	46	7.3	5.3-9.9	44	5.0	3.6-6.8
2020	147	9.7	8.2-11.5	79	12.2	9.6-15.3	68	8.2	6.3-10.4	64	4.2	3.2-5.4	45	7.3	5.3-9.8	19	2.2	1.3-3.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
All Invasive Malignant Tumors**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	8,168	371.2	362.8-379.6	3,874	430.0	415.7-444.6	4,294	337.4	327.1-348.0	4,618	213.5	207.1-220.0	2,383	268.2	256.8-279.9	2,235	177.8	170.3-185.6
1977	7,959	365.0	356.8-373.5	3,860	430.8	416.5-445.4	4,099	326.1	315.8-336.5	4,637	215.1	208.7-221.6	2,382	274.1	262.5-286.0	2,255	177.7	170.2-185.4
1978	8,273	386.7	378.1-395.4	3,967	453.5	438.7-468.6	4,306	348.8	338.2-359.8	4,565	214.7	208.3-221.2	2,344	272.1	260.6-284.0	2,221	178.4	170.8-186.2
1979	8,481	398.5	389.8-407.3	4,060	470.1	455.0-485.5	4,421	357.6	346.8-368.6	4,624	217.5	211.0-224.0	2,385	278.3	266.7-290.3	2,239	178.9	171.4-186.7
1980	8,994	422.3	413.3-431.3	4,428	512.9	497.2-528.9	4,566	369.0	358.0-380.2	4,704	222.5	216.0-229.2	2,457	291.0	279.1-303.3	2,247	179.8	172.3-187.6
1981	8,683	409.6	400.8-418.5	4,188	494.4	478.8-510.2	4,495	362.5	351.7-373.6	4,659	219.5	213.1-226.1	2,298	273.7	262.1-285.7	2,361	186.9	179.2-194.8
1982	8,657	406.3	397.6-415.2	4,145	483.3	468.1-498.9	4,512	362.4	351.7-373.5	4,423	209.5	203.2-215.9	2,258	270.8	259.2-282.7	2,165	172.5	165.1-180.1
1983	8,728	407.6	398.9-416.5	4,152	486.5	471.2-502.1	4,576	365.2	354.5-376.3	4,615	216.7	210.3-223.2	2,326	277.2	265.5-289.1	2,289	180.3	172.8-188.0
1984	9,024	418.8	410.0-427.7	4,353	503.1	487.7-518.8	4,671	371.5	360.6-382.5	4,507	211.4	205.1-217.8	2,260	267.8	256.4-279.5	2,247	177.3	169.9-184.9
1985	8,980	415.5	406.8-424.4	4,320	501.7	486.3-517.5	4,660	368.3	357.5-379.2	4,406	204.7	198.6-211.0	2,216	261.3	250.1-272.8	2,190	170.9	163.7-178.4
1986	9,103	419.6	410.9-428.5	4,432	511.9	496.4-527.7	4,671	367.2	356.5-378.1	4,475	208.1	201.9-214.4	2,202	258.4	247.4-269.8	2,273	178.1	170.7-185.8
1987	8,759	402.3	393.7-410.9	4,336	500.0	484.7-515.6	4,423	346.3	336.0-356.9	4,454	206.0	199.9-212.3	2,292	271.4	260.0-283.1	2,162	167.3	160.2-174.6
1988	8,739	404.8	396.2-413.5	4,137	484.4	469.3-499.8	4,602	361.3	350.7-372.0	4,424	206.4	200.3-212.7	2,254	269.4	258.1-281.1	2,170	168.7	161.6-176.1
1989	9,112	424.5	415.8-433.5	4,435	516.5	501.0-532.3	4,677	370.7	360.0-381.7	4,318	203.2	197.1-209.4	2,150	259.7	248.5-271.2	2,168	168.9	161.7-176.2
1990	9,215	433.5	424.6-442.6	4,387	520.5	504.9-536.6	4,828	386.5	375.5-397.8	4,348	207.1	200.9-213.4	2,158	266.3	254.9-278.0	2,190	173.0	165.8-180.5
1991	9,398	442.8	433.8-451.9	4,527	537.5	521.5-553.8	4,871	389.6	378.6-400.8	4,221	200.4	194.4-206.6	2,133	260.4	249.2-271.9	2,088	164.5	157.4-171.7
1992	9,719	456.2	447.1-465.4	4,737	555.9	539.8-572.3	4,982	399.5	388.4-410.9	4,208	199.9	193.8-206.0	2,040	248.9	238.0-260.2	2,168	171.4	164.2-178.9
1993	9,499	444.2	435.3-453.3	4,638	544.4	528.6-560.7	4,861	386.4	375.5-397.5	4,318	203.8	197.8-210.0	2,157	262.0	250.8-273.5	2,161	169.4	162.3-176.8
1994	10,349	483.6	474.3-493.1	5,046	587.6	571.2-604.3	5,303	421.6	410.3-433.2	4,339	205.1	199.0-211.3	2,164	265.6	254.3-277.2	2,175	170.4	163.2-177.7
1995	9,914	461.9	452.8-471.1	4,812	558.9	542.9-575.2	5,102	404.6	393.5-415.9	4,391	207.3	201.2-213.6	2,144	262.3	251.1-273.8	2,247	176.2	169.0-183.7
1996	10,039	466.8	457.7-476.0	4,830	560.9	544.9-577.2	5,209	411.5	400.4-422.9	4,260	200.4	194.4-206.6	2,021	245.3	234.5-256.4	2,239	174.9	167.7-182.3
1997	10,067	466.2	457.1-475.4	4,837	560.4	544.5-576.7	5,230	410.7	399.6-422.0	4,127	192.8	186.9-198.7	1,983	239.0	228.4-249.9	2,144	166.2	159.2-173.4
1998	10,353	472.5	463.4-481.7	4,923	556.7	541.0-572.7	5,430	422.0	410.8-433.4	4,087	188.4	182.7-194.3	1,957	230.2	220.0-240.8	2,130	163.4	156.5-170.5
1999	10,340	461.3	452.5-470.3	5,023	554.9	539.4-570.6	5,317	403.9	393.1-415.0	3,971	178.7	173.2-184.4	1,896	217.8	208.0-228.0	2,075	155.2	148.6-162.0
2000	10,642	465.4	456.6-474.3	5,292	572.4	556.9-588.2	5,350	398.8	388.1-409.6	3,970	174.7	169.3-180.3	1,889	212.3	202.7-222.2	2,081	151.8	145.3-158.5
2001	10,388	449.7	441.1-458.5	5,117	548.1	533.0-563.5	5,271	389.1	378.7-399.8	3,968	172.9	167.6-178.4	1,892	211.1	201.5-220.9	2,076	149.9	143.5-156.5
2002	10,243	438.6	430.1-447.1	5,000	527.4	512.7-542.4	5,243	384.1	373.8-394.7	3,812	164.2	159.0-169.5	1,849	203.3	194.0-212.9	1,963	141.0	134.9-147.5
2003	10,316	439.9	431.5-448.5	5,134	538.9	524.1-554.1	5,182	378.2	367.9-388.7	3,758	161.3	156.2-166.6	1,789	195.3	186.3-204.7	1,969	140.7	134.5-147.1
2004	10,382	439.8	431.3-448.3	5,125	530.1	515.5-545.0	5,257	382.7	372.4-393.3	3,691	157.6	152.6-162.8	1,788	195.1	186.0-204.5	1,903	135.7	129.6-142.0
2005	10,535	444.3	435.8-452.9	5,190	532.2	517.6-547.0	5,345	388.4	378.0-399.0	3,598	153.1	148.1-158.2	1,695	182.5	173.8-191.5	1,903	136.0	129.9-142.3
2006	10,681	451.8	443.2-460.5	5,246	537.3	522.6-552.2	5,435	397.7	387.1-408.5	3,446	147.2	142.3-152.2	1,670	179.6	170.9-188.5	1,776	127.4	121.5-133.6
2007	11,019	462.3	453.6-471.0	5,447	550.8	536.0-565.9	5,572	406.4	395.7-417.3	3,584	151.2	146.3-156.3	1,708	181.1	172.4-190.0	1,876	133.5	127.5-139.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
All Invasive Malignant Tumors**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	11,111	459.9	451.3-468.6	5,495	546.4	531.8-561.4	5,616	405.0	394.4-415.9	3,521	147.3	142.5-152.3	1,675	176.1	167.6-184.9	1,846	129.8	123.9-135.9
2009	11,388	467.4	458.7-476.1	5,551	545.8	531.2-560.6	5,837	417.3	406.5-428.3	3,534	146.5	141.7-151.5	1,699	176.9	168.5-185.7	1,835	127.7	121.8-133.8
2010	10,983	443.5	435.1-452.0	5,369	515.5	501.4-529.7	5,614	396.7	386.3-407.4	3,743	152.2	147.3-157.2	1,835	187.1	178.5-196.0	1,908	130.5	124.7-136.6
2011	11,663	462.0	453.5-470.6	5,703	533.3	519.3-547.7	5,960	416.1	405.4-427.0	3,858	154.9	150.0-159.9	1,826	182.9	174.4-191.6	2,032	137.4	131.4-143.6
2012	11,406	443.4	435.2-451.7	5,477	502.1	488.6-515.9	5,929	407.9	397.4-418.6	3,786	149.2	144.5-154.1	1,837	180.0	171.7-188.6	1,949	130.8	125.0-136.9
2013	11,871	452.9	444.6-461.2	5,644	505.9	492.4-519.5	6,227	421.1	410.5-431.9	3,805	146.4	141.7-151.2	1,811	172.6	164.6-181.0	1,994	129.2	123.5-135.1
2014	11,557	434.1	426.1-442.2	5,454	478.2	465.3-491.4	6,103	408.0	397.6-418.5	3,730	140.6	136.0-145.2	1,795	167.6	159.8-175.7	1,935	123.4	117.9-129.2
2015	11,790	434.2	426.3-442.3	5,536	473.1	460.4-486.0	6,254	412.5	402.1-423.1	3,608	134.1	129.7-138.6	1,735	158.0	150.5-165.8	1,873	118.4	113.0-124.0
2016	11,914	435.0	427.1-443.0	5,766	487.9	475.1-501.0	6,148	403.2	392.9-413.6	3,599	131.1	126.8-135.6	1,788	159.2	151.7-166.9	1,811	113.6	108.4-119.1
2017	12,391	446.9	438.9-455.0	5,953	495.6	482.8-508.6	6,438	417.6	407.2-428.1	3,758	135.5	131.2-140.0	1,830	161.8	154.3-169.5	1,928	118.7	113.4-124.3
2018	11,869	425.1	417.4-433.0	5,651	464.2	452.0-476.8	6,218	402.6	392.4-413.0	3,477	123.4	119.3-127.6	1,617	138.5	131.7-145.5	1,860	113.8	108.6-119.2
2019	12,330	436.7	428.9-444.7	5,832	471.2	458.9-483.7	6,498	417.2	406.9-427.8	3,335	116.8	112.8-120.9	1,607	137.0	130.3-144.0	1,728	103.9	98.9-109.0
2020	10,527	372.8	365.5-380.1	4,963	399.2	388.0-410.7	5,564	357.9	348.3-367.7	3,158	108.9	105.1-112.8	1,440	120.5	114.3-127.0	1,718	101.6	96.7-106.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Oral cavity and pharynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	204	9.0	7.7-10.3	146	15.1	12.6-17.9	58	4.5	3.4-5.8	85	3.7	3.0-4.7	59	6.2	4.7-8.1	26	2.0	1.3-2.9
1977	213	9.6	8.3-11.0	137	14.9	12.4-17.7	76	5.7	4.5-7.2	99	4.3	3.5-5.3	69	7.0	5.4-8.9	30	2.3	1.5-3.3
1978	241	11.3	9.9-12.8	166	18.2	15.5-21.4	75	6.3	4.9-8.0	89	4.1	3.2-5.0	65	7.1	5.4-9.2	24	1.8	1.1-2.7
1979	229	10.9	9.5-12.5	165	19.0	16.1-22.2	64	5.2	4.0-6.7	99	4.6	3.7-5.7	72	7.9	6.2-10.1	27	2.2	1.5-3.3
1980	220	10.6	9.2-12.2	154	17.8	15.0-21.0	66	5.6	4.3-7.2	90	4.3	3.4-5.3	72	8.2	6.3-10.4	18	1.6	0.9-2.5
1981	241	11.5	10.0-13.1	172	19.5	16.6-22.7	69	5.6	4.3-7.2	86	4.1	3.3-5.1	60	7.0	5.3-9.1	26	2.1	1.3-3.0
1982	238	11.4	10.0-13.0	173	19.5	16.6-22.8	65	5.7	4.4-7.3	98	4.9	4.0-6.1	75	8.9	6.9-11.2	23	2.1	1.3-3.2
1983	232	11.3	9.9-12.9	155	18.1	15.3-21.3	77	6.4	5.0-8.0	97	4.7	3.8-5.8	75	8.7	6.8-11.0	22	1.9	1.2-2.9
1984	261	12.5	11.0-14.2	179	20.5	17.5-23.8	82	6.7	5.3-8.3	85	4.0	3.2-5.0	63	7.2	5.5-9.3	22	1.7	1.1-2.6
1985	242	11.7	10.2-13.3	160	18.3	15.5-21.4	82	6.7	5.3-8.4	80	3.8	3.0-4.8	53	6.1	4.5-8.1	27	2.1	1.4-3.1
1986	232	10.7	9.3-12.2	155	17.2	14.6-20.2	77	5.9	4.6-7.4	87	4.2	3.3-5.2	58	6.7	5.0-8.7	29	2.3	1.5-3.3
1987	246	11.8	10.3-13.4	176	20.2	17.2-23.4	70	5.7	4.4-7.3	68	3.3	2.6-4.2	53	6.2	4.6-8.1	15	1.2	0.7-2.0
1988	226	10.7	9.4-12.3	148	17.0	14.3-20.0	78	6.2	4.9-7.8	80	3.8	3.0-4.8	59	7.0	5.3-9.0	21	1.7	1.1-2.6
1989	224	10.8	9.4-12.3	150	17.3	14.6-20.3	74	6.0	4.7-7.6	82	4.1	3.3-5.1	57	6.8	5.1-8.8	25	2.1	1.4-3.1
1990	224	10.7	9.3-12.2	145	16.7	14.1-19.7	79	6.3	4.9-7.9	61	3.0	2.3-3.8	44	5.2	3.7-7.0	17	1.3	0.8-2.1
1991	235	11.4	9.9-12.9	159	18.5	15.7-21.7	76	6.1	4.8-7.6	92	4.5	3.6-5.5	59	7.1	5.4-9.2	33	2.6	1.8-3.6
1992	284	13.7	12.1-15.4	189	21.5	18.5-24.9	95	7.9	6.3-9.6	70	3.4	2.6-4.2	48	5.6	4.1-7.4	22	1.7	1.1-2.6
1993	231	11.0	9.6-12.6	155	17.9	15.1-20.9	76	6.2	4.9-7.8	79	3.8	3.0-4.7	58	6.5	4.9-8.4	21	1.7	1.0-2.6
1994	233	11.1	9.7-12.6	157	17.5	14.9-20.5	76	6.2	4.9-7.7	78	3.7	2.9-4.7	61	7.2	5.5-9.3	17	1.3	0.8-2.2
1995	238	11.2	9.8-12.7	176	19.4	16.6-22.5	62	5.1	3.9-6.5	84	4.0	3.2-4.9	60	6.8	5.2-8.8	24	1.9	1.2-2.8
1996	221	10.3	9.0-11.7	148	16.5	13.9-19.4	73	5.8	4.5-7.3	76	3.6	2.8-4.5	50	5.5	4.1-7.3	26	2.1	1.4-3.1
1997	222	10.4	9.0-11.8	140	15.4	12.9-18.2	82	6.6	5.2-8.2	59	2.8	2.1-3.6	47	5.4	4.0-7.2	12	0.9	0.5-1.6
1998	247	11.3	9.9-12.8	165	18.0	15.3-21.0	82	6.4	5.1-8.0	66	3.0	2.4-3.9	45	5.1	3.7-6.8	21	1.7	1.0-2.5
1999	231	10.3	9.0-11.7	159	16.8	14.2-19.6	72	5.5	4.3-6.9	59	2.6	2.0-3.4	42	4.4	3.2-6.0	17	1.3	0.8-2.1
2000	238	10.4	9.1-11.8	161	16.4	14.0-19.2	77	5.8	4.5-7.2	83	3.6	2.9-4.5	55	5.7	4.3-7.4	28	2.1	1.4-3.0
2001	200	8.6	7.5-9.9	127	12.7	10.6-15.2	73	5.5	4.3-6.9	64	2.8	2.1-3.5	43	4.3	3.1-5.8	21	1.5	1.0-2.4
2002	219	9.3	8.1-10.7	141	14.2	11.9-16.7	78	5.7	4.5-7.1	65	2.8	2.1-3.5	50	5.0	3.7-6.5	15	1.1	0.6-1.8
2003	225	9.5	8.3-10.8	145	14.1	11.9-16.6	80	5.9	4.7-7.4	58	2.5	1.9-3.2	37	3.6	2.6-5.0	21	1.6	1.0-2.4
2004	222	9.3	8.2-10.7	139	13.6	11.4-16.1	83	6.0	4.8-7.5	66	2.8	2.1-3.5	48	4.7	3.5-6.3	18	1.3	0.8-2.1
2005	217	9.0	7.8-10.3	148	14.2	11.9-16.7	69	5.1	3.9-6.4	66	2.7	2.1-3.5	43	4.2	3.0-5.6	23	1.7	1.1-2.5
2006	228	9.6	8.4-10.9	147	14.5	12.2-17.0	81	5.9	4.6-7.3	52	2.2	1.6-2.9	39	3.8	2.7-5.2	13	1.0	0.5-1.6
2007	230	9.5	8.3-10.8	137	13.0	10.9-15.4	93	6.7	5.4-8.2	69	2.8	2.2-3.6	52	4.9	3.7-6.5	17	1.2	0.7-2.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Oral cavity and pharynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	216	8.9	7.7-10.2	140	13.2	11.0-15.6	76	5.6	4.4-7.0	51	2.2	1.6-2.8	32	3.3	2.2-4.6	19	1.4	0.8-2.1
2009	226	9.2	8.0-10.4	154	14.3	12.1-16.8	72	5.1	4.0-6.5	50	2.0	1.4-2.6	35	3.2	2.2-4.5	15	1.0	0.6-1.7
2010	238	9.5	8.3-10.8	158	14.4	12.2-16.9	80	5.6	4.4-7.0	71	2.9	2.3-3.7	47	4.5	3.3-6.0	24	1.7	1.1-2.5
2011	228	9.1	7.9-10.4	161	15.1	12.8-17.6	67	4.6	3.5-5.8	60	2.3	1.8-3.0	42	3.9	2.8-5.3	18	1.2	0.7-1.9
2012	242	9.2	8.1-10.5	162	14.3	12.1-16.7	80	5.4	4.2-6.7	64	2.5	1.9-3.2	40	3.8	2.7-5.2	24	1.5	1.0-2.3
2013	241	9.3	8.2-10.6	157	13.8	11.7-16.2	84	5.7	4.6-7.1	65	2.4	1.8-3.1	51	4.3	3.2-5.6	14	0.9	0.5-1.6
2014	258	9.6	8.5-10.9	162	13.6	11.5-15.9	96	6.5	5.2-8.0	57	2.1	1.6-2.7	37	3.1	2.2-4.4	20	1.3	0.8-2.0
2015	235	8.7	7.6-9.9	155	13.0	11.0-15.2	80	5.4	4.2-6.7	50	1.9	1.4-2.5	38	3.2	2.3-4.4	12	0.8	0.4-1.4
2016	258	9.4	8.2-10.6	168	14.1	12.0-16.5	90	5.7	4.6-7.1	50	1.8	1.4-2.4	35	3.1	2.2-4.4	15	0.9	0.5-1.5
2017	284	10.3	9.1-11.6	184	15.4	13.2-17.8	100	6.3	5.1-7.7	54	2.0	1.5-2.6	37	3.2	2.2-4.4	17	1.0	0.6-1.7
2018	246	8.9	7.8-10.1	165	13.5	11.5-15.8	81	5.2	4.1-6.5	53	2.0	1.5-2.6	40	3.5	2.5-4.8	13	0.8	0.4-1.5
2019	268	9.4	8.3-10.7	165	13.3	11.3-15.5	103	6.4	5.2-7.8	53	1.9	1.4-2.4	37	3.1	2.1-4.3	16	0.9	0.5-1.5
2020	218	7.7	6.7-8.8	143	11.7	9.8-13.8	75	4.6	3.6-5.8	59	2.1	1.6-2.7	38	3.1	2.2-4.3	21	1.3	0.8-2.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Esophagus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	111	4.8	3.9-5.9	76	7.9	6.1-10.0	35	2.6	1.8-3.7	123	5.6	4.6-6.7	94	9.8	7.8-12.0	29	2.5	1.6-3.6
1977	110	5.1	4.2-6.2	78	8.5	6.6-10.8	32	2.7	1.8-3.9	109	5.1	4.1-6.2	74	8.6	6.7-11.0	35	2.7	1.9-3.8
1978	145	6.7	5.6-7.9	100	11.1	8.9-13.6	45	3.6	2.6-4.9	112	5.2	4.2-6.3	74	8.3	6.4-10.5	38	3.0	2.1-4.2
1979	138	6.3	5.3-7.5	105	11.5	9.3-14.0	33	2.6	1.8-3.7	124	5.8	4.8-6.9	95	10.4	8.4-12.9	29	2.5	1.6-3.6
1980	152	7.2	6.1-8.5	106	12.1	9.8-14.8	46	3.8	2.7-5.0	125	5.9	4.9-7.1	88	10.2	8.1-12.6	37	3.0	2.1-4.1
1981	153	7.3	6.2-8.6	96	11.0	8.9-13.5	57	4.8	3.6-6.2	136	6.3	5.3-7.5	95	10.7	8.6-13.2	41	3.2	2.3-4.4
1982	135	6.0	5.0-7.2	93	10.1	8.1-12.5	42	3.1	2.3-4.3	115	5.4	4.4-6.5	79	9.3	7.3-11.7	36	2.8	1.9-3.9
1983	108	4.9	4.0-5.9	78	8.4	6.6-10.5	30	2.3	1.5-3.3	112	5.0	4.1-6.1	79	8.5	6.7-10.7	33	2.6	1.8-3.7
1984	144	6.5	5.5-7.7	98	10.7	8.6-13.1	46	3.5	2.6-4.7	111	5.1	4.2-6.2	75	8.1	6.3-10.1	36	2.9	2.0-4.1
1985	124	5.9	4.9-7.0	83	9.6	7.6-12.0	41	3.4	2.4-4.7	138	6.5	5.4-7.7	99	11.5	9.3-14.1	39	3.1	2.2-4.3
1986	142	6.6	5.5-7.8	102	11.8	9.6-14.4	40	3.1	2.2-4.3	110	5.2	4.2-6.3	84	9.9	7.9-12.4	26	2.1	1.3-3.1
1987	126	5.8	4.8-6.9	85	9.4	7.5-11.7	41	3.2	2.3-4.4	110	5.1	4.1-6.1	79	8.9	7.0-11.1	31	2.5	1.7-3.5
1988	109	5.2	4.2-6.3	80	9.4	7.4-11.8	29	2.2	1.5-3.2	116	5.6	4.6-6.7	88	10.5	8.3-12.9	28	2.2	1.4-3.2
1989	139	6.6	5.5-7.8	94	10.7	8.6-13.1	45	3.4	2.5-4.6	97	4.6	3.7-5.7	64	7.3	5.6-9.4	33	2.6	1.8-3.7
1990	115	5.6	4.6-6.8	71	8.4	6.6-10.7	44	3.7	2.6-4.9	93	4.4	3.6-5.4	63	7.6	5.8-9.7	30	2.4	1.6-3.4
1991	113	5.4	4.4-6.4	77	8.9	7.0-11.2	36	2.9	2.0-4.0	90	4.2	3.4-5.2	58	6.6	5.0-8.5	32	2.4	1.7-3.5
1992	116	5.6	4.6-6.7	72	8.5	6.6-10.7	44	3.5	2.6-4.8	76	3.7	2.9-4.6	50	5.8	4.3-7.6	26	2.2	1.4-3.2
1993	132	6.4	5.3-7.6	88	10.5	8.4-12.9	44	3.5	2.5-4.7	110	5.2	4.3-6.3	81	9.6	7.6-11.9	29	2.2	1.5-3.2
1994	133	6.4	5.3-7.5	90	10.6	8.5-13.1	43	3.4	2.5-4.6	102	4.8	3.9-5.9	67	7.7	5.9-9.8	35	2.8	1.9-3.9
1995	118	5.6	4.6-6.7	78	9.0	7.1-11.3	40	3.2	2.3-4.4	107	5.1	4.1-6.1	76	8.8	6.9-11.0	31	2.5	1.7-3.5
1996	112	5.3	4.3-6.3	68	7.9	6.1-10.0	44	3.4	2.5-4.5	85	4.0	3.2-5.0	59	6.8	5.2-8.8	26	2.0	1.3-3.0
1997	106	5.0	4.1-6.0	69	7.8	6.1-9.9	37	2.8	2.0-3.9	93	4.3	3.5-5.3	64	7.4	5.7-9.4	29	2.2	1.5-3.2
1998	104	4.8	3.9-5.8	73	8.2	6.4-10.3	31	2.3	1.6-3.3	92	4.2	3.4-5.2	59	6.6	5.0-8.5	33	2.5	1.7-3.5
1999	115	5.1	4.2-6.2	87	9.4	7.5-11.6	28	2.1	1.4-3.0	92	4.1	3.3-5.1	72	7.9	6.2-10.0	20	1.5	0.9-2.3
2000	98	4.3	3.5-5.3	54	5.8	4.4-7.6	44	3.2	2.3-4.3	87	3.8	3.1-4.7	54	5.8	4.4-7.6	33	2.4	1.7-3.4
2001	108	4.7	3.8-5.7	82	8.5	6.8-10.6	26	1.9	1.3-2.8	87	3.8	3.0-4.7	69	7.3	5.7-9.3	18	1.3	0.8-2.1
2002	91	3.9	3.1-4.8	63	6.7	5.1-8.5	28	2.0	1.4-2.9	68	2.9	2.3-3.7	39	4.2	3.0-5.7	29	2.1	1.4-3.0
2003	108	4.6	3.8-5.6	68	7.2	5.6-9.2	40	2.8	2.0-3.9	81	3.5	2.8-4.3	56	5.8	4.4-7.5	25	1.8	1.1-2.6
2004	90	3.8	3.0-4.6	61	6.3	4.8-8.1	29	2.0	1.4-2.9	70	3.0	2.3-3.8	43	4.6	3.3-6.1	27	1.9	1.3-2.8
2005	106	4.5	3.7-5.5	72	7.2	5.6-9.1	34	2.4	1.7-3.4	63	2.6	2.0-3.3	47	4.6	3.4-6.2	16	1.1	0.6-1.7
2006	84	3.6	2.9-4.5	63	6.7	5.1-8.6	21	1.5	0.9-2.4	83	3.5	2.8-4.4	51	5.3	3.9-6.9	32	2.3	1.6-3.2
2007	99	4.1	3.4-5.0	62	6.2	4.7-7.9	37	2.6	1.8-3.6	74	3.1	2.4-3.9	57	5.7	4.3-7.4	17	1.2	0.7-1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Kings (Brooklyn)

Esophagus

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	103	4.2	3.4-5.1	70	7.0	5.4-8.8	33	2.3	1.5-3.2	79	3.3	2.6-4.1	52	5.3	4.0-7.0	27	1.9	1.2-2.8
2009	97	4.0	3.2-4.9	72	7.0	5.4-8.9	25	1.7	1.1-2.6	77	3.1	2.4-3.9	63	6.1	4.7-7.8	14	0.9	0.5-1.5
2010	73	2.9	2.3-3.7	49	4.9	3.6-6.5	24	1.6	1.0-2.4	66	2.6	2.0-3.3	46	4.6	3.3-6.1	20	1.3	0.8-2.1
2011	90	3.6	2.9-4.4	69	6.7	5.2-8.5	21	1.5	0.9-2.2	64	2.6	2.0-3.3	47	4.5	3.3-6.1	17	1.2	0.7-1.9
2012	77	2.9	2.3-3.7	55	5.0	3.7-6.5	22	1.4	0.9-2.2	55	2.2	1.6-2.8	39	3.8	2.7-5.2	16	1.0	0.5-1.6
2013	105	4.1	3.3-4.9	69	6.3	4.9-8.0	36	2.4	1.6-3.3	66	2.5	1.9-3.2	48	4.3	3.2-5.8	18	1.2	0.7-1.9
2014	88	3.4	2.7-4.2	66	6.2	4.8-7.9	22	1.4	0.9-2.1	68	2.7	2.1-3.4	43	4.3	3.1-5.8	25	1.6	1.0-2.4
2015	95	3.5	2.8-4.3	61	5.3	4.0-6.8	34	2.2	1.5-3.1	54	2.1	1.5-2.7	41	3.8	2.7-5.1	13	0.8	0.4-1.4
2016	113	4.0	3.3-4.8	83	7.0	5.5-8.7	30	1.8	1.2-2.6	65	2.3	1.8-3.0	48	4.1	3.0-5.5	17	1.1	0.6-1.7
2017	74	2.7	2.1-3.5	56	4.9	3.6-6.3	18	1.2	0.7-1.9	69	2.5	1.9-3.1	56	4.7	3.5-6.1	13	0.8	0.4-1.5
2018	72	2.5	2.0-3.2	49	4.1	3.0-5.4	23	1.4	0.9-2.1	45	1.5	1.1-2.0	36	2.8	2.0-4.0	9	0.5	0.2-1.0
2019	83	2.7	2.2-3.4	58	4.4	3.3-5.7	25	1.4	0.9-2.2	48	1.6	1.2-2.2	38	3.2	2.3-4.4	10	0.5	0.2-1.0
2020	77	2.7	2.1-3.3	49	3.9	2.9-5.2	28	1.7	1.1-2.5	48	1.6	1.2-2.2	34	2.6	1.8-3.6	14	0.9	0.5-1.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Stomach**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	350	16.9	15.1-18.9	212	25.2	21.7-29.1	138	11.3	9.4-13.4	265	13.0	11.4-14.7	151	18.5	15.5-21.9	114	9.2	7.6-11.2
1977	325	16.0	14.3-17.9	184	22.1	18.8-25.7	141	11.9	10.0-14.1	250	12.5	11.0-14.3	144	18.2	15.1-21.6	106	8.8	7.2-10.8
1978	313	15.2	13.5-17.1	163	19.7	16.6-23.2	150	12.3	10.4-14.5	231	11.2	9.7-12.8	132	15.9	13.2-19.1	99	8.1	6.5-9.9
1979	335	15.7	14.0-17.6	188	21.4	18.3-24.8	147	11.8	9.9-13.9	235	11.1	9.7-12.7	120	14.1	11.5-17.0	115	9.1	7.5-11.0
1980	350	17.1	15.3-19.1	195	23.9	20.6-27.7	155	12.7	10.7-14.9	227	10.8	9.4-12.4	135	16.0	13.3-19.0	92	7.4	5.9-9.1
1981	329	15.4	13.8-17.2	194	23.1	19.8-26.7	135	10.7	8.9-12.7	234	11.2	9.7-12.7	133	15.7	13.1-18.8	101	8.3	6.7-10.1
1982	338	15.7	14.1-17.6	194	22.8	19.6-26.4	144	11.2	9.4-13.3	203	9.6	8.3-11.1	118	14.4	11.8-17.3	85	6.6	5.2-8.2
1983	358	17.1	15.4-19.1	191	22.8	19.5-26.4	167	13.6	11.5-15.8	249	11.8	10.4-13.4	137	16.6	13.9-19.8	112	8.9	7.3-10.7
1984	319	14.8	13.2-16.5	174	20.6	17.5-24.0	145	11.2	9.4-13.2	195	9.4	8.1-10.8	104	12.7	10.3-15.4	91	7.4	5.9-9.1
1985	318	14.9	13.3-16.7	174	20.9	17.8-24.4	144	11.1	9.3-13.1	213	10.1	8.7-11.5	119	14.1	11.6-17.0	94	7.4	5.9-9.1
1986	322	15.4	13.7-17.2	179	21.9	18.7-25.5	143	11.2	9.4-13.2	197	9.2	8.0-10.6	117	14.0	11.5-16.9	80	6.2	4.9-7.7
1987	313	14.6	13.0-16.4	187	22.2	19.1-25.8	126	9.6	8.0-11.4	209	9.8	8.5-11.2	128	15.8	13.1-18.8	81	6.1	4.8-7.6
1988	306	14.4	12.8-16.1	164	19.7	16.7-23.0	142	11.1	9.3-13.1	220	10.2	8.9-11.7	125	14.8	12.3-17.7	95	7.3	5.9-8.9
1989	303	14.3	12.7-16.0	170	20.3	17.3-23.6	133	10.1	8.5-12.0	212	9.9	8.6-11.4	117	14.4	11.8-17.3	95	7.1	5.7-8.7
1990	315	15.0	13.4-16.8	172	20.7	17.6-24.1	143	11.2	9.4-13.2	188	9.0	7.8-10.4	112	14.0	11.5-16.9	76	5.9	4.6-7.4
1991	313	15.0	13.4-16.8	177	21.7	18.6-25.2	136	10.7	9.0-12.7	200	9.7	8.4-11.1	113	14.3	11.7-17.2	87	6.8	5.4-8.4
1992	308	14.6	13.0-16.4	180	22.0	18.8-25.5	128	10.0	8.3-11.9	199	9.6	8.3-11.1	109	13.7	11.2-16.6	90	7.1	5.7-8.8
1993	319	15.2	13.6-17.0	177	22.1	18.9-25.7	142	11.0	9.2-12.9	197	9.4	8.1-10.8	105	13.0	10.6-15.8	92	7.1	5.7-8.7
1994	338	16.0	14.4-17.8	181	22.3	19.1-25.9	157	12.0	10.2-14.0	227	10.9	9.5-12.4	135	17.5	14.6-20.7	92	7.0	5.7-8.6
1995	304	14.3	12.7-16.0	171	20.5	17.5-23.8	133	10.2	8.5-12.1	205	9.7	8.4-11.1	117	14.2	11.7-17.1	88	6.8	5.4-8.3
1996	321	15.2	13.6-16.9	185	22.6	19.4-26.2	136	10.6	8.9-12.5	194	9.2	7.9-10.6	104	12.8	10.4-15.5	90	7.0	5.6-8.6
1997	294	13.7	12.2-15.4	176	21.4	18.3-24.9	118	9.0	7.5-10.8	190	8.9	7.6-10.2	104	12.4	10.1-15.1	86	6.6	5.3-8.1
1998	321	14.9	13.3-16.6	178	20.9	17.9-24.3	143	10.9	9.2-12.8	188	8.7	7.5-10.1	110	13.4	11.0-16.1	78	5.8	4.6-7.3
1999	318	14.4	12.8-16.0	192	22.1	19.0-25.5	126	9.3	7.7-11.1	156	7.0	6.0-8.2	84	10.1	8.0-12.5	72	5.3	4.2-6.7
2000	293	12.9	11.4-14.4	145	16.1	13.6-19.0	148	10.7	9.0-12.6	186	8.2	7.0-9.4	104	11.6	9.5-14.1	82	6.0	4.8-7.4
2001	313	13.7	12.2-15.3	156	17.0	14.4-19.9	157	11.2	9.5-13.1	178	7.8	6.7-9.0	87	9.5	7.6-11.7	91	6.5	5.2-7.9
2002	289	12.4	11.0-13.9	166	17.8	15.2-20.8	123	8.7	7.2-10.4	149	6.4	5.4-7.5	80	8.7	6.9-10.9	69	4.9	3.8-6.2
2003	309	13.2	11.8-14.7	156	16.7	14.2-19.6	153	10.8	9.1-12.6	163	7.0	6.0-8.2	80	8.7	6.8-10.8	83	5.9	4.7-7.3
2004	318	13.7	12.2-15.2	170	18.5	15.8-21.5	148	10.5	8.9-12.4	166	7.2	6.1-8.3	90	10.0	8.1-12.4	76	5.4	4.3-6.8
2005	354	15.1	13.5-16.7	205	22.0	19.1-25.3	149	10.6	8.9-12.4	151	6.4	5.4-7.5	85	9.4	7.5-11.7	66	4.7	3.6-5.9
2006	329	14.0	12.5-15.6	196	20.7	17.9-23.9	133	9.6	8.0-11.4	168	7.2	6.1-8.3	103	11.4	9.3-13.9	65	4.5	3.5-5.8
2007	304	12.9	11.5-14.4	161	17.1	14.5-19.9	143	10.2	8.6-12.1	130	5.5	4.6-6.5	74	7.9	6.2-10.0	56	3.9	3.0-5.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Stomach**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	271	11.3	10.0-12.8	154	16.2	13.7-19.0	117	8.2	6.8-9.8	144	6.1	5.1-7.1	91	9.5	7.6-11.7	53	3.7	2.8-4.9
2009	294	12.3	10.9-13.8	168	17.3	14.7-20.1	126	8.8	7.3-10.5	131	5.6	4.6-6.6	67	7.1	5.5-9.1	64	4.4	3.4-5.6
2010	295	12.3	10.9-13.8	166	17.0	14.5-19.8	129	9.0	7.5-10.7	141	5.7	4.8-6.7	79	8.2	6.5-10.3	62	4.1	3.2-5.3
2011	310	12.3	11.0-13.8	163	15.7	13.3-18.4	147	9.9	8.3-11.7	136	5.6	4.7-6.6	77	7.9	6.2-9.9	59	3.9	3.0-5.1
2012	308	12.4	11.0-13.9	172	16.9	14.4-19.7	136	9.3	7.8-11.0	142	5.6	4.7-6.6	86	8.6	6.8-10.7	56	3.7	2.8-4.8
2013	326	12.6	11.2-14.0	198	18.1	15.6-20.9	128	8.5	7.1-10.1	143	5.4	4.5-6.4	86	8.0	6.4-9.9	57	3.6	2.7-4.7
2014	278	10.6	9.4-12.0	158	14.8	12.5-17.3	120	7.7	6.4-9.3	167	6.5	5.5-7.6	93	8.6	6.9-10.6	74	4.9	3.8-6.1
2015	331	12.4	11.1-13.8	192	17.1	14.7-19.8	139	8.8	7.4-10.5	129	4.9	4.1-5.9	72	6.8	5.3-8.6	57	3.5	2.6-4.6
2016	340	12.5	11.2-13.9	203	17.6	15.2-20.2	137	8.8	7.4-10.5	191	7.1	6.1-8.2	124	11.2	9.3-13.4	67	4.1	3.1-5.2
2017	313	11.4	10.1-12.7	178	15.3	13.1-17.7	135	8.6	7.2-10.2	143	5.4	4.5-6.3	82	7.6	6.0-9.4	61	3.9	3.0-5.1
2018	291	10.3	9.1-11.6	169	13.9	11.9-16.2	122	7.6	6.3-9.2	111	3.9	3.2-4.8	60	5.1	3.9-6.6	51	3.0	2.2-4.0
2019	304	10.7	9.5-11.9	180	14.6	12.5-17.0	124	7.8	6.4-9.3	142	5.1	4.3-6.0	78	6.6	5.2-8.3	64	4.0	3.0-5.1
2020	261	9.0	7.9-10.2	145	11.9	10.0-14.1	116	6.8	5.6-8.2	120	4.2	3.5-5.0	74	6.1	4.8-7.7	46	2.7	2.0-3.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Colorectal**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,328	61.5	58.1-65.0	639	73.6	67.7-80.0	689	53.8	49.7-58.1	711	34.2	31.6-37.0	371	44.4	39.7-49.4	340	27.7	24.7-30.9
1977	1,359	64.0	60.5-67.6	676	79.5	73.2-86.1	683	54.4	50.2-58.7	721	34.5	31.9-37.2	337	42.2	37.5-47.3	384	30.1	27.1-33.4
1978	1,356	63.2	59.7-66.8	652	76.3	70.2-82.7	704	55.1	51.0-59.4	695	33.9	31.3-36.6	354	44.1	39.4-49.3	341	27.7	24.7-30.9
1979	1,418	67.5	63.9-71.3	675	81.2	74.9-87.9	743	58.8	54.6-63.4	709	33.9	31.4-36.6	327	39.1	34.7-43.8	382	30.5	27.5-33.8
1980	1,504	70.2	66.6-73.9	763	89.2	82.7-96.0	741	58.2	54.0-62.6	709	34.8	32.2-37.6	379	48.3	43.3-53.7	330	26.6	23.7-29.7
1981	1,476	70.6	66.9-74.4	734	90.4	83.7-97.5	742	58.6	54.4-63.1	662	31.9	29.5-34.5	283	36.2	31.9-40.9	379	29.7	26.7-32.9
1982	1,400	66.2	62.7-69.9	672	81.6	75.3-88.2	728	56.6	52.5-60.9	638	30.7	28.3-33.3	322	40.5	36.0-45.4	316	25.0	22.3-28.0
1983	1,388	65.1	61.7-68.7	663	80.3	74.0-86.9	725	56.3	52.2-60.7	619	29.8	27.5-32.3	309	38.5	34.1-43.2	310	24.6	21.9-27.5
1984	1,373	63.9	60.5-67.4	667	78.9	72.8-85.3	706	54.4	50.4-58.6	627	29.9	27.6-32.4	298	37.4	33.2-42.1	329	25.7	22.9-28.7
1985	1,421	66.0	62.6-69.6	688	82.5	76.2-89.1	733	56.4	52.3-60.7	598	27.8	25.6-30.2	276	33.0	29.1-37.3	322	24.6	22.0-27.5
1986	1,415	65.7	62.2-69.3	695	82.5	76.2-89.0	720	55.4	51.4-59.7	586	27.5	25.3-29.9	285	34.6	30.6-39.0	301	23.5	20.9-26.4
1987	1,279	58.9	55.6-62.2	665	79.2	73.1-85.6	614	46.5	42.8-50.3	613	28.8	26.6-31.2	290	35.7	31.5-40.2	323	24.9	22.2-27.8
1988	1,282	59.6	56.3-63.0	619	74.8	68.9-81.1	663	50.2	46.4-54.2	547	25.9	23.8-28.2	294	36.6	32.4-41.2	253	19.5	17.1-22.1
1989	1,277	59.8	56.5-63.2	634	75.4	69.5-81.6	643	49.8	45.9-53.8	542	25.7	23.6-28.0	259	32.6	28.6-36.9	283	21.6	19.1-24.3
1990	1,225	58.1	54.8-61.4	561	68.4	62.7-74.4	664	51.4	47.5-55.5	556	26.8	24.6-29.1	262	32.9	29.0-37.3	294	23.0	20.4-25.8
1991	1,272	60.4	57.1-63.9	628	78.2	72.0-84.7	644	49.8	46.0-53.9	506	24.2	22.2-26.5	256	32.5	28.5-36.8	250	19.3	17.0-21.9
1992	1,260	59.6	56.3-63.0	561	68.8	63.1-74.8	699	54.2	50.2-58.4	509	24.4	22.3-26.6	233	29.1	25.4-33.2	276	21.4	18.9-24.0
1993	1,268	60.0	56.7-63.4	610	74.8	68.9-81.1	658	51.1	47.3-55.2	523	24.9	22.8-27.1	245	30.4	26.6-34.5	278	21.5	19.0-24.2
1994	1,430	67.6	64.1-71.2	694	84.3	78.0-90.9	736	57.0	52.9-61.3	539	25.7	23.5-27.9	264	33.9	29.8-38.3	275	21.1	18.6-23.7
1995	1,299	61.1	57.9-64.6	600	73.3	67.5-79.5	699	53.8	49.9-57.9	572	27.1	25.0-29.5	282	35.7	31.6-40.2	290	22.2	19.7-24.9
1996	1,309	61.7	58.4-65.2	647	78.3	72.3-84.7	662	51.2	47.4-55.3	498	23.5	21.5-25.7	231	28.5	24.9-32.5	267	20.5	18.1-23.1
1997	1,306	61.2	57.9-64.6	628	75.7	69.8-81.9	678	52.1	48.2-56.2	507	23.9	21.8-26.0	258	32.0	28.1-36.1	249	18.9	16.6-21.4
1998	1,459	67.4	64.0-70.9	699	82.8	76.7-89.3	760	57.6	53.6-61.9	503	23.4	21.4-25.5	234	28.3	24.8-32.3	269	20.2	17.9-22.8
1999	1,408	63.4	60.1-66.8	635	72.5	66.9-78.4	773	57.5	53.5-61.7	497	22.5	20.6-24.6	231	27.0	23.6-30.8	266	19.6	17.3-22.1
2000	1,321	58.3	55.1-61.5	636	70.8	65.4-76.6	685	49.6	46.0-53.5	489	21.6	19.7-23.6	230	26.1	22.8-29.7	259	18.4	16.2-20.7
2001	1,404	61.2	58.0-64.5	653	72.9	67.3-78.7	751	54.2	50.3-58.2	483	21.1	19.3-23.1	232	26.4	23.1-30.1	251	17.9	15.7-20.2
2002	1,318	56.9	53.9-60.0	639	69.7	64.3-75.4	679	48.4	44.8-52.2	449	19.4	17.7-21.3	213	24.0	20.9-27.5	236	16.7	14.6-19.0
2003	1,323	56.8	53.8-59.9	670	72.2	66.8-77.9	653	46.6	43.1-50.4	519	22.3	20.4-24.3	243	26.9	23.6-30.5	276	19.4	17.1-21.8
2004	1,290	55.1	52.1-58.2	633	67.5	62.3-73.1	657	47.0	43.5-50.8	456	19.7	17.9-21.6	232	25.9	22.6-29.5	224	15.8	13.8-18.0
2005	1,297	55.1	52.2-58.2	620	65.5	60.4-70.9	677	48.0	44.5-51.8	440	18.8	17.1-20.7	214	23.5	20.5-26.9	226	15.8	13.8-18.0
2006	1,253	53.3	50.4-56.4	579	61.0	56.1-66.3	674	48.3	44.7-52.1	428	18.3	16.6-20.2	208	22.8	19.8-26.2	220	15.5	13.5-17.7
2007	1,280	54.0	51.0-57.0	619	63.8	58.8-69.1	661	47.3	43.8-51.1	406	17.3	15.7-19.1	193	20.6	17.8-23.8	213	15.0	13.0-17.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Colorectal**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	1,172	49.0	46.3-52.0	551	56.3	51.6-61.3	621	44.4	40.9-48.0	431	18.2	16.5-20.0	204	21.6	18.7-24.8	227	16.0	14.0-18.3
2009	1,216	50.3	47.5-53.2	603	60.7	55.9-65.8	613	43.0	39.6-46.6	408	16.9	15.3-18.7	204	21.7	18.8-24.9	204	13.9	12.0-15.9
2010	1,132	46.2	43.5-49.0	571	56.6	51.9-61.5	561	38.9	35.7-42.3	415	16.9	15.3-18.6	234	24.0	21.0-27.3	181	12.1	10.4-14.0
2011	1,109	44.6	42.0-47.3	553	53.5	49.0-58.2	556	38.2	35.1-41.6	419	17.0	15.3-18.7	199	20.2	17.4-23.2	220	14.7	12.8-16.8
2012	1,064	41.1	38.7-43.7	531	48.7	44.5-53.1	533	36.0	33.0-39.2	411	16.1	14.6-17.8	192	18.8	16.2-21.7	219	14.3	12.5-16.4
2013	1,127	43.4	40.9-46.1	587	53.9	49.5-58.5	540	35.9	32.9-39.1	371	14.1	12.7-15.7	177	16.5	14.1-19.2	194	12.5	10.7-14.4
2014	1,078	40.9	38.5-43.5	529	47.9	43.8-52.2	549	36.1	33.1-39.3	360	13.4	12.0-14.8	166	15.4	13.1-18.0	194	11.8	10.2-13.7
2015	1,066	39.2	36.8-41.6	520	45.3	41.4-49.4	546	35.0	32.1-38.1	325	11.8	10.5-13.1	153	13.7	11.5-16.1	172	10.4	8.9-12.1
2016	1,082	39.8	37.4-42.3	570	49.6	45.5-53.9	512	32.8	29.9-35.8	364	13.1	11.8-14.5	175	15.6	13.4-18.2	189	11.4	9.8-13.2
2017	1,116	40.4	38.0-42.9	546	46.0	42.1-50.1	570	36.4	33.4-39.5	385	13.8	12.4-15.3	181	15.9	13.6-18.4	204	12.3	10.7-14.2
2018	1,051	37.7	35.4-40.1	541	45.6	41.7-49.7	510	31.9	29.1-34.8	356	12.9	11.6-14.3	186	16.1	13.9-18.7	170	10.6	9.0-12.3
2019	1,096	39.2	36.8-41.6	597	49.2	45.3-53.4	499	31.6	28.8-34.6	321	11.3	10.1-12.6	161	13.7	11.7-16.1	160	9.5	8.1-11.2
2020	891	31.5	29.5-33.7	438	35.4	32.1-38.9	453	28.4	25.7-31.2	324	11.0	9.8-12.3	157	13.0	11.0-15.2	167	9.5	8.1-11.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Colon excluding rectum**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	950	44.0	41.1-47.0	439	50.7	45.8-56.0	511	39.8	36.3-43.5	563	27.1	24.8-29.6	292	34.8	30.7-39.3	271	22.2	19.5-25.1
1977	967	45.9	42.9-49.0	459	54.9	49.7-60.5	508	40.6	37.0-44.4	590	28.2	25.9-30.7	268	33.7	29.5-38.3	322	25.1	22.4-28.1
1978	961	45.3	42.4-48.4	458	54.6	49.4-60.2	503	39.7	36.3-43.5	581	28.3	26.0-30.8	287	36.2	31.9-40.9	294	23.6	20.9-26.6
1979	1,018	48.8	45.8-52.0	469	57.8	52.4-63.6	549	43.4	39.7-47.3	590	28.1	25.8-30.6	264	31.6	27.8-35.9	326	25.9	23.1-28.9
1980	1,074	50.2	47.2-53.4	531	63.4	57.8-69.3	543	42.3	38.7-46.1	589	29.0	26.7-31.6	310	40.1	35.6-45.1	279	22.3	19.7-25.2
1981	1,066	50.8	47.7-54.0	511	63.1	57.5-69.1	555	43.6	40.0-47.5	586	28.2	25.9-30.7	239	30.4	26.5-34.7	347	27.2	24.4-30.3
1982	1,033	49.2	46.2-52.3	472	58.7	53.3-64.4	561	43.5	39.9-47.3	555	26.7	24.5-29.1	277	35.0	30.8-39.5	278	22.1	19.5-24.9
1983	1,051	49.6	46.5-52.7	479	58.6	53.2-64.3	572	44.5	40.9-48.4	529	25.5	23.3-27.8	254	31.9	27.9-36.2	275	21.6	19.1-24.4
1984	1,010	47.4	44.5-50.5	470	56.9	51.7-62.5	540	41.5	38.1-45.2	522	25.0	22.9-27.3	241	30.6	26.7-34.9	281	22.0	19.4-24.7
1985	1,072	50.1	47.1-53.3	501	61.1	55.6-66.8	571	43.9	40.3-47.7	538	25.0	22.9-27.2	248	29.5	25.8-33.6	290	22.1	19.6-24.9
1986	1,055	49.4	46.4-52.5	497	59.9	54.6-65.6	558	43.0	39.4-46.7	505	23.7	21.7-25.9	239	28.6	25.0-32.6	266	20.9	18.4-23.6
1987	966	44.3	41.5-47.3	478	57.3	52.1-62.9	488	36.5	33.3-39.9	544	25.6	23.5-27.9	265	32.6	28.7-36.9	279	21.6	19.1-24.3
1988	966	45.1	42.3-48.1	445	54.2	49.1-59.6	521	39.5	36.2-43.1	485	22.9	20.9-25.1	258	32.3	28.4-36.6	227	17.4	15.2-19.9
1989	937	43.9	41.1-46.9	470	56.2	51.1-61.7	467	36.2	33.0-39.7	472	22.4	20.4-24.5	229	28.9	25.2-33.0	243	18.5	16.3-21.1
1990	931	44.3	41.5-47.3	424	52.2	47.2-57.5	507	39.4	36.0-43.0	486	23.4	21.3-25.6	229	28.8	25.1-32.8	257	20.1	17.7-22.7
1991	962	45.8	42.9-48.8	467	58.3	53.0-63.9	495	38.3	35.0-41.9	457	21.8	19.9-24.0	227	28.7	25.0-32.8	230	17.7	15.5-20.2
1992	950	44.9	42.1-47.9	412	51.1	46.2-56.3	538	41.6	38.1-45.3	457	21.8	19.8-23.9	212	26.3	22.8-30.1	245	18.9	16.6-21.4
1993	972	46.1	43.2-49.1	451	55.6	50.5-61.1	521	40.4	37.0-44.1	463	22.0	20.1-24.2	209	25.8	22.4-29.6	254	19.6	17.3-22.2
1994	1,088	51.5	48.5-54.6	511	62.6	57.2-68.3	577	44.5	41.0-48.3	480	22.9	20.9-25.0	235	30.3	26.5-34.5	245	18.8	16.5-21.3
1995	1,001	47.2	44.4-50.3	457	56.1	51.0-61.6	544	41.8	38.3-45.5	506	24.0	21.9-26.2	251	31.8	27.9-36.1	255	19.5	17.2-22.0
1996	992	46.8	43.9-49.8	474	57.7	52.5-63.2	518	39.9	36.5-43.5	439	20.8	18.9-22.8	203	25.3	21.9-29.1	236	18.0	15.8-20.5
1997	1,008	47.3	44.4-50.3	469	57.1	52.0-62.6	539	41.3	37.9-44.9	439	20.7	18.8-22.7	222	27.5	23.9-31.4	217	16.4	14.3-18.8
1998	1,090	50.4	47.5-53.5	515	61.7	56.4-67.3	575	43.5	40.0-47.2	449	20.9	19.0-22.9	205	24.9	21.6-28.6	244	18.3	16.0-20.7
1999	1,079	48.7	45.8-51.7	478	55.3	50.4-60.5	601	44.6	41.1-48.4	451	20.4	18.6-22.4	211	24.8	21.5-28.4	240	17.7	15.5-20.1
2000	990	43.7	41.1-46.6	460	51.9	47.2-56.9	530	38.3	35.1-41.7	434	19.1	17.4-21.0	202	23.0	19.9-26.4	232	16.4	14.4-18.7
2001	1,063	46.4	43.7-49.3	482	54.4	49.6-59.6	581	41.7	38.4-45.2	421	18.4	16.7-20.2	196	22.4	19.4-25.8	225	16.0	14.0-18.2
2002	1,008	43.6	40.9-46.4	474	52.4	47.7-57.4	534	38.0	34.9-41.4	402	17.4	15.7-19.2	187	21.2	18.3-24.5	215	15.2	13.2-17.4
2003	942	40.5	38.0-43.2	461	50.5	45.9-55.4	481	34.2	31.2-37.5	456	19.6	17.8-21.4	214	23.7	20.6-27.1	242	16.9	14.8-19.1
2004	915	39.2	36.7-41.8	442	48.0	43.6-52.8	473	33.7	30.7-36.9	391	16.9	15.3-18.7	193	21.6	18.6-24.9	198	14.0	12.1-16.1
2005	944	40.1	37.6-42.8	429	45.8	41.5-50.4	515	36.3	33.2-39.6	383	16.4	14.8-18.2	191	21.1	18.1-24.3	192	13.4	11.5-15.4
2006	912	39.0	36.5-41.6	412	44.1	39.9-48.6	500	35.7	32.6-39.0	377	16.2	14.6-17.9	180	19.9	17.0-23.0	197	13.8	12.0-15.9
2007	910	38.5	36.1-41.1	416	43.6	39.4-48.0	494	35.3	32.3-38.6	348	14.8	13.3-16.5	163	17.5	14.9-20.4	185	13.0	11.1-15.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Colon excluding rectum**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	884	37.0	34.6-39.6	406	42.1	38.0-46.5	478	34.0	31.0-37.2	365	15.5	13.9-17.2	172	18.3	15.6-21.3	193	13.7	11.8-15.8
2009	893	37.0	34.6-39.5	408	41.8	37.8-46.1	485	33.7	30.8-36.9	359	15.0	13.5-16.6	181	19.5	16.7-22.6	178	12.1	10.4-14.1
2010	790	32.3	30.0-34.6	380	37.9	34.1-42.0	410	28.3	25.6-31.2	366	14.9	13.4-16.5	204	21.2	18.3-24.4	162	10.7	9.1-12.5
2011	801	32.3	30.0-34.6	380	37.3	33.6-41.3	421	28.7	26.0-31.6	347	14.1	12.7-15.7	164	16.9	14.3-19.7	183	12.3	10.6-14.3
2012	755	29.4	27.3-31.6	362	33.9	30.4-37.6	393	26.5	23.9-29.3	340	13.3	11.9-14.9	155	15.3	12.9-17.9	185	12.1	10.4-14.0
2013	818	31.6	29.5-33.9	409	38.2	34.5-42.1	409	27.1	24.5-30.0	315	12.0	10.7-13.5	140	13.3	11.1-15.7	175	11.2	9.5-13.0
2014	772	29.3	27.3-31.5	360	33.0	29.6-36.7	412	27.1	24.5-29.9	302	11.2	9.9-12.5	138	12.8	10.7-15.2	164	9.9	8.4-11.6
2015	729	26.7	24.8-28.8	339	29.7	26.5-33.1	390	24.8	22.4-27.5	271	9.8	8.7-11.1	124	11.1	9.2-13.3	147	8.9	7.5-10.5
2016	770	28.3	26.3-30.4	394	34.3	30.9-37.9	376	24.0	21.6-26.6	294	10.5	9.3-11.7	141	12.5	10.5-14.8	153	9.1	7.7-10.7
2017	778	28.1	26.1-30.2	365	31.0	27.8-34.4	413	26.0	23.5-28.7	316	11.3	10.1-12.7	145	12.8	10.8-15.1	171	10.3	8.8-12.1
2018	726	25.9	24.0-27.9	360	30.3	27.2-33.6	366	22.9	20.5-25.4	300	10.8	9.6-12.2	155	13.5	11.4-15.8	145	9.0	7.5-10.6
2019	779	27.8	25.8-29.9	403	33.5	30.3-37.0	376	23.5	21.1-26.1	264	9.3	8.2-10.5	130	11.1	9.2-13.2	134	8.0	6.7-9.5
2020	651	22.9	21.1-24.7	295	23.8	21.1-26.7	356	22.1	19.8-24.5	258	8.6	7.6-9.8	122	10.0	8.3-12.0	136	7.7	6.4-9.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Rectum & rectosigmoid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	378	17.5	15.7-19.5	200	22.9	19.7-26.6	178	14.0	11.9-16.3	148	7.1	6.0-8.4	79	9.6	7.5-12.1	69	5.5	4.2-7.0
1977	392	18.1	16.3-20.0	217	24.6	21.2-28.3	175	13.8	11.7-16.0	131	6.3	5.2-7.5	69	8.5	6.5-10.9	62	4.9	3.8-6.4
1978	395	17.8	16.1-19.8	194	21.7	18.6-25.1	201	15.3	13.2-17.7	114	5.6	4.6-6.7	67	7.9	6.0-10.1	47	4.0	2.9-5.4
1979	400	18.7	16.8-20.7	206	23.4	20.2-26.9	194	15.5	13.3-17.9	119	5.8	4.7-6.9	63	7.4	5.6-9.6	56	4.6	3.5-6.0
1980	430	20.0	18.1-22.0	232	25.8	22.5-29.5	198	15.9	13.7-18.3	120	5.8	4.8-7.0	69	8.2	6.3-10.4	51	4.3	3.1-5.6
1981	410	19.8	17.9-21.8	223	27.3	23.7-31.3	187	15.0	12.9-17.4	76	3.7	2.9-4.6	44	5.8	4.2-7.8	32	2.5	1.7-3.5
1982	367	17.1	15.3-18.9	200	22.9	19.7-26.4	167	13.1	11.1-15.3	83	4.0	3.2-5.0	45	5.6	4.0-7.5	38	3.0	2.1-4.1
1983	337	15.6	13.9-17.4	184	21.7	18.6-25.2	153	11.8	10.0-13.9	90	4.3	3.5-5.4	55	6.6	4.9-8.7	35	2.9	2.0-4.1
1984	363	16.5	14.8-18.3	197	22.0	18.9-25.4	166	12.9	11.0-15.0	105	4.9	4.0-6.0	57	6.8	5.1-8.9	48	3.7	2.7-5.0
1985	349	15.9	14.2-17.7	187	21.4	18.3-24.8	162	12.5	10.6-14.6	60	2.9	2.2-3.7	28	3.5	2.3-5.0	32	2.5	1.7-3.5
1986	360	16.3	14.7-18.1	198	22.6	19.4-26.0	162	12.4	10.6-14.5	81	3.8	3.0-4.8	46	6.0	4.3-8.0	35	2.6	1.8-3.7
1987	313	14.5	12.9-16.3	187	21.9	18.7-25.3	126	10.0	8.3-11.9	69	3.2	2.5-4.1	25	3.1	2.0-4.6	44	3.3	2.4-4.5
1988	316	14.5	12.9-16.2	174	20.6	17.6-24.0	142	10.6	8.9-12.5	62	3.0	2.3-3.8	36	4.4	3.0-6.1	26	2.1	1.4-3.1
1989	340	15.9	14.2-17.7	164	19.1	16.3-22.4	176	13.6	11.6-15.7	70	3.3	2.6-4.2	30	3.7	2.5-5.3	40	3.0	2.2-4.2
1990	294	13.7	12.2-15.4	137	16.3	13.6-19.3	157	12.0	10.2-14.1	70	3.4	2.6-4.3	33	4.2	2.8-5.9	37	2.9	2.0-4.0
1991	310	14.6	13.0-16.4	161	19.9	16.9-23.3	149	11.5	9.7-13.6	49	2.4	1.8-3.2	29	3.8	2.5-5.4	20	1.6	1.0-2.5
1992	310	14.7	13.1-16.4	149	17.7	14.9-20.9	161	12.6	10.7-14.7	52	2.6	1.9-3.4	21	2.9	1.7-4.4	31	2.5	1.7-3.6
1993	296	13.9	12.4-15.6	159	19.2	16.3-22.5	137	10.7	9.0-12.7	60	2.8	2.1-3.6	36	4.6	3.2-6.3	24	1.8	1.2-2.8
1994	342	16.1	14.4-17.9	183	21.7	18.7-25.1	159	12.5	10.6-14.6	59	2.8	2.1-3.6	29	3.6	2.4-5.1	30	2.3	1.5-3.3
1995	298	13.9	12.4-15.6	143	17.2	14.5-20.3	155	12.0	10.2-14.0	66	3.1	2.4-4.0	31	3.9	2.6-5.5	35	2.7	1.9-3.8
1996	317	15.0	13.3-16.7	173	20.6	17.6-24.0	144	11.4	9.6-13.4	59	2.7	2.1-3.5	28	3.2	2.1-4.7	31	2.4	1.6-3.4
1997	298	13.9	12.4-15.6	159	18.5	15.7-21.7	139	10.8	9.1-12.7	68	3.2	2.5-4.0	36	4.5	3.1-6.2	32	2.4	1.7-3.4
1998	369	16.9	15.3-18.8	184	21.1	18.1-24.4	185	14.1	12.1-16.3	54	2.5	1.9-3.3	29	3.4	2.3-4.9	25	1.9	1.3-2.9
1999	329	14.7	13.2-16.4	157	17.2	14.6-20.1	172	12.9	11.0-15.0	46	2.1	1.5-2.8	20	2.2	1.4-3.5	26	1.9	1.2-2.7
2000	331	14.5	13.0-16.2	176	18.9	16.2-22.0	155	11.3	9.6-13.3	55	2.4	1.8-3.2	28	3.1	2.1-4.5	27	1.9	1.3-2.8
2001	341	14.8	13.2-16.4	171	18.4	15.8-21.4	170	12.5	10.7-14.5	62	2.7	2.1-3.5	36	4.0	2.8-5.5	26	1.9	1.2-2.8
2002	310	13.3	11.9-14.9	165	17.3	14.8-20.2	145	10.4	8.8-12.3	47	2.0	1.5-2.7	26	2.8	1.8-4.1	21	1.5	0.9-2.3
2003	381	16.3	14.7-18.0	209	21.7	18.8-24.9	172	12.4	10.6-14.4	63	2.7	2.1-3.5	29	3.2	2.1-4.6	34	2.5	1.7-3.5
2004	375	15.9	14.3-17.6	191	19.5	16.8-22.5	184	13.3	11.4-15.4	65	2.8	2.1-3.5	39	4.2	3.0-5.8	26	1.8	1.2-2.7
2005	353	15.0	13.5-16.7	191	19.7	17.0-22.8	162	11.7	10.0-13.7	57	2.4	1.8-3.1	23	2.5	1.6-3.7	34	2.4	1.7-3.4
2006	341	14.3	12.8-15.9	167	17.0	14.5-19.8	174	12.6	10.7-14.6	51	2.2	1.6-2.8	28	2.9	1.9-4.3	23	1.6	1.0-2.5
2007	370	15.4	13.9-17.1	203	20.2	17.5-23.2	167	12.0	10.2-14.0	58	2.5	1.9-3.2	30	3.1	2.1-4.5	28	2.0	1.3-2.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Rectum & rectosigmoid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	288	12.0	10.7-13.5	145	14.2	12.0-16.8	143	10.4	8.8-12.3	66	2.7	2.1-3.5	32	3.3	2.3-4.7	34	2.3	1.6-3.3
2009	323	13.3	11.9-14.8	195	18.9	16.3-21.8	128	9.3	7.7-11.0	49	2.0	1.4-2.6	23	2.2	1.4-3.4	26	1.7	1.1-2.6
2010	342	13.9	12.5-15.5	191	18.7	16.1-21.6	151	10.6	9.0-12.5	49	2.0	1.5-2.7	30	2.8	1.9-4.0	19	1.4	0.8-2.2
2011	308	12.3	11.0-13.8	173	16.2	13.8-18.8	135	9.5	8.0-11.3	72	2.8	2.2-3.6	35	3.3	2.3-4.6	37	2.4	1.7-3.3
2012	309	11.7	10.5-13.1	169	14.8	12.6-17.2	140	9.5	8.0-11.3	71	2.8	2.2-3.5	37	3.5	2.5-4.9	34	2.2	1.5-3.2
2013	309	11.8	10.5-13.2	178	15.7	13.4-18.2	131	8.8	7.3-10.4	56	2.1	1.6-2.7	37	3.2	2.3-4.5	19	1.3	0.8-2.1
2014	306	11.6	10.3-13.0	169	14.9	12.7-17.3	137	9.0	7.5-10.7	58	2.2	1.7-2.9	28	2.6	1.7-3.8	30	1.9	1.3-2.8
2015	337	12.4	11.1-13.9	181	15.6	13.4-18.1	156	10.1	8.6-11.9	54	2.0	1.5-2.6	29	2.6	1.7-3.7	25	1.5	1.0-2.3
2016	312	11.6	10.3-12.9	176	15.3	13.1-17.8	136	8.8	7.3-10.4	70	2.6	2.0-3.4	34	3.1	2.1-4.4	36	2.4	1.6-3.3
2017	338	12.3	11.0-13.7	181	15.0	12.8-17.4	157	10.3	8.8-12.1	69	2.5	1.9-3.1	36	3.1	2.2-4.3	33	2.0	1.4-2.9
2018	325	11.7	10.5-13.1	181	15.3	13.1-17.7	144	9.0	7.6-10.7	56	2.1	1.5-2.7	31	2.6	1.8-3.8	25	1.6	1.0-2.4
2019	317	11.4	10.1-12.7	194	15.7	13.5-18.1	123	8.1	6.7-9.7	57	2.0	1.5-2.7	31	2.7	1.8-3.8	26	1.6	1.0-2.3
2020	240	8.7	7.6-9.9	143	11.6	9.7-13.7	97	6.3	5.1-7.7	66	2.3	1.8-3.0	35	3.0	2.1-4.2	31	1.8	1.2-2.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Liver / intrahepatic bile duct**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	89	4.1	3.3-5.1	50	5.4	4.0-7.3	39	3.1	2.2-4.3	94	4.5	3.6-5.5	54	5.9	4.4-7.9	40	3.4	2.4-4.7
1977	76	3.6	2.8-4.6	39	4.3	3.0-5.9	37	3.1	2.2-4.4	106	4.9	3.9-5.9	57	6.1	4.5-7.9	49	4.0	2.9-5.3
1978	72	3.4	2.6-4.3	43	4.8	3.4-6.6	29	2.3	1.5-3.3	80	3.7	2.9-4.6	40	4.5	3.2-6.3	40	3.2	2.2-4.3
1979	67	3.2	2.4-4.0	39	4.7	3.3-6.5	28	2.2	1.4-3.1	107	5.1	4.2-6.2	48	5.8	4.2-7.7	59	4.8	3.6-6.2
1980	80	3.9	3.0-4.8	44	5.1	3.6-6.9	36	3.0	2.1-4.2	84	3.9	3.1-4.9	42	4.8	3.4-6.5	42	3.3	2.4-4.5
1981	87	4.2	3.3-5.2	53	6.1	4.6-8.1	34	2.7	1.9-3.8	107	5.2	4.2-6.3	62	7.7	5.8-10.0	45	3.5	2.6-4.8
1982	73	3.4	2.6-4.3	44	5.2	3.7-7.1	29	2.1	1.4-3.1	103	4.9	4.0-6.0	54	6.4	4.7-8.4	49	3.9	2.9-5.2
1983	72	3.4	2.6-4.2	44	5.1	3.7-6.9	28	2.3	1.5-3.3	109	4.9	4.0-6.0	59	6.5	4.9-8.5	50	3.8	2.8-5.1
1984	93	4.1	3.3-5.1	55	6.1	4.6-8.0	38	2.8	1.9-3.8	106	4.8	3.9-5.9	61	7.0	5.3-9.1	45	3.4	2.5-4.6
1985	85	4.0	3.2-5.0	53	6.5	4.8-8.6	32	2.5	1.7-3.6	123	5.9	4.9-7.1	76	9.4	7.3-11.8	47	3.6	2.6-4.8
1986	92	4.3	3.5-5.3	51	5.9	4.3-7.7	41	3.3	2.3-4.4	124	5.8	4.8-6.9	60	6.9	5.2-8.9	64	5.0	3.8-6.4
1987	80	3.7	2.9-4.6	55	6.3	4.7-8.3	25	1.9	1.2-2.8	129	6.1	5.1-7.3	69	8.4	6.5-10.6	60	4.7	3.6-6.1
1988	96	4.6	3.7-5.6	57	6.5	4.9-8.4	39	3.2	2.3-4.4	124	5.8	4.8-7.0	69	7.9	6.1-10.0	55	4.4	3.3-5.7
1989	91	4.3	3.5-5.3	55	6.3	4.7-8.3	36	2.8	2.0-3.9	122	5.8	4.8-7.0	70	8.3	6.4-10.5	52	4.1	3.1-5.4
1990	115	5.4	4.5-6.5	67	7.8	6.0-10.0	48	3.7	2.7-4.9	127	6.1	5.1-7.3	69	8.6	6.6-10.9	58	4.4	3.4-5.7
1991	104	4.8	3.9-5.8	66	7.5	5.8-9.6	38	2.9	2.0-4.0	105	4.8	3.9-5.8	57	6.4	4.8-8.3	48	3.6	2.6-4.8
1992	125	5.9	4.9-7.1	76	9.0	7.1-11.4	49	3.9	2.9-5.2	142	6.7	5.6-7.9	85	10.1	8.1-12.6	57	4.4	3.3-5.7
1993	115	5.4	4.4-6.5	73	8.4	6.5-10.5	42	3.3	2.4-4.5	128	6.0	5.0-7.1	72	8.3	6.5-10.5	56	4.4	3.3-5.7
1994	147	6.9	5.8-8.1	91	10.6	8.5-13.0	56	4.4	3.3-5.7	111	5.3	4.3-6.3	72	8.5	6.6-10.7	39	3.0	2.2-4.2
1995	137	6.4	5.4-7.6	81	9.2	7.3-11.4	56	4.3	3.3-5.6	114	5.4	4.5-6.5	71	8.3	6.4-10.4	43	3.2	2.3-4.4
1996	152	7.1	6.0-8.3	104	11.8	9.6-14.3	48	3.7	2.8-5.0	133	6.2	5.2-7.3	81	9.1	7.2-11.3	52	4.0	3.0-5.3
1997	176	8.2	7.1-9.6	112	12.8	10.5-15.4	64	4.9	3.8-6.3	138	6.5	5.4-7.7	80	9.2	7.3-11.5	58	4.4	3.4-5.7
1998	181	8.2	7.1-9.5	122	13.4	11.1-16.0	59	4.5	3.4-5.8	146	6.7	5.6-7.9	102	11.3	9.2-13.7	44	3.4	2.4-4.5
1999	174	7.8	6.7-9.0	98	10.6	8.6-12.9	76	5.7	4.5-7.2	117	5.3	4.3-6.3	72	7.9	6.1-9.9	45	3.4	2.5-4.5
2000	199	8.7	7.5-10.0	143	14.8	12.4-17.4	56	4.2	3.1-5.4	135	5.9	5.0-7.0	86	9.1	7.2-11.2	49	3.6	2.6-4.7
2001	186	8.0	6.9-9.3	140	14.4	12.1-17.0	46	3.4	2.5-4.5	141	6.1	5.2-7.2	93	9.8	7.9-12.1	48	3.5	2.6-4.6
2002	173	7.4	6.3-8.6	128	13.1	10.9-15.6	45	3.3	2.4-4.4	132	5.7	4.8-6.7	90	9.5	7.6-11.7	42	2.9	2.1-4.0
2003	213	9.0	7.9-10.3	143	14.4	12.1-17.0	70	5.1	4.0-6.5	143	6.1	5.1-7.2	98	10.3	8.3-12.5	45	3.2	2.4-4.3
2004	215	9.1	7.9-10.4	138	13.7	11.5-16.2	77	5.6	4.4-7.0	162	6.9	5.9-8.0	93	9.1	7.3-11.2	69	4.9	3.8-6.2
2005	216	9.0	7.8-10.3	153	14.9	12.6-17.5	63	4.4	3.4-5.6	148	6.3	5.3-7.4	85	8.5	6.7-10.5	63	4.4	3.4-5.7
2006	215	9.0	7.8-10.3	149	14.4	12.1-16.9	66	4.8	3.7-6.1	164	6.8	5.8-8.0	116	11.6	9.6-13.9	48	3.4	2.5-4.5
2007	238	9.8	8.5-11.1	166	15.9	13.6-18.6	72	5.2	4.1-6.6	160	6.6	5.6-7.8	110	10.8	8.9-13.1	50	3.5	2.6-4.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Liver / intrahepatic bile duct**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	223	8.9	7.7-10.1	168	15.4	13.1-17.9	55	3.8	2.9-5.0	147	6.0	5.0-7.0	96	9.1	7.4-11.2	51	3.5	2.6-4.7
2009	235	9.6	8.4-10.9	167	15.7	13.4-18.4	68	4.9	3.8-6.2	156	6.5	5.5-7.6	101	9.8	7.9-11.9	55	3.9	3.0-5.1
2010	267	10.7	9.5-12.1	186	17.3	14.8-20.0	81	5.7	4.5-7.1	176	7.0	6.0-8.2	114	10.9	8.9-13.1	62	4.1	3.2-5.3
2011	271	10.5	9.3-11.9	196	17.3	14.9-19.9	75	5.3	4.1-6.6	170	6.8	5.8-7.9	106	9.7	7.9-11.8	64	4.3	3.3-5.6
2012	293	11.2	9.9-12.5	192	17.1	14.7-19.8	101	6.7	5.4-8.2	176	6.8	5.9-7.9	116	10.6	8.7-12.7	60	4.1	3.1-5.3
2013	310	11.4	10.2-12.8	219	18.4	16.0-21.1	91	6.0	4.8-7.3	186	7.0	6.0-8.1	119	10.5	8.7-12.6	67	4.3	3.3-5.4
2014	272	9.9	8.7-11.2	193	15.8	13.6-18.2	79	5.2	4.1-6.5	180	6.6	5.6-7.6	128	11.0	9.1-13.1	52	3.3	2.5-4.4
2015	294	10.6	9.4-11.9	202	16.6	14.3-19.1	92	5.8	4.6-7.1	200	7.3	6.3-8.4	127	10.8	8.9-12.8	73	4.6	3.6-5.8
2016	319	11.2	9.9-12.5	227	18.2	15.9-20.8	92	5.8	4.6-7.1	177	6.3	5.4-7.3	120	9.7	8.0-11.7	57	3.6	2.7-4.8
2017	292	10.2	9.1-11.5	207	16.5	14.3-19.0	85	5.2	4.2-6.5	186	6.6	5.7-7.6	132	10.9	9.1-13.0	54	3.3	2.5-4.4
2018	269	9.4	8.3-10.6	202	16.5	14.3-19.0	67	4.1	3.1-5.2	154	5.4	4.6-6.3	102	8.3	6.7-10.1	52	3.1	2.3-4.1
2019	264	9.0	8.0-10.2	172	13.6	11.6-15.8	92	5.5	4.4-6.8	181	6.2	5.3-7.2	111	9.1	7.4-11.0	70	4.0	3.1-5.1
2020	238	8.3	7.2-9.4	163	13.2	11.2-15.4	75	4.3	3.4-5.5	141	4.8	4.0-5.7	86	7.0	5.6-8.7	55	3.1	2.3-4.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Pancreas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	289	13.2	11.6-14.9	133	14.1	11.7-16.9	156	12.4	10.5-14.6	250	11.3	9.9-12.9	111	11.5	9.3-13.9	139	11.1	9.2-13.2
1977	307	14.3	12.6-16.0	147	16.7	13.9-19.8	160	12.6	10.7-14.8	302	14.2	12.6-16.0	149	17.5	14.6-20.8	153	12.0	10.2-14.2
1978	293	13.5	12.0-15.2	141	15.7	13.0-18.6	152	12.0	10.1-14.1	259	11.9	10.5-13.5	124	14.1	11.5-17.0	135	10.4	8.7-12.4
1979	282	13.1	11.5-14.7	128	14.1	11.7-16.8	154	12.2	10.3-14.3	275	12.9	11.3-14.5	132	14.9	12.3-17.8	143	11.4	9.6-13.5
1980	295	13.7	12.1-15.4	155	17.9	15.1-21.1	140	10.9	9.1-12.9	254	11.8	10.4-13.4	133	15.4	12.8-18.4	121	9.5	7.8-11.4
1981	279	13.2	11.6-14.9	145	17.2	14.4-20.4	134	10.5	8.7-12.5	271	12.6	11.1-14.2	132	15.2	12.6-18.1	139	10.8	9.0-12.8
1982	280	13.2	11.7-14.9	126	15.3	12.6-18.3	154	11.8	10.0-13.8	234	11.3	9.8-12.8	116	14.7	12.0-17.8	118	9.2	7.6-11.0
1983	294	13.7	12.1-15.4	124	14.7	12.2-17.7	170	12.9	11.0-15.1	266	12.4	10.9-14.0	128	15.4	12.7-18.4	138	10.4	8.7-12.3
1984	302	13.8	12.2-15.4	149	17.4	14.6-20.5	153	11.4	9.6-13.4	242	11.1	9.7-12.6	120	13.9	11.4-16.7	122	9.2	7.6-11.0
1985	276	12.9	11.4-14.5	125	14.9	12.3-17.8	151	11.4	9.6-13.4	228	10.5	9.2-12.0	106	12.5	10.2-15.2	122	9.1	7.5-10.9
1986	265	12.3	10.8-13.9	117	13.8	11.4-16.7	148	11.1	9.4-13.1	244	11.2	9.9-12.8	94	11.1	8.9-13.7	150	11.2	9.4-13.1
1987	280	12.8	11.3-14.4	131	15.7	13.0-18.7	149	10.9	9.2-12.8	233	10.6	9.3-12.1	121	14.3	11.8-17.2	112	8.3	6.8-10.0
1988	260	12.1	10.7-13.7	106	13.2	10.8-16.1	154	11.5	9.7-13.5	237	11.0	9.6-12.5	119	14.5	12.0-17.4	118	8.8	7.2-10.5
1989	259	12.3	10.8-13.9	120	14.4	11.8-17.2	139	10.7	9.0-12.7	245	11.4	10.0-13.0	99	11.9	9.6-14.5	146	10.9	9.2-12.9
1990	248	11.8	10.4-13.4	114	14.4	11.8-17.3	134	10.5	8.8-12.5	212	10.1	8.8-11.6	100	12.8	10.4-15.6	112	8.5	7.0-10.3
1991	261	12.3	10.9-13.9	118	14.3	11.8-17.2	143	10.9	9.2-12.8	224	10.5	9.1-11.9	110	13.2	10.8-16.0	114	8.7	7.2-10.5
1992	281	13.3	11.8-14.9	128	15.5	12.9-18.5	153	11.8	10.0-13.8	222	10.4	9.1-11.9	96	11.4	9.2-13.9	126	9.7	8.0-11.5
1993	272	12.8	11.4-14.5	116	13.9	11.4-16.7	156	12.0	10.2-14.1	242	11.5	10.0-13.0	109	13.5	11.0-16.3	133	10.3	8.6-12.2
1994	318	15.1	13.5-16.9	137	16.8	14.0-19.9	181	13.9	11.9-16.0	255	12.1	10.6-13.7	107	13.4	10.9-16.2	148	11.3	9.6-13.3
1995	238	11.2	9.9-12.8	102	12.7	10.3-15.4	136	10.4	8.7-12.3	214	10.1	8.8-11.6	98	12.3	9.9-15.0	116	8.9	7.3-10.6
1996	284	13.4	11.9-15.1	121	15.0	12.4-17.9	163	12.5	10.6-14.6	247	11.7	10.3-13.3	106	13.4	10.9-16.2	141	10.8	9.1-12.7
1997	253	11.8	10.4-13.3	117	14.1	11.7-17.0	136	10.4	8.7-12.3	218	10.2	8.9-11.7	91	11.1	8.9-13.7	127	9.7	8.1-11.5
1998	276	12.8	11.3-14.4	116	13.7	11.3-16.4	160	12.1	10.3-14.2	224	10.4	9.1-11.8	89	10.8	8.7-13.3	135	10.2	8.6-12.1
1999	275	12.4	11.0-13.9	135	15.6	13.1-18.5	140	10.4	8.7-12.2	222	10.0	8.8-11.4	103	11.8	9.6-14.4	119	8.8	7.3-10.6
2000	290	12.8	11.4-14.4	128	14.5	12.1-17.2	162	11.7	10.0-13.6	227	10.0	8.8-11.4	101	11.4	9.3-13.9	126	8.9	7.4-10.6
2001	281	12.3	10.9-13.8	110	12.4	10.1-14.9	171	12.3	10.5-14.3	252	11.0	9.7-12.5	101	11.6	9.4-14.1	151	10.8	9.1-12.6
2002	285	12.3	11.0-13.9	129	13.9	11.6-16.6	156	11.1	9.4-13.0	224	9.7	8.5-11.1	106	11.5	9.4-13.9	118	8.4	6.9-10.0
2003	268	11.6	10.2-13.1	126	13.7	11.4-16.3	142	10.0	8.4-11.8	209	9.0	7.8-10.3	91	9.9	8.0-12.2	118	8.3	6.9-9.9
2004	292	12.5	11.1-14.0	129	14.0	11.6-16.6	163	11.4	9.7-13.3	236	10.1	8.8-11.5	103	11.2	9.1-13.6	133	9.3	7.8-11.1
2005	277	11.8	10.4-13.3	127	13.6	11.3-16.2	150	10.7	9.1-12.6	199	8.5	7.4-9.8	88	9.4	7.5-11.6	111	7.8	6.4-9.4
2006	314	13.3	11.9-14.9	141	14.9	12.5-17.6	173	12.2	10.4-14.2	232	9.8	8.6-11.2	111	11.9	9.7-14.3	121	8.5	7.1-10.2
2007	317	13.3	11.9-14.9	151	16.3	13.8-19.2	166	11.5	9.8-13.4	250	10.5	9.2-11.9	118	12.7	10.5-15.3	132	9.3	7.8-11.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Kings (Brooklyn)

Pancreas

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	323	13.5	12.0-15.0	153	15.8	13.4-18.6	170	11.7	10.0-13.6	254	10.7	9.4-12.1	105	11.1	9.0-13.4	149	10.2	8.6-12.0
2009	344	14.2	12.8-15.8	160	16.6	14.1-19.5	184	12.7	10.9-14.7	257	10.6	9.3-12.0	126	13.0	10.8-15.5	131	8.9	7.5-10.6
2010	371	15.1	13.6-16.7	175	17.0	14.5-19.7	196	13.4	11.6-15.5	266	10.9	9.6-12.3	123	12.2	10.1-14.6	143	9.6	8.1-11.4
2011	347	14.1	12.6-15.7	159	15.6	13.2-18.3	188	12.7	10.9-14.7	298	12.0	10.6-13.4	142	14.1	11.8-16.6	156	10.4	8.8-12.2
2012	387	15.2	13.7-16.8	176	16.6	14.2-19.3	211	14.2	12.3-16.3	308	12.3	10.9-13.7	138	13.3	11.2-15.8	170	11.7	10.0-13.6
2013	360	13.9	12.5-15.5	173	16.4	14.0-19.1	187	12.3	10.5-14.2	277	10.9	9.6-12.2	134	13.1	11.0-15.6	143	9.3	7.8-11.0
2014	339	12.9	11.5-14.3	161	14.6	12.4-17.1	178	11.5	9.9-13.4	260	9.8	8.7-11.1	114	10.4	8.6-12.6	146	9.3	7.9-11.0
2015	381	14.0	12.6-15.5	179	15.3	13.1-17.8	202	12.8	11.0-14.7	285	10.5	9.3-11.8	132	11.6	9.7-13.8	153	9.5	8.0-11.2
2016	370	13.4	12.0-14.9	170	14.4	12.3-16.8	200	12.5	10.8-14.4	265	9.6	8.5-10.9	131	11.0	9.2-13.1	134	8.4	7.0-10.0
2017	413	14.8	13.4-16.3	211	18.1	15.7-20.8	202	12.4	10.7-14.3	297	10.6	9.4-11.9	140	12.3	10.3-14.5	157	9.4	8.0-11.0
2018	393	13.8	12.5-15.3	168	13.8	11.8-16.1	225	13.7	12.0-15.7	299	10.7	9.5-12.0	142	12.2	10.2-14.4	157	9.5	8.0-11.2
2019	382	13.4	12.1-14.9	187	15.4	13.2-17.8	195	11.8	10.2-13.6	262	9.2	8.1-10.4	121	10.1	8.4-12.1	141	8.5	7.1-10.0
2020	365	12.4	11.2-13.8	165	13.1	11.1-15.3	200	12.0	10.4-13.8	288	9.9	8.8-11.1	140	11.7	9.8-13.9	148	8.6	7.3-10.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Kings (Brooklyn)

Larynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	116	4.9	4.1-6.0	94	9.4	7.5-11.6	22	1.7	1.0-2.6	60	2.5	1.9-3.3	48	4.6	3.4-6.2	12	1.0	0.5-1.7
1977	133	5.8	4.8-6.9	113	11.6	9.5-14.0	20	1.6	1.0-2.5	53	2.4	1.8-3.2	45	5.0	3.6-6.8	8	0.6	0.3-1.2
1978	110	5.2	4.3-6.3	92	10.5	8.4-13.0	18	1.6	0.9-2.5	51	2.4	1.7-3.1	43	4.8	3.4-6.6	8	0.6	0.3-1.3
1979	145	6.4	5.4-7.5	121	12.5	10.3-15.0	24	2.0	1.3-3.0	66	3.0	2.3-3.9	56	6.2	4.6-8.1	10	0.8	0.4-1.5
1980	148	6.9	5.8-8.1	115	12.9	10.6-15.5	33	2.7	1.8-3.8	53	2.5	1.8-3.3	38	4.3	3.0-6.0	15	1.3	0.7-2.1
1981	118	5.4	4.4-6.5	94	10.4	8.3-12.8	24	1.9	1.2-2.9	44	1.9	1.4-2.6	29	3.3	2.2-4.8	15	1.1	0.6-1.9
1982	138	6.3	5.3-7.5	116	12.9	10.6-15.5	22	1.7	1.1-2.6	42	1.9	1.3-2.5	40	4.5	3.2-6.2	2	0.1	0.0-0.5
1983	111	5.1	4.2-6.2	86	9.5	7.5-11.8	25	2.1	1.3-3.1	47	2.2	1.6-3.0	33	3.9	2.6-5.5	14	1.0	0.6-1.7
1984	125	6.0	4.9-7.1	98	11.2	9.1-13.8	27	2.4	1.5-3.5	69	3.2	2.5-4.1	55	6.3	4.7-8.3	14	1.1	0.6-1.8
1985	140	6.3	5.3-7.5	105	11.6	9.4-14.1	35	2.8	1.9-3.9	46	2.1	1.5-2.8	38	4.5	3.1-6.2	8	0.6	0.3-1.2
1986	143	6.5	5.4-7.7	116	12.8	10.5-15.4	27	2.2	1.4-3.2	53	2.4	1.8-3.2	46	5.3	3.8-7.1	7	0.5	0.2-1.1
1987	123	5.8	4.8-7.0	96	10.9	8.8-13.4	27	2.3	1.5-3.3	41	1.9	1.3-2.5	31	3.5	2.3-4.9	10	0.8	0.4-1.5
1988	130	6.1	5.1-7.2	111	12.4	10.1-14.9	19	1.5	0.9-2.3	54	2.6	2.0-3.4	44	5.3	3.8-7.1	10	0.8	0.4-1.4
1989	120	5.8	4.8-6.9	85	9.7	7.7-12.0	35	2.9	2.0-4.1	62	3.0	2.3-3.9	44	5.2	3.8-7.1	18	1.4	0.8-2.3
1990	141	6.7	5.6-7.9	108	12.5	10.2-15.1	33	2.7	1.9-3.8	53	2.5	1.9-3.3	43	5.0	3.6-6.8	10	0.8	0.4-1.4
1991	122	5.9	4.9-7.1	100	11.8	9.6-14.4	22	1.9	1.2-2.9	45	2.3	1.6-3.0	36	4.3	3.0-6.0	9	0.8	0.3-1.5
1992	128	6.1	5.0-7.2	103	11.8	9.6-14.4	25	2.1	1.3-3.1	59	2.9	2.2-3.7	45	5.3	3.9-7.1	14	1.2	0.6-2.0
1993	98	4.7	3.8-5.7	71	8.1	6.3-10.2	27	2.2	1.5-3.2	37	1.7	1.2-2.4	30	3.6	2.4-5.2	7	0.6	0.2-1.2
1994	101	4.7	3.9-5.8	83	9.7	7.7-12.0	18	1.4	0.8-2.2	38	1.8	1.3-2.5	31	3.8	2.6-5.4	7	0.5	0.2-1.1
1995	102	4.8	3.9-5.8	81	9.2	7.3-11.4	21	1.8	1.1-2.7	46	2.2	1.6-2.9	40	4.6	3.3-6.3	6	0.5	0.2-1.0
1996	123	5.8	4.8-6.9	95	10.9	8.8-13.4	28	2.2	1.5-3.3	47	2.2	1.6-3.0	43	4.9	3.5-6.6	4	0.3	0.1-0.8
1997	106	5.0	4.1-6.0	82	9.3	7.4-11.6	24	1.9	1.2-2.9	36	1.7	1.2-2.3	26	3.0	2.0-4.4	10	0.8	0.4-1.5
1998	93	4.2	3.4-5.2	71	8.0	6.2-10.1	22	1.7	1.1-2.6	55	2.5	1.9-3.3	44	5.0	3.6-6.7	11	0.9	0.4-1.5
1999	91	4.1	3.3-5.0	71	7.6	5.9-9.6	20	1.6	0.9-2.4	35	1.6	1.1-2.2	25	2.7	1.7-4.0	10	0.7	0.4-1.4
2000	105	4.6	3.8-5.6	86	9.3	7.4-11.5	19	1.5	0.9-2.3	42	1.9	1.3-2.5	36	3.9	2.7-5.4	6	0.4	0.2-1.0
2001	83	3.6	2.9-4.5	63	6.5	5.0-8.3	20	1.5	0.9-2.3	30	1.3	0.9-1.9	23	2.5	1.6-3.7	7	0.5	0.2-1.0
2002	95	4.1	3.3-5.0	74	7.5	5.9-9.4	21	1.6	1.0-2.4	34	1.5	1.0-2.0	28	2.9	1.9-4.1	6	0.5	0.2-1.0
2003	105	4.5	3.7-5.4	82	8.4	6.6-10.4	23	1.7	1.1-2.6	39	1.7	1.2-2.3	34	3.5	2.4-5.0	5	0.4	0.1-0.9
2004	115	4.8	4.0-5.8	100	10.1	8.2-12.2	15	1.1	0.6-1.8	39	1.6	1.1-2.2	33	3.2	2.2-4.6	6	0.5	0.2-1.0
2005	109	4.5	3.7-5.5	79	7.9	6.2-9.8	30	2.2	1.5-3.1	32	1.4	0.9-1.9	24	2.6	1.7-3.9	8	0.5	0.2-1.1
2006	75	3.1	2.4-3.9	59	5.8	4.4-7.4	16	1.1	0.6-1.9	31	1.3	0.9-1.9	27	2.9	1.9-4.2	4	0.3	0.1-0.7
2007	81	3.3	2.6-4.1	63	6.1	4.6-7.8	18	1.3	0.8-2.1	33	1.4	0.9-1.9	26	2.6	1.7-3.8	7	0.5	0.2-1.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Kings (Brooklyn)

Larynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	89	3.7	2.9-4.5	67	6.7	5.1-8.5	22	1.6	1.0-2.4	24	1.1	0.7-1.6	16	1.8	1.0-2.9	8	0.6	0.2-1.2
2009	83	3.3	2.7-4.1	60	5.6	4.3-7.3	23	1.7	1.0-2.5	37	1.5	1.1-2.1	26	2.6	1.7-3.8	11	0.7	0.4-1.4
2010	90	3.6	2.9-4.4	75	7.1	5.6-8.9	15	1.1	0.6-1.8	30	1.3	0.8-1.8	25	2.5	1.6-3.8	5	0.4	0.1-0.9
2011	76	2.9	2.3-3.6	54	4.8	3.6-6.3	22	1.5	0.9-2.3	34	1.4	0.9-1.9	24	2.3	1.5-3.5	10	0.7	0.3-1.3
2012	69	2.7	2.1-3.4	56	5.1	3.8-6.6	13	0.9	0.5-1.5	20	0.8	0.5-1.2	17	1.5	0.9-2.5	3	0.2	0.0-0.6
2013	82	3.0	2.4-3.8	73	6.5	5.1-8.2	9	0.6	0.3-1.1	27	1.0	0.7-1.5	24	2.2	1.4-3.3	3	0.2	0.0-0.6
2014	71	2.6	2.1-3.3	59	5.1	3.9-6.7	12	0.8	0.4-1.4	22	0.8	0.5-1.2	18	1.5	0.9-2.4	4	0.2	0.1-0.7
2015	89	3.2	2.5-3.9	75	6.4	5.0-8.1	14	0.8	0.5-1.4	19	0.8	0.5-1.2	14	1.3	0.7-2.2	5	0.4	0.1-0.8
2016	77	2.8	2.2-3.5	59	5.2	3.9-6.8	18	1.1	0.6-1.8	35	1.3	0.9-1.8	30	2.7	1.8-3.9	5	0.3	0.1-0.7
2017	87	3.0	2.4-3.8	65	5.3	4.1-6.8	22	1.4	0.9-2.2	18	0.6	0.4-1.0	17	1.5	0.8-2.4	1	0.1	0.0-0.4
2018	54	1.9	1.4-2.4	45	3.6	2.6-4.8	9	0.6	0.3-1.1	22	0.8	0.5-1.2	19	1.6	0.9-2.5	3	0.2	0.0-0.6
2019	69	2.3	1.8-3.0	53	4.3	3.2-5.6	16	0.9	0.5-1.6	18	0.7	0.4-1.1	17	1.5	0.9-2.5	1	0.1	0.0-0.4
2020	42	1.4	1.0-2.0	34	2.7	1.8-3.8	8	0.5	0.2-1.0	15	0.5	0.3-0.8	13	1.0	0.5-1.8	2	0.1	0.0-0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Lung and bronchus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,097	48.3	45.4-51.4	771	81.7	75.8-88.0	326	25.3	22.5-28.3	926	40.9	38.2-43.7	663	70.8	65.2-76.7	263	20.3	17.8-23.0
1977	1,145	50.2	47.3-53.3	811	87.1	80.9-93.6	334	25.1	22.4-28.0	939	41.8	39.1-44.7	691	74.8	69.0-80.8	248	19.4	17.0-22.0
1978	1,142	51.4	48.3-54.5	804	87.0	80.9-93.5	338	26.9	24.0-30.0	924	41.7	38.9-44.5	657	71.9	66.3-77.8	267	21.0	18.5-23.7
1979	1,136	51.3	48.3-54.5	777	86.2	80.0-92.8	359	27.8	24.9-30.9	932	41.7	39.0-44.5	667	74.1	68.4-80.2	265	19.9	17.6-22.6
1980	1,174	53.5	50.4-56.8	815	90.8	84.4-97.5	359	27.9	25.0-31.1	970	44.2	41.4-47.2	686	76.3	70.5-82.5	284	22.4	19.8-25.3
1981	1,152	52.7	49.6-55.9	779	89.0	82.7-95.8	373	28.6	25.7-31.7	950	43.6	40.8-46.6	678	77.6	71.7-84.0	272	20.9	18.4-23.6
1982	1,198	54.2	51.1-57.5	785	87.7	81.5-94.3	413	31.8	28.8-35.2	942	42.7	39.9-45.6	634	71.2	65.6-77.2	308	23.8	21.2-26.7
1983	1,155	52.3	49.3-55.5	789	88.9	82.6-95.5	366	28.1	25.2-31.2	999	45.5	42.7-48.5	667	76.7	70.7-82.9	332	25.3	22.6-28.3
1984	1,223	56.1	52.9-59.4	815	92.7	86.3-99.5	408	32.0	28.9-35.4	958	43.8	41.0-46.7	659	75.3	69.5-81.4	299	22.8	20.2-25.6
1985	1,151	52.3	49.3-55.5	750	85.1	78.9-91.6	401	31.1	28.1-34.4	923	42.0	39.3-44.8	591	67.5	62.0-73.4	332	25.7	23.0-28.7
1986	1,215	55.2	52.1-58.5	770	87.1	80.9-93.6	445	34.4	31.2-37.9	960	44.0	41.2-47.0	616	70.5	64.9-76.5	344	27.3	24.4-30.4
1987	1,232	55.4	52.3-58.6	803	89.4	83.2-96.0	429	32.8	29.7-36.1	981	44.4	41.7-47.4	651	74.4	68.6-80.5	330	25.0	22.3-27.9
1988	1,173	53.7	50.6-56.9	713	82.2	76.2-88.6	460	35.1	31.9-38.5	994	45.1	42.3-48.1	659	75.6	69.8-81.8	335	25.5	22.8-28.5
1989	1,241	57.2	54.0-60.5	763	87.1	81.0-93.7	478	37.1	33.8-40.6	970	45.0	42.1-47.9	606	70.7	65.1-76.7	364	28.1	25.2-31.2
1990	1,243	58.0	54.8-61.4	799	94.3	87.7-101.2	444	34.7	31.5-38.1	1,011	47.4	44.5-50.5	652	77.5	71.5-83.8	359	28.1	25.2-31.2
1991	1,287	61.0	57.7-64.4	773	91.5	85.1-98.3	514	40.8	37.3-44.5	1,033	48.8	45.8-51.8	658	79.1	73.0-85.4	375	29.3	26.4-32.4
1992	1,296	61.0	57.7-64.5	815	96.4	89.8-103.3	481	37.9	34.5-41.4	994	47.0	44.1-50.1	613	73.5	67.7-79.6	381	30.0	27.1-33.3
1993	1,290	60.5	57.2-63.9	760	89.8	83.5-96.6	530	41.8	38.2-45.5	1,038	48.7	45.8-51.8	631	75.8	69.9-82.1	407	31.7	28.6-34.9
1994	1,256	59.1	55.9-62.5	726	85.3	79.1-91.8	530	42.2	38.7-46.0	1,009	47.4	44.6-50.5	629	74.5	68.7-80.6	380	29.8	26.9-33.0
1995	1,210	56.8	53.6-60.1	729	85.7	79.5-92.2	481	37.9	34.6-41.5	976	45.9	43.0-48.9	594	70.7	65.0-76.6	382	29.9	27.0-33.1
1996	1,232	57.6	54.5-61.0	703	82.1	76.1-88.5	529	41.6	38.2-45.4	937	43.9	41.1-46.8	552	64.8	59.4-70.5	385	30.0	27.1-33.2
1997	1,249	58.2	55.0-61.5	707	82.3	76.3-88.7	542	42.2	38.7-45.9	956	44.7	41.9-47.6	586	69.2	63.6-75.1	370	28.8	25.9-31.9
1998	1,227	56.4	53.3-59.6	668	75.7	70.0-81.7	559	43.3	39.7-47.0	945	43.5	40.8-46.4	524	59.9	54.8-65.3	421	32.4	29.4-35.7
1999	1,207	54.2	51.2-57.4	674	74.9	69.3-80.8	533	40.2	36.8-43.7	860	38.7	36.2-41.4	489	54.9	50.1-60.0	371	28.0	25.2-31.0
2000	1,175	51.8	48.8-54.8	665	72.8	67.3-78.6	510	38.0	34.7-41.4	886	39.1	36.6-41.8	504	55.5	50.7-60.6	382	28.0	25.3-31.0
2001	1,145	50.0	47.1-53.0	636	69.3	63.9-74.9	509	37.4	34.2-40.8	833	36.4	33.9-38.9	477	52.7	48.0-57.6	356	25.9	23.3-28.7
2002	1,184	51.0	48.1-54.0	652	69.9	64.5-75.5	532	38.6	35.4-42.0	883	38.0	35.5-40.6	515	55.2	50.4-60.2	368	26.5	23.8-29.3
2003	1,123	48.3	45.5-51.2	630	67.8	62.5-73.3	493	35.5	32.5-38.8	839	36.2	33.8-38.7	473	51.1	46.5-56.0	366	26.3	23.6-29.1
2004	1,150	49.0	46.2-52.0	649	67.8	62.6-73.2	501	35.8	32.8-39.1	822	35.1	32.8-37.6	478	51.8	47.2-56.7	344	24.4	21.9-27.2
2005	1,203	51.1	48.2-54.1	641	67.2	62.0-72.6	562	40.4	37.1-43.9	767	32.7	30.4-35.1	449	47.2	42.9-51.8	318	23.0	20.5-25.7
2006	1,143	49.0	46.2-51.9	625	66.0	60.8-71.4	518	37.4	34.3-40.8	738	31.6	29.4-34.0	419	44.7	40.5-49.2	319	22.7	20.3-25.4
2007	1,210	51.4	48.6-54.4	642	67.3	62.1-72.8	568	41.0	37.7-44.6	812	34.6	32.3-37.1	440	46.8	42.5-51.5	372	26.7	24.0-29.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Lung and bronchus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	1,165	48.8	46.0-51.7	630	65.0	60.0-70.4	535	38.3	35.1-41.7	792	33.2	30.9-35.6	436	45.7	41.4-50.2	356	25.0	22.5-27.8
2009	1,217	51.0	48.1-53.9	644	66.3	61.2-71.7	573	40.8	37.5-44.3	731	30.7	28.5-33.0	402	41.9	37.8-46.2	329	23.2	20.7-25.9
2010	1,198	49.2	46.4-52.1	643	64.4	59.4-69.7	555	38.5	35.3-41.9	806	33.1	30.8-35.5	427	43.5	39.4-47.9	379	26.2	23.6-29.0
2011	1,216	49.0	46.3-51.9	657	64.8	59.8-70.0	559	38.3	35.2-41.7	810	32.5	30.2-34.8	432	43.0	39.0-47.4	378	25.2	22.7-28.0
2012	1,288	50.9	48.2-53.8	684	65.5	60.6-70.7	604	41.3	38.1-44.8	845	33.6	31.3-35.9	491	47.8	43.6-52.3	354	24.0	21.5-26.7
2013	1,236	48.1	45.4-50.9	649	61.2	56.4-66.2	587	39.1	36.0-42.5	836	32.5	30.3-34.8	448	43.2	39.2-47.5	388	25.4	22.9-28.2
2014	1,229	46.5	43.9-49.2	671	61.3	56.6-66.2	558	36.3	33.3-39.5	745	28.4	26.3-30.5	420	39.6	35.9-43.7	325	20.7	18.5-23.1
2015	1,238	45.9	43.3-48.6	671	59.8	55.2-64.6	567	36.4	33.4-39.6	713	26.7	24.7-28.7	411	37.7	34.0-41.6	302	19.1	17.0-21.4
2016	1,255	45.5	43.0-48.2	687	59.9	55.4-64.6	568	35.6	32.7-38.8	665	24.2	22.4-26.2	392	35.0	31.5-38.7	273	16.9	14.9-19.1
2017	1,297	46.7	44.1-49.4	703	60.7	56.2-65.4	594	37.0	34.0-40.2	696	25.0	23.2-27.0	380	33.3	29.9-36.9	316	19.3	17.2-21.6
2018	1,234	43.7	41.3-46.3	623	52.7	48.5-57.1	611	37.4	34.5-40.6	643	22.6	20.9-24.5	347	29.4	26.4-32.8	296	17.9	15.9-20.1
2019	1,226	42.1	39.7-44.6	617	50.3	46.4-54.6	609	36.6	33.7-39.7	583	20.0	18.4-21.8	321	26.9	24.0-30.0	262	15.4	13.6-17.5
2020	1,033	35.5	33.3-37.8	508	41.5	37.9-45.3	525	31.2	28.5-34.0	565	19.4	17.8-21.0	307	25.5	22.7-28.6	258	15.1	13.3-17.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Melanoma of the skin**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	101	4.3	3.5-5.3	45	4.5	3.3-6.2	56	4.2	3.1-5.5	45	2.1	1.5-2.8	31	3.3	2.2-4.8	14	1.2	0.6-2.1
1977	106	4.9	4.0-5.9	51	5.7	4.2-7.6	55	4.2	3.1-5.5	48	2.2	1.6-2.9	18	2.1	1.2-3.4	30	2.3	1.5-3.3
1978	94	4.5	3.6-5.5	49	5.8	4.2-7.7	45	3.6	2.6-4.8	44	1.9	1.4-2.6	24	2.7	1.7-4.1	20	1.4	0.9-2.2
1979	108	5.1	4.1-6.2	50	5.4	3.9-7.1	58	4.9	3.7-6.3	43	2.1	1.5-2.8	25	3.0	1.9-4.4	18	1.4	0.8-2.2
1980	116	5.2	4.3-6.3	54	5.5	4.1-7.3	62	5.0	3.8-6.4	47	2.2	1.6-3.0	25	2.8	1.7-4.1	22	1.8	1.1-2.8
1981	90	4.2	3.4-5.2	40	4.4	3.1-6.0	50	4.2	3.1-5.6	43	2.0	1.5-2.7	24	2.7	1.7-4.1	19	1.5	0.9-2.4
1982	94	4.6	3.7-5.6	47	5.5	4.0-7.4	47	3.9	2.8-5.2	38	1.8	1.2-2.4	22	2.4	1.5-3.8	16	1.2	0.7-2.1
1983	113	5.2	4.3-6.3	51	5.7	4.2-7.6	62	5.0	3.8-6.4	39	1.8	1.3-2.5	18	2.1	1.2-3.3	21	1.7	1.1-2.7
1984	131	5.9	4.9-7.1	65	6.7	5.1-8.6	66	5.4	4.1-6.9	45	2.0	1.5-2.7	25	2.7	1.7-4.0	20	1.5	0.9-2.4
1985	136	6.2	5.2-7.4	66	7.1	5.5-9.1	70	5.6	4.4-7.1	55	2.5	1.9-3.3	30	3.4	2.3-4.9	25	1.9	1.2-2.8
1986	133	6.1	5.1-7.3	64	7.0	5.4-9.0	69	5.6	4.3-7.1	43	1.9	1.4-2.6	25	2.6	1.7-3.8	18	1.4	0.8-2.2
1987	124	5.7	4.8-6.9	62	6.9	5.3-8.9	62	5.0	3.8-6.4	40	1.8	1.3-2.4	21	2.3	1.4-3.5	19	1.4	0.8-2.2
1988	104	4.8	3.9-5.8	52	5.8	4.3-7.7	52	4.2	3.1-5.5	31	1.5	1.0-2.1	13	1.6	0.8-2.7	18	1.4	0.9-2.3
1989	115	5.5	4.5-6.6	50	5.8	4.3-7.7	65	5.2	4.0-6.6	38	1.8	1.3-2.5	22	2.6	1.6-3.9	16	1.3	0.7-2.1
1990	104	4.8	3.9-5.8	57	6.3	4.7-8.2	47	3.7	2.7-4.9	36	1.7	1.2-2.3	22	2.3	1.4-3.5	14	1.2	0.6-2.0
1991	109	5.1	4.2-6.2	57	7.0	5.3-9.2	52	4.0	3.0-5.3	30	1.4	1.0-2.1	16	1.9	1.1-3.2	14	1.1	0.6-1.9
1992	75	3.4	2.7-4.3	35	4.0	2.7-5.5	40	3.0	2.1-4.1	23	1.0	0.7-1.6	12	1.4	0.7-2.4	11	0.8	0.4-1.5
1993	102	4.7	3.8-5.7	52	6.2	4.6-8.2	50	3.8	2.8-5.0	29	1.4	0.9-2.0	18	2.3	1.3-3.6	11	0.9	0.4-1.6
1994	124	5.7	4.7-6.8	65	7.4	5.7-9.5	59	4.6	3.5-6.0	44	2.0	1.5-2.7	32	3.7	2.5-5.2	12	0.9	0.5-1.6
1995	125	5.8	4.9-7.0	62	7.0	5.3-8.9	63	5.0	3.9-6.5	28	1.3	0.9-1.9	12	1.4	0.7-2.4	16	1.2	0.7-2.0
1996	120	5.5	4.5-6.6	60	6.5	5.0-8.4	60	4.7	3.6-6.1	27	1.3	0.8-1.8	18	2.1	1.2-3.3	9	0.7	0.3-1.4
1997	144	6.7	5.6-7.9	66	7.9	6.0-10.0	78	6.1	4.8-7.6	34	1.6	1.1-2.2	17	2.0	1.1-3.1	17	1.3	0.8-2.1
1998	136	6.1	5.2-7.3	57	6.2	4.6-8.0	79	6.0	4.8-7.5	34	1.6	1.1-2.2	15	1.7	0.9-2.8	19	1.4	0.9-2.3
1999	143	6.3	5.3-7.4	83	8.9	7.0-11.0	60	4.5	3.4-5.8	27	1.2	0.8-1.8	13	1.4	0.7-2.3	14	1.0	0.6-1.8
2000	154	6.7	5.7-7.8	86	9.3	7.4-11.5	68	5.0	3.9-6.4	37	1.6	1.2-2.3	23	2.6	1.6-3.8	14	1.0	0.6-1.7
2001	149	6.5	5.5-7.6	80	8.6	6.8-10.7	69	5.1	4.0-6.5	32	1.4	1.0-2.0	16	1.7	1.0-2.8	16	1.1	0.6-1.8
2002	155	6.6	5.6-7.8	73	7.7	6.0-9.7	82	6.0	4.8-7.5	34	1.5	1.0-2.1	15	1.6	0.9-2.7	19	1.4	0.8-2.2
2003	138	5.9	4.9-6.9	62	6.2	4.8-8.0	76	5.6	4.4-7.1	30	1.3	0.9-1.8	13	1.4	0.7-2.3	17	1.2	0.7-2.0
2004	169	7.1	6.0-8.2	82	8.2	6.5-10.2	87	6.3	5.0-7.8	25	1.0	0.7-1.5	11	1.1	0.6-2.0	14	0.9	0.5-1.6
2005	197	8.3	7.2-9.5	109	11.2	9.2-13.6	88	6.5	5.2-8.0	28	1.2	0.8-1.7	13	1.5	0.8-2.6	15	1.1	0.6-1.8
2006	249	10.4	9.1-11.7	136	13.7	11.4-16.2	113	8.1	6.7-9.8	38	1.6	1.2-2.2	21	2.2	1.3-3.3	17	1.2	0.7-2.0
2007	212	8.7	7.5-9.9	106	10.5	8.6-12.8	106	7.6	6.2-9.2	35	1.5	1.0-2.0	13	1.4	0.8-2.5	22	1.5	1.0-2.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Melanoma of the skin**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	223	9.2	8.0-10.5	129	12.9	10.7-15.4	94	6.9	5.6-8.5	29	1.2	0.8-1.8	20	2.0	1.2-3.1	9	0.6	0.3-1.2
2009	257	10.6	9.3-12.0	142	14.4	12.1-17.0	115	8.2	6.8-9.9	31	1.3	0.9-1.8	18	1.8	1.1-2.9	13	0.9	0.5-1.6
2010	206	8.3	7.2-9.6	113	11.1	9.1-13.4	93	6.6	5.3-8.1	36	1.4	1.0-2.0	18	1.7	1.0-2.7	18	1.3	0.8-2.0
2011	233	9.2	8.0-10.4	120	11.0	9.1-13.2	113	8.1	6.6-9.8	31	1.3	0.8-1.8	16	1.6	0.9-2.6	15	1.1	0.6-1.8
2012	219	8.7	7.6-9.9	110	10.5	8.5-12.6	109	7.6	6.2-9.2	40	1.6	1.1-2.1	26	2.5	1.6-3.7	14	0.9	0.5-1.6
2013	242	9.2	8.0-10.4	132	11.8	9.8-14.0	110	7.4	6.1-9.0	31	1.1	0.8-1.6	20	1.7	1.0-2.7	11	0.8	0.4-1.4
2014	220	8.4	7.3-9.6	106	9.6	7.9-11.7	114	7.7	6.3-9.2	21	0.8	0.5-1.2	17	1.6	0.9-2.5	4	0.3	0.1-0.7
2015	223	8.4	7.3-9.6	112	10.0	8.2-12.0	111	7.4	6.0-8.9	21	0.8	0.5-1.3	12	1.1	0.6-1.9	9	0.7	0.3-1.3
2016	239	8.9	7.8-10.1	125	11.2	9.3-13.4	114	7.3	6.0-8.8	19	0.7	0.4-1.1	11	1.2	0.6-2.1	8	0.5	0.2-1.0
2017	243	9.0	7.9-10.3	135	11.7	9.8-13.9	108	7.3	5.9-8.8	25	0.9	0.6-1.3	14	1.2	0.6-2.0	11	0.6	0.3-1.1
2018	230	8.2	7.2-9.4	122	10.3	8.5-12.3	108	6.8	5.5-8.3	20	0.7	0.4-1.1	11	1.0	0.5-1.7	9	0.5	0.2-1.1
2019	240	8.6	7.6-9.8	127	10.6	8.8-12.7	113	7.3	6.0-8.9	22	0.8	0.5-1.2	11	0.9	0.5-1.7	11	0.6	0.3-1.2
2020	199	7.2	6.2-8.3	102	8.5	6.9-10.3	97	6.3	5.1-7.7	22	0.8	0.5-1.2	14	1.2	0.6-2.0	8	0.6	0.2-1.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Kings (Brooklyn)

Female breast

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,234	97.7	92.1-103.5	464	36.5	33.1-40.1
1977	1,056	85.7	80.5-91.2	455	35.8	32.5-39.4
1978	1,190	99.0	93.3-105.0	489	39.7	36.1-43.5
1979	1,150	94.8	89.2-100.6	430	35.0	31.6-38.5
1980	1,161	96.8	91.1-102.7	474	38.7	35.2-42.4
1981	1,212	101.5	95.7-107.6	481	39.0	35.5-42.7
1982	1,178	97.5	91.9-103.4	437	35.3	32.0-38.9
1983	1,173	97.1	91.4-103.0	447	36.5	33.1-40.1
1984	1,254	103.3	97.5-109.4	445	36.5	33.1-40.1
1985	1,265	103.6	97.9-109.7	411	33.1	29.9-36.5
1986	1,210	97.5	92.0-103.3	456	36.8	33.4-40.4
1987	1,195	96.6	91.1-102.4	410	32.7	29.6-36.1
1988	1,292	104.8	99.0-110.8	445	35.8	32.5-39.3
1989	1,320	107.5	101.7-113.6	413	32.9	29.7-36.2
1990	1,394	114.6	108.6-120.9	453	36.9	33.6-40.5
1991	1,409	115.6	109.5-121.9	395	31.7	28.6-35.0
1992	1,426	117.7	111.6-124.1	438	35.4	32.2-38.9
1993	1,387	113.2	107.3-119.4	427	34.2	31.0-37.6
1994	1,511	123.0	116.8-129.4	411	33.0	29.9-36.4
1995	1,483	119.9	113.8-126.2	483	38.9	35.5-42.5
1996	1,619	130.4	124.1-136.9	462	36.6	33.4-40.2
1997	1,501	120.2	114.2-126.5	389	30.9	27.9-34.1
1998	1,611	127.4	121.3-133.8	406	31.8	28.8-35.1
1999	1,513	117.1	111.3-123.2	382	28.9	26.0-31.9
2000	1,483	112.3	106.6-118.2	373	27.6	24.9-30.6
2001	1,455	109.2	103.6-114.9	399	29.2	26.4-32.3
2002	1,479	110.0	104.5-115.8	352	25.6	23.0-28.5
2003	1,462	108.5	103.0-114.2	364	26.4	23.8-29.3
2004	1,468	108.9	103.4-114.7	356	25.7	23.1-28.5
2005	1,367	100.7	95.4-106.2	375	27.2	24.5-30.1
2006	1,475	109.9	104.3-115.7	329	23.9	21.4-26.7
2007	1,498	110.8	105.2-116.6	336	24.0	21.5-26.7
2008	1,474	107.8	102.3-113.5	310	22.0	19.6-24.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Kings (Brooklyn)

Female breast

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	1,571	114.2	108.5-120.0	318	22.6	20.1-25.2
2010	1,522	109.8	104.2-115.5	327	22.3	19.9-24.9
2011	1,721	122.2	116.5-128.2	340	23.7	21.2-26.4
2012	1,632	114.1	108.6-119.9	339	23.3	20.9-26.0
2013	1,783	123.3	117.5-129.2	326	21.1	18.8-23.5
2014	1,684	115.0	109.5-120.8	330	21.6	19.2-24.1
2015	1,808	121.9	116.3-127.8	318	20.8	18.5-23.2
2016	1,743	117.2	111.6-123.0	316	20.2	18.0-22.6
2017	1,840	122.9	117.3-128.8	311	19.8	17.6-22.2
2018	1,902	126.9	121.1-132.9	350	21.5	19.2-23.9
2019	2,128	140.2	134.2-146.5	325	20.0	17.8-22.4
2020	1,681	112.2	106.8-117.9	302	18.5	16.4-20.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Kings (Brooklyn)

Cervix uteri

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	213	18.2	15.7-20.9	72	6.1	4.7-7.8
1977	188	16.0	13.7-18.6	76	6.4	5.0-8.0
1978	198	18.0	15.5-20.8	61	5.1	3.9-6.6
1979	209	18.5	16.0-21.3	57	5.0	3.8-6.6
1980	185	16.9	14.4-19.6	69	5.9	4.6-7.5
1981	180	15.2	13.0-17.7	60	5.3	4.0-6.9
1982	169	15.2	13.0-17.8	58	5.1	3.9-6.7
1983	183	16.1	13.8-18.7	37	3.1	2.2-4.4
1984	187	16.2	13.9-18.8	49	4.3	3.1-5.7
1985	168	14.2	12.1-16.6	53	4.5	3.3-5.9
1986	161	13.7	11.6-16.1	36	3.2	2.2-4.4
1987	171	14.7	12.5-17.2	54	4.5	3.4-5.9
1988	160	13.7	11.6-16.1	70	5.8	4.5-7.4
1989	176	14.8	12.7-17.3	38	3.1	2.2-4.3
1990	196	16.8	14.5-19.4	47	4.3	3.1-5.7
1991	178	15.2	13.0-17.6	48	4.0	2.9-5.3
1992	195	16.1	13.9-18.6	55	4.5	3.3-5.8
1993	171	14.4	12.3-16.7	60	4.9	3.7-6.3
1994	197	16.2	14.0-18.7	48	3.9	2.9-5.2
1995	198	16.2	14.0-18.6	51	4.3	3.2-5.6
1996	194	15.8	13.7-18.2	55	4.5	3.4-5.9
1997	179	14.4	12.4-16.7	51	4.1	3.0-5.4
1998	183	14.6	12.5-16.9	45	3.6	2.6-4.8
1999	166	12.9	11.0-15.0	52	4.0	3.0-5.3
2000	205	15.8	13.7-18.1	52	3.9	2.9-5.2
2001	135	10.4	8.7-12.3	41	3.1	2.2-4.2
2002	178	13.6	11.7-15.8	47	3.5	2.6-4.7
2003	150	11.2	9.4-13.1	46	3.4	2.5-4.5
2004	128	9.5	8.0-11.4	44	3.3	2.4-4.4
2005	161	12.1	10.3-14.2	46	3.4	2.5-4.5
2006	155	11.6	9.9-13.6	45	3.4	2.4-4.5
2007	149	11.2	9.5-13.2	52	3.8	2.8-5.0
2008	147	11.0	9.3-13.0	42	3.1	2.2-4.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Cervix uteri**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	147	11.0	9.3-12.9	53	3.8	2.9-5.0
2010	145	10.7	9.0-12.6	35	2.4	1.7-3.4
2011	133	9.8	8.2-11.6	47	3.4	2.5-4.5
2012	142	10.1	8.4-11.9	45	3.1	2.3-4.2
2013	143	9.8	8.3-11.6	43	2.9	2.1-3.9
2014	154	11.0	9.3-12.9	44	3.0	2.2-4.0
2015	121	8.2	6.8-9.9	48	3.1	2.2-4.1
2016	157	11.1	9.4-13.0	39	2.7	1.9-3.7
2017	141	9.5	8.0-11.3	38	2.6	1.8-3.6
2018	130	9.2	7.6-10.9	40	2.6	1.9-3.6
2019	147	9.9	8.4-11.7	27	1.7	1.1-2.5
2020	97	6.6	5.3-8.1	29	1.9	1.3-2.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Kings (Brooklyn)

Corpus uteris and uterus, not otherwise specified

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	259	19.7	17.3-22.4	94	7.3	5.8-9.0
1977	233	17.7	15.4-20.2	56	4.1	3.1-5.4
1978	241	18.4	16.1-20.9	64	4.8	3.7-6.2
1979	254	20.3	17.8-23.0	59	4.4	3.3-5.7
1980	319	24.7	22.0-27.6	63	4.8	3.6-6.1
1981	248	19.3	16.9-21.9	69	5.5	4.2-6.9
1982	279	21.9	19.3-24.7	60	4.4	3.4-5.7
1983	287	22.3	19.7-25.1	81	6.0	4.8-7.5
1984	271	21.1	18.6-23.9	75	5.6	4.4-7.1
1985	271	20.8	18.3-23.5	64	4.8	3.7-6.2
1986	275	21.4	18.9-24.1	55	4.3	3.3-5.7
1987	237	18.5	16.1-21.0	67	5.1	4.0-6.6
1988	272	20.8	18.3-23.4	64	4.6	3.6-6.0
1989	272	21.2	18.7-23.9	63	4.8	3.6-6.1
1990	248	19.9	17.5-22.6	71	5.4	4.2-6.8
1991	286	22.8	20.2-25.7	56	4.5	3.4-5.8
1992	293	23.9	21.2-26.8	67	5.3	4.1-6.8
1993	270	21.6	19.1-24.4	60	4.7	3.6-6.0
1994	314	25.0	22.3-27.9	85	6.7	5.3-8.3
1995	298	23.8	21.2-26.7	79	6.2	4.9-7.7
1996	319	25.4	22.7-28.4	81	6.3	5.0-7.8
1997	334	26.3	23.6-29.3	84	6.5	5.2-8.1
1998	356	28.0	25.1-31.0	82	6.3	5.0-7.9
1999	351	27.0	24.2-29.9	93	7.0	5.6-8.6
2000	359	27.1	24.4-30.1	75	5.5	4.3-6.9
2001	388	29.0	26.1-32.0	72	5.3	4.1-6.6
2002	357	26.4	23.8-29.3	76	5.6	4.4-7.0
2003	392	28.7	25.9-31.7	89	6.5	5.2-7.9
2004	362	26.5	23.8-29.4	76	5.5	4.3-6.8
2005	400	28.9	26.1-31.9	92	6.6	5.3-8.1
2006	375	27.2	24.5-30.1	79	5.7	4.5-7.1
2007	398	28.9	26.1-31.9	79	5.7	4.5-7.1
2008	492	34.9	31.8-38.1	100	7.1	5.7-8.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Kings (Brooklyn)

Corpus uteris and uterus, not otherwise specified

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	447	31.4	28.6-34.5	87	6.1	4.9-7.6
2010	429	30.1	27.3-33.2	108	7.5	6.2-9.1
2011	456	31.5	28.6-34.6	114	7.7	6.3-9.3
2012	448	30.0	27.3-33.0	100	6.7	5.4-8.1
2013	478	31.2	28.4-34.2	99	6.3	5.1-7.7
2014	528	34.2	31.3-37.3	113	7.1	5.9-8.6
2015	549	35.2	32.2-38.3	130	8.1	6.7-9.7
2016	558	35.8	32.8-38.9	123	7.6	6.3-9.1
2017	507	31.9	29.2-34.9	138	8.4	7.0-10.0
2018	503	31.8	29.0-34.7	128	7.6	6.3-9.1
2019	516	31.8	29.0-34.7	121	7.3	6.1-8.8
2020	442	27.5	25.0-30.3	136	8.0	6.7-9.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Kings (Brooklyn)

Ovary

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	181	13.6	11.7-15.8	132	10.0	8.3-11.9
1977	167	13.2	11.2-15.4	138	10.5	8.8-12.5
1978	176	14.2	12.1-16.5	115	9.3	7.7-11.3
1979	193	15.7	13.5-18.1	109	8.5	7.0-10.4
1980	197	16.1	13.9-18.6	131	10.3	8.6-12.3
1981	183	15.0	12.8-17.4	116	9.1	7.5-11.0
1982	182	14.7	12.6-17.1	104	8.4	6.8-10.2
1983	201	16.0	13.8-18.4	117	9.1	7.5-11.0
1984	190	14.8	12.7-17.1	116	9.3	7.6-11.2
1985	195	15.7	13.5-18.1	113	9.0	7.4-10.8
1986	148	11.8	9.9-13.9	105	8.1	6.6-9.8
1987	187	14.9	12.8-17.2	100	7.7	6.3-9.4
1988	172	13.9	11.8-16.2	114	8.9	7.3-10.7
1989	178	14.1	12.1-16.4	105	8.3	6.8-10.1
1990	189	15.3	13.2-17.7	121	9.6	7.9-11.4
1991	199	16.3	14.1-18.7	125	10.0	8.3-12.0
1992	194	15.7	13.5-18.1	135	10.9	9.1-12.9
1993	189	15.4	13.2-17.7	111	8.7	7.2-10.5
1994	211	16.9	14.7-19.3	104	8.2	6.7-9.9
1995	229	18.4	16.1-21.0	120	9.4	7.8-11.3
1996	163	12.8	10.9-15.0	115	9.1	7.5-10.9
1997	205	16.4	14.2-18.8	131	10.3	8.6-12.2
1998	183	14.4	12.4-16.6	106	8.2	6.7-10.0
1999	179	13.8	11.8-16.0	108	8.2	6.7-9.9
2000	165	12.4	10.6-14.4	112	8.2	6.8-9.9
2001	162	12.1	10.3-14.1	89	6.5	5.2-8.0
2002	174	12.9	11.0-14.9	100	7.3	6.0-8.9
2003	159	11.6	9.9-13.6	110	7.9	6.5-9.6
2004	175	12.9	11.0-15.0	104	7.4	6.0-9.0
2005	151	11.1	9.4-13.1	112	8.2	6.7-9.9
2006	158	11.7	10.0-13.7	95	6.9	5.6-8.4
2007	176	12.7	10.9-14.8	92	6.5	5.2-8.0
2008	180	12.9	11.1-15.0	100	6.9	5.6-8.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Kings (Brooklyn)

Ovary

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	170	12.3	10.5-14.3	96	6.6	5.3-8.1
2010	165	11.7	9.9-13.6	88	6.1	4.9-7.6
2011	166	11.7	10.0-13.7	100	6.8	5.5-8.3
2012	170	11.8	10.0-13.7	114	7.7	6.3-9.3
2013	186	12.6	10.8-14.6	111	7.3	6.0-8.8
2014	177	11.6	9.9-13.5	93	5.8	4.7-7.2
2015	173	11.0	9.4-12.9	111	7.1	5.8-8.6
2016	159	10.6	9.0-12.4	112	7.1	5.8-8.5
2017	194	12.8	11.0-14.8	113	7.1	5.8-8.6
2018	184	12.2	10.4-14.1	121	7.6	6.3-9.1
2019	149	9.8	8.3-11.6	92	5.5	4.4-6.8
2020	147	9.6	8.1-11.4	93	5.6	4.5-6.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Kings (Brooklyn)

Prostate

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	555	68.8	62.8-75.1	195	26.2	22.4-30.5
1977	533	64.3	58.6-70.3	210	27.2	23.4-31.4
1978	594	73.9	67.7-80.4	235	32.0	27.8-36.6
1979	628	78.9	72.6-85.7	231	30.8	26.7-35.2
1980	750	93.0	86.1-100.2	232	31.8	27.6-36.3
1981	683	84.7	78.2-91.6	221	29.2	25.4-33.5
1982	676	84.3	77.8-91.2	204	28.0	24.1-32.3
1983	695	85.4	79.0-92.3	228	30.8	26.7-35.2
1984	732	88.7	82.2-95.6	249	33.5	29.4-38.1
1985	750	90.6	83.9-97.5	226	29.3	25.5-33.6
1986	776	94.7	87.9-101.8	233	29.9	26.0-34.1
1987	741	89.8	83.3-96.8	248	32.2	28.2-36.6
1988	753	93.4	86.7-100.5	260	34.4	30.2-38.9
1989	798	98.6	91.7-105.9	222	30.1	26.2-34.4
1990	906	112.5	105.1-120.3	242	32.7	28.6-37.1
1991	953	116.7	109.2-124.5	237	32.0	28.0-36.5
1992	1,168	140.1	132.0-148.5	252	33.4	29.3-37.8
1993	1,191	141.8	133.7-150.2	262	35.0	30.8-39.6
1994	1,349	159.1	150.6-168.0	261	35.8	31.5-40.5
1995	1,329	155.6	147.2-164.3	289	38.6	34.2-43.4
1996	1,308	153.6	145.2-162.2	270	36.0	31.8-40.6
1997	1,426	166.5	157.8-175.4	246	33.0	28.9-37.4
1998	1,412	161.5	153.1-170.2	243	31.3	27.4-35.5
1999	1,514	168.8	160.3-177.6	237	29.9	26.1-33.9
2000	1,795	194.9	185.9-204.2	255	31.7	27.9-35.8
2001	1,694	182.4	173.7-191.4	236	28.9	25.3-32.8
2002	1,573	166.4	158.2-174.9	237	28.8	25.2-32.8
2003	1,600	168.3	160.1-176.9	234	28.1	24.6-32.0
2004	1,557	160.6	152.6-168.9	218	26.6	23.2-30.4
2005	1,520	155.3	147.5-163.4	216	25.4	22.1-29.1
2006	1,611	164.0	156.0-172.3	180	20.8	17.9-24.1
2007	1,758	176.1	167.8-184.6	203	23.5	20.3-27.0
2008	1,783	174.9	166.7-183.3	190	22.3	19.2-25.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Kings (Brooklyn)

Prostate

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	1,695	162.8	155.0-170.8	208	23.7	20.5-27.1
2010	1,548	144.3	137.0-151.8	219	24.0	20.9-27.4
2011	1,831	166.4	158.7-174.4	236	25.6	22.4-29.1
2012	1,565	139.0	132.0-146.2	212	22.5	19.6-25.8
2013	1,589	138.1	131.2-145.2	207	21.9	19.0-25.1
2014	1,432	120.8	114.5-127.4	218	21.5	18.7-24.6
2015	1,482	121.8	115.6-128.4	207	20.1	17.4-23.1
2016	1,531	125.3	119.0-131.9	194	18.3	15.8-21.1
2017	1,704	136.2	129.7-143.0	249	23.2	20.4-26.3
2018	1,656	130.6	124.3-137.2	186	16.7	14.3-19.3
2019	1,748	135.9	129.4-142.5	190	17.3	14.9-19.9
2020	1,515	116.8	110.9-123.0	153	13.3	11.3-15.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Kings (Brooklyn)

Testis

Year of diagnosis	Incidence		
	Cases	Rate per 100,000	95% CI
1976	21	1.9	1.1-2.9
1977	15	1.2	0.7-2.0
1978	20	1.8	1.1-2.8
1979	30	2.9	1.9-4.2
1980	32	3.1	2.1-4.5
1981	30	3.1	2.0-4.5
1982	29	2.9	1.9-4.2
1983	25	2.1	1.3-3.2
1984	23	2.2	1.3-3.3
1985	29	2.7	1.8-4.0
1986	38	3.3	2.3-4.5
1987	28	2.4	1.6-3.5
1988	30	2.8	1.9-4.1
1989	32	2.8	1.9-4.0
1990	36	3.3	2.3-4.7
1991	33	2.9	1.9-4.1
1992	30	2.5	1.7-3.6
1993	31	2.7	1.8-4.0
1994	29	2.5	1.6-3.6
1995	33	2.8	1.9-3.9
1996	37	3.4	2.4-4.7
1997	26	2.2	1.4-3.3
1998	32	2.9	1.9-4.1
1999	43	3.5	2.5-4.8
2000	46	3.9	2.8-5.2
2001	35	2.9	2.0-4.1
2002	28	2.3	1.5-3.4
2003	39	3.3	2.4-4.6
2004	41	3.4	2.4-4.7
2005	51	4.2	3.1-5.5
2006	58	4.9	3.7-6.4
2007	48	4.0	3.0-5.4
2008	56	4.6	3.4-6.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

Mortality data not provided due to small number of deaths per year.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Testis**

Year of diagnosis	Incidence		
	Cases	Rate per 100,000	95% CI
2009	51	4.2	3.1-5.6
2010	54	4.3	3.2-5.6
2011	60	4.9	3.7-6.3
2012	57	4.2	3.2-5.5
2013	58	4.5	3.4-5.8
2014	65	5.0	3.9-6.5
2015	63	4.7	3.6-6.1
2016	55	4.0	3.0-5.3
2017	73	5.5	4.3-7.0
2018	67	5.4	4.1-6.9
2019	81	6.3	4.9-7.8
2020	74	5.8	4.6-7.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

Mortality data not provided due to small number of deaths per year.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Urinary bladder (incl. in situ)**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	455	20.8	18.8-22.8	338	38.2	34.0-42.8	117	9.1	7.5-11.0	115	5.9	4.8-7.2	85	11.0	8.6-13.7	30	2.7	1.8-3.9
1977	389	17.6	15.8-19.5	274	30.2	26.6-34.3	115	9.0	7.4-10.9	124	6.1	5.1-7.3	87	11.1	8.8-13.9	37	3.0	2.1-4.2
1978	421	19.9	17.9-21.9	304	36.8	32.5-41.4	117	9.0	7.4-10.9	107	5.2	4.2-6.4	73	8.9	6.9-11.3	34	2.9	2.0-4.1
1979	430	20.5	18.5-22.6	297	35.4	31.3-39.9	133	10.8	9.0-12.9	109	5.6	4.6-6.9	75	10.2	7.9-12.8	34	2.9	2.0-4.1
1980	514	24.1	22.0-26.3	391	46.0	41.4-51.0	123	9.8	8.1-11.7	113	5.5	4.5-6.6	78	10.0	7.8-12.5	35	2.8	1.9-3.9
1981	461	21.7	19.7-23.8	339	41.0	36.6-45.8	122	9.5	7.9-11.4	120	5.8	4.8-7.0	76	9.8	7.6-12.3	44	3.4	2.5-4.6
1982	392	18.4	16.6-20.4	274	32.4	28.5-36.6	118	9.5	7.8-11.4	105	5.3	4.3-6.4	80	10.9	8.5-13.6	25	2.1	1.3-3.1
1983	481	22.7	20.6-24.8	338	42.2	37.6-47.1	143	10.7	9.0-12.7	106	5.1	4.1-6.2	72	9.4	7.3-12.0	34	2.6	1.8-3.6
1984	450	20.7	18.8-22.7	326	38.1	33.9-42.6	124	9.6	8.0-11.5	105	5.1	4.1-6.2	66	8.4	6.4-10.7	39	3.1	2.2-4.2
1985	400	18.4	16.6-20.3	274	33.0	29.0-37.3	126	9.4	7.9-11.3	98	4.9	4.0-6.0	61	8.2	6.2-10.7	37	3.0	2.1-4.2
1986	426	19.8	17.9-21.8	307	38.0	33.7-42.6	119	8.9	7.4-10.7	91	4.4	3.5-5.4	55	7.3	5.4-9.5	36	2.8	1.9-3.9
1987	437	20.1	18.2-22.1	323	39.1	34.8-43.7	114	8.6	7.1-10.4	112	5.3	4.3-6.4	82	10.7	8.5-13.4	30	2.2	1.5-3.2
1988	438	20.1	18.3-22.1	305	36.5	32.4-40.9	133	10.1	8.5-12.0	96	4.5	3.7-5.5	55	6.9	5.2-9.1	41	3.1	2.2-4.2
1989	395	18.7	16.8-20.6	289	35.7	31.6-40.1	106	8.3	6.8-10.0	97	4.7	3.8-5.8	59	7.8	5.9-10.1	38	3.0	2.1-4.1
1990	409	19.3	17.4-21.2	292	36.4	32.2-40.9	117	8.8	7.2-10.5	100	5.0	4.0-6.0	71	9.5	7.4-12.0	29	2.4	1.6-3.5
1991	389	18.3	16.5-20.2	281	34.5	30.5-38.9	108	8.4	6.9-10.2	67	3.2	2.5-4.1	44	5.7	4.1-7.7	23	1.8	1.1-2.7
1992	396	18.7	16.9-20.7	286	35.2	31.2-39.6	110	8.8	7.2-10.6	72	3.5	2.7-4.4	47	6.5	4.8-8.7	25	1.9	1.2-2.7
1993	379	17.8	16.0-19.7	281	34.5	30.5-38.8	98	7.6	6.1-9.2	96	4.6	3.7-5.6	66	8.5	6.5-10.8	30	2.4	1.6-3.4
1994	415	19.5	17.7-21.5	294	35.8	31.7-40.2	121	9.4	7.8-11.2	95	4.6	3.7-5.6	65	8.7	6.7-11.1	30	2.3	1.5-3.3
1995	409	19.3	17.5-21.3	285	35.4	31.4-39.8	124	9.6	8.0-11.4	112	5.4	4.4-6.4	72	9.6	7.5-12.1	40	3.1	2.2-4.2
1996	358	16.9	15.2-18.8	264	32.5	28.6-36.7	94	7.2	5.8-8.8	93	4.4	3.6-5.4	62	8.1	6.2-10.4	31	2.3	1.6-3.3
1997	413	19.3	17.5-21.2	280	33.6	29.7-37.8	133	10.2	8.6-12.1	87	4.2	3.3-5.1	50	6.7	5.0-8.9	37	2.8	2.0-3.8
1998	433	20.0	18.1-21.9	312	37.0	32.9-41.4	121	9.2	7.6-11.0	71	3.3	2.6-4.2	41	5.3	3.8-7.1	30	2.2	1.5-3.1
1999	401	18.1	16.3-19.9	292	34.1	30.3-38.3	109	8.0	6.6-9.7	86	3.9	3.2-4.9	56	7.2	5.4-9.4	30	2.2	1.5-3.1
2000	414	18.3	16.6-20.2	304	35.1	31.2-39.3	110	8.1	6.6-9.7	70	3.1	2.4-3.9	44	5.2	3.8-7.0	26	1.8	1.2-2.7
2001	394	17.2	15.5-19.0	277	31.2	27.6-35.1	117	8.4	6.9-10.1	92	4.0	3.2-4.9	57	6.8	5.1-8.8	35	2.4	1.7-3.4
2002	382	16.6	14.9-18.3	270	30.3	26.7-34.2	112	8.0	6.6-9.6	69	3.0	2.3-3.8	49	5.8	4.3-7.7	20	1.4	0.8-2.1
2003	375	16.2	14.6-17.9	272	30.4	26.8-34.2	103	7.3	5.9-8.9	62	2.7	2.1-3.4	42	5.1	3.7-6.9	20	1.4	0.8-2.2
2004	440	18.8	17.1-20.6	298	32.4	28.8-36.3	142	9.9	8.3-11.7	78	3.3	2.6-4.1	53	6.1	4.6-8.0	25	1.7	1.1-2.5
2005	413	17.7	16.0-19.5	280	30.2	26.7-34.0	133	9.3	7.8-11.1	69	3.0	2.4-3.8	43	5.1	3.7-6.8	26	1.8	1.2-2.7
2006	416	17.7	16.1-19.5	303	32.5	28.9-36.5	113	8.1	6.7-9.7	62	2.7	2.1-3.5	39	4.6	3.2-6.2	23	1.6	1.0-2.4
2007	382	16.5	14.9-18.2	271	29.8	26.3-33.6	111	7.9	6.5-9.6	65	2.8	2.1-3.5	39	4.7	3.3-6.4	26	1.7	1.1-2.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Urinary bladder (incl. in situ)**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	445	18.7	17.0-20.6	312	32.7	29.1-36.6	133	9.3	7.8-11.0	68	2.9	2.2-3.7	40	4.6	3.3-6.2	28	1.9	1.2-2.7
2009	413	17.4	15.7-19.1	292	31.4	27.8-35.2	121	8.4	6.9-10.1	83	3.5	2.7-4.3	50	5.9	4.4-7.7	33	2.1	1.5-3.0
2010	377	15.4	13.9-17.1	283	28.9	25.6-32.6	94	6.5	5.2-8.0	77	3.3	2.6-4.1	44	5.1	3.7-6.8	33	2.4	1.6-3.3
2011	426	17.1	15.5-18.9	304	30.9	27.5-34.7	122	8.2	6.8-9.9	96	3.9	3.1-4.7	59	6.2	4.7-8.0	37	2.5	1.7-3.4
2012	414	16.3	14.8-18.0	303	30.0	26.6-33.6	111	7.3	6.0-8.8	82	3.2	2.5-4.0	54	5.4	4.0-7.0	28	1.8	1.2-2.6
2013	410	15.8	14.3-17.4	293	28.1	24.9-31.6	117	7.4	6.1-8.9	88	3.3	2.7-4.1	53	5.3	4.0-7.0	35	2.0	1.4-2.9
2014	411	15.6	14.1-17.2	308	29.0	25.8-32.5	103	6.6	5.3-8.0	115	4.3	3.6-5.2	77	7.7	6.0-9.6	38	2.3	1.6-3.2
2015	466	17.2	15.7-18.9	355	32.3	29.0-35.9	111	7.0	5.7-8.5	83	3.0	2.4-3.8	52	5.0	3.7-6.5	31	1.8	1.2-2.6
2016	418	15.4	14.0-17.0	324	29.4	26.2-32.8	94	6.0	4.8-7.3	96	3.5	2.9-4.3	69	6.6	5.1-8.4	27	1.7	1.1-2.4
2017	452	16.5	14.9-18.1	335	29.7	26.6-33.2	117	7.4	6.1-8.9	83	3.0	2.4-3.7	61	5.7	4.3-7.3	22	1.3	0.8-2.0
2018	430	15.4	13.9-16.9	335	28.9	25.8-32.2	95	6.0	4.9-7.4	78	2.8	2.2-3.5	51	4.7	3.5-6.2	27	1.6	1.0-2.3
2019	439	15.3	13.8-16.8	310	25.7	22.9-28.8	129	7.7	6.4-9.2	80	2.7	2.2-3.4	55	4.9	3.7-6.4	25	1.4	0.9-2.1
2020	400	13.7	12.4-15.2	297	24.4	21.7-27.4	103	6.2	5.1-7.6	67	2.2	1.7-2.8	37	3.2	2.3-4.4	30	1.6	1.1-2.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Kidney and renal pelvis**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	137	6.0	5.0-7.1	96	9.9	8.0-12.2	41	3.2	2.2-4.3	77	3.5	2.7-4.4	50	5.3	3.9-7.1	27	2.1	1.4-3.1
1977	131	5.9	4.9-7.0	86	9.0	7.2-11.2	45	3.6	2.6-4.9	65	2.8	2.2-3.7	44	4.5	3.3-6.1	21	1.6	1.0-2.5
1978	151	6.7	5.6-7.9	97	10.6	8.5-13.1	54	4.0	3.0-5.2	65	2.9	2.2-3.7	46	4.8	3.5-6.6	19	1.5	0.9-2.4
1979	138	6.2	5.2-7.4	87	9.7	7.7-12.0	51	3.8	2.8-5.0	95	4.5	3.6-5.5	62	7.3	5.5-9.5	33	2.6	1.8-3.7
1980	150	6.7	5.6-7.9	88	9.3	7.4-11.6	62	4.8	3.6-6.2	63	2.9	2.2-3.7	40	4.4	3.1-6.1	23	1.8	1.1-2.7
1981	133	6.0	5.0-7.2	68	7.3	5.7-9.4	65	5.1	3.9-6.5	73	3.3	2.6-4.2	37	4.1	2.9-5.7	36	2.8	1.9-3.9
1982	186	8.6	7.4-10.0	108	12.6	10.2-15.3	78	6.1	4.8-7.7	70	3.3	2.6-4.2	44	5.3	3.8-7.1	26	2.1	1.3-3.0
1983	184	8.3	7.1-9.6	111	11.9	9.8-14.4	73	5.8	4.5-7.4	87	3.9	3.1-4.8	50	5.3	3.9-7.1	37	2.9	2.0-4.0
1984	164	7.6	6.4-8.8	101	11.1	9.0-13.5	63	5.0	3.8-6.5	74	3.4	2.7-4.3	35	3.9	2.7-5.4	39	3.1	2.2-4.3
1985	181	8.1	7.0-9.4	118	13.1	10.8-15.7	63	4.9	3.8-6.3	72	3.3	2.6-4.2	44	5.1	3.6-6.8	28	2.2	1.4-3.1
1986	178	7.9	6.7-9.1	102	11.2	9.1-13.7	76	5.6	4.4-7.0	76	3.4	2.6-4.2	46	5.3	3.8-7.1	30	2.1	1.4-3.1
1987	206	9.3	8.0-10.7	110	12.2	10.0-14.7	96	7.4	6.0-9.1	64	2.9	2.2-3.7	38	4.3	3.0-6.0	26	2.0	1.3-2.9
1988	184	8.6	7.3-9.9	114	12.6	10.4-15.2	70	5.6	4.3-7.1	57	2.6	2.0-3.4	33	3.8	2.6-5.4	24	1.9	1.2-2.8
1989	185	8.5	7.3-9.8	115	12.9	10.6-15.5	70	5.4	4.2-6.8	69	3.3	2.5-4.1	42	4.8	3.5-6.6	27	2.1	1.4-3.1
1990	198	9.3	8.0-10.7	115	13.3	10.9-16.0	83	6.6	5.3-8.2	66	3.1	2.4-4.0	37	4.6	3.2-6.4	29	2.2	1.4-3.1
1991	235	10.9	9.6-12.4	149	17.1	14.4-20.2	86	6.7	5.3-8.3	70	3.2	2.5-4.1	37	4.3	3.0-6.0	33	2.5	1.7-3.6
1992	204	9.4	8.2-10.8	115	13.1	10.8-15.8	89	6.8	5.5-8.4	55	2.6	1.9-3.4	31	3.8	2.5-5.4	24	1.9	1.2-2.8
1993	229	10.7	9.3-12.1	158	18.1	15.3-21.1	71	5.6	4.3-7.0	72	3.4	2.7-4.3	54	6.3	4.7-8.3	18	1.4	0.9-2.3
1994	225	10.4	9.1-11.9	146	16.5	13.9-19.4	79	6.2	4.9-7.8	74	3.4	2.7-4.3	48	5.7	4.1-7.5	26	2.0	1.3-3.0
1995	206	9.6	8.4-11.1	128	14.8	12.3-17.6	78	6.2	4.9-7.7	71	3.4	2.7-4.3	48	5.9	4.3-7.9	23	1.7	1.1-2.6
1996	272	12.5	11.1-14.1	165	18.7	15.9-21.8	107	8.3	6.8-10.0	71	3.3	2.6-4.2	36	4.2	2.9-5.9	35	2.7	1.9-3.8
1997	242	11.2	9.8-12.7	142	16.1	13.6-19.0	100	7.8	6.4-9.5	77	3.6	2.8-4.4	47	5.5	4.0-7.3	30	2.3	1.6-3.3
1998	250	11.4	10.0-12.9	146	15.9	13.4-18.7	104	8.1	6.6-9.8	59	2.7	2.0-3.5	42	4.8	3.5-6.5	17	1.3	0.7-2.0
1999	299	13.3	11.8-14.9	175	18.8	16.1-21.8	124	9.4	7.8-11.2	59	2.7	2.0-3.4	33	3.7	2.6-5.2	26	1.9	1.2-2.8
2000	286	12.5	11.1-14.1	159	16.8	14.3-19.6	127	9.5	7.9-11.3	53	2.3	1.7-3.0	33	3.5	2.4-5.0	20	1.5	0.9-2.3
2001	289	12.5	11.1-14.1	160	16.7	14.2-19.6	129	9.5	7.9-11.3	71	3.1	2.4-3.9	37	4.1	2.9-5.6	34	2.5	1.7-3.5
2002	272	11.6	10.3-13.1	167	17.3	14.8-20.2	105	7.7	6.3-9.3	78	3.4	2.7-4.2	42	4.6	3.3-6.2	36	2.5	1.8-3.5
2003	302	12.9	11.5-14.5	187	19.3	16.6-22.3	115	8.4	6.9-10.1	61	2.6	2.0-3.4	37	3.9	2.8-5.4	24	1.7	1.1-2.6
2004	305	12.9	11.5-14.4	197	19.9	17.2-23.0	108	7.8	6.4-9.4	69	2.9	2.3-3.7	43	4.6	3.3-6.2	26	1.9	1.2-2.8
2005	324	13.7	12.2-15.2	189	19.3	16.6-22.2	135	9.8	8.2-11.6	53	2.3	1.7-3.0	33	3.5	2.4-5.0	20	1.4	0.9-2.2
2006	323	13.8	12.3-15.4	196	20.1	17.3-23.1	127	9.3	7.8-11.1	69	3.0	2.3-3.8	42	4.7	3.4-6.3	27	2.0	1.3-2.9
2007	323	13.6	12.1-15.1	207	20.4	17.7-23.5	116	8.5	7.0-10.2	62	2.7	2.0-3.4	33	3.5	2.4-4.9	29	2.0	1.4-3.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Kidney and renal pelvis**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	358	14.9	13.4-16.6	217	21.1	18.3-24.2	141	10.3	8.6-12.1	45	1.9	1.4-2.5	29	2.9	1.9-4.2	16	1.1	0.6-1.8
2009	332	13.5	12.1-15.0	200	19.0	16.4-21.9	132	9.4	7.9-11.2	76	3.1	2.4-3.8	54	5.5	4.1-7.2	22	1.4	0.9-2.2
2010	359	14.4	13.0-16.0	231	22.0	19.2-25.1	128	9.1	7.5-10.8	85	3.4	2.7-4.2	57	5.8	4.4-7.5	28	1.9	1.2-2.7
2011	361	14.3	12.8-15.9	209	19.0	16.5-21.8	152	10.6	9.0-12.5	61	2.3	1.8-3.0	36	3.4	2.4-4.8	25	1.6	1.0-2.3
2012	360	14.2	12.8-15.8	221	20.3	17.7-23.2	139	9.7	8.1-11.4	54	2.1	1.6-2.8	31	3.1	2.1-4.4	23	1.4	0.9-2.2
2013	381	14.5	13.0-16.0	221	19.1	16.6-21.9	160	10.9	9.2-12.7	67	2.5	2.0-3.2	32	3.0	2.0-4.2	35	2.1	1.5-3.0
2014	373	14.0	12.5-15.5	243	20.9	18.3-23.7	130	8.7	7.2-10.3	72	2.8	2.2-3.5	39	3.6	2.6-5.0	33	2.2	1.5-3.1
2015	391	14.3	12.9-15.8	243	20.4	17.8-23.2	148	9.4	7.9-11.1	67	2.4	1.9-3.1	46	4.1	3.0-5.5	21	1.2	0.8-1.9
2016	406	14.6	13.2-16.1	257	20.9	18.3-23.7	149	9.8	8.3-11.6	47	1.7	1.2-2.3	37	3.2	2.3-4.5	10	0.7	0.3-1.2
2017	428	15.2	13.8-16.8	273	22.1	19.5-25.0	155	9.9	8.4-11.6	59	2.1	1.6-2.8	44	3.9	2.8-5.3	15	0.9	0.5-1.5
2018	422	15.2	13.8-16.8	279	23.0	20.4-26.0	143	9.2	7.7-10.9	58	2.0	1.5-2.6	42	3.4	2.5-4.7	16	0.9	0.5-1.5
2019	422	15.0	13.5-16.5	267	21.5	19.0-24.3	155	9.9	8.3-11.6	63	2.2	1.7-2.9	37	3.0	2.1-4.2	26	1.6	1.0-2.4
2020	342	12.4	11.1-13.8	208	17.2	14.9-19.7	134	8.3	7.0-9.9	61	2.2	1.7-2.8	40	3.4	2.4-4.6	21	1.2	0.7-1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Kings (Brooklyn)

Brain and other central nervous system

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	113	4.8	3.9-5.8	56	5.4	4.1-7.1	57	4.3	3.2-5.6	90	4.0	3.2-4.9	57	6.0	4.5-7.9	33	2.4	1.7-3.5
1977	105	4.5	3.6-5.4	58	5.6	4.2-7.3	47	3.6	2.6-4.8	96	4.1	3.3-5.1	49	4.8	3.5-6.4	47	3.6	2.7-4.9
1978	97	4.3	3.5-5.3	56	5.7	4.3-7.5	41	3.3	2.3-4.5	88	3.8	3.0-4.7	46	4.8	3.4-6.4	42	3.1	2.2-4.2
1979	118	5.2	4.3-6.3	63	6.3	4.8-8.2	55	4.4	3.3-5.8	81	3.6	2.8-4.4	47	4.9	3.6-6.6	34	2.6	1.8-3.7
1980	129	5.7	4.8-6.8	74	7.6	5.9-9.6	55	4.3	3.2-5.6	94	4.1	3.3-5.1	55	6.0	4.5-7.9	39	2.8	2.0-3.9
1981	117	5.5	4.5-6.6	71	7.7	6.0-9.8	46	3.7	2.7-5.0	87	4.0	3.2-5.0	49	5.3	3.8-7.1	38	3.1	2.2-4.3
1982	117	5.3	4.3-6.3	65	6.9	5.3-8.9	52	4.1	3.0-5.4	93	4.4	3.5-5.4	54	6.1	4.5-8.0	39	3.2	2.2-4.4
1983	110	4.9	4.0-5.9	60	6.1	4.6-7.9	50	3.9	2.8-5.1	95	4.4	3.6-5.4	47	5.3	3.8-7.1	48	3.9	2.8-5.2
1984	137	5.9	4.9-7.0	65	6.5	5.0-8.4	72	5.4	4.2-6.9	91	4.1	3.2-5.0	48	5.2	3.8-6.9	43	3.2	2.3-4.3
1985	131	6.0	5.0-7.1	66	7.3	5.6-9.3	65	5.0	3.8-6.4	88	3.9	3.1-4.9	47	5.0	3.6-6.7	41	3.0	2.2-4.1
1986	137	6.2	5.2-7.3	70	7.4	5.7-9.4	67	5.4	4.2-6.9	93	4.3	3.5-5.3	51	5.9	4.3-7.7	42	3.2	2.3-4.3
1987	138	6.0	5.0-7.1	81	8.1	6.4-10.2	57	4.3	3.3-5.7	91	4.1	3.3-5.0	48	5.1	3.7-6.7	43	3.3	2.4-4.5
1988	134	6.1	5.1-7.3	67	7.1	5.5-9.1	67	5.4	4.1-6.8	57	2.6	2.0-3.4	30	3.2	2.1-4.6	27	2.2	1.4-3.2
1989	130	5.8	4.8-6.8	73	7.6	5.9-9.6	57	4.4	3.4-5.8	79	3.6	2.8-4.5	35	3.9	2.7-5.4	44	3.4	2.5-4.6
1990	144	6.6	5.6-7.8	75	8.3	6.5-10.5	69	5.6	4.3-7.1	73	3.4	2.7-4.3	44	5.4	3.9-7.3	29	2.3	1.5-3.3
1991	142	6.5	5.4-7.7	76	8.1	6.3-10.2	66	5.2	4.0-6.7	74	3.5	2.7-4.4	40	4.5	3.2-6.1	34	2.7	1.9-3.8
1992	127	5.7	4.7-6.8	65	6.4	4.9-8.2	62	5.0	3.8-6.4	80	3.8	3.0-4.7	40	4.6	3.3-6.3	40	3.1	2.2-4.2
1993	133	5.8	4.9-6.9	63	6.4	4.9-8.2	70	5.4	4.2-6.8	58	2.6	2.0-3.3	31	3.0	2.0-4.3	27	2.2	1.4-3.2
1994	158	7.0	5.9-8.1	80	8.0	6.3-10.0	78	6.1	4.8-7.6	84	3.8	3.0-4.7	37	3.9	2.7-5.4	47	3.7	2.7-4.9
1995	160	7.3	6.2-8.5	75	8.0	6.3-10.1	85	6.8	5.4-8.4	72	3.3	2.6-4.2	29	3.3	2.2-4.8	43	3.4	2.5-4.6
1996	155	6.9	5.9-8.1	69	7.1	5.5-9.1	86	6.7	5.4-8.3	68	3.1	2.4-3.9	36	4.0	2.7-5.5	32	2.6	1.7-3.6
1997	152	6.7	5.7-7.9	77	7.8	6.1-9.8	75	5.9	4.7-7.4	69	3.1	2.4-4.0	29	3.1	2.0-4.4	40	3.1	2.2-4.3
1998	158	6.9	5.9-8.1	76	7.4	5.8-9.3	82	6.5	5.1-8.0	74	3.3	2.6-4.2	34	3.8	2.6-5.3	40	3.1	2.2-4.2
1999	146	6.3	5.4-7.5	72	7.2	5.6-9.2	74	5.7	4.4-7.1	78	3.5	2.7-4.3	43	4.7	3.4-6.3	35	2.6	1.8-3.6
2000	149	6.2	5.3-7.3	71	6.8	5.3-8.6	78	5.8	4.5-7.2	61	2.7	2.0-3.4	33	3.3	2.3-4.7	28	2.1	1.4-3.0
2001	164	6.9	5.9-8.1	78	7.5	5.9-9.4	86	6.4	5.1-7.9	68	2.9	2.3-3.7	40	4.0	2.8-5.5	28	2.1	1.4-3.0
2002	144	6.0	5.1-7.1	69	6.7	5.2-8.5	75	5.6	4.4-7.1	59	2.5	1.9-3.2	27	2.6	1.7-3.8	32	2.4	1.6-3.4
2003	150	6.3	5.3-7.3	73	6.9	5.4-8.7	77	5.7	4.5-7.1	55	2.3	1.7-3.0	31	2.9	2.0-4.2	24	1.7	1.1-2.6
2004	129	5.4	4.5-6.5	69	6.8	5.3-8.6	60	4.6	3.5-5.9	59	2.5	1.9-3.2	34	3.4	2.3-4.8	25	1.8	1.2-2.7
2005	131	5.5	4.6-6.5	67	6.3	4.9-8.0	64	4.8	3.7-6.2	69	2.9	2.3-3.7	39	3.9	2.8-5.4	30	2.2	1.5-3.1
2006	140	5.7	4.8-6.8	66	5.9	4.6-7.6	74	5.6	4.4-7.0	57	2.4	1.8-3.1	30	2.9	2.0-4.2	27	2.0	1.3-2.9
2007	140	5.8	4.9-6.9	70	6.3	4.9-8.0	70	5.4	4.2-6.8	80	3.4	2.7-4.2	33	3.2	2.2-4.5	47	3.5	2.6-4.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Kings (Brooklyn)

Brain and other central nervous system

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	149	6.2	5.2-7.3	81	7.5	6.0-9.4	68	5.0	3.9-6.3	58	2.4	1.8-3.0	37	3.4	2.4-4.7	21	1.6	1.0-2.4
2009	145	5.9	5.0-6.9	75	6.9	5.4-8.7	70	5.1	3.9-6.4	65	2.6	2.0-3.3	34	3.3	2.3-4.6	31	2.2	1.5-3.1
2010	150	5.9	5.0-7.0	79	7.2	5.7-9.0	71	5.0	3.9-6.3	63	2.6	2.0-3.3	32	3.0	2.0-4.2	31	2.3	1.5-3.2
2011	138	5.4	4.5-6.4	75	6.8	5.3-8.6	63	4.4	3.4-5.7	76	3.1	2.4-3.8	45	4.2	3.0-5.7	31	2.2	1.5-3.1
2012	151	6.0	5.0-7.0	86	7.4	5.9-9.1	65	4.6	3.6-5.9	75	2.8	2.2-3.6	31	2.7	1.8-3.9	44	3.0	2.2-4.1
2013	166	6.3	5.3-7.3	90	7.4	5.9-9.1	76	5.2	4.1-6.5	88	3.3	2.6-4.1	43	3.7	2.7-5.0	45	2.8	2.0-3.8
2014	161	6.0	5.1-7.0	86	7.1	5.7-8.8	75	5.1	4.0-6.4	80	3.0	2.4-3.7	45	3.8	2.7-5.1	35	2.3	1.6-3.2
2015	150	5.6	4.7-6.6	83	6.8	5.4-8.4	67	4.7	3.6-6.0	78	2.9	2.2-3.6	38	3.1	2.2-4.3	40	2.7	1.9-3.7
2016	183	6.9	5.9-8.0	102	8.5	6.9-10.4	81	5.6	4.4-7.0	76	2.7	2.1-3.4	47	3.9	2.8-5.2	29	1.8	1.2-2.7
2017	156	5.8	4.9-6.9	80	6.6	5.2-8.3	76	5.1	4.0-6.5	97	3.6	2.9-4.4	58	5.1	3.8-6.6	39	2.5	1.8-3.5
2018	150	5.4	4.5-6.3	84	6.6	5.3-8.2	66	4.4	3.4-5.6	76	2.9	2.3-3.6	38	3.2	2.2-4.4	38	2.6	1.8-3.6
2019	172	6.4	5.4-7.4	101	8.3	6.7-10.1	71	4.8	3.7-6.1	86	3.2	2.5-3.9	46	3.9	2.8-5.2	40	2.7	1.9-3.7
2020	159	6.0	5.1-7.1	74	6.1	4.7-7.6	85	6.0	4.7-7.4	73	2.6	2.0-3.3	35	2.8	1.9-3.9	38	2.4	1.7-3.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Kings (Brooklyn)

Thyroid

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	80	3.7	2.9-4.6	22	2.5	1.5-3.8	58	4.6	3.4-6.0	16	0.8	0.4-1.3	6	0.6	0.2-1.3	10	0.9	0.4-1.6
1977	80	3.5	2.7-4.3	19	1.8	1.1-2.9	61	4.8	3.6-6.2	18	0.9	0.5-1.4	10	1.0	0.5-1.9	8	0.7	0.3-1.4
1978	83	3.8	3.0-4.8	29	3.2	2.1-4.7	54	4.3	3.2-5.7	18	0.9	0.5-1.4	6	0.7	0.2-1.5	12	1.0	0.5-1.7
1979	76	3.4	2.7-4.3	34	3.7	2.5-5.2	42	3.4	2.4-4.6	16	0.7	0.4-1.2	9	1.1	0.5-2.1	7	0.5	0.2-1.1
1980	81	3.8	3.0-4.7	21	2.3	1.4-3.6	60	4.9	3.7-6.3	15	0.8	0.4-1.3	5	0.6	0.2-1.4	10	0.8	0.4-1.6
1981	81	3.8	3.0-4.7	17	1.8	1.0-2.9	64	5.3	4.1-6.8	14	0.6	0.3-1.1	3	0.3	0.1-1.0	11	0.9	0.4-1.6
1982	81	3.9	3.1-4.9	21	2.2	1.3-3.4	60	5.2	3.9-6.7	23	1.0	0.7-1.6	6	0.7	0.2-1.5	17	1.3	0.7-2.1
1983	86	4.1	3.3-5.1	14	1.6	0.9-2.8	72	6.2	4.8-7.8	12	0.6	0.3-1.0	4	0.4	0.1-1.1	8	0.7	0.3-1.4
1984	94	4.3	3.5-5.3	25	2.9	1.8-4.3	69	5.6	4.3-7.1	23	1.1	0.7-1.6	10	1.3	0.6-2.4	13	0.9	0.5-1.6
1985	92	4.2	3.4-5.2	23	2.5	1.6-3.8	69	5.5	4.3-7.0	15	0.7	0.4-1.2	9	1.1	0.5-2.0	6	0.4	0.2-1.0
1986	84	3.8	3.0-4.7	20	2.1	1.3-3.2	64	5.2	4.0-6.7	8	0.4	0.2-0.8	4	0.5	0.1-1.3	4	0.3	0.1-0.8
1987	87	3.9	3.1-4.8	21	2.3	1.4-3.6	66	5.3	4.0-6.7	9	0.4	0.2-0.8	2	0.3	0.0-1.0	7	0.5	0.2-1.0
1988	87	3.9	3.1-4.8	26	2.8	1.8-4.2	61	4.8	3.7-6.2	8	0.3	0.1-0.7	4	0.4	0.1-1.1	4	0.3	0.1-0.8
1989	87	3.9	3.1-4.9	26	2.9	1.8-4.2	61	4.9	3.7-6.4	6	0.3	0.1-0.6	2	0.3	0.0-1.0	4	0.3	0.1-0.8
1990	88	4.0	3.2-5.0	21	2.2	1.4-3.4	67	5.6	4.3-7.1	11	0.5	0.3-0.9	3	0.4	0.1-1.1	8	0.6	0.3-1.3
1991	96	4.3	3.5-5.3	25	2.7	1.7-3.9	71	5.7	4.4-7.2	9	0.4	0.2-0.8	3	0.4	0.1-1.0	6	0.5	0.2-1.0
1992	113	5.1	4.2-6.1	25	2.7	1.7-4.0	88	7.1	5.6-8.7	14	0.7	0.4-1.1	7	0.8	0.3-1.7	7	0.5	0.2-1.1
1993	114	5.2	4.3-6.2	28	3.0	2.0-4.4	86	6.9	5.5-8.5	6	0.3	0.1-0.6	2	0.3	0.0-0.9	4	0.3	0.1-0.8
1994	131	6.0	5.0-7.1	36	3.8	2.6-5.3	95	7.6	6.1-9.3	13	0.6	0.3-1.1	2	0.3	0.0-0.9	11	0.9	0.4-1.6
1995	111	4.9	4.0-5.9	27	2.9	1.9-4.2	84	6.7	5.3-8.3	6	0.3	0.1-0.6	2	0.2	0.0-0.8	4	0.3	0.1-0.8
1996	132	5.8	4.8-6.9	25	2.7	1.7-4.0	107	8.5	7.0-10.3	15	0.7	0.4-1.1	8	0.9	0.4-1.7	7	0.5	0.2-1.1
1997	140	6.2	5.2-7.3	25	2.6	1.7-3.8	115	9.2	7.5-11.0	15	0.7	0.4-1.2	6	0.7	0.3-1.5	9	0.7	0.3-1.3
1998	159	7.0	5.9-8.1	39	4.0	2.8-5.4	120	9.4	7.7-11.2	11	0.5	0.2-0.9	3	0.3	0.1-0.8	8	0.6	0.3-1.2
1999	169	7.2	6.2-8.4	41	3.9	2.8-5.3	128	9.9	8.3-11.8	14	0.6	0.4-1.1	3	0.3	0.1-0.9	11	0.8	0.4-1.5
2000	182	7.7	6.6-8.9	38	3.7	2.6-5.1	144	11.1	9.3-13.1	12	0.5	0.3-0.9	3	0.3	0.1-0.9	9	0.7	0.3-1.3
2001	192	8.1	7.0-9.3	43	4.2	3.0-5.7	149	11.3	9.5-13.3	15	0.7	0.4-1.1	3	0.3	0.1-0.9	12	0.8	0.4-1.4
2002	210	8.7	7.6-10.0	55	5.1	3.9-6.7	155	11.8	10.0-13.8	10	0.4	0.2-0.8	2	0.2	0.0-0.7	8	0.6	0.3-1.2
2003	174	7.2	6.2-8.4	34	3.1	2.2-4.4	140	10.5	8.9-12.5	8	0.3	0.1-0.7	1	0.1	0.0-0.5	7	0.5	0.2-1.0
2004	219	9.0	7.8-10.3	52	5.1	3.8-6.7	167	12.4	10.6-14.5	12	0.5	0.3-0.9	5	0.5	0.2-1.2	7	0.5	0.2-1.0
2005	318	13.0	11.6-14.5	67	6.3	4.9-8.0	251	18.8	16.5-21.3	20	0.8	0.5-1.3	6	0.7	0.2-1.4	14	1.0	0.5-1.7
2006	310	12.9	11.5-14.4	60	5.6	4.2-7.2	250	18.9	16.6-21.4	10	0.4	0.2-0.8	2	0.2	0.0-0.8	8	0.6	0.2-1.2
2007	366	15.2	13.7-16.9	80	7.2	5.7-9.0	286	21.9	19.4-24.6	21	0.9	0.5-1.3	6	0.6	0.2-1.4	15	1.1	0.6-1.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Kings (Brooklyn)

Thyroid

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	454	18.3	16.6-20.1	88	7.9	6.3-9.7	366	27.1	24.4-30.1	11	0.5	0.2-0.8	5	0.5	0.2-1.2	6	0.4	0.2-1.0
2009	493	19.8	18.0-21.6	103	9.4	7.6-11.4	390	28.6	25.8-31.6	12	0.5	0.3-0.9	7	0.9	0.3-1.7	5	0.3	0.1-0.8
2010	482	19.0	17.4-20.8	100	9.0	7.3-11.0	382	27.6	24.8-30.5	22	0.9	0.5-1.3	13	1.3	0.7-2.2	9	0.6	0.3-1.2
2011	520	20.3	18.5-22.1	101	8.8	7.1-10.7	419	29.9	27.1-33.0	21	0.9	0.5-1.3	5	0.6	0.2-1.3	16	1.1	0.6-1.8
2012	541	20.8	19.1-22.7	109	9.2	7.5-11.2	432	30.8	27.9-33.9	13	0.5	0.3-0.9	5	0.5	0.2-1.2	8	0.5	0.2-1.1
2013	637	24.3	22.4-26.3	145	12.3	10.4-14.6	492	34.7	31.6-37.9	18	0.7	0.4-1.2	7	0.7	0.3-1.4	11	0.8	0.4-1.4
2014	597	22.4	20.6-24.3	136	11.3	9.4-13.4	461	32.2	29.3-35.4	14	0.5	0.2-0.8	6	0.4	0.2-1.0	8	0.5	0.2-0.9
2015	624	23.6	21.7-25.5	136	11.2	9.4-13.3	488	34.3	31.3-37.6	17	0.6	0.4-1.0	5	0.4	0.1-1.0	12	0.7	0.4-1.3
2016	546	20.4	18.7-22.3	126	10.2	8.4-12.1	420	29.4	26.6-32.4	14	0.5	0.3-0.9	5	0.5	0.2-1.2	9	0.5	0.2-1.1
2017	588	21.5	19.8-23.3	146	11.8	9.9-13.9	442	30.0	27.2-33.0	17	0.6	0.4-1.0	5	0.5	0.2-1.1	12	0.7	0.4-1.3
2018	534	20.2	18.5-22.1	124	10.2	8.4-12.2	410	29.0	26.2-32.1	13	0.4	0.2-0.7	1	0.1	0.0-0.5	12	0.7	0.3-1.2
2019	588	21.8	20.0-23.7	151	12.2	10.3-14.4	437	30.4	27.6-33.5	16	0.5	0.3-0.9	4	0.3	0.1-0.9	12	0.7	0.3-1.2
2020	459	17.5	15.9-19.2	111	8.9	7.3-10.8	348	25.0	22.3-27.8	14	0.5	0.3-0.9	6	0.5	0.2-1.1	8	0.5	0.2-1.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Hodgkin lymphoma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	88	3.6	2.9-4.5	46	4.5	3.2-6.1	42	3.0	2.1-4.1	35	1.6	1.1-2.2	23	2.3	1.4-3.5	12	1.0	0.5-1.7
1977	71	3.0	2.3-3.8	36	3.3	2.3-4.7	35	2.8	1.9-3.9	38	1.7	1.2-2.4	18	1.8	1.0-2.9	20	1.7	1.0-2.6
1978	67	2.7	2.1-3.4	39	3.6	2.6-5.0	28	1.9	1.3-2.8	23	1.0	0.6-1.6	13	1.3	0.7-2.2	10	0.8	0.4-1.6
1979	69	3.3	2.5-4.2	31	3.3	2.2-4.7	38	3.2	2.2-4.4	33	1.6	1.1-2.3	20	2.2	1.3-3.6	13	1.2	0.6-2.1
1980	78	3.5	2.7-4.4	33	3.5	2.4-5.0	45	3.4	2.5-4.6	27	1.3	0.8-1.9	16	1.9	1.0-3.1	11	0.8	0.4-1.5
1981	74	3.1	2.4-3.9	44	4.0	2.9-5.5	30	2.3	1.5-3.3	31	1.4	1.0-2.1	17	1.9	1.1-3.2	14	1.1	0.6-1.9
1982	77	3.2	2.5-4.0	45	4.0	2.9-5.5	32	2.5	1.7-3.5	22	1.0	0.6-1.5	10	1.0	0.5-2.0	12	0.9	0.5-1.7
1983	70	3.2	2.4-4.0	46	4.5	3.2-6.1	24	2.0	1.3-3.0	21	0.9	0.6-1.4	12	1.3	0.7-2.3	9	0.6	0.3-1.2
1984	68	2.8	2.2-3.6	39	3.8	2.6-5.2	29	2.2	1.5-3.2	23	1.1	0.7-1.6	13	1.3	0.7-2.3	10	0.8	0.4-1.5
1985	66	2.8	2.2-3.6	47	4.6	3.3-6.2	19	1.5	0.9-2.3	27	1.1	0.7-1.6	20	1.8	1.1-2.8	7	0.5	0.2-1.1
1986	78	3.3	2.6-4.1	54	5.0	3.7-6.5	24	1.9	1.2-2.9	22	0.9	0.6-1.4	16	1.5	0.8-2.5	6	0.5	0.2-1.0
1987	64	2.6	2.0-3.4	32	3.1	2.1-4.5	32	2.4	1.6-3.4	14	0.6	0.3-1.0	10	0.9	0.4-1.8	4	0.3	0.1-0.8
1988	85	3.6	2.9-4.5	50	4.9	3.6-6.5	35	2.7	1.9-3.8	18	0.8	0.5-1.3	10	1.2	0.6-2.3	8	0.5	0.2-1.1
1989	87	3.9	3.1-4.8	54	5.5	4.1-7.2	33	2.7	1.8-3.8	26	1.2	0.7-1.7	15	1.5	0.8-2.6	11	0.9	0.4-1.6
1990	48	2.0	1.5-2.7	27	2.6	1.7-3.8	21	1.6	1.0-2.4	30	1.3	0.9-1.9	20	2.1	1.2-3.2	10	0.8	0.4-1.5
1991	65	2.8	2.2-3.6	32	3.2	2.1-4.5	33	2.6	1.8-3.7	17	0.8	0.4-1.3	8	0.9	0.4-1.7	9	0.7	0.3-1.3
1992	57	2.4	1.8-3.1	39	3.5	2.5-4.9	18	1.4	0.8-2.3	16	0.7	0.4-1.2	7	0.7	0.3-1.5	9	0.8	0.3-1.4
1993	54	2.2	1.7-2.9	21	1.9	1.2-2.9	33	2.4	1.7-3.4	15	0.7	0.4-1.1	10	0.9	0.4-1.7	5	0.4	0.1-1.0
1994	64	2.7	2.1-3.5	41	3.9	2.8-5.3	23	1.7	1.1-2.6	15	0.7	0.4-1.1	9	0.9	0.4-1.7	6	0.5	0.2-1.0
1995	79	3.4	2.7-4.2	40	3.8	2.7-5.2	39	3.1	2.2-4.2	12	0.6	0.3-1.0	6	0.7	0.3-1.5	6	0.5	0.2-1.1
1996	62	2.6	2.0-3.4	28	2.7	1.8-3.9	34	2.6	1.8-3.6	15	0.7	0.4-1.1	10	1.1	0.5-2.0	5	0.4	0.1-0.9
1997	67	2.8	2.2-3.6	36	3.4	2.4-4.8	31	2.4	1.6-3.5	14	0.6	0.3-1.0	7	0.6	0.2-1.3	7	0.5	0.2-1.1
1998	71	2.9	2.3-3.7	29	2.6	1.7-3.7	42	3.2	2.3-4.4	19	0.8	0.5-1.3	9	0.9	0.4-1.7	10	0.8	0.4-1.4
1999	61	2.5	1.9-3.2	29	2.5	1.7-3.6	32	2.4	1.6-3.3	10	0.4	0.2-0.8	7	0.7	0.3-1.4	3	0.2	0.0-0.7
2000	81	3.3	2.6-4.1	43	4.0	2.9-5.4	38	2.9	2.1-4.0	9	0.4	0.2-0.7	3	0.4	0.1-1.1	6	0.5	0.2-1.0
2001	79	3.2	2.5-4.0	45	3.9	2.8-5.3	34	2.5	1.7-3.5	10	0.4	0.2-0.7	4	0.4	0.1-0.9	6	0.4	0.2-1.0
2002	87	3.5	2.8-4.3	44	3.9	2.8-5.3	43	3.1	2.3-4.2	16	0.7	0.4-1.1	11	1.1	0.5-1.9	5	0.4	0.1-0.9
2003	85	3.5	2.8-4.3	40	3.7	2.6-5.0	45	3.5	2.5-4.6	7	0.3	0.1-0.6	2	0.2	0.0-0.8	5	0.4	0.1-0.9
2004	82	3.2	2.5-4.0	46	3.8	2.8-5.1	36	2.6	1.8-3.7	12	0.5	0.3-0.9	5	0.5	0.2-1.2	7	0.6	0.2-1.1
2005	84	3.5	2.8-4.3	52	4.7	3.5-6.2	32	2.4	1.7-3.4	3	0.1	0.0-0.3	3	0.2	0.0-0.7	0	0.0	0.0-0.3
2006	78	3.1	2.5-3.9	39	3.5	2.5-4.8	39	2.8	2.0-3.8	12	0.5	0.3-0.9	8	0.8	0.3-1.6	4	0.3	0.1-0.8
2007	91	3.6	2.9-4.4	38	3.3	2.3-4.6	53	3.8	2.8-5.0	14	0.6	0.3-1.0	7	0.7	0.3-1.5	7	0.5	0.2-1.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Hodgkin lymphoma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	88	3.5	2.8-4.3	48	4.2	3.1-5.6	40	2.9	2.0-3.9	10	0.4	0.2-0.7	9	0.8	0.3-1.5	1	0.1	0.0-0.4
2009	75	3.0	2.4-3.8	38	3.4	2.4-4.6	37	2.7	1.9-3.8	10	0.4	0.2-0.8	6	0.6	0.2-1.3	4	0.3	0.1-0.8
2010	79	3.0	2.4-3.8	48	4.1	3.0-5.5	31	2.1	1.4-3.0	14	0.5	0.3-0.9	9	0.8	0.3-1.5	5	0.4	0.1-0.9
2011	79	3.1	2.4-3.9	47	3.8	2.8-5.1	32	2.4	1.6-3.4	12	0.5	0.3-0.9	5	0.5	0.2-1.2	7	0.5	0.2-1.0
2012	76	2.9	2.3-3.6	38	3.2	2.2-4.4	38	2.7	1.9-3.7	5	0.2	0.1-0.4	4	0.4	0.1-0.9	1	0.1	0.0-0.4
2013	81	3.0	2.4-3.8	34	2.8	1.9-3.9	47	3.3	2.4-4.4	6	0.3	0.1-0.6	4	0.4	0.1-1.0	2	0.1	0.0-0.5
2014	89	3.3	2.7-4.1	47	3.8	2.8-5.1	42	2.9	2.1-4.0	7	0.3	0.1-0.6	2	0.2	0.0-0.7	5	0.4	0.1-0.9
2015	89	3.3	2.6-4.1	48	3.8	2.8-5.1	41	2.9	2.1-4.0	7	0.3	0.1-0.5	2	0.2	0.0-0.6	5	0.3	0.1-0.7
2016	91	3.4	2.7-4.2	42	3.5	2.5-4.8	49	3.4	2.5-4.5	6	0.2	0.1-0.5	2	0.2	0.0-0.7	4	0.3	0.1-0.7
2017	90	3.2	2.6-4.0	43	3.3	2.4-4.5	47	3.2	2.3-4.3	3	0.1	0.0-0.3	2	0.2	0.0-0.7	1	0.1	0.0-0.4
2018	65	2.5	1.9-3.2	38	3.2	2.2-4.4	27	1.9	1.2-2.8	11	0.4	0.2-0.8	6	0.6	0.2-1.2	5	0.3	0.1-0.8
2019	99	4.0	3.2-4.9	59	5.1	3.8-6.6	40	2.9	2.1-4.0	5	0.2	0.1-0.4	2	0.2	0.0-0.6	3	0.2	0.0-0.6
2020	65	2.4	1.8-3.1	40	3.2	2.2-4.4	25	1.8	1.1-2.6	7	0.2	0.1-0.5	5	0.4	0.1-0.9	2	0.1	0.0-0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Non-Hodgkin lymphomas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	227	10.4	9.0-11.9	109	12.2	9.9-14.9	118	9.3	7.6-11.1	123	5.5	4.6-6.6	69	7.8	5.9-10.0	54	4.1	3.0-5.3
1977	222	9.9	8.6-11.4	119	12.7	10.4-15.4	103	8.0	6.5-9.7	103	4.7	3.8-5.8	53	5.9	4.3-7.8	50	3.9	2.9-5.2
1978	240	11.0	9.6-12.5	114	12.2	9.9-14.7	126	10.0	8.2-11.9	119	5.6	4.6-6.7	67	7.2	5.5-9.3	52	4.3	3.2-5.6
1979	257	12.6	11.0-14.3	124	15.2	12.5-18.2	133	10.9	9.0-12.9	132	6.6	5.5-7.9	66	8.1	6.2-10.3	66	5.6	4.3-7.1
1980	289	13.3	11.8-15.0	143	15.7	13.1-18.6	146	11.5	9.6-13.6	119	5.5	4.5-6.6	65	7.4	5.7-9.5	54	4.1	3.1-5.4
1981	256	11.7	10.3-13.3	126	14.4	11.9-17.3	130	10.1	8.4-12.0	126	5.8	4.8-6.9	58	7.1	5.3-9.2	68	5.1	3.9-6.5
1982	302	13.9	12.4-15.7	151	16.6	14.0-19.6	151	11.9	10.0-14.0	135	6.2	5.2-7.4	58	6.6	5.0-8.6	77	5.9	4.7-7.5
1983	283	13.3	11.8-15.0	136	15.6	13.0-18.6	147	11.6	9.8-13.7	153	7.2	6.1-8.5	80	9.2	7.2-11.5	73	5.9	4.6-7.5
1984	304	13.7	12.2-15.4	161	17.8	15.1-20.9	143	10.9	9.2-12.9	132	6.1	5.1-7.3	73	8.1	6.3-10.2	59	4.7	3.5-6.0
1985	343	15.4	13.8-17.2	179	20.1	17.1-23.4	164	12.3	10.4-14.3	163	7.4	6.3-8.7	92	10.6	8.5-13.1	71	5.4	4.2-6.9
1986	361	16.4	14.7-18.2	178	19.4	16.5-22.5	183	14.1	12.1-16.3	137	6.4	5.3-7.5	78	9.0	7.1-11.3	59	4.5	3.4-5.9
1987	322	14.6	13.1-16.4	160	18.0	15.2-21.1	162	12.2	10.4-14.3	132	6.2	5.1-7.3	69	7.7	6.0-9.8	63	4.9	3.8-6.4
1988	345	15.9	14.2-17.7	174	19.3	16.4-22.4	171	13.4	11.4-15.6	166	7.7	6.6-9.0	85	9.8	7.7-12.1	81	6.3	5.0-7.9
1989	340	16.0	14.3-17.8	199	22.6	19.4-26.0	141	11.2	9.4-13.3	156	7.4	6.2-8.6	90	10.4	8.3-12.8	66	5.3	4.1-6.7
1990	415	19.1	17.3-21.1	224	24.3	21.1-27.8	191	15.0	12.9-17.3	159	7.5	6.3-8.7	78	9.3	7.2-11.6	81	6.3	5.0-7.8
1991	400	18.6	16.8-20.6	202	23.4	20.2-27.0	198	15.2	13.2-17.6	159	7.4	6.3-8.7	83	9.0	7.1-11.2	76	6.0	4.7-7.5
1992	426	19.8	17.9-21.8	220	24.1	20.9-27.6	206	16.3	14.2-18.8	163	7.6	6.5-8.9	84	9.7	7.6-12.0	79	6.2	4.9-7.8
1993	380	17.6	15.8-19.4	206	22.8	19.7-26.2	174	13.6	11.7-15.8	163	7.6	6.4-8.8	86	10.1	8.1-12.5	77	5.9	4.6-7.4
1994	427	19.7	17.9-21.7	222	23.8	20.7-27.3	205	16.1	14.0-18.5	166	7.9	6.7-9.2	82	9.5	7.5-11.8	84	6.6	5.2-8.1
1995	436	20.1	18.2-22.1	239	25.7	22.5-29.2	197	15.4	13.3-17.8	146	6.8	5.8-8.0	76	8.7	6.8-11.0	70	5.5	4.3-6.9
1996	438	20.1	18.3-22.1	242	27.1	23.7-30.8	196	15.2	13.1-17.5	158	7.4	6.3-8.6	79	9.7	7.6-12.1	79	6.1	4.8-7.6
1997	445	20.5	18.7-22.5	211	23.4	20.3-26.9	234	18.1	15.9-20.6	188	8.7	7.5-10.0	77	8.8	6.9-11.1	111	8.5	7.0-10.2
1998	392	17.7	16.0-19.6	203	21.7	18.7-24.9	189	14.7	12.7-17.0	185	8.5	7.3-9.8	86	9.8	7.8-12.1	99	7.5	6.1-9.1
1999	438	19.4	17.6-21.3	204	21.6	18.7-24.8	234	17.7	15.5-20.2	142	6.4	5.4-7.5	62	6.9	5.2-8.8	80	5.9	4.7-7.4
2000	453	19.6	17.8-21.5	222	23.1	20.2-26.4	231	17.2	15.0-19.6	141	6.1	5.2-7.3	60	6.5	5.0-8.4	81	5.9	4.7-7.4
2001	423	18.1	16.5-20.0	224	22.9	20.0-26.2	199	14.7	12.7-16.9	148	6.4	5.4-7.5	81	8.7	6.9-10.9	67	4.7	3.6-6.0
2002	408	17.4	15.7-19.1	219	22.0	19.1-25.1	189	13.8	11.9-15.9	139	6.0	5.0-7.1	80	8.6	6.8-10.7	59	4.3	3.3-5.5
2003	432	18.3	16.6-20.2	228	23.0	20.0-26.2	204	15.0	13.0-17.2	118	5.0	4.2-6.0	65	6.9	5.3-8.8	53	3.8	2.8-4.9
2004	405	17.2	15.6-19.0	208	21.5	18.6-24.6	197	14.4	12.4-16.5	120	5.1	4.2-6.1	53	5.4	4.0-7.1	67	4.8	3.7-6.1
2005	472	19.9	18.2-21.8	237	23.7	20.8-27.0	235	17.2	15.0-19.6	137	5.8	4.9-6.8	72	7.8	6.1-9.8	65	4.5	3.5-5.8
2006	429	18.2	16.5-20.0	212	21.1	18.3-24.1	217	15.9	13.9-18.2	121	5.3	4.4-6.3	55	6.0	4.5-7.8	66	4.9	3.8-6.3
2007	472	19.9	18.1-21.8	249	25.0	21.9-28.3	223	16.3	14.2-18.6	130	5.4	4.5-6.5	62	6.4	4.9-8.2	68	4.8	3.7-6.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Non-Hodgkin lymphomas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	471	19.4	17.7-21.3	251	24.4	21.4-27.7	220	15.7	13.7-18.0	115	4.9	4.0-5.9	58	6.1	4.6-8.0	57	4.0	3.0-5.2
2009	501	20.7	18.9-22.6	248	24.1	21.1-27.4	253	17.9	15.7-20.3	135	5.6	4.7-6.7	63	6.4	4.9-8.2	72	5.1	3.9-6.4
2010	458	18.7	17.0-20.5	232	22.2	19.4-25.3	226	15.9	13.8-18.1	132	5.5	4.6-6.5	67	7.0	5.4-8.9	65	4.4	3.4-5.7
2011	518	20.7	18.9-22.6	255	23.7	20.9-26.9	263	18.1	16.0-20.5	134	5.3	4.5-6.3	62	5.9	4.5-7.6	72	4.9	3.8-6.2
2012	510	19.8	18.1-21.6	275	25.4	22.4-28.6	235	16.1	14.1-18.4	115	4.5	3.7-5.4	67	6.4	5.0-8.2	48	3.1	2.3-4.1
2013	500	19.1	17.5-20.9	248	22.0	19.3-25.0	252	16.8	14.8-19.1	139	5.5	4.6-6.5	69	6.7	5.2-8.6	70	4.6	3.6-5.8
2014	526	19.7	18.0-21.5	287	24.9	22.1-28.0	239	15.7	13.7-17.8	116	4.4	3.6-5.2	52	4.9	3.6-6.4	64	4.1	3.1-5.2
2015	467	17.3	15.7-19.0	240	20.1	17.6-22.9	227	14.7	12.9-16.8	96	3.7	3.0-4.5	49	4.8	3.5-6.4	47	3.0	2.2-4.0
2016	523	19.1	17.5-20.9	271	22.8	20.1-25.8	252	16.3	14.3-18.6	119	4.5	3.7-5.3	58	5.2	3.9-6.7	61	3.9	3.0-5.0
2017	562	20.8	19.1-22.6	302	26.0	23.1-29.1	260	17.0	14.9-19.2	124	4.5	3.7-5.4	59	5.2	3.9-6.7	65	4.0	3.1-5.1
2018	510	18.5	16.9-20.2	271	22.6	20.0-25.5	239	15.3	13.4-17.5	91	3.3	2.6-4.0	48	4.3	3.1-5.7	43	2.5	1.8-3.4
2019	510	18.4	16.8-20.1	285	23.6	20.9-26.6	225	14.2	12.4-16.2	126	4.4	3.6-5.2	72	6.0	4.7-7.6	54	3.2	2.4-4.3
2020	452	16.3	14.8-17.9	231	19.3	16.8-22.0	221	14.0	12.2-16.1	95	3.3	2.7-4.1	48	4.0	3.0-5.4	47	2.8	2.0-3.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Myeloma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	80	3.7	2.9-4.6	46	5.4	3.9-7.2	34	2.5	1.7-3.6	59	2.7	2.0-3.6	26	3.1	2.0-4.6	33	2.5	1.7-3.6
1977	103	4.6	3.8-5.7	47	5.3	3.8-7.1	56	4.2	3.1-5.4	60	2.7	2.0-3.5	26	3.0	1.9-4.5	34	2.5	1.7-3.6
1978	101	4.7	3.8-5.8	55	6.3	4.7-8.2	46	3.6	2.6-4.9	55	2.4	1.8-3.2	26	2.9	1.9-4.3	29	2.1	1.4-3.1
1979	114	5.3	4.3-6.4	53	6.0	4.4-7.9	61	4.9	3.7-6.3	62	2.8	2.1-3.6	29	3.3	2.1-4.7	33	2.5	1.7-3.5
1980	101	4.7	3.8-5.8	39	4.5	3.1-6.2	62	4.9	3.7-6.3	69	3.1	2.4-3.9	31	3.4	2.3-4.9	38	2.8	2.0-3.9
1981	126	5.9	4.9-7.0	50	5.9	4.3-7.8	76	6.0	4.7-7.5	65	3.0	2.3-3.8	29	3.3	2.2-4.8	36	2.7	1.9-3.8
1982	123	5.8	4.8-7.0	53	6.4	4.7-8.4	70	5.4	4.2-6.9	80	3.7	2.9-4.6	35	4.0	2.8-5.6	45	3.4	2.5-4.6
1983	116	5.5	4.5-6.6	46	5.9	4.2-7.9	70	5.3	4.1-6.8	74	3.5	2.7-4.4	23	2.8	1.7-4.2	51	4.0	2.9-5.3
1984	108	5.1	4.2-6.2	50	5.9	4.4-7.9	58	4.6	3.4-5.9	78	3.6	2.8-4.5	25	2.8	1.8-4.2	53	4.1	3.0-5.4
1985	111	5.2	4.2-6.2	52	6.3	4.7-8.4	59	4.5	3.4-5.9	72	3.2	2.5-4.1	30	3.4	2.3-5.0	42	3.1	2.2-4.3
1986	116	5.4	4.4-6.5	60	7.0	5.3-9.0	56	4.3	3.2-5.6	82	3.8	3.0-4.7	36	4.3	3.0-6.0	46	3.5	2.5-4.7
1987	98	4.5	3.7-5.5	42	4.9	3.5-6.6	56	4.3	3.2-5.6	55	2.5	1.9-3.3	30	3.4	2.2-4.8	25	1.9	1.2-2.8
1988	121	5.7	4.7-6.8	64	7.8	5.9-9.9	57	4.4	3.3-5.8	62	2.9	2.2-3.7	28	3.3	2.2-4.8	34	2.6	1.8-3.6
1989	126	5.9	4.9-7.0	62	7.3	5.5-9.4	64	5.0	3.9-6.4	63	3.0	2.3-3.9	31	3.8	2.6-5.5	32	2.5	1.7-3.6
1990	116	5.5	4.5-6.6	48	5.5	4.0-7.3	68	5.4	4.1-6.8	54	2.5	1.9-3.3	20	2.5	1.5-3.9	34	2.6	1.8-3.7
1991	121	5.8	4.8-6.9	51	6.1	4.5-8.0	70	5.6	4.3-7.1	74	3.6	2.8-4.5	31	4.0	2.7-5.7	43	3.4	2.5-4.6
1992	142	6.6	5.6-7.8	71	8.3	6.4-10.4	71	5.5	4.3-6.9	72	3.4	2.7-4.3	29	3.6	2.4-5.2	43	3.5	2.5-4.7
1993	122	5.7	4.7-6.8	47	5.7	4.2-7.6	75	5.7	4.5-7.2	79	3.7	3.0-4.7	41	5.0	3.6-6.8	38	2.9	2.1-4.0
1994	124	5.9	4.9-7.0	55	6.4	4.8-8.3	69	5.7	4.4-7.2	66	3.1	2.4-4.0	32	3.7	2.5-5.2	34	2.6	1.8-3.7
1995	141	6.6	5.6-7.8	62	7.3	5.6-9.4	79	6.2	4.9-7.7	98	4.6	3.7-5.6	39	4.7	3.3-6.5	59	4.6	3.5-5.9
1996	144	6.7	5.6-7.9	68	7.9	6.1-10.1	76	6.0	4.7-7.5	80	3.7	3.0-4.6	30	3.8	2.5-5.4	50	3.8	2.8-5.0
1997	125	5.8	4.9-7.0	55	6.3	4.7-8.2	70	5.5	4.3-6.9	70	3.3	2.5-4.1	36	4.2	2.9-5.8	34	2.7	1.8-3.7
1998	156	7.1	6.1-8.4	70	7.9	6.1-9.9	86	6.6	5.3-8.1	63	2.9	2.2-3.7	28	3.3	2.2-4.8	35	2.7	1.9-3.8
1999	141	6.3	5.3-7.5	55	6.0	4.5-7.8	86	6.5	5.2-8.0	70	3.1	2.4-4.0	28	3.1	2.0-4.4	42	3.1	2.2-4.2
2000	161	7.1	6.0-8.3	57	6.1	4.6-7.9	104	7.7	6.3-9.3	73	3.2	2.5-4.1	24	2.7	1.7-4.0	49	3.6	2.7-4.8
2001	149	6.5	5.5-7.6	65	7.4	5.7-9.4	84	6.2	4.9-7.7	76	3.3	2.6-4.2	34	3.8	2.7-5.4	42	3.0	2.2-4.1
2002	153	6.6	5.6-7.7	68	7.3	5.6-9.2	85	6.3	5.0-7.7	77	3.3	2.6-4.2	40	4.5	3.2-6.1	37	2.7	1.9-3.7
2003	165	7.0	6.0-8.2	81	8.7	6.9-10.8	84	6.1	4.9-7.6	51	2.2	1.6-2.9	26	2.7	1.8-4.0	25	1.8	1.2-2.7
2004	179	7.6	6.5-8.8	70	7.1	5.6-9.0	109	7.8	6.4-9.4	65	2.8	2.2-3.6	24	2.6	1.7-3.9	41	3.0	2.2-4.1
2005	169	7.2	6.2-8.4	89	9.2	7.4-11.3	80	5.9	4.6-7.3	67	2.9	2.2-3.6	37	3.8	2.7-5.3	30	2.0	1.4-2.9
2006	187	8.1	7.0-9.4	103	10.9	8.9-13.2	84	6.2	4.9-7.6	55	2.4	1.8-3.1	29	3.0	2.0-4.4	26	1.9	1.2-2.8
2007	163	6.9	5.9-8.0	75	7.6	6.0-9.6	88	6.3	5.0-7.8	61	2.6	2.0-3.3	26	2.6	1.7-3.8	35	2.5	1.7-3.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Kings (Brooklyn)

Myeloma

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	192	8.0	6.9-9.2	96	9.7	7.8-11.9	96	6.9	5.6-8.5	59	2.5	1.9-3.2	31	3.3	2.2-4.7	28	2.0	1.3-2.8
2009	233	9.6	8.4-10.9	114	11.1	9.1-13.4	119	8.3	6.9-10.0	62	2.6	2.0-3.3	36	3.6	2.5-5.0	26	1.8	1.2-2.6
2010	215	8.8	7.7-10.1	103	10.2	8.3-12.4	112	8.0	6.6-9.6	70	2.9	2.2-3.6	33	3.2	2.2-4.5	37	2.6	1.8-3.6
2011	225	9.1	8.0-10.4	113	10.4	8.5-12.6	112	8.2	6.7-9.8	80	3.2	2.5-4.0	37	3.6	2.5-5.0	43	2.9	2.1-3.9
2012	236	9.2	8.1-10.5	114	11.1	9.1-13.3	122	8.1	6.7-9.7	59	2.4	1.8-3.0	28	2.8	1.8-4.0	31	2.0	1.4-2.9
2013	255	9.5	8.3-10.7	128	11.8	9.8-14.1	127	8.0	6.7-9.6	82	3.1	2.5-3.9	45	4.0	2.9-5.4	37	2.4	1.7-3.3
2014	284	10.9	9.7-12.3	138	12.5	10.5-14.8	146	9.7	8.2-11.5	78	3.0	2.3-3.7	35	3.3	2.3-4.6	43	2.7	2.0-3.7
2015	256	9.3	8.2-10.5	135	11.6	9.7-13.8	121	7.7	6.3-9.2	87	3.3	2.7-4.1	44	4.2	3.0-5.6	43	2.8	2.0-3.8
2016	243	9.0	7.9-10.2	133	11.7	9.7-13.9	110	7.1	5.8-8.6	89	3.3	2.6-4.0	43	3.8	2.7-5.2	46	2.9	2.1-3.8
2017	284	10.4	9.2-11.7	142	12.2	10.3-14.4	142	9.0	7.5-10.6	98	3.5	2.8-4.3	48	4.4	3.2-5.8	50	2.9	2.2-3.9
2018	254	8.8	7.8-10.0	115	9.5	7.8-11.5	139	8.5	7.1-10.0	67	2.3	1.8-2.9	32	2.6	1.8-3.7	35	2.1	1.5-3.0
2019	241	8.6	7.5-9.7	120	10.0	8.3-12.0	121	7.6	6.3-9.1	76	2.5	2.0-3.2	41	3.3	2.4-4.6	35	2.0	1.4-2.8
2020	221	7.8	6.8-8.9	105	8.6	7.0-10.4	116	7.2	5.9-8.6	63	2.1	1.6-2.8	35	3.0	2.1-4.2	28	1.6	1.0-2.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Leukemias**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	246	11.3	9.9-12.9	137	14.7	12.2-17.5	109	8.8	7.2-10.7	180	8.4	7.2-9.8	98	10.9	8.7-13.4	82	6.6	5.2-8.3
1977	227	10.5	9.1-12.0	113	12.3	10.0-15.0	114	9.1	7.5-11.0	148	6.9	5.8-8.1	81	8.8	6.8-11.0	67	5.5	4.2-7.0
1978	231	10.9	9.5-12.4	129	14.4	11.9-17.3	102	8.4	6.8-10.2	159	7.6	6.4-8.9	86	9.5	7.5-11.9	73	6.1	4.8-7.8
1979	240	11.4	10.0-13.0	111	12.4	10.1-15.0	129	10.8	8.9-12.8	165	8.0	6.8-9.4	83	10.0	7.8-12.5	82	6.8	5.4-8.5
1980	264	12.2	10.7-13.8	139	15.3	12.8-18.2	125	10.1	8.4-12.1	163	7.6	6.4-8.9	89	9.6	7.6-11.9	74	6.1	4.7-7.7
1981	241	11.4	9.9-12.9	125	14.5	12.0-17.4	116	9.4	7.7-11.3	172	8.2	7.0-9.5	96	11.2	9.0-13.8	76	6.1	4.8-7.7
1982	241	11.2	9.8-12.7	124	13.8	11.4-16.6	117	9.4	7.7-11.3	174	8.3	7.1-9.6	94	11.1	8.8-13.6	80	6.5	5.1-8.1
1983	241	11.1	9.7-12.7	121	13.4	11.0-16.2	120	9.8	8.1-11.7	167	8.0	6.8-9.3	88	10.1	8.0-12.5	79	6.5	5.1-8.2
1984	266	12.4	11.0-14.1	148	17.2	14.4-20.4	118	9.3	7.7-11.2	170	7.9	6.7-9.2	86	9.7	7.7-12.1	84	6.7	5.3-8.3
1985	245	11.3	9.9-12.8	143	16.7	14.0-19.8	102	8.0	6.5-9.7	138	6.3	5.3-7.5	71	8.3	6.4-10.6	67	5.3	4.1-6.7
1986	241	10.8	9.5-12.3	125	13.5	11.1-16.2	116	9.1	7.5-10.9	150	6.9	5.8-8.1	77	8.5	6.6-10.7	73	5.7	4.5-7.2
1987	204	9.3	8.1-10.7	94	10.9	8.7-13.3	110	8.5	6.9-10.3	144	6.5	5.4-7.6	71	8.1	6.2-10.3	73	5.6	4.4-7.0
1988	182	8.3	7.1-9.6	81	9.4	7.4-11.7	101	7.7	6.3-9.4	125	5.8	4.8-6.9	57	6.7	5.0-8.7	68	5.2	4.0-6.6
1989	260	11.9	10.5-13.5	132	14.8	12.3-17.7	128	10.1	8.4-12.0	145	6.7	5.7-7.9	83	9.6	7.6-12.0	62	4.9	3.7-6.3
1990	273	12.7	11.2-14.3	132	15.2	12.6-18.1	141	11.2	9.4-13.3	143	6.6	5.6-7.8	67	8.3	6.3-10.6	76	5.8	4.5-7.3
1991	223	10.3	9.0-11.7	129	14.3	11.9-17.1	94	7.4	5.9-9.0	120	5.6	4.6-6.7	67	7.8	6.0-10.0	53	4.2	3.1-5.5
1992	249	11.6	10.2-13.1	126	14.2	11.8-17.0	123	9.9	8.2-11.8	132	6.3	5.2-7.4	68	8.3	6.4-10.6	64	5.0	3.9-6.5
1993	247	11.5	10.1-13.0	125	14.2	11.7-16.9	122	9.7	8.0-11.6	149	6.9	5.8-8.1	81	9.1	7.2-11.4	68	5.3	4.1-6.8
1994	286	13.3	11.8-14.9	151	17.5	14.8-20.6	135	10.5	8.8-12.4	136	6.4	5.3-7.5	68	8.1	6.3-10.3	68	5.3	4.1-6.7
1995	268	12.4	11.0-14.0	135	15.5	12.9-18.4	133	10.5	8.8-12.5	147	6.8	5.8-8.0	70	8.3	6.5-10.6	77	6.1	4.8-7.7
1996	263	12.0	10.6-13.5	143	16.1	13.5-19.0	120	9.3	7.7-11.1	147	6.8	5.7-8.0	66	7.8	6.0-10.0	81	6.3	5.0-7.9
1997	263	12.0	10.6-13.5	132	15.1	12.6-18.0	131	10.1	8.4-11.9	133	6.1	5.1-7.3	63	7.4	5.6-9.5	70	5.4	4.2-6.8
1998	227	10.2	8.9-11.6	117	12.8	10.5-15.4	110	8.4	6.9-10.1	124	5.6	4.7-6.7	70	8.0	6.2-10.1	54	4.1	3.1-5.4
1999	269	11.8	10.4-13.3	139	14.9	12.4-17.6	130	9.7	8.1-11.5	145	6.4	5.4-7.6	79	8.8	6.9-10.9	66	4.9	3.8-6.2
2000	272	11.8	10.4-13.3	131	14.0	11.7-16.7	141	10.3	8.7-12.2	125	5.4	4.5-6.5	52	5.7	4.2-7.5	73	5.3	4.1-6.6
2001	244	10.5	9.2-11.9	124	13.2	10.9-15.8	120	8.6	7.2-10.3	149	6.4	5.4-7.5	78	8.6	6.7-10.7	71	5.0	3.9-6.3
2002	288	12.2	10.8-13.7	154	16.0	13.5-18.8	134	9.7	8.1-11.5	139	5.9	5.0-7.0	68	7.6	5.8-9.6	71	5.0	3.9-6.4
2003	277	11.7	10.4-13.2	151	15.4	13.0-18.1	126	9.2	7.6-10.9	120	5.1	4.2-6.1	71	7.4	5.8-9.4	49	3.5	2.6-4.7
2004	271	11.5	10.1-12.9	130	13.3	11.1-15.8	141	10.2	8.6-12.0	127	5.4	4.5-6.5	71	7.8	6.1-9.9	56	3.9	3.0-5.1
2005	280	11.9	10.5-13.4	153	15.7	13.3-18.4	127	9.3	7.7-11.1	123	5.3	4.4-6.3	62	6.5	5.0-8.3	61	4.4	3.4-5.7
2006	251	10.7	9.4-12.1	131	13.6	11.3-16.1	120	8.7	7.2-10.4	124	5.3	4.4-6.3	69	7.3	5.7-9.3	55	3.9	2.9-5.1
2007	297	12.4	11.0-13.9	164	16.3	13.9-19.0	133	9.6	8.1-11.4	116	4.9	4.0-5.9	66	7.0	5.4-8.9	50	3.7	2.7-4.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Kings (Brooklyn)
Leukemias**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	277	11.6	10.3-13.1	139	14.3	12.0-16.9	138	9.9	8.3-11.8	128	5.3	4.4-6.3	62	6.2	4.7-8.0	66	4.6	3.6-5.9
2009	344	14.0	12.6-15.6	183	17.8	15.3-20.6	161	11.4	9.7-13.3	116	4.7	3.9-5.7	57	5.6	4.2-7.3	59	4.1	3.1-5.3
2010	331	13.5	12.1-15.0	181	17.5	15.0-20.3	150	10.8	9.1-12.7	146	6.0	5.1-7.1	76	7.9	6.2-9.9	70	4.9	3.8-6.2
2011	359	13.9	12.5-15.5	188	17.1	14.7-19.8	171	11.6	9.9-13.5	144	5.6	4.7-6.6	77	7.6	6.0-9.5	67	4.3	3.3-5.5
2012	325	12.5	11.2-14.0	183	16.6	14.3-19.3	142	9.5	7.9-11.2	122	4.8	4.0-5.7	72	7.0	5.5-8.9	50	3.3	2.4-4.4
2013	348	13.5	12.1-15.0	197	17.8	15.4-20.6	151	10.4	8.8-12.3	144	5.7	4.8-6.7	75	7.2	5.7-9.1	69	4.7	3.6-5.9
2014	383	14.6	13.2-16.2	197	17.2	14.9-19.9	186	12.5	10.7-14.5	132	5.0	4.2-6.0	70	6.7	5.2-8.5	62	3.9	3.0-5.0
2015	368	13.8	12.4-15.3	193	17.0	14.7-19.7	175	11.5	9.8-13.3	142	5.3	4.4-6.3	75	6.7	5.2-8.4	67	4.2	3.3-5.4
2016	376	14.0	12.6-15.5	214	18.6	16.2-21.4	162	10.5	8.9-12.3	140	5.2	4.4-6.2	73	6.5	5.1-8.2	67	4.4	3.4-5.6
2017	352	12.9	11.6-14.4	177	15.0	12.8-17.4	175	11.3	9.7-13.2	141	5.2	4.4-6.2	67	6.0	4.6-7.6	74	4.7	3.6-5.9
2018	336	12.2	10.9-13.6	191	16.1	13.8-18.6	145	9.3	7.8-11.0	133	4.8	4.0-5.7	67	5.7	4.4-7.3	66	4.2	3.2-5.4
2019	317	11.7	10.4-13.1	170	14.5	12.4-16.9	147	9.5	8.0-11.2	109	4.0	3.3-4.8	62	5.5	4.2-7.0	47	2.9	2.1-3.9
2020	329	12.1	10.8-13.5	187	15.7	13.5-18.2	142	9.4	7.9-11.2	105	3.7	3.0-4.5	59	5.0	3.8-6.5	46	2.9	2.1-3.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

New York State Department of Health, New York State Cancer Registry
 Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020

New York (Manhattan)
 All Invasive Malignant Tumors

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	6,723	432.3	421.8-443.1	3,177	506.8	488.5-525.6	3,546	395.7	382.3-409.4	3,581	232.9	225.1-241.0	1,804	295.2	281.0-309.8	1,777	196.5	187.2-206.2
1977	6,462	420.5	410.1-431.2	3,047	495.1	476.9-513.7	3,415	384.4	371.2-398.0	3,590	236.0	228.1-244.1	1,826	300.9	286.7-315.7	1,764	196.6	187.3-206.4
1978	6,234	409.6	399.3-420.2	2,958	483.6	465.7-502.0	3,276	372.0	358.9-385.4	3,511	233.1	225.2-241.1	1,789	299.8	285.5-314.6	1,722	193.5	184.2-203.1
1979	6,530	430.5	419.9-441.3	3,127	514.3	495.9-533.3	3,403	387.7	374.3-401.4	3,395	225.1	217.5-233.0	1,691	283.4	269.5-297.7	1,704	190.3	181.1-199.8
1980	6,508	432.2	421.5-443.0	3,172	529.1	510.3-548.5	3,336	383.6	370.3-397.3	3,517	233.5	225.7-241.4	1,754	301.1	286.6-316.0	1,763	195.4	186.1-205.0
1981	6,443	427.2	416.6-438.0	3,120	516.2	497.7-535.2	3,323	378.9	365.7-392.4	3,372	225.5	217.9-233.4	1,625	277.4	263.6-291.6	1,747	194.4	185.2-204.0
1982	6,459	433.7	423.0-444.6	3,072	516.9	498.3-536.1	3,387	390.4	377.0-404.2	3,293	220.6	213.0-228.4	1,631	282.3	268.3-296.7	1,662	185.3	176.3-194.7
1983	6,622	439.9	429.2-450.8	3,227	529.3	510.7-548.5	3,395	389.5	376.2-403.2	3,326	223.2	215.6-231.1	1,645	281.9	268.1-296.3	1,681	187.6	178.5-197.0
1984	6,878	454.5	443.6-465.5	3,359	551.5	532.5-571.1	3,519	401.4	387.9-415.3	3,396	224.2	216.6-231.9	1,691	289.3	275.2-303.8	1,705	187.4	178.5-196.8
1985	6,929	456.8	446.0-467.9	3,482	564.5	545.4-584.2	3,447	393.4	380.1-407.1	3,407	225.4	217.8-233.2	1,728	295.2	281.1-309.8	1,679	184.0	175.2-193.2
1986	6,990	459.4	448.6-470.5	3,594	583.5	563.9-603.5	3,396	386.2	373.1-399.8	3,281	217.1	209.7-224.8	1,647	283.8	269.9-298.2	1,634	178.9	170.2-188.0
1987	6,822	446.6	436.0-457.5	3,468	555.8	536.9-575.2	3,354	378.1	365.2-391.4	3,214	212.6	205.3-220.2	1,612	277.1	263.4-291.3	1,602	175.7	167.1-184.7
1988	6,696	440.4	429.8-451.2	3,318	532.8	514.3-551.8	3,378	383.7	370.7-397.2	3,337	221.4	213.9-229.1	1,661	285.6	271.7-300.0	1,676	183.3	174.6-192.5
1989	6,769	446.5	435.8-457.4	3,401	545.0	526.3-564.1	3,368	386.6	373.4-400.1	3,088	205.2	198.0-212.6	1,502	259.4	246.1-273.1	1,586	173.8	165.2-182.7
1990	6,750	445.3	434.7-456.2	3,417	547.5	528.8-566.7	3,333	383.6	370.4-397.1	3,241	216.4	209.0-224.1	1,572	272.6	258.9-286.7	1,669	184.3	175.5-193.6
1991	8,094	539.3	527.5-551.3	3,987	643.8	623.4-664.7	4,107	476.5	461.8-491.5	3,216	217.0	209.5-224.7	1,538	268.0	254.5-282.1	1,678	188.6	179.5-198.0
1992	7,796	519.6	508.0-531.3	3,968	645.1	624.6-666.1	3,828	443.9	429.7-458.4	3,140	212.2	204.8-219.8	1,535	270.6	256.9-284.8	1,605	179.2	170.4-188.3
1993	7,561	503.3	491.9-514.9	3,851	616.4	596.6-636.8	3,710	432.4	418.4-446.8	2,957	200.0	192.8-207.4	1,429	252.1	238.9-265.8	1,528	171.1	162.5-180.0
1994	8,193	546.7	534.8-558.7	4,088	650.7	630.3-671.5	4,105	478.2	463.5-493.2	3,075	207.8	200.4-215.3	1,462	256.0	242.7-269.7	1,613	180.6	171.9-189.8
1995	7,173	478.1	467.0-489.3	3,623	579.7	560.5-599.4	3,550	411.7	398.1-425.6	3,053	206.4	199.1-213.9	1,457	255.5	242.3-269.3	1,596	177.7	169.0-186.7
1996	7,440	496.0	484.8-507.5	3,622	581.5	562.3-601.2	3,818	444.1	430.0-458.6	3,000	202.7	195.5-210.2	1,462	256.1	242.9-269.9	1,538	172.5	163.9-181.5
1997	7,400	491.9	480.7-503.3	3,585	577.1	558.0-596.7	3,815	440.1	426.1-454.5	2,847	191.8	184.8-199.0	1,348	234.8	222.2-247.9	1,499	167.6	159.1-176.4
1998	7,535	499.4	488.1-510.8	3,670	583.2	564.1-602.8	3,865	446.3	432.2-460.8	2,845	191.3	184.3-198.5	1,307	222.9	210.7-235.5	1,538	171.9	163.3-180.8
1999	7,421	486.5	475.4-497.7	3,589	564.5	545.8-583.7	3,832	437.5	423.6-451.7	2,883	192.3	185.3-199.4	1,391	238.3	225.6-251.4	1,492	165.7	157.3-174.4
2000	7,706	499.6	488.5-511.0	3,724	578.8	560.0-598.1	3,982	451.3	437.3-465.7	2,789	183.4	176.6-190.3	1,325	224.3	212.2-236.9	1,464	159.7	151.5-168.2
2001	7,555	482.4	471.5-493.4	3,649	559.4	541.1-578.2	3,906	434.5	420.9-448.5	2,792	180.4	173.7-187.2	1,305	216.9	205.1-229.2	1,487	159.0	150.9-167.4
2002	7,604	479.3	468.6-490.3	3,823	575.2	556.8-594.1	3,781	416.6	403.3-430.2	2,820	180.0	173.4-186.8	1,365	222.4	210.6-234.8	1,455	154.2	146.3-162.5
2003	7,505	467.9	457.3-478.7	3,647	539.8	522.2-557.9	3,858	422.8	409.4-436.5	2,717	170.6	164.2-177.2	1,303	207.1	195.8-218.8	1,414	148.0	140.3-156.0
2004	7,694	472.1	461.5-482.9	3,810	553.6	535.9-571.8	3,884	420.1	406.9-433.8	2,596	160.4	154.3-166.8	1,258	195.7	184.8-207.0	1,338	138.3	130.9-146.0
2005	7,863	476.5	465.9-487.2	3,844	547.6	530.2-565.5	4,019	431.8	418.3-445.6	2,720	166.7	160.5-173.1	1,291	197.6	186.8-208.9	1,429	148.4	140.7-156.4
2006	8,212	495.4	484.7-506.4	4,168	592.2	574.0-610.7	4,044	433.5	420.0-447.3	2,602	157.7	151.6-163.9	1,257	192.3	181.6-203.3	1,345	137.1	129.8-144.8
2007	8,204	490.9	480.2-501.8	4,165	582.1	564.3-600.4	4,039	431.1	417.6-444.9	2,694	161.5	155.4-167.8	1,363	203.3	192.5-214.6	1,331	135.1	127.8-142.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.
 Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.
 Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
 The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
All Invasive Malignant Tumors**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	8,343	492.4	481.8-503.2	4,076	563.5	546.1-581.4	4,267	448.7	435.1-462.7	2,697	159.4	153.4-165.6	1,312	193.4	183.0-204.3	1,385	138.7	131.3-146.3
2009	8,263	483.3	472.8-494.0	4,066	551.7	534.6-569.3	4,197	439.0	425.5-452.8	2,623	153.2	147.3-159.2	1,295	188.6	178.4-199.3	1,328	130.6	123.5-138.0
2010	8,088	465.7	455.4-476.1	3,904	520.4	504.0-537.3	4,184	433.6	420.3-447.4	2,535	143.9	138.3-149.7	1,260	177.7	167.9-187.9	1,275	122.0	115.2-129.0
2011	8,342	469.4	459.3-479.8	4,035	526.7	510.3-543.5	4,307	436.3	423.0-449.9	2,492	139.3	133.8-145.0	1,176	161.8	152.6-171.5	1,316	124.9	118.1-132.1
2012	7,864	434.5	424.8-444.4	3,808	483.7	468.2-499.6	4,056	404.4	391.7-417.4	2,447	133.6	128.3-139.1	1,145	153.3	144.4-162.5	1,302	120.9	114.3-127.8
2013	8,550	464.6	454.6-474.7	4,122	516.0	500.1-532.3	4,428	435.5	422.4-448.9	2,519	134.6	129.4-140.1	1,213	159.4	150.4-168.7	1,306	119.3	112.8-126.2
2014	8,424	451.3	441.5-461.2	3,964	484.2	469.0-499.7	4,460	435.8	422.7-449.2	2,498	130.6	125.5-135.9	1,200	152.4	143.8-161.4	1,298	116.0	109.7-122.7
2015	8,436	444.2	434.6-454.0	4,052	486.2	471.1-501.7	4,384	422.7	409.9-435.9	2,524	129.3	124.2-134.5	1,234	152.8	144.3-161.7	1,290	113.6	107.3-120.2
2016	8,632	446.4	436.8-456.1	4,145	485.0	470.1-500.2	4,487	427.1	414.3-440.3	2,461	123.0	118.1-128.0	1,207	145.1	136.9-153.7	1,254	108.7	102.6-115.1
2017	8,685	445.8	436.2-455.4	4,080	471.1	456.6-486.0	4,605	434.9	422.0-448.1	2,341	115.2	110.5-120.1	1,124	133.4	125.6-141.5	1,217	104.1	98.2-110.3
2018	8,456	428.0	418.7-437.4	4,095	466.6	452.2-481.4	4,361	405.5	393.1-418.3	2,330	111.5	107.0-116.3	1,097	125.9	118.5-133.7	1,233	102.1	96.3-108.1
2019	8,850	442.4	433.0-451.9	4,276	480.0	465.5-494.8	4,574	422.1	409.5-435.0	2,277	107.8	103.4-112.5	1,087	123.0	115.8-130.7	1,190	97.6	91.9-103.5
2020	7,422	371.6	363.0-380.4	3,519	393.8	380.7-407.2	3,903	363.3	351.5-375.5	2,107	99.3	95.0-103.8	1,004	113.2	106.2-120.5	1,103	90.4	84.9-96.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
Oral cavity and pharynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	238	15.7	13.7-17.8	152	23.7	19.9-27.9	86	9.9	7.8-12.3	106	7.1	5.8-8.7	73	11.4	8.9-14.5	33	4.0	2.7-5.7
1977	230	15.0	13.1-17.2	162	25.5	21.6-29.8	68	7.4	5.7-9.5	129	8.4	7.0-10.1	94	14.5	11.6-17.8	35	3.9	2.7-5.5
1978	210	13.8	12.0-15.9	139	21.5	18.0-25.5	71	7.9	6.1-10.0	94	6.0	4.9-7.4	69	10.5	8.1-13.4	25	2.8	1.8-4.2
1979	224	15.1	13.1-17.3	159	24.8	21.0-29.1	65	7.6	5.8-9.7	93	6.1	4.9-7.5	67	10.5	8.1-13.4	26	2.9	1.9-4.3
1980	228	15.7	13.7-17.9	151	23.8	20.1-28.0	77	9.4	7.4-11.9	90	6.1	4.9-7.5	65	10.6	8.2-13.7	25	2.7	1.7-4.1
1981	243	16.4	14.4-18.7	176	27.5	23.5-32.0	67	7.8	6.0-10.0	86	6.0	4.8-7.5	56	9.0	6.8-11.7	30	3.5	2.3-5.1
1982	240	16.4	14.3-18.6	166	26.9	22.9-31.5	74	8.6	6.7-10.9	91	6.4	5.1-7.9	67	11.0	8.5-14.1	24	2.8	1.8-4.2
1983	227	15.1	13.2-17.3	159	24.5	20.8-28.7	68	7.4	5.7-9.4	88	5.8	4.7-7.2	65	10.4	7.9-13.3	23	2.7	1.7-4.0
1984	234	15.5	13.6-17.7	166	25.9	22.0-30.2	68	8.0	6.2-10.3	82	5.5	4.3-6.8	58	9.1	6.9-11.8	24	2.7	1.7-4.1
1985	216	14.6	12.7-16.8	128	20.4	16.9-24.3	88	10.8	8.6-13.3	79	5.3	4.2-6.6	53	8.6	6.4-11.3	26	3.0	1.9-4.4
1986	229	15.5	13.5-17.7	162	25.6	21.7-30.0	67	7.8	6.0-10.0	66	4.2	3.3-5.4	51	7.8	5.8-10.4	15	1.6	0.9-2.7
1987	218	14.5	12.6-16.6	152	23.4	19.8-27.5	66	7.6	5.8-9.7	68	4.5	3.5-5.8	49	7.7	5.7-10.2	19	2.1	1.3-3.4
1988	175	11.7	10.0-13.6	109	17.1	14.0-20.7	66	7.7	5.9-9.9	79	5.4	4.3-6.8	51	8.2	6.1-10.8	28	3.4	2.2-4.9
1989	185	12.6	10.9-14.6	120	18.8	15.5-22.5	65	7.9	6.1-10.1	59	4.0	3.0-5.1	39	6.6	4.6-9.0	20	2.2	1.4-3.5
1990	221	14.7	12.8-16.8	144	23.1	19.4-27.3	77	8.8	6.9-11.1	67	4.6	3.5-5.8	44	7.2	5.2-9.7	23	2.7	1.7-4.2
1991	283	19.1	16.9-21.5	188	29.4	25.3-34.0	95	11.0	8.9-13.5	62	4.2	3.2-5.4	43	7.3	5.3-9.9	19	2.3	1.4-3.6
1992	260	17.6	15.5-19.9	175	27.9	23.9-32.5	85	10.2	8.1-12.6	66	4.5	3.5-5.8	43	7.3	5.3-9.9	23	2.6	1.6-3.9
1993	240	16.4	14.4-18.7	162	25.6	21.8-30.0	78	9.6	7.6-12.0	51	3.4	2.6-4.5	36	5.9	4.1-8.2	15	1.7	1.0-2.9
1994	235	15.8	13.8-18.0	146	23.1	19.4-27.3	89	10.4	8.3-12.8	60	4.1	3.1-5.3	44	7.2	5.2-9.8	16	1.9	1.1-3.1
1995	180	12.2	10.4-14.1	127	19.8	16.4-23.6	53	6.2	4.6-8.1	51	3.5	2.6-4.5	29	4.9	3.2-7.1	22	2.5	1.5-3.8
1996	236	15.9	13.9-18.1	158	24.2	20.5-28.3	78	9.3	7.3-11.6	55	3.7	2.8-4.9	34	5.5	3.8-7.7	21	2.4	1.5-3.7
1997	180	11.8	10.1-13.7	126	19.2	16.0-23.0	54	6.0	4.5-7.9	60	4.0	3.1-5.2	45	7.1	5.2-9.5	15	1.7	1.0-2.9
1998	199	13.1	11.3-15.0	128	19.1	15.9-22.8	71	8.3	6.5-10.5	49	3.3	2.4-4.3	30	4.9	3.3-7.0	19	2.1	1.2-3.3
1999	199	12.8	11.1-14.7	138	20.4	17.1-24.2	61	6.8	5.2-8.7	53	3.5	2.6-4.6	40	5.9	4.2-8.1	13	1.4	0.7-2.4
2000	161	10.3	8.8-12.1	112	16.3	13.4-19.7	49	5.5	4.1-7.4	44	2.9	2.1-3.9	31	4.7	3.2-6.8	13	1.3	0.7-2.3
2001	193	12.3	10.6-14.2	122	18.3	15.1-21.9	71	7.8	6.1-9.8	47	3.0	2.2-4.0	29	4.5	3.0-6.4	18	1.8	1.0-2.9
2002	160	10.0	8.5-11.6	105	14.8	12.0-17.9	55	6.0	4.5-7.8	40	2.5	1.8-3.5	23	3.6	2.3-5.5	17	1.8	1.0-2.9
2003	161	9.9	8.4-11.5	113	16.0	13.1-19.3	48	5.1	3.7-6.8	36	2.2	1.5-3.0	25	3.6	2.3-5.4	11	1.2	0.6-2.1
2004	194	11.9	10.3-13.7	131	18.3	15.2-21.8	63	7.1	5.4-9.1	50	3.1	2.3-4.1	28	4.1	2.7-6.0	22	2.4	1.5-3.7
2005	158	9.5	8.1-11.1	109	15.1	12.4-18.3	49	5.1	3.7-6.8	40	2.4	1.7-3.3	23	3.5	2.2-5.3	17	1.7	1.0-2.8
2006	185	11.1	9.6-12.9	133	17.9	14.9-21.3	52	5.6	4.2-7.4	35	2.1	1.5-3.0	27	3.9	2.6-5.8	8	0.8	0.3-1.7
2007	201	12.3	10.6-14.1	137	19.0	15.9-22.5	64	7.1	5.5-9.2	37	2.2	1.6-3.1	24	3.5	2.2-5.2	13	1.2	0.6-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
Oral cavity and pharynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	224	13.3	11.6-15.2	157	21.4	18.2-25.1	67	6.9	5.3-8.8	39	2.2	1.6-3.0	30	4.0	2.7-5.8	9	0.9	0.4-1.7
2009	196	11.6	10.0-13.3	144	19.0	16.0-22.5	52	5.7	4.2-7.6	44	2.5	1.8-3.4	30	4.3	2.9-6.1	14	1.4	0.8-2.5
2010	198	11.3	9.8-13.1	120	15.7	13.0-18.9	78	8.2	6.4-10.3	50	2.9	2.2-3.9	34	4.8	3.3-6.7	16	1.6	0.9-2.7
2011	193	10.8	9.3-12.5	135	17.2	14.4-20.5	58	5.6	4.2-7.3	43	2.4	1.8-3.3	29	4.0	2.7-5.8	14	1.3	0.7-2.3
2012	176	9.6	8.2-11.2	119	14.7	12.1-17.6	57	5.4	4.1-7.1	41	2.2	1.6-3.0	29	3.5	2.3-5.1	12	1.1	0.6-2.0
2013	194	10.3	8.9-11.9	134	16.0	13.4-19.1	60	5.6	4.2-7.2	48	2.6	1.9-3.4	34	4.4	3.0-6.1	14	1.2	0.6-2.1
2014	200	10.6	9.1-12.2	115	13.9	11.4-16.7	85	7.9	6.3-9.8	43	2.2	1.6-3.0	29	3.6	2.4-5.2	14	1.3	0.7-2.2
2015	248	13.2	11.6-15.0	164	19.9	16.9-23.2	84	8.0	6.3-10.0	52	2.7	2.0-3.5	37	4.5	3.2-6.3	15	1.3	0.7-2.3
2016	232	12.1	10.5-13.8	169	19.7	16.8-23.0	63	6.0	4.6-7.8	42	2.2	1.5-2.9	29	3.5	2.4-5.1	13	1.1	0.6-2.0
2017	211	10.6	9.2-12.2	140	15.8	13.3-18.8	71	6.5	5.0-8.3	45	2.2	1.6-2.9	35	4.0	2.8-5.6	10	0.8	0.4-1.6
2018	226	11.2	9.8-12.8	150	16.6	14.0-19.6	76	6.9	5.4-8.7	37	1.7	1.2-2.4	21	2.3	1.4-3.6	16	1.3	0.7-2.2
2019	213	10.6	9.2-12.2	137	15.5	12.9-18.3	76	6.6	5.2-8.4	33	1.5	1.1-2.2	22	2.4	1.5-3.7	11	0.8	0.4-1.6
2020	203	9.9	8.6-11.4	128	14.0	11.7-16.8	75	6.6	5.1-8.4	47	2.2	1.6-2.9	32	3.6	2.4-5.1	15	1.2	0.6-2.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
Esophagus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	143	9.2	7.7-10.9	91	14.1	11.2-17.4	52	5.8	4.3-7.7	114	7.1	5.9-8.6	81	12.0	9.5-15.0	33	3.8	2.6-5.4
1977	132	8.7	7.2-10.3	78	12.3	9.6-15.4	54	6.2	4.6-8.2	102	6.8	5.5-8.3	65	10.4	7.9-13.3	37	4.3	3.0-6.0
1978	113	7.3	6.0-8.8	72	11.0	8.6-13.9	41	4.5	3.2-6.2	112	7.5	6.2-9.1	73	11.7	9.1-14.8	39	4.7	3.3-6.5
1979	121	7.8	6.4-9.3	85	13.3	10.6-16.5	36	4.1	2.8-5.7	101	6.6	5.4-8.0	66	10.7	8.2-13.7	35	3.8	2.7-5.4
1980	133	8.8	7.3-10.5	96	15.4	12.4-18.9	37	4.0	2.8-5.6	116	7.6	6.3-9.2	78	12.9	10.1-16.3	38	4.0	2.8-5.5
1981	126	8.3	6.9-9.9	85	13.3	10.6-16.6	41	4.8	3.4-6.6	94	6.1	4.9-7.4	65	10.0	7.7-12.8	29	3.2	2.1-4.6
1982	118	7.8	6.5-9.4	75	12.0	9.4-15.1	43	4.8	3.5-6.6	82	5.4	4.3-6.7	55	8.8	6.6-11.5	27	3.2	2.1-4.7
1983	133	9.0	7.5-10.7	93	14.9	11.9-18.3	40	4.9	3.4-6.7	93	6.2	5.0-7.6	64	10.3	7.9-13.3	29	3.2	2.1-4.7
1984	115	7.4	6.1-9.0	75	11.7	9.2-14.8	40	4.3	3.1-5.9	93	6.2	5.0-7.7	58	9.5	7.1-12.3	35	4.0	2.7-5.6
1985	120	8.2	6.7-9.8	93	15.0	12.1-18.5	27	3.2	2.1-4.7	88	5.9	4.7-7.2	69	10.6	8.2-13.5	19	2.2	1.3-3.6
1986	105	6.7	5.5-8.1	75	11.8	9.2-14.9	30	3.3	2.2-4.8	100	6.5	5.3-8.0	81	13.1	10.4-16.4	19	2.2	1.3-3.5
1987	98	6.4	5.2-7.9	50	7.9	5.8-10.5	48	5.4	3.9-7.2	83	5.5	4.3-6.8	50	8.0	5.9-10.7	33	3.8	2.6-5.4
1988	110	7.3	6.0-8.9	68	10.8	8.4-13.8	42	4.6	3.3-6.3	87	5.7	4.5-7.0	56	9.0	6.7-11.7	31	3.4	2.3-4.9
1989	115	7.8	6.4-9.3	69	11.6	9.0-14.7	46	5.3	3.8-7.1	104	7.1	5.8-8.6	61	10.1	7.7-13.0	43	4.9	3.5-6.6
1990	107	7.2	5.9-8.7	66	10.5	8.1-13.5	41	4.5	3.2-6.1	88	5.9	4.7-7.3	54	9.0	6.7-11.8	34	3.6	2.5-5.1
1991	100	6.7	5.4-8.1	67	11.2	8.7-14.3	33	3.6	2.5-5.1	86	5.8	4.6-7.1	53	8.9	6.6-11.7	33	3.7	2.6-5.3
1992	102	6.8	5.5-8.3	61	9.8	7.5-12.7	41	4.7	3.3-6.4	75	5.0	3.9-6.3	51	8.5	6.3-11.2	24	2.6	1.7-4.0
1993	95	6.5	5.2-7.9	62	10.0	7.6-12.8	33	3.8	2.6-5.4	72	4.8	3.8-6.1	46	7.8	5.7-10.4	26	3.0	1.9-4.4
1994	109	7.4	6.0-8.9	67	10.9	8.4-13.9	42	4.7	3.4-6.4	74	5.0	3.9-6.3	48	8.3	6.1-11.0	26	2.9	1.9-4.3
1995	86	5.8	4.7-7.2	52	8.4	6.2-11.0	34	4.0	2.7-5.6	74	5.0	4.0-6.3	49	8.1	6.0-10.7	25	2.7	1.7-4.0
1996	92	6.2	5.0-7.6	59	9.4	7.1-12.2	33	3.8	2.6-5.3	80	5.4	4.3-6.7	55	9.0	6.8-11.8	25	2.8	1.8-4.1
1997	88	5.9	4.7-7.3	58	9.6	7.2-12.4	30	3.4	2.3-4.8	73	5.0	3.9-6.2	48	7.8	5.7-10.4	25	2.8	1.8-4.2
1998	93	6.2	5.0-7.6	61	10.1	7.7-13.0	32	3.7	2.5-5.3	70	4.7	3.7-5.9	43	7.1	5.1-9.6	27	3.1	2.0-4.5
1999	82	5.4	4.3-6.7	58	9.3	7.0-12.0	24	2.6	1.6-3.9	60	4.0	3.0-5.1	42	6.8	4.8-9.2	18	2.0	1.2-3.2
2000	94	6.2	5.0-7.6	61	9.7	7.4-12.4	33	3.6	2.5-5.1	87	5.7	4.5-7.0	55	8.7	6.5-11.4	32	3.5	2.4-5.0
2001	71	4.5	3.5-5.7	48	7.8	5.7-10.3	23	2.4	1.5-3.7	54	3.5	2.6-4.5	40	6.4	4.6-8.8	14	1.5	0.8-2.6
2002	81	5.1	4.1-6.4	53	8.0	6.0-10.5	28	3.0	2.0-4.3	61	3.9	3.0-5.0	45	7.2	5.2-9.6	16	1.6	0.9-2.7
2003	81	5.0	4.0-6.3	49	7.3	5.4-9.7	32	3.3	2.3-4.8	72	4.5	3.5-5.7	44	6.5	4.7-8.8	28	2.9	1.9-4.3
2004	99	6.0	4.9-7.3	71	9.9	7.7-12.6	28	2.9	1.9-4.3	63	3.8	2.9-4.9	40	5.8	4.2-8.0	23	2.4	1.5-3.6
2005	74	4.5	3.5-5.6	39	5.5	3.9-7.5	35	3.7	2.6-5.2	78	4.7	3.7-5.8	55	7.8	5.8-10.1	23	2.4	1.5-3.6
2006	75	4.5	3.6-5.7	55	8.1	6.1-10.6	20	2.0	1.2-3.2	58	3.6	2.7-4.6	28	4.1	2.7-5.9	30	3.1	2.1-4.5
2007	80	4.8	3.8-6.0	56	7.9	6.0-10.3	24	2.5	1.6-3.7	68	4.1	3.2-5.2	50	7.0	5.2-9.3	18	1.9	1.1-3.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Esophagus

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	82	4.9	3.9-6.1	52	7.4	5.5-9.7	30	3.0	2.0-4.3	57	3.4	2.6-4.4	37	5.4	3.8-7.5	20	2.1	1.3-3.3
2009	79	4.5	3.5-5.6	57	7.7	5.8-10.0	22	2.0	1.3-3.2	57	3.3	2.5-4.3	40	5.7	4.0-7.8	17	1.5	0.9-2.5
2010	80	4.7	3.7-5.8	50	6.7	5.0-8.9	30	3.0	2.0-4.4	74	4.2	3.3-5.3	53	7.3	5.4-9.6	21	2.0	1.2-3.1
2011	71	4.0	3.1-5.1	52	6.9	5.1-9.1	19	1.9	1.1-3.1	62	3.4	2.6-4.4	44	5.7	4.1-7.7	18	1.7	1.0-2.8
2012	74	4.0	3.1-5.1	50	6.5	4.8-8.6	24	2.3	1.4-3.4	49	2.8	2.0-3.7	31	4.2	2.8-6.0	18	1.7	1.0-2.8
2013	68	3.7	2.9-4.8	46	6.0	4.4-8.1	22	2.0	1.3-3.2	49	2.7	2.0-3.6	30	3.9	2.6-5.6	19	1.9	1.1-3.0
2014	83	4.3	3.4-5.4	58	6.8	5.1-8.8	25	2.4	1.5-3.6	54	3.0	2.2-3.9	39	5.0	3.5-6.8	15	1.3	0.7-2.3
2015	58	3.0	2.3-3.9	37	4.4	3.1-6.1	21	1.8	1.1-2.9	53	2.7	2.0-3.6	36	4.3	3.0-6.1	17	1.4	0.8-2.3
2016	67	3.3	2.6-4.3	47	5.7	4.2-7.7	20	1.6	1.0-2.6	45	2.3	1.6-3.1	30	3.6	2.4-5.2	15	1.2	0.7-2.1
2017	58	2.9	2.2-3.8	38	4.5	3.1-6.2	20	1.7	1.0-2.7	47	2.3	1.7-3.1	34	4.0	2.8-5.7	13	1.1	0.6-2.0
2018	65	3.1	2.4-3.9	40	4.5	3.2-6.2	25	2.0	1.3-3.0	38	1.8	1.3-2.6	26	3.0	1.9-4.4	12	1.0	0.5-1.8
2019	65	3.1	2.4-4.0	41	4.4	3.1-6.0	24	1.9	1.2-3.0	52	2.3	1.7-3.1	34	3.7	2.5-5.2	18	1.4	0.8-2.3
2020	64	3.1	2.4-4.0	43	4.8	3.5-6.5	21	1.9	1.1-3.0	38	1.7	1.2-2.4	24	2.6	1.7-3.9	14	1.1	0.6-1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
Stomach**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	216	13.7	11.9-15.8	116	18.0	14.7-21.8	100	10.8	8.7-13.2	177	11.6	9.9-13.5	95	15.3	12.2-18.8	82	9.2	7.3-11.5
1977	233	15.3	13.4-17.5	132	22.4	18.6-26.8	101	10.7	8.7-13.1	188	12.3	10.5-14.2	105	17.4	14.1-21.3	83	8.9	7.0-11.1
1978	230	15.2	13.2-17.3	132	22.2	18.4-26.5	98	10.6	8.6-13.0	164	11.1	9.4-12.9	94	16.2	12.9-20.0	70	7.8	6.0-9.9
1979	222	14.8	12.9-17.0	134	22.8	19.0-27.2	88	9.7	7.8-12.1	152	10.1	8.5-11.9	95	16.5	13.2-20.3	57	6.2	4.6-8.1
1980	245	16.4	14.3-18.6	154	26.4	22.2-31.0	91	10.1	8.1-12.4	165	11.2	9.6-13.1	96	17.2	13.8-21.2	69	7.6	5.9-9.7
1981	192	12.9	11.2-14.9	95	16.8	13.4-20.6	97	10.8	8.8-13.3	171	11.5	9.8-13.4	100	17.8	14.3-21.8	71	8.0	6.2-10.1
1982	187	12.7	11.0-14.7	96	17.0	13.7-20.9	91	10.1	8.1-12.5	123	8.4	7.0-10.1	60	11.2	8.4-14.4	63	6.9	5.3-8.9
1983	211	14.1	12.3-16.2	90	15.5	12.4-19.2	121	13.3	11.0-16.0	120	8.0	6.6-9.6	67	11.5	8.8-14.7	53	5.7	4.3-7.5
1984	198	12.9	11.2-14.9	105	17.8	14.5-21.7	93	9.9	8.0-12.2	130	8.5	7.1-10.2	67	11.8	9.1-15.1	63	6.7	5.1-8.7
1985	198	13.3	11.5-15.4	105	18.1	14.7-22.0	93	10.5	8.4-12.9	149	9.9	8.4-11.7	79	13.9	10.9-17.4	70	7.6	5.9-9.7
1986	205	13.4	11.6-15.4	126	21.4	17.8-25.6	79	8.7	6.8-10.9	141	9.2	7.8-10.9	77	13.3	10.5-16.7	64	6.8	5.2-8.7
1987	204	13.3	11.6-15.3	121	20.4	16.9-24.5	83	8.6	6.8-10.7	127	8.3	6.9-9.9	74	12.8	9.9-16.1	53	5.7	4.2-7.5
1988	169	11.2	9.5-13.0	95	15.9	12.8-19.5	74	7.8	6.1-9.9	133	8.7	7.3-10.4	73	12.4	9.7-15.7	60	6.4	4.9-8.3
1989	162	10.8	9.2-12.6	89	15.0	12.0-18.5	73	7.9	6.2-10.0	116	7.7	6.3-9.2	67	11.6	8.9-14.8	49	5.2	3.8-6.9
1990	194	13.1	11.3-15.0	108	19.0	15.5-23.0	86	9.4	7.5-11.7	123	8.3	6.9-9.9	65	11.5	8.9-14.8	58	6.2	4.7-8.0
1991	222	15.0	13.1-17.2	131	22.6	18.8-26.9	91	10.0	8.1-12.4	136	9.2	7.7-10.9	76	13.8	10.8-17.4	60	6.7	5.1-8.6
1992	206	14.0	12.1-16.0	103	17.5	14.2-21.3	103	11.5	9.3-14.0	127	8.7	7.2-10.4	61	10.6	8.0-13.6	66	7.3	5.6-9.3
1993	207	13.9	12.1-16.0	113	19.7	16.2-23.8	94	10.5	8.5-12.9	115	7.8	6.4-9.3	57	10.5	7.9-13.6	58	6.3	4.8-8.2
1994	211	14.2	12.3-16.2	103	17.4	14.1-21.2	108	12.1	9.9-14.7	123	8.3	6.9-9.9	65	11.3	8.7-14.4	58	6.4	4.9-8.4
1995	186	12.5	10.8-14.4	98	16.9	13.6-20.7	88	9.7	7.7-12.0	108	7.3	6.0-8.8	46	8.7	6.3-11.6	62	6.7	5.1-8.7
1996	195	13.1	11.3-15.1	112	19.2	15.7-23.1	83	8.9	7.1-11.1	133	9.0	7.5-10.7	84	15.0	11.9-18.6	49	5.3	3.9-7.1
1997	160	10.6	9.0-12.4	91	15.2	12.2-18.7	69	7.6	5.9-9.7	106	7.1	5.8-8.6	53	9.1	6.8-11.9	53	5.8	4.4-7.7
1998	192	12.8	11.1-14.8	96	16.1	13.0-19.7	96	10.9	8.8-13.4	123	8.2	6.8-9.7	55	9.4	7.1-12.3	68	7.4	5.8-9.5
1999	177	11.8	10.1-13.6	103	17.2	13.9-20.9	74	8.2	6.5-10.4	97	6.4	5.2-7.8	50	8.6	6.4-11.4	47	5.1	3.7-6.8
2000	168	11.0	9.4-12.8	85	13.8	11.0-17.1	83	9.0	7.2-11.3	102	6.6	5.4-8.1	54	9.2	6.9-12.0	48	4.9	3.6-6.6
2001	177	11.4	9.8-13.2	105	16.5	13.4-20.0	72	7.7	6.0-9.8	79	5.1	4.0-6.3	38	6.4	4.5-8.8	41	4.2	3.0-5.7
2002	163	10.4	8.9-12.2	89	13.8	11.1-17.1	74	8.1	6.4-10.3	99	6.3	5.1-7.7	52	8.2	6.1-10.8	47	5.0	3.7-6.7
2003	168	10.6	9.1-12.4	99	15.1	12.2-18.4	69	7.3	5.6-9.2	102	6.4	5.2-7.8	60	9.4	7.1-12.1	42	4.3	3.1-5.9
2004	177	10.9	9.4-12.7	99	14.8	12.0-18.1	78	8.2	6.4-10.2	79	4.8	3.8-6.0	46	7.1	5.1-9.5	33	3.3	2.3-4.7
2005	164	10.1	8.6-11.8	95	14.0	11.3-17.2	69	7.0	5.4-9.0	98	6.0	4.9-7.3	58	8.9	6.7-11.5	40	3.9	2.8-5.4
2006	171	10.4	8.9-12.1	95	13.9	11.2-17.1	76	7.9	6.2-10.0	70	4.3	3.3-5.4	39	5.7	4.0-7.8	31	3.2	2.2-4.6
2007	159	9.6	8.2-11.3	93	13.5	10.8-16.5	66	6.8	5.2-8.7	88	5.3	4.3-6.6	48	7.1	5.2-9.5	40	4.0	2.8-5.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
Stomach**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	160	9.5	8.1-11.2	101	14.2	11.6-17.4	59	6.1	4.6-7.9	82	4.8	3.8-6.0	45	6.6	4.8-8.9	37	3.6	2.5-5.1
2009	150	8.8	7.4-10.3	92	12.5	10.1-15.4	58	5.9	4.5-7.7	82	4.8	3.8-6.0	47	7.0	5.1-9.3	35	3.2	2.2-4.6
2010	179	10.3	8.8-11.9	96	12.9	10.4-15.8	83	8.2	6.5-10.2	60	3.4	2.6-4.4	34	4.9	3.4-6.9	26	2.4	1.5-3.6
2011	139	7.9	6.6-9.4	86	11.8	9.4-14.6	53	5.2	3.8-6.8	78	4.3	3.4-5.4	37	5.0	3.5-6.9	41	3.6	2.6-5.0
2012	152	8.3	7.0-9.7	96	12.3	9.9-15.1	56	5.3	3.9-6.9	61	3.4	2.6-4.4	33	4.3	3.0-6.1	28	2.7	1.7-3.9
2013	166	8.8	7.5-10.3	101	12.7	10.4-15.5	65	6.0	4.6-7.7	65	3.4	2.6-4.4	42	5.4	3.8-7.3	23	2.1	1.3-3.2
2014	155	8.4	7.1-9.8	94	11.6	9.3-14.2	61	5.7	4.3-7.4	79	4.2	3.4-5.3	40	5.1	3.6-7.0	39	3.7	2.6-5.1
2015	156	8.1	6.9-9.6	91	11.1	8.9-13.7	65	5.9	4.5-7.7	62	3.2	2.4-4.1	42	5.2	3.7-7.0	20	1.7	1.0-2.8
2016	155	8.0	6.7-9.4	84	9.9	7.9-12.3	71	6.6	5.1-8.4	68	3.3	2.5-4.2	46	5.4	3.9-7.2	22	1.9	1.2-3.0
2017	139	7.1	5.9-8.4	68	8.1	6.3-10.4	71	6.2	4.8-7.9	60	2.9	2.2-3.8	31	3.6	2.4-5.2	29	2.4	1.6-3.6
2018	142	7.1	5.9-8.4	90	10.3	8.2-12.7	52	4.7	3.4-6.2	71	3.5	2.7-4.5	45	5.2	3.8-7.0	26	2.2	1.4-3.3
2019	164	8.1	6.9-9.5	90	10.2	8.2-12.6	74	6.5	5.0-8.2	81	3.9	3.1-4.9	50	5.5	4.1-7.3	31	2.6	1.7-3.8
2020	114	5.7	4.7-6.9	67	7.4	5.7-9.5	47	4.4	3.2-6.0	59	3.0	2.2-3.9	37	4.3	3.0-6.0	22	1.9	1.2-3.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
Colorectal**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,067	68.2	64.0-72.5	500	81.0	73.8-88.8	567	60.5	55.5-65.9	522	34.6	31.6-37.8	245	43.0	37.5-49.1	277	29.7	26.2-33.6
1977	1,052	68.3	64.2-72.7	499	82.3	75.0-90.2	553	59.6	54.6-64.9	501	33.3	30.4-36.4	259	44.1	38.7-50.1	242	26.7	23.3-30.4
1978	954	62.4	58.5-66.6	456	75.7	68.7-83.2	498	54.4	49.6-59.5	505	34.1	31.1-37.3	249	43.1	37.7-49.0	256	28.7	25.2-32.6
1979	995	65.7	61.6-70.0	475	81.7	74.3-89.6	520	56.4	51.6-61.7	490	32.7	29.8-35.8	233	41.0	35.7-46.8	257	27.8	24.5-31.5
1980	997	65.9	61.8-70.2	478	82.1	74.7-90.1	519	56.2	51.4-61.4	512	34.4	31.4-37.5	252	45.6	39.9-51.8	260	28.5	25.1-32.3
1981	979	64.9	60.9-69.2	457	80.3	72.9-88.3	522	56.3	51.5-61.4	494	32.9	30.1-36.0	214	37.2	32.2-42.7	280	29.9	26.5-33.7
1982	936	62.0	58.0-66.1	432	75.2	68.1-82.9	504	54.8	50.0-59.9	415	28.1	25.4-31.0	210	38.1	33.0-43.8	205	22.3	19.3-25.7
1983	919	60.7	56.8-64.8	445	76.3	69.1-84.0	474	51.3	46.7-56.3	414	28.0	25.3-30.9	197	34.8	30.0-40.2	217	23.5	20.4-26.9
1984	992	65.6	61.5-69.8	460	79.3	72.1-87.1	532	57.3	52.5-62.5	467	30.9	28.2-33.9	238	42.2	36.8-48.1	229	24.7	21.6-28.2
1985	981	64.4	60.4-68.7	483	84.1	76.6-92.1	498	53.6	48.9-58.6	455	30.5	27.7-33.4	212	37.7	32.7-43.2	243	26.0	22.8-29.5
1986	1,007	66.3	62.2-70.6	484	84.1	76.6-92.1	523	56.6	51.8-61.8	411	27.1	24.6-29.9	202	35.8	30.9-41.2	209	22.3	19.3-25.6
1987	848	55.5	51.8-59.4	385	65.8	59.2-72.9	463	49.1	44.7-53.8	398	26.2	23.6-28.9	186	33.3	28.5-38.5	212	22.2	19.3-25.5
1988	793	52.0	48.4-55.8	334	57.5	51.4-64.2	459	48.8	44.4-53.6	402	26.7	24.1-29.5	181	32.9	28.2-38.2	221	23.4	20.4-26.7
1989	748	49.6	46.1-53.3	354	60.7	54.4-67.5	394	43.2	39.0-47.8	363	24.2	21.7-26.8	162	29.3	24.9-34.3	201	21.3	18.4-24.5
1990	742	49.2	45.7-52.9	353	59.5	53.3-66.1	389	42.3	38.2-46.8	382	25.6	23.1-28.3	161	29.2	24.7-34.1	221	23.6	20.6-27.0
1991	827	56.1	52.3-60.1	382	67.7	60.9-75.0	445	49.6	45.0-54.5	371	24.9	22.4-27.5	177	33.0	28.2-38.3	194	20.6	17.8-23.8
1992	791	53.4	49.7-57.2	350	61.5	55.1-68.4	441	49.1	44.6-54.0	336	22.7	20.3-25.2	162	30.3	25.7-35.4	174	18.8	16.1-21.9
1993	836	56.2	52.4-60.1	392	67.3	60.7-74.5	444	49.2	44.7-54.0	363	24.4	22.0-27.1	161	29.1	24.7-34.1	202	21.8	18.9-25.1
1994	894	60.3	56.4-64.4	389	66.7	60.1-73.8	505	56.3	51.4-61.4	340	22.9	20.6-25.5	160	29.0	24.6-33.9	180	19.3	16.6-22.4
1995	778	52.6	48.9-56.4	370	64.1	57.6-71.1	408	45.6	41.2-50.3	341	23.2	20.8-25.8	149	27.2	22.9-32.0	192	21.0	18.1-24.2
1996	784	52.9	49.3-56.8	361	61.2	54.9-68.0	423	47.4	43.0-52.2	332	22.5	20.2-25.1	159	28.8	24.4-33.7	173	18.9	16.2-22.0
1997	840	56.7	52.9-60.7	372	63.1	56.7-69.9	468	51.4	46.9-56.4	337	22.7	20.4-25.3	149	26.5	22.3-31.2	188	20.1	17.3-23.3
1998	831	55.6	51.8-59.5	377	63.5	57.1-70.3	454	50.5	46.0-55.5	317	21.4	19.1-23.9	119	20.9	17.2-25.1	198	21.3	18.4-24.5
1999	778	51.7	48.1-55.5	372	62.3	56.0-69.1	406	45.1	40.8-49.7	328	22.0	19.6-24.5	139	24.8	20.8-29.3	189	20.5	17.7-23.7
2000	793	51.9	48.3-55.6	367	59.6	53.5-66.1	426	46.9	42.5-51.6	300	19.7	17.5-22.0	143	25.1	21.1-29.7	157	16.5	14.0-19.3
2001	754	48.6	45.2-52.2	347	56.2	50.4-62.6	407	43.9	39.7-48.4	282	18.3	16.2-20.6	154	25.7	21.8-30.2	128	13.4	11.1-15.9
2002	831	52.9	49.3-56.6	405	63.2	57.1-69.8	426	45.7	41.4-50.3	334	21.4	19.1-23.8	166	27.8	23.7-32.4	168	17.6	15.0-20.6
2003	751	47.0	43.7-50.5	363	55.4	49.8-61.5	388	41.3	37.2-45.7	288	18.0	16.0-20.3	134	21.6	18.1-25.7	154	15.8	13.4-18.6
2004	758	46.7	43.4-50.1	360	53.8	48.3-59.8	398	41.5	37.5-45.8	260	16.0	14.1-18.1	110	16.9	13.8-20.4	150	14.9	12.6-17.6
2005	759	46.4	43.2-49.9	358	53.1	47.7-59.0	401	41.4	37.4-45.7	256	15.7	13.8-17.8	121	19.2	15.9-23.0	135	13.4	11.2-15.9
2006	707	42.8	39.6-46.1	340	49.7	44.5-55.4	367	37.9	34.0-42.0	268	16.3	14.4-18.4	118	18.4	15.2-22.1	150	15.2	12.9-18.0
2007	670	40.0	37.0-43.2	334	48.4	43.2-53.9	336	34.2	30.6-38.1	246	14.6	12.9-16.6	113	16.5	13.5-19.8	133	13.0	10.9-15.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
Colorectal**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	684	40.7	37.7-43.9	336	47.4	42.4-52.9	348	35.4	31.7-39.5	272	16.0	14.1-18.0	145	21.8	18.3-25.7	127	12.2	10.2-14.6
2009	703	41.1	38.1-44.3	338	46.3	41.5-51.6	365	36.7	33.0-40.8	235	13.7	12.0-15.7	100	14.8	12.0-18.1	135	12.7	10.6-15.1
2010	605	35.1	32.4-38.1	289	39.3	34.8-44.1	316	32.1	28.6-36.0	229	12.8	11.2-14.6	108	15.1	12.4-18.3	121	11.0	9.1-13.2
2011	627	35.5	32.7-38.4	317	42.8	38.1-47.8	310	30.2	26.9-33.9	222	12.4	10.8-14.2	117	16.1	13.3-19.3	105	9.9	8.1-12.1
2012	624	34.6	31.9-37.5	312	40.5	36.1-45.4	312	29.9	26.6-33.5	213	11.4	9.9-13.1	102	13.6	11.1-16.6	111	9.6	7.9-11.7
2013	641	35.0	32.3-37.9	312	39.7	35.4-44.5	329	31.1	27.8-34.8	233	12.3	10.8-14.0	106	13.7	11.2-16.6	127	11.1	9.2-13.2
2014	659	35.5	32.8-38.3	328	40.6	36.3-45.3	331	31.3	28.0-35.0	214	11.3	9.8-12.9	96	12.3	9.9-15.0	118	10.3	8.5-12.5
2015	677	35.8	33.1-38.7	359	43.7	39.2-48.6	318	29.8	26.5-33.4	235	12.1	10.6-13.8	114	14.1	11.6-16.9	121	10.6	8.7-12.8
2016	661	34.3	31.7-37.1	350	41.8	37.5-46.6	311	28.6	25.5-32.1	218	10.9	9.5-12.5	111	13.5	11.0-16.3	107	9.1	7.4-11.1
2017	664	34.2	31.6-37.0	348	40.8	36.6-45.4	316	29.3	26.0-32.8	213	10.4	9.0-11.9	109	12.9	10.6-15.6	104	8.7	7.0-10.6
2018	583	29.5	27.1-32.1	299	34.8	30.9-39.0	284	25.2	22.3-28.5	181	8.8	7.6-10.3	91	10.6	8.5-13.1	90	7.4	5.9-9.2
2019	639	31.7	29.2-34.3	320	36.4	32.5-40.7	319	27.9	24.8-31.3	174	8.3	7.1-9.7	91	10.5	8.4-13.0	83	6.7	5.3-8.4
2020	526	26.2	23.9-28.6	267	30.2	26.6-34.1	259	22.9	20.1-26.1	189	9.2	7.9-10.7	111	12.8	10.5-15.5	78	6.3	5.0-8.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
Colon excluding rectum**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	779	49.5	46.0-53.2	353	56.7	50.7-63.2	426	45.2	40.9-49.9	422	28.2	25.5-31.1	191	34.2	29.2-39.8	231	24.9	21.7-28.4
1977	775	50.5	46.9-54.3	359	59.7	53.4-66.4	416	44.7	40.4-49.3	408	27.1	24.5-29.9	205	35.2	30.3-40.6	203	22.3	19.3-25.8
1978	698	45.9	42.4-49.5	327	54.7	48.7-61.2	371	40.7	36.6-45.2	419	28.2	25.5-31.1	211	36.4	31.5-41.9	208	23.2	20.1-26.7
1979	732	48.4	44.9-52.1	344	59.5	53.2-66.4	388	42.1	37.9-46.6	401	26.7	24.1-29.5	179	31.5	26.9-36.7	222	24.0	20.9-27.5
1980	734	48.6	45.1-52.3	342	59.8	53.4-66.7	392	42.2	38.0-46.7	439	29.6	26.9-32.6	210	38.5	33.2-44.2	229	25.2	21.9-28.7
1981	703	46.7	43.3-50.4	327	58.2	51.9-65.0	376	40.0	36.0-44.4	415	27.6	25.0-30.5	181	31.3	26.8-36.4	234	25.0	21.9-28.5
1982	669	44.3	41.0-47.8	292	51.9	45.9-58.4	377	40.6	36.6-45.0	348	23.5	21.0-26.1	181	33.1	28.3-38.5	167	18.0	15.3-21.0
1983	666	44.0	40.7-47.5	310	53.7	47.7-60.2	356	38.3	34.3-42.5	340	23.0	20.6-25.7	162	28.7	24.3-33.6	178	19.3	16.5-22.4
1984	736	48.5	45.0-52.2	339	59.1	52.8-65.9	397	42.5	38.4-47.0	404	26.7	24.1-29.5	204	36.1	31.2-41.6	200	21.6	18.7-24.9
1985	729	48.1	44.6-51.8	339	60.7	54.3-67.7	390	41.8	37.7-46.2	399	26.7	24.1-29.5	178	31.8	27.2-37.0	221	23.6	20.5-26.9
1986	766	50.4	46.9-54.2	356	62.1	55.6-69.0	410	44.3	40.0-48.9	353	23.3	20.9-25.9	167	29.5	25.1-34.4	186	19.8	17.1-23.0
1987	624	41.0	37.8-44.3	266	46.0	40.5-52.1	358	37.8	34.0-42.0	339	22.3	20.0-24.8	156	28.0	23.6-32.8	183	19.2	16.5-22.2
1988	588	38.7	35.6-42.0	245	42.1	36.9-47.9	343	36.7	32.9-40.9	352	23.3	20.9-25.9	163	29.3	24.9-34.3	189	20.0	17.3-23.2
1989	558	37.1	34.1-40.4	249	43.2	37.9-49.0	309	34.1	30.3-38.2	333	22.1	19.8-24.6	146	26.4	22.2-31.1	187	19.7	17.0-22.8
1990	548	36.4	33.4-39.6	249	42.9	37.6-48.6	299	32.5	28.9-36.5	337	22.6	20.3-25.2	138	25.1	21.0-29.7	199	21.2	18.3-24.4
1991	625	42.4	39.1-45.8	287	51.0	45.1-57.4	338	37.5	33.6-41.8	315	21.1	18.8-23.6	143	26.7	22.4-31.5	172	18.3	15.6-21.3
1992	599	40.5	37.3-43.9	260	46.4	40.8-52.5	339	37.7	33.8-42.0	292	19.8	17.5-22.2	141	26.5	22.2-31.3	151	16.4	13.9-19.3
1993	631	42.3	39.0-45.7	292	50.1	44.4-56.4	339	37.4	33.5-41.7	312	21.1	18.8-23.6	138	25.2	21.1-29.8	174	18.9	16.2-22.0
1994	681	45.9	42.5-49.5	290	49.8	44.2-56.0	391	43.6	39.3-48.2	301	20.3	18.1-22.8	141	25.2	21.1-29.8	160	17.3	14.7-20.2
1995	571	38.7	35.6-42.1	254	45.4	39.9-51.4	317	35.1	31.3-39.2	291	19.8	17.6-22.3	127	23.5	19.5-28.0	164	17.8	15.2-20.8
1996	601	40.5	37.3-43.9	276	47.2	41.7-53.3	325	35.9	32.1-40.1	294	19.9	17.7-22.3	141	25.4	21.3-30.1	153	16.7	14.1-19.6
1997	641	43.4	40.1-46.9	268	46.9	41.4-53.0	373	40.8	36.8-45.3	293	19.9	17.6-22.3	124	22.3	18.5-26.7	169	18.1	15.5-21.2
1998	589	39.5	36.3-42.8	260	44.6	39.2-50.4	329	36.1	32.3-40.3	286	19.3	17.1-21.6	109	19.3	15.7-23.3	177	19.0	16.3-22.0
1999	583	38.8	35.7-42.1	262	44.9	39.5-50.8	321	35.5	31.7-39.7	277	18.6	16.4-20.9	118	21.5	17.7-25.8	159	17.2	14.6-20.1
2000	577	37.9	34.8-41.1	254	42.2	37.1-47.8	323	35.4	31.6-39.6	259	17.0	15.0-19.2	123	22.0	18.2-26.3	136	14.3	11.9-16.9
2001	545	35.2	32.3-38.3	251	41.5	36.4-47.0	294	31.4	27.8-35.2	244	15.9	13.9-18.0	133	22.4	18.7-26.6	111	11.6	9.5-14.0
2002	627	39.9	36.8-43.2	289	45.6	40.4-51.3	338	36.1	32.3-40.2	283	18.1	16.1-20.4	143	24.0	20.1-28.3	140	14.7	12.4-17.4
2003	545	34.2	31.3-37.2	256	40.0	35.2-45.3	289	30.4	27.0-34.2	247	15.5	13.6-17.5	110	18.0	14.8-21.8	137	14.0	11.8-16.7
2004	573	35.3	32.4-38.3	259	38.7	34.0-43.8	314	32.4	28.9-36.3	226	13.9	12.1-15.8	96	14.7	11.9-18.0	130	12.9	10.7-15.3
2005	574	35.1	32.3-38.1	260	39.1	34.4-44.2	314	32.3	28.8-36.1	213	13.1	11.4-15.0	98	15.5	12.5-18.9	115	11.4	9.4-13.7
2006	499	30.2	27.6-33.0	238	35.5	31.1-40.4	261	26.7	23.5-30.3	231	14.0	12.3-16.0	100	15.6	12.7-19.1	131	13.3	11.1-15.8
2007	469	28.1	25.6-30.7	233	34.4	30.1-39.2	236	23.8	20.8-27.1	200	12.0	10.4-13.8	90	13.3	10.7-16.4	110	10.9	8.9-13.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Colon excluding rectum

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	508	30.5	27.9-33.3	236	33.9	29.7-38.6	272	27.8	24.5-31.4	215	12.7	11.0-14.5	110	16.8	13.8-20.3	105	10.1	8.2-12.3
2009	507	29.6	27.0-32.3	234	32.8	28.7-37.4	273	27.0	23.8-30.5	197	11.5	9.9-13.2	82	12.2	9.7-15.2	115	10.7	8.8-13.0
2010	449	26.0	23.6-28.5	209	28.4	24.7-32.6	240	24.1	21.1-27.5	194	10.8	9.3-12.4	89	12.4	9.9-15.3	105	9.4	7.7-11.5
2011	445	25.0	22.7-27.5	223	30.2	26.3-34.5	222	21.3	18.5-24.4	183	10.2	8.8-11.8	96	13.3	10.7-16.2	87	8.2	6.5-10.1
2012	448	24.7	22.4-27.1	211	27.7	24.0-31.8	237	22.4	19.6-25.5	177	9.4	8.0-10.9	85	11.4	9.1-14.2	92	7.8	6.2-9.6
2013	447	24.2	22.0-26.6	211	26.9	23.4-30.9	236	22.1	19.3-25.2	186	9.8	8.4-11.3	78	10.2	8.1-12.8	108	9.3	7.6-11.3
2014	473	25.4	23.1-27.8	226	28.2	24.6-32.2	247	23.2	20.3-26.4	165	8.6	7.3-10.1	71	9.2	7.2-11.6	94	8.2	6.6-10.1
2015	460	24.4	22.2-26.8	234	29.1	25.5-33.2	226	20.9	18.2-24.0	192	9.9	8.5-11.4	94	11.5	9.2-14.1	98	8.7	7.0-10.7
2016	455	23.5	21.3-25.8	227	27.4	23.9-31.3	228	20.7	18.1-23.7	163	8.1	6.9-9.5	79	9.6	7.6-12.1	84	7.2	5.7-8.9
2017	456	23.3	21.1-25.6	226	26.7	23.2-30.4	230	20.9	18.2-23.9	168	8.1	6.9-9.5	84	9.9	7.9-12.3	84	7.0	5.5-8.7
2018	406	20.3	18.3-22.4	201	23.4	20.2-27.0	205	17.8	15.4-20.5	132	6.4	5.3-7.6	64	7.4	5.7-9.5	68	5.5	4.2-7.1
2019	440	21.6	19.6-23.8	216	24.6	21.4-28.2	224	19.1	16.6-21.9	130	6.1	5.1-7.3	68	7.8	6.1-10.0	62	4.9	3.7-6.4
2020	350	17.2	15.4-19.2	166	18.7	15.9-21.8	184	16.1	13.7-18.7	138	6.7	5.6-7.9	81	9.3	7.4-11.7	57	4.6	3.5-6.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
Rectum & rectosigmoid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	288	18.7	16.5-21.0	147	24.3	20.4-28.8	141	15.3	12.8-18.2	100	6.4	5.1-7.8	54	8.8	6.5-11.6	46	4.9	3.5-6.6
1977	277	17.8	15.8-20.1	140	22.6	18.9-26.9	137	14.9	12.4-17.7	93	6.2	5.0-7.6	54	9.0	6.6-11.8	39	4.3	3.0-6.0
1978	256	16.6	14.6-18.8	129	21.0	17.5-25.1	127	13.7	11.3-16.3	86	5.9	4.7-7.4	38	6.6	4.6-9.2	48	5.5	4.0-7.3
1979	263	17.3	15.2-19.5	131	22.1	18.4-26.4	132	14.3	12.0-17.1	89	6.0	4.8-7.4	54	9.5	7.0-12.4	35	3.8	2.6-5.4
1980	263	17.3	15.2-19.5	136	22.4	18.7-26.6	127	14.1	11.7-16.8	73	4.8	3.7-6.0	42	7.2	5.1-9.7	31	3.4	2.3-4.9
1981	276	18.2	16.1-20.5	130	22.1	18.4-26.4	146	16.3	13.7-19.2	79	5.3	4.2-6.6	33	5.9	4.0-8.3	46	4.9	3.6-6.6
1982	267	17.7	15.6-20.0	140	23.3	19.5-27.6	127	14.2	11.8-17.0	67	4.6	3.6-5.9	29	5.0	3.3-7.2	38	4.4	3.1-6.1
1983	253	16.7	14.7-19.0	135	22.6	18.8-26.9	118	13.1	10.8-15.7	74	5.0	3.9-6.3	35	6.2	4.2-8.7	39	4.2	3.0-5.8
1984	256	17.0	15.0-19.3	121	20.2	16.7-24.3	135	14.8	12.4-17.7	63	4.2	3.2-5.4	34	6.1	4.2-8.5	29	3.1	2.1-4.5
1985	252	16.3	14.4-18.5	144	23.4	19.6-27.6	108	11.8	9.6-14.3	56	3.8	2.8-4.9	34	5.8	4.0-8.2	22	2.4	1.5-3.7
1986	241	15.8	13.9-18.0	128	22.0	18.3-26.3	113	12.4	10.2-14.9	58	3.8	2.9-5.0	35	6.3	4.4-8.8	23	2.4	1.5-3.7
1987	224	14.6	12.7-16.6	119	19.8	16.3-23.7	105	11.3	9.2-13.7	59	3.9	2.9-5.0	30	5.3	3.5-7.6	29	3.0	2.0-4.4
1988	205	13.3	11.5-15.3	89	15.4	12.3-19.0	116	12.2	10.0-14.6	50	3.4	2.5-4.5	18	3.6	2.1-5.7	32	3.3	2.3-4.8
1989	190	12.5	10.7-14.4	105	17.5	14.3-21.3	85	9.2	7.3-11.4	30	2.1	1.4-2.9	16	3.0	1.7-4.8	14	1.6	0.9-2.7
1990	194	12.7	11.0-14.7	104	16.6	13.5-20.2	90	9.8	7.8-12.1	45	3.0	2.2-4.0	23	4.1	2.6-6.2	22	2.4	1.5-3.7
1991	202	13.7	11.9-15.8	95	16.7	13.5-20.5	107	12.1	9.9-14.6	56	3.7	2.8-4.9	34	6.3	4.3-8.9	22	2.3	1.5-3.6
1992	192	12.9	11.1-14.8	90	15.1	12.0-18.6	102	11.4	9.3-13.9	44	2.9	2.1-3.9	21	3.8	2.3-5.8	23	2.4	1.5-3.7
1993	205	13.9	12.0-15.9	100	17.2	13.9-21.0	105	11.8	9.6-14.3	51	3.4	2.5-4.4	23	4.0	2.5-6.0	28	3.0	2.0-4.3
1994	213	14.3	12.5-16.4	99	16.9	13.6-20.6	114	12.7	10.5-15.3	39	2.6	1.9-3.6	19	3.8	2.3-5.9	20	2.1	1.3-3.2
1995	207	13.8	12.0-15.9	116	18.7	15.4-22.5	91	10.5	8.5-13.0	50	3.3	2.5-4.4	22	3.7	2.3-5.6	28	3.1	2.1-4.5
1996	183	12.4	10.7-14.3	85	14.0	11.1-17.3	98	11.5	9.3-14.0	38	2.6	1.8-3.6	18	3.3	2.0-5.3	20	2.3	1.4-3.5
1997	199	13.3	11.5-15.3	104	16.2	13.2-19.6	95	10.6	8.6-13.0	44	2.9	2.1-3.9	25	4.1	2.7-6.1	19	2.0	1.2-3.1
1998	242	16.1	14.1-18.3	117	18.9	15.5-22.7	125	14.4	12.0-17.2	31	2.1	1.4-3.0	10	1.6	0.8-3.0	21	2.3	1.4-3.6
1999	195	12.9	11.1-14.8	110	17.4	14.3-21.1	85	9.5	7.6-11.8	51	3.4	2.5-4.5	21	3.3	2.0-5.1	30	3.3	2.2-4.8
2000	216	14.0	12.2-16.0	113	17.4	14.3-21.0	103	11.4	9.3-13.9	41	2.7	1.9-3.6	20	3.2	1.9-4.9	21	2.2	1.4-3.4
2001	209	13.4	11.6-15.3	96	14.8	11.9-18.1	113	12.6	10.3-15.1	38	2.5	1.7-3.4	21	3.3	2.1-5.1	17	1.8	1.0-2.9
2002	204	13.0	11.2-14.9	116	17.6	14.5-21.1	88	9.6	7.7-11.9	51	3.2	2.4-4.3	23	3.9	2.4-5.8	28	2.9	1.9-4.3
2003	206	12.9	11.2-14.8	107	15.4	12.6-18.7	99	10.9	8.8-13.3	41	2.6	1.9-3.5	24	3.6	2.3-5.4	17	1.8	1.0-2.9
2004	185	11.4	9.8-13.1	101	15.2	12.3-18.5	84	9.0	7.2-11.2	34	2.1	1.4-2.9	14	2.2	1.2-3.7	20	2.1	1.3-3.3
2005	185	11.3	9.7-13.1	98	14.0	11.4-17.2	87	9.1	7.3-11.3	43	2.7	1.9-3.6	23	3.7	2.3-5.5	20	2.0	1.2-3.2
2006	208	12.6	10.9-14.4	102	14.2	11.5-17.3	106	11.1	9.1-13.5	37	2.2	1.6-3.1	18	2.8	1.6-4.4	19	2.0	1.2-3.1
2007	201	11.9	10.3-13.7	101	13.9	11.3-17.0	100	10.4	8.4-12.7	46	2.7	1.9-3.6	23	3.2	2.0-4.8	23	2.1	1.3-3.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
Rectum & rectosigmoid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	176	10.2	8.8-11.9	100	13.5	11.0-16.5	76	7.7	6.0-9.7	57	3.3	2.5-4.3	35	5.0	3.5-7.0	22	2.1	1.3-3.3
2009	196	11.5	9.9-13.3	104	13.5	11.0-16.4	92	9.8	7.8-12.0	38	2.3	1.6-3.1	18	2.6	1.5-4.2	20	1.9	1.2-3.1
2010	156	9.2	7.8-10.8	80	10.8	8.6-13.5	76	8.0	6.2-10.0	35	2.0	1.4-2.8	19	2.7	1.6-4.2	16	1.6	0.9-2.6
2011	182	10.5	9.0-12.2	94	12.6	10.1-15.5	88	8.9	7.1-11.1	39	2.2	1.5-3.0	21	2.8	1.7-4.4	18	1.7	1.0-2.8
2012	176	9.9	8.5-11.5	101	12.8	10.4-15.7	75	7.5	5.9-9.5	36	2.0	1.4-2.8	17	2.2	1.3-3.6	19	1.8	1.1-2.9
2013	194	10.8	9.3-12.5	101	12.8	10.4-15.6	93	9.1	7.3-11.2	47	2.6	1.9-3.4	28	3.5	2.3-5.1	19	1.7	1.0-2.8
2014	186	10.1	8.7-11.7	102	12.4	10.1-15.1	84	8.1	6.4-10.1	49	2.7	2.0-3.6	25	3.1	2.0-4.6	24	2.1	1.3-3.3
2015	217	11.4	9.9-13.1	125	14.6	12.1-17.4	92	8.9	7.1-11.0	43	2.2	1.6-3.1	20	2.6	1.6-4.0	23	1.9	1.2-3.0
2016	206	10.8	9.4-12.5	123	14.5	12.0-17.3	83	7.9	6.3-9.9	55	2.8	2.1-3.7	32	3.8	2.6-5.4	23	1.9	1.2-3.0
2017	208	10.9	9.5-12.6	122	14.1	11.7-17.0	86	8.4	6.6-10.5	45	2.2	1.6-3.0	25	3.0	1.9-4.5	20	1.7	1.0-2.8
2018	177	9.2	7.9-10.7	98	11.3	9.2-13.9	79	7.4	5.8-9.3	49	2.5	1.8-3.3	27	3.2	2.1-4.7	22	1.9	1.1-2.9
2019	199	10.1	8.7-11.6	104	11.8	9.6-14.4	95	8.8	7.1-10.9	44	2.2	1.6-3.0	23	2.7	1.7-4.1	21	1.8	1.1-2.9
2020	176	8.9	7.6-10.4	101	11.5	9.4-14.1	75	6.9	5.3-8.7	51	2.6	1.9-3.4	30	3.4	2.3-5.0	21	1.7	1.0-2.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
Liver / intrahepatic bile duct**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	80	4.9	3.9-6.2	59	9.0	6.8-11.7	21	2.2	1.3-3.4	76	4.8	3.7-6.0	53	7.9	5.9-10.4	23	2.6	1.6-3.9
1977	71	4.5	3.5-5.8	54	8.4	6.3-11.1	17	1.8	1.0-2.9	70	4.4	3.4-5.6	46	7.2	5.2-9.7	24	2.5	1.6-3.8
1978	79	5.1	4.1-6.5	51	8.1	6.0-10.8	28	3.0	2.0-4.5	75	5.1	4.0-6.4	51	8.7	6.4-11.6	24	2.8	1.8-4.2
1979	69	4.5	3.5-5.7	49	8.0	5.8-10.6	20	2.1	1.3-3.3	86	5.8	4.6-7.1	54	9.3	6.9-12.2	32	3.6	2.4-5.1
1980	83	5.5	4.4-6.9	54	8.8	6.6-11.6	29	3.4	2.2-4.9	71	4.7	3.6-5.9	43	7.0	5.0-9.5	28	3.1	2.0-4.6
1981	74	5.0	3.9-6.3	44	6.9	5.0-9.3	30	3.3	2.2-4.8	77	5.2	4.1-6.5	42	7.0	5.0-9.6	35	3.8	2.6-5.4
1982	66	4.3	3.3-5.5	55	8.8	6.6-11.6	11	1.1	0.6-2.1	78	5.0	3.9-6.2	55	8.7	6.5-11.3	23	2.4	1.5-3.7
1983	96	6.4	5.2-7.9	63	10.4	7.9-13.3	33	3.7	2.5-5.3	94	6.3	5.1-7.7	62	10.0	7.6-12.9	32	3.5	2.4-5.0
1984	83	5.5	4.4-6.9	55	9.1	6.8-11.9	28	3.2	2.1-4.6	95	6.3	5.1-7.7	56	9.5	7.1-12.4	39	4.1	2.9-5.6
1985	83	5.5	4.3-6.8	59	9.4	7.1-12.2	24	2.6	1.7-4.0	77	5.0	3.9-6.3	41	6.9	4.9-9.4	36	3.9	2.7-5.4
1986	85	5.7	4.5-7.0	62	10.5	8.0-13.5	23	2.5	1.6-3.8	91	6.1	4.9-7.5	58	9.7	7.4-12.6	33	3.6	2.4-5.1
1987	98	6.5	5.2-7.9	59	9.8	7.4-12.8	39	4.3	3.1-6.0	78	5.1	4.0-6.3	49	8.2	6.0-10.9	29	3.2	2.1-4.6
1988	89	6.0	4.8-7.4	59	9.9	7.5-12.8	30	3.2	2.2-4.7	95	6.4	5.2-7.9	60	10.1	7.6-13.0	35	3.8	2.6-5.3
1989	81	5.4	4.3-6.7	55	9.2	6.9-12.1	26	2.8	1.8-4.2	90	5.9	4.8-7.3	56	9.4	7.1-12.2	34	3.6	2.5-5.1
1990	116	7.7	6.3-9.2	75	12.0	9.4-15.1	41	4.6	3.3-6.3	96	6.3	5.1-7.7	61	9.8	7.5-12.7	35	3.9	2.7-5.4
1991	108	7.2	5.9-8.8	79	12.8	10.1-16.1	29	3.3	2.2-4.9	102	6.9	5.6-8.4	59	10.3	7.8-13.4	43	4.8	3.4-6.5
1992	109	7.4	6.1-8.9	64	10.5	8.0-13.5	45	5.1	3.7-6.9	104	7.1	5.8-8.6	64	10.7	8.2-13.7	40	4.6	3.3-6.3
1993	92	6.3	5.1-7.8	63	10.1	7.7-13.0	29	3.3	2.2-4.8	91	6.2	5.0-7.6	59	10.0	7.5-12.9	32	3.5	2.4-5.0
1994	103	6.9	5.6-8.4	58	9.0	6.8-11.7	45	5.0	3.7-6.8	83	5.6	4.4-6.9	54	8.6	6.4-11.3	29	3.2	2.1-4.6
1995	112	7.5	6.2-9.1	74	11.8	9.2-14.8	38	4.5	3.2-6.2	94	6.4	5.1-7.8	57	9.3	7.0-12.0	37	4.2	3.0-5.8
1996	134	9.0	7.6-10.7	100	16.0	13.0-19.6	34	3.9	2.7-5.4	94	6.3	5.1-7.8	62	10.3	7.9-13.3	32	3.6	2.4-5.1
1997	135	8.9	7.5-10.6	90	14.0	11.2-17.2	45	4.9	3.6-6.6	94	6.3	5.1-7.8	65	10.8	8.3-13.8	29	3.2	2.1-4.7
1998	154	10.3	8.7-12.0	113	17.9	14.7-21.5	41	4.6	3.3-6.3	112	7.5	6.1-9.0	66	10.6	8.2-13.6	46	5.1	3.8-6.9
1999	156	10.3	8.7-12.0	106	16.1	13.1-19.5	50	5.6	4.2-7.5	100	6.6	5.4-8.0	70	11.2	8.7-14.2	30	3.2	2.2-4.7
2000	144	9.4	7.9-11.0	94	13.9	11.2-17.1	50	5.6	4.2-7.5	109	7.1	5.8-8.6	66	10.0	7.7-12.8	43	4.8	3.5-6.5
2001	156	10.1	8.6-11.8	112	16.6	13.6-20.0	44	4.8	3.5-6.6	106	6.8	5.6-8.3	68	10.3	8.0-13.1	38	4.1	2.9-5.6
2002	163	10.3	8.7-12.0	117	16.7	13.8-20.1	46	5.0	3.6-6.7	107	6.8	5.6-8.3	68	10.1	7.8-12.8	39	4.1	2.9-5.6
2003	149	9.4	7.9-11.0	110	15.6	12.8-18.8	39	4.3	3.0-5.9	103	6.4	5.2-7.8	66	9.9	7.6-12.6	37	3.8	2.7-5.3
2004	172	10.6	9.1-12.4	130	18.5	15.4-22.1	42	4.7	3.3-6.4	118	7.3	6.1-8.8	84	12.3	9.8-15.3	34	3.6	2.5-5.1
2005	186	11.3	9.8-13.1	141	19.4	16.3-23.0	45	4.9	3.6-6.7	112	6.7	5.5-8.1	81	11.2	8.9-14.0	31	3.3	2.2-4.7
2006	206	12.5	10.8-14.3	148	20.4	17.2-24.0	58	6.2	4.7-8.0	117	7.1	5.9-8.5	84	11.9	9.5-14.8	33	3.4	2.4-4.9
2007	228	13.5	11.8-15.4	172	23.0	19.6-26.7	56	5.6	4.2-7.4	129	7.8	6.5-9.3	90	12.4	9.9-15.3	39	4.3	3.0-6.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
Liver / intrahepatic bile duct**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	208	12.2	10.6-14.1	156	20.9	17.7-24.5	52	5.4	4.0-7.2	115	6.7	5.5-8.1	81	11.1	8.8-13.9	34	3.4	2.4-4.9
2009	202	11.6	10.1-13.4	147	19.3	16.3-22.8	55	5.5	4.1-7.3	134	7.9	6.6-9.3	97	13.1	10.6-16.0	37	3.8	2.7-5.3
2010	216	12.4	10.7-14.2	141	18.0	15.1-21.3	75	7.6	6.0-9.6	135	7.7	6.4-9.1	97	12.9	10.4-15.8	38	3.5	2.4-4.8
2011	196	10.8	9.3-12.5	135	17.0	14.2-20.2	61	5.7	4.3-7.4	131	7.2	6.0-8.6	82	10.5	8.3-13.0	49	4.6	3.4-6.2
2012	225	12.1	10.6-13.9	164	20.1	17.1-23.5	61	5.7	4.3-7.4	129	7.0	5.9-8.4	87	11.2	8.9-13.9	42	3.8	2.8-5.3
2013	199	10.5	9.1-12.1	133	16.1	13.5-19.2	66	6.0	4.6-7.7	136	7.1	5.9-8.4	87	10.8	8.6-13.4	49	4.3	3.2-5.8
2014	217	11.1	9.7-12.7	143	16.5	13.9-19.5	74	6.4	5.0-8.2	115	6.0	5.0-7.3	74	8.8	6.9-11.1	41	3.6	2.6-5.0
2015	219	11.1	9.6-12.7	141	16.5	13.9-19.6	78	6.8	5.3-8.6	133	6.7	5.6-7.9	90	10.7	8.5-13.2	43	3.6	2.6-5.0
2016	237	11.7	10.3-13.4	161	18.0	15.3-21.1	76	6.6	5.2-8.3	135	6.6	5.5-7.8	93	10.5	8.5-13.0	42	3.5	2.5-4.8
2017	223	11.2	9.7-12.8	155	17.7	15.0-20.8	68	5.8	4.5-7.4	117	5.8	4.8-7.0	77	8.9	7.0-11.2	40	3.4	2.4-4.7
2018	181	8.7	7.5-10.1	124	13.3	11.0-15.9	57	5.0	3.7-6.6	115	5.4	4.5-6.6	71	8.0	6.2-10.1	44	3.7	2.7-5.1
2019	188	9.2	7.9-10.6	123	13.9	11.6-16.7	65	5.2	4.0-6.8	119	5.6	4.6-6.7	71	8.0	6.3-10.2	48	3.9	2.8-5.2
2020	156	7.5	6.4-8.9	92	10.1	8.1-12.5	64	5.5	4.2-7.1	89	4.1	3.3-5.1	55	6.0	4.5-7.8	34	2.7	1.8-3.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Pancreas

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	237	15.3	13.4-17.5	128	21.5	17.7-25.8	109	11.5	9.4-14.0	217	14.1	12.2-16.2	116	19.9	16.3-24.1	101	10.9	8.9-13.4
1977	220	14.6	12.7-16.7	90	15.5	12.3-19.2	130	14.1	11.7-16.8	205	13.5	11.7-15.5	84	14.2	11.2-17.8	121	13.0	10.7-15.6
1978	211	13.5	11.7-15.5	92	15.4	12.3-19.0	119	12.4	10.2-14.9	194	12.4	10.7-14.3	90	14.6	11.7-18.1	104	11.0	8.9-13.4
1979	208	13.6	11.8-15.6	101	16.3	13.2-19.9	107	11.6	9.4-14.0	180	11.8	10.1-13.7	84	14.0	11.1-17.4	96	10.3	8.3-12.7
1980	201	13.5	11.7-15.6	100	17.7	14.3-21.6	101	10.7	8.7-13.1	206	13.8	11.9-15.8	101	17.2	13.9-21.0	105	11.4	9.3-13.9
1981	208	13.5	11.7-15.6	108	18.3	14.9-22.2	100	10.4	8.4-12.7	151	9.9	8.4-11.7	77	13.2	10.3-16.5	74	7.6	6.0-9.6
1982	218	14.5	12.6-16.6	93	16.2	13.0-19.9	125	13.3	11.0-15.9	188	12.4	10.7-14.4	75	13.2	10.3-16.6	113	11.8	9.7-14.2
1983	229	15.1	13.2-17.3	112	18.6	15.2-22.4	117	12.7	10.5-15.3	187	12.4	10.6-14.3	90	15.0	12.0-18.5	97	10.6	8.5-13.0
1984	228	15.2	13.3-17.3	104	17.9	14.5-21.7	124	13.3	11.1-16.0	220	14.7	12.8-16.8	100	17.8	14.4-21.8	120	12.9	10.7-15.5
1985	178	11.6	9.9-13.4	82	13.2	10.5-16.5	96	10.3	8.3-12.6	155	10.3	8.7-12.1	73	12.2	9.5-15.4	82	8.9	7.0-11.1
1986	211	14.1	12.3-16.2	103	17.8	14.4-21.7	108	11.8	9.6-14.3	182	11.9	10.3-13.8	79	13.5	10.6-16.9	103	10.9	8.9-13.2
1987	179	11.8	10.1-13.7	73	12.2	9.5-15.4	106	11.4	9.3-13.8	151	10.1	8.5-11.8	67	11.5	8.9-14.6	84	9.2	7.3-11.5
1988	202	13.5	11.7-15.5	84	14.5	11.5-18.1	118	12.5	10.3-15.0	138	9.1	7.6-10.8	62	10.5	8.0-13.5	76	8.0	6.3-10.0
1989	201	13.5	11.7-15.5	91	16.1	12.9-19.8	110	12.0	9.8-14.5	174	11.6	10.0-13.5	72	12.7	9.9-16.1	102	10.8	8.8-13.2
1990	179	11.9	10.3-13.8	82	14.4	11.4-17.9	97	10.6	8.5-12.9	179	12.0	10.3-13.9	85	15.4	12.2-19.1	94	10.3	8.3-12.7
1991	201	13.6	11.8-15.6	85	14.6	11.6-18.2	116	12.7	10.5-15.3	169	11.5	9.8-13.4	72	12.5	9.7-15.7	97	10.7	8.7-13.1
1992	197	13.4	11.6-15.4	98	17.0	13.7-20.8	99	11.0	8.9-13.4	153	10.4	8.8-12.2	73	12.6	9.8-15.9	80	8.7	6.9-10.9
1993	174	11.7	10.0-13.6	79	13.5	10.6-16.9	95	10.2	8.2-12.5	140	9.4	7.9-11.1	70	12.2	9.4-15.4	70	7.4	5.8-9.4
1994	199	13.6	11.8-15.6	85	14.1	11.2-17.5	114	12.7	10.5-15.3	178	12.2	10.5-14.1	68	12.0	9.3-15.2	110	11.9	9.8-14.4
1995	179	12.1	10.4-14.0	97	16.8	13.6-20.6	82	9.0	7.1-11.2	160	10.8	9.2-12.6	78	13.8	10.9-17.3	82	8.9	7.1-11.2
1996	186	12.5	10.8-14.4	83	14.2	11.3-17.7	103	11.4	9.3-13.9	164	11.0	9.4-12.9	74	12.7	10.0-16.0	90	9.9	7.9-12.2
1997	208	14.1	12.2-16.2	99	16.8	13.6-20.5	109	12.1	9.9-14.6	163	11.1	9.4-12.9	78	13.7	10.8-17.2	85	9.3	7.4-11.6
1998	198	13.5	11.7-15.5	91	15.2	12.2-18.8	107	12.0	9.8-14.5	159	10.7	9.1-12.5	75	12.9	10.1-16.2	84	9.2	7.3-11.4
1999	208	13.9	12.1-15.9	86	14.6	11.6-18.1	122	13.7	11.3-16.4	187	12.4	10.7-14.3	87	14.5	11.6-18.0	100	10.8	8.8-13.2
2000	204	13.5	11.7-15.5	80	13.3	10.5-16.6	124	13.6	11.3-16.3	159	10.5	8.9-12.3	61	10.4	7.9-13.4	98	10.6	8.6-13.0
2001	206	13.4	11.6-15.3	85	13.6	10.8-16.9	121	12.9	10.7-15.4	175	11.5	9.8-13.3	63	10.6	8.1-13.6	112	11.8	9.7-14.2
2002	207	13.3	11.5-15.2	97	15.1	12.2-18.5	110	11.6	9.5-14.0	168	10.8	9.2-12.5	66	10.6	8.2-13.5	102	10.4	8.5-12.7
2003	223	14.0	12.2-16.0	105	16.5	13.5-20.0	118	12.3	10.2-14.8	174	11.0	9.4-12.8	88	13.9	11.1-17.1	86	8.9	7.1-11.0
2004	205	12.6	10.9-14.4	91	13.5	10.9-16.7	114	11.7	9.7-14.2	184	11.3	9.8-13.1	82	12.6	10.0-15.6	102	10.2	8.3-12.4
2005	189	11.6	10.0-13.3	93	14.0	11.3-17.2	96	9.9	8.0-12.1	182	11.1	9.5-12.8	86	13.1	10.4-16.2	96	9.8	7.9-12.1
2006	252	15.3	13.4-17.3	128	18.8	15.6-22.4	124	12.8	10.6-15.3	183	11.2	9.6-13.0	90	13.5	10.9-16.7	93	9.4	7.5-11.6
2007	234	14.0	12.2-15.9	113	16.3	13.4-19.7	121	12.3	10.2-14.7	190	11.3	9.8-13.1	102	15.6	12.7-19.0	88	8.8	7.0-10.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Pancreas

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	238	14.1	12.4-16.1	98	14.2	11.5-17.4	140	14.0	11.7-16.6	211	12.5	10.8-14.3	81	11.7	9.3-14.6	130	12.9	10.8-15.4
2009	233	13.5	11.8-15.4	121	17.1	14.2-20.5	112	10.8	8.8-13.0	204	11.9	10.3-13.7	96	13.6	11.0-16.6	108	10.4	8.5-12.6
2010	249	14.4	12.6-16.3	129	17.9	14.9-21.3	120	11.7	9.7-14.1	191	10.8	9.3-12.4	89	12.3	9.9-15.2	102	9.5	7.7-11.6
2011	226	12.6	11.0-14.4	110	14.6	11.9-17.6	116	11.1	9.1-13.4	170	9.5	8.1-11.1	79	10.8	8.6-13.6	91	8.4	6.7-10.4
2012	223	12.2	10.6-14.0	104	13.7	11.2-16.7	119	11.2	9.2-13.5	180	9.9	8.5-11.5	79	10.4	8.2-13.0	101	9.5	7.7-11.6
2013	273	14.6	12.9-16.4	143	18.0	15.1-21.3	130	11.9	9.9-14.3	189	10.0	8.6-11.6	94	12.1	9.7-14.8	95	8.8	7.1-10.8
2014	253	13.3	11.7-15.0	135	16.8	14.1-19.9	118	10.5	8.6-12.6	206	10.6	9.2-12.2	110	13.9	11.4-16.7	96	8.2	6.6-10.2
2015	257	13.4	11.8-15.2	136	16.6	13.9-19.7	121	10.8	8.9-13.0	208	10.6	9.2-12.1	103	12.6	10.3-15.4	105	8.9	7.2-10.9
2016	244	12.2	10.7-13.9	115	13.6	11.2-16.4	129	11.1	9.2-13.3	204	10.1	8.7-11.6	105	12.7	10.3-15.4	99	8.3	6.7-10.2
2017	310	15.3	13.7-17.2	151	17.6	14.9-20.8	159	13.6	11.5-16.0	218	10.7	9.3-12.3	105	12.6	10.3-15.3	113	9.5	7.8-11.5
2018	276	13.4	11.9-15.2	133	15.3	12.8-18.2	143	12.1	10.1-14.3	219	10.1	8.8-11.6	92	10.2	8.2-12.6	127	10.1	8.4-12.1
2019	236	11.3	9.9-12.9	118	13.3	11.0-16.0	118	9.6	7.9-11.6	183	8.7	7.4-10.1	91	10.3	8.3-12.7	92	7.6	6.1-9.4
2020	250	11.8	10.4-13.4	128	14.2	11.8-16.9	122	10.0	8.2-12.0	192	8.8	7.6-10.2	95	10.5	8.5-12.9	97	7.6	6.1-9.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Larynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	113	7.3	6.0-8.8	87	13.4	10.7-16.7	26	3.2	2.0-4.7	53	3.3	2.4-4.3	41	6.0	4.3-8.3	12	1.4	0.7-2.5
1977	106	6.7	5.5-8.1	90	13.7	10.9-16.9	16	1.8	1.0-3.0	42	2.7	2.0-3.7	36	5.7	3.9-8.0	6	0.9	0.3-1.9
1978	105	7.0	5.7-8.5	84	13.2	10.4-16.4	21	2.3	1.4-3.6	55	3.6	2.7-4.7	42	6.6	4.7-9.0	13	1.4	0.7-2.5
1979	115	7.4	6.1-8.9	92	14.1	11.3-17.3	23	2.5	1.5-3.8	38	2.5	1.7-3.4	32	5.2	3.5-7.4	6	0.7	0.2-1.6
1980	101	6.8	5.5-8.3	84	13.5	10.7-16.8	17	2.2	1.2-3.5	47	3.0	2.2-4.0	41	6.5	4.6-8.9	6	0.7	0.3-1.6
1981	97	6.7	5.4-8.2	80	12.8	10.1-15.9	17	2.0	1.2-3.3	34	2.3	1.6-3.3	25	4.1	2.6-6.1	9	1.1	0.5-2.1
1982	84	5.7	4.5-7.0	68	10.7	8.3-13.6	16	1.9	1.0-3.1	42	2.8	2.0-3.9	29	4.7	3.1-6.8	13	1.5	0.8-2.6
1983	112	7.3	6.0-8.9	88	13.3	10.7-16.5	24	2.8	1.8-4.2	43	3.0	2.1-4.0	38	6.2	4.4-8.6	5	0.6	0.2-1.5
1984	130	8.5	7.1-10.2	99	15.4	12.5-18.8	31	3.4	2.3-4.9	35	2.3	1.6-3.2	29	4.4	2.9-6.3	6	0.8	0.3-1.8
1985	107	7.2	5.8-8.7	89	13.7	10.9-16.9	18	2.1	1.2-3.4	51	3.3	2.4-4.3	42	6.3	4.5-8.5	9	0.9	0.4-1.8
1986	118	7.9	6.5-9.5	85	13.5	10.8-16.8	33	4.0	2.7-5.7	43	2.9	2.1-3.9	32	5.2	3.5-7.4	11	1.4	0.7-2.5
1987	110	7.4	6.1-8.9	85	13.6	10.8-16.8	25	3.1	2.0-4.7	46	3.1	2.2-4.1	35	5.7	3.9-8.0	11	1.1	0.6-2.1
1988	132	9.1	7.6-10.8	94	14.5	11.7-17.8	38	5.2	3.6-7.3	45	3.0	2.2-4.0	37	6.0	4.2-8.3	8	0.9	0.4-1.9
1989	153	10.6	8.9-12.4	96	14.9	12.0-18.3	57	7.3	5.5-9.6	37	2.5	1.7-3.4	25	3.9	2.5-5.8	12	1.4	0.7-2.5
1990	102	6.8	5.5-8.3	82	13.0	10.3-16.2	20	2.4	1.4-3.7	37	2.4	1.7-3.4	32	5.1	3.5-7.2	5	0.6	0.2-1.4
1991	112	7.5	6.2-9.1	79	12.3	9.7-15.3	33	4.1	2.8-5.8	48	3.2	2.3-4.2	35	5.2	3.6-7.3	13	1.6	0.8-2.7
1992	115	7.8	6.4-9.3	79	12.5	9.9-15.7	36	4.3	3.0-6.0	39	2.7	1.9-3.6	32	5.3	3.6-7.5	7	0.9	0.3-1.9
1993	99	6.7	5.4-8.1	79	12.9	10.2-16.1	20	2.3	1.4-3.6	55	3.7	2.8-4.8	40	6.7	4.7-9.1	15	1.7	1.0-2.9
1994	107	7.3	6.0-8.8	75	12.4	9.7-15.6	32	3.9	2.7-5.6	35	2.4	1.7-3.3	21	3.4	2.1-5.3	14	1.7	0.9-2.9
1995	74	5.0	3.9-6.2	57	8.8	6.6-11.5	17	2.0	1.2-3.3	46	3.1	2.3-4.2	35	5.7	3.9-8.0	11	1.3	0.6-2.3
1996	94	6.4	5.2-7.8	78	12.4	9.8-15.5	16	1.9	1.1-3.1	37	2.5	1.8-3.4	30	5.0	3.4-7.2	7	0.8	0.3-1.7
1997	84	5.6	4.5-7.0	70	11.0	8.5-13.9	14	1.6	0.9-2.7	34	2.3	1.6-3.2	27	4.3	2.8-6.3	7	0.8	0.3-1.7
1998	82	5.4	4.3-6.8	57	9.2	6.9-12.0	25	2.9	1.9-4.3	28	1.9	1.2-2.7	24	3.7	2.4-5.6	4	0.5	0.1-1.3
1999	72	4.7	3.7-5.9	56	8.7	6.5-11.3	16	1.9	1.1-3.2	28	1.9	1.2-2.7	20	3.3	2.0-5.2	8	0.9	0.4-1.8
2000	71	4.6	3.6-5.8	54	8.3	6.2-10.9	17	2.0	1.1-3.2	33	2.2	1.5-3.1	24	3.9	2.5-5.8	9	1.0	0.5-2.0
2001	53	3.4	2.5-4.4	39	5.9	4.2-8.1	14	1.6	0.8-2.6	18	1.2	0.7-1.8	15	2.3	1.3-3.8	3	0.3	0.1-1.0
2002	61	3.9	3.0-5.0	53	8.0	6.0-10.5	8	1.0	0.4-1.9	19	1.2	0.7-1.8	14	2.0	1.1-3.5	5	0.5	0.2-1.2
2003	77	4.9	3.8-6.1	56	8.6	6.5-11.2	21	2.4	1.5-3.7	28	1.8	1.2-2.6	20	3.1	1.9-4.8	8	0.8	0.3-1.6
2004	64	3.9	3.0-5.0	54	7.8	5.8-10.2	10	1.1	0.5-2.1	25	1.5	1.0-2.3	21	3.1	1.9-4.7	4	0.5	0.1-1.2
2005	63	3.9	3.0-4.9	48	6.9	5.1-9.2	15	1.7	0.9-2.8	26	1.6	1.0-2.3	20	2.9	1.7-4.4	6	0.6	0.2-1.4
2006	54	3.3	2.4-4.3	45	6.5	4.7-8.8	9	0.9	0.4-1.8	16	1.0	0.6-1.6	14	2.2	1.2-3.6	2	0.2	0.0-0.8
2007	61	3.6	2.7-4.6	47	6.3	4.6-8.4	14	1.4	0.8-2.4	22	1.3	0.8-2.0	17	2.5	1.4-3.9	5	0.5	0.2-1.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Larynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	69	4.1	3.2-5.2	49	6.8	5.0-9.0	20	2.1	1.3-3.3	18	1.1	0.7-1.8	15	2.2	1.2-3.6	3	0.3	0.1-1.0
2009	68	3.8	3.0-4.9	54	7.1	5.3-9.3	14	1.5	0.8-2.6	26	1.5	1.0-2.3	19	2.7	1.6-4.2	7	0.7	0.3-1.5
2010	55	3.1	2.3-4.1	47	6.1	4.5-8.2	8	0.8	0.3-1.7	25	1.4	0.9-2.0	18	2.5	1.5-4.0	7	0.6	0.3-1.4
2011	49	2.7	2.0-3.5	36	4.5	3.1-6.2	13	1.2	0.6-2.1	18	1.0	0.6-1.6	13	1.9	1.0-3.2	5	0.5	0.2-1.3
2012	64	3.4	2.6-4.3	52	6.3	4.7-8.4	12	1.1	0.6-2.0	14	0.7	0.4-1.3	11	1.4	0.7-2.6	3	0.3	0.1-1.0
2013	61	3.3	2.5-4.3	52	6.7	5.0-8.8	9	0.8	0.4-1.7	18	1.0	0.6-1.6	14	1.9	1.0-3.2	4	0.3	0.1-0.9
2014	55	2.9	2.2-3.8	43	5.2	3.7-7.0	12	1.2	0.6-2.1	22	1.2	0.7-1.8	18	2.3	1.3-3.6	4	0.4	0.1-1.1
2015	51	2.6	1.9-3.5	42	4.8	3.5-6.6	9	0.9	0.4-1.8	13	0.7	0.3-1.2	10	1.2	0.6-2.2	3	0.3	0.1-0.9
2016	48	2.4	1.8-3.3	37	4.3	3.0-6.0	11	0.9	0.5-1.8	23	1.0	0.7-1.6	20	2.2	1.3-3.5	3	0.2	0.0-0.8
2017	49	2.4	1.7-3.2	36	3.9	2.7-5.4	13	1.2	0.6-2.1	16	0.8	0.4-1.3	11	1.3	0.6-2.4	5	0.4	0.1-1.0
2018	43	2.2	1.6-2.9	33	3.8	2.6-5.3	10	0.9	0.4-1.7	10	0.5	0.2-0.9	9	1.0	0.4-1.9	1	0.1	0.0-0.6
2019	40	2.0	1.4-2.8	34	3.9	2.7-5.4	6	0.6	0.2-1.3	10	0.5	0.2-0.9	6	0.7	0.2-1.5	4	0.3	0.1-1.0
2020	40	2.0	1.4-2.7	33	3.7	2.5-5.2	7	0.7	0.3-1.6	9	0.4	0.2-0.8	6	0.6	0.2-1.5	3	0.2	0.0-0.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
Lung and bronchus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	927	57.7	54.0-61.7	632	95.9	88.4-103.9	295	31.5	27.9-35.5	655	41.0	37.8-44.3	447	69.2	62.7-76.2	208	22.1	19.1-25.4
1977	842	53.4	49.8-57.2	546	85.9	78.6-93.7	296	31.7	28.1-35.7	682	43.1	39.8-46.5	467	73.1	66.4-80.2	215	22.9	19.9-26.3
1978	837	53.9	50.2-57.7	557	88.0	80.6-95.8	280	31.1	27.5-35.1	704	45.8	42.4-49.4	472	75.9	69.1-83.3	232	25.9	22.5-29.6
1979	930	60.1	56.3-64.2	607	96.7	89.0-104.9	323	35.2	31.4-39.4	696	45.2	41.9-48.8	467	74.8	68.0-82.1	229	25.0	21.8-28.6
1980	853	55.6	51.9-59.6	557	89.3	81.9-97.3	296	33.3	29.5-37.5	693	45.2	41.9-48.8	444	72.6	65.8-79.9	249	27.5	24.1-31.3
1981	880	57.1	53.3-61.0	545	87.3	79.9-95.1	335	36.4	32.5-40.7	680	44.6	41.2-48.1	434	70.1	63.6-77.2	246	26.9	23.6-30.6
1982	913	61.1	57.1-65.3	557	93.6	85.8-101.9	356	40.4	36.2-44.9	700	46.2	42.8-49.8	427	70.8	64.1-78.0	273	30.5	26.9-34.5
1983	913	60.3	56.4-64.4	563	92.0	84.4-100.1	350	39.2	35.1-43.7	727	48.4	44.9-52.1	425	71.5	64.6-78.8	302	33.3	29.6-37.4
1984	940	61.2	57.3-65.3	577	92.5	84.9-100.5	363	40.1	36.0-44.5	721	47.2	43.8-50.8	442	73.0	66.2-80.3	279	30.6	27.1-34.6
1985	923	60.1	56.3-64.2	566	92.6	84.9-100.7	357	39.3	35.3-43.8	743	48.5	45.0-52.2	447	73.6	66.8-80.9	296	32.0	28.4-36.0
1986	865	57.2	53.4-61.2	499	81.7	74.6-89.4	366	40.9	36.7-45.4	704	46.3	42.9-49.9	409	68.7	62.1-75.9	295	32.0	28.4-36.0
1987	943	61.8	57.9-65.9	595	97.3	89.5-105.6	348	37.5	33.6-41.7	754	49.7	46.2-53.4	465	77.8	70.8-85.4	289	31.9	28.3-35.9
1988	875	57.6	53.8-61.6	508	83.6	76.3-91.4	367	40.4	36.3-44.8	758	50.0	46.5-53.7	451	75.3	68.4-82.8	307	33.5	29.8-37.6
1989	858	56.8	53.1-60.8	507	83.5	76.3-91.3	351	39.3	35.2-43.7	673	44.5	41.2-48.0	401	67.3	60.7-74.4	272	30.0	26.5-33.9
1990	870	57.6	53.8-61.6	497	80.2	73.2-87.8	373	42.2	37.9-46.7	722	47.9	44.5-51.6	430	70.7	64.0-77.8	292	32.3	28.6-36.3
1991	961	64.8	60.8-69.1	524	87.1	79.6-95.0	437	49.2	44.6-54.1	723	48.6	45.1-52.3	398	67.2	60.6-74.2	325	36.3	32.4-40.5
1992	977	65.9	61.8-70.2	560	93.2	85.4-101.4	417	47.7	43.2-52.6	741	50.1	46.5-53.8	400	68.3	61.6-75.4	341	37.9	34.0-42.2
1993	876	59.1	55.3-63.2	486	80.4	73.2-88.0	390	44.6	40.3-49.3	669	45.2	41.8-48.8	387	66.0	59.5-73.1	282	31.9	28.2-35.9
1994	967	65.5	61.5-69.8	514	85.8	78.4-93.7	453	52.2	47.5-57.3	723	48.9	45.4-52.6	423	72.4	65.5-79.8	300	34.0	30.2-38.1
1995	918	61.9	57.9-66.0	508	83.3	76.1-91.0	410	46.6	42.2-51.4	711	47.9	44.4-51.5	383	64.3	57.9-71.2	328	36.8	32.9-41.1
1996	921	62.2	58.2-66.4	506	84.2	76.9-92.0	415	48.0	43.5-52.9	712	48.3	44.8-51.9	401	67.4	60.8-74.4	311	35.1	31.3-39.2
1997	883	59.6	55.7-63.6	475	78.0	71.0-85.4	408	46.8	42.4-51.7	600	40.6	37.4-44.0	327	55.6	49.6-62.1	273	30.9	27.3-34.8
1998	847	57.2	53.4-61.2	441	71.6	65.0-78.7	406	46.4	42.0-51.2	605	40.8	37.6-44.2	309	51.4	45.7-57.6	296	33.5	29.8-37.6
1999	886	58.9	55.1-63.0	436	69.8	63.3-76.8	450	50.8	46.2-55.8	652	43.5	40.2-47.0	339	56.1	50.2-62.5	313	35.0	31.2-39.1
2000	923	60.9	57.0-64.9	435	70.0	63.5-77.0	488	54.6	49.9-59.7	616	40.6	37.5-44.0	302	49.8	44.2-55.8	314	34.5	30.8-38.6
2001	951	61.6	57.8-65.7	453	71.8	65.2-78.8	498	54.5	49.7-59.5	669	43.4	40.2-46.9	330	53.6	47.9-59.8	339	36.3	32.5-40.4
2002	873	55.7	52.1-59.5	447	69.2	62.9-76.0	426	46.5	42.2-51.2	629	40.2	37.1-43.5	324	51.1	45.6-57.0	305	32.4	28.8-36.3
2003	895	56.7	53.0-60.5	457	69.5	63.2-76.3	438	47.2	42.9-51.9	658	41.6	38.4-44.9	337	52.3	46.8-58.2	321	33.8	30.2-37.8
2004	921	57.4	53.8-61.3	451	68.1	61.9-74.8	470	50.4	45.9-55.3	576	35.8	33.0-38.9	325	49.6	44.3-55.4	251	26.3	23.1-29.8
2005	897	55.0	51.4-58.7	482	70.9	64.6-77.6	415	44.1	39.9-48.6	652	40.3	37.3-43.6	323	49.0	43.7-54.7	329	34.5	30.8-38.5
2006	888	54.3	50.7-58.0	450	66.5	60.5-73.0	438	45.6	41.4-50.2	552	33.7	30.9-36.6	285	42.7	37.8-48.0	267	27.4	24.1-30.9
2007	907	54.7	51.1-58.4	472	67.8	61.8-74.3	435	45.4	41.2-50.0	599	36.1	33.2-39.1	309	45.8	40.8-51.2	290	29.5	26.1-33.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Lung and bronchus

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	919	54.4	50.9-58.1	418	59.7	54.0-65.8	501	50.9	46.5-55.7	625	37.2	34.3-40.2	325	47.0	42.0-52.5	300	30.3	26.9-34.0
2009	880	51.4	48.0-54.9	429	60.5	54.8-66.6	451	45.3	41.2-49.8	564	32.7	30.0-35.6	284	40.7	36.1-45.8	280	27.4	24.2-30.9
2010	862	49.5	46.2-53.0	407	55.9	50.5-61.7	455	45.3	41.1-49.8	590	33.5	30.9-36.4	297	41.4	36.8-46.5	293	28.1	24.9-31.6
2011	935	52.5	49.2-56.0	409	55.4	50.1-61.1	526	50.9	46.6-55.6	528	29.6	27.1-32.2	242	33.1	29.0-37.6	286	27.1	24.0-30.5
2012	896	49.0	45.8-52.4	379	50.0	45.0-55.4	517	48.9	44.7-53.4	535	29.3	26.8-31.9	253	33.8	29.7-38.3	282	26.2	23.2-29.6
2013	945	50.9	47.6-54.3	432	56.1	50.9-61.7	513	47.7	43.6-52.1	585	31.3	28.8-34.0	276	36.3	32.1-40.9	309	28.2	25.1-31.6
2014	912	48.1	45.0-51.4	434	54.1	49.1-59.5	478	44.3	40.4-48.6	519	27.0	24.7-29.4	254	32.3	28.4-36.6	265	23.3	20.5-26.3
2015	892	45.6	42.7-48.8	405	49.1	44.4-54.2	487	43.6	39.8-47.8	523	26.4	24.2-28.8	254	31.0	27.3-35.1	269	23.2	20.5-26.3
2016	907	45.8	42.9-49.0	383	45.7	41.2-50.6	524	46.0	42.1-50.2	471	23.6	21.5-25.9	224	26.9	23.5-30.8	247	21.3	18.7-24.3
2017	954	47.2	44.2-50.3	425	49.8	45.1-54.9	529	45.4	41.5-49.5	441	21.5	19.5-23.6	212	24.8	21.6-28.5	229	19.0	16.6-21.7
2018	881	42.4	39.6-45.3	394	44.5	40.2-49.2	487	40.9	37.2-44.8	434	20.7	18.8-22.8	195	22.4	19.4-25.8	239	19.4	17.0-22.1
2019	922	43.9	41.1-46.9	424	47.5	43.1-52.4	498	41.3	37.6-45.2	414	19.2	17.4-21.2	185	20.8	17.9-24.0	229	18.0	15.7-20.7
2020	747	35.3	32.8-38.0	348	39.2	35.2-43.6	399	32.7	29.5-36.2	354	16.1	14.5-18.0	174	19.2	16.4-22.3	180	13.9	11.9-16.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
Melanoma of the skin**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	99	6.3	5.1-7.8	44	7.0	5.0-9.5	55	6.3	4.7-8.3	31	2.0	1.3-2.9	13	2.1	1.1-3.7	18	1.9	1.1-3.1
1977	88	5.5	4.4-6.8	46	6.6	4.8-8.8	42	4.6	3.3-6.4	34	2.3	1.6-3.2	19	2.9	1.7-4.6	15	1.8	1.0-3.0
1978	99	6.7	5.4-8.2	48	7.3	5.3-9.9	51	6.5	4.7-8.6	22	1.5	0.9-2.2	16	2.7	1.5-4.5	6	0.7	0.2-1.6
1979	107	7.0	5.7-8.5	60	8.7	6.6-11.2	47	5.6	4.1-7.6	35	2.3	1.6-3.3	19	3.0	1.8-4.7	16	2.0	1.1-3.3
1980	122	8.1	6.7-9.7	69	10.7	8.2-13.6	53	6.3	4.6-8.3	29	1.9	1.3-2.7	19	3.0	1.8-4.8	10	1.1	0.5-2.1
1981	103	6.9	5.6-8.5	50	7.8	5.7-10.3	53	6.5	4.8-8.6	36	2.4	1.7-3.4	16	2.6	1.4-4.3	20	2.1	1.3-3.3
1982	113	7.6	6.2-9.2	60	9.6	7.3-12.4	53	6.2	4.6-8.2	35	2.3	1.6-3.3	23	3.7	2.3-5.7	12	1.4	0.7-2.6
1983	99	6.5	5.3-8.0	53	8.6	6.4-11.3	46	5.3	3.9-7.2	43	2.9	2.1-4.0	25	4.0	2.6-6.0	18	2.1	1.2-3.3
1984	120	7.6	6.3-9.2	56	8.4	6.3-11.0	64	7.3	5.5-9.4	42	2.8	2.0-3.8	28	4.6	3.0-6.6	14	1.6	0.9-2.8
1985	105	6.8	5.5-8.2	72	10.7	8.3-13.6	33	4.0	2.7-5.7	37	2.4	1.7-3.4	19	2.9	1.7-4.6	18	2.1	1.2-3.4
1986	108	6.7	5.4-8.1	53	8.2	6.1-10.9	55	5.8	4.4-7.7	33	2.2	1.5-3.1	18	3.1	1.8-4.9	15	1.7	0.9-2.8
1987	105	6.9	5.6-8.3	58	9.3	7.0-12.2	47	5.4	3.9-7.2	31	2.0	1.4-2.9	16	2.7	1.5-4.4	15	1.7	0.9-2.8
1988	94	6.2	5.0-7.6	59	8.7	6.6-11.4	35	4.0	2.8-5.6	37	2.5	1.8-3.5	22	3.7	2.3-5.6	15	1.8	1.0-3.0
1989	97	6.2	5.0-7.5	52	7.9	5.8-10.4	45	4.9	3.6-6.6	26	1.7	1.1-2.5	21	3.5	2.1-5.3	5	0.5	0.2-1.3
1990	102	6.7	5.4-8.1	55	9.0	6.7-11.8	47	5.4	3.9-7.2	26	1.7	1.1-2.5	14	2.2	1.2-3.8	12	1.3	0.7-2.3
1991	132	8.5	7.1-10.1	79	11.7	9.2-14.7	53	6.1	4.6-8.1	31	2.0	1.4-2.9	22	3.3	2.0-5.0	9	1.0	0.4-1.9
1992	99	6.5	5.3-7.9	57	8.6	6.5-11.2	42	4.7	3.4-6.5	24	1.5	1.0-2.3	16	2.3	1.3-3.9	8	0.8	0.3-1.7
1993	111	7.1	5.8-8.6	65	10.2	7.8-13.0	46	5.3	3.8-7.1	22	1.5	0.9-2.3	11	1.6	0.8-2.9	11	1.4	0.7-2.5
1994	126	8.3	6.9-9.9	77	11.8	9.3-14.9	49	5.4	4.0-7.2	27	1.8	1.2-2.7	12	2.2	1.1-3.8	15	1.7	0.9-2.8
1995	131	8.7	7.2-10.3	77	12.0	9.4-15.1	54	6.1	4.6-8.1	29	2.0	1.3-2.8	16	2.7	1.5-4.5	13	1.3	0.7-2.4
1996	160	10.3	8.8-12.1	92	15.2	12.1-18.7	68	7.5	5.8-9.6	24	1.6	1.0-2.4	10	1.6	0.8-3.1	14	1.6	0.9-2.7
1997	162	10.4	8.8-12.1	94	14.7	11.8-18.1	68	7.6	5.9-9.7	29	1.9	1.3-2.8	17	2.9	1.7-4.7	12	1.3	0.7-2.4
1998	209	13.2	11.4-15.1	115	17.5	14.4-21.1	94	10.3	8.3-12.7	41	2.8	2.0-3.8	28	4.7	3.1-6.8	13	1.5	0.8-2.6
1999	165	10.2	8.7-12.0	108	16.2	13.2-19.6	57	6.3	4.7-8.2	26	1.7	1.1-2.5	19	2.9	1.7-4.6	7	0.8	0.3-1.7
2000	197	12.3	10.6-14.2	110	17.0	13.9-20.6	87	9.5	7.6-11.8	36	2.4	1.7-3.3	19	3.4	2.1-5.4	17	2.0	1.1-3.2
2001	238	14.5	12.7-16.5	125	18.1	15.0-21.6	113	12.2	10.0-14.7	27	1.8	1.2-2.6	17	2.9	1.7-4.6	10	1.1	0.5-2.1
2002	259	15.7	13.8-17.8	142	21.0	17.6-24.9	117	12.5	10.3-15.0	27	1.7	1.1-2.5	12	1.8	0.9-3.2	15	1.7	0.9-2.8
2003	189	11.3	9.7-13.0	100	13.8	11.2-16.9	89	9.7	7.8-12.0	35	2.1	1.5-3.0	23	3.9	2.5-5.9	12	1.2	0.6-2.2
2004	329	19.3	17.2-21.5	176	24.7	21.1-28.8	153	15.9	13.4-18.7	27	1.6	1.1-2.4	11	1.5	0.8-2.8	16	1.7	0.9-2.8
2005	362	21.1	19.0-23.5	192	26.4	22.7-30.6	170	18.0	15.3-21.0	23	1.4	0.9-2.1	14	2.3	1.2-3.8	9	0.9	0.4-1.8
2006	458	26.6	24.2-29.2	255	33.8	29.7-38.4	203	21.3	18.5-24.6	31	1.9	1.3-2.7	18	2.9	1.7-4.6	13	1.4	0.7-2.5
2007	335	19.5	17.4-21.7	177	23.4	20.0-27.2	158	16.8	14.3-19.8	40	2.3	1.7-3.2	30	4.4	3.0-6.4	10	1.0	0.5-2.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
Melanoma of the skin**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	333	19.2	17.1-21.4	176	23.8	20.4-27.7	157	16.7	14.1-19.6	31	1.8	1.2-2.6	15	2.3	1.3-3.9	16	1.6	0.9-2.7
2009	356	20.0	17.9-22.2	205	27.0	23.4-31.1	151	15.1	12.7-17.8	24	1.4	0.9-2.0	18	2.5	1.5-4.0	6	0.6	0.2-1.4
2010	314	17.9	15.9-20.0	164	21.6	18.3-25.2	150	15.6	13.1-18.4	25	1.5	0.9-2.2	17	2.4	1.4-3.8	8	0.9	0.4-1.8
2011	402	22.1	19.9-24.4	220	28.6	24.9-32.7	182	17.6	15.1-20.5	20	1.1	0.6-1.7	13	1.7	0.9-2.9	7	0.6	0.2-1.3
2012	295	16.2	14.4-18.2	172	21.7	18.5-25.2	123	12.1	10.0-14.6	35	1.9	1.3-2.7	11	1.6	0.8-2.8	24	2.2	1.4-3.3
2013	344	18.7	16.7-20.8	192	24.3	21.0-28.1	152	14.5	12.3-17.2	29	1.5	1.0-2.1	17	2.1	1.2-3.4	12	1.0	0.5-1.9
2014	344	18.3	16.4-20.4	198	24.5	21.2-28.2	146	14.0	11.8-16.6	19	1.0	0.6-1.6	14	1.7	0.9-2.9	5	0.4	0.1-1.1
2015	334	17.3	15.4-19.3	201	24.4	21.1-28.0	133	12.1	10.1-14.5	23	1.2	0.7-1.8	16	1.9	1.1-3.2	7	0.7	0.3-1.5
2016	341	17.5	15.7-19.5	191	22.6	19.4-26.1	150	14.0	11.8-16.5	28	1.4	0.9-2.0	17	2.0	1.1-3.2	11	0.9	0.4-1.7
2017	334	17.0	15.2-19.0	202	23.7	20.5-27.3	132	12.1	10.0-14.5	26	1.3	0.8-1.9	16	1.9	1.1-3.2	10	0.8	0.4-1.6
2018	349	17.2	15.4-19.2	202	22.8	19.7-26.2	147	13.2	11.1-15.6	22	1.0	0.6-1.6	14	1.6	0.9-2.8	8	0.6	0.3-1.4
2019	409	20.9	18.9-23.1	229	26.3	23.0-30.1	180	17.1	14.6-20.0	20	0.9	0.6-1.5	15	1.7	0.9-2.9	5	0.4	0.1-1.0
2020	268	13.1	11.6-14.8	152	17.1	14.4-20.1	116	10.2	8.4-12.4	18	0.9	0.5-1.5	8	1.0	0.4-2.0	10	0.8	0.4-1.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Female breast

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	959	111.5	104.3-119.1	361	40.3	36.1-44.9
1977	925	108.7	101.6-116.2	333	38.7	34.5-43.2
1978	844	98.7	91.9-105.9	343	39.4	35.2-44.0
1979	916	110.1	102.9-117.8	329	38.0	33.9-42.5
1980	857	104.0	96.9-111.5	353	40.4	36.2-45.0
1981	914	109.0	101.8-116.6	343	39.7	35.5-44.3
1982	926	113.1	105.7-120.9	318	37.0	32.9-41.4
1983	907	109.3	102.1-116.9	331	38.6	34.5-43.1
1984	1,005	121.0	113.4-128.9	329	37.2	33.2-41.5
1985	979	117.4	110.0-125.3	320	36.1	32.2-40.4
1986	963	114.6	107.3-122.3	310	35.1	31.2-39.4
1987	961	113.7	106.4-121.3	314	35.5	31.6-39.7
1988	995	117.2	109.9-124.9	335	37.4	33.4-41.7
1989	1,022	121.3	113.8-129.1	348	39.4	35.3-43.8
1990	1,023	122.8	115.3-130.7	316	35.8	31.9-40.1
1991	1,208	145.5	137.3-154.0	315	36.6	32.6-40.9
1992	1,113	133.5	125.7-141.7	310	35.8	31.9-40.1
1993	1,032	125.2	117.6-133.2	284	32.8	29.1-36.9
1994	1,162	139.3	131.4-147.7	320	36.9	32.9-41.2
1995	1,083	129.8	122.2-137.9	315	35.4	31.6-39.6
1996	1,189	142.9	134.9-151.3	303	35.0	31.1-39.2
1997	1,190	141.1	133.1-149.4	287	32.5	28.8-36.5
1998	1,244	146.1	138.0-154.5	259	29.2	25.7-33.1
1999	1,181	137.1	129.3-145.2	257	28.7	25.3-32.5
2000	1,261	146.2	138.2-154.6	245	26.7	23.5-30.3
2001	1,162	132.1	124.6-140.0	264	28.5	25.1-32.2
2002	1,164	130.9	123.4-138.7	244	26.3	23.0-29.8
2003	1,147	128.5	121.1-136.3	235	24.4	21.4-27.9
2004	1,082	120.4	113.3-127.9	234	24.6	21.5-28.1
2005	1,187	130.4	122.9-138.1	245	26.0	22.8-29.5
2006	1,200	132.7	125.2-140.5	227	23.4	20.4-26.7
2007	1,280	140.6	132.9-148.7	209	21.7	18.8-24.9
2008	1,316	143.7	135.9-151.9	235	23.5	20.5-26.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Female breast

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	1,296	139.3	131.6-147.3	224	22.4	19.5-25.6
2010	1,249	132.2	124.8-139.9	198	19.4	16.7-22.4
2011	1,297	135.6	128.2-143.4	214	20.5	17.8-23.5
2012	1,249	129.4	122.2-137.0	210	19.9	17.3-22.9
2013	1,380	141.2	133.7-149.1	208	19.8	17.1-22.8
2014	1,442	146.2	138.6-154.2	206	18.9	16.4-21.8
2015	1,395	140.3	132.8-148.1	196	18.2	15.7-21.0
2016	1,382	138.6	131.2-146.4	207	18.7	16.2-21.6
2017	1,457	145.0	137.4-153.0	171	14.9	12.7-17.4
2018	1,424	139.2	131.8-146.9	186	16.1	13.8-18.7
2019	1,475	142.3	134.8-150.0	192	16.9	14.5-19.6
2020	1,260	125.5	118.4-132.9	173	14.9	12.7-17.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Cervix uteri

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	150	17.7	14.9-20.9	58	6.7	5.0-8.8
1977	151	18.5	15.6-21.9	62	7.2	5.4-9.3
1978	133	16.8	13.9-20.0	43	5.0	3.6-6.8
1979	138	17.3	14.4-20.6	47	5.8	4.2-7.8
1980	133	16.6	13.8-19.8	50	6.0	4.4-8.0
1981	113	15.1	12.3-18.2	41	4.8	3.4-6.7
1982	112	14.1	11.5-17.1	49	6.1	4.5-8.2
1983	119	14.6	12.0-17.6	40	4.8	3.4-6.6
1984	100	12.2	9.9-15.0	32	3.6	2.4-5.2
1985	112	12.9	10.6-15.6	29	3.5	2.3-5.1
1986	121	15.2	12.6-18.3	31	3.5	2.4-5.0
1987	111	13.0	10.7-15.8	32	3.8	2.5-5.4
1988	85	10.3	8.2-12.8	23	2.8	1.8-4.3
1989	87	10.0	7.9-12.4	25	2.9	1.9-4.3
1990	94	11.0	8.9-13.6	37	4.3	3.0-5.9
1991	116	14.1	11.7-17.0	27	3.2	2.1-4.8
1992	112	13.3	10.9-16.0	26	3.1	2.0-4.6
1993	122	14.7	12.1-17.6	27	3.3	2.2-4.9
1994	98	11.9	9.6-14.5	37	4.3	3.0-6.0
1995	113	13.4	11.0-16.1	36	4.2	2.9-5.8
1996	105	12.3	10.1-15.0	27	3.1	2.0-4.5
1997	116	13.6	11.2-16.4	31	3.7	2.5-5.3
1998	81	9.3	7.4-11.6	33	3.9	2.7-5.5
1999	101	11.9	9.7-14.5	35	4.0	2.8-5.6
2000	97	11.2	9.1-13.7	28	3.2	2.1-4.7
2001	89	10.2	8.2-12.6	34	4.0	2.8-5.6
2002	71	8.0	6.3-10.2	29	3.2	2.1-4.6
2003	81	9.0	7.2-11.3	27	3.1	2.1-4.6
2004	83	9.1	7.3-11.4	28	3.1	2.1-4.6
2005	89	9.8	7.8-12.1	25	2.6	1.7-3.9
2006	67	7.6	5.8-9.7	31	3.3	2.3-4.8
2007	61	7.0	5.3-9.0	31	3.4	2.3-4.9
2008	66	7.0	5.4-9.0	25	2.7	1.7-4.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Cervix uteri

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	65	7.5	5.7-9.6	22	2.1	1.3-3.3
2010	60	6.7	5.1-8.7	20	2.2	1.3-3.4
2011	74	7.9	6.1-10.0	25	2.6	1.7-4.0
2012	56	6.0	4.5-7.9	21	2.1	1.3-3.3
2013	83	8.5	6.7-10.6	27	2.8	1.8-4.1
2014	51	5.4	4.0-7.2	25	2.4	1.5-3.6
2015	78	7.7	6.0-9.7	11	1.1	0.5-2.0
2016	63	6.7	5.1-8.6	19	1.7	1.0-2.8
2017	74	7.9	6.2-10.0	24	2.3	1.4-3.5
2018	67	6.8	5.2-8.8	16	1.6	0.9-2.6
2019	60	6.3	4.8-8.3	16	1.6	0.9-2.6
2020	46	4.7	3.4-6.4	12	1.0	0.5-1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Corpus uteri and uterus, not otherwise specified

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	270	28.9	25.5-32.8	70	7.1	5.5-9.0
1977	224	24.7	21.5-28.3	73	7.5	5.9-9.5
1978	159	17.6	14.9-20.7	70	7.4	5.7-9.4
1979	184	20.4	17.5-23.8	49	5.7	4.2-7.6
1980	195	22.0	18.9-25.4	66	7.1	5.5-9.1
1981	173	19.5	16.6-22.7	48	5.1	3.7-6.8
1982	169	19.7	16.7-23.0	44	4.6	3.3-6.3
1983	169	19.0	16.2-22.3	49	5.3	3.9-7.1
1984	165	18.7	15.8-21.8	51	5.3	4.0-7.0
1985	176	19.5	16.7-22.7	42	4.4	3.2-6.0
1986	147	16.6	14.0-19.6	37	3.8	2.7-5.4
1987	161	18.3	15.5-21.4	34	3.5	2.4-4.9
1988	148	17.0	14.3-20.0	34	3.7	2.5-5.2
1989	130	14.8	12.3-17.6	40	4.3	3.0-5.9
1990	140	16.5	13.8-19.5	33	3.5	2.4-5.0
1991	180	21.1	18.1-24.5	49	5.7	4.2-7.5
1992	147	17.4	14.7-20.5	43	4.6	3.3-6.3
1993	177	20.8	17.8-24.2	58	6.4	4.9-8.3
1994	185	22.1	19.0-25.6	33	3.7	2.5-5.2
1995	212	25.4	22.0-29.1	38	4.1	2.9-5.7
1996	204	23.9	20.7-27.5	48	5.5	4.0-7.3
1997	217	25.5	22.2-29.1	57	6.5	4.9-8.4
1998	218	25.5	22.2-29.1	56	6.3	4.8-8.3
1999	220	25.3	22.0-28.9	44	5.1	3.7-6.8
2000	209	23.9	20.8-27.4	45	4.9	3.6-6.6
2001	230	26.0	22.8-29.7	48	5.2	3.8-7.0
2002	256	28.6	25.2-32.4	38	4.1	2.9-5.7
2003	240	26.5	23.2-30.1	46	5.1	3.7-6.9
2004	252	27.5	24.2-31.2	55	5.8	4.3-7.6
2005	243	26.1	22.9-29.7	44	4.6	3.4-6.3
2006	280	29.6	26.2-33.3	58	6.0	4.6-7.9
2007	218	22.8	19.8-26.1	58	5.9	4.5-7.7
2008	299	30.8	27.4-34.6	53	5.2	3.8-6.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Corpus uteris and uterus, not otherwise specified

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	266	27.7	24.4-31.3	52	5.1	3.8-6.7
2010	289	29.1	25.8-32.8	47	4.5	3.3-6.0
2011	291	29.2	25.9-32.8	60	5.7	4.3-7.4
2012	295	28.3	25.1-31.9	66	6.1	4.7-7.8
2013	300	29.2	25.9-32.8	54	4.9	3.7-6.5
2014	293	27.8	24.6-31.3	72	6.5	5.0-8.2
2015	314	30.0	26.7-33.7	76	6.6	5.1-8.3
2016	340	32.0	28.6-35.7	54	4.8	3.6-6.4
2017	327	30.9	27.6-34.6	63	5.4	4.1-7.0
2018	324	29.4	26.2-33.0	62	5.3	4.0-6.8
2019	292	27.5	24.4-31.0	58	4.8	3.6-6.3
2020	295	26.8	23.8-30.2	59	4.9	3.7-6.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Ovary

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	150	17.0	14.3-20.1	81	9.0	7.0-11.3
1977	133	15.7	13.0-18.7	93	10.6	8.5-13.0
1978	140	16.4	13.7-19.6	86	9.6	7.6-11.9
1979	131	15.6	12.9-18.6	84	9.6	7.6-12.0
1980	141	16.4	13.7-19.5	80	8.9	7.0-11.2
1981	148	17.0	14.3-20.2	74	8.4	6.6-10.7
1982	146	17.1	14.3-20.2	92	10.1	8.1-12.5
1983	146	17.2	14.5-20.4	93	10.4	8.4-12.9
1984	149	17.1	14.4-20.2	80	8.6	6.8-10.8
1985	163	19.4	16.4-22.7	88	10.0	8.0-12.4
1986	128	14.7	12.2-17.5	96	11.3	9.1-13.8
1987	133	16.0	13.3-19.0	83	9.2	7.3-11.5
1988	122	14.2	11.8-17.1	90	10.1	8.1-12.5
1989	129	15.0	12.5-18.0	72	7.9	6.1-10.0
1990	127	14.7	12.2-17.6	80	8.8	7.0-11.1
1991	148	17.7	14.9-20.9	98	11.2	9.1-13.7
1992	143	17.0	14.3-20.1	90	10.5	8.4-12.9
1993	143	16.8	14.1-19.8	79	9.0	7.1-11.2
1994	157	18.1	15.4-21.3	89	10.2	8.2-12.6
1995	164	19.5	16.6-22.7	91	10.5	8.4-13.0
1996	152	17.8	15.1-20.9	74	8.6	6.7-10.8
1997	148	17.5	14.7-20.5	72	8.3	6.5-10.5
1998	140	16.7	14.1-19.8	86	9.8	7.9-12.2
1999	130	15.0	12.6-17.9	87	10.0	8.0-12.3
2000	116	13.3	11.0-16.0	70	7.9	6.1-10.0
2001	148	17.0	14.4-20.0	76	8.7	6.8-10.9
2002	147	16.4	13.8-19.3	78	8.4	6.6-10.5
2003	124	13.8	11.5-16.5	76	8.3	6.5-10.4
2004	150	16.7	14.1-19.6	76	8.1	6.4-10.2
2005	120	13.3	11.0-15.9	67	7.0	5.4-9.0
2006	119	12.8	10.6-15.4	70	7.3	5.6-9.3
2007	127	13.6	11.3-16.3	75	7.7	6.0-9.7
2008	117	12.3	10.1-14.8	75	7.6	5.9-9.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Ovary

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	133	14.2	11.9-17.0	72	7.3	5.7-9.3
2010	135	14.4	12.1-17.2	81	7.9	6.3-9.9
2011	143	14.5	12.2-17.2	74	7.1	5.6-9.0
2012	121	11.7	9.6-14.0	75	7.2	5.6-9.1
2013	126	12.6	10.4-15.1	79	7.4	5.8-9.3
2014	121	12.5	10.3-15.0	72	6.8	5.3-8.6
2015	136	14.1	11.7-16.8	76	7.3	5.7-9.2
2016	99	9.8	7.9-12.0	69	5.8	4.5-7.5
2017	117	10.8	8.9-13.1	80	7.1	5.6-8.9
2018	108	9.9	8.1-12.0	65	5.6	4.3-7.3
2019	106	10.1	8.2-12.4	81	6.9	5.4-8.7
2020	98	9.5	7.6-11.8	69	5.8	4.5-7.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Prostate

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	507	86.0	78.3-94.3	168	31.6	26.7-37.1
1977	529	88.2	80.6-96.4	178	32.2	27.4-37.5
1978	493	85.4	77.7-93.6	182	34.1	29.1-39.7
1979	505	87.0	79.3-95.2	152	28.1	23.6-33.1
1980	567	100.0	91.7-108.9	173	32.7	27.9-38.1
1981	581	100.1	91.9-108.8	199	39.0	33.6-45.0
1982	507	92.1	84.0-100.7	180	34.8	29.8-40.4
1983	533	92.4	84.5-100.8	201	38.8	33.5-44.7
1984	543	95.9	87.8-104.6	198	37.9	32.7-43.6
1985	552	95.7	87.6-104.2	207	39.7	34.3-45.6
1986	594	105.2	96.7-114.2	201	39.4	34.0-45.3
1987	574	99.0	90.9-107.6	194	37.0	31.9-42.6
1988	561	100.3	92.0-109.1	192	37.0	31.8-42.7
1989	557	96.4	88.4-104.9	186	35.3	30.3-40.8
1990	605	106.7	98.2-115.8	231	45.1	39.4-51.4
1991	765	133.2	123.7-143.2	184	35.5	30.5-41.1
1992	965	165.7	155.2-176.7	238	45.9	40.2-52.2
1993	882	148.8	139.0-159.2	215	42.9	37.2-49.0
1994	997	166.7	156.3-177.5	199	39.4	34.0-45.3
1995	870	144.0	134.4-154.0	221	42.9	37.4-49.0
1996	919	151.1	141.4-161.4	206	40.8	35.4-46.9
1997	1,026	166.8	156.6-177.4	173	33.4	28.5-38.8
1998	1,071	173.4	163.0-184.2	190	36.0	31.0-41.5
1999	1,068	169.7	159.5-180.3	175	33.3	28.5-38.6
2000	1,216	191.1	180.3-202.3	171	32.6	27.9-37.9
2001	1,064	163.0	153.3-173.3	186	34.6	29.8-40.0
2002	1,195	179.9	169.7-190.6	209	37.5	32.5-43.0
2003	1,002	148.1	139.0-157.7	161	28.4	24.2-33.2
2004	1,050	152.7	143.5-162.4	158	27.8	23.5-32.5
2005	1,041	146.9	138.1-156.3	157	26.3	22.3-30.8
2006	1,228	173.6	163.9-183.8	174	29.8	25.5-34.6
2007	1,282	177.7	168.0-187.8	169	27.4	23.4-31.8
2008	1,201	163.3	154.1-172.9	140	22.8	19.2-27.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Prostate

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	1,168	156.7	147.7-166.1	190	29.8	25.7-34.4
2010	1,115	145.2	136.7-154.1	160	24.3	20.7-28.4
2011	1,177	149.7	141.1-158.6	147	21.6	18.2-25.4
2012	1,064	131.2	123.3-139.4	139	19.6	16.4-23.1
2013	1,087	131.6	123.8-139.8	136	18.7	15.7-22.2
2014	992	116.8	109.5-124.4	155	20.2	17.2-23.7
2015	1,022	117.9	110.7-125.5	142	18.0	15.2-21.3
2016	1,125	127.5	120.0-135.2	157	19.6	16.6-22.9
2017	1,107	123.2	116.0-130.8	131	15.6	13.0-18.6
2018	1,170	127.9	120.7-135.6	150	17.3	14.6-20.4
2019	1,287	140.6	132.9-148.6	134	14.9	12.5-17.7
2020	1,039	112.1	105.3-119.2	115	12.9	10.6-15.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Testis

Year of diagnosis	Incidence		
	Cases	Rate per 100,000	95% CI
1976	21	2.8	1.7-4.4
1977	25	3.2	2.0-4.8
1978	17	2.3	1.3-3.7
1979	23	3.2	2.0-4.9
1980	26	3.1	2.0-4.7
1981	35	3.9	2.7-5.6
1982	37	4.4	3.1-6.3
1983	30	3.8	2.5-5.5
1984	29	3.4	2.2-5.0
1985	30	3.7	2.5-5.4
1986	26	2.9	1.9-4.4
1987	38	4.6	3.2-6.4
1988	32	3.7	2.5-5.4
1989	41	4.6	3.3-6.4
1990	42	5.2	3.7-7.2
1991	39	4.4	3.1-6.2
1992	46	5.5	4.0-7.5
1993	33	3.5	2.3-5.0
1994	41	4.9	3.5-6.8
1995	28	3.2	2.1-4.8
1996	35	3.9	2.7-5.6
1997	26	2.7	1.8-4.1
1998	45	4.9	3.5-6.7
1999	44	4.8	3.4-6.5
2000	59	6.4	4.9-8.4
2001	46	4.8	3.5-6.5
2002	49	5.4	4.0-7.3
2003	45	5.0	3.6-6.9
2004	60	6.5	4.9-8.5
2005	57	5.9	4.5-7.8
2006	47	5.1	3.7-6.8
2007	71	8.0	6.2-10.1
2008	40	4.4	3.1-6.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

Mortality data not provided due to small number of deaths per year.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Testis

Year of diagnosis	Incidence		
	Cases	Rate per 100,000	95% CI
2009	63	7.0	5.3-9.1
2010	61	6.9	5.2-8.9
2011	62	6.7	5.1-8.7
2012	61	6.9	5.2-9.0
2013	63	6.6	5.0-8.6
2014	52	5.9	4.3-7.8
2015	57	5.8	4.3-7.6
2016	74	8.1	6.3-10.3
2017	63	6.7	5.1-8.7
2018	74	8.5	6.6-10.8
2019	62	6.3	4.7-8.2
2020	54	6.2	4.6-8.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

Mortality data not provided due to small number of deaths per year.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
Urinary bladder (incl. in situ)**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	272	17.5	15.4-19.8	192	32.2	27.5-37.3	80	8.4	6.6-10.6	81	5.5	4.3-6.9	58	10.3	7.7-13.4	23	2.7	1.7-4.2
1977	280	18.3	16.1-20.6	200	34.8	29.9-40.3	80	8.4	6.6-10.5	98	7.0	5.7-8.6	67	13.2	10.0-16.9	31	3.5	2.4-5.1
1978	302	19.7	17.5-22.1	198	33.0	28.4-38.0	104	11.3	9.2-13.8	100	7.1	5.7-8.7	60	11.0	8.3-14.3	40	4.8	3.4-6.6
1979	320	20.7	18.4-23.1	210	35.8	31.0-41.2	110	11.5	9.4-14.0	86	5.7	4.6-7.1	51	9.2	6.8-12.2	35	3.7	2.5-5.2
1980	296	19.5	17.3-21.9	197	33.9	29.2-39.1	99	10.8	8.7-13.2	96	6.5	5.2-8.0	61	11.0	8.3-14.3	35	3.9	2.7-5.5
1981	289	19.1	17.0-21.5	198	34.0	29.3-39.2	91	9.8	7.9-12.2	79	5.5	4.3-6.8	41	7.8	5.5-10.7	38	4.2	3.0-5.8
1982	292	19.3	17.1-21.6	184	32.1	27.5-37.3	108	11.5	9.4-13.9	91	6.1	4.9-7.6	50	9.4	6.9-12.5	41	4.3	3.1-5.9
1983	289	18.9	16.8-21.3	190	33.0	28.3-38.2	99	10.5	8.5-12.9	73	4.9	3.8-6.1	46	8.3	6.1-11.2	27	2.9	1.9-4.2
1984	270	17.4	15.4-19.6	191	32.4	27.8-37.4	79	8.3	6.5-10.4	68	4.6	3.6-5.9	42	7.8	5.6-10.6	26	2.8	1.8-4.2
1985	258	17.3	15.2-19.5	169	29.7	25.2-34.6	89	9.4	7.6-11.7	85	5.7	4.5-7.0	59	11.2	8.5-14.5	26	2.7	1.7-4.0
1986	283	18.7	16.6-21.1	191	33.6	28.9-38.9	92	9.9	8.0-12.3	75	5.1	4.0-6.4	48	9.4	6.9-12.5	27	2.8	1.8-4.1
1987	241	15.7	13.8-17.9	168	28.7	24.4-33.5	73	8.0	6.2-10.1	46	3.1	2.2-4.1	26	5.0	3.2-7.3	20	2.1	1.3-3.3
1988	237	15.6	13.7-17.8	162	28.0	23.8-32.8	75	7.9	6.2-9.9	76	5.0	3.9-6.3	50	9.2	6.8-12.2	26	2.7	1.8-4.1
1989	253	16.9	14.8-19.1	159	28.1	23.8-32.9	94	10.2	8.2-12.5	60	4.0	3.1-5.2	34	6.5	4.4-9.0	26	2.7	1.7-4.0
1990	224	14.9	13.0-17.0	151	26.8	22.6-31.6	73	7.9	6.2-10.0	52	3.5	2.6-4.6	28	5.5	3.6-8.0	24	2.5	1.6-3.8
1991	303	20.6	18.3-23.0	188	33.6	28.9-38.8	115	12.6	10.4-15.2	68	4.6	3.5-5.8	42	7.7	5.5-10.4	26	2.7	1.8-4.0
1992	277	18.8	16.6-21.1	180	32.4	27.8-37.6	97	10.5	8.5-12.9	52	3.5	2.6-4.6	30	5.8	3.9-8.3	22	2.3	1.4-3.5
1993	281	19.1	16.9-21.5	184	32.4	27.8-37.5	97	10.9	8.8-13.3	68	4.6	3.6-5.8	34	6.3	4.3-8.8	34	3.6	2.5-5.1
1994	295	19.7	17.5-22.1	194	34.2	29.5-39.5	101	11.2	9.1-13.7	69	4.7	3.6-5.9	37	7.1	4.9-9.8	32	3.3	2.3-4.8
1995	267	18.0	15.9-20.3	194	33.2	28.6-38.3	73	8.1	6.4-10.3	72	4.9	3.8-6.2	39	7.4	5.2-10.1	33	3.4	2.3-4.8
1996	233	15.7	13.8-17.9	151	26.3	22.2-31.0	82	9.0	7.2-11.3	53	3.6	2.7-4.7	38	7.5	5.3-10.3	15	1.5	0.9-2.6
1997	308	20.9	18.6-23.3	213	36.4	31.6-41.7	95	10.7	8.6-13.1	56	3.8	2.8-4.9	36	7.0	4.8-9.6	20	2.1	1.3-3.3
1998	285	19.4	17.2-21.8	195	32.5	28.1-37.5	90	10.2	8.2-12.6	83	5.6	4.5-7.0	52	9.7	7.2-12.7	31	3.3	2.2-4.7
1999	263	17.5	15.5-19.8	166	27.8	23.7-32.5	97	10.9	8.9-13.4	68	4.5	3.5-5.7	49	9.0	6.6-11.9	19	1.9	1.2-3.1
2000	300	19.7	17.5-22.0	205	34.1	29.5-39.2	95	10.4	8.4-12.8	73	4.8	3.8-6.1	52	9.5	7.1-12.5	21	2.2	1.3-3.4
2001	292	18.8	16.7-21.1	208	33.7	29.2-38.7	84	9.0	7.2-11.2	64	4.1	3.2-5.2	36	6.4	4.5-8.9	28	2.9	1.9-4.2
2002	262	16.9	14.9-19.0	188	30.4	26.1-35.1	74	7.9	6.2-9.9	74	4.7	3.7-5.9	40	6.8	4.8-9.3	34	3.3	2.3-4.7
2003	306	19.4	17.3-21.8	216	33.9	29.5-38.8	90	9.5	7.6-11.7	56	3.5	2.7-4.6	36	6.2	4.3-8.6	20	1.9	1.2-3.0
2004	264	16.4	14.5-18.5	184	28.4	24.4-32.9	80	8.3	6.5-10.3	63	3.9	3.0-5.0	40	6.4	4.5-8.7	23	2.1	1.3-3.3
2005	298	18.2	16.2-20.5	199	30.0	25.9-34.6	99	10.2	8.2-12.4	66	4.1	3.1-5.2	41	6.6	4.7-8.9	25	2.5	1.6-3.7
2006	303	18.4	16.3-20.6	224	34.6	30.1-39.5	79	7.9	6.2-9.9	70	4.2	3.2-5.3	46	7.5	5.5-10.0	24	2.2	1.4-3.3
2007	316	18.9	16.8-21.1	219	32.5	28.3-37.2	97	9.9	8.0-12.1	86	5.2	4.2-6.5	56	8.8	6.7-11.5	30	2.8	1.9-4.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
Urinary bladder (incl. in situ)**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	314	18.5	16.5-20.6	240	34.6	30.3-39.3	74	7.5	5.9-9.5	67	4.0	3.1-5.1	48	7.4	5.4-9.8	19	1.8	1.1-2.9
2009	296	17.4	15.4-19.5	215	30.9	26.9-35.4	81	8.0	6.3-10.0	61	3.5	2.7-4.5	48	7.4	5.4-9.8	13	1.2	0.6-2.1
2010	318	18.4	16.4-20.6	238	34.0	29.8-38.7	80	7.7	6.1-9.7	78	4.4	3.5-5.5	57	8.2	6.2-10.7	21	1.8	1.1-2.9
2011	296	16.5	14.7-18.6	215	29.8	25.9-34.1	81	7.5	6.0-9.5	64	3.5	2.7-4.5	44	6.3	4.6-8.5	20	1.7	1.0-2.7
2012	281	15.5	13.7-17.4	201	27.3	23.6-31.4	80	7.2	5.7-9.0	67	3.7	2.9-4.8	48	6.8	5.0-9.0	19	1.8	1.1-2.9
2013	353	19.1	17.1-21.2	260	34.3	30.2-38.8	93	8.7	7.0-10.7	58	3.0	2.3-3.9	42	5.8	4.1-7.8	16	1.3	0.7-2.2
2014	310	16.3	14.5-18.3	233	29.6	25.8-33.7	77	6.9	5.4-8.7	65	3.3	2.5-4.2	41	5.5	3.9-7.4	24	2.0	1.2-3.0
2015	358	18.5	16.6-20.5	270	33.9	29.9-38.2	88	7.8	6.2-9.7	71	3.5	2.7-4.5	49	6.3	4.7-8.4	22	1.7	1.1-2.7
2016	350	17.2	15.5-19.2	261	31.4	27.7-35.5	89	7.3	5.9-9.1	65	3.1	2.4-4.0	41	5.1	3.6-6.9	24	1.8	1.1-2.8
2017	372	18.4	16.5-20.4	263	31.1	27.4-35.2	109	9.3	7.6-11.3	71	3.3	2.6-4.2	50	6.0	4.5-8.0	21	1.6	1.0-2.6
2018	356	17.0	15.3-18.9	252	28.9	25.4-32.8	104	8.8	7.2-10.8	67	3.1	2.4-4.0	45	5.2	3.8-7.0	22	1.8	1.1-2.8
2019	359	17.2	15.5-19.2	260	29.6	26.1-33.5	99	8.4	6.8-10.3	69	3.0	2.3-3.8	45	4.9	3.6-6.7	24	1.7	1.1-2.6
2020	288	13.7	12.2-15.4	189	21.5	18.5-24.8	99	8.2	6.6-10.0	51	2.3	1.7-3.0	28	3.2	2.1-4.7	23	1.6	1.0-2.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Kidney and renal pelvis

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	113	7.1	5.8-8.6	69	10.1	7.9-12.9	44	4.7	3.4-6.4	50	3.1	2.2-4.1	36	5.4	3.7-7.5	14	1.4	0.8-2.5
1977	102	6.6	5.4-8.1	73	12.1	9.4-15.4	29	2.9	2.0-4.3	50	3.3	2.5-4.4	33	5.6	3.7-7.9	17	1.9	1.1-3.1
1978	113	7.5	6.2-9.1	68	10.9	8.4-13.9	45	4.9	3.5-6.6	62	4.1	3.1-5.3	41	6.5	4.6-8.8	21	2.5	1.5-3.8
1979	105	6.6	5.4-8.1	62	9.4	7.1-12.0	43	4.7	3.4-6.4	43	2.9	2.0-3.9	31	4.9	3.3-7.0	12	1.4	0.7-2.5
1980	106	7.0	5.7-8.5	67	10.7	8.2-13.6	39	4.7	3.3-6.5	44	2.9	2.1-3.9	28	4.9	3.2-7.1	16	1.7	0.9-2.8
1981	111	7.4	6.1-8.9	68	11.4	8.7-14.5	43	5.0	3.6-6.9	52	3.4	2.6-4.6	32	5.3	3.6-7.6	20	2.2	1.3-3.4
1982	107	7.3	6.0-8.9	71	11.4	8.8-14.4	36	4.1	2.8-5.8	44	2.9	2.1-3.9	25	4.2	2.7-6.3	19	2.0	1.2-3.1
1983	110	7.3	6.0-8.8	70	11.2	8.7-14.3	40	4.5	3.2-6.2	39	2.6	1.8-3.6	23	3.9	2.4-5.8	16	1.8	1.0-3.0
1984	128	8.7	7.3-10.4	82	14.1	11.1-17.6	46	5.3	3.8-7.1	51	3.3	2.5-4.4	35	5.7	3.9-8.0	16	1.8	1.0-3.0
1985	112	7.3	6.0-8.8	68	10.8	8.3-13.7	44	4.7	3.4-6.4	44	2.8	2.1-3.8	26	4.5	2.9-6.6	18	1.9	1.1-3.1
1986	141	9.4	7.9-11.1	86	13.7	10.9-17.0	55	6.0	4.5-7.9	54	3.6	2.7-4.7	30	5.1	3.4-7.3	24	2.7	1.7-4.1
1987	125	8.3	6.9-9.9	84	13.8	10.9-17.1	41	4.6	3.3-6.3	51	3.4	2.5-4.5	34	6.0	4.1-8.4	17	1.8	1.0-2.9
1988	128	8.5	7.1-10.2	79	13.3	10.5-16.7	49	5.8	4.3-7.8	53	3.5	2.6-4.6	36	6.4	4.4-8.8	17	1.8	1.1-3.0
1989	130	8.6	7.1-10.2	77	12.8	10.0-16.0	53	5.9	4.4-7.7	45	3.0	2.2-4.0	21	3.9	2.4-5.9	24	2.5	1.6-3.8
1990	110	7.3	6.0-8.9	67	10.7	8.3-13.7	43	4.9	3.5-6.7	51	3.4	2.5-4.5	24	4.1	2.6-6.1	27	3.0	1.9-4.4
1991	152	10.3	8.7-12.0	87	14.1	11.3-17.5	65	7.4	5.7-9.5	44	3.0	2.2-4.0	27	4.5	3.0-6.6	17	1.9	1.1-3.2
1992	145	9.9	8.3-11.6	87	14.6	11.6-18.0	58	6.9	5.2-9.0	36	2.5	1.7-3.4	20	3.7	2.3-5.8	16	1.7	1.0-2.9
1993	134	9.0	7.6-10.7	87	14.2	11.3-17.6	47	5.4	3.9-7.2	42	2.9	2.1-3.9	24	4.1	2.6-6.1	18	2.1	1.2-3.3
1994	155	10.6	9.0-12.4	86	14.1	11.2-17.5	69	8.0	6.2-10.2	44	3.0	2.2-4.0	30	5.1	3.4-7.3	14	1.6	0.9-2.7
1995	138	9.3	7.8-11.1	82	13.3	10.5-16.6	56	6.4	4.8-8.4	49	3.4	2.5-4.4	29	4.9	3.3-7.1	20	2.2	1.3-3.5
1996	153	10.4	8.8-12.2	98	15.6	12.6-19.0	55	6.5	4.9-8.5	42	2.8	2.0-3.8	21	3.5	2.1-5.3	21	2.2	1.4-3.5
1997	163	11.0	9.3-12.8	90	14.9	11.9-18.4	73	8.5	6.7-10.8	53	3.6	2.7-4.7	28	4.7	3.1-6.8	25	2.7	1.8-4.1
1998	159	10.8	9.2-12.6	99	16.0	13.0-19.6	60	7.0	5.4-9.1	39	2.6	1.9-3.6	23	3.9	2.4-5.9	16	1.8	1.0-3.0
1999	188	12.6	10.8-14.5	122	19.0	15.7-22.7	66	7.9	6.1-10.0	47	3.2	2.3-4.3	30	5.1	3.4-7.3	17	1.9	1.1-3.1
2000	184	12.0	10.3-13.9	98	15.3	12.4-18.7	86	9.6	7.6-11.9	48	3.1	2.3-4.1	28	4.5	3.0-6.6	20	2.2	1.3-3.4
2001	192	12.3	10.6-14.2	129	19.4	16.2-23.2	63	6.9	5.3-8.8	37	2.3	1.6-3.2	23	3.6	2.3-5.5	14	1.4	0.8-2.4
2002	178	11.4	9.7-13.2	104	15.0	12.2-18.2	74	8.2	6.4-10.4	42	2.7	1.9-3.7	28	4.3	2.9-6.3	14	1.5	0.8-2.6
2003	215	13.4	11.7-15.3	142	20.8	17.4-24.5	73	7.9	6.2-10.0	37	2.3	1.6-3.2	25	3.8	2.4-5.6	12	1.2	0.6-2.2
2004	204	12.6	10.9-14.4	133	18.9	15.7-22.4	71	7.7	6.0-9.7	38	2.4	1.7-3.3	21	3.1	1.9-4.8	17	1.8	1.0-2.9
2005	205	12.6	11.0-14.5	123	17.4	14.4-20.8	82	8.8	7.0-11.0	38	2.3	1.6-3.2	22	3.2	2.0-4.9	16	1.5	0.9-2.6
2006	243	14.9	13.1-16.9	169	23.9	20.4-27.9	74	7.9	6.2-10.0	42	2.5	1.8-3.4	24	3.5	2.2-5.3	18	1.7	1.0-2.8
2007	213	13.0	11.3-14.9	136	19.3	16.2-22.9	77	8.2	6.5-10.4	50	3.0	2.2-4.0	33	4.8	3.3-6.7	17	1.6	0.9-2.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Kidney and renal pelvis

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	217	13.0	11.3-14.9	131	18.3	15.3-21.8	86	9.2	7.3-11.4	38	2.2	1.6-3.1	22	3.3	2.1-5.1	16	1.6	0.9-2.6
2009	220	13.2	11.5-15.1	144	20.0	16.8-23.6	76	7.9	6.1-9.9	45	2.7	2.0-3.7	30	4.7	3.1-6.6	15	1.5	0.8-2.5
2010	198	11.7	10.1-13.5	130	17.3	14.4-20.6	68	7.2	5.6-9.2	29	1.7	1.1-2.4	18	2.5	1.5-4.0	11	1.1	0.5-2.0
2011	218	12.5	10.9-14.3	141	18.2	15.3-21.6	77	7.9	6.2-9.9	34	1.8	1.3-2.6	24	3.1	2.0-4.7	10	0.9	0.4-1.8
2012	218	12.5	10.9-14.4	151	19.4	16.4-22.8	67	7.1	5.5-9.1	30	1.7	1.1-2.4	21	2.7	1.7-4.2	9	0.9	0.4-1.8
2013	259	14.0	12.3-15.8	162	20.0	17.0-23.4	97	9.3	7.5-11.4	33	1.8	1.2-2.5	20	2.6	1.6-4.0	13	1.1	0.6-2.0
2014	226	12.3	10.7-14.0	151	18.2	15.4-21.4	75	7.3	5.7-9.3	35	1.8	1.2-2.5	27	3.2	2.1-4.8	8	0.7	0.3-1.4
2015	241	12.9	11.3-14.6	151	18.0	15.2-21.2	90	8.6	6.9-10.7	53	2.7	2.1-3.6	37	4.6	3.2-6.4	16	1.4	0.8-2.3
2016	214	11.2	9.7-12.8	130	15.1	12.6-18.0	84	8.1	6.4-10.1	36	1.7	1.2-2.5	20	2.4	1.4-3.7	16	1.3	0.7-2.2
2017	243	12.9	11.3-14.6	164	19.2	16.3-22.4	79	7.7	6.0-9.8	38	1.9	1.3-2.7	24	2.8	1.8-4.2	14	1.2	0.6-2.2
2018	255	13.0	11.4-14.8	169	19.3	16.5-22.6	86	7.9	6.2-9.9	40	1.9	1.4-2.7	25	2.9	1.9-4.3	15	1.1	0.6-2.0
2019	252	13.0	11.4-14.7	163	18.7	15.9-21.9	89	8.7	6.9-10.8	31	1.4	0.9-2.0	18	1.9	1.1-3.1	13	0.9	0.5-1.7
2020	226	11.4	10.0-13.1	143	16.3	13.7-19.2	83	7.6	6.0-9.5	36	1.8	1.3-2.6	23	2.8	1.8-4.2	13	1.1	0.6-2.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
Brain and other central nervous system**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	104	6.6	5.4-8.0	63	9.1	7.0-11.8	41	4.7	3.3-6.5	82	5.3	4.2-6.7	44	6.5	4.7-8.8	38	4.6	3.2-6.4
1977	85	5.8	4.6-7.2	49	7.5	5.5-10.0	36	4.4	3.0-6.2	79	5.2	4.1-6.6	43	6.4	4.6-8.8	36	4.4	3.0-6.2
1978	85	5.8	4.6-7.2	45	6.9	5.0-9.2	40	5.0	3.5-6.9	64	4.0	3.1-5.1	35	5.2	3.6-7.3	29	2.9	1.9-4.2
1979	82	5.3	4.2-6.6	43	6.7	4.8-9.2	39	4.2	2.9-5.9	60	3.8	2.9-4.9	35	5.6	3.9-7.8	25	2.6	1.6-3.9
1980	81	5.2	4.1-6.5	37	5.8	4.0-8.0	44	4.8	3.4-6.5	61	4.0	3.0-5.1	28	4.3	2.8-6.3	33	3.8	2.5-5.4
1981	77	5.4	4.3-6.8	49	7.8	5.8-10.4	28	3.5	2.3-5.1	60	4.1	3.1-5.3	33	5.2	3.6-7.4	27	3.1	2.0-4.6
1982	87	5.9	4.7-7.4	48	7.3	5.3-9.8	39	4.7	3.3-6.6	76	4.9	3.9-6.2	45	6.7	4.8-9.0	31	3.3	2.3-4.8
1983	81	5.5	4.4-7.0	31	4.9	3.3-7.0	50	6.3	4.6-8.4	58	3.9	3.0-5.1	31	5.2	3.5-7.4	27	3.0	2.0-4.5
1984	83	5.7	4.5-7.1	38	5.7	4.0-7.9	45	5.9	4.2-8.0	64	4.1	3.1-5.3	30	4.7	3.1-6.7	34	3.7	2.5-5.3
1985	118	8.2	6.7-9.8	60	9.4	7.1-12.2	58	7.5	5.6-9.9	66	4.3	3.3-5.6	37	6.1	4.2-8.4	29	3.4	2.2-5.0
1986	75	5.0	3.9-6.3	53	8.1	6.0-10.7	22	2.6	1.6-4.1	67	4.4	3.4-5.7	38	6.0	4.2-8.3	29	3.4	2.2-4.9
1987	88	6.0	4.8-7.4	39	6.0	4.2-8.3	49	5.9	4.3-7.9	60	4.0	3.1-5.2	35	5.8	4.0-8.1	25	2.8	1.8-4.2
1988	116	7.7	6.3-9.2	67	10.0	7.7-12.7	49	6.0	4.4-8.0	39	2.5	1.8-3.5	22	3.4	2.1-5.2	17	2.0	1.2-3.3
1989	109	7.5	6.1-9.1	53	8.2	6.1-10.8	56	6.8	5.1-8.9	45	3.0	2.2-4.0	28	4.3	2.8-6.2	17	1.9	1.1-3.1
1990	108	7.3	5.9-8.8	50	7.7	5.6-10.2	58	6.9	5.2-9.1	52	3.6	2.7-4.8	26	4.4	2.8-6.4	26	3.1	2.0-4.6
1991	111	7.6	6.2-9.1	61	9.7	7.3-12.5	50	6.2	4.5-8.2	65	4.4	3.4-5.6	26	4.1	2.6-6.0	39	4.7	3.4-6.5
1992	114	7.8	6.4-9.4	59	8.8	6.6-11.4	55	6.9	5.1-9.0	47	3.2	2.4-4.3	23	3.8	2.4-5.8	24	2.6	1.7-4.0
1993	107	7.2	5.8-8.7	55	8.4	6.2-11.0	52	6.4	4.7-8.5	45	3.1	2.2-4.2	25	4.3	2.7-6.3	20	2.5	1.5-3.9
1994	121	8.1	6.7-9.8	62	9.1	6.9-11.8	59	7.4	5.6-9.6	46	3.1	2.3-4.2	28	4.5	2.9-6.5	18	2.1	1.2-3.3
1995	109	7.3	6.0-8.8	67	10.0	7.6-12.8	42	5.1	3.7-7.0	50	3.2	2.4-4.3	33	5.1	3.5-7.3	17	1.9	1.1-3.1
1996	141	9.4	7.9-11.2	71	10.3	8.0-13.1	70	8.4	6.5-10.7	44	2.9	2.1-3.9	21	3.2	2.0-5.0	23	2.7	1.7-4.1
1997	103	7.0	5.7-8.6	47	7.0	5.1-9.4	56	6.9	5.2-9.1	66	4.4	3.4-5.6	31	5.0	3.4-7.1	35	4.0	2.8-5.6
1998	149	9.8	8.2-11.5	88	12.2	9.7-15.2	61	7.8	5.9-10.1	58	3.9	3.0-5.1	26	3.9	2.5-5.7	32	3.7	2.5-5.3
1999	113	7.5	6.2-9.1	61	8.8	6.7-11.4	52	6.4	4.7-8.5	55	3.7	2.8-4.8	33	5.3	3.6-7.5	22	2.6	1.6-3.9
2000	116	7.7	6.4-9.3	58	8.5	6.4-11.1	58	7.1	5.4-9.3	42	2.8	2.0-3.8	27	4.1	2.7-6.0	15	1.9	1.0-3.1
2001	109	7.1	5.8-8.6	50	7.0	5.1-9.3	59	7.4	5.5-9.6	49	3.2	2.4-4.3	21	3.4	2.1-5.1	28	3.3	2.2-4.8
2002	101	6.5	5.3-7.9	58	9.0	6.8-11.7	43	4.9	3.5-6.7	54	3.5	2.6-4.6	31	4.8	3.3-6.9	23	2.7	1.7-4.1
2003	117	7.3	6.0-8.8	58	8.0	6.0-10.4	59	6.7	5.1-8.8	53	3.3	2.4-4.3	33	4.7	3.2-6.6	20	2.2	1.3-3.4
2004	112	7.1	5.8-8.5	62	9.0	6.8-11.5	50	5.6	4.1-7.5	48	2.9	2.1-3.8	25	3.4	2.2-5.0	23	2.5	1.6-3.7
2005	115	7.1	5.8-8.6	58	8.0	6.0-10.4	57	6.5	4.8-8.6	57	3.5	2.6-4.5	30	4.4	2.9-6.3	27	2.8	1.8-4.1
2006	99	6.3	5.1-7.7	56	8.1	6.0-10.5	43	4.9	3.5-6.6	53	3.2	2.4-4.2	32	4.4	3.0-6.3	21	2.1	1.3-3.3
2007	110	7.2	5.8-8.7	54	7.7	5.7-10.1	56	6.7	5.0-8.9	57	3.3	2.5-4.4	33	4.4	3.0-6.2	24	2.4	1.5-3.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
Brain and other central nervous system**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	118	7.3	6.0-8.8	56	7.5	5.7-9.9	62	6.9	5.2-8.9	50	2.9	2.2-3.9	26	3.5	2.3-5.2	24	2.5	1.6-3.8
2009	104	6.3	5.1-7.7	58	7.8	5.9-10.2	46	4.8	3.5-6.6	64	3.8	2.9-4.8	34	4.5	3.1-6.4	30	3.1	2.0-4.5
2010	87	5.6	4.5-7.0	41	5.8	4.1-7.9	46	5.5	3.9-7.5	53	3.0	2.2-4.0	23	3.1	2.0-4.7	30	2.9	1.9-4.2
2011	94	5.6	4.5-7.0	46	6.2	4.5-8.4	48	5.3	3.8-7.1	60	3.5	2.6-4.5	27	3.6	2.4-5.3	33	3.3	2.2-4.7
2012	86	5.0	4.0-6.3	44	5.8	4.1-7.8	42	4.4	3.1-6.1	44	2.4	1.7-3.3	22	2.9	1.8-4.4	22	2.1	1.3-3.3
2013	129	7.6	6.3-9.1	73	9.4	7.3-11.9	56	6.2	4.6-8.1	56	3.1	2.3-4.1	31	4.4	2.9-6.3	25	2.3	1.4-3.5
2014	126	7.2	6.0-8.7	76	9.7	7.5-12.2	50	5.3	3.9-7.1	59	3.2	2.4-4.2	32	4.0	2.7-5.8	27	2.7	1.7-4.0
2015	134	8.1	6.7-9.7	71	9.8	7.6-12.5	63	6.6	5.0-8.6	56	2.9	2.2-3.8	29	3.5	2.3-5.0	27	2.4	1.6-3.6
2016	113	6.5	5.3-7.9	75	9.2	7.1-11.6	38	4.2	2.9-6.0	66	3.5	2.7-4.5	35	4.3	3.0-6.1	31	2.8	1.9-4.1
2017	124	6.8	5.6-8.2	70	8.4	6.5-10.8	54	5.5	4.1-7.4	60	3.4	2.6-4.4	33	4.1	2.8-5.8	27	3.0	1.9-4.4
2018	110	6.5	5.3-8.0	70	9.0	6.9-11.5	40	4.4	3.0-6.3	59	3.0	2.3-3.9	41	4.6	3.3-6.3	18	1.8	1.0-3.0
2019	128	6.8	5.6-8.2	59	7.0	5.2-9.1	69	6.6	5.0-8.5	69	3.6	2.8-4.6	37	4.5	3.1-6.3	32	2.8	1.9-4.0
2020	111	5.9	4.8-7.2	69	7.8	6.0-10.0	42	4.4	3.1-6.0	70	3.5	2.7-4.5	43	4.8	3.5-6.6	27	2.4	1.5-3.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Thyroid

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	51	3.4	2.5-4.5	17	2.7	1.5-4.4	34	4.1	2.8-5.8	12	0.8	0.4-1.5	6	1.1	0.4-2.5	6	0.7	0.2-1.5
1977	65	4.0	3.1-5.2	11	1.7	0.8-3.1	54	6.0	4.4-8.0	7	0.4	0.2-0.9	4	0.7	0.2-1.7	3	0.3	0.1-1.0
1978	66	4.2	3.2-5.3	16	2.3	1.3-3.9	50	5.6	4.1-7.5	12	0.9	0.4-1.6	3	0.6	0.1-1.7	9	1.1	0.5-2.2
1979	49	3.3	2.4-4.4	17	2.7	1.5-4.3	32	3.9	2.6-5.6	10	0.7	0.3-1.3	3	0.6	0.1-1.6	7	0.8	0.3-1.6
1980	54	3.5	2.6-4.6	17	2.4	1.3-3.9	37	4.5	3.1-6.4	12	0.7	0.4-1.3	4	0.7	0.2-1.7	8	0.8	0.3-1.6
1981	42	2.7	1.9-3.6	16	2.3	1.3-3.8	26	3.0	1.9-4.5	7	0.4	0.2-0.9	1	0.1	0.0-0.9	6	0.7	0.2-1.5
1982	45	3.0	2.1-4.0	16	2.2	1.3-3.7	29	3.5	2.3-5.1	9	0.6	0.3-1.2	2	0.4	0.0-1.5	7	0.8	0.3-1.8
1983	63	4.3	3.3-5.6	20	2.9	1.8-4.6	43	5.6	4.0-7.6	7	0.4	0.2-1.0	1	0.2	0.0-1.0	6	0.7	0.2-1.6
1984	74	4.9	3.8-6.2	23	3.3	2.1-5.1	51	6.4	4.7-8.4	8	0.6	0.2-1.1	2	0.4	0.0-1.4	6	0.8	0.3-1.7
1985	76	5.0	3.9-6.2	24	3.4	2.1-5.1	52	6.2	4.6-8.2	13	0.9	0.5-1.5	7	1.4	0.5-2.8	6	0.6	0.2-1.4
1986	63	4.0	3.0-5.1	22	3.4	2.1-5.2	41	4.7	3.4-6.5	9	0.6	0.3-1.2	2	0.4	0.0-1.4	7	0.7	0.3-1.6
1987	76	4.6	3.6-5.8	21	2.9	1.7-4.5	55	6.2	4.6-8.1	13	0.8	0.4-1.5	5	0.9	0.3-2.1	8	0.8	0.4-1.7
1988	67	4.4	3.4-5.6	26	3.8	2.5-5.6	41	4.9	3.5-6.7	7	0.5	0.2-1.0	2	0.3	0.0-1.2	5	0.5	0.2-1.3
1989	56	3.5	2.6-4.6	10	1.3	0.6-2.5	46	5.4	3.9-7.2	6	0.4	0.1-0.9	0	0.0	0.0-0.6	6	0.6	0.2-1.5
1990	77	4.9	3.9-6.2	20	2.9	1.7-4.5	57	6.7	5.0-8.7	10	0.7	0.3-1.3	0	0.0	0.0-0.6	10	1.1	0.5-2.1
1991	87	5.4	4.3-6.7	20	2.9	1.7-4.5	67	7.7	5.9-9.8	9	0.6	0.3-1.1	1	0.2	0.0-0.9	8	0.8	0.4-1.7
1992	83	5.4	4.3-6.7	28	4.1	2.7-6.0	55	6.6	4.9-8.7	6	0.4	0.1-0.9	3	0.5	0.1-1.5	3	0.4	0.1-1.1
1993	84	5.4	4.3-6.7	25	3.6	2.3-5.3	59	7.2	5.4-9.3	8	0.5	0.2-1.1	3	0.5	0.1-1.5	5	0.5	0.2-1.3
1994	109	6.7	5.5-8.1	28	3.9	2.5-5.8	81	9.4	7.4-11.7	6	0.4	0.1-0.9	2	0.3	0.0-1.1	4	0.4	0.1-1.2
1995	104	6.4	5.2-7.8	28	4.0	2.6-5.8	76	8.6	6.7-10.8	5	0.3	0.1-0.8	1	0.2	0.0-0.9	4	0.5	0.1-1.3
1996	111	6.7	5.5-8.1	29	4.0	2.7-5.9	82	9.2	7.3-11.5	10	0.7	0.3-1.3	3	0.5	0.1-1.5	7	0.8	0.3-1.7
1997	117	7.1	5.8-8.5	35	4.6	3.2-6.5	82	9.4	7.4-11.7	10	0.6	0.3-1.2	4	0.6	0.2-1.5	6	0.7	0.2-1.5
1998	144	8.8	7.4-10.4	37	5.1	3.6-7.1	107	12.2	10.0-14.9	7	0.5	0.2-1.0	3	0.6	0.1-1.7	4	0.5	0.1-1.2
1999	158	9.6	8.1-11.3	33	4.2	2.9-6.0	125	14.3	11.8-17.1	5	0.3	0.1-0.8	3	0.5	0.1-1.4	2	0.2	0.0-0.8
2000	150	8.7	7.3-10.2	44	5.3	3.8-7.2	106	11.6	9.5-14.1	10	0.7	0.3-1.2	1	0.2	0.0-1.1	9	1.0	0.4-1.9
2001	168	10.1	8.6-11.8	49	6.7	5.0-9.0	119	13.1	10.8-15.8	8	0.5	0.2-1.0	4	0.6	0.2-1.5	4	0.4	0.1-1.1
2002	152	9.0	7.6-10.6	48	6.3	4.6-8.4	104	11.5	9.4-14.0	8	0.5	0.2-1.0	5	0.7	0.2-1.8	3	0.3	0.1-0.9
2003	163	9.6	8.1-11.2	38	5.0	3.5-6.9	125	13.6	11.3-16.3	5	0.3	0.1-0.8	2	0.4	0.0-1.2	3	0.3	0.1-1.0
2004	197	11.4	9.9-13.2	52	7.0	5.2-9.2	145	15.7	13.2-18.5	9	0.5	0.2-1.0	6	1.0	0.4-2.2	3	0.3	0.1-0.9
2005	256	15.0	13.2-17.0	60	7.5	5.7-9.8	196	21.8	18.8-25.2	16	1.0	0.6-1.6	4	0.6	0.2-1.5	12	1.2	0.6-2.1
2006	273	15.7	13.9-17.7	77	9.5	7.4-11.9	196	21.6	18.7-25.0	13	0.8	0.4-1.3	6	1.0	0.3-2.1	7	0.6	0.3-1.4
2007	266	15.2	13.4-17.2	64	7.9	6.1-10.2	202	21.7	18.8-25.1	11	0.7	0.3-1.2	5	0.7	0.2-1.8	6	0.6	0.2-1.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Thyroid

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	317	18.0	16.0-20.1	80	10.3	8.1-12.9	237	25.3	22.1-28.8	12	0.7	0.4-1.3	5	0.8	0.2-1.8	7	0.7	0.3-1.6
2009	360	20.6	18.5-22.9	96	11.5	9.2-14.1	264	29.0	25.5-32.9	10	0.6	0.3-1.1	6	0.9	0.3-1.9	4	0.4	0.1-1.2
2010	363	20.9	18.7-23.2	89	11.3	9.1-14.0	274	29.7	26.2-33.6	5	0.3	0.1-0.7	2	0.3	0.0-1.1	3	0.3	0.1-0.9
2011	346	19.5	17.5-21.7	71	8.8	6.8-11.2	275	29.4	26.0-33.3	6	0.3	0.1-0.7	2	0.2	0.0-0.9	4	0.4	0.1-1.0
2012	324	18.4	16.4-20.6	96	11.4	9.2-14.0	228	25.0	21.7-28.6	7	0.4	0.2-0.8	2	0.3	0.0-1.1	5	0.5	0.2-1.3
2013	372	20.6	18.5-22.9	98	11.7	9.4-14.3	274	28.9	25.5-32.7	16	0.9	0.5-1.5	10	1.4	0.7-2.5	6	0.5	0.2-1.2
2014	392	22.0	19.8-24.3	124	14.9	12.3-17.9	268	28.7	25.2-32.5	14	0.7	0.4-1.3	4	0.5	0.1-1.4	10	0.9	0.4-1.7
2015	328	17.9	15.9-20.0	92	10.5	8.4-13.0	236	24.9	21.7-28.5	7	0.3	0.1-0.7	5	0.6	0.2-1.4	2	0.1	0.0-0.7
2016	368	20.3	18.2-22.5	94	10.9	8.8-13.5	274	28.9	25.4-32.7	8	0.4	0.2-0.8	1	0.1	0.0-0.8	7	0.6	0.3-1.4
2017	311	17.0	15.1-19.1	81	9.1	7.2-11.4	230	24.4	21.2-28.0	8	0.4	0.2-0.9	5	0.6	0.2-1.5	3	0.3	0.1-0.9
2018	345	19.0	17.0-21.2	97	11.6	9.4-14.3	248	25.6	22.4-29.2	9	0.4	0.2-0.8	4	0.5	0.1-1.3	5	0.4	0.1-1.0
2019	348	18.8	16.8-20.9	96	10.9	8.8-13.4	252	26.3	23.1-30.0	5	0.2	0.1-0.6	2	0.2	0.0-0.9	3	0.2	0.0-0.8
2020	260	14.4	12.7-16.4	79	9.4	7.4-11.8	181	19.3	16.4-22.5	9	0.4	0.2-0.8	2	0.2	0.0-0.9	7	0.5	0.2-1.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Hodgkin lymphoma

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	54	3.4	2.5-4.4	29	4.1	2.7-5.9	25	2.7	1.7-4.1	13	0.9	0.5-1.6	8	1.1	0.5-2.2	5	0.6	0.2-1.5
1977	46	2.8	2.1-3.8	28	3.9	2.5-5.7	18	2.1	1.2-3.4	26	1.7	1.1-2.5	14	2.2	1.1-3.7	12	1.4	0.7-2.6
1978	60	3.7	2.8-4.8	28	3.8	2.5-5.5	32	3.5	2.3-5.0	17	1.1	0.6-1.7	9	1.4	0.6-2.7	8	0.9	0.4-1.8
1979	59	3.8	2.9-5.0	27	4.1	2.6-6.0	32	3.5	2.3-5.0	12	0.8	0.4-1.4	7	0.9	0.3-1.9	5	0.6	0.2-1.5
1980	49	3.1	2.2-4.1	26	3.5	2.3-5.3	23	2.7	1.7-4.2	17	1.1	0.6-1.7	6	0.9	0.3-2.0	11	1.2	0.6-2.2
1981	41	2.6	1.8-3.6	20	2.8	1.6-4.4	21	2.4	1.4-3.8	17	1.1	0.7-1.9	7	1.0	0.4-2.1	10	1.2	0.6-2.3
1982	52	3.5	2.6-4.6	34	5.0	3.4-7.1	18	2.2	1.3-3.6	11	0.7	0.3-1.3	4	0.6	0.1-1.5	7	0.8	0.3-1.7
1983	66	3.8	2.9-4.9	38	5.2	3.6-7.2	28	2.8	1.8-4.1	21	1.3	0.8-2.0	13	1.8	0.9-3.2	8	0.9	0.4-1.8
1984	57	3.5	2.6-4.5	30	4.3	2.8-6.2	27	2.9	1.9-4.3	16	1.0	0.5-1.6	12	1.6	0.8-2.9	4	0.5	0.1-1.2
1985	68	4.1	3.2-5.3	38	5.1	3.5-7.1	30	3.1	2.1-4.5	21	1.3	0.8-2.0	10	1.5	0.7-2.8	11	1.1	0.6-2.1
1986	69	4.3	3.3-5.5	48	6.7	4.9-9.0	21	2.3	1.4-3.6	22	1.4	0.9-2.1	17	2.5	1.4-4.1	5	0.5	0.2-1.2
1987	47	3.0	2.2-4.0	32	4.6	3.1-6.6	15	1.6	0.9-2.8	17	1.1	0.7-1.9	12	1.8	0.9-3.2	5	0.6	0.2-1.5
1988	52	3.1	2.3-4.2	36	4.8	3.3-6.8	16	1.7	1.0-2.9	8	0.5	0.2-1.0	4	0.6	0.2-1.7	4	0.5	0.1-1.3
1989	52	3.1	2.3-4.2	32	4.1	2.8-5.9	20	2.2	1.3-3.6	13	0.8	0.4-1.5	11	1.6	0.8-2.9	2	0.2	0.0-0.9
1990	44	2.8	2.0-3.8	26	3.7	2.4-5.5	18	2.1	1.2-3.5	15	0.9	0.5-1.6	9	1.3	0.6-2.6	6	0.6	0.2-1.3
1991	69	4.6	3.5-5.9	39	5.4	3.7-7.5	30	3.9	2.6-5.6	9	0.6	0.3-1.1	3	0.4	0.1-1.2	6	0.6	0.2-1.5
1992	54	3.4	2.5-4.5	31	4.3	2.9-6.2	23	2.6	1.6-4.1	13	0.8	0.4-1.4	8	1.1	0.4-2.2	5	0.6	0.2-1.4
1993	64	4.0	3.0-5.1	37	5.0	3.5-7.0	27	3.1	2.0-4.6	11	0.6	0.3-1.2	8	1.1	0.5-2.2	3	0.3	0.1-0.9
1994	55	3.5	2.6-4.6	28	3.9	2.6-5.7	27	3.1	2.0-4.6	10	0.7	0.3-1.3	4	0.7	0.2-1.7	6	0.7	0.2-1.5
1995	65	3.8	2.9-5.0	32	4.2	2.8-6.0	33	3.5	2.4-5.1	5	0.3	0.1-0.8	3	0.5	0.1-1.5	2	0.2	0.0-0.8
1996	40	2.5	1.8-3.5	23	3.1	1.9-4.8	17	1.9	1.1-3.1	7	0.4	0.2-0.9	3	0.4	0.1-1.2	4	0.4	0.1-1.1
1997	45	2.7	1.9-3.6	28	3.6	2.4-5.4	17	1.8	1.0-3.0	10	0.6	0.3-1.2	3	0.5	0.1-1.6	7	0.8	0.3-1.7
1998	58	3.5	2.6-4.6	32	4.1	2.8-5.9	26	3.0	1.9-4.5	4	0.3	0.1-0.7	2	0.3	0.0-1.1	2	0.2	0.0-0.9
1999	63	3.9	3.0-5.0	33	4.2	2.9-6.0	30	3.5	2.3-5.1	7	0.5	0.2-0.9	4	0.6	0.1-1.5	3	0.3	0.1-1.0
2000	68	4.1	3.1-5.2	31	3.9	2.6-5.7	37	4.3	3.0-6.0	12	0.7	0.4-1.3	6	0.7	0.3-1.7	6	0.6	0.2-1.5
2001	66	3.8	2.9-4.9	32	4.1	2.8-5.9	34	3.6	2.4-5.1	10	0.6	0.3-1.2	5	0.7	0.2-1.7	5	0.6	0.2-1.4
2002	67	4.1	3.1-5.2	37	4.8	3.3-6.7	30	3.4	2.2-4.9	8	0.5	0.2-1.0	7	1.0	0.4-2.2	1	0.1	0.0-0.7
2003	48	2.9	2.1-3.9	23	2.9	1.8-4.4	25	2.9	1.8-4.4	10	0.6	0.3-1.2	6	0.9	0.3-1.9	4	0.4	0.1-1.2
2004	49	2.8	2.0-3.7	26	3.0	1.9-4.5	23	2.5	1.6-3.8	6	0.4	0.1-0.8	3	0.4	0.1-1.2	3	0.3	0.1-1.0
2005	68	3.9	3.0-5.0	40	5.1	3.6-7.1	28	2.8	1.8-4.1	7	0.5	0.2-0.9	4	0.5	0.1-1.4	3	0.4	0.1-1.1
2006	59	3.5	2.7-4.6	40	5.2	3.7-7.1	19	2.0	1.2-3.2	1	0.0	0.0-0.3	0	0.0	0.0-0.6	1	0.1	0.0-0.6
2007	51	2.9	2.1-3.9	24	3.0	1.9-4.5	27	2.8	1.8-4.1	6	0.3	0.1-0.7	1	0.1	0.0-0.7	5	0.5	0.1-1.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Hodgkin lymphoma

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	67	3.7	2.9-4.8	34	4.1	2.8-5.8	33	3.2	2.2-4.6	10	0.6	0.3-1.1	6	0.9	0.3-2.0	4	0.4	0.1-1.1
2009	66	3.8	2.9-4.9	41	5.1	3.6-7.0	25	2.6	1.6-4.0	4	0.2	0.1-0.6	2	0.3	0.0-1.0	2	0.2	0.0-0.9
2010	67	3.6	2.8-4.7	37	4.3	3.0-6.0	30	3.0	2.0-4.4	6	0.3	0.1-0.7	4	0.4	0.1-1.2	2	0.2	0.0-0.9
2011	79	4.2	3.3-5.3	44	5.0	3.6-6.8	35	3.5	2.4-5.0	14	0.8	0.4-1.3	7	0.9	0.4-1.9	7	0.7	0.3-1.4
2012	51	2.8	2.1-3.8	24	2.8	1.8-4.3	27	2.9	1.8-4.3	11	0.6	0.3-1.1	5	0.6	0.2-1.5	6	0.5	0.2-1.3
2013	65	3.7	2.8-4.8	32	4.0	2.7-5.8	33	3.4	2.3-4.9	11	0.6	0.3-1.1	7	1.0	0.4-2.0	4	0.4	0.1-1.1
2014	70	3.9	3.0-5.1	35	4.3	2.9-6.1	35	3.7	2.5-5.2	5	0.3	0.1-0.7	3	0.4	0.1-1.1	2	0.2	0.0-0.8
2015	49	2.7	2.0-3.7	25	3.0	1.9-4.6	24	2.4	1.5-3.7	7	0.3	0.1-0.8	5	0.6	0.2-1.5	2	0.2	0.0-0.8
2016	52	2.9	2.1-3.8	21	2.6	1.6-4.1	31	3.0	2.0-4.4	5	0.2	0.1-0.5	2	0.2	0.0-0.8	3	0.2	0.0-0.8
2017	55	3.1	2.3-4.1	24	2.7	1.7-4.1	31	3.5	2.3-5.1	5	0.3	0.1-0.7	2	0.2	0.0-0.9	3	0.3	0.1-0.9
2018	47	2.5	1.8-3.4	28	3.3	2.1-4.9	19	1.9	1.1-3.1	3	0.2	0.0-0.5	3	0.4	0.1-1.1	0	0.0	0.0-0.5
2019	50	2.7	1.9-3.6	28	3.2	2.0-4.7	22	2.2	1.3-3.5	2	0.1	0.0-0.4	1	0.1	0.0-0.7	1	0.1	0.0-0.6
2020	51	2.9	2.1-3.9	27	3.2	2.1-4.8	24	2.6	1.6-3.9	6	0.2	0.1-0.6	4	0.4	0.1-1.1	2	0.1	0.0-0.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
Non-Hodgkin lymphomas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	184	11.6	9.9-13.4	84	12.9	10.2-16.1	100	10.8	8.7-13.2	92	6.1	4.9-7.5	49	8.1	5.9-10.8	43	4.7	3.4-6.4
1977	173	11.4	9.7-13.3	78	12.6	9.8-15.8	95	10.7	8.6-13.1	87	5.7	4.5-7.0	37	6.1	4.2-8.6	50	5.4	3.9-7.2
1978	194	13.0	11.2-15.0	98	15.4	12.4-18.8	96	11.1	8.9-13.6	93	6.3	5.0-7.7	47	7.8	5.7-10.5	46	5.2	3.8-7.1
1979	191	12.4	10.6-14.3	85	14.1	11.1-17.5	106	11.2	9.1-13.7	103	7.0	5.7-8.5	54	9.1	6.7-12.0	49	5.3	3.9-7.1
1980	208	13.9	12.0-16.0	103	16.8	13.7-20.5	105	11.4	9.3-13.9	95	6.3	5.1-7.8	44	7.0	5.0-9.5	51	5.7	4.2-7.5
1981	190	12.7	10.9-14.7	87	13.8	11.0-17.1	103	11.7	9.5-14.3	102	6.9	5.6-8.5	44	7.2	5.2-9.8	58	6.6	5.0-8.6
1982	206	13.5	11.7-15.5	100	15.3	12.4-18.8	106	11.4	9.3-13.8	101	6.9	5.6-8.4	51	8.6	6.4-11.4	50	5.5	4.1-7.4
1983	251	17.3	15.2-19.6	142	23.0	19.3-27.3	109	12.5	10.2-15.2	119	8.1	6.7-9.7	57	9.4	7.1-12.3	62	6.7	5.1-8.6
1984	247	16.6	14.5-18.8	136	21.7	18.1-25.8	111	12.5	10.2-15.2	97	6.4	5.2-7.9	53	8.4	6.2-11.1	44	4.8	3.5-6.5
1985	296	19.5	17.3-21.9	184	27.8	23.8-32.3	112	12.3	10.1-14.9	126	8.3	6.9-9.9	72	11.5	8.9-14.6	54	5.9	4.4-7.8
1986	294	19.1	16.9-21.5	181	27.1	23.1-31.5	113	12.7	10.4-15.4	128	8.7	7.3-10.4	68	11.4	8.7-14.5	60	6.7	5.1-8.7
1987	318	21.3	19.0-23.8	214	32.6	28.2-37.4	104	11.7	9.5-14.2	112	7.4	6.1-9.0	62	10.1	7.7-13.1	50	5.4	4.0-7.2
1988	317	20.5	18.3-23.0	211	30.7	26.5-35.3	106	11.9	9.7-14.5	130	8.7	7.3-10.4	80	12.6	9.9-15.8	50	5.3	4.0-7.1
1989	350	22.9	20.5-25.5	239	34.6	30.2-39.4	111	12.6	10.3-15.2	137	9.1	7.6-10.8	87	13.9	11.1-17.3	50	5.4	4.0-7.2
1990	369	23.9	21.5-26.6	251	36.1	31.6-41.0	118	13.5	11.1-16.2	132	8.6	7.2-10.3	68	10.7	8.2-13.7	64	6.8	5.2-8.8
1991	445	28.8	26.1-31.6	304	44.4	39.4-49.9	141	15.8	13.3-18.8	123	8.2	6.8-9.8	76	12.0	9.4-15.1	47	5.1	3.7-6.8
1992	428	27.6	25.0-30.4	272	38.9	34.2-44.0	156	17.7	15.0-20.7	134	8.9	7.4-10.6	74	12.0	9.4-15.2	60	6.4	4.8-8.2
1993	434	28.3	25.6-31.1	295	42.8	37.9-48.2	139	16.1	13.5-19.1	113	7.6	6.3-9.2	56	9.2	6.9-12.1	57	6.4	4.9-8.4
1994	517	33.7	30.8-36.8	350	49.2	44.0-54.8	167	19.8	16.9-23.1	129	8.6	7.2-10.2	68	11.0	8.5-14.0	61	6.7	5.1-8.6
1995	414	26.7	24.2-29.5	279	40.4	35.7-45.6	135	15.1	12.6-17.9	135	9.2	7.7-10.8	70	11.7	9.1-14.9	65	7.2	5.5-9.2
1996	418	27.0	24.4-29.7	237	34.2	29.8-39.0	181	20.9	17.9-24.2	125	8.2	6.9-9.8	65	10.9	8.4-14.0	60	6.6	5.0-8.5
1997	357	23.1	20.7-25.6	218	32.6	28.2-37.3	139	15.9	13.3-18.8	118	7.9	6.5-9.4	78	13.3	10.4-16.7	40	4.4	3.2-6.1
1998	351	22.7	20.4-25.2	216	31.9	27.7-36.5	135	15.3	12.8-18.1	110	7.3	6.0-8.9	55	8.7	6.5-11.3	55	6.1	4.6-7.9
1999	359	23.1	20.8-25.7	205	30.2	26.1-34.8	154	17.2	14.6-20.2	121	8.1	6.7-9.7	66	11.6	8.9-14.8	55	6.0	4.5-7.9
2000	335	21.5	19.3-24.0	188	27.1	23.3-31.4	147	16.7	14.1-19.7	102	6.6	5.4-8.1	63	10.2	7.8-13.1	39	4.2	3.0-5.8
2001	354	22.6	20.3-25.1	188	28.7	24.6-33.2	166	18.7	15.9-21.8	116	7.4	6.1-8.9	66	10.6	8.1-13.5	50	5.2	3.8-6.9
2002	321	20.4	18.2-22.8	194	28.5	24.6-32.9	127	14.2	11.8-17.0	104	6.6	5.4-8.0	57	9.0	6.8-11.7	47	5.1	3.7-6.8
2003	403	25.1	22.7-27.8	208	29.9	25.9-34.4	195	21.2	18.3-24.5	109	6.8	5.6-8.2	56	8.6	6.4-11.2	53	5.5	4.1-7.2
2004	392	24.2	21.8-26.7	225	32.2	28.1-36.8	167	17.9	15.3-20.9	106	6.5	5.4-7.9	64	10.2	7.8-13.1	42	4.2	3.0-5.7
2005	387	23.7	21.4-26.3	207	29.3	25.4-33.7	180	19.3	16.5-22.4	113	7.0	5.7-8.4	67	10.4	8.0-13.2	46	4.7	3.4-6.3
2006	355	21.5	19.3-23.9	202	28.3	24.5-32.5	153	16.4	13.8-19.2	123	7.5	6.2-9.0	70	10.4	8.1-13.2	53	5.5	4.1-7.2
2007	391	23.7	21.3-26.2	220	30.4	26.5-34.8	171	17.7	15.1-20.7	109	6.7	5.5-8.1	64	9.6	7.3-12.2	45	4.7	3.4-6.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
Non-Hodgkin lymphomas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	407	24.0	21.7-26.4	232	31.6	27.6-36.0	175	18.1	15.5-21.1	98	5.7	4.6-7.0	54	7.9	5.9-10.3	44	4.2	3.0-5.7
2009	410	24.2	21.9-26.7	214	29.3	25.4-33.5	196	20.2	17.4-23.3	93	5.5	4.4-6.7	56	7.8	5.9-10.2	37	3.7	2.6-5.2
2010	399	23.1	20.9-25.6	221	29.8	25.9-34.1	178	18.3	15.7-21.3	79	4.6	3.6-5.7	46	6.6	4.8-8.8	33	3.2	2.2-4.6
2011	397	22.7	20.5-25.2	214	27.9	24.3-32.0	183	19.0	16.3-22.1	97	5.5	4.5-6.8	56	7.7	5.8-10.0	41	3.9	2.8-5.3
2012	390	22.0	19.8-24.4	209	27.1	23.5-31.2	181	18.1	15.5-21.0	87	4.8	3.8-5.9	55	7.5	5.6-9.8	32	2.8	1.9-4.1
2013	425	23.5	21.3-25.9	241	30.5	26.7-34.7	184	17.5	15.0-20.3	73	4.0	3.1-5.0	41	5.5	3.9-7.5	32	2.9	2.0-4.2
2014	429	23.2	21.0-25.5	228	28.6	25.0-32.7	201	18.9	16.4-21.9	95	4.9	4.0-6.1	46	5.9	4.3-8.0	49	4.1	3.0-5.5
2015	424	22.5	20.4-24.8	236	28.5	24.9-32.5	188	17.9	15.3-20.8	80	4.1	3.2-5.1	43	5.2	3.8-7.1	37	3.4	2.3-4.7
2016	441	22.7	20.5-24.9	227	27.0	23.5-30.8	214	19.4	16.8-22.3	91	4.6	3.7-5.7	49	5.8	4.3-7.7	42	3.6	2.6-4.9
2017	445	23.0	20.9-25.3	219	25.8	22.4-29.5	226	20.9	18.1-23.9	89	4.4	3.5-5.4	46	5.5	4.0-7.4	43	3.6	2.5-4.9
2018	425	22.3	20.2-24.6	236	28.7	25.1-32.7	189	17.3	14.8-20.1	100	4.8	3.9-5.9	60	7.1	5.4-9.1	40	3.1	2.2-4.4
2019	465	22.9	20.8-25.2	238	26.9	23.5-30.6	227	20.0	17.4-22.9	70	3.3	2.6-4.2	38	4.4	3.1-6.1	32	2.5	1.7-3.7
2020	348	17.6	15.7-19.6	171	19.4	16.5-22.6	177	16.4	14.0-19.2	85	3.8	3.1-4.8	53	5.9	4.4-7.8	32	2.4	1.6-3.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
Myeloma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	89	5.9	4.7-7.3	42	6.9	4.9-9.4	47	5.5	4.0-7.3	39	2.7	1.9-3.7	16	2.8	1.5-4.6	23	2.7	1.7-4.1
1977	87	5.8	4.6-7.2	38	6.5	4.5-9.1	49	5.5	4.0-7.3	50	3.4	2.5-4.5	22	3.9	2.4-6.0	28	3.0	2.0-4.4
1978	77	5.0	3.9-6.3	40	6.6	4.6-9.1	37	3.9	2.8-5.5	45	3.1	2.2-4.2	21	3.5	2.1-5.5	24	2.8	1.8-4.3
1979	90	6.1	4.9-7.5	43	7.0	5.0-9.5	47	5.2	3.8-7.0	57	3.8	2.8-4.9	24	3.9	2.4-5.8	33	3.7	2.5-5.2
1980	83	5.6	4.5-7.0	44	7.6	5.5-10.3	39	4.3	3.1-6.0	61	4.0	3.0-5.2	30	5.3	3.5-7.7	31	3.3	2.2-4.7
1981	101	6.5	5.3-8.0	56	9.3	7.0-12.2	45	4.7	3.4-6.4	59	3.9	3.0-5.0	31	5.5	3.7-7.8	28	3.0	2.0-4.4
1982	87	5.7	4.6-7.1	34	6.0	4.1-8.4	53	5.7	4.2-7.5	60	4.1	3.1-5.3	25	4.5	2.9-6.8	35	4.0	2.7-5.6
1983	77	5.2	4.1-6.5	38	6.4	4.5-8.9	39	4.2	3.0-5.9	42	2.8	2.0-3.9	20	3.3	2.0-5.1	22	2.4	1.5-3.8
1984	81	5.3	4.2-6.6	33	5.7	3.9-8.1	48	5.0	3.7-6.7	50	3.3	2.4-4.4	20	3.5	2.1-5.4	30	3.3	2.2-4.7
1985	73	4.8	3.7-6.0	36	6.3	4.4-8.9	37	4.0	2.8-5.6	42	2.8	2.0-3.8	23	4.4	2.8-6.6	19	2.0	1.2-3.1
1986	83	5.5	4.4-6.9	44	7.7	5.5-10.3	39	4.3	3.0-5.9	40	2.6	1.8-3.5	18	3.1	1.8-4.9	22	2.2	1.4-3.4
1987	93	6.1	4.9-7.5	55	9.4	7.1-12.3	38	4.0	2.8-5.6	47	3.2	2.3-4.2	27	4.9	3.2-7.2	20	2.1	1.3-3.3
1988	83	5.6	4.4-6.9	38	6.7	4.7-9.2	45	4.8	3.5-6.5	52	3.4	2.5-4.5	28	5.2	3.4-7.5	24	2.5	1.6-3.7
1989	90	6.0	4.8-7.4	38	6.7	4.7-9.2	52	5.7	4.3-7.6	52	3.4	2.5-4.5	23	4.1	2.5-6.1	29	3.1	2.0-4.4
1990	96	6.4	5.2-7.8	50	8.8	6.5-11.7	46	5.4	3.9-7.2	60	4.0	3.1-5.2	29	5.4	3.6-7.8	31	3.3	2.3-4.8
1991	91	6.1	4.9-7.5	42	6.8	4.8-9.2	49	5.6	4.1-7.5	49	3.4	2.5-4.4	27	4.6	3.0-6.7	22	2.5	1.6-3.9
1992	99	6.6	5.3-8.0	46	8.3	6.0-11.1	53	5.8	4.3-7.6	51	3.4	2.5-4.5	27	5.1	3.3-7.4	24	2.6	1.7-4.0
1993	96	6.5	5.3-8.0	38	6.4	4.5-8.9	58	6.5	4.9-8.4	57	3.8	2.9-5.0	25	4.4	2.8-6.5	32	3.4	2.3-4.9
1994	98	6.6	5.4-8.1	47	7.7	5.6-10.2	51	5.9	4.4-7.8	44	2.9	2.1-4.0	19	3.3	1.9-5.1	25	2.7	1.7-4.0
1995	85	5.8	4.6-7.1	40	6.7	4.7-9.2	45	5.2	3.8-7.0	60	4.0	3.1-5.2	30	5.1	3.4-7.3	30	3.3	2.2-4.7
1996	109	7.3	6.0-8.9	48	8.5	6.2-11.3	61	7.1	5.4-9.1	53	3.6	2.7-4.7	18	3.2	1.9-5.1	35	3.9	2.7-5.5
1997	85	5.7	4.6-7.1	43	7.4	5.3-10.0	42	4.8	3.4-6.5	42	2.8	2.1-3.9	15	2.9	1.6-4.8	27	3.0	2.0-4.4
1998	102	6.9	5.6-8.4	50	8.1	6.0-10.7	52	5.9	4.4-7.8	43	2.9	2.1-4.0	17	2.6	1.5-4.2	26	3.1	2.0-4.5
1999	104	6.9	5.6-8.3	47	7.9	5.7-10.5	57	6.4	4.8-8.3	52	3.5	2.6-4.6	25	4.4	2.8-6.5	27	3.0	2.0-4.5
2000	101	6.5	5.3-8.0	43	6.8	4.9-9.2	58	6.5	4.9-8.4	55	3.6	2.7-4.8	26	4.4	2.9-6.5	29	3.0	2.0-4.4
2001	101	6.6	5.4-8.0	46	7.3	5.3-9.8	55	6.1	4.6-8.0	51	3.3	2.4-4.3	22	3.9	2.4-5.9	29	3.0	2.0-4.4
2002	89	5.7	4.6-7.0	47	7.6	5.5-10.1	42	4.6	3.3-6.3	59	3.8	2.9-4.9	29	4.8	3.2-6.9	30	3.2	2.1-4.6
2003	110	6.8	5.6-8.3	53	8.3	6.2-10.9	57	6.0	4.5-7.9	54	3.4	2.6-4.5	21	3.5	2.1-5.3	33	3.5	2.4-5.0
2004	100	6.1	4.9-7.4	57	8.5	6.4-11.0	43	4.6	3.3-6.2	49	3.1	2.3-4.1	22	3.7	2.3-5.5	27	2.8	1.8-4.1
2005	138	8.5	7.1-10.0	65	9.5	7.3-12.1	73	7.7	6.0-9.8	48	3.0	2.2-4.0	18	2.9	1.7-4.6	30	3.2	2.1-4.6
2006	121	7.4	6.1-8.9	58	8.7	6.6-11.3	63	6.6	5.0-8.4	52	3.2	2.4-4.2	32	4.8	3.3-6.8	20	2.1	1.3-3.4
2007	107	6.5	5.3-7.8	54	7.7	5.8-10.1	53	5.5	4.1-7.3	48	2.9	2.2-3.9	22	3.3	2.1-5.0	26	2.7	1.7-3.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

New York (Manhattan)

Myeloma

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	146	8.7	7.3-10.2	79	11.2	8.9-14.0	67	6.9	5.3-8.8	50	3.0	2.2-3.9	22	3.2	2.0-4.8	28	2.8	1.9-4.2
2009	141	8.3	7.0-9.8	71	9.8	7.6-12.4	70	7.1	5.5-9.0	49	2.8	2.1-3.8	26	3.8	2.5-5.6	23	2.3	1.4-3.5
2010	148	8.6	7.3-10.2	80	10.8	8.5-13.5	68	7.0	5.4-9.0	62	3.6	2.8-4.7	34	5.1	3.5-7.1	28	2.7	1.8-4.0
2011	165	9.2	7.8-10.7	102	13.4	10.9-16.3	63	6.3	4.8-8.1	42	2.3	1.7-3.2	17	2.3	1.3-3.7	25	2.3	1.5-3.5
2012	146	7.9	6.7-9.3	89	11.4	9.1-14.1	57	5.3	4.0-6.9	53	2.9	2.2-3.8	26	3.4	2.2-5.0	27	2.5	1.6-3.7
2013	148	8.0	6.7-9.4	68	8.4	6.5-10.7	80	7.5	5.9-9.4	55	2.9	2.2-3.8	35	4.5	3.1-6.3	20	1.7	1.1-2.8
2014	159	8.5	7.2-9.9	93	11.7	9.4-14.4	66	6.0	4.6-7.7	65	3.5	2.7-4.4	36	4.5	3.1-6.3	29	2.7	1.8-3.9
2015	164	8.5	7.2-9.9	82	9.9	7.9-12.4	82	7.4	5.9-9.3	52	2.7	2.0-3.6	29	3.7	2.5-5.3	23	1.9	1.2-3.0
2016	160	8.0	6.8-9.4	92	10.7	8.6-13.2	68	6.1	4.7-7.8	60	3.0	2.3-3.9	41	5.0	3.6-6.8	19	1.7	1.0-2.8
2017	169	8.4	7.1-9.8	82	9.7	7.7-12.2	87	7.3	5.9-9.2	45	2.2	1.6-3.0	21	2.5	1.5-3.9	24	2.0	1.3-3.1
2018	147	7.4	6.2-8.7	78	9.0	7.1-11.3	69	5.9	4.5-7.5	40	1.9	1.3-2.6	15	1.7	0.9-2.8	25	2.1	1.3-3.2
2019	164	8.0	6.8-9.4	95	10.5	8.5-12.9	69	6.2	4.8-7.9	52	2.4	1.8-3.2	25	2.8	1.8-4.1	27	2.1	1.3-3.1
2020	136	6.7	5.6-7.9	84	9.5	7.5-11.8	52	4.6	3.4-6.1	40	1.8	1.3-2.5	25	2.8	1.8-4.2	15	1.2	0.7-2.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
Leukemias**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	163	11.2	9.5-13.2	90	15.1	12.0-18.8	73	8.4	6.5-10.6	135	9.4	7.8-11.2	68	11.5	8.8-14.7	67	7.9	6.1-10.2
1977	145	9.6	8.1-11.4	76	11.9	9.3-15.0	69	8.0	6.1-10.2	120	8.2	6.7-9.8	66	10.8	8.3-13.9	54	6.5	4.8-8.5
1978	166	11.4	9.7-13.3	86	14.5	11.5-18.0	80	9.3	7.3-11.7	139	9.6	8.0-11.4	72	11.9	9.2-15.1	67	8.0	6.1-10.3
1979	168	11.7	10.0-13.7	88	14.8	11.8-18.4	80	9.6	7.5-12.1	113	7.9	6.5-9.5	55	10.0	7.4-13.1	58	6.9	5.2-9.0
1980	199	13.6	11.7-15.6	100	16.6	13.4-20.3	99	11.6	9.3-14.2	115	7.7	6.3-9.3	57	10.0	7.5-13.1	58	6.4	4.8-8.4
1981	177	12.1	10.4-14.1	99	16.7	13.5-20.5	78	8.7	6.8-11.0	128	9.0	7.4-10.7	58	9.8	7.3-12.7	70	8.5	6.5-10.9
1982	188	12.9	11.1-14.9	96	16.4	13.2-20.1	92	10.7	8.6-13.3	125	8.6	7.2-10.3	74	13.2	10.3-16.7	51	5.8	4.3-7.7
1983	169	11.7	9.9-13.6	78	13.3	10.4-16.8	91	10.7	8.5-13.3	98	6.9	5.6-8.4	46	8.3	6.0-11.2	52	6.1	4.5-8.1
1984	153	10.6	8.9-12.5	83	14.5	11.5-18.1	70	8.2	6.3-10.5	113	7.6	6.3-9.2	54	9.5	7.1-12.5	59	6.7	5.0-8.8
1985	168	11.4	9.7-13.3	87	14.9	11.8-18.5	81	9.2	7.2-11.5	119	8.2	6.8-9.8	61	10.9	8.3-14.1	58	6.3	4.8-8.2
1986	189	12.7	11.0-14.8	98	16.4	13.2-20.1	91	10.6	8.5-13.2	107	7.1	5.8-8.6	57	9.4	7.1-12.3	50	5.4	4.0-7.1
1987	174	11.5	9.8-13.4	77	11.9	9.3-15.0	97	10.6	8.6-13.0	96	6.4	5.2-7.8	43	7.2	5.2-9.8	53	5.8	4.3-7.7
1988	184	12.9	11.1-14.9	93	16.5	13.3-20.3	91	10.8	8.6-13.4	118	7.9	6.5-9.4	65	11.1	8.5-14.2	53	5.6	4.2-7.4
1989	186	12.8	11.0-14.9	100	17.9	14.5-21.9	86	10.1	8.0-12.6	104	7.0	5.7-8.5	51	9.3	6.9-12.3	53	5.7	4.2-7.5
1990	189	13.1	11.2-15.1	109	18.1	14.7-21.9	80	9.5	7.5-11.9	114	8.0	6.5-9.6	51	9.2	6.8-12.2	63	7.3	5.6-9.5
1991	202	13.7	11.9-15.8	111	17.9	14.6-21.6	91	10.6	8.5-13.0	98	6.8	5.5-8.4	56	9.8	7.4-12.8	42	5.1	3.6-7.0
1992	196	13.6	11.7-15.6	104	17.8	14.5-21.7	92	10.5	8.4-12.9	106	7.3	6.0-8.9	47	8.1	5.9-10.8	59	6.5	4.9-8.5
1993	188	12.7	10.9-14.7	87	14.1	11.2-17.5	101	12.1	9.8-14.8	101	6.9	5.6-8.4	53	9.1	6.7-11.9	48	5.3	3.9-7.1
1994	198	13.5	11.7-15.5	97	15.4	12.4-18.9	101	11.9	9.6-14.5	124	8.4	7.0-10.0	56	9.3	6.9-12.1	68	7.5	5.8-9.5
1995	145	10.0	8.4-11.8	77	13.0	10.2-16.3	68	7.7	6.0-9.8	106	7.2	5.9-8.7	63	10.8	8.2-13.9	43	4.8	3.4-6.5
1996	190	12.8	11.1-14.8	105	16.9	13.8-20.6	85	9.8	7.8-12.2	93	6.3	5.1-7.7	44	7.5	5.4-10.2	49	5.4	4.0-7.3
1997	188	12.8	11.0-14.8	90	14.6	11.7-18.0	98	11.6	9.4-14.2	99	6.7	5.4-8.2	54	9.6	7.2-12.6	45	5.0	3.6-6.8
1998	172	11.8	10.1-13.7	99	15.8	12.8-19.3	73	8.8	6.8-11.1	95	6.3	5.1-7.7	51	8.2	6.1-10.9	44	4.9	3.6-6.7
1999	199	13.2	11.5-15.2	96	15.1	12.2-18.5	103	11.9	9.7-14.5	95	6.5	5.2-7.9	48	8.3	6.1-11.1	47	5.5	4.0-7.4
2000	186	12.5	10.8-14.5	90	14.8	11.8-18.2	96	11.0	8.8-13.5	106	7.0	5.7-8.5	50	8.0	5.9-10.6	56	6.0	4.5-7.9
2001	195	12.9	11.1-14.8	105	16.4	13.4-20.0	90	10.2	8.2-12.6	99	6.3	5.1-7.7	47	7.6	5.5-10.1	52	5.3	3.9-7.0
2002	151	9.8	8.3-11.6	84	13.3	10.5-16.5	67	7.2	5.6-9.2	114	7.5	6.1-9.0	53	8.8	6.5-11.5	61	6.8	5.1-8.8
2003	222	14.5	12.6-16.6	112	17.2	14.1-20.8	110	12.4	10.2-15.0	82	5.3	4.2-6.6	38	6.2	4.3-8.5	44	4.8	3.5-6.5
2004	192	12.4	10.6-14.3	109	16.8	13.7-20.3	83	9.2	7.3-11.4	98	6.1	4.9-7.4	52	8.1	6.0-10.7	46	4.7	3.4-6.3
2005	202	12.5	10.8-14.4	116	17.1	14.0-20.6	86	9.2	7.3-11.5	121	7.4	6.1-8.8	59	9.0	6.8-11.6	62	6.5	4.9-8.3
2006	196	12.3	10.6-14.1	111	15.9	13.0-19.2	85	9.4	7.5-11.8	113	6.8	5.6-8.2	58	8.5	6.4-11.1	55	5.6	4.2-7.4
2007	224	13.9	12.1-15.9	123	17.3	14.4-20.8	101	11.2	9.1-13.8	96	5.9	4.8-7.2	61	9.2	7.0-11.9	35	3.8	2.6-5.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**New York (Manhattan)
Leukemias**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	217	13.4	11.6-15.4	126	18.3	15.2-21.9	91	9.9	7.9-12.2	105	6.3	5.2-7.7	61	9.1	6.9-11.7	44	4.6	3.3-6.2
2009	233	14.7	12.8-16.8	127	18.7	15.6-22.4	106	11.4	9.2-14.0	106	6.5	5.3-7.9	48	7.1	5.2-9.5	58	6.0	4.5-7.8
2010	270	16.4	14.4-18.5	154	21.0	17.8-24.7	116	12.6	10.3-15.2	97	5.6	4.5-6.9	50	7.0	5.2-9.2	47	4.7	3.4-6.4
2011	265	15.8	13.9-17.9	145	19.9	16.7-23.6	120	12.7	10.4-15.3	105	6.1	5.0-7.5	55	7.8	5.8-10.2	50	5.1	3.7-6.8
2012	243	13.9	12.1-15.8	139	18.2	15.2-21.5	104	11.2	9.0-13.7	103	5.8	4.7-7.0	52	7.2	5.3-9.4	51	4.8	3.6-6.4
2013	256	14.5	12.7-16.5	162	21.7	18.4-25.4	94	9.2	7.4-11.4	103	5.8	4.7-7.1	63	8.8	6.7-11.3	40	3.6	2.5-5.0
2014	313	17.2	15.3-19.3	170	22.2	19.0-25.9	143	13.6	11.4-16.1	96	5.3	4.3-6.5	46	6.2	4.5-8.3	50	4.7	3.5-6.3
2015	267	14.4	12.7-16.3	160	20.1	17.0-23.5	107	10.2	8.3-12.5	119	6.4	5.3-7.7	61	8.2	6.2-10.6	58	5.1	3.8-6.7
2016	302	15.7	13.9-17.7	177	21.2	18.2-24.7	125	11.5	9.5-13.9	97	4.9	4.0-6.0	45	5.6	4.1-7.6	52	4.6	3.4-6.1
2017	242	12.7	11.1-14.5	143	17.4	14.6-20.5	99	9.2	7.4-11.4	94	4.7	3.8-5.9	48	5.9	4.3-7.9	46	4.0	2.9-5.4
2018	285	15.2	13.4-17.2	159	19.5	16.5-22.9	126	11.8	9.6-14.3	102	5.0	4.1-6.2	51	6.1	4.5-8.1	51	4.2	3.1-5.7
2019	254	13.2	11.6-15.1	148	17.8	15.0-21.0	106	9.7	7.8-11.9	95	4.5	3.6-5.6	56	6.7	5.0-8.8	39	3.0	2.1-4.2
2020	247	12.8	11.2-14.6	125	14.8	12.3-17.8	122	11.2	9.1-13.5	95	4.6	3.7-5.7	47	5.6	4.1-7.6	48	4.1	2.9-5.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
All Invasive Malignant Tumors**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	8,108	387.4	378.7-396.3	3,880	451.2	436.1-466.7	4,228	353.8	342.8-365.0	4,573	220.6	214.0-227.3	2,366	283.8	271.6-296.4	2,207	182.1	174.4-190.2
1977	8,025	383.7	375.0-392.5	3,860	452.4	437.3-467.9	4,165	348.3	337.4-359.5	4,525	215.2	208.7-221.8	2,355	281.4	269.4-293.8	2,170	176.2	168.6-184.1
1978	8,031	381.0	372.4-389.7	3,915	458.1	443.0-473.5	4,116	339.8	329.1-350.7	4,422	208.3	202.0-214.8	2,271	267.0	255.5-278.9	2,151	173.0	165.5-180.7
1979	8,109	383.4	374.9-392.2	3,904	454.5	439.6-469.8	4,205	346.9	336.1-358.0	4,330	205.9	199.6-212.4	2,229	266.0	254.5-278.0	2,101	170.9	163.4-178.6
1980	8,426	398.7	390.0-407.6	4,075	476.7	461.5-492.3	4,351	358.9	347.9-370.1	4,458	210.1	203.8-216.6	2,224	266.1	254.6-278.0	2,234	177.3	169.8-185.0
1981	8,525	402.9	394.1-411.8	4,063	475.6	460.4-491.2	4,462	364.7	353.7-376.0	4,272	201.9	195.8-208.3	2,183	260.6	249.2-272.3	2,089	167.5	160.2-175.1
1982	8,727	410.9	402.0-419.8	4,123	485.7	470.3-501.4	4,604	375.2	364.1-386.7	4,476	209.4	203.1-215.8	2,274	271.9	260.3-283.9	2,202	174.1	166.7-181.8
1983	8,728	407.7	399.0-416.6	4,228	494.6	479.2-510.4	4,500	362.0	351.2-373.1	4,309	200.9	194.8-207.1	2,154	258.3	247.0-269.9	2,155	168.5	161.2-175.9
1984	8,764	410.4	401.7-419.3	4,162	482.2	467.0-497.7	4,602	374.7	363.6-386.1	4,380	204.3	198.2-210.6	2,214	263.5	252.2-275.2	2,166	169.7	162.5-177.3
1985	8,949	416.5	407.7-425.4	4,294	494.4	479.2-510.0	4,655	374.0	363.0-385.2	4,326	200.6	194.6-206.8	2,194	259.7	248.5-271.2	2,132	167.1	159.9-174.6
1986	8,925	419.6	410.8-428.6	4,248	497.1	481.7-512.9	4,677	380.4	369.2-391.8	4,235	196.9	190.9-203.0	2,083	247.5	236.5-258.8	2,152	168.1	160.9-175.5
1987	9,093	427.9	418.9-436.9	4,253	495.4	480.1-511.0	4,840	394.1	382.8-405.7	4,322	202.7	196.6-209.0	2,121	254.6	243.5-266.0	2,201	173.3	165.9-180.9
1988	8,518	403.5	394.9-412.3	3,956	465.4	450.5-480.7	4,562	374.2	363.1-385.5	4,183	196.8	190.8-203.0	2,003	242.0	231.2-253.2	2,180	172.8	165.4-180.4
1989	8,445	401.4	392.8-410.2	3,994	468.2	453.4-483.4	4,451	366.3	355.3-377.5	4,105	193.9	187.9-200.0	2,030	245.6	234.7-256.9	2,075	165.0	157.8-172.4
1990	9,388	446.6	437.4-455.8	4,491	529.2	513.4-545.3	4,897	403.9	392.4-415.7	4,209	199.3	193.3-205.5	2,042	248.0	237.1-259.3	2,167	172.8	165.4-180.4
1991	9,199	437.8	428.7-446.9	4,442	523.5	507.8-539.5	4,757	391.8	380.5-403.3	4,163	197.5	191.5-203.7	2,002	244.1	233.2-255.3	2,161	172.1	164.8-179.7
1992	9,715	460.6	451.4-470.0	4,815	561.2	545.1-577.7	4,900	404.8	393.3-416.5	4,073	192.9	187.0-199.0	1,992	242.8	232.0-253.9	2,081	165.7	158.6-173.1
1993	9,423	445.0	436.0-454.2	4,733	546.9	531.2-563.1	4,690	385.9	374.8-397.4	4,011	189.5	183.6-195.5	2,011	243.6	232.9-254.8	2,000	159.1	152.1-166.3
1994	9,997	471.8	462.5-481.2	4,939	568.6	552.5-584.9	5,058	415.1	403.6-426.9	3,949	186.2	180.4-192.1	1,987	238.1	227.5-249.0	1,962	155.6	148.8-162.8
1995	9,673	454.5	445.5-463.7	4,661	534.0	518.5-549.8	5,012	407.8	396.4-419.4	3,882	182.6	176.9-188.4	1,827	218.1	208.0-228.5	2,055	162.4	155.3-169.6
1996	9,671	453.6	444.6-462.8	4,649	531.7	516.3-547.4	5,022	408.0	396.7-419.5	3,821	179.1	173.5-184.9	1,835	216.9	207.0-227.2	1,986	156.2	149.4-163.3
1997	9,677	451.0	442.0-460.1	4,719	535.9	520.5-551.6	4,958	399.9	388.7-411.3	3,725	173.9	168.4-179.6	1,854	219.6	209.6-230.0	1,871	146.5	139.9-153.3
1998	9,669	445.1	436.3-454.1	4,682	521.4	506.4-536.7	4,987	399.8	388.8-411.2	3,650	168.4	163.0-173.9	1,780	207.2	197.5-217.1	1,870	145.8	139.2-152.6
1999	9,794	442.8	434.1-451.7	4,731	519.1	504.3-534.3	5,063	398.5	387.6-409.7	3,660	165.5	160.2-170.9	1,777	202.7	193.3-212.4	1,883	144.2	137.7-150.9
2000	9,844	439.1	430.4-447.8	4,836	523.3	508.6-538.4	5,008	389.4	378.6-400.4	3,682	164.0	158.8-169.4	1,786	200.9	191.6-210.5	1,896	142.2	135.9-148.8
2001	9,722	430.1	421.6-438.8	4,868	520.8	506.1-535.8	4,854	375.3	364.7-386.0	3,580	158.3	153.1-163.5	1,781	197.5	188.4-207.0	1,799	134.5	128.3-140.9
2002	9,632	422.7	414.3-431.2	4,818	507.1	492.7-521.7	4,814	369.1	358.7-379.7	3,424	149.9	144.9-155.0	1,679	184.3	175.5-193.4	1,745	128.9	122.9-135.2
2003	9,600	419.1	410.8-427.6	4,699	494.0	479.8-508.4	4,901	374.4	363.9-385.1	3,428	149.4	144.4-154.5	1,658	179.8	171.2-188.8	1,770	130.7	124.6-137.0
2004	9,823	427.5	419.1-436.1	4,792	498.6	484.4-513.1	5,031	384.1	373.5-394.9	3,468	150.9	145.9-156.0	1,688	182.1	173.4-191.1	1,780	131.5	125.4-137.8
2005	9,936	432.4	423.9-441.0	4,953	513.4	499.1-528.1	4,983	382.9	372.2-393.8	3,322	143.9	139.0-148.9	1,635	176.0	167.5-184.8	1,687	124.3	118.4-130.4
2006	10,152	442.1	433.5-450.9	5,004	517.6	503.2-532.4	5,148	394.7	383.9-405.8	3,389	147.6	142.7-152.7	1,578	171.0	162.6-179.8	1,811	134.1	127.9-140.5
2007	10,268	444.3	435.6-453.0	5,162	526.7	512.2-541.5	5,106	391.7	380.9-402.7	3,373	145.6	140.7-150.6	1,604	171.6	163.2-180.3	1,769	129.8	123.8-136.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
All Invasive Malignant Tumors**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	10,497	448.1	439.5-456.8	5,265	527.2	512.9-541.9	5,232	396.9	386.1-407.9	3,175	136.0	131.3-140.9	1,644	172.7	164.4-181.4	1,531	111.6	106.0-117.4
2009	10,641	448.7	440.2-457.4	5,337	524.8	510.5-539.3	5,304	400.2	389.4-411.3	3,259	137.7	132.9-142.5	1,616	168.4	160.2-177.0	1,643	118.1	112.4-124.1
2010	10,522	437.2	428.8-445.7	5,246	505.6	491.7-519.7	5,276	394.4	383.7-405.4	3,321	138.4	133.6-143.2	1,612	165.1	157.0-173.5	1,709	121.7	115.9-127.7
2011	10,811	441.0	432.6-449.5	5,434	511.1	497.4-525.2	5,377	396.2	385.5-407.2	3,337	135.4	130.7-140.1	1,645	163.4	155.5-171.7	1,692	117.1	111.5-122.9
2012	10,480	418.5	410.4-426.7	5,138	470.8	457.8-484.1	5,342	387.3	376.8-398.0	3,483	139.0	134.4-143.8	1,719	167.0	159.0-175.2	1,764	121.4	115.7-127.3
2013	10,939	428.3	420.2-436.5	5,280	472.5	459.5-485.7	5,659	403.6	393.0-414.5	3,341	130.3	125.8-134.8	1,691	159.2	151.6-167.1	1,650	111.2	105.8-116.8
2014	10,825	415.6	407.7-423.7	5,176	449.4	436.9-462.1	5,649	397.2	386.7-407.9	3,355	128.4	124.1-132.9	1,669	152.9	145.5-160.6	1,686	112.6	107.2-118.2
2015	10,966	412.2	404.4-420.2	5,288	447.1	434.8-459.6	5,678	393.8	383.4-404.4	3,196	119.0	114.8-123.3	1,543	137.4	130.5-144.6	1,653	107.7	102.5-113.2
2016	11,317	418.6	410.8-426.6	5,495	455.1	442.9-467.6	5,822	399.0	388.6-409.7	3,345	122.1	117.9-126.3	1,628	141.8	134.8-149.0	1,717	109.4	104.2-114.8
2017	11,421	417.4	409.6-425.3	5,496	446.9	434.9-459.1	5,925	403.0	392.5-413.7	3,228	115.9	111.9-120.1	1,619	138.3	131.6-145.4	1,609	101.1	96.1-106.3
2018	11,387	413.8	406.1-421.6	5,399	431.6	419.9-443.5	5,988	407.9	397.3-418.7	2,981	105.4	101.6-109.3	1,477	123.5	117.2-130.1	1,504	93.6	88.8-98.6
2019	11,952	428.7	420.8-436.6	5,719	447.2	435.4-459.2	6,233	421.6	410.9-432.6	3,475	120.9	116.9-125.1	1,713	139.7	133.1-146.6	1,762	108.4	103.2-113.7
2020	10,121	360.9	353.7-368.2	4,855	378.2	367.4-389.3	5,266	354.3	344.5-364.4	2,943	102.1	98.3-105.9	1,528	123.1	116.9-129.6	1,415	87.6	83.0-92.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Oral cavity and pharynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	241	11.2	9.8-12.8	163	18.9	15.9-22.3	78	5.9	4.7-7.5	110	5.2	4.3-6.4	83	9.6	7.6-12.1	27	2.3	1.5-3.4
1977	222	10.2	8.8-11.6	146	16.4	13.7-19.4	76	5.8	4.6-7.4	97	4.8	3.8-5.8	71	8.8	6.7-11.3	26	2.1	1.4-3.1
1978	203	9.3	8.0-10.8	129	13.9	11.5-16.6	74	6.0	4.6-7.6	86	3.9	3.1-4.9	55	6.0	4.5-8.0	31	2.4	1.6-3.5
1979	215	10.3	8.9-11.9	145	17.1	14.3-20.3	70	5.5	4.2-7.0	84	4.1	3.2-5.1	62	7.3	5.5-9.5	22	1.8	1.1-2.8
1980	205	9.9	8.6-11.4	122	14.1	11.6-17.0	83	7.0	5.5-8.7	77	3.7	2.9-4.7	48	5.7	4.1-7.6	29	2.3	1.5-3.3
1981	224	10.7	9.3-12.3	158	18.4	15.5-21.6	66	5.5	4.2-7.1	79	3.6	2.8-4.5	49	5.3	3.8-7.1	30	2.4	1.6-3.4
1982	230	11.0	9.5-12.5	148	17.3	14.5-20.5	82	6.7	5.3-8.4	86	4.0	3.2-5.0	72	8.0	6.2-10.2	14	1.2	0.6-2.1
1983	246	11.5	10.0-13.0	183	20.7	17.7-24.1	63	4.9	3.8-6.4	75	3.3	2.6-4.2	56	5.9	4.4-7.7	19	1.4	0.8-2.2
1984	221	10.4	9.1-12.0	142	16.2	13.5-19.2	79	6.5	5.1-8.1	93	4.3	3.5-5.4	69	8.0	6.2-10.3	24	1.9	1.2-2.9
1985	249	11.7	10.2-13.3	160	18.2	15.4-21.4	89	7.1	5.7-8.8	81	3.7	2.9-4.7	55	6.2	4.6-8.2	26	2.1	1.3-3.1
1986	211	10.1	8.8-11.7	142	16.4	13.7-19.4	69	5.6	4.3-7.1	77	3.5	2.7-4.4	52	6.0	4.4-8.0	25	1.9	1.2-2.9
1987	219	10.5	9.2-12.1	141	15.9	13.3-18.8	78	6.6	5.2-8.3	63	3.0	2.3-3.9	38	4.6	3.2-6.3	25	1.9	1.2-2.9
1988	187	8.9	7.7-10.4	115	12.8	10.5-15.4	72	5.7	4.4-7.2	78	3.9	3.1-4.9	46	5.7	4.1-7.6	32	2.5	1.7-3.6
1989	212	10.4	9.0-11.9	143	16.6	13.9-19.6	69	5.9	4.5-7.5	65	3.1	2.4-4.0	42	4.8	3.4-6.5	23	2.0	1.2-3.0
1990	226	11.3	9.8-12.9	156	18.5	15.6-21.7	70	5.8	4.5-7.4	64	3.2	2.4-4.1	40	4.9	3.5-6.8	24	2.0	1.3-3.0
1991	233	11.5	10.0-13.1	144	16.9	14.2-20.0	89	7.5	6.0-9.3	51	2.5	1.8-3.3	38	4.3	3.0-5.9	13	1.1	0.6-1.9
1992	225	10.8	9.4-12.3	149	16.5	14.0-19.5	76	6.4	5.0-8.0	64	3.1	2.4-3.9	42	4.7	3.4-6.4	22	1.7	1.1-2.7
1993	218	10.5	9.2-12.1	137	15.8	13.3-18.8	81	6.8	5.4-8.5	67	3.3	2.5-4.2	51	5.9	4.4-7.8	16	1.2	0.7-1.9
1994	211	10.2	8.9-11.7	138	15.7	13.1-18.5	73	6.1	4.8-7.7	50	2.4	1.8-3.2	36	4.0	2.8-5.6	14	1.3	0.7-2.1
1995	183	8.9	7.6-10.3	107	12.1	9.9-14.6	76	6.2	4.8-7.7	56	2.7	2.0-3.5	36	4.2	2.9-5.8	20	1.6	0.9-2.4
1996	196	9.3	8.1-10.8	126	14.0	11.7-16.8	70	5.6	4.4-7.1	50	2.4	1.8-3.2	34	3.7	2.5-5.1	16	1.3	0.7-2.2
1997	193	9.0	7.8-10.4	121	13.1	10.8-15.6	72	5.7	4.5-7.2	43	2.0	1.5-2.8	31	3.5	2.4-5.0	12	1.0	0.5-1.7
1998	174	8.0	6.9-9.3	101	10.5	8.6-12.8	73	5.9	4.6-7.4	44	2.0	1.5-2.7	27	2.9	1.9-4.2	17	1.2	0.7-2.0
1999	208	9.4	8.2-10.8	137	14.3	12.0-16.9	71	5.7	4.4-7.2	47	2.2	1.6-2.9	31	3.5	2.3-4.9	16	1.3	0.7-2.1
2000	188	8.4	7.3-9.7	124	12.8	10.6-15.3	64	5.1	4.0-6.6	34	1.5	1.0-2.1	21	2.2	1.4-3.4	13	1.0	0.5-1.7
2001	191	8.5	7.3-9.7	128	13.1	11.0-15.6	63	4.9	3.8-6.3	52	2.3	1.7-3.0	39	4.0	2.9-5.5	13	1.0	0.5-1.7
2002	197	8.7	7.5-10.0	133	13.6	11.4-16.2	64	5.0	3.8-6.4	33	1.4	1.0-2.0	23	2.3	1.4-3.4	10	0.8	0.4-1.4
2003	200	8.7	7.5-10.0	127	12.7	10.6-15.2	73	5.6	4.4-7.0	44	1.9	1.4-2.6	34	3.4	2.4-4.8	10	0.7	0.4-1.4
2004	195	8.5	7.3-9.7	123	12.3	10.2-14.7	72	5.5	4.3-7.0	46	2.0	1.5-2.7	28	3.0	2.0-4.3	18	1.3	0.8-2.1
2005	195	8.4	7.3-9.7	138	13.5	11.3-16.0	57	4.5	3.4-5.8	41	1.8	1.3-2.4	27	2.7	1.8-4.0	14	1.0	0.5-1.7
2006	177	7.5	6.4-8.7	110	10.6	8.7-12.8	67	5.0	3.8-6.3	37	1.6	1.1-2.2	25	2.6	1.7-3.8	12	0.8	0.4-1.5
2007	210	8.9	7.7-10.2	135	13.1	11.0-15.6	75	5.6	4.4-7.0	40	1.7	1.2-2.3	28	2.7	1.8-4.0	12	0.9	0.4-1.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Oral cavity and pharynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	204	8.8	7.6-10.1	120	11.8	9.8-14.2	84	6.5	5.2-8.1	52	2.2	1.7-2.9	36	3.7	2.6-5.1	16	1.2	0.7-2.0
2009	222	9.3	8.1-10.6	153	14.2	12.0-16.7	69	5.3	4.1-6.8	48	2.0	1.5-2.7	34	3.3	2.3-4.6	14	1.1	0.6-1.8
2010	221	9.1	7.9-10.4	134	12.4	10.3-14.7	87	6.6	5.3-8.2	44	1.8	1.3-2.4	21	2.1	1.3-3.2	23	1.6	1.0-2.4
2011	227	9.2	8.0-10.5	153	13.6	11.5-16.0	74	5.7	4.4-7.1	46	1.8	1.3-2.5	29	2.6	1.7-3.8	17	1.2	0.7-2.0
2012	236	9.2	8.1-10.5	145	12.6	10.6-14.8	91	6.5	5.2-8.0	47	1.9	1.4-2.5	33	3.0	2.1-4.3	14	1.1	0.6-1.8
2013	222	8.6	7.5-9.8	155	13.4	11.3-15.7	67	4.7	3.6-6.0	54	2.0	1.5-2.7	35	2.9	2.0-4.1	19	1.3	0.8-2.0
2014	242	9.3	8.2-10.6	145	12.3	10.3-14.5	97	6.8	5.5-8.3	48	1.9	1.4-2.5	34	3.1	2.1-4.3	14	0.9	0.5-1.6
2015	246	9.2	8.0-10.4	154	12.4	10.5-14.6	92	6.4	5.2-7.9	51	1.9	1.4-2.5	39	3.3	2.3-4.5	12	0.8	0.4-1.5
2016	259	9.5	8.4-10.8	167	13.8	11.7-16.1	92	6.3	5.0-7.7	49	1.8	1.3-2.4	30	2.6	1.7-3.7	19	1.2	0.7-2.0
2017	260	9.4	8.3-10.7	163	13.0	11.0-15.2	97	6.5	5.2-8.0	56	1.9	1.5-2.5	39	3.1	2.2-4.3	17	1.0	0.6-1.6
2018	245	8.7	7.6-9.9	159	12.2	10.4-14.3	86	5.6	4.5-7.0	49	1.7	1.3-2.3	32	2.6	1.8-3.7	17	1.1	0.6-1.8
2019	258	9.2	8.1-10.5	156	11.8	10.0-13.9	102	7.1	5.8-8.7	55	1.9	1.4-2.5	40	3.1	2.2-4.2	15	0.9	0.5-1.5
2020	223	8.0	6.9-9.1	130	10.1	8.4-12.1	93	6.3	5.0-7.8	53	1.9	1.4-2.5	33	2.6	1.8-3.7	20	1.3	0.8-2.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Esophagus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	103	4.7	3.8-5.8	77	8.3	6.5-10.6	26	2.1	1.3-3.1	92	4.2	3.4-5.3	65	7.0	5.3-9.1	27	2.3	1.5-3.4
1977	104	4.9	3.9-5.9	74	8.2	6.3-10.4	30	2.4	1.6-3.4	100	4.4	3.6-5.4	71	7.7	5.9-9.8	29	2.2	1.4-3.2
1978	97	4.5	3.6-5.5	67	7.3	5.6-9.3	30	2.4	1.6-3.5	88	4.0	3.2-5.0	57	6.3	4.7-8.3	31	2.3	1.6-3.3
1979	104	4.7	3.8-5.7	67	7.1	5.4-9.1	37	2.9	2.0-4.0	103	4.6	3.8-5.7	70	7.3	5.6-9.3	33	2.6	1.8-3.7
1980	121	5.6	4.6-6.8	87	10.4	8.2-13.0	34	2.6	1.8-3.6	95	4.3	3.5-5.3	58	6.6	4.9-8.6	37	2.8	1.9-3.9
1981	94	4.5	3.6-5.6	68	7.8	5.9-10.0	26	2.2	1.4-3.3	100	4.7	3.8-5.8	77	8.8	6.8-11.1	23	1.8	1.1-2.8
1982	96	4.3	3.4-5.2	69	7.3	5.6-9.3	27	2.1	1.4-3.2	97	4.4	3.5-5.4	72	7.9	6.1-10.0	25	1.9	1.2-2.9
1983	109	5.0	4.1-6.0	66	7.3	5.6-9.4	43	3.3	2.4-4.6	87	3.9	3.1-4.9	61	6.7	5.0-8.7	26	2.0	1.3-3.0
1984	92	4.1	3.3-5.1	63	7.1	5.4-9.2	29	2.2	1.4-3.1	97	4.3	3.5-5.3	64	6.7	5.1-8.6	33	2.6	1.7-3.6
1985	96	4.3	3.5-5.3	63	6.6	5.1-8.6	33	2.7	1.8-3.8	68	3.1	2.4-3.9	47	4.9	3.6-6.6	21	1.7	1.0-2.6
1986	104	5.1	4.1-6.1	66	7.9	6.1-10.2	38	3.0	2.1-4.2	98	4.6	3.7-5.6	66	7.8	6.0-10.0	32	2.5	1.7-3.6
1987	99	4.6	3.8-5.7	59	6.8	5.2-8.9	40	3.1	2.2-4.3	78	3.7	2.9-4.6	54	6.3	4.7-8.3	24	1.9	1.2-2.8
1988	88	4.2	3.4-5.3	53	6.2	4.6-8.2	35	2.9	2.0-4.0	79	3.7	2.9-4.6	57	6.8	5.1-8.8	22	1.7	1.0-2.6
1989	105	5.0	4.1-6.1	77	9.2	7.2-11.5	28	2.2	1.5-3.3	77	3.6	2.8-4.6	57	6.5	4.9-8.5	20	1.6	1.0-2.5
1990	90	4.2	3.4-5.2	61	6.9	5.2-8.9	29	2.3	1.5-3.4	88	4.2	3.4-5.2	57	6.9	5.2-9.0	31	2.5	1.7-3.5
1991	92	4.4	3.5-5.4	63	7.4	5.7-9.6	29	2.2	1.5-3.2	71	3.2	2.5-4.1	52	5.8	4.3-7.7	19	1.5	0.9-2.4
1992	99	4.8	3.9-5.9	68	7.9	6.1-10.1	31	2.6	1.8-3.7	82	4.0	3.1-4.9	64	7.5	5.7-9.6	18	1.4	0.8-2.3
1993	80	3.7	2.9-4.6	59	6.5	4.9-8.4	21	1.7	1.1-2.7	67	3.2	2.5-4.1	48	5.6	4.1-7.4	19	1.6	0.9-2.5
1994	100	4.6	3.8-5.7	80	9.1	7.2-11.4	20	1.7	1.0-2.6	75	3.5	2.7-4.4	56	6.5	4.9-8.4	19	1.5	0.9-2.4
1995	91	4.3	3.5-5.3	51	5.9	4.4-7.8	40	3.1	2.2-4.3	70	3.3	2.5-4.1	50	5.8	4.3-7.7	20	1.6	1.0-2.5
1996	85	4.0	3.2-4.9	59	6.9	5.2-8.9	26	1.9	1.3-2.9	81	3.8	3.0-4.8	51	6.0	4.5-7.9	30	2.3	1.6-3.4
1997	82	3.8	3.0-4.7	58	6.7	5.1-8.7	24	1.8	1.2-2.8	63	2.9	2.3-3.8	45	5.2	3.8-7.0	18	1.3	0.8-2.1
1998	89	4.1	3.3-5.1	65	7.3	5.6-9.3	24	1.9	1.2-2.9	64	3.0	2.3-3.8	50	5.8	4.3-7.7	14	1.1	0.6-1.8
1999	88	4.0	3.2-4.9	60	6.6	5.0-8.5	28	2.1	1.4-3.1	67	3.0	2.3-3.8	49	5.3	3.9-7.0	18	1.3	0.8-2.1
2000	77	3.4	2.7-4.3	49	5.2	3.9-6.9	28	2.1	1.4-3.1	82	3.7	2.9-4.5	54	5.9	4.4-7.7	28	2.0	1.4-3.0
2001	98	4.3	3.5-5.3	68	7.3	5.7-9.3	30	2.2	1.5-3.2	61	2.7	2.1-3.4	50	5.4	4.0-7.1	11	0.8	0.4-1.5
2002	102	4.5	3.7-5.5	69	7.3	5.6-9.2	33	2.5	1.7-3.5	78	3.4	2.7-4.3	58	6.3	4.8-8.2	20	1.5	0.9-2.3
2003	102	4.4	3.6-5.4	71	7.3	5.7-9.2	31	2.3	1.5-3.2	70	3.0	2.4-3.9	50	5.1	3.8-6.8	20	1.5	0.9-2.3
2004	84	3.7	2.9-4.5	49	5.1	3.7-6.7	35	2.5	1.7-3.5	77	3.4	2.6-4.2	55	5.8	4.3-7.5	22	1.5	1.0-2.4
2005	107	4.7	3.8-5.7	64	6.7	5.2-8.6	43	3.2	2.3-4.3	71	3.0	2.4-3.8	46	4.9	3.6-6.6	25	1.8	1.1-2.6
2006	102	4.5	3.7-5.5	72	7.7	6.0-9.7	30	2.3	1.5-3.2	90	4.0	3.2-4.9	56	5.9	4.5-7.7	34	2.5	1.7-3.5
2007	67	2.9	2.2-3.7	45	4.5	3.2-6.0	22	1.7	1.0-2.6	71	3.1	2.4-4.0	48	5.1	3.8-6.8	23	1.8	1.1-2.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Esophagus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	97	4.1	3.3-5.1	74	7.6	5.9-9.5	23	1.7	1.1-2.6	61	2.6	2.0-3.3	48	4.9	3.6-6.5	13	1.0	0.5-1.7
2009	70	2.9	2.2-3.6	49	4.6	3.4-6.1	21	1.5	0.9-2.4	73	3.1	2.4-3.9	54	5.5	4.1-7.2	19	1.4	0.8-2.2
2010	72	2.9	2.3-3.7	54	5.2	3.9-6.8	18	1.2	0.7-2.0	49	2.0	1.5-2.6	34	3.3	2.3-4.6	15	1.1	0.6-1.8
2011	88	3.6	2.8-4.4	70	6.5	5.0-8.3	18	1.3	0.7-2.0	60	2.5	1.9-3.2	42	4.1	2.9-5.6	18	1.2	0.7-1.9
2012	76	3.0	2.3-3.7	52	4.7	3.5-6.2	24	1.6	1.0-2.4	47	1.8	1.3-2.4	37	3.4	2.4-4.7	10	0.7	0.3-1.3
2013	95	3.8	3.0-4.6	74	6.7	5.2-8.4	21	1.5	0.9-2.3	62	2.3	1.8-3.0	47	3.8	2.8-5.1	15	1.0	0.6-1.8
2014	94	3.6	2.9-4.4	67	5.9	4.6-7.6	27	1.7	1.1-2.5	68	2.6	2.0-3.3	48	4.4	3.3-5.9	20	1.3	0.8-2.1
2015	88	3.1	2.5-3.9	64	5.3	4.0-6.8	24	1.5	0.9-2.3	51	1.8	1.4-2.4	33	2.8	1.9-3.9	18	1.0	0.6-1.7
2016	86	3.1	2.4-3.8	66	5.3	4.1-6.8	20	1.3	0.8-2.1	64	2.3	1.8-3.0	47	4.0	2.9-5.4	17	1.0	0.6-1.7
2017	100	3.5	2.8-4.2	74	5.8	4.5-7.3	26	1.6	1.0-2.4	69	2.4	1.8-3.0	49	3.9	2.8-5.2	20	1.2	0.8-2.0
2018	98	3.4	2.7-4.1	78	6.1	4.8-7.7	20	1.1	0.7-1.8	62	2.1	1.6-2.8	53	4.2	3.2-5.6	9	0.5	0.2-1.1
2019	89	3.2	2.5-3.9	62	5.1	3.9-6.6	27	1.7	1.1-2.5	63	2.2	1.7-2.9	47	3.8	2.8-5.1	16	1.0	0.5-1.6
2020	87	2.9	2.3-3.6	58	4.3	3.2-5.6	29	1.7	1.1-2.5	63	2.1	1.6-2.8	47	3.7	2.7-5.0	16	0.9	0.5-1.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Stomach**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	296	14.6	12.9-16.4	172	21.0	17.7-24.6	124	10.4	8.6-12.4	220	10.8	9.3-12.4	133	16.2	13.4-19.5	87	7.3	5.8-9.0
1977	293	14.5	12.8-16.3	159	19.4	16.3-22.9	134	11.2	9.3-13.4	196	9.9	8.5-11.4	104	13.4	10.7-16.4	92	7.7	6.2-9.5
1978	265	13.0	11.5-14.7	155	19.1	16.1-22.6	110	9.0	7.4-11.0	226	10.9	9.5-12.5	128	15.5	12.8-18.6	98	8.1	6.5-9.9
1979	299	14.7	13.0-16.5	156	18.5	15.5-21.8	143	11.9	10.0-14.1	202	10.2	8.8-11.7	113	14.3	11.7-17.4	89	7.5	6.0-9.3
1980	300	14.3	12.7-16.0	174	20.5	17.5-24.0	126	10.2	8.5-12.3	233	11.2	9.8-12.8	129	15.6	12.9-18.6	104	8.3	6.7-10.1
1981	319	15.3	13.7-17.2	180	22.0	18.7-25.6	139	10.9	9.2-13.0	207	9.9	8.6-11.4	120	14.8	12.2-17.9	87	6.8	5.4-8.5
1982	276	13.0	11.5-14.7	151	18.6	15.6-22.0	125	9.9	8.2-11.8	220	10.6	9.2-12.1	136	17.5	14.5-20.8	84	6.5	5.2-8.1
1983	319	15.1	13.5-16.9	183	22.2	18.9-25.8	136	10.3	8.6-12.2	202	9.6	8.3-11.0	108	13.7	11.1-16.6	94	7.0	5.7-8.6
1984	278	12.9	11.4-14.6	159	18.4	15.6-21.7	119	9.2	7.6-11.0	188	8.8	7.6-10.2	103	12.4	10.0-15.1	85	6.4	5.1-8.0
1985	254	12.2	10.7-13.8	142	17.3	14.5-20.5	112	8.8	7.2-10.6	194	8.9	7.7-10.3	109	12.8	10.4-15.6	85	6.4	5.1-7.9
1986	274	12.8	11.3-14.5	146	17.0	14.2-20.0	128	10.1	8.4-12.0	155	7.3	6.2-8.6	94	11.9	9.6-14.7	61	4.5	3.5-5.9
1987	306	14.2	12.6-15.9	169	19.3	16.4-22.6	137	10.4	8.7-12.3	178	8.4	7.2-9.7	96	11.1	8.9-13.6	82	6.3	5.0-7.9
1988	271	12.8	11.3-14.4	163	19.8	16.8-23.2	108	8.5	6.9-10.3	149	6.9	5.9-8.2	81	9.3	7.3-11.6	68	5.5	4.2-7.0
1989	262	12.4	10.9-14.0	137	15.7	13.1-18.6	125	9.8	8.1-11.7	166	7.9	6.8-9.2	95	11.1	9.0-13.7	71	5.8	4.5-7.3
1990	316	15.2	13.5-16.9	187	23.0	19.7-26.7	129	10.2	8.5-12.2	175	8.5	7.3-9.9	95	11.5	9.3-14.1	80	6.4	5.0-8.0
1991	289	13.9	12.4-15.7	164	20.0	17.0-23.3	125	9.9	8.3-11.9	158	7.5	6.4-8.8	95	11.4	9.2-14.0	63	4.9	3.8-6.3
1992	286	13.4	11.9-15.1	163	19.4	16.5-22.7	123	9.3	7.7-11.2	203	9.7	8.4-11.2	114	14.1	11.5-17.0	89	7.0	5.6-8.7
1993	299	14.0	12.5-15.7	181	21.3	18.3-24.7	118	9.2	7.6-11.1	190	8.9	7.7-10.3	117	14.4	11.8-17.3	73	5.8	4.5-7.3
1994	321	15.2	13.6-17.0	179	20.8	17.8-24.1	142	11.3	9.5-13.3	191	9.0	7.8-10.4	108	12.5	10.2-15.1	83	6.5	5.2-8.1
1995	292	13.8	12.2-15.4	166	19.4	16.5-22.6	126	9.7	8.1-11.6	179	8.5	7.3-9.8	104	12.8	10.4-15.5	75	5.8	4.5-7.3
1996	293	13.8	12.2-15.5	157	18.3	15.5-21.5	136	10.5	8.8-12.4	176	8.3	7.1-9.6	82	9.4	7.5-11.7	94	7.2	5.8-8.8
1997	307	14.3	12.7-16.0	183	21.5	18.5-24.9	124	9.5	7.9-11.3	158	7.4	6.2-8.6	93	11.0	8.9-13.5	65	5.1	3.9-6.5
1998	318	14.6	13.1-16.4	176	20.3	17.3-23.5	142	11.1	9.3-13.1	148	6.8	5.8-8.0	79	9.3	7.3-11.6	69	5.2	4.1-6.7
1999	283	12.8	11.3-14.4	161	18.2	15.5-21.2	122	9.1	7.5-10.9	174	7.9	6.8-9.1	94	10.7	8.7-13.2	80	5.9	4.7-7.4
2000	297	13.2	11.7-14.8	165	18.6	15.9-21.7	132	9.8	8.2-11.6	142	6.3	5.3-7.4	63	7.1	5.4-9.1	79	5.8	4.6-7.2
2001	262	11.5	10.2-13.0	157	17.3	14.7-20.3	105	7.9	6.4-9.6	151	6.6	5.6-7.8	97	10.7	8.6-13.0	54	4.1	3.0-5.3
2002	275	12.0	10.6-13.5	154	16.5	14.0-19.4	121	9.0	7.5-10.8	134	5.8	4.9-6.9	79	8.8	7.0-11.0	55	4.1	3.1-5.3
2003	307	13.3	11.8-14.9	179	18.9	16.2-21.9	128	9.5	7.9-11.3	148	6.4	5.4-7.6	79	8.7	6.9-10.8	69	5.0	3.9-6.4
2004	275	11.9	10.5-13.4	163	17.1	14.5-19.9	112	8.3	6.8-10.0	154	6.6	5.6-7.8	77	8.2	6.5-10.3	77	5.7	4.5-7.2
2005	291	12.7	11.3-14.2	160	16.9	14.4-19.8	131	9.8	8.2-11.7	157	6.8	5.7-7.9	92	9.8	7.9-12.0	65	4.6	3.5-5.9
2006	273	11.9	10.5-13.4	152	15.7	13.3-18.4	121	8.9	7.4-10.7	145	6.4	5.4-7.5	71	7.7	6.0-9.7	74	5.6	4.4-7.0
2007	311	13.5	12.0-15.1	189	19.5	16.8-22.5	122	9.2	7.7-11.1	120	5.2	4.3-6.2	65	6.8	5.3-8.7	55	4.0	3.0-5.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Stomach**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	282	12.0	10.7-13.6	165	17.1	14.5-19.9	117	8.5	7.0-10.2	126	5.3	4.4-6.3	80	8.2	6.5-10.2	46	3.3	2.4-4.5
2009	299	12.5	11.2-14.1	190	19.0	16.3-21.9	109	8.0	6.5-9.6	153	6.4	5.5-7.6	92	9.3	7.5-11.5	61	4.5	3.4-5.8
2010	275	11.4	10.0-12.8	159	15.2	12.9-17.8	116	8.3	6.8-10.0	120	5.0	4.1-6.0	60	6.2	4.7-8.0	60	4.2	3.2-5.4
2011	271	11.0	9.7-12.4	167	16.1	13.7-18.8	104	7.3	5.9-8.9	129	5.3	4.4-6.3	75	7.4	5.8-9.4	54	3.8	2.8-5.0
2012	301	12.1	10.7-13.5	174	16.3	13.9-19.0	127	8.9	7.4-10.6	153	6.1	5.1-7.1	84	8.0	6.3-9.9	69	4.7	3.7-6.0
2013	331	13.1	11.7-14.6	182	17.1	14.7-19.8	149	10.2	8.6-12.1	137	5.4	4.5-6.4	81	7.7	6.1-9.6	56	3.9	2.9-5.1
2014	335	12.9	11.6-14.4	214	19.2	16.6-22.0	121	8.2	6.8-9.8	134	5.2	4.4-6.2	83	7.6	6.0-9.5	51	3.5	2.6-4.6
2015	318	11.8	10.5-13.1	178	14.6	12.5-17.0	140	9.5	8.0-11.3	134	5.0	4.2-6.0	73	6.6	5.1-8.3	61	4.0	3.0-5.2
2016	336	12.4	11.1-13.8	207	17.5	15.1-20.1	129	8.6	7.1-10.2	130	4.7	3.9-5.6	92	7.8	6.2-9.6	38	2.4	1.7-3.3
2017	368	13.5	12.1-14.9	205	17.1	14.8-19.7	163	10.7	9.1-12.5	143	5.1	4.3-6.1	85	7.1	5.6-8.8	58	3.6	2.7-4.7
2018	302	11.0	9.7-12.3	166	13.4	11.4-15.7	136	9.0	7.5-10.8	123	4.4	3.7-5.3	68	5.8	4.5-7.4	55	3.4	2.5-4.5
2019	333	12.2	10.9-13.6	194	15.8	13.6-18.3	139	9.4	7.9-11.2	138	5.1	4.2-6.0	81	6.9	5.5-8.6	57	3.8	2.8-4.9
2020	283	10.3	9.1-11.6	155	12.3	10.4-14.5	128	8.7	7.2-10.4	105	3.7	3.0-4.5	60	4.7	3.6-6.1	45	2.9	2.1-4.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Colorectal**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,454	69.6	65.9-73.4	709	83.6	77.1-90.5	745	60.6	56.2-65.3	770	38.2	35.4-41.1	401	51.1	45.8-56.8	369	30.4	27.3-33.8
1977	1,368	65.1	61.6-68.8	682	81.9	75.4-88.7	686	54.4	50.3-58.8	693	32.9	30.4-35.5	359	44.2	39.4-49.4	334	26.2	23.4-29.2
1978	1,380	65.8	62.2-69.4	667	81.4	75.0-88.2	713	56.4	52.2-60.8	708	34.3	31.7-37.0	349	43.6	38.8-48.7	359	28.7	25.7-31.9
1979	1,319	60.9	57.6-64.4	628	73.1	67.2-79.3	691	53.4	49.4-57.7	673	32.3	29.9-35.0	316	39.5	35.1-44.4	357	28.2	25.3-31.4
1980	1,396	65.9	62.4-69.5	715	84.8	78.4-91.5	681	53.7	49.6-58.0	674	32.3	29.8-34.9	320	39.2	34.9-44.0	354	27.7	24.9-30.9
1981	1,466	68.4	64.8-72.1	700	82.4	76.1-89.0	766	59.4	55.2-63.9	628	30.1	27.8-32.7	318	40.0	35.5-44.8	310	24.4	21.7-27.3
1982	1,556	72.4	68.8-76.2	741	88.1	81.6-95.0	815	63.0	58.7-67.6	671	31.4	29.1-34.0	334	40.3	35.9-45.1	337	26.1	23.4-29.2
1983	1,531	70.3	66.8-74.0	745	87.4	81.0-94.2	786	59.8	55.6-64.2	673	31.5	29.2-34.1	331	40.8	36.3-45.6	342	26.1	23.4-29.1
1984	1,414	66.0	62.6-69.6	679	79.0	73.0-85.5	735	57.3	53.1-61.7	701	33.1	30.7-35.7	344	42.4	37.9-47.3	357	27.5	24.7-30.6
1985	1,455	66.2	62.7-69.7	719	82.7	76.5-89.2	736	55.7	51.7-60.0	610	28.2	25.9-30.5	300	35.3	31.2-39.7	310	23.7	21.1-26.6
1986	1,507	69.5	66.0-73.2	763	90.8	84.3-97.8	744	56.8	52.7-61.1	633	29.6	27.3-32.0	319	38.6	34.3-43.3	314	24.1	21.5-27.0
1987	1,363	63.2	59.8-66.7	670	79.7	73.6-86.2	693	52.8	48.9-57.0	604	28.4	26.1-30.8	287	35.7	31.5-40.2	317	24.0	21.4-26.8
1988	1,342	62.8	59.4-66.3	639	75.9	69.9-82.2	703	54.2	50.2-58.4	579	27.3	25.1-29.6	299	37.7	33.4-42.4	280	21.5	19.0-24.3
1989	1,254	59.0	55.8-62.5	617	74.1	68.2-80.4	637	49.0	45.2-53.1	558	26.3	24.1-28.6	267	33.1	29.2-37.5	291	22.2	19.7-24.9
1990	1,347	63.1	59.7-66.6	629	75.3	69.3-81.6	718	55.2	51.2-59.5	550	26.0	23.8-28.3	271	33.4	29.4-37.8	279	21.5	19.0-24.3
1991	1,290	60.9	57.6-64.3	630	76.7	70.7-83.1	660	50.9	47.0-55.0	570	27.2	25.0-29.6	278	35.7	31.5-40.2	292	22.4	19.8-25.1
1992	1,198	56.7	53.5-60.0	554	67.2	61.6-73.1	644	50.3	46.5-54.4	511	24.1	22.1-26.3	240	29.5	25.8-33.5	271	20.7	18.3-23.4
1993	1,249	58.6	55.3-61.9	595	70.9	65.2-76.9	654	50.1	46.3-54.1	474	22.4	20.4-24.6	237	29.5	25.8-33.6	237	18.1	15.8-20.6
1994	1,301	61.0	57.7-64.4	612	73.1	67.3-79.2	689	53.2	49.3-57.4	512	24.0	21.9-26.1	252	30.9	27.1-35.0	260	19.6	17.2-22.1
1995	1,306	61.0	57.7-64.4	616	72.5	66.8-78.5	690	53.1	49.2-57.3	523	24.4	22.4-26.6	243	29.0	25.4-33.0	280	21.3	18.9-24.0
1996	1,240	58.1	54.9-61.4	585	68.8	63.3-74.7	655	50.8	46.9-54.9	465	21.7	19.7-23.7	243	29.6	25.9-33.6	222	16.7	14.5-19.1
1997	1,316	61.2	58.0-64.6	613	71.4	65.8-77.4	703	54.2	50.3-58.4	524	24.4	22.4-26.6	267	32.2	28.4-36.4	257	19.5	17.2-22.1
1998	1,189	54.8	51.7-58.0	549	63.2	58.0-68.8	640	49.2	45.4-53.1	446	20.5	18.7-22.5	215	25.3	22.0-28.9	231	17.5	15.3-20.0
1999	1,267	57.3	54.2-60.6	613	68.2	62.8-73.8	654	50.1	46.3-54.1	451	20.4	18.5-22.3	221	25.6	22.3-29.2	230	17.3	15.1-19.7
2000	1,296	57.7	54.6-60.9	640	70.5	65.1-76.2	656	49.0	45.3-52.9	440	19.6	17.8-21.5	221	24.9	21.7-28.4	219	16.1	14.0-18.4
2001	1,232	54.4	51.4-57.6	619	68.0	62.7-73.6	613	45.6	42.1-49.4	432	19.1	17.3-21.0	195	22.1	19.0-25.4	237	17.3	15.1-19.6
2002	1,177	51.6	48.7-54.7	539	57.6	52.8-62.7	638	46.9	43.3-50.7	410	17.9	16.2-19.7	210	23.1	20.1-26.5	200	14.3	12.3-16.4
2003	1,220	53.1	50.2-56.2	593	63.3	58.2-68.6	627	46.3	42.7-50.1	411	17.9	16.2-19.7	198	21.5	18.6-24.8	213	15.3	13.3-17.5
2004	1,223	53.0	50.1-56.1	558	59.4	54.5-64.6	665	48.7	45.1-52.6	365	15.8	14.2-17.5	184	19.8	17.0-22.9	181	12.9	11.1-15.0
2005	1,075	46.4	43.7-49.3	524	55.1	50.5-60.1	551	40.2	36.9-43.8	369	16.0	14.4-17.7	174	19.1	16.3-22.1	195	13.9	12.0-16.1
2006	1,157	50.3	47.5-53.3	552	58.7	53.8-63.8	605	44.9	41.3-48.6	394	17.1	15.4-18.9	196	21.4	18.5-24.6	198	14.2	12.2-16.3
2007	1,148	49.4	46.5-52.3	551	56.3	51.6-61.3	597	44.1	40.6-47.9	369	15.8	14.2-17.5	170	18.3	15.6-21.2	199	14.1	12.2-16.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Colorectal**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	1,111	47.1	44.3-50.0	553	55.5	50.9-60.4	558	40.3	37.0-43.9	323	13.7	12.2-15.2	159	16.7	14.2-19.5	164	11.5	9.8-13.5
2009	1,086	45.7	43.0-48.6	533	53.4	48.8-58.2	553	40.0	36.7-43.5	363	15.2	13.7-16.9	167	17.5	15.0-20.5	196	13.4	11.5-15.4
2010	1,092	45.2	42.5-47.9	552	53.9	49.4-58.7	540	38.8	35.6-42.3	369	15.3	13.8-17.0	173	18.1	15.4-21.0	196	13.6	11.7-15.7
2011	1,055	43.0	40.4-45.7	537	51.8	47.4-56.5	518	36.7	33.5-40.0	372	14.7	13.3-16.3	175	17.2	14.7-20.0	197	13.1	11.3-15.1
2012	996	39.9	37.4-42.5	524	49.5	45.2-54.0	472	32.8	29.9-36.0	362	14.5	13.0-16.1	190	18.7	16.1-21.6	172	11.4	9.7-13.3
2013	1,029	40.0	37.6-42.6	500	46.1	42.1-50.4	529	35.8	32.7-39.0	340	13.0	11.7-14.5	170	15.9	13.6-18.5	170	10.9	9.3-12.7
2014	1,056	40.4	38.0-43.0	546	47.3	43.3-51.6	510	35.1	32.1-38.4	321	12.2	10.9-13.6	168	15.4	13.1-17.9	153	10.0	8.4-11.8
2015	1,038	39.2	36.8-41.7	534	46.1	42.2-50.3	504	34.0	31.1-37.2	328	12.2	10.9-13.7	178	16.5	14.1-19.1	150	9.5	8.0-11.2
2016	1,042	39.0	36.6-41.5	547	46.5	42.6-50.7	495	33.2	30.3-36.4	351	12.7	11.3-14.1	177	15.2	13.0-17.6	174	10.8	9.2-12.6
2017	1,008	36.8	34.5-39.2	530	42.9	39.2-46.8	478	32.0	29.1-35.1	329	11.8	10.6-13.2	143	12.0	10.1-14.2	186	11.7	10.0-13.6
2018	1,021	36.8	34.5-39.2	526	42.2	38.6-46.0	495	32.3	29.4-35.3	271	9.6	8.5-10.8	130	10.6	8.8-12.7	141	8.7	7.3-10.4
2019	1,035	37.1	34.8-39.5	527	41.8	38.2-45.6	508	33.5	30.5-36.6	320	11.2	10.0-12.6	160	13.3	11.3-15.6	160	9.5	8.1-11.2
2020	895	32.1	29.9-34.3	467	36.9	33.6-40.6	428	27.9	25.2-30.8	300	10.3	9.2-11.6	155	12.4	10.5-14.6	145	8.8	7.4-10.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Colon excluding rectum**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	996	47.9	44.9-51.1	465	56.2	50.8-61.9	531	42.9	39.3-46.9	621	31.0	28.6-33.7	311	40.4	35.7-45.6	310	25.6	22.8-28.7
1977	927	44.7	41.7-47.7	446	54.6	49.3-60.2	481	38.6	35.1-42.3	568	26.9	24.7-29.3	289	35.4	31.2-40.1	279	21.9	19.3-24.7
1978	961	46.0	43.0-49.1	448	54.6	49.4-60.3	513	40.8	37.3-44.6	579	27.9	25.6-30.4	278	34.8	30.6-39.4	301	23.9	21.2-26.8
1979	927	43.0	40.2-45.9	420	48.6	43.9-53.8	507	39.5	36.0-43.2	563	27.0	24.7-29.4	255	31.6	27.6-35.9	308	24.4	21.7-27.4
1980	958	45.7	42.8-48.8	474	57.6	52.3-63.3	484	38.2	34.8-41.8	570	27.2	24.9-29.6	269	32.9	28.9-37.2	301	23.5	20.8-26.4
1981	1,030	48.2	45.2-51.3	476	56.6	51.4-62.2	554	43.0	39.4-46.8	536	25.8	23.7-28.2	272	34.3	30.1-38.8	264	20.9	18.4-23.6
1982	1,073	50.3	47.3-53.5	491	59.5	54.1-65.3	582	45.0	41.4-49.0	597	28.0	25.7-30.4	293	35.3	31.2-39.8	304	23.7	21.1-26.6
1983	1,060	49.2	46.3-52.4	498	59.3	54.0-65.0	562	43.1	39.6-46.9	570	26.6	24.4-28.9	283	34.8	30.7-39.3	287	21.8	19.3-24.5
1984	1,005	46.9	44.0-50.0	472	55.4	50.4-60.9	533	41.3	37.8-45.0	606	28.6	26.3-31.0	297	36.7	32.4-41.3	309	23.8	21.2-26.7
1985	1,048	47.9	45.0-51.0	500	58.2	53.0-63.8	548	41.8	38.3-45.5	530	24.5	22.4-26.7	256	30.1	26.4-34.2	274	21.1	18.6-23.8
1986	1,110	51.2	48.2-54.4	550	66.1	60.5-72.1	560	42.5	39.0-46.3	553	25.9	23.7-28.2	278	33.8	29.8-38.1	275	21.2	18.7-23.9
1987	972	44.9	42.1-47.9	456	54.5	49.5-60.0	516	39.0	35.6-42.5	528	24.8	22.7-27.1	252	31.4	27.5-35.6	276	20.9	18.5-23.5
1988	969	45.3	42.4-48.3	445	53.3	48.3-58.7	524	40.1	36.7-43.8	518	24.4	22.3-26.6	265	33.4	29.4-37.8	253	19.4	17.1-22.0
1989	920	43.5	40.7-46.4	431	52.4	47.4-57.7	489	37.7	34.4-41.3	498	23.5	21.5-25.7	236	29.6	25.9-33.8	262	20.0	17.6-22.6
1990	987	46.4	43.5-49.4	436	53.0	48.0-58.3	551	42.5	39.0-46.3	486	23.0	21.0-25.2	230	28.5	24.8-32.6	256	19.8	17.4-22.4
1991	958	45.2	42.4-48.2	447	54.9	49.8-60.3	511	39.2	35.8-42.8	497	23.8	21.7-26.0	242	31.4	27.5-35.7	255	19.5	17.1-22.1
1992	874	41.3	38.6-44.2	392	48.0	43.2-53.1	482	37.4	34.1-40.9	449	21.1	19.2-23.1	208	25.6	22.1-29.4	241	18.3	16.1-20.8
1993	898	42.1	39.3-44.9	404	48.6	43.9-53.7	494	37.7	34.4-41.3	413	19.5	17.7-21.5	202	25.3	21.9-29.2	211	16.1	14.0-18.5
1994	948	44.5	41.7-47.4	430	52.1	47.2-57.3	518	39.8	36.4-43.4	446	20.9	19.0-22.9	215	26.5	23.0-30.4	231	17.4	15.2-19.8
1995	948	44.3	41.5-47.2	429	51.3	46.5-56.5	519	39.7	36.3-43.3	463	21.6	19.7-23.7	215	25.8	22.4-29.6	248	18.8	16.5-21.4
1996	869	40.7	38.0-43.5	393	46.6	42.0-51.5	476	36.5	33.3-40.0	406	19.0	17.2-20.9	216	26.4	23.0-30.3	190	14.2	12.3-16.4
1997	962	44.7	42.0-47.7	443	51.9	47.1-57.1	519	39.9	36.5-43.5	456	21.3	19.3-23.3	226	27.4	23.9-31.3	230	17.5	15.3-19.9
1998	868	40.0	37.4-42.7	391	45.6	41.1-50.4	477	36.7	33.4-40.1	396	18.2	16.5-20.1	194	23.0	19.8-26.5	202	15.3	13.3-17.6
1999	900	40.7	38.1-43.5	420	47.1	42.7-51.9	480	36.7	33.5-40.1	388	17.5	15.8-19.4	183	21.3	18.3-24.7	205	15.4	13.4-17.7
2000	942	41.9	39.3-44.7	446	49.6	45.0-54.4	496	36.8	33.6-40.2	390	17.3	15.6-19.1	195	22.0	19.0-25.4	195	14.3	12.3-16.4
2001	874	38.6	36.1-41.2	426	46.9	42.5-51.6	448	33.0	30.0-36.2	377	16.6	15.0-18.4	163	18.6	15.8-21.7	214	15.5	13.5-17.8
2002	825	36.2	33.8-38.8	360	39.1	35.1-43.4	465	34.0	30.9-37.3	351	15.3	13.7-17.0	179	19.9	17.1-23.1	172	12.3	10.5-14.3
2003	890	38.8	36.2-41.4	419	45.2	40.9-49.7	471	34.6	31.5-37.9	364	15.8	14.2-17.5	175	19.0	16.3-22.0	189	13.5	11.6-15.6
2004	892	38.7	36.2-41.3	400	42.6	38.5-47.1	492	35.8	32.7-39.1	316	13.7	12.2-15.2	155	16.7	14.1-19.5	161	11.5	9.8-13.4
2005	789	34.1	31.7-36.5	369	39.0	35.1-43.2	420	30.5	27.7-33.7	333	14.4	12.9-16.0	152	16.7	14.2-19.6	181	12.9	11.1-15.0
2006	809	35.2	32.8-37.8	367	39.3	35.3-43.5	442	32.7	29.7-35.9	351	15.2	13.7-16.9	176	19.2	16.4-22.3	175	12.5	10.7-14.5
2007	812	35.0	32.6-37.5	379	39.3	35.4-43.5	433	31.8	28.8-35.0	328	14.0	12.5-15.6	153	16.3	13.8-19.2	175	12.4	10.6-14.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Colon excluding rectum**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	767	32.7	30.4-35.1	369	37.9	34.1-42.0	398	28.5	25.8-31.5	276	11.7	10.4-13.2	132	14.0	11.7-16.6	144	10.2	8.6-12.1
2009	771	32.7	30.4-35.1	358	36.8	33.0-40.8	413	29.5	26.7-32.6	319	13.4	11.9-14.9	150	15.7	13.3-18.5	169	11.4	9.8-13.4
2010	780	32.4	30.2-34.8	377	37.1	33.4-41.2	403	28.7	26.0-31.8	317	13.2	11.7-14.7	146	15.2	12.8-17.9	171	11.9	10.2-13.9
2011	749	30.7	28.6-33.1	384	37.6	33.9-41.6	365	25.9	23.2-28.7	315	12.5	11.1-13.9	144	14.1	11.9-16.7	171	11.4	9.7-13.3
2012	718	28.9	26.8-31.1	346	33.2	29.8-37.0	372	25.8	23.2-28.7	301	12.0	10.7-13.5	157	15.4	13.1-18.1	144	9.5	8.0-11.3
2013	720	28.1	26.0-30.2	332	31.0	27.7-34.6	388	26.1	23.5-28.9	295	11.3	10.1-12.7	146	13.7	11.6-16.2	149	9.6	8.1-11.3
2014	746	28.7	26.6-30.9	379	33.5	30.1-37.1	367	25.1	22.6-27.9	277	10.5	9.3-11.8	142	13.1	11.0-15.4	135	8.8	7.3-10.5
2015	716	27.1	25.1-29.2	352	31.1	27.8-34.6	364	24.4	21.9-27.1	268	9.9	8.7-11.2	145	13.3	11.2-15.7	123	7.7	6.4-9.2
2016	757	28.3	26.3-30.4	390	33.3	30.0-36.9	367	24.4	21.9-27.1	294	10.7	9.5-12.0	147	12.7	10.7-15.0	147	9.2	7.7-10.9
2017	683	24.9	23.0-26.9	344	28.1	25.2-31.4	339	22.3	19.9-24.9	273	9.7	8.6-11.0	115	9.6	7.9-11.6	158	9.8	8.3-11.5
2018	725	26.2	24.2-28.2	364	29.1	26.1-32.3	361	23.6	21.2-26.3	232	8.2	7.2-9.3	111	9.2	7.5-11.1	121	7.4	6.1-8.9
2019	742	26.4	24.5-28.5	354	28.2	25.2-31.3	388	25.1	22.6-27.9	261	9.3	8.2-10.5	120	10.2	8.4-12.2	141	8.5	7.1-10.1
2020	619	22.0	20.2-23.8	310	24.7	22.0-27.7	309	19.6	17.4-22.1	241	8.3	7.3-9.4	121	9.7	8.0-11.6	120	7.3	6.0-8.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Rectum & rectosigmoid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	458	21.7	19.7-23.9	244	27.4	23.9-31.4	214	17.7	15.3-20.3	149	7.1	6.0-8.4	90	10.7	8.4-13.3	59	4.8	3.6-6.3
1977	441	20.4	18.5-22.5	236	27.3	23.7-31.3	205	15.9	13.7-18.3	125	5.9	4.9-7.1	70	8.8	6.7-11.3	55	4.3	3.2-5.7
1978	419	19.8	17.9-21.8	219	26.8	23.2-30.8	200	15.5	13.4-17.9	129	6.3	5.3-7.6	71	8.8	6.7-11.2	58	4.8	3.6-6.2
1979	392	17.9	16.2-19.9	208	24.4	21.1-28.1	184	13.9	12.0-16.2	110	5.3	4.4-6.5	61	8.0	6.0-10.4	49	3.8	2.8-5.1
1980	438	20.2	18.3-22.2	241	27.1	23.7-31.0	197	15.5	13.4-17.9	104	5.1	4.1-6.2	51	6.4	4.7-8.5	53	4.3	3.2-5.6
1981	436	20.2	18.3-22.2	224	25.8	22.4-29.5	212	16.4	14.2-18.8	92	4.3	3.4-5.3	46	5.7	4.1-7.6	46	3.5	2.5-4.7
1982	483	22.1	20.2-24.2	250	28.5	25.0-32.5	233	18.0	15.7-20.5	74	3.5	2.7-4.4	41	5.1	3.6-6.9	33	2.4	1.7-3.5
1983	471	21.1	19.2-23.1	247	28.1	24.6-32.0	224	16.7	14.6-19.1	103	5.0	4.0-6.1	48	5.9	4.3-8.0	55	4.4	3.3-5.7
1984	409	19.1	17.2-21.1	207	23.6	20.4-27.2	202	16.0	13.8-18.5	95	4.5	3.7-5.6	47	5.7	4.2-7.7	48	3.7	2.7-4.9
1985	407	18.2	16.5-20.1	219	24.4	21.2-28.0	188	14.0	12.0-16.2	80	3.7	2.9-4.6	44	5.2	3.7-7.0	36	2.7	1.9-3.7
1986	397	18.3	16.5-20.2	213	24.7	21.4-28.4	184	14.2	12.2-16.5	80	3.7	2.9-4.6	41	4.9	3.4-6.7	39	3.0	2.1-4.1
1987	391	18.2	16.4-20.2	214	25.2	21.8-28.9	177	13.9	11.8-16.1	76	3.6	2.8-4.5	35	4.3	2.9-6.0	41	3.1	2.2-4.2
1988	373	17.5	15.8-19.4	194	22.5	19.4-26.0	179	14.1	12.0-16.3	61	2.9	2.2-3.7	34	4.3	3.0-6.1	27	2.1	1.4-3.1
1989	334	15.6	13.9-17.3	186	21.7	18.6-25.2	148	11.3	9.6-13.4	60	2.8	2.1-3.6	31	3.5	2.4-5.0	29	2.2	1.5-3.2
1990	360	16.7	15.0-18.6	193	22.3	19.2-25.8	167	12.7	10.8-14.8	64	3.0	2.3-3.8	41	4.9	3.5-6.7	23	1.7	1.1-2.6
1991	332	15.7	14.0-17.5	183	21.9	18.8-25.4	149	11.7	9.8-13.7	73	3.4	2.7-4.3	36	4.3	3.0-6.0	37	2.9	2.0-4.0
1992	324	15.3	13.7-17.1	162	19.2	16.3-22.5	162	12.9	11.0-15.1	62	3.1	2.3-3.9	32	3.9	2.6-5.5	30	2.4	1.6-3.5
1993	351	16.5	14.8-18.3	191	22.2	19.1-25.7	160	12.4	10.5-14.5	61	2.9	2.2-3.7	35	4.2	2.9-5.8	26	2.0	1.3-2.9
1994	353	16.5	14.8-18.4	182	21.0	18.0-24.3	171	13.4	11.5-15.6	66	3.1	2.4-3.9	37	4.4	3.1-6.1	29	2.2	1.5-3.2
1995	358	16.7	15.0-18.6	187	21.2	18.2-24.5	171	13.4	11.5-15.7	60	2.8	2.2-3.6	28	3.2	2.1-4.6	32	2.5	1.7-3.6
1996	371	17.4	15.7-19.3	192	22.2	19.2-25.6	179	14.3	12.2-16.5	59	2.7	2.1-3.5	27	3.2	2.1-4.6	32	2.5	1.7-3.5
1997	354	16.5	14.8-18.3	170	19.5	16.6-22.7	184	14.3	12.3-16.6	68	3.2	2.5-4.0	41	4.8	3.4-6.5	27	2.1	1.4-3.0
1998	321	14.8	13.2-16.5	158	17.6	14.9-20.6	163	12.5	10.6-14.6	50	2.3	1.7-3.0	21	2.3	1.4-3.5	29	2.2	1.5-3.2
1999	367	16.6	15.0-18.4	193	21.0	18.2-24.3	174	13.4	11.4-15.5	63	2.8	2.2-3.6	38	4.3	3.0-5.9	25	1.9	1.2-2.8
2000	354	15.8	14.2-17.5	194	21.0	18.1-24.1	160	12.2	10.4-14.2	50	2.2	1.7-3.0	26	2.8	1.8-4.1	24	1.8	1.1-2.7
2001	358	15.9	14.3-17.6	193	21.1	18.2-24.3	165	12.6	10.8-14.7	55	2.4	1.8-3.2	32	3.5	2.4-4.9	23	1.7	1.1-2.6
2002	352	15.4	13.8-17.1	179	18.5	15.9-21.4	173	12.9	11.0-15.0	59	2.6	2.0-3.3	31	3.2	2.1-4.5	28	2.0	1.3-2.9
2003	330	14.4	12.9-16.0	174	18.1	15.5-21.0	156	11.7	9.9-13.7	47	2.1	1.5-2.7	23	2.5	1.6-3.8	24	1.8	1.1-2.7
2004	331	14.3	12.8-16.0	158	16.7	14.2-19.6	173	12.9	11.1-15.0	49	2.1	1.6-2.8	29	3.1	2.1-4.5	20	1.4	0.9-2.2
2005	286	12.3	11.0-13.9	155	16.1	13.7-18.9	131	9.7	8.1-11.5	36	1.6	1.1-2.2	22	2.3	1.5-3.5	14	1.0	0.6-1.7
2006	348	15.1	13.5-16.8	185	19.4	16.7-22.4	163	12.2	10.4-14.3	43	1.9	1.4-2.5	20	2.2	1.3-3.4	23	1.7	1.1-2.5
2007	336	14.4	12.9-16.1	172	17.0	14.5-19.8	164	12.3	10.5-14.4	41	1.8	1.3-2.4	17	1.9	1.1-3.1	24	1.7	1.1-2.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Rectum & rectosigmoid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	344	14.4	12.9-16.0	184	17.6	15.1-20.4	160	11.8	10.0-13.8	47	2.0	1.4-2.6	27	2.7	1.8-3.9	20	1.3	0.8-2.1
2009	315	13.1	11.6-14.6	175	16.6	14.2-19.3	140	10.4	8.7-12.3	44	1.9	1.4-2.5	17	1.8	1.0-2.9	27	1.9	1.2-2.8
2010	312	12.7	11.4-14.3	175	16.7	14.3-19.5	137	10.1	8.4-12.0	52	2.1	1.6-2.8	27	2.9	1.9-4.2	25	1.7	1.1-2.5
2011	306	12.3	10.9-13.8	153	14.2	12.0-16.7	153	10.8	9.1-12.7	57	2.3	1.7-3.0	31	3.0	2.0-4.3	26	1.7	1.1-2.5
2012	278	11.0	9.8-12.4	178	16.2	13.9-18.8	100	7.0	5.7-8.5	61	2.4	1.8-3.1	33	3.2	2.2-4.6	28	1.9	1.3-2.8
2013	309	12.0	10.6-13.4	168	15.1	12.9-17.7	141	9.7	8.1-11.4	45	1.7	1.2-2.3	24	2.2	1.4-3.2	21	1.3	0.8-2.0
2014	310	11.7	10.5-13.2	167	13.8	11.8-16.2	143	10.0	8.4-11.9	44	1.7	1.2-2.2	26	2.3	1.5-3.4	18	1.2	0.7-1.9
2015	322	12.1	10.8-13.6	182	15.1	12.9-17.5	140	9.7	8.1-11.5	60	2.3	1.8-3.0	33	3.1	2.2-4.4	27	1.8	1.2-2.6
2016	285	10.7	9.5-12.1	157	13.2	11.2-15.5	128	8.8	7.3-10.6	57	2.0	1.5-2.6	30	2.4	1.6-3.5	27	1.6	1.0-2.4
2017	325	11.9	10.6-13.3	186	14.7	12.6-17.1	139	9.7	8.1-11.5	56	2.1	1.6-2.7	28	2.4	1.6-3.5	28	1.9	1.2-2.8
2018	296	10.6	9.4-12.0	162	13.1	11.1-15.3	134	8.7	7.2-10.3	39	1.4	1.0-1.9	19	1.5	0.9-2.3	20	1.3	0.8-2.1
2019	293	10.6	9.4-12.0	173	13.6	11.6-15.9	120	8.4	6.9-10.1	59	2.0	1.5-2.6	40	3.1	2.2-4.3	19	1.0	0.6-1.7
2020	276	10.1	8.9-11.4	157	12.2	10.4-14.4	119	8.3	6.8-10.0	59	2.0	1.5-2.7	34	2.7	1.9-3.9	25	1.5	0.9-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Liver / intrahepatic bile duct**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	88	4.3	3.4-5.4	45	5.2	3.7-7.1	43	3.6	2.6-5.0	102	4.9	4.0-6.0	53	5.6	4.1-7.5	49	4.2	3.0-5.6
1977	88	4.1	3.2-5.1	59	6.8	5.1-8.9	29	2.2	1.5-3.2	99	4.6	3.7-5.6	55	6.1	4.6-8.1	44	3.5	2.5-4.7
1978	86	4.1	3.3-5.1	49	5.6	4.1-7.6	37	3.0	2.1-4.2	90	4.2	3.3-5.2	58	6.5	4.9-8.6	32	2.6	1.7-3.7
1979	71	3.4	2.7-4.4	40	4.8	3.4-6.7	31	2.6	1.8-3.8	86	4.0	3.2-5.0	48	5.6	4.0-7.5	38	3.1	2.2-4.3
1980	73	3.5	2.7-4.4	42	4.9	3.5-6.7	31	2.6	1.7-3.7	92	4.2	3.4-5.2	51	5.8	4.2-7.7	41	3.3	2.3-4.5
1981	83	4.0	3.2-5.0	51	6.3	4.6-8.3	32	2.6	1.7-3.7	98	4.7	3.8-5.7	57	6.9	5.2-9.1	41	3.2	2.3-4.4
1982	80	3.6	2.8-4.5	53	6.0	4.4-7.9	27	2.1	1.4-3.2	93	4.3	3.4-5.2	60	7.0	5.3-9.1	33	2.5	1.7-3.6
1983	68	3.3	2.5-4.2	40	4.7	3.3-6.5	28	2.2	1.4-3.2	111	5.3	4.3-6.4	62	7.1	5.4-9.3	49	4.0	3.0-5.4
1984	90	4.4	3.5-5.4	61	7.3	5.5-9.4	29	2.5	1.6-3.6	90	4.3	3.5-5.4	47	5.4	3.9-7.2	43	3.5	2.5-4.8
1985	86	4.0	3.2-5.0	55	6.3	4.7-8.3	31	2.4	1.6-3.5	129	6.1	5.1-7.3	81	10.0	7.9-12.5	48	3.6	2.6-4.8
1986	106	5.0	4.1-6.0	76	8.7	6.8-11.0	30	2.5	1.7-3.7	112	5.2	4.3-6.3	67	7.6	5.9-9.8	45	3.6	2.6-4.9
1987	82	3.9	3.1-4.8	48	5.5	4.0-7.4	34	2.6	1.8-3.7	106	5.0	4.1-6.1	60	7.3	5.5-9.5	46	3.5	2.6-4.8
1988	93	4.5	3.6-5.5	59	6.8	5.1-8.8	34	2.8	1.9-3.9	108	5.0	4.1-6.1	64	7.4	5.6-9.5	44	3.4	2.4-4.6
1989	75	3.5	2.8-4.4	51	5.8	4.3-7.7	24	1.9	1.2-2.9	74	3.6	2.8-4.5	52	6.0	4.4-7.9	22	1.8	1.1-2.7
1990	101	4.8	3.9-5.8	61	7.2	5.5-9.4	40	3.1	2.2-4.3	126	5.8	4.8-6.9	67	8.0	6.2-10.3	59	4.5	3.4-5.9
1991	149	7.2	6.1-8.5	94	11.2	9.0-13.7	55	4.4	3.3-5.8	153	7.2	6.1-8.5	96	11.2	9.0-13.7	57	4.4	3.3-5.7
1992	125	5.9	4.9-7.1	82	9.2	7.3-11.4	43	3.4	2.4-4.6	140	6.7	5.6-7.9	83	9.7	7.7-12.1	57	4.6	3.5-6.0
1993	137	6.5	5.4-7.6	91	10.5	8.5-13.0	46	3.6	2.6-4.8	130	6.0	5.0-7.2	67	7.7	6.0-9.9	63	4.9	3.8-6.3
1994	138	6.6	5.6-7.8	94	10.6	8.5-13.0	44	3.5	2.6-4.8	117	5.6	4.6-6.7	79	9.5	7.4-11.8	38	3.0	2.1-4.2
1995	118	5.7	4.7-6.8	74	8.3	6.5-10.5	44	3.5	2.5-4.7	113	5.4	4.4-6.5	62	7.3	5.6-9.4	51	4.0	2.9-5.2
1996	148	7.0	5.9-8.2	104	11.6	9.5-14.1	44	3.5	2.6-4.8	114	5.3	4.4-6.4	76	8.6	6.8-10.8	38	2.9	2.0-4.0
1997	146	6.9	5.8-8.1	90	10.1	8.1-12.4	56	4.3	3.3-5.6	120	5.6	4.6-6.7	71	7.9	6.1-10.0	49	3.8	2.8-5.0
1998	195	9.0	7.8-10.4	119	12.8	10.6-15.3	76	6.0	4.7-7.5	117	5.4	4.5-6.5	76	8.4	6.6-10.5	41	3.1	2.2-4.3
1999	190	8.6	7.5-10.0	134	14.2	11.9-16.8	56	4.3	3.2-5.6	136	6.2	5.2-7.3	87	9.6	7.6-11.8	49	3.8	2.8-5.0
2000	180	8.1	6.9-9.3	122	13.0	10.8-15.5	58	4.4	3.3-5.7	136	6.1	5.1-7.2	95	10.4	8.4-12.7	41	3.1	2.2-4.2
2001	160	7.1	6.1-8.3	113	11.6	9.5-13.9	47	3.5	2.6-4.7	115	5.1	4.2-6.1	80	8.3	6.6-10.4	35	2.6	1.8-3.6
2002	172	7.5	6.5-8.8	121	12.2	10.1-14.6	51	3.8	2.8-5.0	129	5.6	4.7-6.7	84	8.6	6.9-10.7	45	3.2	2.3-4.3
2003	206	9.0	7.9-10.4	136	13.7	11.5-16.2	70	5.3	4.1-6.7	125	5.5	4.6-6.5	75	7.9	6.2-10.0	50	3.7	2.7-4.9
2004	194	8.4	7.3-9.7	142	14.2	11.9-16.7	52	3.9	2.9-5.1	136	5.9	5.0-7.0	82	8.3	6.6-10.3	54	4.1	3.1-5.3
2005	222	9.5	8.3-10.9	161	15.7	13.4-18.4	61	4.6	3.5-5.9	148	6.4	5.4-7.5	101	10.3	8.4-12.5	47	3.3	2.4-4.5
2006	204	8.8	7.6-10.1	146	14.3	12.0-16.8	58	4.4	3.4-5.8	136	5.9	5.0-7.0	88	9.1	7.3-11.3	48	3.6	2.6-4.8
2007	222	9.5	8.3-10.8	157	15.0	12.7-17.5	65	4.9	3.8-6.3	170	7.3	6.2-8.5	112	11.4	9.3-13.7	58	4.2	3.1-5.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Liver / intrahepatic bile duct**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	229	9.6	8.4-10.9	163	15.1	12.9-17.7	66	5.0	3.9-6.4	131	5.5	4.6-6.5	96	9.3	7.5-11.4	35	2.4	1.7-3.4
2009	235	9.7	8.5-11.0	166	15.3	13.0-17.8	69	5.0	3.9-6.3	142	6.0	5.1-7.1	97	9.4	7.6-11.5	45	3.4	2.4-4.5
2010	237	9.7	8.5-11.0	172	15.8	13.5-18.4	65	4.9	3.8-6.3	148	6.1	5.1-7.2	100	9.3	7.6-11.4	48	3.5	2.5-4.6
2011	242	9.6	8.5-11.0	166	15.0	12.7-17.5	76	5.4	4.2-6.8	139	5.5	4.6-6.6	88	8.2	6.5-10.1	51	3.6	2.7-4.8
2012	272	10.4	9.2-11.7	192	16.3	14.0-18.8	80	5.6	4.4-7.0	161	6.4	5.4-7.4	103	9.3	7.6-11.3	58	4.1	3.1-5.4
2013	254	9.7	8.5-11.0	184	15.6	13.4-18.1	70	4.7	3.7-6.0	167	6.5	5.5-7.6	109	9.5	7.7-11.5	58	4.0	3.1-5.3
2014	264	9.7	8.6-11.0	174	14.0	12.0-16.4	90	6.0	4.8-7.4	182	6.9	5.9-8.0	110	9.4	7.7-11.4	72	4.9	3.8-6.2
2015	284	10.4	9.2-11.7	198	16.3	14.0-18.8	86	5.7	4.5-7.1	155	5.7	4.8-6.6	97	8.1	6.5-9.9	58	3.7	2.8-4.9
2016	281	10.0	8.9-11.3	214	16.9	14.7-19.4	67	4.5	3.5-5.8	176	6.4	5.4-7.4	124	10.2	8.4-12.2	52	3.3	2.5-4.4
2017	302	10.6	9.4-11.9	214	16.6	14.4-19.0	88	5.5	4.4-6.8	175	6.2	5.3-7.3	110	8.8	7.2-10.6	65	4.3	3.3-5.5
2018	297	10.3	9.2-11.6	205	15.8	13.7-18.2	92	5.9	4.7-7.3	163	5.6	4.8-6.6	108	8.4	6.9-10.2	55	3.4	2.6-4.5
2019	273	9.5	8.4-10.8	193	14.9	12.8-17.2	80	5.3	4.2-6.7	182	6.2	5.3-7.2	120	9.3	7.7-11.2	62	3.7	2.9-4.9
2020	224	7.6	6.6-8.7	148	11.3	9.5-13.3	76	4.6	3.6-5.8	159	5.5	4.6-6.4	113	8.9	7.3-10.8	46	2.8	2.0-3.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Pancreas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	272	13.0	11.4-14.7	129	15.5	12.8-18.7	143	11.4	9.5-13.5	250	12.1	10.6-13.7	121	14.5	11.9-17.5	129	10.3	8.6-12.4
1977	269	12.9	11.4-14.6	143	17.2	14.4-20.5	126	10.1	8.3-12.1	226	10.8	9.4-12.3	119	14.4	11.7-17.4	107	8.5	6.9-10.3
1978	287	13.2	11.7-14.9	134	14.5	12.1-17.4	153	12.1	10.2-14.2	264	12.1	10.6-13.7	125	13.8	11.4-16.6	139	10.8	9.0-12.8
1979	254	11.8	10.4-13.4	131	14.5	12.0-17.3	123	9.7	8.1-11.7	254	11.9	10.4-13.5	120	13.5	11.1-16.3	134	10.6	8.8-12.6
1980	298	13.7	12.2-15.4	135	15.3	12.7-18.3	163	12.6	10.7-14.7	249	11.5	10.1-13.1	125	14.4	11.9-17.3	124	9.6	8.0-11.5
1981	288	13.6	12.0-15.3	149	17.3	14.6-20.5	139	11.0	9.2-13.1	269	12.7	11.2-14.4	129	15.4	12.7-18.5	140	10.8	9.0-12.8
1982	298	14.1	12.5-15.9	132	16.5	13.7-19.7	166	12.7	10.8-14.9	262	12.3	10.8-13.9	121	14.7	12.2-17.7	141	10.7	9.0-12.7
1983	253	11.5	10.1-13.0	126	14.2	11.7-17.0	127	9.5	7.9-11.3	245	11.2	9.8-12.7	129	14.3	11.9-17.1	116	8.9	7.3-10.7
1984	261	11.9	10.4-13.4	110	12.8	10.4-15.5	151	11.3	9.5-13.3	242	11.1	9.7-12.6	104	12.3	10.0-15.0	138	10.3	8.7-12.3
1985	301	13.8	12.2-15.5	143	16.3	13.7-19.3	158	11.8	10.0-13.9	243	11.1	9.7-12.6	114	13.1	10.7-15.8	129	9.6	8.0-11.5
1986	291	13.7	12.2-15.4	122	14.9	12.3-17.9	169	12.9	11.0-15.1	237	11.0	9.6-12.5	111	13.3	10.8-16.1	126	9.4	7.8-11.3
1987	278	13.0	11.5-14.7	124	14.5	12.0-17.4	154	11.7	9.9-13.7	237	11.0	9.6-12.5	100	11.3	9.2-13.9	137	10.4	8.7-12.4
1988	265	12.2	10.8-13.8	115	13.3	10.9-16.1	150	11.2	9.5-13.2	215	10.0	8.7-11.5	93	11.0	8.8-13.5	122	9.2	7.6-11.0
1989	258	12.2	10.7-13.8	116	13.8	11.3-16.6	142	11.1	9.3-13.1	224	10.7	9.3-12.2	110	13.3	10.9-16.1	114	8.8	7.3-10.7
1990	296	13.8	12.2-15.5	130	14.8	12.3-17.7	166	12.8	10.9-14.9	233	10.9	9.5-12.4	102	11.7	9.5-14.3	131	10.0	8.4-11.9
1991	239	11.2	9.8-12.8	106	13.3	10.8-16.2	133	10.2	8.5-12.2	219	10.2	8.9-11.7	95	11.4	9.2-14.0	124	9.4	7.8-11.3
1992	267	12.4	11.0-14.0	105	12.8	10.4-15.6	162	12.3	10.4-14.4	202	9.5	8.2-10.9	81	9.9	7.8-12.4	121	9.2	7.6-11.0
1993	265	12.4	11.0-14.0	133	15.7	13.1-18.7	132	10.2	8.5-12.2	242	11.3	9.9-12.8	123	14.7	12.2-17.6	119	9.1	7.5-10.9
1994	268	12.7	11.2-14.3	115	13.8	11.4-16.6	153	11.8	10.0-13.9	226	10.6	9.2-12.0	92	10.9	8.7-13.4	134	10.3	8.6-12.2
1995	266	12.5	11.0-14.1	101	12.1	9.8-14.8	165	12.7	10.8-14.8	214	10.0	8.7-11.4	88	10.5	8.4-12.9	126	9.4	7.8-11.2
1996	271	12.6	11.2-14.2	120	13.8	11.4-16.5	151	11.8	10.0-13.8	251	11.7	10.3-13.3	107	12.5	10.2-15.1	144	11.1	9.4-13.1
1997	267	12.4	11.0-14.0	119	13.5	11.2-16.2	148	11.3	9.5-13.3	199	9.3	8.0-10.7	85	9.9	7.9-12.3	114	8.7	7.2-10.5
1998	244	11.2	9.9-12.7	132	15.4	12.9-18.3	112	8.5	7.0-10.3	195	8.9	7.7-10.3	100	11.7	9.5-14.2	95	7.2	5.8-8.9
1999	261	11.8	10.4-13.3	126	14.0	11.7-16.7	135	10.0	8.4-11.9	216	9.8	8.5-11.2	115	12.9	10.6-15.4	101	7.4	6.1-9.1
2000	322	14.3	12.8-16.0	139	15.4	12.9-18.2	183	13.6	11.7-15.8	223	9.9	8.7-11.3	100	11.1	9.0-13.5	123	9.0	7.5-10.8
2001	264	11.7	10.3-13.2	122	13.4	11.1-16.0	142	10.5	8.8-12.4	232	10.2	8.9-11.6	111	12.5	10.3-15.1	121	8.8	7.3-10.6
2002	244	10.7	9.4-12.1	116	12.5	10.3-15.1	128	9.2	7.7-11.0	218	9.5	8.3-10.9	92	10.1	8.1-12.4	126	9.2	7.6-10.9
2003	284	12.4	11.0-13.9	133	14.7	12.3-17.4	151	10.7	9.1-12.6	220	9.6	8.4-11.0	110	11.9	9.8-14.4	110	7.8	6.4-9.4
2004	288	12.5	11.1-14.1	122	13.0	10.8-15.5	166	12.1	10.3-14.1	243	10.5	9.2-11.9	114	12.1	10.0-14.6	129	9.3	7.8-11.1
2005	260	11.2	9.8-12.6	120	12.4	10.3-14.9	140	10.2	8.5-12.0	224	9.6	8.4-11.0	109	11.5	9.4-13.9	115	8.3	6.8-10.0
2006	296	12.7	11.3-14.3	116	11.9	9.8-14.3	180	13.0	11.2-15.1	218	9.4	8.2-10.8	95	10.1	8.2-12.4	123	8.9	7.4-10.6
2007	275	11.8	10.4-13.3	136	14.3	11.9-16.9	139	9.9	8.3-11.7	232	9.9	8.7-11.3	99	10.3	8.4-12.6	133	9.6	8.0-11.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Pancreas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	303	13.0	11.6-14.6	156	16.2	13.7-18.9	147	10.5	8.9-12.4	236	10.0	8.8-11.4	121	12.8	10.6-15.3	115	8.0	6.6-9.7
2009	314	13.2	11.8-14.8	164	16.9	14.3-19.7	150	10.7	9.0-12.6	241	10.3	9.0-11.6	133	14.1	11.8-16.7	108	7.6	6.2-9.2
2010	316	13.4	11.9-14.9	163	16.4	14.0-19.2	153	11.2	9.4-13.1	247	10.5	9.2-11.9	127	13.0	10.8-15.5	120	8.5	7.1-10.3
2011	322	13.0	11.6-14.5	151	14.4	12.2-16.9	171	12.1	10.3-14.1	257	10.5	9.2-11.9	129	13.0	10.8-15.5	128	8.9	7.4-10.6
2012	316	12.6	11.2-14.1	141	13.4	11.2-15.8	175	12.0	10.3-14.0	258	10.1	8.9-11.5	118	11.2	9.2-13.4	140	9.5	7.9-11.2
2013	347	13.3	11.9-14.8	162	14.4	12.2-16.9	185	12.2	10.5-14.2	261	10.1	8.9-11.4	124	11.5	9.5-13.7	137	8.9	7.4-10.5
2014	344	13.2	11.8-14.7	169	15.0	12.8-17.5	175	11.7	10.0-13.6	264	10.1	8.9-11.4	137	12.3	10.3-14.6	127	8.3	6.9-10.0
2015	336	12.5	11.2-13.9	164	14.5	12.3-17.0	172	11.0	9.4-12.8	243	9.1	7.9-10.3	116	10.5	8.6-12.6	127	8.2	6.8-9.8
2016	353	12.7	11.4-14.2	169	14.5	12.3-16.9	184	11.5	9.8-13.3	273	9.9	8.7-11.2	137	11.7	9.8-13.9	136	8.4	7.0-10.0
2017	362	13.1	11.8-14.6	170	14.3	12.2-16.7	192	12.2	10.5-14.2	268	9.5	8.4-10.8	123	10.4	8.6-12.5	145	8.8	7.4-10.4
2018	376	13.2	11.8-14.6	160	12.9	10.9-15.1	216	13.1	11.4-15.1	220	7.7	6.7-8.8	92	7.6	6.1-9.3	128	7.7	6.4-9.2
2019	369	12.9	11.6-14.3	169	13.5	11.5-15.7	200	12.1	10.5-14.0	314	10.6	9.5-11.9	144	10.8	9.1-12.8	170	10.0	8.5-11.7
2020	386	13.1	11.8-14.5	180	13.7	11.7-15.9	206	12.3	10.7-14.2	264	8.9	7.9-10.1	127	9.7	8.1-11.6	137	8.2	6.9-9.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Larynx**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	125	5.5	4.5-6.6	104	10.9	8.8-13.3	21	1.7	1.0-2.6	37	1.8	1.2-2.5	28	3.6	2.3-5.3	9	0.7	0.3-1.4
1977	100	4.5	3.6-5.5	87	9.2	7.3-11.5	13	1.0	0.5-1.8	54	2.4	1.8-3.2	47	5.3	3.8-7.1	7	0.5	0.2-1.2
1978	125	5.5	4.6-6.7	98	10.7	8.6-13.1	27	2.0	1.3-3.0	52	2.2	1.7-3.0	46	4.7	3.4-6.3	6	0.5	0.2-1.0
1979	101	4.6	3.7-5.6	82	8.8	7.0-11.1	19	1.6	0.9-2.6	45	1.9	1.4-2.5	33	3.5	2.4-5.0	12	0.8	0.4-1.5
1980	129	5.9	4.9-7.0	102	11.2	9.0-13.7	27	2.1	1.4-3.1	38	1.8	1.3-2.5	30	3.6	2.4-5.2	8	0.6	0.3-1.3
1981	110	4.9	4.0-5.9	88	9.0	7.1-11.2	22	1.8	1.1-2.8	44	2.0	1.4-2.7	38	4.1	2.8-5.7	6	0.5	0.2-1.2
1982	105	4.7	3.9-5.8	80	8.8	6.9-11.1	25	2.1	1.3-3.1	45	1.9	1.4-2.5	35	3.6	2.4-5.1	10	0.7	0.3-1.4
1983	105	4.8	3.9-5.9	80	8.7	6.8-10.9	25	2.0	1.3-3.0	29	1.3	0.9-1.9	22	2.6	1.6-4.0	7	0.5	0.2-1.1
1984	132	6.0	5.0-7.1	104	11.1	9.0-13.5	28	2.4	1.5-3.5	39	1.7	1.2-2.3	31	3.2	2.2-4.7	8	0.6	0.2-1.2
1985	141	6.3	5.3-7.5	108	11.5	9.4-14.0	33	2.5	1.7-3.6	43	1.9	1.4-2.6	36	4.0	2.8-5.6	7	0.5	0.2-1.1
1986	101	4.8	3.9-5.9	76	8.6	6.7-10.9	25	2.2	1.4-3.3	27	1.3	0.8-1.9	22	2.8	1.7-4.2	5	0.4	0.1-1.0
1987	95	4.4	3.5-5.4	79	8.6	6.8-10.8	16	1.2	0.7-2.0	38	1.8	1.2-2.4	30	3.4	2.2-4.8	8	0.6	0.3-1.3
1988	108	5.1	4.1-6.2	84	9.2	7.3-11.5	24	2.1	1.3-3.1	31	1.4	0.9-2.0	24	2.6	1.6-3.9	7	0.6	0.2-1.3
1989	109	5.3	4.4-6.5	86	10.3	8.2-12.7	23	1.9	1.2-2.9	29	1.3	0.9-1.9	22	2.4	1.5-3.6	7	0.5	0.2-1.2
1990	130	6.1	5.1-7.3	108	12.1	9.8-14.6	22	1.9	1.2-3.0	29	1.4	0.9-2.0	26	3.1	2.0-4.6	3	0.3	0.1-0.8
1991	98	4.7	3.8-5.8	79	9.1	7.2-11.4	19	1.6	1.0-2.6	30	1.5	1.0-2.1	23	2.9	1.8-4.4	7	0.6	0.2-1.3
1992	124	6.0	5.0-7.2	102	11.5	9.4-14.0	22	1.9	1.2-2.8	40	1.9	1.4-2.6	28	3.5	2.3-5.1	12	0.9	0.5-1.6
1993	114	5.3	4.4-6.4	94	10.4	8.4-12.8	20	1.8	1.1-2.8	49	2.3	1.7-3.0	38	4.2	2.9-5.7	11	0.9	0.4-1.6
1994	94	4.4	3.6-5.4	77	8.7	6.8-10.9	17	1.4	0.8-2.2	39	1.8	1.3-2.5	32	3.6	2.4-5.0	7	0.6	0.2-1.2
1995	89	4.3	3.4-5.3	71	8.0	6.2-10.1	18	1.5	0.9-2.3	44	2.1	1.5-2.9	30	3.4	2.3-4.8	14	1.1	0.6-1.9
1996	104	4.9	4.0-5.9	81	9.0	7.1-11.2	23	1.9	1.2-2.8	29	1.3	0.9-1.9	22	2.6	1.6-4.0	7	0.6	0.2-1.2
1997	94	4.4	3.6-5.4	74	8.3	6.5-10.4	20	1.7	1.0-2.6	30	1.4	1.0-2.0	22	2.4	1.5-3.6	8	0.6	0.3-1.3
1998	103	4.7	3.9-5.8	82	8.8	7.0-10.9	21	1.7	1.0-2.6	20	0.9	0.6-1.5	17	1.9	1.1-3.0	3	0.2	0.1-0.7
1999	93	4.2	3.4-5.2	82	9.0	7.1-11.1	11	0.8	0.4-1.5	29	1.3	0.9-1.9	23	2.5	1.6-3.7	6	0.5	0.2-1.1
2000	66	3.0	2.3-3.8	49	5.2	3.8-6.9	17	1.3	0.8-2.1	31	1.4	0.9-2.0	28	3.1	2.1-4.5	3	0.2	0.1-0.7
2001	95	4.2	3.4-5.1	72	7.7	6.0-9.7	23	1.8	1.1-2.7	21	0.9	0.6-1.4	16	1.8	1.0-2.9	5	0.3	0.1-0.8
2002	80	3.5	2.8-4.4	66	7.0	5.4-8.9	14	1.0	0.6-1.7	31	1.4	0.9-1.9	23	2.5	1.6-3.8	8	0.6	0.2-1.2
2003	81	3.5	2.8-4.4	67	6.9	5.3-8.8	14	1.0	0.5-1.7	25	1.1	0.7-1.6	21	2.1	1.3-3.3	4	0.3	0.1-0.8
2004	81	3.5	2.8-4.4	66	6.5	5.0-8.3	15	1.2	0.6-1.9	24	1.0	0.7-1.5	20	2.0	1.2-3.2	4	0.3	0.1-0.8
2005	73	3.2	2.5-4.0	62	6.2	4.8-8.0	11	0.8	0.4-1.5	22	0.9	0.6-1.4	19	1.9	1.1-2.9	3	0.2	0.0-0.7
2006	84	3.6	2.8-4.4	67	6.8	5.2-8.6	17	1.3	0.8-2.1	30	1.3	0.9-1.8	23	2.4	1.5-3.6	7	0.5	0.2-1.1
2007	78	3.4	2.7-4.2	59	6.0	4.5-7.7	19	1.4	0.8-2.2	30	1.3	0.9-1.8	25	2.6	1.7-3.8	5	0.4	0.1-0.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Queens

Larynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	86	3.6	2.9-4.5	65	6.4	4.9-8.2	21	1.6	1.0-2.4	23	1.0	0.6-1.5	19	1.9	1.1-2.9	4	0.3	0.1-0.8
2009	80	3.4	2.7-4.2	63	6.2	4.8-8.0	17	1.3	0.7-2.0	24	1.0	0.6-1.5	20	2.0	1.2-3.1	4	0.3	0.1-0.7
2010	75	3.1	2.4-3.9	60	5.9	4.5-7.6	15	1.1	0.6-1.8	16	0.6	0.4-1.1	12	1.2	0.6-2.1	4	0.2	0.1-0.6
2011	67	2.6	2.0-3.3	58	5.3	4.0-6.9	9	0.6	0.3-1.2	23	0.9	0.6-1.4	16	1.6	0.9-2.6	7	0.5	0.2-1.0
2012	73	2.8	2.2-3.5	63	5.5	4.2-7.0	10	0.7	0.3-1.3	25	1.0	0.6-1.4	22	2.1	1.3-3.2	3	0.2	0.0-0.6
2013	73	2.8	2.2-3.5	59	5.3	4.0-6.9	14	0.9	0.5-1.6	26	1.1	0.7-1.6	23	2.2	1.4-3.3	3	0.2	0.0-0.7
2014	82	3.1	2.5-3.9	63	5.5	4.2-7.0	19	1.2	0.7-2.0	21	0.8	0.5-1.2	16	1.3	0.7-2.2	5	0.3	0.1-0.8
2015	66	2.5	1.9-3.2	57	4.8	3.6-6.3	9	0.6	0.3-1.2	23	0.8	0.5-1.3	17	1.4	0.8-2.2	6	0.4	0.1-0.9
2016	53	2.0	1.5-2.6	47	4.0	2.9-5.4	6	0.4	0.1-0.9	19	0.7	0.4-1.1	15	1.3	0.7-2.1	4	0.2	0.1-0.7
2017	64	2.2	1.7-2.9	52	4.2	3.1-5.5	12	0.8	0.4-1.5	14	0.5	0.2-0.8	11	0.8	0.4-1.6	3	0.2	0.0-0.6
2018	63	2.1	1.6-2.8	54	4.1	3.1-5.4	9	0.5	0.2-1.1	15	0.5	0.3-0.9	13	1.2	0.6-2.1	2	0.1	0.0-0.5
2019	67	2.3	1.8-2.9	56	4.4	3.3-5.8	11	0.7	0.3-1.3	25	0.8	0.5-1.2	22	1.6	1.0-2.5	3	0.2	0.0-0.5
2020	56	1.9	1.4-2.5	46	3.5	2.5-4.7	10	0.6	0.3-1.2	10	0.3	0.2-0.6	8	0.6	0.3-1.3	2	0.1	0.0-0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Lung and bronchus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,124	50.5	47.5-53.7	808	87.7	81.4-94.3	316	25.2	22.4-28.3	897	40.4	37.7-43.3	656	72.3	66.6-78.4	241	19.1	16.7-21.8
1977	1,122	51.1	48.1-54.3	784	87.3	81.0-94.1	338	27.0	24.1-30.2	959	43.8	40.9-46.7	695	77.9	71.9-84.3	264	21.3	18.8-24.2
1978	1,151	51.7	48.7-54.9	837	93.2	86.7-100.1	314	24.3	21.6-27.2	965	43.6	40.8-46.5	692	77.9	71.9-84.3	273	21.3	18.8-24.1
1979	1,206	54.7	51.5-57.9	828	92.9	86.4-99.8	378	29.8	26.7-33.0	953	43.5	40.7-46.4	675	76.4	70.4-82.7	278	22.3	19.6-25.1
1980	1,170	53.5	50.4-56.7	789	89.6	83.2-96.4	381	30.2	27.1-33.5	1,024	46.4	43.5-49.4	688	78.4	72.3-84.8	336	25.9	23.1-28.9
1981	1,135	51.8	48.8-55.0	735	84.1	77.8-90.7	400	31.0	28.0-34.4	928	42.3	39.6-45.2	636	72.3	66.6-78.4	292	23.1	20.4-26.0
1982	1,223	55.4	52.2-58.6	784	88.8	82.4-95.5	439	33.9	30.7-37.3	976	44.6	41.8-47.6	652	75.0	69.1-81.2	324	25.5	22.7-28.6
1983	1,216	54.8	51.7-58.1	784	88.7	82.4-95.4	432	33.1	29.9-36.4	935	42.3	39.6-45.2	591	68.1	62.5-74.0	344	26.0	23.2-29.0
1984	1,207	55.1	52.0-58.4	791	89.7	83.3-96.4	416	33.0	29.8-36.5	937	42.7	39.9-45.6	629	72.3	66.6-78.4	308	23.8	21.1-26.7
1985	1,271	57.7	54.5-61.0	818	92.8	86.3-99.6	453	34.8	31.6-38.3	967	43.5	40.8-46.4	622	71.5	65.8-77.6	345	26.2	23.4-29.2
1986	1,213	55.4	52.3-58.7	760	86.4	80.1-93.0	453	35.5	32.2-39.0	919	41.5	38.9-44.4	572	64.9	59.5-70.6	347	26.5	23.7-29.5
1987	1,274	58.1	54.9-61.5	750	85.6	79.4-92.1	524	40.8	37.3-44.6	992	45.6	42.7-48.5	620	72.3	66.6-78.5	372	29.0	26.0-32.2
1988	1,236	57.2	54.0-60.5	724	83.4	77.2-89.9	512	40.8	37.3-44.6	975	45.2	42.4-48.2	586	68.6	63.0-74.6	389	31.0	27.9-34.3
1989	1,253	58.4	55.2-61.8	735	85.5	79.3-92.1	518	41.7	38.1-45.6	981	45.4	42.5-48.3	604	71.8	66.0-78.0	377	29.5	26.5-32.7
1990	1,294	60.8	57.5-64.3	729	85.9	79.6-92.5	565	45.2	41.5-49.2	984	45.7	42.9-48.7	594	70.4	64.7-76.4	390	30.6	27.6-33.8
1991	1,216	57.2	54.0-60.6	709	82.4	76.3-88.8	507	40.7	37.2-44.5	990	46.7	43.8-49.7	555	66.1	60.5-71.9	435	34.7	31.5-38.2
1992	1,299	61.0	57.7-64.4	762	89.0	82.7-95.7	537	43.0	39.4-46.9	939	43.9	41.1-46.8	556	65.2	59.8-70.9	383	30.5	27.5-33.8
1993	1,218	57.4	54.2-60.8	704	82.1	76.0-88.4	514	41.3	37.8-45.1	960	45.2	42.4-48.2	554	65.6	60.2-71.4	406	32.4	29.3-35.8
1994	1,186	55.7	52.5-58.9	662	76.3	70.5-82.4	524	41.7	38.2-45.5	880	41.3	38.6-44.2	536	63.0	57.7-68.6	344	26.9	24.1-30.0
1995	1,168	54.5	51.4-57.7	659	76.2	70.4-82.3	509	40.1	36.7-43.8	885	41.3	38.7-44.2	490	57.1	52.1-62.4	395	31.2	28.2-34.5
1996	1,210	56.6	53.4-59.9	666	76.7	70.9-82.8	544	42.9	39.4-46.7	881	41.1	38.5-44.0	512	59.3	54.2-64.8	369	28.8	25.9-31.9
1997	1,273	59.4	56.2-62.8	720	82.3	76.4-88.6	553	44.0	40.4-47.9	879	41.0	38.3-43.8	530	61.1	56.0-66.6	349	27.2	24.4-30.2
1998	1,247	57.5	54.3-60.8	690	76.9	71.2-82.9	557	43.6	40.1-47.4	859	39.6	37.0-42.3	484	54.9	50.0-60.0	375	29.3	26.4-32.4
1999	1,135	51.3	48.4-54.4	604	67.3	62.0-72.9	531	40.8	37.4-44.5	871	39.4	36.8-42.1	476	53.6	48.8-58.6	395	30.3	27.4-33.5
2000	1,195	53.4	50.4-56.5	667	73.2	67.7-79.0	528	40.1	36.8-43.7	873	39.0	36.4-41.6	516	57.3	52.4-62.5	357	26.7	24.0-29.7
2001	1,192	52.8	49.8-55.9	646	70.6	65.2-76.3	546	40.9	37.5-44.5	838	37.1	34.7-39.7	469	51.6	47.0-56.6	369	27.6	24.8-30.6
2002	1,231	54.1	51.1-57.2	678	72.4	67.0-78.1	553	41.4	38.0-45.1	823	36.1	33.7-38.7	446	48.4	44.0-53.2	377	27.9	25.1-30.9
2003	1,130	49.5	46.6-52.5	619	65.9	60.8-71.4	511	38.2	35.0-41.7	793	34.7	32.4-37.2	444	48.2	43.7-52.9	349	25.9	23.2-28.8
2004	1,148	50.1	47.2-53.1	632	67.0	61.9-72.5	516	38.3	35.1-41.8	794	34.6	32.2-37.1	425	45.3	41.0-49.8	369	27.1	24.4-30.0
2005	1,154	50.4	47.5-53.4	637	67.7	62.5-73.2	517	38.9	35.6-42.4	748	32.6	30.3-35.0	433	46.2	41.9-50.8	315	23.5	20.9-26.3
2006	1,137	49.6	46.8-52.6	576	60.6	55.7-65.9	561	42.3	38.8-46.0	789	34.5	32.1-37.0	438	47.0	42.7-51.7	351	26.1	23.4-29.0
2007	1,179	51.6	48.7-54.7	633	66.6	61.5-72.1	546	41.3	37.8-44.9	765	33.1	30.8-35.6	415	44.5	40.2-49.0	350	25.6	23.0-28.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Lung and bronchus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	1,208	52.5	49.5-55.5	629	65.7	60.6-71.1	579	43.2	39.7-47.0	704	30.6	28.4-33.0	405	42.5	38.4-46.9	299	22.0	19.5-24.7
2009	1,181	50.3	47.5-53.3	627	64.3	59.3-69.6	554	40.6	37.2-44.2	720	30.4	28.2-32.8	383	39.7	35.7-43.9	337	24.1	21.6-26.9
2010	1,164	48.9	46.1-51.8	604	60.5	55.7-65.7	560	40.9	37.5-44.5	729	30.7	28.5-33.1	392	40.2	36.2-44.4	337	24.4	21.8-27.2
2011	1,195	49.5	46.7-52.4	630	62.0	57.2-67.1	565	40.5	37.2-44.1	734	30.0	27.9-32.3	409	40.6	36.7-44.8	325	22.1	19.8-24.8
2012	1,156	46.4	43.8-49.2	601	57.6	53.0-62.5	555	39.2	35.9-42.6	730	29.5	27.3-31.7	420	40.9	37.0-45.1	310	21.5	19.1-24.1
2013	1,155	45.2	42.6-47.9	594	54.5	50.1-59.2	561	38.8	35.6-42.2	734	28.8	26.7-31.0	420	39.7	35.9-43.7	314	21.3	19.0-23.9
2014	1,116	42.4	39.9-45.0	564	50.3	46.1-54.7	552	36.9	33.8-40.2	676	25.6	23.7-27.7	377	34.2	30.7-37.9	299	19.8	17.5-22.2
2015	1,184	43.9	41.4-46.5	640	55.5	51.2-60.1	544	35.4	32.4-38.6	647	23.9	22.0-25.8	333	28.9	25.8-32.3	314	20.2	18.0-22.6
2016	1,183	43.3	40.9-45.9	604	51.3	47.2-55.7	579	37.8	34.7-41.1	616	22.4	20.6-24.3	319	27.6	24.6-30.9	297	18.5	16.4-20.8
2017	1,232	44.2	41.7-46.7	649	54.3	50.1-58.8	583	36.8	33.8-40.0	586	21.1	19.4-22.9	328	28.1	25.1-31.4	258	16.1	14.1-18.2
2018	1,231	43.5	41.0-46.0	636	51.7	47.7-56.0	595	37.9	34.8-41.1	574	20.0	18.4-21.7	333	27.7	24.8-31.0	241	14.5	12.7-16.6
2019	1,293	44.8	42.4-47.4	631	50.0	46.1-54.2	662	40.7	37.6-44.0	593	20.3	18.7-22.1	320	26.0	23.2-29.1	273	16.3	14.4-18.4
2020	1,011	34.1	32.0-36.4	550	43.1	39.5-47.0	461	27.4	24.9-30.1	497	17.0	15.5-18.6	288	23.0	20.4-25.9	209	12.5	10.8-14.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Melanoma of the skin**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	108	5.4	4.4-6.6	57	6.7	5.0-8.8	51	4.6	3.3-6.1	44	2.1	1.5-2.9	22	2.5	1.5-3.9	22	1.9	1.2-3.0
1977	118	5.5	4.5-6.6	37	4.0	2.7-5.6	81	6.7	5.3-8.4	42	1.9	1.4-2.6	22	2.3	1.4-3.5	20	1.6	1.0-2.6
1978	114	5.6	4.6-6.8	47	5.4	3.9-7.3	67	5.7	4.4-7.3	33	1.6	1.1-2.3	18	2.2	1.3-3.6	15	1.3	0.7-2.2
1979	135	6.5	5.4-7.8	63	7.0	5.3-9.1	72	6.3	4.8-8.0	44	2.0	1.5-2.8	22	2.4	1.5-3.7	22	1.7	1.1-2.7
1980	123	6.1	5.1-7.4	56	6.3	4.7-8.3	67	6.0	4.6-7.8	47	2.3	1.7-3.1	19	2.3	1.3-3.6	28	2.3	1.5-3.5
1981	130	6.2	5.2-7.4	65	7.2	5.5-9.2	65	5.4	4.2-7.0	63	3.2	2.4-4.2	38	4.4	3.1-6.2	25	2.3	1.5-3.4
1982	127	6.2	5.1-7.4	71	8.1	6.3-10.3	56	4.7	3.5-6.2	43	2.1	1.5-2.8	18	2.1	1.2-3.4	25	2.0	1.3-3.1
1983	160	7.5	6.4-8.9	78	8.7	6.8-11.0	82	6.8	5.3-8.5	44	2.1	1.5-2.8	32	3.8	2.6-5.5	12	0.9	0.5-1.7
1984	146	7.0	5.9-8.3	75	8.4	6.6-10.7	71	6.2	4.8-7.8	52	2.3	1.7-3.1	28	2.9	1.9-4.2	24	1.8	1.1-2.7
1985	126	6.0	5.0-7.2	69	7.5	5.8-9.6	57	5.1	3.8-6.7	56	2.7	2.0-3.6	33	4.2	2.9-6.0	23	1.8	1.1-2.8
1986	123	6.0	5.0-7.2	58	6.7	5.0-8.7	65	5.6	4.3-7.2	51	2.5	1.8-3.2	23	2.6	1.6-4.0	28	2.4	1.6-3.5
1987	157	7.5	6.4-8.8	85	9.9	7.8-12.3	72	5.9	4.6-7.4	50	2.4	1.8-3.2	28	3.2	2.1-4.7	22	1.8	1.1-2.7
1988	138	6.5	5.5-7.8	69	8.2	6.3-10.4	69	5.6	4.3-7.1	52	2.5	1.8-3.3	30	3.5	2.3-5.0	22	1.8	1.1-2.8
1989	137	6.6	5.6-7.9	69	7.8	6.0-9.9	68	5.7	4.4-7.3	38	1.9	1.3-2.6	23	2.7	1.7-4.0	15	1.4	0.8-2.4
1990	143	7.0	5.9-8.2	72	8.3	6.5-10.5	71	6.1	4.7-7.8	43	2.1	1.5-2.8	19	2.4	1.4-3.8	24	2.0	1.3-3.0
1991	129	6.0	5.0-7.2	67	7.5	5.8-9.7	62	5.0	3.8-6.4	37	1.8	1.2-2.4	13	1.7	0.9-2.9	24	1.9	1.2-2.9
1992	129	6.1	5.1-7.3	67	7.7	5.9-9.8	62	5.0	3.8-6.5	41	1.9	1.3-2.6	25	2.6	1.7-3.9	16	1.3	0.7-2.1
1993	105	5.0	4.1-6.0	57	6.7	5.1-8.7	48	3.8	2.8-5.0	31	1.4	1.0-2.0	17	2.2	1.2-3.5	14	1.0	0.6-1.8
1994	145	6.9	5.8-8.1	77	8.8	6.9-11.1	68	5.6	4.3-7.1	27	1.2	0.8-1.8	20	2.1	1.3-3.3	7	0.6	0.2-1.2
1995	142	6.6	5.5-7.8	75	8.2	6.4-10.3	67	5.4	4.2-6.9	32	1.5	1.1-2.2	17	1.9	1.1-3.1	15	1.3	0.7-2.1
1996	145	6.8	5.7-8.0	70	8.0	6.2-10.1	75	6.0	4.7-7.5	48	2.2	1.6-2.9	23	2.4	1.5-3.7	25	2.0	1.3-3.0
1997	152	7.0	5.9-8.2	86	9.8	7.8-12.2	66	5.3	4.1-6.7	33	1.5	1.1-2.2	19	2.0	1.2-3.2	14	1.2	0.6-2.0
1998	144	6.6	5.5-7.8	78	8.7	6.9-10.9	66	5.3	4.1-6.8	39	1.8	1.3-2.4	21	2.5	1.5-3.8	18	1.4	0.8-2.2
1999	134	6.1	5.1-7.2	61	6.7	5.1-8.7	73	5.8	4.6-7.4	45	2.1	1.5-2.8	26	2.9	1.9-4.3	19	1.5	0.9-2.3
2000	179	7.9	6.8-9.1	81	8.9	7.0-11.1	98	7.7	6.2-9.4	33	1.5	1.0-2.1	20	2.3	1.4-3.5	13	1.0	0.5-1.7
2001	186	8.2	7.0-9.4	95	10.0	8.0-12.2	91	7.0	5.6-8.6	34	1.5	1.0-2.1	16	1.8	1.0-3.0	18	1.4	0.8-2.2
2002	168	7.4	6.3-8.6	90	9.5	7.6-11.7	78	6.1	4.8-7.6	29	1.3	0.9-1.8	16	1.7	1.0-2.8	13	1.0	0.5-1.7
2003	146	6.4	5.4-7.5	72	7.9	6.2-10.0	74	5.6	4.4-7.1	29	1.3	0.8-1.8	9	1.0	0.5-2.0	20	1.5	0.9-2.4
2004	165	7.2	6.1-8.4	96	10.2	8.2-12.5	69	5.4	4.2-6.9	37	1.6	1.1-2.2	24	2.7	1.7-4.0	13	1.0	0.5-1.7
2005	217	9.5	8.3-10.9	112	11.6	9.6-14.0	105	8.1	6.6-9.8	33	1.4	1.0-2.0	21	2.2	1.4-3.4	12	0.9	0.5-1.6
2006	235	10.2	8.9-11.6	119	12.2	10.1-14.7	116	9.0	7.4-10.8	23	1.0	0.7-1.6	12	1.3	0.7-2.3	11	0.9	0.4-1.6
2007	204	8.9	7.7-10.2	93	9.6	7.7-11.8	111	8.4	6.9-10.2	28	1.2	0.8-1.8	16	1.7	1.0-2.8	12	0.9	0.5-1.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Melanoma of the skin**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	219	9.5	8.3-10.8	111	11.3	9.3-13.7	108	8.2	6.7-10.0	27	1.2	0.8-1.7	11	1.2	0.6-2.1	16	1.2	0.7-2.0
2009	224	9.4	8.2-10.7	114	11.2	9.2-13.5	110	8.1	6.7-9.8	33	1.5	1.0-2.1	20	2.1	1.3-3.2	13	1.0	0.5-1.8
2010	197	8.2	7.1-9.5	106	10.3	8.4-12.5	91	6.7	5.4-8.3	35	1.5	1.0-2.1	16	1.6	0.9-2.5	19	1.4	0.8-2.2
2011	221	9.0	7.9-10.3	114	10.7	8.8-12.9	107	7.9	6.5-9.6	35	1.4	1.0-2.0	16	1.7	0.9-2.7	19	1.3	0.8-2.1
2012	221	8.8	7.7-10.1	121	11.5	9.5-13.7	100	7.3	5.9-8.9	35	1.4	1.0-1.9	19	1.8	1.1-2.8	16	1.1	0.6-1.9
2013	230	9.0	7.9-10.3	114	10.5	8.6-12.7	116	8.2	6.7-9.9	33	1.2	0.8-1.7	22	2.0	1.2-3.0	11	0.7	0.3-1.2
2014	204	8.0	6.9-9.2	103	9.7	7.9-11.9	101	7.0	5.7-8.6	40	1.5	1.1-2.1	20	1.9	1.1-2.9	20	1.2	0.8-2.0
2015	215	8.2	7.1-9.4	116	10.6	8.7-12.7	99	6.7	5.4-8.3	19	0.7	0.4-1.1	7	0.6	0.2-1.3	12	0.8	0.4-1.5
2016	211	8.0	6.9-9.1	120	10.5	8.6-12.6	91	6.2	5.0-7.7	30	1.2	0.8-1.7	16	1.4	0.8-2.2	14	1.0	0.5-1.7
2017	197	7.3	6.3-8.5	114	9.7	7.9-11.6	83	5.7	4.5-7.1	30	1.1	0.8-1.6	21	1.9	1.2-2.9	9	0.6	0.3-1.2
2018	193	7.0	6.0-8.0	105	8.4	6.8-10.2	88	5.9	4.7-7.4	20	0.7	0.4-1.1	13	1.1	0.6-1.8	7	0.4	0.2-0.9
2019	217	7.8	6.8-8.9	112	9.2	7.5-11.1	105	7.0	5.7-8.6	23	0.8	0.5-1.3	14	1.2	0.6-2.0	9	0.6	0.3-1.2
2020	174	6.2	5.3-7.2	104	8.6	7.0-10.5	70	4.6	3.6-5.9	22	0.8	0.5-1.2	11	1.0	0.5-1.7	11	0.7	0.3-1.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Female breast**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,170	101.2	95.2-107.5	457	38.3	34.7-42.1
1977	1,150	100.1	94.1-106.3	478	39.7	36.1-43.6
1978	1,168	100.1	94.2-106.3	438	36.4	33.0-40.1
1979	1,153	98.4	92.6-104.6	408	34.7	31.3-38.4
1980	1,184	101.5	95.5-107.7	454	37.1	33.6-40.8
1981	1,179	102.5	96.4-108.8	443	36.8	33.3-40.5
1982	1,205	103.1	97.1-109.4	443	36.3	32.9-40.0
1983	1,176	98.3	92.5-104.3	413	33.6	30.4-37.2
1984	1,310	111.1	104.9-117.5	438	36.2	32.8-39.9
1985	1,310	109.3	103.3-115.7	418	34.0	30.7-37.6
1986	1,280	109.1	103.0-115.5	425	34.6	31.3-38.2
1987	1,386	117.9	111.5-124.5	446	37.2	33.7-40.9
1988	1,267	108.9	102.8-115.3	459	37.7	34.2-41.4
1989	1,239	105.7	99.7-111.9	448	37.3	33.8-41.0
1990	1,401	120.7	114.3-127.4	393	33.0	29.8-36.5
1991	1,343	115.1	108.9-121.6	406	33.5	30.2-37.0
1992	1,395	119.8	113.5-126.4	400	33.4	30.2-37.0
1993	1,369	116.4	110.2-122.9	415	33.9	30.7-37.4
1994	1,484	126.6	120.1-133.3	377	31.3	28.1-34.6
1995	1,444	121.7	115.4-128.2	387	32.0	28.8-35.4
1996	1,528	128.4	121.9-135.0	396	32.3	29.1-35.7
1997	1,443	119.9	113.7-126.3	366	29.4	26.4-32.6
1998	1,502	123.2	117.1-129.7	346	27.5	24.6-30.5
1999	1,525	123.0	116.9-129.4	320	25.0	22.3-27.9
2000	1,450	115.7	109.8-121.8	336	25.7	23.0-28.6
2001	1,369	108.0	102.3-113.9	353	27.1	24.3-30.1
2002	1,333	104.6	99.1-110.5	307	22.8	20.3-25.6
2003	1,386	108.4	102.7-114.3	295	22.2	19.7-24.9
2004	1,398	109.0	103.3-114.9	326	24.6	22.0-27.4
2005	1,377	108.4	102.7-114.3	319	23.9	21.3-26.7
2006	1,423	111.7	105.9-117.7	333	24.9	22.2-27.7
2007	1,493	116.5	110.6-122.6	292	21.8	19.3-24.5
2008	1,467	113.2	107.5-119.3	241	18.0	15.7-20.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Female breast**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	1,519	116.2	110.4-122.3	260	19.1	16.9-21.7
2010	1,522	115.3	109.5-121.4	236	17.0	14.8-19.3
2011	1,519	114.2	108.4-120.2	251	17.7	15.6-20.1
2012	1,457	106.8	101.3-112.5	281	19.4	17.1-21.8
2013	1,612	117.5	111.7-123.5	260	17.6	15.5-20.0
2014	1,642	117.9	112.1-123.8	258	17.5	15.4-19.9
2015	1,621	114.3	108.6-120.1	262	17.4	15.3-19.8
2016	1,734	120.3	114.6-126.3	270	17.5	15.4-19.8
2017	1,798	125.3	119.4-131.4	226	14.4	12.6-16.5
2018	1,921	134.5	128.4-140.9	271	17.2	15.2-19.5
2019	1,910	131.2	125.1-137.4	284	18.5	16.4-20.9
2020	1,679	116.9	111.1-122.8	220	14.2	12.3-16.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Cervix uteri**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	122	11.3	9.3-13.7	45	3.8	2.8-5.2
1977	148	13.7	11.5-16.2	49	3.9	2.9-5.3
1978	136	12.6	10.5-15.1	56	4.8	3.6-6.3
1979	143	13.1	11.0-15.6	41	3.5	2.5-4.8
1980	138	13.0	10.8-15.5	45	3.8	2.7-5.1
1981	114	10.7	8.8-13.0	46	4.1	3.0-5.5
1982	156	14.7	12.4-17.4	39	3.3	2.3-4.6
1983	120	11.2	9.2-13.5	57	4.7	3.5-6.1
1984	120	11.2	9.2-13.5	40	3.6	2.5-4.9
1985	115	10.6	8.7-12.8	40	3.6	2.5-5.0
1986	120	11.4	9.4-13.7	38	3.3	2.3-4.6
1987	148	13.8	11.6-16.3	20	1.6	1.0-2.5
1988	145	13.3	11.1-15.7	47	4.1	3.0-5.5
1989	109	10.3	8.4-12.4	35	3.0	2.1-4.3
1990	158	14.5	12.3-17.1	39	3.3	2.3-4.6
1991	146	13.6	11.4-16.0	33	3.1	2.1-4.3
1992	151	13.6	11.5-16.0	29	2.6	1.7-3.8
1993	146	13.2	11.1-15.5	35	3.2	2.2-4.5
1994	183	16.4	14.0-18.9	39	3.5	2.5-4.8
1995	153	13.4	11.3-15.7	51	4.4	3.3-5.8
1996	162	14.1	12.0-16.5	35	2.9	2.0-4.0
1997	150	12.7	10.7-14.9	33	2.8	1.9-3.9
1998	139	11.7	9.8-13.8	36	2.9	2.0-4.1
1999	132	10.9	9.1-13.0	33	2.8	1.9-3.9
2000	128	10.5	8.8-12.6	36	2.8	2.0-3.9
2001	120	9.7	8.1-11.6	26	2.1	1.4-3.1
2002	124	10.1	8.4-12.1	39	3.1	2.2-4.3
2003	115	9.3	7.7-11.2	46	3.7	2.7-5.0
2004	115	9.3	7.7-11.2	25	2.0	1.3-2.9
2005	136	11.3	9.4-13.3	27	2.1	1.4-3.1
2006	127	10.1	8.4-12.1	34	2.6	1.8-3.6
2007	123	9.8	8.1-11.7	35	2.7	1.9-3.8
2008	136	10.9	9.1-12.9	28	2.1	1.4-3.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Cervix uteri**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	127	10.0	8.4-12.0	29	2.1	1.4-3.1
2010	130	10.5	8.8-12.5	36	2.6	1.8-3.6
2011	125	10.0	8.3-11.9	48	3.6	2.6-4.7
2012	140	11.1	9.3-13.1	37	2.8	2.0-3.9
2013	125	9.8	8.1-11.7	41	2.9	2.1-4.0
2014	119	9.1	7.5-10.9	35	2.4	1.7-3.4
2015	127	9.7	8.0-11.6	36	2.5	1.7-3.5
2016	114	8.6	7.1-10.4	43	3.0	2.1-4.1
2017	145	11.2	9.4-13.2	31	2.1	1.4-3.1
2018	123	8.7	7.1-10.4	31	2.1	1.4-3.1
2019	127	9.6	8.0-11.5	16	1.1	0.6-1.8
2020	114	8.5	7.0-10.3	34	2.3	1.6-3.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Queens

Corpus uteris and uterus, not otherwise specified

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	280	23.1	20.4-26.1	68	5.5	4.2-7.0
1977	247	20.7	18.1-23.6	68	5.7	4.4-7.3
1978	248	19.4	17.0-22.1	66	5.1	3.9-6.6
1979	252	19.6	17.2-22.3	69	5.2	4.0-6.7
1980	274	22.0	19.4-24.9	76	5.6	4.4-7.1
1981	274	21.7	19.1-24.5	61	4.6	3.5-5.9
1982	254	20.4	17.9-23.2	67	5.4	4.2-7.0
1983	265	20.7	18.2-23.4	64	4.7	3.6-6.1
1984	276	21.5	19.0-24.3	56	4.3	3.2-5.6
1985	273	21.8	19.2-24.7	57	4.4	3.3-5.8
1986	224	18.0	15.7-20.6	71	5.3	4.1-6.7
1987	271	21.8	19.2-24.7	65	5.0	3.8-6.4
1988	248	20.2	17.7-22.9	55	4.4	3.3-5.8
1989	242	19.6	17.1-22.3	49	3.8	2.8-5.1
1990	232	18.9	16.5-21.6	66	5.0	3.8-6.3
1991	274	22.9	20.2-25.9	72	5.6	4.3-7.0
1992	288	24.3	21.5-27.3	62	4.8	3.7-6.2
1993	277	23.2	20.5-26.2	51	4.0	3.0-5.3
1994	320	26.2	23.3-29.2	64	5.1	3.9-6.5
1995	285	23.1	20.5-26.0	50	3.9	2.9-5.2
1996	305	24.9	22.1-27.9	67	5.1	4.0-6.6
1997	310	25.4	22.6-28.4	55	4.3	3.2-5.6
1998	311	25.5	22.7-28.5	63	5.0	3.8-6.4
1999	354	28.3	25.4-31.4	69	5.3	4.1-6.8
2000	295	23.1	20.6-26.0	55	4.1	3.1-5.4
2001	312	24.7	22.0-27.6	68	5.2	4.0-6.6
2002	334	25.9	23.2-28.8	66	5.0	3.9-6.4
2003	331	25.4	22.7-28.3	66	4.9	3.8-6.2
2004	374	28.8	26.0-31.9	60	4.5	3.4-5.8
2005	369	28.3	25.5-31.4	74	5.5	4.3-7.0
2006	368	28.2	25.4-31.3	75	5.6	4.4-7.1
2007	347	26.4	23.7-29.3	66	4.8	3.7-6.1
2008	410	31.0	28.1-34.2	38	2.8	1.9-3.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Queens

Corpus uteris and uterus, not otherwise specified

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	361	27.2	24.4-30.2	65	4.8	3.7-6.1
2010	367	26.9	24.2-29.9	85	6.0	4.8-7.5
2011	426	30.7	27.8-33.8	76	5.3	4.2-6.7
2012	371	26.3	23.7-29.2	67	4.5	3.5-5.7
2013	459	31.7	28.8-34.7	75	5.0	4.0-6.4
2014	390	26.5	23.9-29.3	92	6.4	5.1-7.8
2015	446	30.2	27.5-33.3	95	6.1	4.9-7.5
2016	461	30.3	27.6-33.3	114	7.3	6.0-8.9
2017	461	30.1	27.4-33.1	92	5.7	4.5-7.0
2018	451	29.5	26.8-32.5	71	4.4	3.4-5.6
2019	475	31.1	28.3-34.2	130	8.1	6.7-9.7
2020	407	26.3	23.8-29.1	90	5.6	4.5-6.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Ovary**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	192	15.8	13.6-18.3	145	11.7	9.8-13.8
1977	193	16.1	13.8-18.6	139	10.9	9.2-13.0
1978	187	15.8	13.5-18.3	121	9.5	7.8-11.4
1979	189	16.7	14.3-19.4	125	10.0	8.3-12.0
1980	177	14.6	12.4-17.0	138	11.0	9.2-13.1
1981	204	16.8	14.5-19.4	124	9.9	8.1-11.9
1982	211	18.0	15.5-20.7	124	10.0	8.2-12.0
1983	206	17.2	14.9-19.9	115	9.1	7.4-10.9
1984	207	17.8	15.4-20.6	119	9.2	7.6-11.1
1985	208	16.7	14.4-19.2	108	8.8	7.1-10.7
1986	222	18.9	16.4-21.6	128	9.8	8.2-11.8
1987	208	17.6	15.2-20.3	134	10.9	9.1-13.0
1988	193	16.6	14.2-19.2	136	10.8	9.0-12.8
1989	190	15.8	13.6-18.3	97	7.6	6.1-9.4
1990	197	17.0	14.6-19.6	117	9.5	7.8-11.4
1991	188	15.8	13.6-18.3	107	8.8	7.2-10.7
1992	212	18.4	16.0-21.1	115	9.6	7.9-11.5
1993	185	15.8	13.6-18.3	118	9.6	7.9-11.5
1994	179	15.1	12.9-17.5	108	8.6	7.1-10.5
1995	203	17.2	14.8-19.7	123	10.0	8.3-12.0
1996	202	16.9	14.6-19.4	96	7.8	6.3-9.6
1997	176	14.6	12.5-17.0	110	8.6	7.1-10.4
1998	176	14.5	12.4-16.8	94	7.4	6.0-9.1
1999	185	14.7	12.7-17.0	94	7.3	5.9-9.0
2000	183	14.6	12.5-16.9	107	8.1	6.6-9.8
2001	205	16.3	14.1-18.7	86	6.5	5.2-8.0
2002	188	14.6	12.6-16.9	108	8.1	6.7-9.8
2003	182	14.1	12.1-16.4	105	7.9	6.5-9.6
2004	177	13.8	11.8-16.0	109	8.2	6.8-10.0
2005	185	14.4	12.4-16.7	109	8.3	6.8-10.0
2006	166	12.8	10.9-14.9	102	7.6	6.2-9.3
2007	151	11.7	9.9-13.8	91	6.9	5.5-8.5
2008	160	12.1	10.3-14.2	94	6.9	5.6-8.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Ovary**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	178	13.5	11.6-15.7	91	6.5	5.2-8.0
2010	176	13.4	11.5-15.6	106	7.6	6.2-9.2
2011	190	13.8	11.9-16.0	112	7.7	6.3-9.4
2012	165	11.9	10.1-13.9	121	8.6	7.1-10.3
2013	155	11.6	9.8-13.6	78	5.4	4.2-6.7
2014	175	12.3	10.5-14.3	89	6.1	4.9-7.6
2015	180	13.2	11.3-15.4	78	5.3	4.1-6.6
2016	178	12.6	10.8-14.7	82	5.5	4.3-6.8
2017	168	11.6	9.8-13.5	89	5.7	4.5-7.1
2018	128	9.1	7.5-10.9	77	4.9	3.8-6.1
2019	169	12.4	10.6-14.6	106	6.7	5.5-8.2
2020	163	11.4	9.6-13.4	64	4.0	3.0-5.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Prostate**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	521	67.0	60.9-73.5	201	28.1	24.1-32.6
1977	551	71.2	64.9-77.8	205	28.6	24.6-33.1
1978	551	70.4	64.3-77.0	188	25.2	21.4-29.3
1979	631	78.7	72.3-85.5	197	28.9	24.7-33.4
1980	661	82.0	75.5-88.9	211	29.4	25.3-33.8
1981	664	83.9	77.3-90.8	224	31.1	26.9-35.6
1982	631	78.5	72.2-85.2	196	26.7	22.9-31.0
1983	645	80.3	74.0-87.0	224	30.4	26.4-34.9
1984	714	86.6	80.1-93.5	234	32.5	28.3-37.1
1985	699	83.0	76.7-89.6	220	28.6	24.8-32.8
1986	700	85.4	79.0-92.3	200	26.6	22.9-30.7
1987	765	91.6	85.0-98.5	219	28.4	24.7-32.6
1988	691	85.9	79.4-92.8	197	26.1	22.4-30.1
1989	731	88.5	82.0-95.4	228	30.4	26.5-34.8
1990	984	119.7	112.1-127.6	260	34.3	30.1-38.8
1991	1,056	125.9	118.3-134.0	238	32.0	27.9-36.4
1992	1,354	159.2	150.7-168.1	257	34.7	30.5-39.3
1993	1,337	154.1	145.8-162.7	256	33.7	29.6-38.2
1994	1,399	161.4	152.9-170.2	253	33.3	29.3-37.8
1995	1,356	156.0	147.7-164.6	233	30.2	26.4-34.4
1996	1,302	149.3	141.2-157.7	213	27.5	23.9-31.5
1997	1,319	150.1	142.0-158.5	220	29.0	25.2-33.1
1998	1,297	144.6	136.8-152.8	249	32.2	28.3-36.5
1999	1,414	156.0	147.9-164.4	186	23.4	20.1-27.0
2000	1,499	162.6	154.4-171.1	200	24.3	21.0-28.0
2001	1,493	159.5	151.4-167.8	203	24.3	21.0-27.8
2002	1,474	155.5	147.6-163.7	195	23.1	19.9-26.6
2003	1,346	141.8	134.3-149.7	183	21.6	18.5-24.9
2004	1,372	141.9	134.4-149.6	187	21.8	18.8-25.2
2005	1,480	152.9	145.1-161.0	162	19.0	16.2-22.2
2006	1,596	164.9	156.8-173.3	160	18.9	16.1-22.1
2007	1,609	162.8	154.8-171.1	172	20.1	17.2-23.3
2008	1,610	157.8	150.1-165.8	165	18.7	15.9-21.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Prostate**

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	1,689	162.1	154.4-170.2	170	19.3	16.5-22.4
2010	1,599	150.0	142.6-157.6	195	21.6	18.7-24.9
2011	1,690	153.5	146.2-161.2	181	19.4	16.7-22.5
2012	1,460	128.4	121.8-135.4	180	18.7	16.0-21.7
2013	1,518	129.8	123.2-136.6	160	16.4	13.9-19.1
2014	1,415	117.5	111.3-124.0	167	16.4	14.0-19.1
2015	1,379	110.3	104.4-116.4	139	13.1	11.0-15.6
2016	1,514	118.4	112.4-124.7	176	16.4	14.1-19.1
2017	1,473	114.3	108.4-120.4	180	16.3	14.0-18.9
2018	1,537	117.0	111.1-123.2	180	16.0	13.7-18.5
2019	1,668	121.6	115.7-127.8	195	16.6	14.4-19.2
2020	1,306	94.4	89.3-99.8	153	12.3	10.4-14.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Testis**

Year of diagnosis	Incidence		
	Cases	Rate per 100,000	95% CI
1976	28	3.0	1.9-4.4
1977	24	2.8	1.7-4.2
1978	20	2.3	1.4-3.5
1979	23	2.6	1.6-4.0
1980	30	3.3	2.2-4.8
1981	28	2.8	1.8-4.1
1982	28	2.8	1.8-4.1
1983	33	3.3	2.2-4.7
1984	31	3.2	2.1-4.6
1985	34	3.2	2.2-4.6
1986	34	3.2	2.2-4.6
1987	34	3.1	2.2-4.5
1988	32	3.1	2.1-4.5
1989	31	3.1	2.1-4.5
1990	33	3.0	2.0-4.3
1991	34	3.1	2.1-4.4
1992	44	4.1	2.9-5.6
1993	36	3.4	2.3-4.7
1994	42	3.8	2.7-5.2
1995	33	3.0	2.0-4.2
1996	29	2.5	1.7-3.6
1997	37	3.3	2.3-4.6
1998	33	2.8	1.9-4.0
1999	44	3.8	2.7-5.1
2000	30	2.7	1.8-3.8
2001	33	2.7	1.9-3.8
2002	45	3.7	2.7-5.0
2003	52	4.5	3.4-6.0
2004	49	4.1	3.1-5.5
2005	48	4.2	3.1-5.6
2006	43	3.7	2.7-5.0
2007	47	4.1	3.0-5.4
2008	36	3.1	2.1-4.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

Mortality data not provided due to small number of deaths per year.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Testis**

Year of diagnosis	Incidence		
	Cases	Rate per 100,000	95% CI
2009	55	4.8	3.6-6.3
2010	45	3.9	2.9-5.3
2011	44	3.7	2.7-5.1
2012	56	4.8	3.6-6.3
2013	46	3.8	2.8-5.1
2014	42	3.5	2.5-4.8
2015	45	3.8	2.7-5.1
2016	52	4.4	3.3-5.8
2017	60	5.0	3.8-6.4
2018	43	3.8	2.7-5.1
2019	56	5.1	3.9-6.7
2020	61	5.7	4.3-7.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

Mortality data not provided due to small number of deaths per year.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Urinary bladder (incl. in situ)**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	361	17.0	15.3-19.0	251	29.9	26.0-34.1	110	8.8	7.2-10.7	133	6.9	5.8-8.3	82	11.0	8.6-13.9	51	4.4	3.3-5.9
1977	427	19.9	18.0-21.9	295	35.0	30.9-39.5	132	10.3	8.6-12.3	134	6.6	5.5-7.8	94	12.0	9.5-14.9	40	3.3	2.3-4.5
1978	426	20.0	18.1-22.0	326	38.7	34.4-43.4	100	7.8	6.3-9.5	118	5.8	4.8-7.0	78	10.4	8.1-13.1	40	3.3	2.3-4.5
1979	402	18.8	17.0-20.8	290	34.8	30.7-39.3	112	9.0	7.3-10.9	126	6.1	5.1-7.3	89	11.6	9.2-14.5	37	3.0	2.1-4.1
1980	463	21.4	19.4-23.5	355	41.8	37.4-46.6	108	8.4	6.9-10.2	110	5.4	4.4-6.5	78	10.3	8.0-12.9	32	2.5	1.7-3.6
1981	473	21.8	19.9-24.0	337	39.5	35.2-44.1	136	10.8	9.0-12.8	92	4.5	3.6-5.5	67	8.4	6.4-10.7	25	2.1	1.3-3.1
1982	440	20.0	18.1-22.0	312	36.5	32.4-41.0	128	9.8	8.1-11.7	117	5.5	4.5-6.6	78	9.8	7.6-12.3	39	3.0	2.1-4.1
1983	485	22.2	20.2-24.3	342	40.2	35.8-44.9	143	10.9	9.2-13.0	95	4.6	3.7-5.6	60	8.4	6.3-10.9	35	2.6	1.8-3.7
1984	442	19.9	18.0-21.9	321	36.8	32.8-41.3	121	9.1	7.5-10.9	106	4.9	4.0-6.0	75	9.1	7.1-11.5	31	2.3	1.6-3.4
1985	382	17.4	15.7-19.3	282	33.2	29.3-37.5	100	7.6	6.2-9.3	107	5.0	4.1-6.1	71	9.5	7.3-12.1	36	2.7	1.9-3.8
1986	407	19.0	17.1-21.0	284	34.0	30.0-38.4	123	9.8	8.1-11.7	97	4.5	3.6-5.5	65	8.1	6.1-10.4	32	2.4	1.6-3.4
1987	431	19.8	17.9-21.8	302	35.5	31.5-39.9	129	9.9	8.2-11.8	97	4.7	3.8-5.7	64	8.6	6.5-11.0	33	2.6	1.8-3.7
1988	394	17.9	16.2-19.8	277	32.2	28.4-36.4	117	8.7	7.2-10.4	95	4.3	3.5-5.3	52	6.4	4.7-8.5	43	3.2	2.3-4.3
1989	375	17.4	15.6-19.3	256	30.3	26.6-34.4	119	9.2	7.6-11.0	110	5.2	4.3-6.3	72	9.4	7.3-11.9	38	2.8	2.0-3.9
1990	427	19.8	17.9-21.8	298	35.5	31.4-39.8	129	9.7	8.1-11.5	89	4.2	3.4-5.2	53	7.1	5.3-9.4	36	2.7	1.9-3.8
1991	425	19.9	18.0-21.9	311	37.4	33.3-41.9	114	8.8	7.3-10.6	82	4.0	3.1-4.9	52	7.1	5.2-9.3	30	2.2	1.5-3.2
1992	455	21.2	19.3-23.2	307	36.7	32.6-41.1	148	11.5	9.7-13.5	100	4.7	3.8-5.8	66	8.7	6.7-11.1	34	2.6	1.8-3.7
1993	414	19.4	17.5-21.3	294	35.3	31.3-39.7	120	9.3	7.7-11.2	107	5.1	4.2-6.2	81	10.6	8.3-13.2	26	2.0	1.3-2.9
1994	469	21.8	19.8-23.8	308	36.8	32.7-41.2	161	12.4	10.6-14.5	84	3.9	3.1-4.9	58	7.8	5.9-10.1	26	2.0	1.3-2.9
1995	443	20.6	18.7-22.6	298	35.6	31.6-40.0	145	11.2	9.4-13.2	91	4.2	3.4-5.2	55	7.1	5.3-9.2	36	2.7	1.9-3.7
1996	409	19.1	17.3-21.1	289	34.9	31.0-39.2	120	9.2	7.6-11.0	93	4.3	3.5-5.3	57	7.3	5.5-9.5	36	2.7	1.9-3.8
1997	437	20.3	18.4-22.3	314	36.9	32.9-41.2	123	9.5	7.8-11.3	81	3.7	3.0-4.7	58	7.6	5.7-9.8	23	1.7	1.1-2.6
1998	432	19.9	18.1-21.9	291	34.4	30.5-38.6	141	10.7	9.0-12.7	98	4.5	3.7-5.5	60	7.4	5.6-9.5	38	2.7	1.9-3.8
1999	416	18.8	17.0-20.7	299	34.3	30.5-38.5	117	8.7	7.2-10.5	88	4.0	3.2-4.9	54	6.5	4.9-8.5	34	2.5	1.7-3.5
2000	421	18.7	17.0-20.6	283	32.1	28.5-36.1	138	10.1	8.5-11.9	89	3.9	3.2-4.9	60	7.1	5.4-9.2	29	2.1	1.4-3.0
2001	403	17.8	16.1-19.7	286	32.0	28.4-36.0	117	8.6	7.1-10.3	92	4.1	3.3-5.0	70	8.1	6.3-10.2	22	1.6	1.0-2.4
2002	414	18.2	16.5-20.1	294	33.0	29.3-37.0	120	8.8	7.2-10.5	71	3.1	2.4-3.9	44	5.2	3.7-6.9	27	1.9	1.2-2.7
2003	424	18.5	16.8-20.4	305	33.7	30.0-37.8	119	8.7	7.2-10.4	74	3.2	2.5-4.0	53	6.0	4.5-7.9	21	1.5	0.9-2.3
2004	442	19.2	17.5-21.1	325	35.4	31.6-39.5	117	8.5	7.0-10.2	84	3.6	2.9-4.5	61	6.8	5.2-8.8	23	1.6	1.0-2.4
2005	420	18.3	16.5-20.1	306	33.2	29.5-37.1	114	8.5	7.0-10.2	90	3.9	3.1-4.8	55	6.4	4.8-8.3	35	2.4	1.6-3.3
2006	430	18.6	16.9-20.5	311	33.7	30.1-37.7	119	8.7	7.2-10.4	96	4.1	3.3-5.0	61	7.1	5.4-9.1	35	2.5	1.7-3.5
2007	450	19.6	17.8-21.5	333	36.4	32.6-40.6	117	8.6	7.1-10.4	87	3.7	2.9-4.5	58	6.4	4.8-8.2	29	2.0	1.4-3.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Urinary bladder (incl. in situ)**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	442	19.0	17.3-20.9	341	36.2	32.4-40.3	101	7.5	6.1-9.1	101	4.4	3.6-5.3	70	7.8	6.1-9.9	31	2.3	1.5-3.3
2009	452	19.2	17.4-21.0	328	34.3	30.6-38.3	124	8.8	7.3-10.6	78	3.3	2.6-4.1	47	5.1	3.7-6.7	31	2.1	1.4-3.0
2010	438	18.4	16.7-20.2	321	33.5	29.8-37.4	117	8.4	7.0-10.2	104	4.2	3.4-5.1	73	7.6	5.9-9.6	31	2.0	1.4-2.9
2011	428	17.8	16.2-19.6	318	31.9	28.4-35.7	110	8.0	6.6-9.7	85	3.4	2.7-4.3	50	5.2	3.9-6.9	35	2.4	1.7-3.4
2012	446	18.0	16.3-19.8	308	30.7	27.3-34.3	138	9.4	7.9-11.2	104	4.1	3.3-4.9	68	6.8	5.3-8.7	36	2.2	1.5-3.1
2013	417	16.5	15.0-18.2	305	28.8	25.6-32.3	112	7.8	6.4-9.4	83	3.2	2.5-3.9	55	5.6	4.2-7.3	28	1.7	1.1-2.6
2014	464	17.8	16.2-19.5	327	30.3	27.0-33.8	137	9.0	7.5-10.7	91	3.5	2.8-4.3	60	6.1	4.7-7.9	31	1.8	1.2-2.7
2015	435	16.5	14.9-18.1	327	29.7	26.5-33.2	108	7.1	5.8-8.6	87	3.3	2.6-4.1	60	5.7	4.3-7.3	27	1.7	1.1-2.5
2016	437	16.0	14.5-17.6	325	28.6	25.5-31.9	112	7.1	5.8-8.6	76	2.8	2.2-3.6	57	5.4	4.0-7.0	19	1.3	0.8-2.1
2017	471	16.9	15.4-18.5	351	29.9	26.8-33.2	120	7.5	6.2-9.1	85	3.1	2.4-3.8	64	5.7	4.4-7.3	21	1.3	0.8-2.1
2018	424	15.0	13.6-16.5	319	26.4	23.5-29.5	105	6.4	5.2-7.8	78	2.7	2.2-3.4	47	4.2	3.1-5.6	31	1.8	1.2-2.7
2019	456	16.0	14.5-17.5	333	27.3	24.4-30.5	123	7.4	6.1-8.9	90	3.0	2.4-3.7	62	5.2	4.0-6.7	28	1.6	1.0-2.4
2020	370	12.6	11.3-14.0	283	22.6	20.0-25.5	87	5.3	4.2-6.6	62	2.1	1.6-2.7	48	4.0	2.9-5.3	14	0.8	0.4-1.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Kidney and renal pelvis**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	173	8.2	6.9-9.5	104	11.6	9.3-14.2	69	5.9	4.5-7.5	87	4.4	3.5-5.4	56	7.3	5.4-9.7	31	2.6	1.7-3.7
1977	173	8.1	6.9-9.4	107	12.0	9.7-14.6	66	5.4	4.1-7.0	89	4.1	3.3-5.1	56	6.4	4.8-8.5	33	2.5	1.7-3.5
1978	144	6.7	5.6-7.9	98	10.9	8.8-13.4	46	3.7	2.7-5.0	86	4.1	3.3-5.1	60	7.1	5.3-9.2	26	2.1	1.4-3.2
1979	157	7.3	6.2-8.6	100	11.2	9.0-13.8	57	4.5	3.4-5.9	68	3.3	2.6-4.3	41	5.1	3.6-7.0	27	2.2	1.4-3.2
1980	147	6.7	5.7-8.0	92	10.1	8.0-12.5	55	4.5	3.4-5.9	70	3.2	2.5-4.0	47	5.3	3.8-7.1	23	1.8	1.1-2.8
1981	160	7.2	6.1-8.5	103	11.8	9.5-14.4	57	4.1	3.1-5.4	60	2.7	2.0-3.5	38	4.3	3.0-6.0	22	1.6	1.0-2.5
1982	171	8.0	6.8-9.4	108	12.2	10.0-14.9	63	5.0	3.8-6.5	86	3.9	3.1-4.9	48	5.7	4.1-7.7	38	2.8	2.0-3.9
1983	157	7.5	6.4-8.8	100	11.9	9.6-14.6	57	4.7	3.6-6.2	67	3.1	2.4-4.0	43	5.1	3.7-7.0	24	1.8	1.1-2.7
1984	190	8.7	7.5-10.0	118	12.8	10.5-15.4	72	5.8	4.5-7.3	80	3.7	2.9-4.6	57	6.6	5.0-8.7	23	1.8	1.1-2.8
1985	180	8.5	7.3-9.9	104	12.1	9.8-14.7	76	6.1	4.8-7.7	72	3.5	2.7-4.4	39	4.6	3.2-6.4	33	2.8	1.9-4.0
1986	177	8.0	6.9-9.4	104	11.7	9.5-14.3	73	5.7	4.4-7.2	78	3.6	2.9-4.6	54	6.5	4.8-8.6	24	1.8	1.2-2.8
1987	225	10.5	9.1-12.0	142	16.1	13.5-19.1	83	6.6	5.2-8.2	59	2.8	2.1-3.6	35	4.2	2.9-5.9	24	1.9	1.2-2.8
1988	163	7.8	6.6-9.1	103	11.9	9.7-14.5	60	4.9	3.7-6.4	66	3.2	2.4-4.0	39	4.5	3.1-6.1	27	2.2	1.5-3.3
1989	190	9.2	7.9-10.6	106	12.5	10.1-15.2	84	7.1	5.6-8.8	72	3.4	2.6-4.3	40	5.0	3.5-6.8	32	2.5	1.7-3.6
1990	216	10.2	8.8-11.6	116	13.6	11.2-16.3	100	8.2	6.7-10.1	81	3.9	3.1-4.9	46	5.7	4.1-7.7	35	2.9	2.0-4.0
1991	182	8.6	7.4-10.0	97	11.1	9.0-13.6	85	7.0	5.6-8.8	62	2.9	2.2-3.7	35	3.8	2.6-5.3	27	2.1	1.4-3.2
1992	202	9.6	8.3-11.1	127	14.7	12.2-17.5	75	6.1	4.8-7.7	65	3.1	2.4-3.9	37	4.6	3.2-6.3	28	2.1	1.4-3.1
1993	219	10.3	9.0-11.8	126	14.2	11.8-16.9	93	7.6	6.1-9.4	67	3.1	2.4-4.0	45	5.0	3.6-6.7	22	1.7	1.0-2.6
1994	236	11.3	9.9-12.9	143	16.7	14.0-19.7	93	7.4	6.0-9.2	65	3.0	2.3-3.9	37	4.4	3.1-6.1	28	2.1	1.4-3.1
1995	225	10.6	9.2-12.0	133	14.7	12.3-17.4	92	7.7	6.2-9.4	71	3.4	2.6-4.2	44	5.1	3.7-6.9	27	2.2	1.4-3.2
1996	246	11.4	10.0-13.0	148	16.7	14.1-19.6	98	7.7	6.3-9.4	68	3.2	2.5-4.0	47	5.5	4.0-7.3	21	1.6	1.0-2.5
1997	242	11.3	9.9-12.8	146	16.6	14.0-19.5	96	7.8	6.3-9.5	65	3.0	2.4-3.9	45	5.5	4.0-7.3	20	1.5	0.9-2.4
1998	269	12.5	11.0-14.0	172	18.9	16.2-21.9	97	7.8	6.3-9.6	71	3.3	2.6-4.2	40	4.7	3.4-6.4	31	2.5	1.7-3.5
1999	259	11.7	10.3-13.2	153	16.5	14.0-19.3	106	8.2	6.7-9.9	60	2.7	2.1-3.5	38	4.2	3.0-5.8	22	1.6	1.0-2.5
2000	243	10.9	9.6-12.3	154	16.5	13.9-19.3	89	6.8	5.5-8.4	61	2.7	2.1-3.5	39	4.4	3.1-6.0	22	1.6	1.0-2.5
2001	256	11.4	10.0-12.9	161	16.8	14.3-19.6	95	7.3	5.9-9.0	59	2.6	2.0-3.4	33	3.6	2.5-5.0	26	2.0	1.3-3.0
2002	254	11.2	9.9-12.7	173	17.6	15.0-20.4	81	6.2	4.9-7.7	58	2.6	1.9-3.3	43	4.7	3.4-6.3	15	1.1	0.6-1.8
2003	273	12.0	10.6-13.5	173	17.4	14.9-20.2	100	7.7	6.2-9.3	57	2.5	1.9-3.2	35	3.7	2.5-5.1	22	1.6	1.0-2.4
2004	325	14.2	12.7-15.8	196	20.0	17.3-23.0	129	10.0	8.3-11.8	82	3.6	2.9-4.5	52	5.7	4.3-7.5	30	2.3	1.5-3.2
2005	302	13.3	11.8-14.9	192	19.6	16.9-22.6	110	8.6	7.0-10.4	49	2.2	1.6-2.9	29	3.1	2.1-4.5	20	1.5	0.9-2.3
2006	337	14.8	13.2-16.4	202	20.5	17.7-23.5	135	10.5	8.8-12.5	55	2.4	1.8-3.1	29	3.0	2.0-4.3	26	1.9	1.3-2.9
2007	333	14.4	12.9-16.0	220	22.0	19.1-25.1	113	8.6	7.1-10.4	62	2.6	2.0-3.4	43	4.6	3.3-6.2	19	1.4	0.8-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Kidney and renal pelvis**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	349	14.7	13.2-16.4	233	22.6	19.8-25.8	116	8.7	7.2-10.5	69	2.9	2.2-3.7	40	4.0	2.8-5.5	29	2.0	1.4-3.0
2009	302	12.9	11.5-14.4	177	17.2	14.7-19.9	125	9.5	7.8-11.3	73	3.1	2.4-3.9	51	5.1	3.8-6.7	22	1.6	1.0-2.5
2010	346	14.3	12.9-16.0	230	21.5	18.8-24.6	116	8.7	7.2-10.5	66	2.7	2.1-3.5	37	3.8	2.6-5.2	29	2.0	1.3-2.9
2011	324	13.2	11.8-14.8	208	18.8	16.3-21.6	116	8.7	7.2-10.5	52	2.1	1.5-2.7	36	3.5	2.4-4.8	16	1.1	0.6-1.8
2012	321	12.6	11.3-14.1	203	18.0	15.6-20.7	118	8.4	6.9-10.1	61	2.4	1.8-3.1	39	3.6	2.5-5.0	22	1.5	0.9-2.3
2013	349	13.7	12.3-15.2	232	20.6	18.0-23.5	117	8.2	6.8-9.9	53	2.1	1.5-2.7	33	3.2	2.2-4.5	20	1.3	0.8-2.0
2014	346	13.0	11.7-14.5	240	19.8	17.4-22.6	106	7.4	6.0-9.0	58	2.2	1.7-2.9	33	3.0	2.1-4.3	25	1.6	1.0-2.4
2015	350	13.1	11.7-14.6	222	18.1	15.8-20.8	128	8.9	7.4-10.6	69	2.5	2.0-3.2	46	4.0	2.9-5.4	23	1.4	0.9-2.2
2016	385	14.3	12.9-15.9	255	21.1	18.5-23.9	130	8.8	7.3-10.5	58	2.2	1.6-2.8	34	3.1	2.1-4.4	24	1.5	1.0-2.3
2017	379	13.8	12.4-15.3	226	18.1	15.8-20.7	153	10.3	8.7-12.2	71	2.6	2.0-3.3	46	3.9	2.8-5.2	25	1.7	1.1-2.5
2018	379	13.8	12.4-15.3	241	19.6	17.1-22.3	138	9.1	7.6-10.8	59	2.1	1.6-2.7	40	3.3	2.3-4.5	19	1.1	0.7-1.8
2019	457	16.6	15.1-18.2	293	23.1	20.4-26.0	164	11.1	9.4-13.0	53	1.8	1.4-2.4	32	2.5	1.7-3.5	21	1.3	0.8-2.0
2020	369	13.4	12.0-14.9	242	19.0	16.6-21.7	127	8.5	7.1-10.2	55	2.0	1.5-2.6	41	3.4	2.4-4.7	14	1.0	0.5-1.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Brain and other central nervous system**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	110	5.1	4.1-6.2	60	6.3	4.8-8.3	50	4.1	3.0-5.4	78	3.7	2.9-4.7	44	4.9	3.5-6.7	34	2.9	1.9-4.1
1977	135	6.5	5.4-7.7	71	7.5	5.8-9.6	64	5.5	4.2-7.1	82	3.8	3.0-4.8	43	4.5	3.2-6.1	39	3.3	2.3-4.5
1978	120	5.6	4.6-6.8	55	5.7	4.2-7.5	65	5.5	4.2-7.1	89	4.2	3.3-5.2	46	5.0	3.6-6.7	43	3.6	2.5-4.9
1979	124	6.0	4.9-7.2	63	7.2	5.5-9.3	61	5.2	3.9-6.8	94	4.4	3.5-5.4	50	5.6	4.1-7.4	44	3.5	2.5-4.8
1980	113	5.4	4.4-6.5	60	6.8	5.1-8.8	53	4.5	3.3-5.9	91	4.1	3.3-5.0	53	5.4	4.0-7.2	38	3.1	2.1-4.3
1981	125	6.1	5.1-7.3	69	8.0	6.1-10.2	56	4.6	3.4-6.1	83	3.9	3.1-4.9	43	5.0	3.6-6.8	40	3.2	2.2-4.4
1982	129	6.2	5.2-7.5	66	7.4	5.6-9.4	63	5.3	4.0-6.9	97	4.5	3.6-5.5	40	4.4	3.1-6.0	57	4.6	3.4-6.0
1983	123	6.1	5.0-7.3	73	8.0	6.2-10.1	50	4.4	3.2-5.9	90	4.1	3.3-5.0	48	5.3	3.9-7.1	42	3.1	2.2-4.2
1984	126	6.2	5.1-7.4	62	7.1	5.4-9.1	64	5.5	4.2-7.0	74	3.3	2.6-4.2	47	4.9	3.6-6.7	27	2.2	1.4-3.3
1985	115	5.5	4.5-6.7	60	6.6	5.0-8.5	55	4.7	3.5-6.2	81	3.8	3.0-4.7	35	4.0	2.7-5.6	46	3.6	2.6-4.9
1986	116	5.6	4.6-6.8	67	7.8	6.0-10.0	49	4.1	3.0-5.4	78	3.6	2.8-4.5	46	5.3	3.8-7.1	32	2.4	1.7-3.5
1987	134	6.7	5.6-7.9	69	7.7	5.9-9.8	65	5.8	4.4-7.5	77	3.7	2.9-4.6	38	4.2	3.0-5.8	39	3.2	2.3-4.4
1988	117	5.5	4.5-6.6	55	5.9	4.4-7.7	62	5.2	3.9-6.7	72	3.4	2.7-4.3	38	4.2	3.0-5.9	34	2.9	2.0-4.1
1989	140	7.1	6.0-8.4	77	9.2	7.2-11.5	63	5.7	4.3-7.4	70	3.3	2.6-4.2	38	4.4	3.1-6.1	32	2.5	1.7-3.5
1990	120	6.0	4.9-7.2	60	6.6	5.0-8.6	60	5.5	4.1-7.1	71	3.5	2.7-4.4	39	4.5	3.2-6.2	32	2.7	1.8-3.8
1991	127	6.2	5.2-7.4	52	5.9	4.4-7.8	75	6.4	5.0-8.0	71	3.3	2.6-4.2	37	4.1	2.8-5.7	34	2.7	1.9-3.9
1992	143	6.9	5.8-8.2	71	7.6	5.9-9.7	72	6.3	4.9-8.0	55	2.6	1.9-3.3	29	3.3	2.2-4.8	26	2.1	1.4-3.1
1993	139	6.8	5.7-8.0	72	8.2	6.4-10.4	67	5.8	4.5-7.4	66	3.2	2.5-4.1	34	3.9	2.7-5.5	32	2.7	1.8-3.8
1994	141	6.9	5.8-8.1	71	7.7	6.0-9.8	70	6.1	4.7-7.7	64	3.0	2.3-3.9	33	3.8	2.6-5.3	31	2.5	1.7-3.6
1995	150	7.1	6.0-8.4	72	7.8	6.1-9.8	78	6.6	5.2-8.2	67	3.2	2.5-4.1	35	3.9	2.7-5.5	32	2.5	1.7-3.6
1996	159	7.5	6.4-8.7	80	8.4	6.6-10.5	79	6.8	5.4-8.5	75	3.6	2.8-4.5	40	4.4	3.1-6.0	35	2.8	2.0-3.9
1997	128	6.1	5.1-7.2	66	7.2	5.6-9.2	62	5.4	4.1-6.9	70	3.3	2.6-4.1	29	3.3	2.2-4.8	41	3.4	2.4-4.6
1998	152	7.0	6.0-8.3	85	9.0	7.1-11.1	67	5.6	4.4-7.2	65	3.0	2.4-3.9	38	4.1	2.9-5.7	27	2.2	1.5-3.3
1999	123	5.6	4.7-6.7	62	6.3	4.8-8.1	61	4.9	3.8-6.3	84	3.8	3.0-4.7	42	4.3	3.1-5.9	42	3.3	2.4-4.5
2000	146	6.6	5.5-7.7	73	7.4	5.8-9.3	73	5.7	4.5-7.2	74	3.4	2.6-4.2	29	3.1	2.1-4.4	45	3.6	2.6-4.8
2001	117	5.3	4.4-6.4	59	6.0	4.6-7.8	58	4.8	3.6-6.2	63	2.8	2.2-3.6	36	3.7	2.6-5.1	27	2.1	1.4-3.0
2002	147	6.6	5.6-7.8	79	7.9	6.3-9.9	68	5.6	4.3-7.1	76	3.4	2.7-4.3	34	3.5	2.4-4.9	42	3.4	2.4-4.6
2003	111	4.8	4.0-5.8	56	5.3	4.0-6.9	55	4.3	3.3-5.7	52	2.2	1.7-2.9	33	3.2	2.2-4.5	19	1.4	0.8-2.2
2004	137	6.1	5.1-7.2	78	7.8	6.2-9.8	59	4.6	3.5-6.0	64	2.8	2.2-3.6	36	3.6	2.5-5.1	28	2.2	1.4-3.1
2005	154	7.0	5.9-8.2	88	9.1	7.3-11.3	66	5.6	4.3-7.1	71	3.1	2.4-3.9	40	4.2	3.0-5.7	31	2.4	1.6-3.4
2006	140	6.2	5.2-7.4	83	8.4	6.7-10.4	57	4.5	3.4-5.9	71	3.2	2.5-4.0	37	3.9	2.7-5.3	34	2.7	1.9-3.8
2007	114	5.1	4.2-6.1	62	6.0	4.6-7.8	52	4.2	3.1-5.6	67	2.9	2.3-3.7	38	3.9	2.8-5.4	29	2.2	1.4-3.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Queens

Brain and other central nervous system

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	138	6.2	5.2-7.3	76	7.5	5.9-9.4	62	5.2	4.0-6.7	72	3.1	2.4-4.0	42	4.1	3.0-5.6	30	2.3	1.6-3.4
2009	142	6.2	5.2-7.3	75	7.2	5.6-9.1	67	5.4	4.1-6.9	71	3.1	2.4-3.9	43	4.4	3.1-5.9	28	2.2	1.5-3.2
2010	142	6.2	5.2-7.3	79	7.7	6.1-9.7	63	5.1	3.9-6.6	68	2.8	2.2-3.6	36	3.4	2.4-4.8	32	2.5	1.7-3.5
2011	153	6.5	5.5-7.7	82	7.6	6.0-9.5	71	5.5	4.2-6.9	78	3.3	2.6-4.1	44	4.2	3.0-5.6	34	2.6	1.8-3.7
2012	140	5.8	4.9-6.9	76	7.0	5.5-8.8	64	5.0	3.8-6.4	79	3.1	2.5-3.9	29	2.6	1.7-3.8	50	3.6	2.6-4.8
2013	112	4.7	3.9-5.7	55	5.0	3.7-6.5	57	4.5	3.4-5.9	63	2.5	1.9-3.2	33	3.0	2.0-4.2	30	2.1	1.4-3.0
2014	125	5.2	4.3-6.2	73	6.4	5.0-8.1	52	4.2	3.1-5.6	75	2.9	2.3-3.7	41	3.6	2.6-4.9	34	2.4	1.6-3.4
2015	127	5.1	4.2-6.1	67	5.8	4.5-7.4	60	4.3	3.3-5.6	67	2.6	2.0-3.4	35	3.1	2.1-4.3	32	2.4	1.6-3.4
2016	136	5.3	4.4-6.2	76	6.4	5.0-8.0	60	4.1	3.1-5.4	81	3.1	2.5-3.9	41	3.5	2.5-4.8	40	2.8	2.0-3.9
2017	138	5.4	4.5-6.4	74	6.3	4.9-8.0	64	4.6	3.5-6.0	75	2.7	2.1-3.4	38	3.3	2.3-4.5	37	2.3	1.6-3.2
2018	144	5.8	4.8-6.8	82	6.9	5.5-8.7	62	4.8	3.6-6.2	79	3.0	2.4-3.8	39	3.4	2.4-4.6	40	2.7	1.9-3.8
2019	124	4.9	4.1-5.9	70	5.8	4.5-7.4	54	4.2	3.1-5.5	71	2.7	2.1-3.4	39	3.2	2.3-4.5	32	2.3	1.5-3.3
2020	118	4.8	4.0-5.8	61	5.2	4.0-6.8	57	4.4	3.3-5.8	79	2.9	2.3-3.6	50	4.0	2.9-5.3	29	1.8	1.2-2.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Thyroid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	62	3.0	2.3-3.9	16	1.7	0.9-2.8	46	4.1	3.0-5.6	17	0.7	0.4-1.2	6	0.6	0.2-1.4	11	0.8	0.4-1.5
1977	62	3.2	2.4-4.1	14	1.5	0.8-2.5	48	4.5	3.3-6.0	20	0.9	0.6-1.5	9	1.0	0.4-2.0	11	0.9	0.4-1.6
1978	69	3.5	2.7-4.5	22	2.5	1.5-3.8	47	4.4	3.2-5.8	8	0.4	0.2-0.8	1	0.2	0.0-1.1	7	0.5	0.2-1.2
1979	67	3.6	2.7-4.6	25	2.7	1.7-4.1	42	4.3	3.1-5.9	9	0.4	0.2-0.8	5	0.5	0.2-1.3	4	0.3	0.1-0.8
1980	71	3.8	2.9-4.8	17	2.2	1.3-3.5	54	5.1	3.8-6.7	16	0.8	0.5-1.3	6	0.7	0.3-1.6	10	0.8	0.4-1.5
1981	73	3.7	2.9-4.7	18	2.0	1.2-3.2	55	5.1	3.8-6.7	12	0.6	0.3-1.0	5	0.5	0.2-1.2	7	0.6	0.2-1.2
1982	65	3.3	2.5-4.2	22	2.9	1.8-4.4	43	3.7	2.6-5.0	15	0.7	0.4-1.1	8	0.9	0.4-1.8	7	0.5	0.2-1.1
1983	75	3.6	2.8-4.6	16	1.5	0.8-2.5	59	5.4	4.0-7.0	9	0.4	0.2-0.9	1	0.1	0.0-0.6	8	0.7	0.3-1.4
1984	93	4.5	3.6-5.6	24	2.7	1.7-4.1	69	6.1	4.7-7.7	12	0.6	0.3-1.0	3	0.4	0.1-1.2	9	0.7	0.3-1.3
1985	107	5.2	4.2-6.3	33	3.3	2.3-4.7	74	6.7	5.2-8.4	23	1.0	0.7-1.6	10	1.2	0.5-2.3	13	1.0	0.5-1.8
1986	85	4.0	3.1-4.9	21	2.3	1.4-3.5	64	5.4	4.1-7.0	12	0.6	0.3-1.0	2	0.2	0.0-0.7	10	0.8	0.4-1.5
1987	80	4.2	3.3-5.2	23	2.8	1.7-4.2	57	5.4	4.1-7.1	9	0.4	0.2-0.7	3	0.3	0.1-0.9	6	0.4	0.2-1.0
1988	84	4.2	3.3-5.2	25	3.1	2.0-4.6	59	5.4	4.1-7.0	13	0.6	0.3-1.0	3	0.5	0.1-1.4	10	0.7	0.3-1.4
1989	67	3.4	2.6-4.4	14	1.5	0.8-2.6	53	5.1	3.8-6.7	15	0.7	0.4-1.1	4	0.4	0.1-1.1	11	0.8	0.4-1.5
1990	87	4.2	3.3-5.2	21	2.4	1.5-3.7	66	5.7	4.4-7.4	11	0.5	0.2-0.9	4	0.5	0.1-1.2	7	0.5	0.2-1.2
1991	96	4.7	3.8-5.7	22	2.6	1.6-3.9	74	6.5	5.1-8.3	19	0.9	0.5-1.4	4	0.6	0.1-1.4	15	1.2	0.7-2.0
1992	103	5.0	4.1-6.1	23	2.6	1.6-3.9	80	7.1	5.6-8.9	9	0.4	0.2-0.8	3	0.4	0.1-1.1	6	0.4	0.2-1.0
1993	117	5.7	4.7-6.9	27	2.9	1.9-4.3	90	8.2	6.6-10.1	12	0.6	0.3-1.0	4	0.4	0.1-1.1	8	0.6	0.2-1.2
1994	104	5.1	4.1-6.1	32	3.4	2.3-4.8	72	6.6	5.1-8.3	16	0.8	0.4-1.3	9	1.0	0.5-1.9	7	0.5	0.2-1.1
1995	151	6.9	5.9-8.1	36	3.5	2.4-4.8	115	10.0	8.3-12.1	13	0.6	0.3-1.0	3	0.3	0.1-0.9	10	0.7	0.3-1.4
1996	127	5.9	4.9-7.0	34	3.6	2.5-5.1	93	7.9	6.4-9.7	16	0.7	0.4-1.2	7	0.9	0.3-1.8	9	0.7	0.3-1.3
1997	152	6.9	5.9-8.1	36	3.7	2.6-5.1	116	9.8	8.0-11.7	14	0.6	0.4-1.1	3	0.3	0.1-1.0	11	0.8	0.4-1.5
1998	149	6.7	5.6-7.8	33	3.3	2.3-4.7	116	9.7	8.0-11.7	8	0.4	0.2-0.7	2	0.2	0.0-0.7	6	0.5	0.2-1.1
1999	160	7.1	6.1-8.3	36	3.6	2.5-5.0	124	10.4	8.6-12.4	10	0.5	0.2-0.8	6	0.7	0.2-1.4	4	0.3	0.1-0.8
2000	182	8.0	6.9-9.3	34	3.2	2.2-4.6	148	12.3	10.4-14.5	19	0.9	0.5-1.3	6	0.7	0.2-1.5	13	1.0	0.5-1.7
2001	206	8.9	7.7-10.2	42	4.1	2.9-5.6	164	13.5	11.5-15.7	11	0.5	0.2-0.9	2	0.2	0.0-0.8	9	0.7	0.3-1.3
2002	203	8.8	7.7-10.2	51	4.9	3.7-6.5	152	12.5	10.6-14.7	10	0.4	0.2-0.8	4	0.4	0.1-1.1	6	0.4	0.1-0.9
2003	198	8.6	7.4-9.9	45	4.3	3.1-5.7	153	12.6	10.6-14.7	9	0.4	0.2-0.8	5	0.6	0.2-1.4	4	0.3	0.1-0.8
2004	221	9.6	8.4-11.0	48	4.5	3.3-6.0	173	14.3	12.2-16.6	9	0.4	0.2-0.7	4	0.4	0.1-1.1	5	0.3	0.1-0.8
2005	276	12.0	10.6-13.5	72	7.0	5.4-8.8	204	16.7	14.5-19.2	12	0.5	0.3-0.9	7	0.8	0.3-1.6	5	0.4	0.1-1.0
2006	267	11.6	10.2-13.1	55	5.1	3.9-6.7	212	17.3	15.1-19.9	19	0.9	0.5-1.3	6	0.7	0.2-1.4	13	1.0	0.5-1.8
2007	300	13.0	11.6-14.6	58	5.4	4.1-7.0	242	20.1	17.6-22.8	12	0.5	0.3-0.9	2	0.2	0.0-0.8	10	0.7	0.3-1.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Thyroid**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	349	14.7	13.2-16.4	76	7.0	5.5-8.7	273	21.9	19.3-24.7	11	0.5	0.2-0.9	5	0.6	0.2-1.4	6	0.4	0.2-1.0
2009	393	16.6	14.9-18.3	90	8.0	6.4-9.9	303	24.5	21.8-27.5	16	0.7	0.4-1.1	4	0.5	0.1-1.2	12	0.9	0.4-1.5
2010	403	17.0	15.3-18.7	93	8.2	6.6-10.0	310	25.3	22.5-28.3	17	0.7	0.4-1.1	6	0.6	0.2-1.2	11	0.8	0.4-1.4
2011	420	17.5	15.8-19.3	91	8.1	6.5-9.9	329	26.1	23.3-29.2	8	0.3	0.1-0.7	2	0.2	0.0-0.8	6	0.4	0.1-0.9
2012	475	19.4	17.7-21.3	102	8.8	7.1-10.7	373	29.4	26.5-32.6	16	0.7	0.4-1.1	10	1.0	0.5-1.8	6	0.4	0.1-1.0
2013	507	20.7	18.9-22.6	112	9.7	8.0-11.7	395	31.2	28.1-34.5	12	0.5	0.2-0.9	4	0.4	0.1-1.0	8	0.5	0.2-1.1
2014	503	20.1	18.4-22.0	111	9.2	7.6-11.2	392	30.4	27.4-33.6	13	0.5	0.3-0.8	2	0.1	0.0-0.6	11	0.7	0.4-1.3
2015	528	21.2	19.4-23.1	125	10.3	8.5-12.3	403	31.6	28.5-34.9	14	0.5	0.3-0.9	4	0.3	0.1-0.9	10	0.7	0.3-1.3
2016	533	21.6	19.8-23.5	115	9.6	7.9-11.6	418	32.9	29.8-36.3	14	0.5	0.3-0.8	7	0.5	0.2-1.1	7	0.4	0.2-0.9
2017	495	19.7	17.9-21.5	120	10.0	8.3-12.0	375	28.9	25.9-32.0	17	0.6	0.4-1.0	6	0.5	0.2-1.1	11	0.7	0.3-1.2
2018	484	19.8	18.1-21.7	113	9.4	7.7-11.4	371	29.8	26.8-33.1	11	0.4	0.2-0.7	3	0.2	0.0-0.7	8	0.5	0.2-1.1
2019	536	22.1	20.2-24.1	123	10.1	8.4-12.2	413	33.6	30.3-37.1	18	0.6	0.4-1.0	9	0.8	0.3-1.5	9	0.5	0.2-1.0
2020	414	17.0	15.4-18.8	107	9.3	7.6-11.3	307	24.5	21.8-27.5	16	0.6	0.3-1.0	8	0.7	0.3-1.4	8	0.5	0.2-1.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Hodgkin lymphoma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	62	3.0	2.3-3.9	37	3.8	2.7-5.4	25	2.4	1.5-3.5	26	1.3	0.8-2.0	16	1.9	1.0-3.2	10	0.9	0.4-1.6
1977	77	3.6	2.8-4.5	44	4.4	3.2-6.0	33	2.9	2.0-4.1	24	1.2	0.8-1.8	14	1.7	0.9-2.9	10	0.9	0.4-1.6
1978	60	2.9	2.2-3.8	32	3.4	2.3-4.8	28	2.6	1.7-3.8	29	1.3	0.9-1.9	14	1.3	0.7-2.3	15	1.2	0.7-2.1
1979	66	3.5	2.7-4.4	44	5.0	3.6-6.8	22	2.2	1.4-3.4	18	0.9	0.5-1.4	10	1.2	0.5-2.2	8	0.7	0.3-1.5
1980	66	3.1	2.4-4.0	34	3.7	2.5-5.2	32	2.7	1.9-3.9	23	1.1	0.7-1.6	12	1.3	0.7-2.4	11	0.9	0.4-1.6
1981	74	3.5	2.7-4.4	43	4.6	3.3-6.2	31	2.7	1.8-3.8	18	1.0	0.6-1.5	10	1.2	0.6-2.2	8	0.8	0.3-1.5
1982	66	3.2	2.4-4.1	39	4.3	3.0-6.0	27	2.4	1.5-3.5	20	0.8	0.5-1.3	13	1.2	0.6-2.2	7	0.5	0.2-1.1
1983	63	3.1	2.4-4.0	33	3.8	2.6-5.5	30	2.6	1.7-3.8	21	0.9	0.6-1.5	12	1.3	0.7-2.3	9	0.7	0.3-1.3
1984	56	2.7	2.0-3.5	33	3.4	2.3-4.9	23	2.0	1.3-3.1	22	1.0	0.6-1.6	14	1.6	0.8-2.7	8	0.6	0.3-1.2
1985	80	3.8	3.0-4.7	48	4.9	3.6-6.6	32	2.9	1.9-4.1	18	0.9	0.5-1.5	12	1.2	0.6-2.2	6	0.7	0.2-1.4
1986	65	3.1	2.4-4.0	40	4.2	2.9-5.8	25	2.1	1.3-3.2	20	0.9	0.5-1.4	10	0.9	0.4-1.8	10	0.9	0.4-1.6
1987	57	2.7	2.0-3.5	31	3.1	2.1-4.5	26	2.2	1.4-3.3	15	0.7	0.4-1.1	7	0.7	0.3-1.5	8	0.6	0.3-1.3
1988	61	2.9	2.2-3.8	23	2.2	1.4-3.4	38	3.5	2.4-4.8	12	0.5	0.3-0.9	4	0.3	0.1-0.9	8	0.6	0.3-1.3
1989	53	2.5	1.8-3.2	31	2.9	2.0-4.2	22	2.0	1.2-3.0	21	1.0	0.6-1.6	13	1.4	0.7-2.5	8	0.6	0.3-1.3
1990	67	3.2	2.5-4.1	38	4.0	2.8-5.6	29	2.6	1.7-3.7	16	0.7	0.4-1.2	12	1.3	0.7-2.3	4	0.3	0.1-0.8
1991	70	3.4	2.6-4.3	39	4.3	3.0-6.0	31	2.7	1.8-3.9	24	1.1	0.7-1.7	17	1.8	1.0-3.0	7	0.6	0.2-1.2
1992	67	3.0	2.3-3.9	37	3.6	2.5-5.0	30	2.6	1.7-3.7	14	0.6	0.3-1.1	8	0.8	0.3-1.6	6	0.5	0.2-1.1
1993	58	2.7	2.1-3.5	34	3.6	2.5-5.1	24	1.9	1.2-2.9	13	0.6	0.3-1.0	7	0.8	0.3-1.6	6	0.5	0.2-1.1
1994	66	3.1	2.4-4.0	40	4.1	2.9-5.6	26	2.3	1.5-3.4	15	0.7	0.4-1.1	7	0.7	0.3-1.5	8	0.6	0.3-1.3
1995	62	2.9	2.2-3.7	34	3.3	2.3-4.6	28	2.5	1.6-3.6	10	0.5	0.2-0.9	6	0.6	0.2-1.3	4	0.4	0.1-0.9
1996	50	2.2	1.7-3.0	28	2.7	1.8-3.9	22	1.8	1.1-2.7	8	0.3	0.1-0.7	3	0.3	0.1-0.8	5	0.4	0.1-0.9
1997	61	2.8	2.2-3.7	31	3.0	2.1-4.3	30	2.6	1.8-3.8	15	0.7	0.4-1.2	6	0.7	0.2-1.5	9	0.7	0.3-1.4
1998	76	3.4	2.7-4.2	40	3.7	2.6-5.1	36	3.0	2.1-4.2	10	0.4	0.2-0.8	5	0.5	0.2-1.1	5	0.4	0.1-0.9
1999	53	2.3	1.7-3.0	29	2.6	1.7-3.8	24	2.0	1.3-3.0	5	0.2	0.1-0.5	2	0.2	0.0-0.8	3	0.3	0.1-0.8
2000	67	2.9	2.3-3.7	41	3.7	2.6-5.0	26	2.2	1.5-3.3	8	0.3	0.1-0.7	5	0.6	0.2-1.3	3	0.2	0.0-0.7
2001	53	2.3	1.7-3.0	32	2.9	2.0-4.1	21	1.7	1.0-2.6	5	0.2	0.1-0.5	2	0.2	0.0-0.7	3	0.2	0.0-0.7
2002	60	2.6	2.0-3.4	24	2.2	1.4-3.3	36	3.1	2.1-4.3	5	0.2	0.1-0.5	3	0.3	0.1-0.9	2	0.2	0.0-0.6
2003	66	2.9	2.3-3.7	34	3.2	2.2-4.5	32	2.7	1.9-3.9	16	0.7	0.4-1.1	7	0.7	0.3-1.4	9	0.7	0.3-1.4
2004	65	2.9	2.2-3.7	34	3.2	2.2-4.5	31	2.6	1.8-3.7	8	0.4	0.2-0.7	6	0.6	0.2-1.2	2	0.2	0.0-0.6
2005	59	2.6	2.0-3.4	35	3.4	2.3-4.7	24	1.9	1.2-2.9	12	0.5	0.3-0.9	6	0.6	0.2-1.4	6	0.4	0.2-1.0
2006	72	3.2	2.5-4.0	39	3.5	2.5-4.8	33	2.9	2.0-4.1	3	0.1	0.0-0.4	3	0.3	0.1-0.9	0	0.0	0.0-0.3
2007	66	2.9	2.3-3.8	39	3.7	2.6-5.0	27	2.3	1.5-3.4	12	0.5	0.3-0.9	6	0.6	0.2-1.3	6	0.5	0.2-1.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Hodgkin lymphoma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	74	3.3	2.6-4.2	37	3.5	2.4-4.8	37	3.2	2.2-4.4	8	0.4	0.2-0.7	4	0.4	0.1-1.1	4	0.3	0.1-0.9
2009	74	3.3	2.6-4.1	36	3.3	2.3-4.6	38	3.3	2.3-4.5	8	0.3	0.1-0.7	4	0.4	0.1-1.0	4	0.3	0.1-0.8
2010	65	2.8	2.1-3.6	33	3.0	2.0-4.2	32	2.6	1.8-3.8	8	0.3	0.1-0.7	5	0.4	0.1-1.1	3	0.2	0.0-0.7
2011	58	2.4	1.8-3.1	31	2.7	1.8-3.9	27	2.2	1.4-3.2	8	0.3	0.1-0.6	4	0.3	0.1-0.9	4	0.3	0.1-0.8
2012	53	2.3	1.7-3.0	29	2.6	1.7-3.8	24	2.1	1.3-3.1	5	0.2	0.1-0.5	3	0.3	0.1-0.8	2	0.1	0.0-0.5
2013	68	2.8	2.2-3.6	37	3.1	2.2-4.3	31	2.6	1.7-3.7	6	0.2	0.1-0.5	4	0.4	0.1-0.9	2	0.2	0.0-0.6
2014	63	2.6	2.0-3.4	41	3.5	2.5-4.7	22	1.8	1.1-2.8	10	0.4	0.2-0.7	5	0.4	0.1-1.0	5	0.4	0.1-0.9
2015	70	3.0	2.3-3.9	42	3.9	2.8-5.2	28	2.3	1.5-3.4	9	0.4	0.2-0.7	6	0.5	0.2-1.2	3	0.2	0.0-0.7
2016	75	3.2	2.5-4.0	42	3.6	2.6-4.9	33	2.8	1.9-4.0	5	0.2	0.1-0.4	4	0.3	0.1-0.9	1	0.0	0.0-0.4
2017	63	2.6	2.0-3.3	38	3.2	2.2-4.4	25	2.1	1.3-3.1	7	0.3	0.1-0.6	3	0.2	0.0-0.7	4	0.3	0.1-0.8
2018	50	2.2	1.6-2.9	28	2.6	1.7-3.7	22	1.8	1.1-2.8	4	0.1	0.0-0.3	3	0.2	0.0-0.6	1	0.1	0.0-0.4
2019	52	2.2	1.6-2.9	28	2.5	1.6-3.6	24	2.0	1.2-3.0	3	0.1	0.0-0.3	1	0.1	0.0-0.4	2	0.1	0.0-0.5
2020	64	2.7	2.1-3.5	37	3.3	2.3-4.6	27	2.3	1.5-3.4	5	0.2	0.1-0.4	2	0.1	0.0-0.6	3	0.2	0.0-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Non-Hodgkin lymphomas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	244	11.6	10.1-13.2	124	14.1	11.6-17.0	120	10.0	8.3-12.1	140	6.7	5.6-7.9	72	8.5	6.5-10.9	68	5.6	4.3-7.2
1977	221	10.5	9.1-12.1	111	12.2	9.9-14.8	110	9.3	7.6-11.3	127	6.1	5.1-7.3	62	7.2	5.5-9.4	65	5.4	4.1-6.9
1978	218	10.4	9.0-11.9	124	13.8	11.3-16.6	94	8.0	6.4-9.9	117	5.4	4.5-6.6	67	7.5	5.8-9.7	50	4.0	2.9-5.3
1979	259	12.3	10.8-14.0	122	13.7	11.3-16.4	137	11.4	9.5-13.6	118	5.7	4.7-6.8	60	7.0	5.3-9.1	58	4.7	3.5-6.1
1980	260	12.2	10.7-13.8	117	12.7	10.4-15.4	143	11.7	9.8-13.8	128	6.0	5.0-7.2	67	7.7	5.9-9.8	61	4.9	3.7-6.4
1981	301	14.2	12.6-16.0	132	14.7	12.2-17.5	169	13.6	11.5-15.8	125	5.9	4.9-7.1	53	6.2	4.6-8.2	72	5.6	4.4-7.1
1982	326	15.5	13.9-17.4	147	16.8	14.0-19.8	179	14.6	12.4-17.0	140	6.6	5.6-7.9	71	8.4	6.5-10.7	69	5.3	4.1-6.8
1983	290	13.8	12.2-15.5	152	17.6	14.8-20.8	138	11.0	9.2-13.1	135	6.4	5.4-7.7	63	7.2	5.4-9.2	72	5.8	4.5-7.4
1984	302	14.4	12.8-16.2	144	16.6	13.9-19.6	158	12.7	10.7-14.9	140	6.6	5.5-7.8	68	8.0	6.1-10.2	72	5.7	4.4-7.2
1985	328	15.4	13.8-17.2	164	18.9	16.0-22.1	164	13.1	11.1-15.3	163	7.7	6.5-9.0	81	9.2	7.3-11.6	82	6.4	5.1-8.0
1986	362	16.9	15.2-18.8	188	20.9	17.9-24.2	174	13.6	11.6-15.9	132	6.2	5.1-7.3	76	8.6	6.8-10.9	56	4.3	3.2-5.6
1987	368	17.5	15.8-19.5	188	21.6	18.5-25.0	180	14.6	12.5-17.0	159	7.3	6.2-8.6	84	9.4	7.5-11.8	75	5.8	4.5-7.3
1988	293	13.9	12.3-15.6	150	17.2	14.5-20.3	143	11.1	9.4-13.2	158	7.5	6.3-8.8	68	8.2	6.3-10.4	90	6.9	5.5-8.5
1989	316	15.2	13.6-17.0	151	16.5	14.0-19.5	165	13.6	11.6-16.0	139	6.7	5.6-7.9	70	8.3	6.4-10.5	69	5.5	4.3-7.0
1990	371	17.9	16.1-19.9	211	24.4	21.1-28.0	160	12.9	10.9-15.1	153	7.2	6.1-8.5	77	8.8	6.9-11.0	76	6.0	4.7-7.5
1991	393	18.6	16.8-20.6	207	23.2	20.1-26.7	186	14.7	12.7-17.1	143	6.9	5.8-8.1	73	8.6	6.7-10.8	70	5.5	4.2-6.9
1992	394	19.1	17.3-21.1	209	24.2	20.9-27.7	185	15.3	13.2-17.8	160	7.6	6.5-8.9	78	9.5	7.5-12.0	82	6.4	5.1-8.0
1993	366	17.3	15.6-19.2	206	22.3	19.3-25.7	160	13.1	11.1-15.4	143	6.7	5.6-7.9	74	8.3	6.5-10.5	69	5.5	4.2-6.9
1994	436	20.6	18.7-22.7	247	27.5	24.1-31.2	189	15.1	13.0-17.4	158	7.6	6.4-8.8	86	10.1	8.1-12.6	72	5.7	4.5-7.3
1995	420	19.8	17.9-21.8	231	25.6	22.3-29.1	189	15.3	13.1-17.6	141	6.6	5.6-7.8	77	9.1	7.2-11.4	64	5.0	3.8-6.4
1996	404	19.0	17.2-21.0	230	25.5	22.3-29.1	174	14.2	12.1-16.5	147	6.9	5.8-8.1	77	8.8	7.0-11.1	70	5.5	4.3-6.9
1997	407	19.0	17.2-20.9	213	23.3	20.2-26.7	194	15.4	13.3-17.7	149	6.9	5.9-8.2	82	9.5	7.5-11.8	67	5.3	4.1-6.7
1998	375	17.1	15.4-19.0	209	22.4	19.4-25.7	166	13.2	11.3-15.4	157	7.2	6.2-8.5	80	8.9	7.0-11.1	77	6.0	4.7-7.5
1999	395	17.8	16.1-19.7	201	21.6	18.7-24.9	194	15.2	13.1-17.5	138	6.3	5.3-7.4	77	8.6	6.8-10.8	61	4.7	3.6-6.1
2000	369	16.5	14.8-18.3	184	19.1	16.4-22.1	185	14.3	12.3-16.6	159	7.1	6.0-8.3	81	9.0	7.2-11.2	78	5.9	4.6-7.3
2001	387	17.1	15.4-18.9	215	22.3	19.3-25.5	172	13.0	11.1-15.2	144	6.4	5.4-7.5	87	9.6	7.7-11.9	57	4.2	3.1-5.4
2002	389	17.0	15.4-18.8	198	19.8	17.1-22.8	191	14.5	12.5-16.8	137	5.9	5.0-7.0	77	8.1	6.4-10.2	60	4.3	3.3-5.6
2003	393	17.2	15.5-19.0	182	18.6	15.9-21.5	211	16.3	14.1-18.6	149	6.5	5.5-7.6	71	7.4	5.7-9.3	78	5.8	4.5-7.2
2004	400	17.5	15.8-19.3	196	19.9	17.2-22.9	204	15.6	13.5-17.9	132	5.8	4.8-6.8	69	7.5	5.8-9.5	63	4.6	3.5-5.9
2005	433	18.9	17.1-20.7	213	21.5	18.7-24.6	220	16.8	14.6-19.2	111	4.8	4.0-5.8	52	5.4	4.0-7.1	59	4.4	3.4-5.7
2006	411	18.0	16.3-19.8	228	23.3	20.3-26.5	183	13.9	12.0-16.2	117	5.1	4.2-6.1	60	6.5	5.0-8.4	57	4.2	3.1-5.4
2007	387	16.7	15.1-18.4	198	20.0	17.3-23.0	189	14.4	12.4-16.7	118	5.1	4.2-6.1	62	6.5	5.0-8.4	56	4.1	3.1-5.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Non-Hodgkin lymphomas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	442	18.8	17.1-20.7	223	22.3	19.4-25.5	219	16.4	14.3-18.8	117	4.9	4.1-5.9	66	6.8	5.2-8.6	51	3.6	2.7-4.8
2009	468	20.0	18.2-21.9	229	23.0	20.1-26.3	239	18.1	15.8-20.6	103	4.2	3.4-5.1	54	5.6	4.2-7.4	49	3.4	2.5-4.5
2010	424	17.6	15.9-19.3	222	21.0	18.3-24.1	202	14.8	12.8-17.0	107	4.5	3.7-5.5	55	5.6	4.2-7.3	52	3.7	2.7-4.8
2011	470	19.2	17.5-21.1	242	22.8	20.0-25.9	228	16.3	14.2-18.6	108	4.4	3.6-5.3	54	5.3	3.9-6.9	54	3.8	2.8-5.0
2012	473	19.2	17.5-21.1	263	24.2	21.3-27.4	210	15.2	13.2-17.4	122	4.9	4.0-5.8	59	5.8	4.4-7.5	63	4.2	3.2-5.4
2013	517	20.7	18.9-22.6	283	26.2	23.2-29.5	234	16.4	14.3-18.7	127	5.0	4.1-5.9	69	6.4	4.9-8.1	58	3.9	2.9-5.1
2014	452	17.7	16.1-19.5	231	21.0	18.3-23.9	221	15.3	13.3-17.5	128	4.9	4.0-5.8	76	6.9	5.4-8.6	52	3.3	2.5-4.4
2015	492	18.7	17.1-20.5	271	23.4	20.7-26.5	221	15.2	13.2-17.4	116	4.4	3.7-5.3	73	6.6	5.2-8.3	43	2.9	2.1-3.9
2016	504	18.8	17.2-20.6	269	22.5	19.8-25.4	235	16.0	14.0-18.2	137	5.0	4.2-5.9	77	6.9	5.4-8.6	60	3.6	2.8-4.7
2017	477	17.6	16.0-19.3	256	21.3	18.7-24.1	221	14.9	12.9-17.1	120	4.4	3.6-5.3	67	5.8	4.5-7.4	53	3.3	2.4-4.4
2018	482	17.9	16.3-19.7	255	21.3	18.7-24.2	227	15.4	13.4-17.6	93	3.3	2.6-4.0	56	4.7	3.5-6.1	37	2.1	1.5-3.0
2019	543	19.6	17.9-21.3	302	24.4	21.7-27.4	241	15.8	13.8-18.0	133	4.6	3.8-5.5	81	6.8	5.4-8.5	52	3.0	2.2-3.9
2020	456	16.5	15.0-18.1	238	19.1	16.7-21.7	218	14.2	12.3-16.3	83	2.9	2.3-3.6	52	4.3	3.2-5.7	31	1.8	1.2-2.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Myeloma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	101	4.7	3.8-5.8	46	5.4	3.8-7.3	55	4.4	3.3-5.8	61	3.0	2.2-3.8	29	3.7	2.4-5.5	32	2.5	1.7-3.6
1977	102	4.7	3.8-5.8	52	5.7	4.2-7.7	50	4.0	2.9-5.3	69	3.0	2.3-3.9	36	3.8	2.6-5.3	33	2.5	1.7-3.6
1978	93	4.4	3.5-5.4	45	5.5	3.9-7.4	48	3.7	2.7-5.0	56	2.5	1.9-3.3	29	3.1	2.1-4.6	27	2.1	1.3-3.0
1979	106	4.9	4.0-6.0	61	7.1	5.4-9.3	45	3.5	2.5-4.7	63	3.0	2.3-3.8	26	2.8	1.8-4.3	37	2.9	2.0-4.1
1980	100	4.6	3.7-5.6	42	5.1	3.6-6.9	58	4.4	3.3-5.7	71	3.4	2.6-4.3	31	3.7	2.5-5.3	40	3.1	2.2-4.3
1981	118	5.2	4.3-6.3	62	6.6	5.0-8.6	56	4.2	3.2-5.5	53	2.4	1.8-3.1	24	2.7	1.7-4.1	29	2.2	1.5-3.2
1982	103	4.7	3.8-5.7	50	5.8	4.2-7.7	53	3.9	2.9-5.2	61	2.8	2.1-3.6	36	4.2	2.8-5.8	25	2.0	1.3-3.0
1983	130	6.2	5.1-7.4	68	8.0	6.2-10.3	62	5.1	3.9-6.6	75	3.5	2.7-4.4	32	3.9	2.6-5.5	43	3.2	2.3-4.4
1984	117	5.6	4.6-6.7	56	6.9	5.2-9.1	61	4.7	3.5-6.0	66	3.0	2.3-3.8	34	3.9	2.7-5.5	32	2.3	1.6-3.3
1985	113	5.3	4.3-6.3	61	7.1	5.4-9.2	52	4.2	3.1-5.5	74	3.3	2.6-4.2	41	5.0	3.5-6.9	33	2.4	1.6-3.4
1986	111	5.2	4.2-6.3	52	5.9	4.3-7.8	59	4.6	3.5-5.9	71	3.3	2.6-4.2	37	4.4	3.0-6.1	34	2.7	1.9-3.9
1987	112	5.3	4.4-6.4	65	7.9	6.1-10.2	47	3.7	2.7-4.9	85	4.0	3.2-5.0	54	6.6	4.9-8.7	31	2.4	1.6-3.4
1988	98	4.5	3.7-5.5	51	6.0	4.4-7.9	47	3.6	2.6-4.8	54	2.4	1.8-3.2	32	3.7	2.5-5.3	22	1.6	1.0-2.4
1989	112	5.3	4.3-6.4	49	6.0	4.4-7.9	63	4.9	3.8-6.4	66	3.0	2.3-3.9	30	3.4	2.3-4.9	36	2.8	1.9-3.9
1990	119	5.6	4.6-6.7	59	6.9	5.2-9.0	60	4.8	3.7-6.3	69	3.2	2.5-4.0	30	3.4	2.3-4.9	39	3.1	2.2-4.2
1991	113	5.3	4.4-6.4	43	5.1	3.7-6.9	70	5.6	4.3-7.1	69	3.1	2.4-4.0	34	4.0	2.7-5.6	35	2.7	1.9-3.8
1992	122	5.7	4.8-6.9	66	7.8	6.0-10.0	56	4.4	3.3-5.8	63	3.0	2.3-3.9	31	3.9	2.6-5.6	32	2.5	1.7-3.6
1993	109	5.2	4.2-6.2	53	6.3	4.7-8.3	56	4.4	3.3-5.8	76	3.6	2.8-4.5	38	4.6	3.2-6.3	38	3.0	2.1-4.1
1994	104	4.9	4.0-5.9	51	5.8	4.3-7.7	53	4.2	3.1-5.5	71	3.3	2.6-4.2	32	3.6	2.5-5.2	39	3.0	2.1-4.2
1995	117	5.4	4.5-6.5	57	6.8	5.1-8.8	60	4.7	3.6-6.0	66	3.1	2.4-3.9	37	4.4	3.1-6.1	29	2.3	1.5-3.3
1996	107	5.0	4.1-6.1	47	5.4	3.9-7.1	60	4.7	3.6-6.1	54	2.5	1.9-3.3	25	2.9	1.9-4.4	29	2.2	1.5-3.2
1997	119	5.6	4.6-6.7	59	6.7	5.1-8.7	60	4.7	3.6-6.1	54	2.5	1.9-3.3	24	2.8	1.8-4.1	30	2.3	1.6-3.4
1998	134	6.2	5.2-7.3	76	8.6	6.8-10.8	58	4.6	3.5-6.0	74	3.4	2.7-4.3	34	3.8	2.6-5.4	40	3.0	2.2-4.2
1999	137	6.2	5.2-7.3	53	5.7	4.2-7.4	84	6.4	5.1-8.0	59	2.7	2.0-3.4	23	2.6	1.6-3.9	36	2.7	1.9-3.8
2000	111	5.0	4.1-6.0	50	5.4	4.0-7.2	61	4.6	3.5-5.9	55	2.5	1.8-3.2	22	2.5	1.5-3.7	33	2.5	1.7-3.5
2001	151	6.7	5.7-7.8	77	8.1	6.4-10.2	74	5.6	4.4-7.1	61	2.7	2.1-3.5	35	3.8	2.6-5.3	26	1.9	1.3-2.8
2002	151	6.6	5.6-7.8	79	8.4	6.7-10.5	72	5.4	4.2-6.8	61	2.7	2.0-3.4	21	2.3	1.4-3.5	40	2.9	2.1-4.0
2003	156	6.8	5.8-8.0	77	8.3	6.6-10.4	79	5.9	4.6-7.3	69	3.0	2.3-3.8	29	3.1	2.1-4.5	40	2.8	2.0-3.9
2004	153	6.7	5.7-7.8	77	8.1	6.4-10.1	76	5.7	4.5-7.1	56	2.4	1.8-3.2	31	3.4	2.3-4.9	25	1.8	1.2-2.7
2005	157	6.8	5.8-8.0	67	7.0	5.4-9.0	90	6.7	5.4-8.3	58	2.5	1.9-3.3	31	3.3	2.2-4.7	27	2.0	1.3-2.9
2006	142	6.2	5.2-7.3	64	6.7	5.2-8.6	78	5.7	4.5-7.1	52	2.3	1.7-3.0	29	3.2	2.1-4.6	23	1.7	1.1-2.6
2007	154	6.8	5.7-7.9	88	9.1	7.3-11.2	66	4.9	3.8-6.2	77	3.4	2.7-4.2	38	4.1	2.9-5.6	39	2.8	2.0-3.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Myeloma**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	165	7.0	5.9-8.1	95	9.7	7.8-11.9	70	5.1	4.0-6.5	63	2.7	2.1-3.5	28	3.1	2.0-4.4	35	2.5	1.7-3.5
2009	186	7.8	6.7-9.1	108	10.9	8.9-13.1	78	5.7	4.5-7.2	54	2.3	1.7-3.0	33	3.4	2.3-4.8	21	1.5	0.9-2.3
2010	191	7.8	6.8-9.1	105	10.4	8.4-12.6	86	6.1	4.9-7.6	73	3.0	2.4-3.8	40	4.1	2.9-5.6	33	2.3	1.6-3.3
2011	218	8.9	7.8-10.2	126	12.2	10.1-14.6	92	6.7	5.4-8.3	58	2.4	1.8-3.1	32	3.2	2.2-4.5	26	1.9	1.2-2.8
2012	203	8.0	6.9-9.2	99	8.8	7.1-10.8	104	7.3	6.0-8.9	66	2.6	2.0-3.4	30	2.8	1.9-4.1	36	2.5	1.7-3.5
2013	208	8.1	7.0-9.3	104	9.5	7.7-11.6	104	7.2	5.9-8.8	64	2.4	1.9-3.1	41	3.8	2.7-5.1	23	1.5	1.0-2.3
2014	215	8.4	7.3-9.6	109	9.6	7.9-11.7	106	7.5	6.1-9.1	73	2.9	2.2-3.6	35	3.2	2.2-4.5	38	2.6	1.8-3.6
2015	232	8.8	7.7-10.0	112	9.7	8.0-11.8	120	8.1	6.7-9.7	85	3.1	2.5-3.9	45	3.9	2.8-5.3	40	2.6	1.8-3.5
2016	221	8.0	6.9-9.1	106	8.6	7.0-10.4	115	7.4	6.1-8.9	78	2.9	2.2-3.6	35	3.0	2.1-4.2	43	2.8	2.0-3.8
2017	227	8.2	7.2-9.4	117	9.8	8.0-11.8	110	7.0	5.7-8.5	71	2.5	2.0-3.2	32	2.8	1.9-3.9	39	2.4	1.7-3.3
2018	210	7.4	6.4-8.5	104	8.1	6.6-9.9	106	6.8	5.6-8.3	62	2.1	1.6-2.7	30	2.4	1.6-3.5	32	1.9	1.3-2.7
2019	241	8.5	7.4-9.7	134	10.6	8.8-12.6	107	6.9	5.6-8.4	71	2.4	1.8-3.0	37	3.0	2.1-4.2	34	1.9	1.3-2.7
2020	181	6.2	5.3-7.2	97	7.3	5.9-8.9	84	5.2	4.1-6.5	60	2.0	1.5-2.6	34	2.8	1.9-3.9	26	1.6	1.0-2.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Leukemias**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	238	11.8	10.3-13.5	129	15.2	12.5-18.2	109	9.6	7.8-11.6	161	8.3	7.0-9.7	87	10.9	8.6-13.6	74	6.4	5.0-8.1
1977	206	10.5	9.1-12.1	122	14.8	12.1-17.8	84	7.5	5.9-9.3	162	8.4	7.1-9.8	86	11.4	8.9-14.2	76	6.6	5.2-8.3
1978	263	13.1	11.5-14.8	164	19.5	16.4-22.9	99	8.5	6.9-10.5	144	7.1	5.9-8.4	83	9.9	7.8-12.4	61	5.1	3.9-6.7
1979	200	9.9	8.5-11.4	113	13.3	10.9-16.1	87	7.7	6.1-9.6	166	8.1	6.9-9.5	100	11.2	9.0-13.8	66	5.9	4.5-7.6
1980	262	12.8	11.2-14.5	134	15.7	13.0-18.7	128	11.0	9.1-13.1	128	6.4	5.3-7.6	79	9.7	7.6-12.2	49	4.4	3.2-5.9
1981	216	10.6	9.2-12.2	130	15.2	12.6-18.2	86	7.2	5.7-9.0	143	6.9	5.7-8.1	78	8.8	6.9-11.1	65	5.4	4.2-7.0
1982	239	11.6	10.1-13.2	144	17.3	14.5-20.4	95	7.9	6.3-9.7	137	6.6	5.5-7.9	79	9.4	7.4-11.8	58	4.7	3.5-6.2
1983	216	10.5	9.1-12.0	116	13.7	11.2-16.5	100	8.7	7.0-10.7	147	7.1	5.9-8.3	82	9.8	7.7-12.3	65	5.5	4.2-7.1
1984	251	12.2	10.7-13.8	126	15.2	12.6-18.3	125	10.2	8.4-12.2	155	7.4	6.3-8.7	80	9.7	7.6-12.2	75	5.9	4.6-7.4
1985	261	12.6	11.1-14.3	127	14.5	12.0-17.4	134	11.2	9.3-13.3	141	6.7	5.6-7.9	72	8.2	6.4-10.5	69	5.5	4.2-7.0
1986	265	12.9	11.4-14.6	141	16.7	13.9-19.8	124	10.2	8.5-12.3	156	7.5	6.3-8.8	81	9.5	7.5-11.9	75	6.1	4.7-7.7
1987	203	9.9	8.6-11.4	113	13.9	11.4-16.8	90	7.3	5.8-9.1	131	6.4	5.3-7.6	70	8.9	6.8-11.3	61	5.0	3.8-6.5
1988	210	10.0	8.7-11.5	131	15.4	12.8-18.4	79	6.7	5.3-8.4	125	6.0	4.9-7.1	83	9.7	7.6-12.1	42	3.4	2.4-4.6
1989	244	11.8	10.4-13.5	126	14.7	12.1-17.5	118	9.8	8.1-11.9	138	6.4	5.4-7.6	75	8.6	6.7-10.8	63	5.1	3.9-6.5
1990	244	11.7	10.3-13.3	128	15.0	12.4-17.9	116	9.4	7.8-11.4	140	6.7	5.6-7.9	79	9.5	7.5-11.9	61	4.8	3.7-6.2
1991	228	11.1	9.7-12.7	121	14.3	11.8-17.1	107	9.0	7.3-10.9	131	6.4	5.3-7.6	70	8.7	6.7-11.0	61	5.0	3.8-6.4
1992	241	11.6	10.1-13.1	127	14.8	12.3-17.7	114	9.7	7.9-11.7	126	6.1	5.1-7.3	65	8.1	6.2-10.4	61	5.0	3.8-6.5
1993	219	10.4	9.1-11.9	105	12.0	9.8-14.6	114	9.3	7.6-11.2	131	6.1	5.1-7.3	72	8.6	6.7-10.9	59	4.8	3.6-6.2
1994	248	12.0	10.5-13.6	121	14.3	11.8-17.1	127	10.5	8.7-12.6	148	7.0	5.9-8.3	76	8.8	6.9-11.1	72	5.8	4.5-7.4
1995	249	12.0	10.5-13.6	123	14.1	11.6-16.8	126	10.3	8.5-12.3	148	7.0	5.9-8.2	80	9.4	7.4-11.7	68	5.3	4.1-6.7
1996	227	10.7	9.3-12.2	127	14.6	12.2-17.5	100	8.1	6.6-9.9	135	6.3	5.3-7.5	68	7.6	5.9-9.7	67	5.5	4.2-7.0
1997	229	10.7	9.4-12.2	112	12.4	10.2-15.0	117	9.5	7.9-11.4	130	6.1	5.1-7.2	80	9.3	7.3-11.5	50	4.0	2.9-5.2
1998	275	12.7	11.3-14.3	142	15.5	13.1-18.4	133	10.6	8.8-12.6	127	5.9	4.9-7.0	60	6.8	5.2-8.8	67	5.3	4.1-6.7
1999	264	12.0	10.6-13.5	149	16.3	13.7-19.1	115	8.9	7.3-10.7	130	5.8	4.9-6.9	72	8.1	6.3-10.2	58	4.3	3.3-5.6
2000	275	12.3	10.9-13.9	151	16.1	13.6-18.9	124	9.7	8.0-11.6	143	6.3	5.3-7.5	66	7.1	5.5-9.1	77	5.9	4.7-7.5
2001	282	12.6	11.2-14.1	136	14.5	12.1-17.1	146	11.3	9.5-13.4	130	5.8	4.8-6.9	67	7.2	5.5-9.1	63	4.7	3.6-6.0
2002	244	10.9	9.6-12.4	136	14.3	12.0-17.0	108	8.5	6.9-10.3	122	5.3	4.4-6.4	68	7.4	5.7-9.4	54	4.0	3.0-5.3
2003	253	11.2	9.8-12.7	132	13.8	11.6-16.4	121	9.1	7.5-10.9	138	6.0	5.1-7.1	73	7.6	5.9-9.6	65	4.8	3.7-6.1
2004	265	11.7	10.3-13.2	153	16.4	13.9-19.2	112	8.5	7.0-10.2	137	6.1	5.1-7.2	75	8.3	6.5-10.4	62	4.7	3.6-6.1
2005	274	12.1	10.7-13.6	159	16.5	14.0-19.3	115	8.7	7.2-10.5	124	5.5	4.5-6.5	79	8.5	6.7-10.6	45	3.3	2.4-4.4
2006	287	12.9	11.4-14.4	170	17.8	15.2-20.8	117	9.1	7.5-11.0	126	5.5	4.6-6.6	70	7.5	5.8-9.4	56	4.2	3.1-5.4
2007	266	11.9	10.5-13.4	146	15.2	12.8-17.9	120	9.6	7.9-11.6	145	6.2	5.3-7.3	78	8.4	6.6-10.5	67	5.0	3.9-6.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Queens
Leukemias**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	292	12.8	11.4-14.4	159	16.5	14.0-19.3	133	10.5	8.8-12.5	147	6.4	5.4-7.6	77	8.4	6.6-10.5	70	5.2	4.0-6.6
2009	266	11.4	10.0-12.9	145	14.6	12.3-17.2	121	9.1	7.6-11.0	119	5.0	4.2-6.1	56	5.8	4.4-7.6	63	4.6	3.5-5.9
2010	309	13.2	11.7-14.8	159	16.1	13.7-18.8	150	11.3	9.5-13.3	123	5.3	4.4-6.3	68	7.0	5.4-8.9	55	4.1	3.1-5.4
2011	312	13.2	11.7-14.7	190	19.0	16.3-21.9	122	9.1	7.5-10.9	140	5.7	4.8-6.8	80	8.2	6.5-10.2	60	4.1	3.1-5.3
2012	315	13.0	11.6-14.6	179	16.9	14.4-19.6	136	10.2	8.5-12.1	154	6.3	5.3-7.4	95	9.5	7.7-11.7	59	4.0	3.0-5.2
2013	341	13.6	12.2-15.2	176	16.3	13.9-18.9	165	11.7	9.9-13.7	142	5.7	4.8-6.7	84	8.1	6.4-10.1	58	4.0	3.0-5.2
2014	365	14.5	13.0-16.1	185	17.0	14.6-19.7	180	12.8	11.0-14.9	115	4.6	3.8-5.5	58	5.6	4.2-7.2	57	3.9	2.9-5.1
2015	366	14.0	12.6-15.6	212	18.5	16.0-21.2	154	10.5	8.9-12.4	128	4.9	4.0-5.8	81	7.4	5.8-9.2	47	3.1	2.3-4.2
2016	379	14.4	13.0-16.0	214	18.3	15.9-21.0	165	11.3	9.6-13.2	141	5.3	4.4-6.3	69	6.2	4.8-7.9	72	4.7	3.7-6.0
2017	349	13.1	11.7-14.6	207	17.6	15.3-20.3	142	9.6	8.1-11.4	136	4.9	4.1-5.8	79	6.8	5.3-8.5	57	3.6	2.7-4.7
2018	324	12.7	11.3-14.2	174	15.3	13.1-17.8	150	10.8	9.0-12.8	120	4.4	3.6-5.2	70	5.9	4.6-7.5	50	3.3	2.4-4.5
2019	313	11.8	10.5-13.3	184	15.2	13.0-17.6	129	9.1	7.5-10.9	131	4.7	3.9-5.6	78	6.7	5.3-8.4	53	3.2	2.4-4.2
2020	295	11.5	10.1-12.9	183	15.9	13.6-18.4	112	7.8	6.4-9.5	112	4.0	3.3-4.9	69	6.0	4.6-7.6	43	2.6	1.9-3.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Richmond (Staten Island)
All Invasive Malignant Tumors**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1,157	440.1	414.4-466.9	603	558.7	512.1-608.2	554	371.3	340.5-404.0	634	247.4	228.0-267.9	357	341.9	305.1-381.6	277	186.8	165.2-210.4
1977	1,102	407.4	383.0-432.8	526	474.2	431.8-519.4	576	373.4	343.2-405.6	635	237.7	219.1-257.4	316	290.3	257.0-326.5	319	207.1	184.7-231.4
1978	1,131	400.6	377.1-425.2	566	484.0	442.5-528.0	565	353.3	324.4-384.0	598	218.1	200.6-236.7	324	290.1	257.5-325.3	274	172.9	152.8-194.9
1979	1,244	436.9	412.4-462.4	622	544.9	500.1-592.4	622	383.7	353.7-415.5	620	221.9	204.4-240.5	342	308.7	274.7-345.5	278	172.5	152.6-194.2
1980	1,244	419.9	396.4-444.5	643	540.0	496.3-586.4	601	354.2	326.0-384.1	662	226.6	209.3-245.0	343	302.2	268.8-338.2	319	187.0	166.8-209.0
1981	1,335	441.4	417.5-466.3	632	511.7	470.1-555.8	703	407.6	377.6-439.3	707	240.7	222.9-259.5	377	324.5	290.4-361.2	330	192.8	172.3-215.0
1982	1,289	425.8	402.4-450.2	677	559.6	515.6-606.1	612	351.6	323.9-381.0	643	216.3	199.6-234.0	338	288.6	256.8-323.0	305	173.9	154.8-194.8
1983	1,357	438.1	414.7-462.6	687	543.6	501.1-588.6	670	379.9	351.3-410.1	690	225.9	209.0-243.7	363	303.2	270.6-338.2	327	184.2	164.6-205.6
1984	1,437	461.4	437.4-486.3	695	555.0	511.9-600.5	742	415.9	386.1-447.3	724	234.1	217.0-252.1	386	309.0	277.3-343.2	338	187.8	168.1-209.1
1985	1,564	490.8	466.3-516.1	805	620.2	575.5-667.3	759	418.2	388.6-449.4	738	236.1	219.1-254.1	408	331.7	298.4-367.4	330	180.7	161.5-201.5
1986	1,530	468.0	444.4-492.4	731	548.1	506.9-591.6	799	427.2	397.8-458.3	750	233.5	216.8-251.1	390	302.7	271.6-336.0	360	192.3	172.8-213.5
1987	1,604	486.2	462.4-511.0	759	561.1	519.7-604.7	845	449.2	419.1-480.8	738	226.8	210.5-244.0	382	290.5	260.3-322.9	356	188.3	169.1-209.1
1988	1,740	521.8	497.2-547.3	807	598.3	555.5-643.4	933	486.1	455.1-518.6	769	238.0	221.2-255.6	409	321.5	289.3-356.0	360	186.8	167.9-207.2
1989	1,717	516.3	491.8-541.6	808	607.8	564.0-653.9	909	475.4	444.7-507.6	807	248.2	231.2-266.1	418	325.1	292.9-359.6	389	204.6	184.6-226.1
1990	1,689	493.4	469.9-517.8	770	543.3	503.6-585.0	919	468.0	438.0-499.5	796	239.4	222.8-256.8	406	303.7	273.2-336.4	390	199.8	180.3-220.7
1991	1,733	496.0	472.7-520.2	864	606.8	564.9-650.8	869	437.2	408.4-467.5	774	227.2	211.3-243.9	407	306.1	275.5-338.8	367	182.8	164.5-202.6
1992	1,957	548.3	524.0-573.3	994	688.5	644.4-734.7	963	474.0	444.3-505.0	844	243.0	226.8-260.1	418	308.5	278.2-341.0	426	207.4	188.1-228.2
1993	1,831	501.4	478.5-525.0	926	625.6	584.1-669.2	905	433.3	405.4-462.6	853	238.8	222.9-255.5	423	302.4	272.8-334.1	430	205.1	186.1-225.5
1994	1,813	483.9	461.8-506.9	857	546.4	509.1-585.6	956	448.3	420.3-477.8	848	232.1	216.6-248.3	416	289.6	261.1-320.1	432	202.2	183.5-222.2
1995	1,942	508.6	486.2-531.9	923	579.7	541.6-619.8	1,019	469.7	441.2-499.6	823	220.9	206.0-236.5	402	265.2	238.9-293.4	421	192.0	174.0-211.3
1996	1,929	493.8	471.9-516.4	920	559.2	522.6-597.7	1,009	457.3	429.4-486.5	833	218.3	203.7-233.7	401	258.8	233.2-286.3	432	192.5	174.8-211.6
1997	2,070	518.1	496.0-541.0	988	587.7	550.5-626.6	1,082	476.4	448.4-505.7	832	212.6	198.3-227.5	407	261.7	236.1-289.3	425	184.6	167.4-203.1
1998	2,206	535.2	513.0-558.0	1,091	631.4	593.5-671.0	1,115	478.3	450.6-507.3	864	213.4	199.4-228.2	450	277.8	251.9-305.5	414	174.8	158.4-192.6
1999	2,237	527.0	505.3-549.4	1,097	614.5	577.7-653.1	1,140	477.1	449.8-505.7	845	203.5	190.0-217.7	412	249.1	224.8-275.1	433	179.1	162.6-196.9
2000	2,260	520.9	499.6-542.9	1,201	655.9	618.3-695.0	1,059	431.8	406.1-458.7	834	196.9	183.7-210.8	420	250.2	226.1-276.0	414	166.1	150.4-183.0
2001	2,304	521.6	500.4-543.4	1,171	623.4	587.4-661.1	1,133	454.8	428.6-482.1	805	185.8	173.1-199.1	390	223.5	201.2-247.5	415	164.7	149.2-181.5
2002	2,334	516.3	495.5-537.8	1,191	605.6	570.9-641.8	1,143	448.0	422.3-474.9	844	189.7	177.1-203.0	415	228.0	206.0-251.6	429	165.7	150.3-182.3
2003	2,237	489.9	469.7-510.8	1,147	590.1	555.6-626.2	1,090	422.1	397.3-448.1	840	186.3	173.9-199.4	391	210.2	189.3-232.8	449	171.3	155.8-188.1
2004	2,339	502.5	482.2-523.5	1,138	568.7	535.2-603.6	1,201	460.7	434.8-487.7	864	190.2	177.6-203.4	434	236.6	214.2-260.6	430	160.4	145.5-176.5
2005	2,419	514.2	493.8-535.3	1,162	571.5	538.2-606.2	1,257	478.1	451.9-505.6	822	178.6	166.5-191.3	399	209.1	188.6-231.2	423	158.2	143.4-174.2
2006	2,481	522.4	501.8-543.6	1,255	609.6	575.4-645.2	1,226	463.2	437.4-490.2	893	192.6	180.1-205.8	455	237.6	215.7-261.1	438	161.8	146.9-177.9
2007	2,623	539.0	518.3-560.3	1,302	623.4	589.0-659.2	1,321	489.2	462.9-516.7	775	162.7	151.3-174.7	379	195.8	176.1-217.2	396	144.0	130.0-159.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

All Invasive Malignant Tumors

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	2,696	541.8	521.2-562.9	1,375	627.6	593.9-662.7	1,321	482.4	456.4-509.5	834	171.2	159.7-183.4	410	205.0	185.1-226.5	424	151.0	136.8-166.3
2009	2,706	534.3	514.0-555.2	1,324	591.7	559.3-625.5	1,382	496.9	470.7-524.3	800	162.1	150.9-173.9	387	186.9	168.1-207.1	413	145.7	131.8-160.7
2010	2,563	499.5	480.0-519.6	1,244	548.4	517.3-580.8	1,319	467.2	441.9-493.7	838	167.3	156.0-179.2	424	203.4	183.8-224.4	414	144.0	130.3-158.9
2011	2,761	526.5	506.7-547.0	1,385	595.3	563.3-628.6	1,376	479.7	454.1-506.3	771	149.8	139.2-161.0	373	176.5	158.5-196.0	398	133.2	120.1-147.3
2012	2,679	502.4	483.1-522.2	1,288	539.7	509.7-571.0	1,391	482.2	456.6-508.9	784	147.7	137.4-158.7	386	169.3	152.3-187.7	398	132.7	119.7-146.7
2013	2,876	526.9	507.4-547.0	1,417	575.6	545.1-607.5	1,459	495.8	470.1-522.7	846	156.7	146.1-167.9	447	195.9	177.6-215.6	399	130.2	117.5-144.0
2014	2,870	518.5	499.3-538.4	1,394	556.9	527.1-588.0	1,476	494.4	468.8-521.1	832	151.3	141.0-162.2	447	193.5	175.4-212.9	385	121.3	109.2-134.4
2015	2,881	516.1	496.9-535.8	1,425	566.2	536.3-597.4	1,456	485.3	460.0-511.7	853	150.4	140.3-161.1	428	177.9	160.9-196.2	425	132.7	120.1-146.4
2016	2,946	515.9	497.0-535.5	1,409	543.4	514.5-573.6	1,537	502.0	476.4-528.8	855	148.9	138.9-159.5	418	174.9	158.1-193.0	437	132.0	119.6-145.4
2017	2,996	514.9	496.2-534.3	1,476	554.9	526.1-585.0	1,520	488.4	463.3-514.4	862	147.3	137.4-157.7	417	167.2	151.1-184.6	445	134.1	121.6-147.6
2018	2,972	502.6	484.2-521.6	1,441	533.6	505.6-562.8	1,531	482.4	457.8-508.2	868	143.6	134.0-153.7	419	161.9	146.3-178.6	449	130.6	118.5-143.7
2019	3,162	530.6	511.7-550.0	1,536	564.0	535.3-593.9	1,626	512.6	487.1-539.2	835	138.5	129.1-148.4	422	161.8	146.4-178.6	413	120.5	108.9-133.1
2020	2,763	458.9	441.4-477.0	1,327	475.5	449.5-502.8	1,436	449.5	425.6-474.4	801	130.8	121.7-140.4	380	145.6	131.0-161.5	421	120.9	109.3-133.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Oral cavity and pharynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	39	14.7	10.3-20.2	29	25.9	16.6-38.2	10	6.6	3.1-12.2	19	6.5	3.9-10.4	11	8.4	3.9-15.7	8	5.0	2.1-9.9
1977	47	17.4	12.7-23.2	34	30.1	20.3-42.8	13	8.3	4.4-14.1	20	7.8	4.7-12.1	15	15.7	8.4-26.2	5	2.9	0.9-7.0
1978	31	10.8	7.3-15.4	25	20.3	12.8-30.4	6	4.0	1.4-8.7	14	4.7	2.5-7.9	6	5.2	1.7-11.4	8	4.8	2.0-9.5
1979	31	10.7	7.2-15.3	20	16.6	9.7-26.2	11	7.2	3.5-12.9	13	4.3	2.3-7.4	9	6.5	2.7-13.0	4	2.7	0.7-6.8
1980	43	14.2	10.2-19.3	28	22.7	14.6-33.3	15	8.8	4.9-14.6	22	7.9	4.9-12.0	16	14.1	7.8-23.2	6	3.9	1.4-8.4
1981	31	9.6	6.5-13.8	19	12.0	7.2-19.0	12	6.8	3.5-12.0	17	5.5	3.1-8.8	10	6.9	3.3-12.9	7	4.0	1.6-8.3
1982	40	12.7	9.0-17.3	25	19.5	12.3-29.2	15	8.1	4.5-13.4	9	3.1	1.4-5.9	5	4.2	1.3-9.6	4	2.5	0.7-6.2
1983	35	11.0	7.6-15.5	23	18.4	11.1-28.4	12	6.9	3.5-12.1	11	3.3	1.6-6.1	10	7.9	3.5-14.9	1	0.4	0.0-2.8
1984	39	11.9	8.4-16.3	21	14.6	8.9-22.6	18	9.4	5.6-15.0	14	4.2	2.2-7.1	10	7.7	3.4-14.7	4	2.2	0.6-5.7
1985	55	16.1	12.0-21.0	40	26.1	18.4-36.0	15	7.8	4.3-13.0	13	3.9	2.0-6.6	10	6.9	3.2-12.9	3	1.7	0.3-4.9
1986	43	12.4	9.0-16.8	31	20.9	13.9-30.0	12	6.2	3.2-11.0	8	2.4	1.0-4.8	3	1.9	0.4-5.8	5	2.7	0.9-6.4
1987	51	15.2	11.2-20.0	30	21.6	14.2-31.4	21	10.7	6.6-16.4	14	4.3	2.3-7.3	11	8.8	4.2-16.0	3	1.4	0.3-4.3
1988	59	17.2	13.0-22.2	46	30.4	22.0-40.9	13	6.7	3.5-11.4	22	6.6	4.1-10.1	14	10.5	5.5-17.9	8	3.9	1.7-7.8
1989	45	13.4	9.7-18.0	30	23.1	15.0-33.6	15	7.7	4.3-12.7	12	3.5	1.8-6.2	11	7.4	3.5-13.7	1	0.5	0.0-3.0
1990	47	13.6	10.0-18.1	34	22.7	15.6-31.8	13	6.3	3.4-10.9	13	4.0	2.1-6.8	10	8.0	3.5-15.1	3	1.6	0.3-4.6
1991	44	13.1	9.5-17.5	29	21.5	14.0-31.4	15	8.2	4.6-13.5	14	4.0	2.2-6.8	11	8.0	3.8-14.5	3	1.5	0.3-4.4
1992	42	11.8	8.5-16.0	30	20.2	13.3-29.2	12	5.8	3.0-10.1	14	4.1	2.2-6.9	9	6.6	2.8-12.7	5	2.5	0.8-5.9
1993	40	10.9	7.8-14.8	28	17.4	11.4-25.3	12	5.6	2.9-9.9	13	3.6	1.9-6.1	10	6.8	3.1-12.7	3	1.4	0.3-4.2
1994	31	8.0	5.5-11.4	22	12.8	8.0-19.4	9	4.2	1.9-8.1	7	1.8	0.7-3.8	5	2.7	0.9-6.5	2	0.9	0.1-3.3
1995	37	9.6	6.8-13.3	27	16.2	10.6-23.7	10	4.7	2.2-8.6	3	0.8	0.2-2.2	2	1.1	0.1-4.0	1	0.5	0.0-2.6
1996	44	11.1	8.1-14.9	28	15.4	10.2-22.3	16	7.2	4.1-11.7	8	2.0	0.9-4.0	5	3.1	1.0-7.1	3	1.3	0.3-3.9
1997	39	9.9	7.0-13.6	21	13.9	8.4-21.3	18	8.0	4.7-12.7	9	2.3	1.0-4.3	6	4.0	1.4-8.7	3	1.3	0.3-3.9
1998	40	9.6	6.8-13.1	29	15.5	10.3-22.4	11	4.7	2.3-8.4	7	1.7	0.7-3.5	3	1.4	0.3-4.3	4	1.6	0.4-4.2
1999	37	8.6	6.0-11.8	28	14.6	9.6-21.2	9	3.8	1.7-7.2	9	2.1	1.0-4.1	4	2.1	0.6-5.5	5	2.1	0.7-5.0
2000	34	7.7	5.3-10.8	24	12.4	7.9-18.6	10	3.9	1.9-7.3	5	1.1	0.4-2.7	3	1.9	0.4-5.5	2	0.8	0.1-3.0
2001	38	8.6	6.1-11.9	21	11.1	6.8-17.1	17	7.0	4.1-11.2	8	1.8	0.8-3.6	4	2.5	0.7-6.2	4	1.5	0.4-3.9
2002	47	10.1	7.4-13.4	24	11.6	7.4-17.5	23	9.0	5.7-13.5	5	1.1	0.3-2.5	4	1.9	0.5-4.9	1	0.4	0.0-2.4
2003	45	9.4	6.8-12.6	32	14.7	10.0-20.9	13	4.7	2.5-8.2	9	2.1	1.0-4.0	6	3.6	1.3-7.7	3	1.1	0.2-3.4
2004	48	10.2	7.5-13.6	26	12.8	8.2-18.9	22	8.7	5.5-13.3	16	3.5	2.0-5.7	7	3.8	1.5-7.8	9	3.5	1.6-6.7
2005	55	11.4	8.6-14.9	40	18.7	13.2-25.6	15	5.8	3.2-9.6	10	2.1	1.0-3.9	6	2.9	1.0-6.3	4	1.4	0.4-3.8
2006	49	10.0	7.4-13.3	35	15.6	10.8-21.9	14	5.0	2.7-8.5	12	2.7	1.4-4.7	7	3.8	1.5-7.8	5	1.9	0.6-4.5
2007	32	6.3	4.3-8.9	21	9.3	5.7-14.4	11	4.1	2.0-7.5	9	1.7	0.8-3.3	5	2.1	0.6-5.0	4	1.4	0.4-3.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Oral cavity and pharynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	49	9.5	7.0-12.7	30	12.5	8.3-18.0	19	6.9	4.1-10.9	3	0.6	0.1-1.7	3	1.2	0.2-3.6	0	0.0	0.0-1.6
2009	34	6.9	4.8-9.7	27	12.0	7.8-17.7	7	2.4	1.0-5.2	16	3.1	1.7-5.0	11	4.5	2.2-8.3	5	1.8	0.6-4.4
2010	45	8.5	6.2-11.5	27	10.7	7.0-15.7	18	6.4	3.7-10.3	11	2.1	1.0-3.9	6	2.3	0.8-5.2	5	1.9	0.6-4.5
2011	58	10.6	8.0-13.9	39	15.3	10.8-21.2	19	6.6	3.9-10.4	8	1.6	0.7-3.2	6	2.7	1.0-6.0	2	0.7	0.1-2.7
2012	56	10.1	7.6-13.3	42	17.1	12.2-23.3	14	4.5	2.4-7.8	14	2.6	1.4-4.4	10	4.2	2.0-7.8	4	1.4	0.4-3.8
2013	59	10.3	7.8-13.4	39	14.4	10.2-20.0	20	6.8	4.1-10.7	6	1.1	0.4-2.4	2	1.0	0.1-3.4	4	1.1	0.3-3.2
2014	60	10.5	8.0-13.6	35	13.4	9.2-18.9	25	8.2	5.3-12.3	6	1.0	0.4-2.3	3	1.0	0.2-3.2	3	1.0	0.2-3.2
2015	51	8.7	6.4-11.6	38	14.2	9.9-19.8	13	4.2	2.2-7.4	12	1.9	1.0-3.5	9	3.1	1.4-6.2	3	0.9	0.2-2.9
2016	56	9.7	7.3-12.8	38	14.5	10.1-20.1	18	5.6	3.3-9.1	13	2.1	1.1-3.8	9	3.2	1.4-6.4	4	1.2	0.3-3.3
2017	61	10.1	7.7-13.1	47	17.3	12.6-23.3	14	4.1	2.2-7.2	7	1.1	0.4-2.4	4	1.4	0.4-3.9	3	0.8	0.2-2.7
2018	70	11.5	8.9-14.7	46	16.4	11.8-22.1	24	7.1	4.5-10.8	20	3.0	1.8-4.7	15	5.1	2.8-8.7	5	1.3	0.4-3.4
2019	62	9.8	7.5-12.7	37	12.6	8.7-17.6	25	7.3	4.7-11.1	10	1.6	0.8-3.1	9	3.4	1.5-6.7	1	0.2	0.0-1.9
2020	53	8.7	6.4-11.5	42	14.6	10.4-20.1	11	3.7	1.8-6.9	8	1.1	0.5-2.4	4	1.1	0.3-3.3	4	1.1	0.3-3.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Esophagus

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	23	8.4	5.2-12.7	16	13.4	7.3-22.4	7	4.7	1.9-9.7	14	5.4	2.9-9.0	8	7.5	2.9-15.1	6	4.0	1.5-8.8
1977	16	6.1	3.4-9.9	12	11.9	5.8-21.0	4	2.4	0.6-6.2	15	5.6	3.1-9.4	10	9.4	4.0-17.9	5	3.3	1.1-7.6
1978	16	5.8	3.3-9.4	11	9.5	4.6-17.2	5	3.2	1.0-7.5	17	6.5	3.7-10.4	12	11.9	6.0-20.9	5	3.1	1.0-7.2
1979	17	6.0	3.4-9.6	11	9.6	4.5-17.5	6	3.8	1.4-8.2	15	5.1	2.8-8.5	12	10.2	4.9-18.3	3	1.8	0.4-5.2
1980	8	2.9	1.2-5.7	4	4.5	1.2-10.8	4	2.2	0.6-5.8	3	1.2	0.2-3.3	3	3.2	0.6-8.7	0	0.0	0.0-2.1
1981	15	5.0	2.8-8.3	9	7.0	3.1-13.5	6	3.4	1.2-7.5	14	4.4	2.4-7.4	9	6.8	3.0-13.1	5	2.6	0.8-6.3
1982	12	4.5	2.3-7.8	7	7.5	2.8-15.3	5	3.2	1.0-7.3	10	3.6	1.7-6.5	7	7.0	2.6-14.3	3	1.8	0.4-5.1
1983	12	4.2	2.1-7.3	9	7.8	3.3-15.2	3	1.8	0.4-5.2	9	3.4	1.5-6.4	7	8.2	3.0-16.6	2	1.2	0.1-4.3
1984	10	3.1	1.5-5.8	8	6.6	2.7-13.2	2	1.1	0.1-3.9	9	3.1	1.4-5.8	6	5.0	1.6-11.3	3	1.8	0.4-5.1
1985	17	5.1	3.0-8.3	11	8.0	3.8-14.7	6	3.2	1.2-7.0	12	3.7	1.9-6.5	12	9.5	4.7-16.9	0	0.0	0.0-2.0
1986	17	5.2	3.0-8.3	10	7.2	3.4-13.3	7	3.8	1.5-7.9	16	4.7	2.7-7.7	10	7.3	3.4-13.6	6	3.2	1.1-6.9
1987	16	4.9	2.8-8.0	10	7.4	3.4-13.7	6	3.3	1.2-7.1	12	3.6	1.8-6.3	8	5.8	2.4-11.6	4	2.1	0.6-5.4
1988	11	3.6	1.8-6.3	8	6.9	2.8-13.6	3	1.5	0.3-4.5	18	5.9	3.5-9.2	14	11.8	6.2-19.8	4	2.3	0.6-5.7
1989	17	5.0	2.9-8.0	15	10.8	5.7-18.4	2	1.0	0.1-3.7	13	3.7	1.9-6.4	9	6.9	2.8-13.6	4	2.1	0.6-5.5
1990	17	5.0	2.9-8.0	14	9.8	5.1-16.7	3	1.6	0.3-4.6	15	4.5	2.5-7.4	12	8.4	4.2-14.8	3	1.6	0.3-4.7
1991	15	4.1	2.3-6.8	11	6.9	3.4-12.4	4	2.0	0.5-5.1	10	3.0	1.4-5.5	7	5.4	2.0-11.3	3	1.5	0.3-4.4
1992	11	3.2	1.6-5.8	8	7.1	2.9-13.9	3	1.3	0.3-4.0	19	5.4	3.3-8.5	15	10.7	5.8-17.8	4	1.9	0.5-4.9
1993	19	5.3	3.2-8.2	16	10.1	5.7-16.5	3	1.3	0.3-4.0	13	3.5	1.9-6.1	12	8.3	4.1-14.7	1	0.5	0.0-2.6
1994	18	4.8	2.8-7.5	12	6.9	3.5-12.1	6	2.8	1.0-6.2	21	5.5	3.4-8.5	16	9.4	5.3-15.3	5	2.2	0.7-5.3
1995	13	3.5	1.8-5.9	9	5.0	2.3-9.5	4	1.9	0.5-4.8	11	2.9	1.4-5.1	7	3.7	1.5-7.8	4	1.8	0.5-4.6
1996	15	4.0	2.2-6.6	12	7.3	3.7-12.8	3	1.3	0.3-4.0	13	3.4	1.8-5.8	10	5.9	2.8-10.8	3	1.3	0.3-3.9
1997	17	4.3	2.5-6.8	12	7.7	3.8-13.6	5	2.2	0.7-5.1	12	3.1	1.6-5.3	10	6.4	3.0-11.8	2	0.8	0.1-3.1
1998	19	4.8	2.9-7.4	12	7.3	3.7-12.7	7	2.8	1.1-6.0	18	4.5	2.7-7.1	14	8.2	4.4-13.9	4	1.6	0.4-4.1
1999	12	2.8	1.5-4.9	11	6.3	3.1-11.4	1	0.4	0.0-2.5	14	3.3	1.8-5.6	10	6.0	2.8-11.0	4	1.7	0.5-4.4
2000	9	2.1	0.9-3.9	7	3.6	1.4-7.5	2	0.8	0.1-3.0	10	2.4	1.1-4.4	9	5.4	2.4-10.1	1	0.4	0.0-2.4
2001	15	3.5	1.9-5.7	11	6.6	3.2-11.8	4	1.6	0.4-4.2	9	2.0	0.9-3.9	8	4.2	1.8-8.3	1	0.4	0.0-2.4
2002	18	4.0	2.3-6.3	15	7.6	4.2-12.6	3	1.1	0.2-3.4	19	4.1	2.5-6.5	15	7.6	4.2-12.6	4	1.5	0.4-4.0
2003	16	3.5	2.0-5.7	8	3.9	1.6-7.8	8	3.2	1.4-6.4	14	3.2	1.7-5.3	10	5.0	2.4-9.2	4	1.6	0.4-4.2
2004	25	5.3	3.4-7.9	16	7.8	4.4-12.7	9	3.4	1.6-6.6	21	4.5	2.7-6.8	18	9.2	5.4-14.8	3	1.1	0.2-3.4
2005	21	4.4	2.7-6.8	9	4.5	2.0-8.7	12	4.5	2.3-8.0	28	6.0	4.0-8.8	20	10.0	6.0-15.5	8	2.9	1.3-5.9
2006	22	4.6	2.9-7.0	17	8.2	4.7-13.3	5	1.7	0.6-4.2	19	4.0	2.4-6.2	14	6.9	3.7-11.8	5	1.8	0.6-4.3
2007	22	4.7	2.9-7.1	18	8.9	5.2-14.1	4	1.4	0.4-3.8	16	3.1	1.8-5.2	12	5.3	2.7-9.5	4	1.5	0.4-4.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Esophagus

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	29	5.3	3.5-7.7	20	7.4	4.5-11.7	9	3.4	1.5-6.5	23	4.6	2.9-7.0	16	7.2	4.0-11.8	7	2.5	1.0-5.4
2009	22	4.3	2.7-6.6	15	6.3	3.4-10.5	7	2.5	1.0-5.3	21	4.2	2.6-6.5	14	6.4	3.4-10.9	7	2.5	1.0-5.3
2010	18	3.3	1.9-5.3	14	6.0	3.2-10.3	4	1.3	0.3-3.6	17	3.0	1.7-4.8	12	4.5	2.3-8.1	5	1.7	0.5-4.1
2011	17	3.4	2.0-5.6	14	6.3	3.4-10.7	3	1.1	0.2-3.3	11	2.3	1.1-4.2	6	3.1	1.1-6.7	5	1.8	0.6-4.3
2012	15	2.7	1.5-4.6	13	5.4	2.8-9.4	2	0.5	0.1-2.3	15	2.7	1.5-4.5	14	5.5	2.9-9.5	1	0.4	0.0-2.4
2013	17	3.0	1.7-4.9	14	5.8	3.1-10.0	3	0.9	0.2-2.9	16	2.8	1.6-4.7	14	5.6	3.0-9.7	2	0.7	0.1-2.7
2014	19	3.4	2.0-5.5	13	5.0	2.6-8.8	6	2.2	0.8-5.0	21	3.8	2.3-6.0	18	7.3	4.2-11.8	3	1.1	0.2-3.4
2015	22	3.8	2.4-5.9	18	7.3	4.2-11.8	4	1.2	0.3-3.3	21	3.9	2.4-6.0	14	5.5	3.0-9.5	7	2.5	1.0-5.3
2016	34	5.5	3.8-7.8	24	8.2	5.2-12.4	10	3.3	1.5-6.2	18	3.0	1.7-4.8	15	5.1	2.8-8.7	3	0.9	0.2-3.0
2017	27	4.5	2.9-6.6	21	8.1	5.0-12.6	6	1.7	0.6-4.0	20	3.3	2.0-5.1	15	5.7	3.1-9.6	5	1.4	0.4-3.6
2018	27	4.1	2.7-6.1	22	7.2	4.4-11.1	5	1.5	0.5-3.8	22	3.5	2.2-5.4	16	5.5	3.1-9.2	6	1.8	0.6-4.1
2019	31	5.1	3.4-7.3	25	9.0	5.8-13.6	6	1.9	0.7-4.4	12	1.8	0.9-3.3	9	3.0	1.3-6.0	3	0.9	0.2-2.8
2020	23	3.4	2.1-5.2	19	6.3	3.7-10.1	4	1.0	0.3-3.0	16	2.5	1.4-4.2	15	5.1	2.8-8.7	1	0.3	0.0-2.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Richmond (Staten Island)
Stomach**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	32	13.0	8.8-18.3	20	20.4	12.1-31.8	12	8.2	4.2-14.2	29	12.9	8.6-18.4	16	19.9	11.1-32.3	13	9.3	4.9-15.8
1977	33	13.9	9.5-19.5	20	24.2	14.4-37.5	13	8.6	4.5-14.6	36	15.1	10.5-20.9	23	26.9	16.6-40.6	13	8.8	4.7-15.1
1978	38	13.8	9.7-19.1	18	15.7	8.8-25.4	20	12.5	7.6-19.3	28	10.9	7.2-15.8	11	11.9	5.5-21.5	17	10.8	6.3-17.2
1979	38	14.9	10.5-20.4	19	20.2	11.8-31.7	19	12.3	7.4-19.2	26	10.1	6.6-14.8	15	15.1	8.1-25.1	11	7.4	3.7-13.1
1980	36	13.0	9.0-18.1	20	17.2	10.1-27.1	16	10.1	5.8-16.4	23	8.5	5.3-12.7	12	11.2	5.4-19.9	11	6.6	3.3-11.8
1981	42	14.6	10.4-19.8	21	18.4	11.0-28.6	21	12.2	7.5-18.7	31	11.6	7.8-16.5	19	19.8	11.5-31.1	12	7.2	3.7-12.5
1982	29	9.8	6.5-14.1	14	12.8	6.7-21.7	15	8.4	4.7-13.9	13	4.6	2.4-7.8	7	7.1	2.7-14.6	6	3.5	1.2-7.5
1983	40	13.7	9.7-18.6	24	22.1	13.8-33.2	16	8.7	4.9-14.2	28	9.8	6.4-14.1	16	15.4	8.5-25.1	12	6.6	3.4-11.6
1984	33	11.2	7.7-15.8	17	14.8	8.3-24.1	16	8.8	5.0-14.3	36	11.4	7.9-15.9	19	15.9	9.2-25.2	17	9.0	5.2-14.5
1985	32	10.0	6.8-14.2	19	15.3	8.8-24.4	13	6.9	3.7-11.9	15	5.2	2.9-8.5	9	7.1	3.1-13.7	6	3.4	1.2-7.4
1986	33	10.2	7.0-14.4	18	13.9	8.1-22.2	15	7.7	4.3-12.8	20	6.3	3.8-9.8	6	5.2	1.8-11.4	14	7.3	4.0-12.2
1987	36	10.9	7.6-15.2	24	17.3	10.9-26.0	12	6.3	3.2-11.0	15	4.6	2.5-7.6	9	6.0	2.7-11.6	6	3.1	1.1-6.8
1988	50	15.5	11.4-20.5	29	21.3	13.9-31.1	21	10.7	6.6-16.3	36	11.5	8.0-15.9	19	14.6	8.6-23.0	17	8.8	5.1-14.1
1989	38	11.9	8.4-16.4	19	13.4	7.8-21.2	19	10.0	6.0-15.6	28	8.9	5.9-12.8	16	11.6	6.5-18.9	12	6.4	3.3-11.2
1990	41	13.2	9.4-17.8	25	19.5	12.4-29.0	16	8.2	4.7-13.3	32	10.4	7.1-14.7	19	15.1	8.7-23.8	13	6.9	3.6-11.7
1991	45	13.5	9.8-18.1	32	22.3	15.0-31.8	13	6.6	3.5-11.3	23	6.8	4.3-10.3	14	10.0	5.2-17.0	9	4.4	2.0-8.4
1992	39	11.4	8.0-15.5	18	13.5	7.7-21.6	21	10.3	6.4-15.8	33	9.7	6.6-13.5	16	11.9	6.5-19.6	17	8.1	4.7-13.0
1993	45	12.4	9.0-16.7	27	19.4	12.4-28.6	18	8.4	5.0-13.4	27	7.6	5.0-11.0	17	12.6	7.1-20.4	10	4.8	2.3-8.9
1994	31	8.7	5.9-12.4	19	13.3	7.8-21.0	12	5.5	2.9-9.7	20	5.9	3.6-9.1	11	9.4	4.5-16.7	9	4.3	1.9-8.1
1995	43	11.6	8.4-15.6	22	15.0	9.1-22.9	21	9.4	5.8-14.4	26	7.2	4.7-10.5	14	9.3	4.9-15.8	12	5.4	2.8-9.4
1996	51	13.3	9.9-17.5	30	17.7	11.9-25.4	21	9.3	5.8-14.3	20	5.3	3.3-8.2	13	8.7	4.6-14.9	7	3.2	1.3-6.5
1997	47	12.1	8.9-16.0	27	17.4	11.4-25.4	20	8.5	5.2-13.2	26	6.7	4.4-9.8	17	11.2	6.4-17.9	9	3.8	1.7-7.3
1998	48	12.1	8.9-16.0	25	15.5	9.9-23.0	23	9.4	5.9-14.2	27	6.7	4.4-9.8	18	10.6	6.2-16.7	9	3.6	1.7-7.0
1999	49	12.0	8.8-15.8	25	14.6	9.3-21.6	24	9.8	6.3-14.6	22	5.4	3.4-8.2	9	5.5	2.4-10.5	13	5.3	2.8-9.1
2000	35	8.0	5.5-11.1	17	8.7	5.0-14.1	18	7.1	4.2-11.2	14	3.2	1.8-5.4	8	4.7	1.9-9.4	6	2.4	0.9-5.3
2001	52	12.0	8.9-15.7	25	14.4	9.2-21.4	27	10.8	7.1-15.8	17	3.8	2.2-6.0	11	5.4	2.6-9.8	6	2.4	0.9-5.3
2002	39	8.8	6.2-12.0	27	13.9	9.1-20.4	12	4.6	2.4-8.1	24	5.3	3.4-7.9	11	5.2	2.5-9.4	13	4.9	2.6-8.5
2003	42	9.4	6.7-12.7	24	12.9	8.1-19.4	18	7.1	4.2-11.3	16	3.5	2.0-5.8	8	4.1	1.7-8.3	8	3.1	1.3-6.2
2004	49	10.8	7.9-14.2	32	15.8	10.7-22.5	17	6.3	3.7-10.3	28	6.3	4.2-9.1	12	7.1	3.6-12.4	16	5.9	3.4-9.8
2005	53	11.3	8.5-14.9	30	15.6	10.4-22.4	23	8.1	5.1-12.3	30	6.6	4.5-9.5	14	7.3	3.9-12.3	16	5.5	3.1-9.1
2006	36	7.8	5.4-10.8	19	10.0	5.9-15.7	17	6.4	3.7-10.4	20	4.1	2.5-6.3	12	5.8	2.9-10.2	8	2.9	1.2-5.9
2007	49	10.2	7.6-13.6	30	15.2	10.1-21.9	19	6.6	4.0-10.5	23	5.0	3.2-7.6	11	6.3	3.1-11.2	12	4.2	2.2-7.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Stomach

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	43	8.8	6.3-11.9	25	11.6	7.4-17.3	18	6.5	3.9-10.5	19	4.0	2.4-6.3	11	5.8	2.8-10.3	8	2.9	1.2-5.9
2009	40	8.1	5.7-11.0	26	12.7	8.2-18.8	14	4.9	2.7-8.4	25	5.3	3.4-7.9	15	8.7	4.8-14.2	10	3.4	1.6-6.3
2010	48	9.5	7.0-12.7	31	14.1	9.5-20.2	17	5.7	3.3-9.3	13	2.4	1.3-4.2	4	2.0	0.5-5.1	9	2.9	1.3-5.6
2011	32	6.5	4.4-9.3	16	7.8	4.4-12.8	16	5.5	3.1-9.1	22	4.1	2.6-6.3	14	6.0	3.2-10.3	8	2.6	1.1-5.3
2012	48	9.4	6.9-12.5	29	11.8	7.8-17.2	19	7.5	4.4-11.8	18	3.4	2.0-5.4	13	5.6	2.9-9.7	5	1.6	0.5-4.0
2013	38	6.8	4.8-9.4	19	7.6	4.5-12.0	19	5.8	3.5-9.4	17	2.9	1.7-4.8	13	5.1	2.7-8.9	4	1.2	0.3-3.4
2014	52	9.5	7.0-12.5	32	13.4	9.1-19.2	20	6.4	3.9-10.1	26	4.2	2.7-6.2	14	5.7	3.0-9.7	12	3.2	1.6-5.9
2015	54	9.6	7.2-12.7	29	11.9	7.8-17.3	25	7.9	5.1-11.9	18	3.1	1.8-5.0	8	3.5	1.5-7.0	10	2.8	1.3-5.5
2016	45	7.7	5.6-10.5	34	13.4	9.1-19.0	11	3.3	1.6-6.2	25	4.7	3.0-7.1	14	6.7	3.6-11.2	11	3.7	1.8-6.9
2017	50	8.5	6.2-11.3	29	10.8	7.1-15.8	21	6.4	3.9-10.0	21	3.6	2.2-5.6	11	3.9	1.9-7.1	10	3.3	1.5-6.3
2018	52	8.8	6.5-11.7	28	10.9	7.1-15.9	24	7.3	4.6-11.1	22	3.6	2.3-5.6	9	3.4	1.5-6.6	13	3.9	2.0-6.9
2019	49	8.4	6.2-11.3	30	12.1	8.1-17.4	19	5.9	3.5-9.4	23	3.7	2.3-5.6	14	4.9	2.6-8.5	9	2.6	1.2-5.2
2020	39	7.1	5.0-9.8	24	10.3	6.5-15.4	15	4.4	2.4-7.6	19	3.1	1.9-5.0	13	5.0	2.6-8.8	6	1.7	0.6-4.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
Richmond (Staten Island)
Colorectal

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	174	70.5	60.2-82.0	87	84.2	66.4-105.0	87	60.7	48.5-75.0	107	45.2	36.8-54.8	56	59.7	44.2-78.3	51	36.3	26.9-47.7
1977	173	65.6	56.0-76.4	73	66.5	51.3-84.6	100	65.4	53.1-79.7	96	37.3	30.1-45.7	42	39.6	28.0-54.0	54	35.6	26.6-46.5
1978	195	72.7	62.6-83.8	92	85.9	68.2-106.5	103	65.2	53.1-79.1	91	34.7	27.8-42.7	50	47.6	34.5-63.5	41	26.5	19.0-36.0
1979	219	79.1	68.8-90.5	112	99.0	80.3-120.3	107	67.3	55.0-81.4	104	39.0	31.7-47.4	54	51.6	37.9-68.3	50	31.7	23.4-41.8
1980	227	78.0	68.0-89.1	110	94.0	76.1-114.6	117	68.5	56.5-82.3	96	34.5	27.8-42.2	40	38.0	26.5-52.3	56	33.1	24.9-43.1
1981	223	76.5	66.6-87.4	95	79.4	63.4-97.9	128	73.7	61.4-87.7	97	34.7	28.0-42.5	41	36.1	25.3-49.6	56	33.5	25.2-43.5
1982	230	78.8	68.8-89.9	125	108.8	89.5-130.8	105	60.7	49.6-73.6	90	31.5	25.2-38.8	42	38.8	27.4-52.9	48	27.5	20.2-36.6
1983	236	79.3	69.3-90.3	117	97.1	79.2-117.6	119	68.0	56.3-81.5	102	35.0	28.5-42.6	49	43.0	31.2-57.5	53	30.7	23.0-40.2
1984	243	80.7	70.7-91.6	132	111.7	92.3-133.8	111	62.4	51.2-75.2	103	33.4	27.1-40.6	54	42.7	31.6-56.3	49	26.8	19.8-35.5
1985	241	77.4	67.8-88.0	134	110.7	91.7-132.2	107	58.0	47.5-70.2	102	34.2	27.8-41.6	56	51.0	37.8-66.9	46	25.2	18.4-33.7
1986	248	78.3	68.7-88.8	115	88.5	72.3-107.1	133	70.7	59.1-83.9	115	37.2	30.6-44.7	61	52.9	39.8-68.6	54	28.6	21.5-37.4
1987	214	67.1	58.3-76.8	107	80.3	65.2-97.8	107	56.1	46.0-67.9	95	30.0	24.1-36.7	46	36.4	26.0-49.3	49	26.4	19.4-34.9
1988	237	74.1	64.8-84.2	118	89.8	73.6-108.3	119	63.8	52.7-76.4	96	31.3	25.2-38.2	47	37.9	27.3-50.9	49	26.0	19.2-34.4
1989	233	71.9	62.9-81.9	118	92.6	75.8-111.8	115	59.1	48.7-71.0	92	29.1	23.4-35.8	36	27.0	18.6-37.7	56	29.3	22.1-38.1
1990	213	64.5	56.0-73.9	90	66.6	52.9-82.7	123	62.3	51.7-74.4	111	33.7	27.7-40.6	57	41.8	31.1-54.7	54	27.3	20.5-35.7
1991	192	55.7	48.0-64.3	106	76.8	62.2-93.7	86	42.0	33.6-52.0	88	27.0	21.6-33.2	41	33.6	23.7-45.9	47	23.6	17.3-31.4
1992	246	71.4	62.7-81.0	135	98.7	82.0-117.5	111	54.0	44.4-65.1	98	29.7	24.1-36.2	46	37.5	27.0-50.3	52	25.6	19.1-33.7
1993	216	60.5	52.6-69.1	108	73.7	60.0-89.5	108	51.1	41.9-61.7	77	21.8	17.2-27.3	40	29.1	20.4-40.0	37	17.6	12.4-24.3
1994	232	63.6	55.6-72.3	116	75.8	62.1-91.4	116	53.2	44.0-63.9	103	28.8	23.4-34.9	57	42.5	31.7-55.4	46	21.3	15.6-28.5
1995	254	68.7	60.5-77.8	130	89.9	74.4-107.3	124	56.1	46.6-66.9	114	31.7	26.1-38.1	62	44.6	33.7-57.5	52	23.3	17.4-30.6
1996	252	65.6	57.7-74.2	124	78.4	64.8-93.9	128	57.4	47.8-68.3	96	25.3	20.5-30.9	48	29.8	21.7-39.7	48	20.9	15.4-27.7
1997	246	63.2	55.5-71.6	102	64.4	52.1-78.5	144	61.9	52.1-72.9	103	26.6	21.7-32.3	45	31.8	22.9-42.8	58	24.9	18.9-32.2
1998	299	73.9	65.8-82.8	156	96.3	81.3-113.1	143	60.1	50.7-70.9	121	30.2	25.1-36.1	60	36.5	27.7-47.3	61	25.2	19.3-32.5
1999	288	69.8	61.9-78.3	149	89.7	75.4-105.7	139	57.4	48.2-67.8	101	24.8	20.1-30.1	62	38.6	29.3-49.6	39	15.9	11.3-21.8
2000	276	64.6	57.2-72.7	152	84.7	71.4-99.7	124	49.8	41.4-59.5	123	29.4	24.4-35.1	55	33.8	25.2-44.2	68	27.0	20.9-34.3
2001	257	59.9	52.8-67.7	131	73.0	60.7-86.9	126	50.0	41.6-59.6	98	23.3	18.9-28.4	47	28.3	20.6-37.7	51	20.0	14.9-26.4
2002	266	59.8	52.8-67.4	137	71.3	59.5-84.6	129	49.6	41.3-59.0	92	20.9	16.9-25.7	46	26.1	18.9-35.0	46	17.3	12.7-23.2
2003	261	58.3	51.4-65.8	128	66.8	55.4-79.7	133	50.4	42.2-59.8	90	20.2	16.2-24.9	42	23.7	16.8-32.2	48	18.0	13.3-24.0
2004	224	49.8	43.5-56.8	118	62.5	51.4-75.2	106	40.2	32.9-48.7	98	22.1	17.9-27.0	52	29.5	21.8-38.8	46	16.9	12.3-22.6
2005	253	55.0	48.4-62.3	122	63.5	52.4-76.1	131	49.3	41.2-58.6	84	18.7	14.9-23.2	35	19.1	13.2-26.7	49	18.3	13.5-24.2
2006	249	53.1	46.6-60.1	122	62.0	51.2-74.3	127	46.5	38.7-55.5	92	19.8	16.0-24.4	51	27.9	20.6-36.8	41	14.9	10.7-20.4
2007	232	48.8	42.6-55.5	131	64.0	53.2-76.3	101	37.4	30.4-45.6	74	15.9	12.5-20.1	38	19.9	13.9-27.5	36	13.2	9.2-18.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Colorectal

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	252	52.5	46.1-59.5	125	61.5	50.9-73.6	127	46.0	38.3-54.9	94	19.8	15.9-24.2	48	25.2	18.5-33.6	46	16.0	11.7-21.5
2009	229	46.5	40.6-53.0	112	53.2	43.5-64.3	117	41.8	34.5-50.3	84	16.8	13.4-20.9	40	19.8	13.9-27.2	44	15.7	11.4-21.2
2010	233	46.0	40.2-52.4	120	55.5	45.7-66.7	113	38.4	31.5-46.3	88	18.0	14.4-22.3	56	29.5	22.1-38.4	32	10.5	7.1-15.0
2011	257	49.7	43.7-56.3	125	56.1	46.4-67.2	132	45.6	38.0-54.3	83	16.1	12.7-20.0	40	18.6	13.1-25.6	43	13.9	10.0-19.0
2012	224	43.1	37.5-49.3	106	46.7	38.0-56.8	118	40.2	33.2-48.5	79	15.0	11.8-18.8	27	11.1	7.2-16.4	52	17.7	13.2-23.5
2013	258	47.1	41.4-53.4	136	56.7	47.3-67.5	122	38.9	32.1-46.6	83	16.0	12.7-19.9	43	20.1	14.4-27.2	40	12.9	9.1-17.8
2014	233	41.5	36.2-47.3	127	50.0	41.4-59.9	106	33.7	27.5-41.1	97	17.7	14.3-21.7	59	26.2	19.8-34.0	38	11.6	8.2-16.2
2015	196	35.9	31.0-41.5	88	37.2	29.6-46.1	108	34.7	28.3-42.2	82	14.5	11.5-18.1	33	14.2	9.7-20.1	49	15.0	11.0-20.0
2016	234	41.3	36.0-47.1	131	51.7	43.0-61.7	103	32.4	26.3-39.6	81	14.2	11.2-17.7	44	18.8	13.5-25.4	37	10.7	7.5-15.1
2017	244	41.9	36.7-47.6	126	48.1	39.8-57.6	118	36.6	30.1-44.1	91	15.1	12.1-18.7	46	17.9	13.0-24.1	45	13.0	9.4-17.6
2018	248	42.4	37.2-48.2	140	54.3	45.5-64.4	108	32.4	26.5-39.5	86	14.6	11.6-18.1	42	16.5	11.8-22.5	44	12.6	9.1-17.3
2019	234	39.4	34.4-44.9	119	44.2	36.4-53.3	115	35.6	29.2-43.1	65	11.0	8.4-14.1	38	14.5	10.2-20.2	27	8.0	5.3-11.9
2020	243	40.5	35.4-46.1	135	49.1	40.9-58.5	108	33.2	27.1-40.5	80	12.6	9.9-15.8	43	15.7	11.2-21.4	37	10.3	7.2-14.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Colon excluding rectum

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	114	47.4	38.9-57.1	54	55.8	41.0-73.6	60	42.0	31.9-54.1	91	37.4	29.9-46.1	48	49.2	35.5-66.0	43	30.0	21.6-40.5
1977	121	45.4	37.5-54.3	49	41.7	30.3-55.8	72	47.3	36.9-59.6	76	29.1	22.8-36.6	30	26.9	17.8-38.9	46	30.2	22.0-40.3
1978	149	55.7	46.9-65.5	71	65.8	50.5-84.0	78	49.5	39.0-61.9	75	28.6	22.4-36.0	37	35.1	24.1-49.1	38	24.5	17.3-33.7
1979	171	62.9	53.6-73.2	82	75.2	58.8-94.5	89	56.2	45.0-69.2	89	33.3	26.6-41.1	44	40.8	28.9-55.6	45	28.6	20.8-38.2
1980	161	56.8	48.2-66.5	73	66.8	51.3-85.0	88	51.9	41.6-64.1	84	30.3	24.1-37.7	35	34.4	23.4-48.4	49	29.0	21.4-38.5
1981	144	50.4	42.3-59.4	60	51.2	38.4-66.6	84	49.2	39.2-61.0	86	30.8	24.5-38.1	34	29.3	19.8-41.5	52	31.2	23.2-41.0
1982	150	52.0	43.9-61.2	80	68.6	53.7-86.2	70	41.1	31.9-52.0	83	28.9	22.9-35.9	40	37.3	26.0-51.3	43	24.6	17.7-33.2
1983	169	57.1	48.7-66.6	86	70.5	55.5-88.1	83	48.1	38.2-59.7	89	30.6	24.4-37.7	45	38.3	27.4-51.9	44	25.9	18.8-34.8
1984	173	58.2	49.8-67.7	93	82.2	65.3-101.7	80	45.4	35.9-56.6	94	30.3	24.3-37.1	48	37.9	27.5-50.9	46	25.0	18.3-33.5
1985	180	58.4	50.0-67.7	98	82.4	66.0-101.3	82	44.5	35.3-55.4	85	28.8	22.9-35.7	47	43.0	30.9-57.8	38	21.1	14.9-29.1
1986	174	55.5	47.5-64.5	68	54.1	41.4-69.3	106	56.9	46.5-68.9	104	33.3	27.1-40.4	55	46.5	34.4-61.2	49	25.9	19.1-34.3
1987	155	49.1	41.6-57.5	74	57.1	44.3-72.2	81	42.4	33.6-52.8	80	25.3	20.0-31.6	41	33.1	23.1-45.6	39	21.0	14.9-28.8
1988	157	49.1	41.6-57.5	68	51.5	39.4-65.9	89	47.1	37.7-58.0	87	28.5	22.7-35.2	42	34.4	24.3-47.0	45	23.9	17.4-32.0
1989	154	48.0	40.7-56.3	70	57.6	44.1-73.5	84	43.0	34.3-53.3	81	25.6	20.3-31.9	28	20.6	13.4-30.1	53	27.8	20.8-36.4
1990	164	50.0	42.5-58.3	68	50.2	38.3-64.5	96	48.7	39.4-59.5	105	32.1	26.2-38.9	52	38.6	28.3-51.1	53	26.8	20.1-35.1
1991	137	39.8	33.4-47.2	70	50.2	38.7-63.8	67	33.0	25.5-42.0	84	25.8	20.5-31.9	41	33.6	23.7-45.9	43	21.6	15.6-29.1
1992	178	51.7	44.4-60.0	98	72.7	58.4-89.2	80	38.8	30.8-48.4	86	26.2	20.9-32.4	41	33.7	23.7-46.0	45	22.4	16.3-30.0
1993	154	43.2	36.6-50.7	75	51.5	40.1-65.0	79	37.5	29.7-46.8	68	19.4	15.0-24.6	36	26.4	18.1-37.0	32	15.3	10.5-21.7
1994	166	45.6	38.9-53.1	82	53.1	41.8-66.3	84	38.6	30.8-47.8	97	26.9	21.8-32.9	54	39.4	29.3-51.8	43	19.9	14.4-26.9
1995	191	52.2	45.0-60.2	97	68.9	55.3-84.5	94	42.7	34.5-52.3	105	29.2	23.9-35.4	58	42.3	31.7-55.0	47	21.0	15.5-28.0
1996	188	49.0	42.2-56.6	89	57.3	45.6-70.9	99	44.3	36.0-54.0	87	23.0	18.4-28.4	43	26.9	19.2-36.5	44	19.1	13.9-25.8
1997	183	47.3	40.7-54.7	77	49.1	38.5-61.7	106	45.5	37.2-55.1	90	23.2	18.6-28.5	41	27.8	19.7-37.9	49	21.0	15.5-27.8
1998	208	51.4	44.7-58.9	95	57.9	46.4-71.1	113	47.4	39.1-57.1	108	26.9	22.1-32.5	51	31.3	23.1-41.3	57	23.8	18.0-30.8
1999	200	48.8	42.3-56.1	107	65.3	53.2-79.3	93	38.3	30.9-46.9	86	21.0	16.8-25.9	53	32.8	24.3-43.0	33	13.5	9.3-19.0
2000	201	47.2	40.8-54.2	100	56.1	45.3-68.5	101	40.5	33.0-49.3	103	24.6	20.1-29.9	46	27.8	20.1-37.3	57	22.6	17.1-29.4
2001	192	44.9	38.8-51.7	90	51.3	41.0-63.3	102	40.6	33.1-49.3	87	20.7	16.6-25.5	40	23.8	16.9-32.5	47	18.5	13.6-24.7
2002	195	43.9	37.9-50.6	91	48.1	38.4-59.4	104	40.0	32.6-48.5	80	18.2	14.4-22.7	39	21.9	15.4-30.1	41	15.4	11.1-21.0
2003	198	44.3	38.4-51.0	95	49.1	39.5-60.3	103	39.1	31.9-47.5	77	17.3	13.7-21.7	34	19.4	13.3-27.3	43	16.3	11.8-22.0
2004	162	36.3	30.9-42.4	83	45.1	35.7-56.1	79	29.8	23.6-37.3	90	20.3	16.3-25.0	45	25.4	18.3-34.1	45	16.5	12.0-22.2
2005	174	38.0	32.5-44.1	76	39.5	30.9-49.7	98	36.6	29.7-44.7	71	16.0	12.5-20.2	28	15.8	10.4-22.8	43	16.2	11.7-21.9
2006	179	38.5	33.0-44.6	86	44.1	35.1-54.7	93	34.1	27.5-42.0	71	15.2	11.8-19.2	39	20.7	14.6-28.5	32	11.4	7.8-16.3
2007	168	35.2	30.1-41.1	95	46.0	37.0-56.6	73	27.0	21.1-34.1	68	14.6	11.3-18.5	35	18.2	12.5-25.4	33	11.9	8.2-16.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Colon excluding rectum

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	175	36.7	31.4-42.6	89	44.5	35.5-55.0	86	31.2	24.9-38.6	79	16.7	13.2-20.8	43	23.0	16.5-31.0	36	12.6	8.7-17.5
2009	172	35.1	30.0-40.8	81	38.8	30.6-48.5	91	32.9	26.4-40.5	67	13.7	10.5-17.4	32	16.0	10.8-22.8	35	12.9	8.9-18.0
2010	174	34.1	29.2-39.7	83	38.3	30.3-47.8	91	30.3	24.3-37.4	78	16.0	12.6-20.0	50	26.8	19.8-35.4	28	9.1	6.0-13.3
2011	177	34.5	29.5-40.1	85	38.9	30.8-48.4	92	31.7	25.4-39.1	72	14.1	11.0-17.8	36	16.7	11.5-23.3	36	11.7	8.2-16.4
2012	164	31.9	27.2-37.4	75	33.2	25.9-42.0	89	30.9	24.7-38.2	70	13.4	10.4-17.0	21	9.0	5.5-13.9	49	16.5	12.2-22.0
2013	190	35.0	30.1-40.5	94	40.4	32.4-49.7	96	30.4	24.5-37.3	70	13.5	10.5-17.2	38	17.8	12.5-24.6	32	10.3	7.0-14.8
2014	162	28.8	24.4-33.7	84	33.7	26.7-42.0	78	24.2	19.0-30.5	75	13.7	10.7-17.3	44	19.7	14.2-26.6	31	9.5	6.4-13.6
2015	142	26.3	22.1-31.1	61	25.7	19.5-33.3	81	26.2	20.7-32.8	71	12.7	9.9-16.1	29	12.7	8.4-18.4	42	12.9	9.3-17.7
2016	167	29.3	25.0-34.3	91	36.9	29.5-45.6	76	23.2	18.2-29.2	67	11.7	9.1-15.0	37	15.6	10.9-21.7	30	8.9	6.0-13.0
2017	178	30.7	26.3-35.7	82	31.4	24.8-39.3	96	29.9	24.1-36.8	75	12.5	9.8-15.8	34	13.5	9.2-19.1	41	11.9	8.5-16.4
2018	178	30.4	26.0-35.3	101	39.4	31.9-48.2	77	23.2	18.2-29.3	70	11.7	9.1-15.0	33	12.7	8.7-18.1	37	10.6	7.4-14.9
2019	187	31.2	26.8-36.1	96	35.6	28.6-43.8	91	27.6	22.2-34.2	52	8.9	6.6-11.8	29	11.3	7.5-16.5	23	7.1	4.4-10.8
2020	177	29.1	24.9-33.9	95	34.5	27.8-42.5	82	24.7	19.5-30.9	68	10.7	8.3-13.7	35	12.7	8.8-18.0	33	9.2	6.2-13.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Rectum & rectosigmoid

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	60	23.1	17.5-29.9	33	28.5	19.1-40.5	27	18.7	12.3-27.3	16	7.8	4.4-12.5	8	10.4	4.3-20.1	8	6.2	2.7-12.1
1977	52	20.3	15.0-26.7	24	24.8	15.4-37.4	28	18.2	12.0-26.3	20	8.2	4.9-12.6	12	12.6	6.3-22.2	8	5.4	2.3-10.7
1978	46	17.0	12.3-22.8	21	20.1	12.0-31.2	25	15.7	10.1-23.2	16	6.0	3.4-9.9	13	12.4	6.2-21.6	3	2.0	0.4-5.8
1979	48	16.3	11.9-21.7	30	23.7	15.6-34.4	18	11.1	6.5-17.6	15	5.7	3.2-9.5	10	10.8	4.9-20.0	5	3.1	1.0-7.2
1980	66	21.2	16.3-27.1	37	27.2	18.9-38.0	29	16.6	11.0-23.9	12	4.1	2.1-7.2	5	3.5	1.1-8.4	7	4.1	1.6-8.4
1981	79	26.1	20.6-32.7	35	28.2	19.2-39.8	44	24.4	17.7-32.9	11	3.9	1.9-7.0	7	6.8	2.6-14.0	4	2.3	0.6-5.9
1982	80	26.8	21.2-33.5	45	40.2	28.6-54.5	35	19.6	13.6-27.4	7	2.6	1.0-5.3	2	1.5	0.2-5.4	5	2.9	0.9-6.8
1983	67	22.2	17.1-28.3	31	26.6	17.6-38.3	36	19.9	13.9-27.7	13	4.5	2.3-7.7	4	4.7	1.2-11.5	9	4.8	2.2-9.3
1984	70	22.4	17.4-28.4	39	29.5	20.5-41.0	31	17.0	11.5-24.1	9	3.1	1.4-5.8	6	4.8	1.7-10.5	3	1.7	0.4-5.0
1985	61	19.0	14.5-24.5	36	28.3	19.3-39.8	25	13.5	8.7-19.9	17	5.4	3.1-8.6	9	8.0	3.5-15.2	8	4.1	1.7-8.1
1986	74	22.8	17.8-28.7	47	34.4	24.9-46.2	27	13.8	9.1-20.2	11	3.9	1.9-6.9	6	6.4	2.2-13.4	5	2.7	0.9-6.3
1987	59	18.0	13.6-23.3	33	23.3	15.7-33.1	26	13.7	8.9-20.2	15	4.6	2.6-7.7	5	3.3	1.0-7.9	10	5.3	2.5-9.8
1988	80	25.0	19.7-31.1	50	38.3	28.0-50.9	30	16.7	11.2-23.8	9	2.8	1.3-5.3	5	3.4	1.1-8.1	4	2.1	0.6-5.4
1989	79	23.9	18.9-29.8	48	35.1	25.5-46.8	31	16.1	10.9-22.9	11	3.5	1.7-6.2	8	6.4	2.7-12.5	3	1.6	0.3-4.6
1990	49	14.6	10.7-19.3	22	16.4	10.1-25.0	27	13.6	9.0-19.9	6	1.6	0.6-3.6	5	3.2	1.0-7.8	1	0.5	0.0-2.8
1991	55	15.9	11.9-20.7	36	26.7	18.1-37.6	19	9.0	5.4-14.2	4	1.2	0.3-3.1	0	0.0	0.0-2.4	4	1.9	0.5-5.0
1992	68	19.7	15.2-24.9	37	26.0	18.0-36.1	31	15.2	10.3-21.6	12	3.5	1.8-6.0	5	3.8	1.2-8.7	7	3.3	1.3-6.8
1993	62	17.2	13.2-22.1	33	22.2	15.1-31.4	29	13.5	9.1-19.5	9	2.4	1.1-4.6	4	2.7	0.7-6.8	5	2.3	0.7-5.4
1994	66	18.0	13.9-22.9	34	22.7	15.4-32.0	32	14.7	10.0-20.8	6	1.8	0.7-3.9	3	3.0	0.6-8.3	3	1.4	0.3-4.1
1995	63	16.5	12.7-21.1	33	21.0	14.2-29.8	30	13.4	9.0-19.2	9	2.5	1.1-4.7	4	2.2	0.6-5.8	5	2.2	0.7-5.3
1996	64	16.6	12.7-21.2	35	21.1	14.6-29.5	29	13.0	8.7-18.7	9	2.3	1.1-4.4	5	2.8	0.9-6.6	4	1.7	0.5-4.5
1997	63	15.9	12.2-20.4	25	15.3	9.7-22.7	38	16.4	11.6-22.5	13	3.4	1.8-5.8	4	4.0	1.1-9.5	9	3.9	1.8-7.5
1998	91	22.5	18.1-27.6	61	38.4	29.1-49.6	30	12.7	8.6-18.2	13	3.3	1.8-5.6	9	5.3	2.4-10.0	4	1.5	0.4-4.0
1999	88	21.0	16.8-25.8	42	24.3	17.3-33.1	46	19.1	14.0-25.5	15	3.7	2.1-6.1	9	5.8	2.6-10.9	6	2.4	0.9-5.4
2000	75	17.5	13.7-21.9	52	28.6	21.1-37.8	23	9.3	5.9-14.0	20	4.8	2.9-7.4	9	6.0	2.7-11.3	11	4.4	2.2-7.9
2001	65	15.0	11.6-19.1	41	21.7	15.5-29.6	24	9.4	6.0-14.1	11	2.6	1.3-4.6	7	4.5	1.7-9.1	4	1.5	0.4-4.0
2002	71	15.8	12.4-20.0	46	23.2	16.9-31.0	25	9.6	6.2-14.2	12	2.7	1.4-4.8	7	4.2	1.6-8.6	5	1.9	0.6-4.5
2003	63	13.9	10.7-17.9	33	17.6	12.0-24.9	30	11.3	7.6-16.2	13	2.9	1.5-4.9	8	4.3	1.8-8.5	5	1.7	0.6-4.2
2004	62	13.5	10.3-17.3	35	17.4	12.0-24.4	27	10.4	6.8-15.2	8	1.8	0.8-3.6	7	4.1	1.6-8.4	1	0.4	0.0-2.2
2005	79	17.1	13.5-21.3	46	24.0	17.4-32.1	33	12.7	8.7-17.9	13	2.7	1.4-4.7	7	3.3	1.3-7.0	6	2.0	0.7-4.6
2006	70	14.6	11.3-18.5	36	17.9	12.4-24.9	34	12.3	8.5-17.4	21	4.6	2.9-7.1	12	7.1	3.6-12.4	9	3.5	1.6-6.7
2007	64	13.5	10.4-17.3	36	17.9	12.4-25.0	28	10.4	6.9-15.1	6	1.4	0.5-3.0	3	1.8	0.3-5.0	3	1.2	0.3-3.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Rectum & rectosigmoid

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	77	15.8	12.4-19.8	36	17.0	11.8-23.8	41	14.8	10.6-20.3	15	3.1	1.7-5.1	5	2.3	0.7-5.4	10	3.4	1.6-6.5
2009	57	11.4	8.6-14.8	31	14.4	9.6-20.5	26	9.0	5.8-13.3	17	3.2	1.8-5.1	8	3.8	1.6-7.6	9	2.8	1.3-5.6
2010	59	11.8	9.0-15.4	37	17.2	11.9-23.9	22	8.1	5.0-12.4	10	2.1	1.0-3.8	6	2.7	1.0-5.9	4	1.4	0.4-3.8
2011	80	15.2	12.0-19.1	40	17.2	12.1-23.6	40	13.9	9.9-19.1	11	2.0	1.0-3.6	4	2.0	0.5-5.1	7	2.2	0.9-4.7
2012	60	11.1	8.4-14.5	31	13.5	9.0-19.3	29	9.4	6.2-13.7	9	1.6	0.7-3.2	6	2.1	0.7-5.0	3	1.2	0.3-3.6
2013	68	12.1	9.4-15.5	42	16.3	11.6-22.3	26	8.5	5.5-12.7	13	2.4	1.3-4.2	5	2.3	0.7-5.4	8	2.6	1.1-5.3
2014	71	12.7	9.9-16.2	43	16.4	11.7-22.3	28	9.5	6.3-14.0	22	4.0	2.5-6.1	15	6.5	3.6-10.9	7	2.2	0.9-4.7
2015	54	9.7	7.2-12.7	27	11.4	7.4-16.8	27	8.5	5.5-12.6	11	1.8	0.9-3.4	4	1.5	0.4-4.1	7	2.0	0.8-4.4
2016	67	11.9	9.2-15.3	40	14.8	10.5-20.4	27	9.2	6.0-13.7	14	2.4	1.3-4.2	7	3.2	1.2-6.6	7	1.8	0.7-4.1
2017	66	11.2	8.6-14.3	44	16.6	12.0-22.6	22	6.7	4.1-10.4	16	2.6	1.5-4.3	12	4.4	2.2-7.9	4	1.1	0.3-3.1
2018	70	12.0	9.3-15.4	39	14.9	10.5-20.6	31	9.2	6.2-13.4	16	2.8	1.6-4.7	9	3.7	1.7-7.2	7	2.0	0.8-4.5
2019	47	8.2	6.0-11.0	23	8.6	5.4-13.2	24	8.0	5.0-12.1	13	2.1	1.1-3.7	9	3.2	1.4-6.3	4	1.0	0.3-2.9
2020	66	11.4	8.7-14.6	40	14.6	10.3-20.2	26	8.5	5.5-12.8	12	1.9	1.0-3.4	8	2.9	1.2-6.0	4	1.1	0.3-3.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Liver / intrahepatic bile duct

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	15	5.8	3.2-9.6	7	6.0	2.3-12.5	8	5.3	2.3-10.4	11	4.5	2.2-8.0	7	6.4	2.5-13.2	4	2.8	0.8-7.1
1977	15	5.6	3.1-9.3	9	7.3	3.2-14.0	6	4.0	1.5-8.7	16	5.9	3.3-9.7	7	5.6	2.2-11.8	9	5.8	2.6-11.1
1978	11	4.2	2.1-7.6	6	5.9	1.9-12.9	5	3.3	1.1-7.7	11	3.9	1.9-7.0	6	5.6	1.9-12.1	5	2.9	0.9-6.9
1979	12	4.2	2.1-7.3	7	5.7	2.2-11.8	5	2.9	0.9-6.9	17	6.5	3.8-10.4	10	9.3	4.3-17.2	7	4.5	1.8-9.1
1980	9	2.6	1.2-5.1	6	4.4	1.5-10.0	3	1.5	0.3-4.5	10	3.5	1.7-6.5	7	6.4	2.4-13.3	3	1.7	0.3-5.0
1981	12	4.0	2.0-7.0	5	3.4	1.1-8.2	7	4.1	1.6-8.5	15	5.3	3.0-8.8	9	7.5	3.3-14.4	6	3.9	1.4-8.3
1982	8	2.8	1.2-5.4	6	5.4	1.9-11.8	2	1.3	0.2-4.6	13	4.1	2.2-7.1	7	6.2	2.3-12.9	6	3.2	1.2-7.1
1983	16	5.4	3.1-8.9	8	6.8	2.6-13.8	8	4.8	2.1-9.4	19	6.6	4.0-10.4	9	7.9	3.3-15.3	10	6.0	2.9-11.0
1984	15	4.8	2.7-8.0	8	6.3	2.6-12.7	7	3.7	1.5-7.7	26	8.8	5.7-12.9	15	12.2	6.5-20.4	11	6.3	3.1-11.3
1985	13	4.3	2.3-7.4	8	6.8	2.7-13.5	5	2.7	0.9-6.4	16	5.5	3.1-8.8	9	7.6	3.3-14.4	7	4.0	1.6-8.3
1986	14	4.6	2.5-7.6	11	9.0	4.2-16.5	3	1.8	0.4-5.2	17	5.5	3.2-8.8	14	12.3	6.5-20.7	3	1.7	0.3-4.9
1987	12	3.8	1.9-6.6	5	3.3	1.0-7.8	7	3.8	1.5-7.7	17	5.3	3.0-8.5	10	6.5	3.1-12.1	7	3.7	1.5-7.7
1988	20	6.3	3.8-9.7	14	10.1	5.4-17.2	6	3.2	1.2-7.0	27	8.7	5.7-12.6	18	15.4	8.8-24.7	9	4.7	2.2-9.0
1989	16	5.1	2.9-8.3	7	4.7	1.9-9.7	9	4.8	2.2-9.1	21	6.6	4.1-10.1	7	5.7	2.2-11.7	14	7.3	4.0-12.3
1990	15	4.3	2.4-7.1	10	6.9	3.2-12.7	5	2.3	0.8-5.6	24	7.3	4.6-10.8	16	12.2	6.7-20.0	8	4.0	1.7-8.0
1991	18	5.1	3.0-8.1	12	8.0	4.0-14.0	6	2.9	1.1-6.4	28	8.0	5.3-11.6	16	11.5	6.2-19.1	12	5.8	3.0-10.2
1992	17	4.7	2.7-7.6	13	8.1	4.3-13.8	4	1.9	0.5-4.8	21	6.0	3.7-9.2	12	7.5	3.8-13.2	9	4.3	2.0-8.2
1993	16	4.6	2.6-7.5	8	5.8	2.5-11.4	8	3.8	1.6-7.4	24	7.0	4.5-10.4	16	11.1	6.2-18.0	8	3.8	1.6-7.5
1994	12	3.3	1.7-5.7	9	5.8	2.6-11.1	3	1.4	0.3-4.2	15	4.1	2.3-6.7	11	7.2	3.5-12.8	4	2.0	0.5-5.0
1995	22	5.7	3.6-8.6	13	7.4	3.9-12.8	9	4.0	1.8-7.6	25	6.6	4.3-9.8	12	7.3	3.6-13.0	13	5.9	3.1-10.1
1996	25	6.5	4.2-9.6	17	11.3	6.4-18.2	8	3.4	1.5-6.9	19	5.1	3.1-8.0	10	7.2	3.3-13.3	9	3.8	1.7-7.2
1997	23	5.6	3.6-8.5	16	8.7	4.9-14.1	7	3.1	1.3-6.5	16	4.0	2.3-6.4	9	5.0	2.3-9.5	7	3.0	1.2-6.2
1998	22	5.4	3.4-8.2	18	11.2	6.5-17.7	4	1.9	0.5-4.8	21	5.2	3.2-7.9	21	13.1	7.9-20.1	0	0.0	0.0-1.7
1999	27	6.5	4.3-9.4	17	8.8	5.1-14.2	10	3.9	1.9-7.3	23	5.6	3.5-8.4	10	5.1	2.4-9.5	13	5.4	2.8-9.2
2000	38	8.9	6.3-12.3	27	14.8	9.7-21.6	11	4.3	2.1-7.7	33	7.8	5.3-10.9	22	12.2	7.5-18.5	11	4.3	2.1-7.8
2001	34	7.6	5.2-10.6	24	11.5	7.3-17.2	10	3.9	1.9-7.3	27	6.2	4.1-9.1	18	9.9	5.8-15.8	9	3.5	1.6-6.7
2002	25	5.6	3.6-8.2	16	8.3	4.7-13.6	9	3.6	1.6-6.8	28	6.4	4.2-9.2	21	11.7	7.1-17.9	7	2.8	1.1-5.8
2003	37	8.2	5.7-11.3	30	15.5	10.4-22.2	7	2.7	1.1-5.6	25	5.4	3.5-8.0	17	8.4	4.9-13.6	8	3.0	1.3-6.0
2004	35	7.4	5.1-10.3	25	12.7	8.1-19.0	10	3.8	1.8-7.1	32	7.0	4.8-9.9	21	10.1	6.2-15.5	11	4.2	2.1-7.6
2005	43	9.1	6.6-12.3	36	17.8	12.3-24.9	7	2.5	1.0-5.4	30	6.4	4.3-9.2	17	8.4	4.8-13.6	13	5.0	2.6-8.6
2006	51	10.4	7.7-13.7	34	15.2	10.4-21.5	17	5.9	3.4-9.6	37	7.8	5.5-10.8	23	10.9	6.8-16.5	14	5.2	2.8-8.9
2007	46	9.0	6.5-12.1	36	16.0	11.1-22.4	10	3.7	1.8-7.0	22	4.5	2.8-6.8	14	6.7	3.6-11.4	8	3.1	1.3-6.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Liver / intrahepatic bile duct

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	61	11.9	9.0-15.3	46	18.8	13.6-25.3	15	5.7	3.2-9.6	33	6.8	4.6-9.6	22	10.1	6.3-15.5	11	3.9	1.9-7.1
2009	53	9.8	7.3-13.0	35	13.7	9.4-19.3	18	6.0	3.5-9.6	38	7.4	5.2-10.3	21	8.9	5.4-13.8	17	6.2	3.6-10.1
2010	52	9.8	7.2-12.9	39	16.5	11.6-22.8	13	4.3	2.3-7.6	41	7.8	5.5-10.6	27	11.7	7.6-17.3	14	4.6	2.5-8.0
2011	59	10.7	8.1-13.8	42	16.6	11.8-22.8	17	5.5	3.2-9.0	32	5.9	4.0-8.5	22	9.2	5.7-14.2	10	3.3	1.5-6.2
2012	67	11.7	9.0-15.0	50	19.0	14.0-25.4	17	5.7	3.3-9.3	45	8.3	6.0-11.2	32	13.3	9.0-19.1	13	4.4	2.3-7.8
2013	57	9.6	7.2-12.5	43	15.4	11.1-21.1	14	4.6	2.5-7.9	35	6.2	4.3-8.7	29	11.3	7.5-16.5	6	2.0	0.7-4.5
2014	70	12.1	9.4-15.5	52	19.7	14.6-26.1	18	5.8	3.3-9.3	48	8.5	6.2-11.3	32	12.1	8.2-17.4	16	5.4	3.0-8.9
2015	61	10.6	8.0-13.7	42	16.0	11.4-21.9	19	6.4	3.8-10.3	38	6.4	4.5-8.9	31	11.4	7.7-16.5	7	2.4	0.9-5.1
2016	60	9.6	7.3-12.6	45	16.3	11.7-22.0	15	4.2	2.3-7.3	35	5.7	4.0-8.1	20	7.3	4.4-11.5	15	4.2	2.3-7.2
2017	60	9.6	7.2-12.4	43	14.4	10.3-19.7	17	5.3	3.1-8.7	38	6.2	4.3-8.6	24	8.9	5.6-13.5	14	4.1	2.2-7.1
2018	79	12.9	10.2-16.3	54	19.5	14.5-25.8	25	7.2	4.6-10.8	52	8.4	6.2-11.1	33	11.2	7.6-16.0	19	5.6	3.4-9.0
2019	70	10.8	8.4-13.8	51	17.0	12.6-22.6	19	5.7	3.4-9.2	40	6.5	4.6-9.0	30	10.9	7.2-15.8	10	3.0	1.4-5.9
2020	57	8.7	6.5-11.3	43	14.5	10.4-19.9	14	3.9	2.1-6.8	31	4.8	3.3-7.0	20	7.2	4.3-11.4	11	3.1	1.5-5.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Pancreas

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	32	12.4	8.4-17.6	18	15.5	8.8-25.0	14	9.9	5.4-16.5	33	13.0	8.9-18.4	16	13.6	7.5-22.6	17	12.0	6.9-19.1
1977	25	9.2	5.9-13.7	14	12.7	6.7-21.6	11	7.2	3.5-12.8	32	12.0	8.2-17.1	17	15.0	8.5-24.4	15	9.9	5.5-16.4
1978	21	6.6	4.0-10.1	15	9.8	5.4-16.4	6	3.6	1.3-7.9	21	7.2	4.4-11.0	15	11.6	6.4-19.5	6	3.7	1.3-8.1
1979	31	11.6	7.8-16.5	13	12.6	6.4-21.9	18	10.9	6.5-17.3	27	10.3	6.7-15.0	13	13.7	7.0-23.6	14	8.7	4.7-14.5
1980	41	14.5	10.3-19.7	22	20.1	12.2-30.7	19	11.2	6.7-17.6	34	11.6	7.9-16.2	17	15.0	8.4-24.3	17	9.4	5.4-15.1
1981	30	11.0	7.3-15.6	16	13.6	7.5-22.2	14	9.0	4.9-15.0	34	12.2	8.4-17.1	13	11.9	6.1-20.4	21	12.4	7.6-19.0
1982	43	14.8	10.6-20.0	19	17.1	9.8-27.1	24	14.1	8.9-21.0	29	9.7	6.4-14.0	12	10.2	5.0-18.0	17	9.8	5.7-15.8
1983	37	12.2	8.5-16.9	17	13.1	7.4-21.4	20	11.4	6.9-17.6	32	10.4	7.1-14.7	12	9.3	4.5-16.6	20	11.6	7.0-18.0
1984	33	11.5	7.9-16.2	13	11.6	5.8-20.1	20	11.3	6.9-17.5	35	11.8	8.1-16.4	19	14.8	8.6-23.5	16	9.3	5.3-15.1
1985	36	12.0	8.4-16.7	18	15.4	8.9-24.5	18	9.8	5.8-15.6	34	11.5	7.9-16.1	18	16.0	9.2-25.5	16	8.7	5.0-14.2
1986	44	13.5	9.8-18.1	22	16.8	10.2-25.7	22	11.2	7.0-17.1	41	12.5	9.0-17.1	21	15.8	9.5-24.4	20	10.4	6.3-16.2
1987	44	14.0	10.1-18.8	23	17.5	10.8-26.5	21	11.6	7.1-17.8	36	11.0	7.7-15.3	21	14.6	8.9-22.5	15	8.2	4.6-13.6
1988	53	16.6	12.4-21.7	25	19.9	12.5-29.8	28	14.6	9.6-21.1	41	13.2	9.4-17.9	24	19.9	12.4-29.9	17	9.1	5.3-14.6
1989	47	14.5	10.6-19.3	20	17.3	10.1-27.2	27	13.7	9.0-20.0	45	14.1	10.2-18.9	20	17.3	10.2-27.0	25	12.6	8.1-18.6
1990	55	16.8	12.6-22.0	24	16.8	10.7-25.1	31	15.6	10.6-22.2	44	13.3	9.6-17.9	22	16.3	10.0-24.9	22	11.2	7.0-16.9
1991	60	17.5	13.3-22.6	32	22.6	15.2-32.1	28	13.6	9.0-19.7	44	13.3	9.6-17.9	24	16.9	10.5-25.5	20	10.2	6.2-15.7
1992	54	15.7	11.8-20.5	31	22.8	15.2-32.8	23	11.2	7.1-16.8	46	13.4	9.8-17.9	26	19.3	12.3-28.5	20	9.4	5.7-14.5
1993	50	13.9	10.3-18.3	18	11.2	6.6-17.8	32	15.1	10.3-21.3	55	15.3	11.5-20.0	23	15.2	9.5-22.8	32	15.1	10.3-21.3
1994	50	14.1	10.5-18.6	24	16.3	10.3-24.4	26	12.0	7.9-17.7	45	12.5	9.1-16.7	20	12.7	7.6-19.8	25	11.4	7.4-16.9
1995	46	12.6	9.2-16.8	17	11.0	6.3-17.6	29	13.2	8.9-19.0	41	11.1	8.0-15.1	18	11.3	6.6-17.9	23	10.5	6.7-15.8
1996	38	9.9	7.0-13.6	17	10.5	6.0-16.8	21	9.3	5.7-14.2	45	11.8	8.6-15.8	20	12.3	7.4-19.1	25	11.2	7.2-16.5
1997	60	15.2	11.6-19.5	30	16.7	11.2-24.0	30	13.0	8.8-18.6	46	11.7	8.6-15.6	26	15.2	9.8-22.3	20	8.4	5.1-13.1
1998	69	16.9	13.2-21.4	39	23.0	16.2-31.6	30	12.5	8.5-18.0	57	14.3	10.8-18.5	26	15.5	9.9-22.8	31	12.9	8.7-18.3
1999	67	16.0	12.4-20.3	32	18.7	12.6-26.6	35	14.2	9.9-19.9	56	13.3	10.0-17.3	29	16.5	10.9-23.9	27	11.1	7.3-16.2
2000	78	18.1	14.3-22.7	41	22.7	16.1-31.1	37	14.5	10.2-20.1	62	14.5	11.1-18.6	29	16.1	10.6-23.3	33	13.0	8.9-18.3
2001	56	12.9	9.7-16.7	23	12.8	8.0-19.2	33	12.8	8.8-18.1	61	14.1	10.8-18.2	28	16.1	10.5-23.4	33	12.9	8.9-18.2
2002	61	13.7	10.4-17.6	30	14.9	9.9-21.5	31	12.1	8.2-17.3	44	10.1	7.3-13.6	21	11.6	7.1-17.8	23	8.9	5.6-13.4
2003	61	13.6	10.4-17.5	28	14.6	9.6-21.2	33	12.5	8.6-17.6	63	13.9	10.7-17.8	34	17.3	11.8-24.3	29	10.8	7.2-15.6
2004	58	12.8	9.7-16.6	28	14.9	9.7-21.6	30	11.6	7.8-16.6	52	11.7	8.7-15.4	31	17.4	11.7-24.8	21	7.9	4.9-12.2
2005	66	14.1	10.9-18.0	28	13.6	8.9-19.9	38	14.1	9.9-19.5	56	12.0	9.0-15.6	30	14.8	9.9-21.3	26	9.6	6.2-14.2
2006	70	15.0	11.6-19.0	35	17.1	11.8-24.0	35	12.9	8.9-18.0	53	11.5	8.6-15.1	20	10.5	6.3-16.4	33	11.9	8.2-16.9
2007	71	14.6	11.3-18.4	36	16.7	11.5-23.4	35	12.4	8.6-17.4	56	11.6	8.8-15.2	23	11.6	7.2-17.6	33	11.6	7.9-16.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Pancreas

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	71	14.3	11.1-18.1	40	18.1	12.8-24.9	31	10.9	7.3-15.6	61	12.3	9.3-15.8	40	19.0	13.4-26.2	21	7.3	4.5-11.2
2009	85	17.0	13.5-21.1	35	15.2	10.5-21.4	50	18.1	13.4-24.0	65	13.0	9.9-16.6	30	12.6	8.4-18.3	35	12.8	8.8-17.9
2010	75	14.6	11.4-18.4	31	14.6	9.7-20.9	44	14.7	10.6-19.9	62	11.9	9.0-15.3	30	13.2	8.7-19.0	32	10.4	7.0-14.9
2011	77	14.9	11.7-18.8	42	18.5	13.2-25.3	35	12.0	8.3-16.8	67	13.1	10.1-16.7	30	13.6	9.1-19.6	37	12.7	8.9-17.7
2012	67	12.5	9.6-16.0	38	16.1	11.3-22.4	29	10.0	6.6-14.5	59	10.8	8.2-14.0	32	14.3	9.6-20.4	27	8.7	5.7-12.9
2013	61	10.7	8.1-13.8	28	10.6	6.9-15.5	33	10.7	7.3-15.2	47	8.5	6.2-11.5	23	8.9	5.5-13.6	24	7.9	5.0-12.0
2014	90	16.1	12.9-19.9	44	17.4	12.4-23.6	46	14.5	10.6-19.6	55	9.9	7.4-13.0	26	10.8	7.0-16.1	29	9.2	6.1-13.4
2015	85	14.8	11.7-18.4	42	16.8	12.0-23.0	43	13.2	9.5-18.0	60	10.5	8.0-13.6	28	10.7	7.0-15.8	32	9.9	6.8-14.2
2016	85	14.4	11.5-18.0	41	16.2	11.5-22.2	44	13.1	9.4-17.8	71	11.7	9.1-14.9	32	12.4	8.4-17.8	39	11.2	7.9-15.6
2017	97	15.9	12.9-19.6	47	16.1	11.7-21.7	50	15.5	11.4-20.6	77	12.5	9.8-15.8	37	13.6	9.4-19.0	40	11.9	8.5-16.5
2018	93	15.7	12.6-19.3	53	20.3	15.1-26.8	40	11.8	8.4-16.3	65	10.8	8.3-13.9	34	13.1	9.0-18.5	31	9.3	6.2-13.4
2019	102	16.9	13.7-20.6	44	15.7	11.3-21.3	58	17.4	13.1-22.8	79	13.1	10.3-16.4	37	13.8	9.6-19.2	42	12.3	8.8-16.9
2020	94	14.6	11.7-18.0	48	16.8	12.2-22.6	46	13.0	9.4-17.6	75	12.0	9.4-15.2	35	13.2	9.1-18.6	40	10.9	7.7-15.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Larynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	19	6.1	3.6-9.6	13	9.5	5.0-16.6	6	3.4	1.2-7.5	11	3.8	1.9-6.9	10	8.5	3.9-16.0	1	0.6	0.0-3.4
1977	21	7.4	4.5-11.4	17	14.1	7.8-23.3	4	2.8	0.7-7.1	10	3.4	1.6-6.3	9	6.9	2.9-13.8	1	0.7	0.0-3.7
1978	20	6.6	4.0-10.3	19	14.9	8.7-23.7	1	0.5	0.0-3.1	7	2.5	1.0-5.2	6	5.1	1.7-11.5	1	0.9	0.0-4.2
1979	30	10.4	6.9-14.8	22	19.0	11.5-29.2	8	4.8	2.1-9.5	5	1.5	0.5-3.7	4	3.0	0.7-7.9	1	0.5	0.0-3.1
1980	26	8.5	5.5-12.5	22	16.0	9.9-24.3	4	2.6	0.7-6.6	11	3.6	1.8-6.5	8	6.2	2.5-12.5	3	1.8	0.4-5.2
1981	23	7.7	4.8-11.6	17	14.2	8.0-23.1	6	3.3	1.2-7.3	12	4.2	2.2-7.4	10	9.0	4.1-16.7	2	1.4	0.2-4.9
1982	21	7.2	4.4-11.0	17	15.8	8.9-25.3	4	2.0	0.5-5.3	8	2.7	1.2-5.3	6	4.4	1.5-9.7	2	1.2	0.2-4.4
1983	30	9.0	6.1-13.0	25	16.8	10.7-25.0	5	3.0	0.9-6.9	7	2.2	0.9-4.5	4	3.2	0.8-8.3	3	1.6	0.3-4.7
1984	24	7.6	4.8-11.3	17	12.7	7.2-20.5	7	4.0	1.6-8.3	9	2.9	1.3-5.6	7	5.2	2.0-10.9	2	1.1	0.1-4.0
1985	28	8.7	5.7-12.6	23	16.8	10.4-25.5	5	2.8	0.9-6.5	5	1.6	0.5-3.6	2	1.8	0.2-6.3	3	1.7	0.4-5.0
1986	29	8.4	5.6-12.1	23	15.6	9.8-23.6	6	3.3	1.2-7.2	7	1.8	0.7-3.7	5	2.7	0.9-6.7	2	1.0	0.1-3.6
1987	27	8.2	5.4-12.0	19	13.6	7.9-21.7	8	4.9	2.1-9.5	8	2.3	1.0-4.5	8	5.3	2.2-10.7	0	0.0	0.0-2.0
1988	27	7.9	5.2-11.5	23	16.3	10.1-24.9	4	2.2	0.6-5.6	9	2.8	1.3-5.4	9	7.7	3.2-14.9	0	0.0	0.0-1.9
1989	19	5.4	3.2-8.5	14	9.3	5.0-15.7	5	2.7	0.9-6.3	11	3.3	1.7-6.0	10	7.5	3.6-13.8	1	0.7	0.0-3.4
1990	25	7.1	4.6-10.5	19	12.6	7.4-20.1	6	3.1	1.1-6.8	4	1.1	0.3-2.9	3	1.8	0.4-5.4	1	0.5	0.0-2.8
1991	23	6.8	4.3-10.2	19	13.7	8.0-21.5	4	2.0	0.6-5.2	13	3.5	1.8-5.9	8	5.2	2.2-10.3	5	2.5	0.8-5.8
1992	20	5.2	3.1-8.0	13	7.5	4.0-12.9	7	3.3	1.3-6.9	10	2.8	1.3-5.1	5	3.1	1.0-7.3	5	2.4	0.8-5.6
1993	22	5.9	3.7-9.0	18	11.1	6.5-17.6	4	1.8	0.5-4.8	3	0.7	0.2-2.2	3	1.7	0.3-5.1	0	0.0	0.0-1.8
1994	19	5.1	3.1-7.9	14	9.4	5.0-15.8	5	2.3	0.7-5.4	9	2.5	1.1-4.7	7	5.4	2.1-11.0	2	1.0	0.1-3.6
1995	18	4.6	2.7-7.2	14	7.7	4.2-13.0	4	1.8	0.5-4.7	1	0.2	0.0-1.4	1	0.5	0.0-3.1	0	0.0	0.0-1.7
1996	15	3.9	2.2-6.3	11	6.4	3.2-11.5	4	1.9	0.5-4.8	7	1.8	0.7-3.7	7	4.3	1.7-8.9	0	0.0	0.0-1.7
1997	23	5.8	3.7-8.7	20	11.5	6.9-17.8	3	1.3	0.3-3.9	8	2.0	0.9-4.0	7	4.4	1.7-9.0	1	0.5	0.0-2.6
1998	15	3.6	2.0-5.9	11	5.7	2.8-10.3	4	1.8	0.5-4.5	7	1.7	0.7-3.6	4	2.7	0.7-6.8	3	1.2	0.3-3.7
1999	15	3.4	1.9-5.7	11	5.4	2.7-9.8	4	1.6	0.4-4.3	6	1.4	0.5-3.1	5	2.9	0.9-6.7	1	0.4	0.0-2.4
2000	29	6.3	4.2-9.1	22	10.5	6.5-16.1	7	2.9	1.2-6.1	9	2.1	1.0-4.0	6	3.6	1.3-7.9	3	1.2	0.3-3.7
2001	25	5.4	3.5-8.1	20	10.3	6.2-16.1	5	2.0	0.7-4.8	7	1.6	0.6-3.3	7	4.0	1.6-8.3	0	0.0	0.0-1.6
2002	18	4.0	2.3-6.3	15	7.5	4.1-12.5	3	1.2	0.3-3.6	11	2.3	1.2-4.2	10	5.0	2.4-9.4	1	0.3	0.0-2.2
2003	21	4.5	2.8-7.0	15	7.2	4.0-12.0	6	2.5	0.9-5.4	4	0.9	0.2-2.3	2	1.1	0.1-3.9	2	0.9	0.1-3.1
2004	28	5.6	3.7-8.2	22	10.5	6.5-16.2	6	2.2	0.8-4.9	5	1.0	0.3-2.5	5	2.6	0.8-6.2	0	0.0	0.0-1.6
2005	22	4.5	2.8-6.8	18	8.5	5.0-13.6	4	1.4	0.4-3.7	2	0.5	0.1-1.7	2	1.1	0.1-3.8	0	0.0	0.0-1.6
2006	24	4.9	3.1-7.3	19	8.7	5.2-13.7	5	1.9	0.6-4.5	8	1.8	0.8-3.5	7	3.8	1.5-7.8	1	0.4	0.0-2.2
2007	29	5.8	3.8-8.4	25	11.0	7.0-16.4	4	1.4	0.4-3.8	8	1.5	0.6-3.1	6	2.5	0.9-5.6	2	0.7	0.1-2.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Larynx

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	27	5.5	3.6-8.0	20	9.3	5.6-14.5	7	2.6	1.0-5.5	14	2.7	1.5-4.6	13	5.7	3.0-9.9	1	0.4	0.0-2.2
2009	21	3.9	2.4-6.0	18	7.3	4.3-11.8	3	0.9	0.2-3.0	4	0.7	0.2-2.0	4	1.6	0.4-4.3	0	0.0	0.0-1.6
2010	26	5.1	3.3-7.6	17	7.4	4.2-12.0	9	3.4	1.5-6.5	9	1.7	0.8-3.3	6	2.8	1.0-6.1	3	1.0	0.2-3.2
2011	21	3.8	2.3-5.8	19	7.8	4.6-12.4	2	0.6	0.1-2.5	8	1.5	0.6-3.1	8	3.7	1.5-7.4	0	0.0	0.0-1.5
2012	26	4.5	2.9-6.7	21	8.2	4.9-12.7	5	1.7	0.5-4.1	3	0.6	0.1-1.8	1	0.5	0.0-2.8	2	0.7	0.1-2.7
2013	25	4.4	2.8-6.5	17	6.4	3.7-10.5	8	2.6	1.1-5.4	7	1.3	0.5-2.8	4	2.0	0.5-5.2	3	1.0	0.2-3.1
2014	20	3.8	2.3-5.9	13	5.4	2.8-9.4	7	2.4	1.0-5.2	5	0.9	0.3-2.2	5	2.2	0.7-5.2	0	0.0	0.0-1.5
2015	23	3.8	2.4-5.8	19	7.5	4.4-11.9	4	1.2	0.3-3.3	6	1.0	0.4-2.4	5	2.1	0.7-5.1	1	0.3	0.0-2.0
2016	20	2.9	1.8-4.6	18	6.0	3.5-9.8	2	0.5	0.1-2.2	3	0.4	0.1-1.4	2	0.6	0.1-2.5	1	0.2	0.0-1.9
2017	22	3.9	2.4-6.0	19	8.1	4.8-12.8	3	1.0	0.2-3.2	2	0.3	0.0-1.3	1	0.4	0.0-2.3	1	0.2	0.0-1.9
2018	19	3.0	1.8-4.8	13	4.5	2.3-7.9	6	1.8	0.6-4.1	7	1.1	0.4-2.4	3	0.9	0.2-3.1	4	1.1	0.3-3.2
2019	16	3.0	1.7-5.0	12	4.9	2.5-8.7	4	1.4	0.4-3.7	6	1.0	0.4-2.3	4	1.5	0.4-4.1	2	0.7	0.1-2.7
2020	11	1.5	0.7-2.9	8	2.5	1.0-5.2	3	0.8	0.2-2.7	3	0.4	0.1-1.5	2	0.8	0.1-3.0	1	0.3	0.0-2.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Richmond (Staten Island)
Lung and bronchus**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	172	63.0	53.8-73.4	123	107.1	88.3-128.6	49	30.7	22.6-40.7	153	56.8	48.0-66.7	116	102.8	84.2-124.1	37	23.9	16.7-33.0
1977	159	57.4	48.6-67.3	114	100.6	81.9-122.1	45	27.8	20.2-37.4	136	47.8	39.9-56.7	95	80.3	64.1-99.1	41	24.6	17.6-33.5
1978	175	61.9	52.9-72.0	125	109.7	90.4-131.7	50	30.6	22.6-40.5	146	50.8	42.7-59.9	113	93.8	76.6-113.6	33	20.0	13.7-28.2
1979	203	69.4	60.0-79.8	140	116.2	96.8-138.2	63	38.9	29.8-49.8	152	52.4	44.2-61.5	109	93.4	75.9-113.6	43	26.0	18.7-35.1
1980	220	71.5	62.2-81.8	150	121.5	101.6-144.0	70	40.2	31.2-51.0	175	58.2	49.8-67.7	121	102.4	83.8-123.7	54	32.2	24.1-42.2
1981	213	69.5	60.3-79.7	148	120.4	100.7-142.5	65	36.3	27.9-46.4	194	62.9	54.1-72.5	129	102.6	84.6-123.0	65	37.6	28.8-48.0
1982	231	75.1	65.6-85.6	160	127.8	107.8-150.2	71	39.6	30.8-50.1	193	62.5	53.9-72.1	135	105.1	87.3-125.3	58	32.8	24.8-42.4
1983	220	68.8	59.9-78.7	158	122.8	103.3-144.7	62	33.7	25.7-43.3	190	59.6	51.3-68.9	125	98.9	81.3-119.0	65	35.1	27.0-44.9
1984	225	72.0	62.7-82.2	148	115.5	96.7-136.8	77	42.6	33.5-53.3	167	53.8	45.8-62.7	117	92.9	76.1-112.2	50	27.5	20.4-36.4
1985	259	79.9	70.3-90.4	179	134.7	114.6-157.1	80	43.8	34.6-54.7	208	64.5	55.9-74.0	146	112.0	93.7-132.7	62	32.9	25.1-42.2
1986	232	71.1	62.1-81.0	150	112.4	94.5-132.6	82	42.8	34.0-53.2	204	62.4	54.1-71.7	130	98.1	81.2-117.3	74	38.7	30.3-48.7
1987	256	77.5	68.2-87.8	162	122.9	103.7-144.4	94	49.0	39.5-60.1	194	58.2	50.2-67.1	128	94.5	78.0-113.3	66	34.5	26.6-44.1
1988	281	84.0	74.3-94.5	191	143.1	122.6-165.9	90	45.9	36.8-56.5	171	50.7	43.3-59.0	124	91.2	75.1-109.6	47	24.1	17.6-32.1
1989	254	76.2	67.0-86.2	157	112.9	95.3-132.7	97	51.9	42.0-63.4	224	67.2	58.6-76.7	155	114.9	96.6-135.4	69	35.5	27.5-45.0
1990	259	75.7	66.6-85.6	143	101.9	85.2-120.8	116	58.6	48.3-70.3	202	58.6	50.7-67.3	105	73.6	59.8-89.4	97	49.0	39.7-59.9
1991	250	72.1	63.3-81.7	139	97.6	81.4-115.9	111	55.8	45.9-67.3	207	59.5	51.6-68.2	130	95.7	79.3-114.3	77	38.2	30.1-47.8
1992	329	91.4	81.7-101.9	194	128.5	110.5-148.5	135	65.9	55.2-78.1	234	66.6	58.3-75.8	139	99.5	83.0-118.1	95	46.3	37.4-56.6
1993	305	84.3	75.0-94.4	179	125.6	107.0-146.2	126	60.1	50.0-71.6	264	73.2	64.6-82.6	155	108.5	91.4-127.8	109	51.9	42.6-62.6
1994	307	82.6	73.6-92.4	166	104.9	89.1-122.6	141	67.4	56.7-79.5	233	62.9	55.1-71.6	125	82.8	68.4-99.1	108	51.8	42.5-62.6
1995	290	75.9	67.4-85.2	157	97.5	82.5-114.4	133	60.9	51.0-72.2	218	57.7	50.2-65.9	127	80.6	66.8-96.3	91	41.3	33.2-50.7
1996	308	79.4	70.7-88.8	157	98.1	83.0-115.1	151	68.2	57.7-80.0	228	59.2	51.7-67.4	118	73.1	60.2-87.8	110	49.4	40.6-59.6
1997	297	74.7	66.5-83.8	161	98.6	83.6-115.5	136	60.4	50.6-71.4	227	57.6	50.4-65.6	124	77.0	63.7-92.2	103	45.5	37.2-55.3
1998	306	74.8	66.6-83.6	167	98.9	84.1-115.5	139	59.6	50.1-70.4	249	61.2	53.8-69.3	142	87.7	73.5-103.8	107	45.7	37.5-55.3
1999	294	70.0	62.2-78.5	151	85.8	72.3-100.9	143	60.0	50.6-70.7	243	58.2	51.1-66.0	132	77.9	64.9-92.7	111	46.5	38.2-56.0
2000	334	77.6	69.5-86.5	185	103.9	89.1-120.3	149	61.0	51.5-71.6	229	54.0	47.3-61.5	138	81.1	67.7-96.2	91	36.7	29.5-45.1
2001	304	68.6	61.0-76.8	168	88.5	75.3-103.3	136	54.4	45.7-64.5	227	51.9	45.3-59.1	124	67.9	56.2-81.3	103	41.1	33.5-49.9
2002	335	74.5	66.7-82.9	170	87.7	74.7-102.2	165	64.6	55.1-75.4	250	55.9	49.2-63.3	135	72.3	60.4-85.9	115	44.9	37.1-54.0
2003	301	66.9	59.5-75.0	174	92.2	78.8-107.2	127	49.1	40.9-58.5	238	52.3	45.9-59.5	127	65.1	54.0-77.7	111	42.7	35.1-51.5
2004	333	72.0	64.4-80.3	183	95.4	81.7-110.7	150	57.0	48.2-67.0	247	53.9	47.3-61.1	148	79.3	66.7-93.5	99	37.4	30.4-45.7
2005	337	72.1	64.5-80.3	173	87.5	74.7-101.8	164	61.7	52.6-72.1	234	51.2	44.8-58.3	138	71.9	60.1-85.2	96	37.3	30.2-45.6
2006	326	70.0	62.5-78.1	164	85.2	72.3-99.6	162	60.4	51.3-70.5	267	57.9	51.1-65.4	144	73.1	61.4-86.4	123	46.5	38.6-55.6
2007	337	69.6	62.3-77.5	161	79.5	67.3-93.1	176	63.3	54.2-73.6	224	46.8	40.8-53.5	113	59.8	49.0-72.2	111	39.7	32.6-48.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Lung and bronchus

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	323	66.3	59.2-74.1	166	79.4	67.5-92.8	157	57.4	48.7-67.3	230	46.5	40.6-53.0	119	57.3	47.2-69.0	111	39.3	32.2-47.5
2009	376	75.3	67.8-83.5	202	95.9	82.7-110.5	174	61.2	52.3-71.2	229	47.0	41.0-53.5	123	60.5	50.0-72.5	106	37.5	30.6-45.5
2010	347	69.1	61.9-76.9	188	85.9	73.7-99.6	159	57.5	48.8-67.4	238	48.5	42.4-55.2	125	61.1	50.5-73.1	113	40.5	33.2-48.9
2011	327	62.8	56.0-70.1	170	74.5	63.4-87.1	157	54.1	45.8-63.5	224	42.9	37.3-49.0	118	54.5	44.8-65.7	106	35.1	28.7-42.7
2012	318	59.7	53.2-66.8	151	65.2	54.9-76.9	167	57.0	48.5-66.6	200	37.6	32.5-43.3	106	44.5	36.1-54.2	94	32.0	25.7-39.4
2013	367	68.3	61.3-75.8	200	84.9	73.2-98.0	167	55.7	47.4-65.1	237	44.5	38.9-50.7	127	57.1	47.4-68.3	110	36.3	29.7-44.0
2014	390	68.3	61.6-75.7	198	79.5	68.5-91.9	192	59.7	51.4-69.1	199	35.9	31.0-41.4	113	46.3	37.9-56.1	86	27.7	22.1-34.5
2015	376	66.7	60.0-74.0	190	77.1	66.2-89.3	186	59.2	50.8-68.6	238	41.3	36.1-47.0	122	50.4	41.5-60.5	116	35.1	28.9-42.3
2016	367	63.5	57.1-70.5	169	68.8	58.5-80.3	198	60.4	52.1-69.7	213	37.5	32.6-43.1	117	49.6	40.8-59.7	96	29.5	23.8-36.4
2017	408	70.5	63.7-77.8	196	77.8	67.0-89.9	212	65.8	57.1-75.5	201	34.4	29.7-39.7	96	37.9	30.5-46.6	105	31.5	25.7-38.5
2018	380	62.0	55.8-68.8	170	62.3	53.1-72.8	210	62.6	54.2-71.9	185	31.2	26.8-36.2	82	32.9	26.0-41.1	103	30.2	24.5-36.9
2019	418	66.8	60.4-73.7	200	73.3	63.2-84.6	218	63.2	55.0-72.5	202	32.7	28.3-37.7	105	40.0	32.5-48.7	97	27.6	22.3-34.0
2020	351	55.9	50.1-62.3	175	64.2	54.8-74.8	176	49.9	42.7-58.1	184	30.0	25.7-34.8	97	37.1	29.9-45.6	87	25.4	20.3-31.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Melanoma of the skin

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	25	9.1	5.8-13.6	14	11.8	6.0-20.6	11	7.1	3.5-12.8	5	2.4	0.8-5.5	1	0.8	0.0-4.6	4	3.2	0.9-8.0
1977	21	7.1	4.3-10.9	9	6.9	3.0-13.4	12	7.2	3.7-12.7	2	0.9	0.1-3.0	2	2.4	0.2-8.4	0	0.0	0.0-2.3
1978	16	5.6	3.2-9.1	10	6.8	3.2-12.7	6	4.3	1.6-9.1	7	2.5	1.0-5.1	4	3.4	0.8-8.9	3	1.9	0.4-5.6
1979	21	7.1	4.4-11.0	11	7.7	3.8-13.9	10	6.6	3.1-12.1	6	2.0	0.7-4.4	4	2.4	0.6-6.6	2	1.3	0.2-4.6
1980	23	7.3	4.5-11.0	9	6.6	2.9-12.9	14	8.2	4.4-14.0	6	1.9	0.7-4.2	3	1.6	0.3-5.3	3	1.9	0.4-5.5
1981	33	9.8	6.6-13.8	13	8.7	4.3-15.5	20	11.0	6.7-17.2	9	3.0	1.3-5.7	4	2.7	0.7-7.0	5	3.1	1.0-7.2
1982	14	4.7	2.5-7.9	10	7.4	3.4-13.7	4	2.6	0.7-6.4	7	1.9	0.8-4.0	6	3.4	1.2-7.8	1	0.5	0.0-2.9
1983	21	7.1	4.3-10.8	13	8.7	4.6-15.1	8	5.1	2.2-10.0	8	2.3	1.0-4.7	5	2.8	0.9-6.9	3	2.0	0.4-5.5
1984	30	9.7	6.5-13.9	18	14.1	8.1-22.5	12	6.8	3.5-12.0	7	2.2	0.9-4.5	6	4.3	1.5-9.5	1	0.7	0.0-3.4
1985	31	10.0	6.7-14.2	20	15.5	9.3-24.2	11	6.1	3.0-10.9	13	4.5	2.4-7.6	5	3.8	1.2-8.8	8	5.0	2.2-9.8
1986	36	10.4	7.2-14.5	18	11.2	6.5-18.0	18	9.3	5.4-14.7	9	2.8	1.2-5.3	5	3.0	0.9-7.2	4	2.3	0.6-5.9
1987	25	7.5	4.8-11.1	14	10.2	5.3-17.4	11	5.9	2.9-10.5	7	2.3	0.9-4.6	6	5.2	1.7-11.3	1	0.5	0.0-2.9
1988	39	11.3	8.0-15.5	19	13.6	7.9-21.6	20	10.0	6.1-15.6	8	2.2	0.9-4.4	8	4.8	2.1-9.7	0	0.0	0.0-1.9
1989	38	11.3	8.0-15.5	26	18.8	12.0-28.0	12	6.2	3.2-10.9	6	1.8	0.6-3.8	3	1.9	0.4-5.6	3	1.5	0.3-4.5
1990	31	8.8	5.9-12.5	17	11.2	6.4-18.2	14	7.4	4.0-12.4	4	1.2	0.3-3.1	3	1.8	0.4-5.5	1	0.7	0.0-3.3
1991	35	9.7	6.7-13.6	23	13.9	8.8-21.1	12	6.1	3.1-10.7	11	2.8	1.4-5.0	7	4.1	1.6-8.5	4	1.7	0.5-4.6
1992	28	7.7	5.1-11.1	12	7.7	3.9-13.5	16	8.1	4.6-13.2	3	0.9	0.2-2.7	2	1.9	0.2-6.6	1	0.5	0.0-2.8
1993	27	7.3	4.8-10.6	18	11.6	6.7-18.5	9	4.3	2.0-8.2	3	0.9	0.2-2.5	3	2.1	0.4-6.1	0	0.0	0.0-1.8
1994	31	8.2	5.6-11.7	15	9.9	5.3-16.6	16	7.5	4.3-12.3	8	2.2	0.9-4.2	4	2.6	0.7-6.6	4	2.0	0.5-5.1
1995	30	7.5	5.0-10.7	25	14.1	9.1-21.0	5	2.3	0.7-5.4	8	2.1	0.9-4.2	6	3.6	1.3-7.8	2	1.0	0.1-3.6
1996	33	8.3	5.7-11.6	15	8.5	4.6-14.2	18	8.3	4.9-13.1	8	2.1	0.9-4.2	3	1.6	0.3-4.8	5	2.3	0.8-5.5
1997	42	10.3	7.4-13.9	19	9.9	6.0-15.6	23	10.0	6.3-15.1	11	2.7	1.3-4.8	6	3.2	1.2-7.1	5	2.2	0.7-5.2
1998	31	7.4	5.0-10.4	21	11.1	6.8-17.1	10	4.2	2.0-7.9	6	1.4	0.5-3.1	3	1.6	0.3-4.7	3	1.2	0.3-3.7
1999	45	10.5	7.6-14.0	23	12.3	7.7-18.6	22	9.2	5.8-14.0	7	1.6	0.6-3.3	4	1.8	0.5-4.8	3	1.3	0.3-3.7
2000	49	10.8	8.0-14.3	27	13.3	8.7-19.5	22	9.0	5.7-13.7	12	2.7	1.4-4.8	7	3.5	1.4-7.2	5	2.1	0.7-4.9
2001	52	11.6	8.6-15.2	27	13.1	8.6-19.2	25	10.2	6.6-15.1	8	1.8	0.8-3.5	6	3.1	1.1-6.8	2	0.8	0.1-2.9
2002	52	11.3	8.4-14.8	27	12.8	8.4-18.8	25	10.1	6.6-15.0	5	1.1	0.3-2.5	0	0.0	0.0-1.9	5	1.9	0.6-4.5
2003	61	13.1	10.0-16.9	37	18.4	12.8-25.6	24	9.3	5.9-13.9	13	2.8	1.5-4.8	5	2.4	0.8-5.7	8	3.2	1.4-6.3
2004	67	14.5	11.2-18.5	37	18.6	13.0-25.7	30	11.6	7.8-16.7	5	1.0	0.3-2.4	3	1.2	0.3-3.8	2	0.8	0.1-2.9
2005	56	11.7	8.8-15.2	25	11.4	7.3-17.0	31	11.8	8.0-16.8	11	2.3	1.2-4.2	5	2.5	0.8-6.0	6	2.2	0.8-4.9
2006	70	14.9	11.6-18.9	37	17.8	12.5-24.7	33	12.6	8.6-17.8	11	2.3	1.1-4.2	8	4.4	1.8-8.6	3	1.1	0.2-3.4
2007	73	15.2	11.9-19.2	45	21.5	15.5-29.1	28	11.2	7.4-16.3	6	1.1	0.4-2.4	4	1.8	0.5-4.9	2	0.6	0.1-2.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Melanoma of the skin

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	75	15.0	11.8-18.9	46	20.4	14.8-27.4	29	10.6	7.1-15.4	12	2.5	1.3-4.4	9	4.1	1.8-7.9	3	1.2	0.2-3.5
2009	78	15.6	12.3-19.5	33	13.6	9.3-19.3	45	17.4	12.6-23.4	7	1.3	0.5-2.8	6	2.5	0.9-5.7	1	0.3	0.0-2.0
2010	62	12.2	9.3-15.8	34	15.7	10.7-22.1	28	9.7	6.4-14.2	8	1.7	0.7-3.4	5	2.6	0.8-5.9	3	1.1	0.2-3.4
2011	103	20.3	16.5-24.8	59	26.1	19.7-33.9	44	16.4	11.8-22.1	4	0.8	0.2-2.1	1	0.4	0.0-2.3	3	1.0	0.2-3.2
2012	97	17.9	14.4-21.9	45	18.4	13.3-24.9	52	18.1	13.4-23.9	8	1.3	0.6-2.7	6	2.3	0.8-5.2	2	0.7	0.1-2.7
2013	116	21.4	17.6-25.8	81	34.5	27.2-43.2	35	12.0	8.3-16.9	13	2.6	1.4-4.6	8	3.8	1.6-7.5	5	1.9	0.6-4.6
2014	92	16.3	13.0-20.1	44	18.1	13.0-24.5	48	15.9	11.6-21.4	15	2.6	1.4-4.4	11	4.4	2.1-8.1	4	1.3	0.3-3.5
2015	92	17.2	13.8-21.2	55	23.3	17.4-30.6	37	12.8	8.9-17.9	8	1.5	0.7-3.1	5	2.5	0.8-5.8	3	1.0	0.2-3.2
2016	84	14.7	11.7-18.4	45	17.0	12.3-23.1	39	13.0	9.1-18.0	8	1.3	0.6-2.7	4	2.0	0.6-5.1	4	1.1	0.3-3.1
2017	79	14.3	11.2-17.9	48	19.1	13.9-25.6	31	10.8	7.2-15.6	11	1.9	1.0-3.6	4	1.9	0.5-4.9	7	2.1	0.8-4.6
2018	98	16.1	13.0-19.7	56	20.2	15.1-26.5	42	12.6	9.0-17.2	6	1.0	0.3-2.2	4	1.5	0.4-4.1	2	0.6	0.1-2.5
2019	76	13.1	10.2-16.5	38	15.2	10.6-21.0	38	12.0	8.4-16.9	9	1.7	0.8-3.4	6	2.7	1.0-6.0	3	1.0	0.2-3.2
2020	84	13.8	10.9-17.2	41	15.7	11.1-21.6	43	12.3	8.9-16.9	8	1.6	0.7-3.2	3	1.3	0.3-3.9	5	1.8	0.5-4.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Female breast

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	149	102.4	86.3-120.5	52	34.2	25.4-45.0
1977	169	110.8	94.5-129.1	72	47.8	37.2-60.3
1978	173	109.4	93.5-127.3	68	43.4	33.6-55.1
1979	163	101.2	86.0-118.2	54	33.1	24.8-43.4
1980	138	85.2	71.4-100.9	59	34.1	25.9-44.2
1981	193	115.4	99.4-133.2	53	32.0	23.9-42.0
1982	133	78.1	65.2-92.8	54	31.0	23.2-40.5
1983	185	107.1	92.0-123.9	49	28.3	20.9-37.5
1984	218	124.9	108.6-142.9	68	38.5	29.8-49.0
1985	235	131.7	115.2-149.9	68	37.6	29.1-47.8
1986	215	117.2	101.9-134.2	58	32.1	24.3-41.6
1987	263	142.9	125.9-161.4	77	40.9	32.2-51.2
1988	281	147.4	130.5-165.9	81	41.6	33.0-51.8
1989	282	149.9	132.8-168.6	71	39.0	30.4-49.2
1990	269	139.3	123.0-157.1	65	33.8	26.0-43.2
1991	253	128.7	113.3-145.7	71	34.9	27.2-44.1
1992	296	148.5	132.0-166.5	78	38.3	30.2-47.8
1993	277	135.0	119.5-151.9	96	46.5	37.6-56.8
1994	265	125.5	110.8-141.6	83	38.5	30.6-47.8
1995	308	144.5	128.7-161.6	73	33.8	26.4-42.5
1996	293	133.6	118.7-149.9	77	35.0	27.6-43.8
1997	335	148.9	133.3-165.7	75	32.9	25.9-41.3
1998	331	143.3	128.2-159.6	73	31.1	24.4-39.1
1999	343	144.4	129.5-160.6	88	36.4	29.2-44.9
2000	302	124.0	110.4-138.9	61	24.7	18.9-31.8
2001	327	131.1	117.2-146.1	77	30.8	24.3-38.6
2002	330	130.2	116.5-145.2	73	28.2	22.1-35.5
2003	321	124.7	111.4-139.3	81	30.5	24.2-38.0
2004	352	135.0	121.2-150.0	86	31.4	25.1-38.9
2005	340	130.1	116.5-144.8	73	26.9	21.1-34.0
2006	349	131.9	118.3-146.6	67	24.8	19.2-31.6
2007	379	139.4	125.6-154.4	55	19.5	14.6-25.5
2008	349	127.1	113.9-141.3	81	28.4	22.5-35.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Female breast

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	378	134.4	121.0-148.9	62	21.7	16.6-28.1
2010	365	126.5	113.6-140.5	72	25.3	19.7-32.1
2011	397	136.2	122.9-150.7	61	20.6	15.7-26.7
2012	389	134.3	121.0-148.7	57	18.0	13.6-23.6
2013	398	135.1	121.8-149.5	61	19.7	15.0-25.6
2014	394	131.6	118.6-145.8	54	16.1	12.0-21.3
2015	414	140.0	126.4-154.7	52	15.7	11.7-20.9
2016	423	138.4	125.0-152.8	56	17.0	12.7-22.4
2017	393	130.1	117.1-144.2	65	19.0	14.6-24.6
2018	416	132.3	119.5-146.2	59	16.9	12.8-22.1
2019	471	151.5	137.6-166.5	61	17.6	13.4-22.9
2020	450	144.2	130.6-158.8	61	17.4	13.2-22.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Cervix uteri

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	23	15.5	9.7-23.4	5	3.0	1.0-7.1
1977	20	12.4	7.5-19.3	7	4.6	1.8-9.5
1978	15	9.8	5.4-16.2	7	4.8	1.9-9.8
1979	14	8.6	4.6-14.5	3	2.0	0.4-5.7
1980	9	5.3	2.4-10.1	3	2.0	0.4-5.7
1981	22	13.1	8.1-20.0	5	2.8	0.9-6.5
1982	20	11.4	6.9-17.8	6	3.5	1.3-7.6
1983	13	7.1	3.7-12.2	4	2.0	0.5-5.1
1984	17	8.9	5.1-14.5	1	0.7	0.0-3.6
1985	15	7.9	4.3-13.1	5	2.3	0.7-5.5
1986	18	9.6	5.6-15.3	2	1.0	0.1-3.7
1987	23	12.6	7.9-18.9	4	2.1	0.6-5.3
1988	22	11.0	6.9-16.8	6	3.1	1.1-6.9
1989	39	20.3	14.4-27.9	6	3.2	1.2-7.1
1990	32	15.7	10.7-22.2	8	4.2	1.8-8.3
1991	28	14.0	9.3-20.3	2	0.9	0.1-3.4
1992	32	15.2	10.4-21.6	3	1.5	0.3-4.3
1993	17	8.3	4.8-13.3	4	2.0	0.5-5.1
1994	35	16.1	11.2-22.5	10	4.7	2.2-8.7
1995	30	14.0	9.5-20.1	11	5.3	2.6-9.4
1996	33	15.0	10.3-21.1	11	4.8	2.4-8.6
1997	23	10.0	6.3-15.1	5	2.2	0.7-5.3
1998	25	10.9	7.1-16.1	6	2.6	0.9-5.7
1999	25	10.5	6.8-15.6	6	2.6	1.0-5.7
2000	19	7.9	4.8-12.4	9	3.7	1.7-7.1
2001	22	9.2	5.7-13.9	3	1.3	0.3-3.7
2002	25	10.0	6.5-14.8	4	1.5	0.4-4.0
2003	22	8.4	5.2-12.8	6	2.2	0.8-4.9
2004	27	10.5	6.9-15.3	5	2.1	0.7-4.8
2005	31	12.3	8.3-17.5	11	4.0	2.0-7.3
2006	17	7.0	4.0-11.2	7	2.7	1.1-5.6
2007	35	13.8	9.6-19.3	3	1.3	0.3-3.7
2008	29	11.4	7.6-16.4	3	1.0	0.2-3.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Cervix uteri

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	23	8.9	5.6-13.5	8	2.8	1.2-5.7
2010	10	3.6	1.7-6.8	4	1.2	0.3-3.4
2011	17	5.5	3.2-9.0	5	1.6	0.5-3.9
2012	20	7.3	4.4-11.5	8	2.6	1.1-5.4
2013	25	9.9	6.3-14.7	7	2.3	0.9-5.0
2014	20	7.6	4.5-11.9	10	3.6	1.7-6.8
2015	16	6.5	3.6-10.6	7	2.4	0.9-5.2
2016	29	10.4	6.9-15.2	6	2.0	0.7-4.6
2017	22	8.3	5.1-12.8	4	1.1	0.3-3.2
2018	28	9.3	6.1-13.8	7	1.9	0.7-4.2
2019	14	5.3	2.8-9.0	4	1.6	0.4-4.1
2020	18	6.0	3.4-9.9	6	1.7	0.6-4.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Corpus uteris and uterus, not otherwise specified

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	36	22.7	15.8-31.5	8	5.3	2.3-10.5
1977	27	16.6	10.9-24.2	12	7.4	3.8-12.9
1978	33	20.2	13.9-28.5	10	6.3	3.0-11.6
1979	38	22.5	15.9-31.1	9	5.6	2.5-10.5
1980	29	17.3	11.5-24.9	11	6.6	3.2-11.8
1981	33	18.1	12.4-25.5	8	4.6	1.9-9.0
1982	33	19.1	13.1-27.0	9	5.6	2.6-10.5
1983	46	25.8	18.8-34.5	4	2.0	0.5-5.3
1984	41	22.9	16.4-31.2	5	2.7	0.9-6.3
1985	46	25.5	18.6-34.0	5	2.9	0.9-6.7
1986	46	23.8	17.3-31.9	7	3.9	1.6-8.1
1987	45	23.4	17.0-31.4	11	5.8	2.9-10.5
1988	53	27.0	20.1-35.3	7	3.6	1.4-7.4
1989	61	30.9	23.6-39.8	16	8.0	4.6-13.1
1990	47	23.7	17.4-31.6	3	1.4	0.3-4.3
1991	58	30.1	22.8-39.0	7	3.6	1.4-7.4
1992	45	22.8	16.6-30.5	6	2.8	1.0-6.2
1993	63	30.2	23.2-38.7	22	10.3	6.4-15.6
1994	57	26.8	20.3-34.7	15	6.9	3.9-11.5
1995	53	25.2	18.9-33.0	14	6.2	3.4-10.5
1996	47	21.5	15.8-28.7	11	4.8	2.4-8.7
1997	65	29.2	22.5-37.2	11	4.8	2.4-8.7
1998	81	34.9	27.7-43.4	10	4.3	2.1-8.0
1999	70	29.3	22.8-37.1	10	4.1	2.0-7.6
2000	76	31.1	24.5-39.0	11	4.3	2.2-7.8
2001	79	31.5	24.9-39.3	14	5.4	3.0-9.2
2002	53	20.8	15.5-27.2	15	5.7	3.2-9.5
2003	68	26.2	20.3-33.3	14	5.5	3.0-9.3
2004	102	38.1	31.1-46.4	17	6.3	3.7-10.2
2005	81	29.9	23.7-37.2	16	6.0	3.4-9.9
2006	80	29.1	23.0-36.4	16	5.7	3.2-9.3
2007	89	31.9	25.6-39.4	14	5.6	3.0-9.4
2008	102	35.3	28.7-43.1	19	7.0	4.2-11.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Corpus uteris and uterus, not otherwise specified

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	96	32.4	26.1-39.8	10	3.5	1.7-6.6
2010	98	34.0	27.5-41.7	16	5.7	3.2-9.5
2011	101	34.1	27.6-41.6	15	5.2	2.9-8.8
2012	108	35.6	29.1-43.3	14	4.5	2.4-7.7
2013	113	36.7	30.1-44.4	17	5.7	3.3-9.4
2014	105	33.9	27.6-41.3	17	5.1	2.9-8.5
2015	116	36.6	30.1-44.3	24	7.6	4.9-11.6
2016	109	34.6	28.3-42.1	24	7.9	5.0-12.0
2017	129	38.7	32.1-46.3	19	5.7	3.4-9.1
2018	121	35.6	29.4-42.8	22	6.0	3.8-9.4
2019	132	39.8	33.1-47.6	20	5.7	3.4-9.1
2020	91	25.9	20.7-32.1	25	7.2	4.6-10.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Ovary

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	24	15.3	9.7-22.9	12	8.3	4.3-14.4
1977	22	14.2	8.8-21.5	22	14.0	8.7-21.3
1978	22	12.8	7.9-19.4	11	6.8	3.4-12.3
1979	26	15.8	10.2-23.2	16	9.0	5.1-14.8
1980	20	11.7	7.1-18.1	21	12.2	7.5-18.7
1981	23	14.4	9.1-21.7	12	6.6	3.4-11.6
1982	26	14.9	9.6-21.9	6	3.1	1.1-7.0
1983	28	15.4	10.2-22.3	15	8.4	4.6-13.9
1984	32	17.3	11.8-24.6	19	11.0	6.6-17.2
1985	35	19.4	13.5-27.1	20	11.1	6.8-17.2
1986	30	16.3	10.9-23.4	19	9.6	5.7-15.1
1987	35	18.5	12.8-25.9	13	7.3	3.8-12.4
1988	51	26.1	19.4-34.4	26	12.9	8.4-19.0
1989	43	22.9	16.5-30.9	19	9.6	5.7-15.0
1990	40	20.4	14.5-27.8	23	12.0	7.6-18.0
1991	57	28.9	21.8-37.5	28	14.1	9.3-20.4
1992	43	21.8	15.7-29.3	32	15.7	10.7-22.2
1993	35	17.2	12.0-23.9	24	11.7	7.5-17.4
1994	43	19.9	14.4-26.9	26	12.2	7.9-17.9
1995	44	20.5	14.9-27.5	15	7.0	3.9-11.6
1996	37	17.3	12.1-23.8	25	11.6	7.5-17.1
1997	47	20.8	15.3-27.7	17	7.5	4.3-12.0
1998	40	17.4	12.4-23.7	12	5.1	2.6-8.9
1999	42	17.9	12.9-24.2	27	11.3	7.4-16.5
2000	40	16.2	11.6-22.1	20	8.1	5.0-12.6
2001	36	14.3	10.0-19.9	28	11.3	7.5-16.4
2002	44	17.3	12.5-23.2	23	8.8	5.6-13.3
2003	25	9.7	6.3-14.4	28	10.6	7.0-15.3
2004	33	12.4	8.5-17.5	28	10.4	6.9-15.2
2005	25	9.3	6.0-13.8	20	7.5	4.5-11.6
2006	24	9.2	5.9-13.7	33	12.3	8.4-17.3
2007	48	17.5	12.9-23.4	22	7.9	4.9-12.1
2008	45	15.9	11.5-21.4	31	10.9	7.4-15.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Ovary

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	35	12.3	8.5-17.2	23	7.9	5.0-12.0
2010	36	12.8	8.9-17.9	19	6.3	3.8-10.1
2011	40	14.1	10.0-19.4	25	8.3	5.3-12.5
2012	31	10.6	7.1-15.2	30	9.9	6.6-14.2
2013	43	14.1	10.1-19.2	30	9.4	6.3-13.6
2014	40	13.9	9.8-19.2	12	3.7	1.9-6.7
2015	38	12.7	8.9-17.7	18	5.9	3.5-9.6
2016	48	15.6	11.4-21.0	20	5.5	3.3-8.7
2017	45	13.3	9.7-18.1	26	7.7	4.9-11.5
2018	38	12.8	9.0-17.9	27	7.5	4.9-11.1
2019	45	14.9	10.7-20.3	24	7.2	4.6-11.0
2020	52	18.1	13.3-24.1	17	5.2	3.0-8.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Prostate

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	99	105.1	84.5-128.8	17	19.4	10.9-31.3
1977	65	64.6	49.0-83.2	23	26.2	16.2-39.6
1978	69	67.4	51.8-86.0	31	31.8	21.2-45.4
1979	88	92.3	72.9-114.8	30	33.4	22.0-48.0
1980	76	76.5	59.3-96.5	23	25.4	15.6-38.3
1981	88	78.5	62.1-97.7	38	40.1	27.7-55.4
1982	102	90.6	72.8-111.0	35	37.4	25.5-52.3
1983	79	72.5	56.6-91.1	38	35.5	24.6-49.1
1984	76	68.8	53.4-86.9	31	29.7	19.8-42.4
1985	96	82.5	65.9-101.7	34	28.4	19.4-39.9
1986	104	91.0	73.4-111.1	27	23.9	15.4-35.1
1987	95	79.5	63.5-98.0	35	34.2	23.2-48.0
1988	96	75.5	60.5-92.8	30	29.1	19.2-41.7
1989	111	100.4	81.5-122.0	46	39.7	28.6-53.2
1990	126	97.0	79.9-116.4	47	44.2	31.7-59.4
1991	168	128.9	109.3-150.8	38	34.8	24.1-48.0
1992	262	190.0	166.8-215.3	46	39.8	28.7-53.4
1993	243	174.0	151.9-198.2	35	31.6	21.6-44.1
1994	181	118.9	101.6-138.1	49	39.3	28.6-52.4
1995	199	125.2	107.8-144.4	43	34.6	24.6-46.8
1996	226	139.6	121.5-159.5	54	41.5	30.8-54.4
1997	263	157.1	138.2-177.8	48	36.3	26.5-48.3
1998	294	168.1	149.1-188.8	43	30.5	21.8-41.2
1999	306	171.1	152.0-191.8	38	26.3	18.4-36.1
2000	382	208.9	188.1-231.5	41	28.7	20.4-38.9
2001	351	187.7	168.3-208.8	36	25.2	17.5-34.9
2002	345	172.7	154.7-192.2	38	25.5	17.9-35.0
2003	304	155.4	138.0-174.3	36	22.7	15.8-31.5
2004	285	137.6	121.7-155.0	39	24.1	17.0-33.0
2005	294	141.0	125.0-158.6	36	21.7	15.1-30.2
2006	371	175.6	157.7-194.9	36	22.1	15.4-30.5
2007	374	176.2	158.3-195.6	29	16.0	10.6-23.0
2008	403	179.5	161.9-198.5	37	21.5	15.0-29.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Prostate

Year of diagnosis/death	Incidence			Mortality		
	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	352	149.4	133.7-166.4	24	13.8	8.8-20.5
2010	318	133.5	118.7-149.6	37	20.3	14.2-28.0
2011	349	141.3	126.4-157.5	30	16.7	11.2-23.9
2012	336	132.7	118.5-148.3	26	14.0	9.1-20.5
2013	321	122.9	109.4-137.7	43	21.8	15.7-29.4
2014	330	122.7	109.4-137.2	35	17.2	11.9-23.9
2015	315	116.1	103.3-130.2	44	19.8	14.3-26.7
2016	349	125.2	112.0-139.6	39	18.3	13.0-25.2
2017	325	112.6	100.4-126.0	30	13.5	9.0-19.4
2018	326	112.4	100.2-125.7	40	17.1	12.1-23.4
2019	411	140.4	126.8-155.1	34	14.5	9.9-20.3
2020	326	107.0	95.4-119.7	36	14.8	10.3-20.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Testis

Year of diagnosis	Incidence		
	Cases	Rate per 100,000	95% CI
1976	7	4.1	1.6-9.0
1977	5	2.7	0.9-6.8
1978	9	5.1	2.3-10.1
1979	3	1.4	0.3-4.9
1980	4	2.4	0.6-6.5
1981	2	0.9	0.1-4.1
1982	11	5.3	2.6-10.0
1983	11	6.0	2.9-11.1
1984	10	5.1	2.4-9.8
1985	8	3.9	1.6-8.1
1986	12	5.8	3.0-10.6
1987	8	3.8	1.6-8.0
1988	6	3.0	1.1-7.0
1989	6	2.7	1.0-6.4
1990	12	5.6	2.9-10.3
1991	13	6.2	3.2-11.0
1992	9	4.6	2.1-9.0
1993	19	8.7	5.2-13.9
1994	9	4.0	1.8-7.9
1995	15	7.0	3.9-11.8
1996	13	5.9	3.1-10.3
1997	15	6.8	3.8-11.4
1998	10	4.6	2.2-8.6
1999	27	12.1	7.9-17.7
2000	10	4.7	2.2-8.8
2001	16	7.2	4.1-11.8
2002	13	5.9	3.2-10.2
2003	15	6.7	3.7-11.1
2004	7	3.1	1.2-6.5
2005	17	7.6	4.4-12.2
2006	11	4.9	2.4-8.8
2007	15	7.0	3.9-11.6
2008	15	7.1	4.0-11.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

Mortality data not provided due to small number of deaths per year.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Testis

Year of diagnosis	Incidence		
	Cases	Rate per 100,000	95% CI
2009	16	6.9	3.9-11.3
2010	13	5.9	3.1-10.2
2011	16	7.3	4.2-11.9
2012	12	5.7	2.9-9.9
2013	10	4.6	2.2-8.5
2014	12	4.7	2.4-8.4
2015	20	9.4	5.7-14.5
2016	12	5.9	3.0-10.3
2017	9	4.2	1.9-8.1
2018	13	5.9	3.1-10.0
2019	14	6.3	3.4-10.7
2020	23	10.4	6.6-15.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

Mortality data not provided due to small number of deaths per year.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Richmond (Staten Island)
Urinary bladder (incl. in situ)**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	51	20.1	14.9-26.5	39	36.7	25.4-50.9	12	8.4	4.3-14.6	22	9.4	5.8-14.2	18	19.0	10.9-30.3	4	3.2	0.9-7.9
1977	64	24.2	18.5-31.0	43	37.5	26.5-51.3	21	14.3	8.8-21.8	14	5.7	3.1-9.6	6	5.9	2.1-12.7	8	5.3	2.3-10.5
1978	56	19.8	14.9-25.8	42	33.3	23.5-45.6	14	9.5	5.2-15.8	14	5.4	2.9-9.1	9	8.5	3.7-16.3	5	3.4	1.1-7.7
1979	81	29.9	23.6-37.2	58	55.1	41.1-71.8	23	14.1	8.9-21.2	20	7.4	4.5-11.5	12	10.4	5.1-18.6	8	5.4	2.3-10.4
1980	79	27.4	21.6-34.3	58	52.5	39.0-68.7	21	12.1	7.4-18.5	19	7.4	4.4-11.5	14	14.1	7.4-23.7	5	3.2	1.0-7.4
1981	82	27.6	21.8-34.3	65	54.7	41.3-70.7	17	9.6	5.6-15.4	16	6.0	3.4-9.7	10	10.4	4.8-19.0	6	3.5	1.3-7.7
1982	73	23.6	18.4-29.8	54	44.3	32.5-58.5	19	10.6	6.3-16.6	15	5.1	2.8-8.4	12	11.2	5.5-19.9	3	1.7	0.3-5.0
1983	73	24.3	18.9-30.6	56	46.9	34.8-61.4	17	9.3	5.4-15.0	18	6.7	3.9-10.5	9	9.3	4.0-17.6	9	5.3	2.4-10.0
1984	73	24.0	18.7-30.3	52	44.9	32.9-59.5	21	11.4	7.0-17.5	14	4.7	2.5-7.9	9	9.1	3.9-17.3	5	2.5	0.8-6.0
1985	81	25.2	20.0-31.4	61	45.4	34.3-58.8	20	10.9	6.7-16.9	21	6.8	4.2-10.4	18	15.0	8.6-24.0	3	1.6	0.3-4.7
1986	77	23.9	18.8-30.0	57	44.2	32.7-58.1	20	11.1	6.7-17.2	12	3.9	2.0-6.8	8	6.7	2.7-13.3	4	2.2	0.6-5.5
1987	87	26.2	20.9-32.4	62	44.4	33.7-57.4	25	13.1	8.4-19.3	16	5.4	3.1-8.8	11	8.6	4.1-15.5	5	2.8	0.9-6.5
1988	90	27.9	22.3-34.4	63	49.6	37.4-64.2	27	14.1	9.2-20.5	24	7.9	5.1-11.8	19	17.3	10.2-26.9	5	2.6	0.9-6.2
1989	79	24.3	19.2-30.4	62	48.0	36.2-62.2	17	9.0	5.2-14.4	27	9.2	6.0-13.3	19	19.5	11.3-30.5	8	4.4	1.9-8.6
1990	83	25.3	20.1-31.4	53	40.2	29.6-53.2	30	15.6	10.5-22.2	19	5.7	3.4-8.9	11	8.2	3.9-14.7	8	4.1	1.8-8.1
1991	86	24.8	19.8-30.7	64	46.3	35.1-59.6	22	10.9	6.8-16.5	20	6.0	3.6-9.3	15	12.5	6.7-20.9	5	2.4	0.8-5.6
1992	102	29.2	23.7-35.5	72	53.6	41.3-68.1	30	14.3	9.6-20.5	23	6.7	4.3-10.1	15	12.1	6.5-20.2	8	3.7	1.6-7.4
1993	83	23.3	18.5-28.9	58	40.2	30.1-52.5	25	12.2	7.9-18.0	29	8.3	5.5-11.9	19	14.6	8.5-23.0	10	4.8	2.3-8.8
1994	85	22.8	18.2-28.2	61	37.7	28.6-48.6	24	11.2	7.2-16.7	23	6.3	4.0-9.4	19	13.2	7.8-20.7	4	1.9	0.5-4.8
1995	99	26.4	21.5-32.2	72	48.6	37.5-61.8	27	12.1	7.9-17.6	30	8.3	5.6-11.9	20	14.6	8.8-22.4	10	4.4	2.1-8.1
1996	78	20.4	16.1-25.5	57	37.2	27.8-48.5	21	9.7	6.0-14.8	22	6.0	3.7-9.1	11	7.3	3.5-13.3	11	4.6	2.3-8.2
1997	90	23.0	18.5-28.3	65	39.3	30.1-50.4	25	10.6	6.8-15.7	31	8.1	5.5-11.5	19	12.6	7.4-19.8	12	5.0	2.6-8.9
1998	129	31.6	26.3-37.5	87	49.5	39.4-61.4	42	17.5	12.6-23.7	25	6.4	4.1-9.4	19	12.2	7.2-19.0	6	2.5	0.9-5.5
1999	120	28.5	23.6-34.1	89	52.1	41.5-64.4	31	12.7	8.6-18.1	27	6.8	4.4-9.8	18	12.5	7.3-19.6	9	3.6	1.6-6.8
2000	102	23.8	19.4-28.9	77	45.6	35.7-57.1	25	10.1	6.5-14.9	21	5.1	3.1-7.7	15	9.2	5.1-15.0	6	2.3	0.8-5.1
2001	129	30.0	25.1-35.7	88	50.2	40.0-62.0	41	16.8	12.1-22.9	22	5.1	3.2-7.7	13	7.8	4.0-13.3	9	3.6	1.6-6.9
2002	100	22.5	18.3-27.4	79	44.2	34.7-55.4	21	7.9	4.9-12.2	23	5.2	3.3-7.9	16	9.7	5.4-15.9	7	2.8	1.1-5.8
2003	127	28.7	23.9-34.2	95	52.3	42.1-64.2	32	12.2	8.3-17.3	22	5.1	3.2-7.7	14	8.9	4.8-15.0	8	3.2	1.4-6.3
2004	127	27.8	23.2-33.1	91	47.3	37.9-58.3	36	13.6	9.5-18.9	21	4.8	3.0-7.3	15	9.1	5.0-15.0	6	2.3	0.9-5.2
2005	126	27.6	23.0-32.9	86	45.2	36.0-56.1	40	15.1	10.8-20.7	25	5.8	3.8-8.6	17	9.6	5.6-15.4	8	3.0	1.3-6.0
2006	118	25.8	21.3-31.0	90	47.3	37.9-58.3	28	10.5	7.0-15.3	25	5.7	3.7-8.4	19	10.6	6.3-16.5	6	2.0	0.7-4.5
2007	161	33.6	28.6-39.3	115	57.3	47.0-69.0	46	16.5	12.0-22.1	23	4.8	3.0-7.2	19	10.3	6.1-16.2	4	1.5	0.4-3.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Richmond (Staten Island)
Urinary bladder (incl. in situ)**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	135	27.5	23.0-32.6	105	49.8	40.5-60.6	30	10.6	7.1-15.3	17	3.5	2.0-5.7	14	7.4	3.9-12.4	3	1.1	0.2-3.3
2009	123	25.2	20.9-30.1	86	41.4	32.9-51.5	37	13.7	9.6-19.0	17	3.6	2.1-5.8	8	4.3	1.8-8.4	9	3.2	1.4-6.2
2010	113	22.2	18.2-26.7	88	39.8	31.7-49.4	25	8.3	5.4-12.4	23	4.4	2.8-6.7	15	7.0	3.8-11.7	8	2.6	1.1-5.3
2011	143	27.0	22.7-31.9	110	50.1	40.9-60.8	33	10.8	7.4-15.4	22	4.4	2.7-6.7	18	9.2	5.3-14.5	4	1.3	0.3-3.6
2012	133	25.3	21.1-30.1	105	46.7	37.9-56.8	28	9.4	6.2-13.8	26	4.9	3.2-7.3	20	9.1	5.4-14.1	6	1.7	0.6-4.0
2013	137	25.3	21.1-30.0	101	43.7	35.3-53.4	36	11.9	8.3-16.8	27	4.8	3.2-7.1	21	9.4	5.7-14.5	6	1.8	0.6-4.2
2014	129	23.4	19.5-27.9	100	42.6	34.4-52.1	29	8.6	5.7-12.7	31	5.9	4.0-8.4	19	9.8	5.9-15.3	12	3.6	1.8-6.5
2015	171	30.0	25.6-35.0	142	57.3	48.0-67.9	29	9.5	6.4-13.9	30	5.4	3.6-7.8	23	9.8	6.1-14.9	7	2.2	0.9-4.7
2016	140	24.7	20.7-29.2	106	43.4	35.3-52.8	34	10.7	7.3-15.1	31	5.6	3.8-8.0	19	8.9	5.3-13.9	12	3.7	1.9-6.7
2017	153	25.4	21.5-29.9	111	43.0	35.1-52.1	42	12.1	8.7-16.7	30	5.3	3.6-7.7	25	11.3	7.2-16.7	5	1.3	0.4-3.4
2018	148	24.7	20.8-29.1	113	44.2	36.2-53.4	35	9.8	6.8-13.8	29	4.8	3.2-7.0	19	7.9	4.7-12.5	10	3.0	1.4-5.7
2019	153	24.9	21.0-29.3	107	40.0	32.6-48.6	46	13.3	9.7-17.9	24	4.0	2.5-6.0	18	7.4	4.4-11.9	6	1.6	0.6-3.7
2020	128	20.4	17.0-24.4	94	33.6	27.0-41.4	34	10.1	6.9-14.3	20	3.2	2.0-5.1	12	4.6	2.3-8.2	8	2.3	1.0-4.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Kidney and renal pelvis

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	30	11.4	7.6-16.3	15	14.2	7.5-23.9	15	9.9	5.5-16.5	13	4.8	2.5-8.2	10	9.3	4.1-17.5	3	1.9	0.4-5.5
1977	18	6.5	3.8-10.3	12	9.5	4.8-16.8	6	4.1	1.5-9.0	6	2.6	0.9-5.5	3	3.4	0.7-9.5	3	2.3	0.5-6.4
1978	29	10.1	6.7-14.5	15	11.8	6.4-19.7	14	8.7	4.7-14.7	13	4.8	2.5-8.2	6	5.5	1.9-12.0	7	4.3	1.7-8.9
1979	31	10.7	7.2-15.3	23	19.1	11.8-29.2	8	4.7	2.0-9.4	12	4.5	2.3-7.9	9	8.7	3.9-16.4	3	1.8	0.4-5.2
1980	17	5.2	3.0-8.4	13	9.3	4.8-16.1	4	2.1	0.6-5.4	9	2.6	1.2-5.0	4	2.3	0.6-6.3	5	2.6	0.8-6.2
1981	24	7.8	5.0-11.7	17	13.2	7.4-21.6	7	3.9	1.6-8.1	15	5.2	2.8-8.5	11	8.4	4.1-15.1	4	2.4	0.6-6.1
1982	24	8.4	5.3-12.5	10	7.9	3.6-14.8	14	8.3	4.5-13.9	13	4.3	2.2-7.4	6	6.2	2.1-13.2	7	3.7	1.5-7.7
1983	23	7.3	4.6-11.1	15	11.3	6.1-19.0	8	4.3	1.8-8.6	9	2.8	1.3-5.4	5	4.0	1.1-9.7	4	2.1	0.6-5.5
1984	30	9.0	6.0-12.9	19	13.6	8.0-21.5	11	5.7	2.8-10.3	9	3.1	1.4-5.9	6	4.9	1.7-10.5	3	1.8	0.4-5.1
1985	32	9.6	6.5-13.6	22	16.2	10.0-24.7	10	4.9	2.4-9.2	11	3.5	1.7-6.3	8	6.0	2.5-11.9	3	1.6	0.3-4.6
1986	26	7.6	4.9-11.2	17	11.0	6.3-17.9	9	4.9	2.2-9.3	13	3.9	2.0-6.6	6	3.5	1.3-7.9	7	3.7	1.5-7.6
1987	28	8.3	5.5-12.1	16	10.6	6.0-17.4	12	6.4	3.3-11.2	19	5.3	3.2-8.2	14	8.9	4.8-15.0	5	2.5	0.8-5.9
1988	22	6.9	4.3-10.4	10	9.6	4.3-17.8	12	6.2	3.2-10.9	11	3.2	1.6-5.8	7	5.2	2.1-10.7	4	1.9	0.5-5.1
1989	35	10.4	7.2-14.5	24	16.8	10.6-25.1	11	5.8	2.9-10.4	14	4.5	2.5-7.6	7	6.4	2.4-13.1	7	3.7	1.5-7.6
1990	42	11.9	8.6-16.1	27	17.2	11.2-25.2	15	7.7	4.3-12.7	11	3.2	1.6-5.7	6	4.4	1.6-9.5	5	2.5	0.8-6.0
1991	38	10.5	7.4-14.4	25	14.6	9.4-21.7	13	6.4	3.4-11.0	8	2.3	1.0-4.5	5	3.0	1.0-7.2	3	1.6	0.3-4.6
1992	34	9.1	6.3-12.8	20	11.3	6.9-17.6	14	6.8	3.7-11.4	10	2.9	1.4-5.4	6	4.1	1.3-9.2	4	1.9	0.5-5.0
1993	36	9.8	6.8-13.5	22	14.3	8.8-21.9	14	6.7	3.7-11.3	12	3.2	1.7-5.6	10	6.5	3.1-12.1	2	0.9	0.1-3.4
1994	52	13.9	10.4-18.2	30	19.9	13.2-28.6	22	10.3	6.5-15.7	14	3.7	2.0-6.2	7	4.7	1.8-9.9	7	3.1	1.2-6.4
1995	51	12.8	9.5-16.9	29	16.8	11.2-24.2	22	10.0	6.3-15.2	17	4.4	2.5-7.0	9	5.4	2.4-10.3	8	3.6	1.6-7.1
1996	46	12.1	8.9-16.2	27	17.9	11.6-26.2	19	8.4	5.1-13.2	15	4.0	2.2-6.5	7	4.5	1.8-9.3	8	3.4	1.5-6.8
1997	40	9.9	7.1-13.5	22	12.5	7.7-19.0	18	8.1	4.8-12.8	18	4.6	2.7-7.3	6	3.9	1.4-8.4	12	5.1	2.6-9.0
1998	52	12.4	9.3-16.3	25	13.8	8.8-20.4	27	11.7	7.7-17.0	19	4.7	2.8-7.4	8	5.2	2.2-10.1	11	4.7	2.3-8.4
1999	55	12.8	9.7-16.7	38	19.4	13.7-26.8	17	7.2	4.2-11.6	12	2.9	1.5-5.1	7	3.8	1.5-8.1	5	2.0	0.7-4.8
2000	60	13.7	10.4-17.6	36	18.8	13.1-26.3	24	9.9	6.3-14.8	16	3.9	2.2-6.3	11	7.0	3.4-12.4	5	2.0	0.6-4.7
2001	76	16.8	13.3-21.1	45	22.4	16.2-30.2	31	12.5	8.5-17.9	15	3.4	1.9-5.6	9	5.2	2.3-9.8	6	2.5	0.9-5.5
2002	82	17.9	14.2-22.2	53	25.1	18.7-32.9	29	11.3	7.5-16.3	16	3.7	2.1-6.0	9	5.0	2.2-9.6	7	2.8	1.1-5.8
2003	79	17.3	13.7-21.5	59	28.6	21.6-37.1	20	8.2	5.0-12.6	18	4.0	2.4-6.3	11	5.8	2.8-10.3	7	2.5	1.0-5.3
2004	77	16.7	13.2-20.9	54	26.0	19.4-34.0	23	9.1	5.7-13.7	15	3.3	1.8-5.4	9	4.2	1.9-8.0	6	2.3	0.8-5.1
2005	83	17.6	14.0-21.9	51	23.6	17.5-31.2	32	12.1	8.2-17.2	12	2.5	1.3-4.5	5	2.3	0.7-5.4	7	2.5	1.0-5.3
2006	82	17.3	13.7-21.5	53	24.3	18.1-32.0	29	11.3	7.5-16.3	24	5.1	3.3-7.7	16	7.6	4.3-12.5	8	3.0	1.3-6.0
2007	92	18.7	15.0-23.0	58	25.9	19.5-33.8	34	13.2	9.1-18.5	17	3.7	2.1-5.9	10	5.1	2.4-9.5	7	2.6	1.0-5.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Kidney and renal pelvis

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	100	19.2	15.6-23.5	65	27.5	21.1-35.3	35	11.9	8.3-16.8	13	2.7	1.4-4.7	9	4.5	2.0-8.6	4	1.4	0.4-3.7
2009	90	17.8	14.3-22.0	60	25.9	19.6-33.5	30	10.8	7.3-15.6	16	3.1	1.7-5.1	9	3.6	1.6-7.0	7	2.4	1.0-5.1
2010	90	17.1	13.7-21.2	54	21.7	16.1-28.6	36	13.0	9.0-18.2	16	3.1	1.8-5.1	12	5.5	2.7-9.7	4	1.5	0.4-4.0
2011	99	18.6	15.0-22.7	70	29.0	22.4-36.9	29	10.0	6.7-14.6	20	3.7	2.2-5.8	9	3.4	1.5-6.6	11	3.6	1.8-6.6
2012	99	18.4	14.9-22.6	59	23.5	17.7-30.6	40	13.7	9.7-18.8	16	3.2	1.8-5.2	9	4.2	1.9-8.1	7	2.4	0.9-5.1
2013	102	18.0	14.6-22.0	75	27.9	21.8-35.2	27	9.4	6.1-13.8	16	2.9	1.6-4.8	13	5.1	2.6-8.9	3	1.2	0.2-3.5
2014	110	20.0	16.3-24.2	68	26.3	20.3-33.7	42	14.6	10.5-20.0	19	3.7	2.2-5.8	8	3.7	1.6-7.4	11	3.6	1.8-6.6
2015	115	19.6	16.1-23.7	79	29.3	23.0-36.9	36	11.6	8.0-16.3	30	5.3	3.6-7.7	21	8.3	5.0-12.9	9	3.0	1.3-5.8
2016	118	20.6	17.0-24.9	71	26.9	20.8-34.2	47	15.3	11.1-20.7	16	2.7	1.5-4.5	13	5.1	2.7-8.9	3	0.9	0.2-3.0
2017	156	26.5	22.4-31.2	104	39.2	31.8-47.9	52	15.6	11.6-20.7	21	3.7	2.2-5.7	16	6.3	3.5-10.4	5	1.8	0.6-4.4
2018	122	20.8	17.2-25.0	90	33.1	26.4-41.0	32	10.2	6.9-14.6	23	3.8	2.4-5.9	15	5.8	3.2-9.8	8	2.1	0.9-4.5
2019	128	21.5	17.8-25.8	92	34.0	27.2-42.1	36	11.0	7.6-15.5	16	2.6	1.4-4.3	14	5.4	2.9-9.2	2	0.5	0.1-2.2
2020	106	16.7	13.6-20.4	70	23.6	18.2-30.2	36	10.9	7.5-15.4	18	3.0	1.7-4.8	8	3.5	1.5-6.9	10	2.7	1.3-5.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Brain and other central nervous system

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	26	8.9	5.7-13.2	17	12.2	6.9-20.0	9	6.0	2.7-11.4	21	7.2	4.4-11.0	11	8.5	4.1-15.5	10	6.2	3.0-11.5
1977	23	7.2	4.5-11.0	11	6.9	3.4-12.8	12	7.2	3.7-12.7	8	2.3	1.0-4.7	5	2.9	0.9-7.3	3	1.9	0.4-5.6
1978	14	4.3	2.3-7.3	8	5.1	2.1-10.4	6	3.7	1.3-8.1	13	4.3	2.3-7.5	8	5.8	2.5-11.7	5	2.9	0.9-6.9
1979	20	6.2	3.8-9.7	14	9.6	5.1-16.5	6	3.4	1.2-7.6	11	3.9	1.9-6.9	8	6.1	2.5-12.3	3	2.2	0.4-6.1
1980	13	4.1	2.1-7.0	8	6.1	2.5-12.4	5	2.8	0.9-6.6	13	3.8	2.0-6.6	10	7.0	3.2-13.2	3	1.5	0.3-4.7
1981	26	7.3	4.7-10.8	11	6.2	3.1-11.4	15	8.1	4.5-13.5	13	3.7	1.9-6.4	6	3.8	1.4-8.5	7	3.5	1.4-7.3
1982	19	6.1	3.6-9.5	10	7.6	3.4-14.3	9	5.1	2.3-9.7	10	3.2	1.5-6.0	4	3.4	0.8-8.6	6	3.2	1.2-7.0
1983	23	6.9	4.3-10.5	14	9.0	4.7-15.6	9	5.2	2.3-9.9	15	4.7	2.6-7.8	9	7.6	3.2-14.8	6	3.1	1.1-6.7
1984	18	5.0	2.9-8.0	9	5.9	2.6-11.4	9	4.5	2.0-8.7	17	5.2	3.0-8.4	7	5.3	2.0-11.2	10	5.5	2.6-10.2
1985	26	7.5	4.9-11.1	14	9.5	5.1-16.2	12	6.5	3.3-11.4	19	5.4	3.2-8.5	9	6.0	2.6-11.6	10	5.3	2.5-9.8
1986	25	7.3	4.7-10.9	16	10.9	6.0-18.1	9	4.6	2.1-8.8	16	4.8	2.7-7.8	11	7.8	3.7-14.4	5	2.4	0.8-5.7
1987	28	7.9	5.2-11.5	11	6.8	3.3-12.3	17	8.4	4.9-13.6	17	5.2	3.0-8.4	8	6.2	2.6-12.3	9	4.9	2.2-9.3
1988	26	7.8	5.0-11.4	15	9.9	5.4-16.6	11	6.0	3.0-10.8	17	5.2	3.0-8.3	8	5.2	2.2-10.3	9	4.8	2.2-9.2
1989	26	7.2	4.7-10.6	16	9.6	5.4-15.9	10	5.1	2.4-9.4	15	4.3	2.4-7.2	9	6.2	2.8-11.9	6	3.0	1.1-6.6
1990	26	7.1	4.6-10.5	14	8.2	4.5-13.9	12	6.0	3.1-10.6	9	2.7	1.2-5.1	3	2.5	0.4-7.5	6	3.2	1.2-7.0
1991	28	7.4	4.9-10.8	18	10.1	5.9-16.1	10	4.8	2.3-8.9	14	3.9	2.1-6.6	10	6.6	3.1-12.2	4	2.0	0.5-5.2
1992	25	7.0	4.5-10.3	16	11.0	6.0-18.2	9	4.4	2.0-8.3	15	4.4	2.4-7.2	7	4.8	1.7-10.1	8	4.1	1.8-8.1
1993	26	6.9	4.5-10.2	17	10.1	5.8-16.3	9	4.2	1.9-8.0	13	3.7	1.9-6.3	9	6.2	2.7-11.9	4	1.8	0.5-4.8
1994	24	6.2	4.0-9.3	12	6.7	3.4-11.8	12	5.7	2.9-9.9	12	3.1	1.6-5.4	6	3.9	1.4-8.4	6	2.8	1.0-6.2
1995	42	11.0	7.9-14.8	22	13.8	8.5-21.2	20	9.3	5.7-14.4	24	6.3	4.0-9.4	14	8.4	4.5-14.2	10	4.6	2.2-8.4
1996	31	7.7	5.3-11.0	20	11.4	6.8-17.9	11	5.1	2.5-9.2	17	4.5	2.6-7.1	9	6.5	2.8-12.3	8	3.6	1.6-7.2
1997	40	9.9	7.1-13.6	17	9.2	5.3-14.9	23	10.4	6.6-15.6	24	6.1	3.9-9.1	7	3.9	1.5-8.0	17	7.3	4.3-11.8
1998	28	6.6	4.4-9.6	12	6.3	3.2-11.1	16	7.2	4.1-11.7	17	4.0	2.3-6.4	6	2.9	1.1-6.4	11	4.8	2.4-8.6
1999	29	6.6	4.4-9.5	13	6.3	3.3-10.9	16	6.9	3.9-11.2	11	2.6	1.3-4.6	5	2.4	0.8-5.6	6	2.6	1.0-5.7
2000	32	7.5	5.1-10.6	13	7.5	3.9-12.8	19	8.0	4.8-12.6	9	2.1	0.9-3.9	6	3.3	1.2-7.2	3	1.2	0.2-3.6
2001	35	7.9	5.5-11.0	22	10.8	6.7-16.5	13	5.4	2.9-9.3	18	4.1	2.5-6.6	11	5.7	2.8-10.2	7	2.8	1.1-5.8
2002	27	5.9	3.9-8.5	16	7.4	4.2-12.1	11	4.4	2.2-7.9	18	3.9	2.3-6.2	11	5.1	2.5-9.2	7	2.8	1.1-5.8
2003	27	6.0	3.9-8.7	12	5.7	2.9-10.0	15	6.0	3.3-10.0	12	2.6	1.3-4.5	6	2.8	1.0-6.3	6	2.4	0.9-5.2
2004	24	5.2	3.3-7.8	15	7.0	3.9-11.7	9	3.7	1.7-7.0	15	3.3	1.8-5.4	5	2.5	0.8-5.9	10	3.9	1.9-7.3
2005	36	7.6	5.3-10.6	20	9.5	5.7-14.9	16	6.3	3.6-10.3	17	3.6	2.1-5.8	9	4.5	2.0-8.7	8	3.1	1.3-6.3
2006	44	9.6	7.0-12.9	21	9.5	5.8-14.6	23	9.6	6.0-14.4	23	4.9	3.1-7.3	15	6.8	3.8-11.3	8	3.0	1.3-6.0
2007	36	7.6	5.3-10.6	18	8.7	5.1-13.8	18	6.9	4.0-11.0	22	4.4	2.8-6.8	13	5.9	3.1-10.2	9	3.1	1.4-6.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Brain and other central nervous system

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	29	5.9	3.9-8.6	12	5.3	2.7-9.4	17	6.7	3.9-10.9	13	2.8	1.5-4.8	4	2.2	0.6-5.6	9	3.3	1.5-6.3
2009	30	6.0	4.0-8.6	17	6.9	3.9-11.2	13	5.2	2.7-9.0	13	2.8	1.5-4.9	6	2.9	1.0-6.3	7	2.9	1.2-6.1
2010	33	6.7	4.6-9.4	18	7.4	4.3-11.8	15	6.2	3.4-10.3	19	3.5	2.1-5.5	11	4.0	2.0-7.5	8	2.8	1.2-5.7
2011	30	6.5	4.3-9.2	16	7.7	4.4-12.6	14	5.4	2.9-9.2	21	4.1	2.5-6.3	9	4.4	1.9-8.3	12	4.3	2.2-7.7
2012	37	7.6	5.3-10.5	17	7.5	4.4-12.1	20	7.5	4.5-11.7	14	2.8	1.5-4.8	8	3.5	1.5-7.1	6	2.2	0.8-4.8
2013	34	6.7	4.6-9.5	19	8.4	5.0-13.2	15	5.4	3.0-9.1	21	4.3	2.6-6.6	12	5.2	2.6-9.2	9	3.2	1.4-6.3
2014	40	7.7	5.5-10.6	24	10.2	6.5-15.3	16	5.5	3.0-9.1	33	6.2	4.2-8.8	20	8.6	5.2-13.5	13	4.1	2.2-7.3
2015	35	6.4	4.4-9.0	15	5.3	2.9-9.1	20	7.3	4.4-11.4	17	2.8	1.6-4.6	8	2.9	1.2-6.0	9	2.7	1.2-5.5
2016	39	6.8	4.8-9.5	18	7.0	4.1-11.3	21	6.7	4.1-10.6	20	3.5	2.1-5.5	10	3.9	1.8-7.4	10	3.2	1.5-6.1
2017	46	8.7	6.3-11.7	28	11.8	7.8-17.2	18	5.7	3.3-9.3	27	4.7	3.1-6.9	14	5.1	2.7-8.8	13	4.4	2.3-7.7
2018	30	5.3	3.5-7.7	15	5.2	2.9-8.9	15	5.3	2.9-8.9	19	3.2	1.9-5.2	14	5.2	2.8-9.0	5	1.4	0.5-3.6
2019	36	7.1	4.9-10.0	18	7.8	4.5-12.5	18	6.6	3.8-10.7	23	3.9	2.4-6.0	11	4.2	2.0-7.8	12	3.6	1.8-6.6
2020	30	5.5	3.6-7.9	16	5.9	3.3-9.9	14	5.1	2.7-8.7	19	3.1	1.8-4.9	8	3.0	1.3-6.2	11	3.1	1.5-5.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Thyroid

Year of diagnosis	Incidence								
	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	10	3.4	1.6-6.3	2	1.3	0.2-5.1	8	5.1	2.2-10.1
1977	16	5.1	2.8-8.4	4	2.6	0.7-7.1	12	6.8	3.5-12.1
1978	10	3.8	1.8-7.0	0	0.0	0.0-3.0	10	6.8	3.2-12.5
1979	13	4.7	2.4-8.0	5	4.1	1.3-9.6	8	5.2	2.2-10.2
1980	17	5.0	2.9-8.1	3	2.1	0.4-6.3	14	7.3	3.9-12.4
1981	9	2.6	1.2-5.1	3	1.6	0.3-5.3	6	3.6	1.3-7.9
1982	10	3.2	1.5-5.8	4	3.0	0.8-7.9	6	3.5	1.2-7.6
1983	9	2.3	1.0-4.5	4	2.2	0.6-6.0	5	2.4	0.8-5.8
1984	13	4.0	2.1-7.0	5	3.0	1.0-7.3	8	4.7	2.0-9.3
1985	17	4.8	2.7-7.8	6	3.4	1.2-7.8	11	5.7	2.8-10.3
1986	17	5.1	2.9-8.3	2	1.1	0.1-4.4	15	8.0	4.4-13.2
1987	15	4.2	2.3-7.0	7	4.5	1.8-9.4	8	4.0	1.7-8.0
1988	13	3.6	1.9-6.3	2	0.8	0.1-3.8	11	6.0	3.0-10.7
1989	16	4.4	2.5-7.2	5	3.1	1.0-7.4	11	5.5	2.7-9.9
1990	20	5.0	3.1-7.9	8	4.3	1.8-8.8	12	5.7	2.9-10.0
1991	17	4.2	2.4-6.8	5	2.4	0.8-6.0	12	5.7	2.9-10.0
1992	31	7.6	5.1-10.8	5	2.7	0.9-6.4	26	12.0	7.8-17.7
1993	26	6.7	4.4-9.8	5	3.3	1.0-7.7	21	10.0	6.1-15.3
1994	30	7.5	5.1-10.7	6	3.2	1.1-7.0	24	11.3	7.2-16.8
1995	29	7.4	4.9-10.6	13	7.2	3.8-12.5	16	7.2	4.1-11.7
1996	28	6.9	4.6-10.0	6	3.1	1.1-6.9	22	10.4	6.5-15.8
1997	31	7.3	4.9-10.3	15	7.6	4.2-12.7	16	7.2	4.1-11.7
1998	20	4.7	2.8-7.2	8	3.9	1.7-7.7	12	5.3	2.7-9.2
1999	32	7.2	4.9-10.1	7	3.4	1.4-7.1	25	10.6	6.9-15.7
2000	32	7.0	4.8-9.9	11	5.1	2.5-9.2	21	8.9	5.5-13.6
2001	40	8.5	6.1-11.6	10	4.4	2.1-8.2	30	12.4	8.3-17.7
2002	58	12.4	9.4-16.1	14	6.4	3.5-10.8	44	17.9	13.0-24.1
2003	54	11.4	8.5-14.9	13	6.0	3.2-10.4	41	16.1	11.5-21.9
2004	75	15.7	12.3-19.7	20	8.5	5.2-13.3	55	22.0	16.6-28.7
2005	97	20.5	16.6-25.0	25	11.1	7.2-16.5	72	29.1	22.8-36.8
2006	91	18.6	14.9-22.9	22	9.2	5.7-14.1	69	27.2	21.1-34.5
2007	111	22.8	18.7-27.5	27	12.4	8.1-18.1	84	32.7	26.0-40.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

Mortality data not provided due to small number of deaths per year.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
Richmond (Staten Island)
Thyroid

Year of diagnosis	Incidence								
	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
2008	139	28.2	23.6-33.4	35	14.6	10.1-20.4	104	40.8	33.2-49.5
2009	179	35.5	30.4-41.2	44	17.2	12.4-23.3	135	51.8	43.3-61.6
2010	171	33.9	28.9-39.5	42	17.0	12.1-23.1	129	49.4	41.1-58.9
2011	141	27.6	23.2-32.7	38	15.2	10.6-21.1	103	39.3	31.9-47.9
2012	158	31.2	26.4-36.6	31	13.4	9.0-19.2	127	47.6	39.5-56.9
2013	202	39.5	34.1-45.5	60	24.6	18.6-31.9	142	53.3	44.7-63.1
2014	175	34.6	29.5-40.3	46	18.2	13.2-24.5	129	50.1	41.6-59.8
2015	172	33.4	28.4-38.9	50	20.2	14.9-26.9	122	45.4	37.5-54.6
2016	192	37.2	32.0-43.1	49	19.0	13.9-25.3	143	54.0	45.2-64.0
2017	164	31.1	26.4-36.4	51	19.4	14.3-25.8	113	42.0	34.4-50.9
2018	163	32.1	27.2-37.7	37	14.7	10.2-20.5	126	48.5	40.1-58.2
2019	177	34.6	29.5-40.3	63	25.8	19.6-33.3	114	42.9	35.1-51.9
2020	126	24.1	20.0-29.0	43	17.2	12.3-23.4	83	30.7	24.2-38.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

Mortality data not provided due to small number of deaths per year.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Hodgkin lymphoma

Year of diagnosis	Incidence								
	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	10	3.2	1.5-6.0	8	5.1	2.2-10.5	2	1.5	0.2-5.3
1977	12	3.9	1.9-6.9	8	6.1	2.3-12.8	4	2.4	0.6-6.2
1978	16	4.4	2.5-7.3	10	6.1	2.9-11.6	6	3.0	1.1-6.8
1979	8	2.4	1.0-4.9	4	3.0	0.8-7.9	4	1.8	0.5-5.0
1980	22	6.0	3.7-9.3	14	9.3	4.7-16.2	8	3.6	1.6-7.4
1981	19	5.2	3.1-8.2	11	6.5	3.1-12.2	8	4.1	1.7-8.2
1982	20	5.5	3.3-8.5	10	5.8	2.7-11.0	10	5.2	2.4-9.7
1983	11	3.3	1.6-6.0	7	4.0	1.6-8.7	4	2.5	0.7-6.3
1984	21	5.9	3.6-9.1	14	7.7	4.1-13.3	7	4.1	1.6-8.5
1985	15	3.9	2.2-6.6	9	4.8	2.1-9.6	6	3.1	1.1-6.8
1986	16	4.3	2.4-7.0	6	3.4	1.2-7.7	10	4.9	2.3-9.2
1987	16	4.1	2.3-6.7	8	4.5	1.9-9.2	8	3.5	1.5-7.1
1988	25	6.2	4.0-9.3	11	5.8	2.8-10.7	14	6.5	3.5-11.1
1989	23	5.8	3.6-8.8	10	5.1	2.4-9.7	13	6.3	3.3-10.9
1990	16	4.3	2.4-7.0	10	5.5	2.6-10.3	6	3.0	1.1-6.7
1991	19	4.8	2.9-7.6	7	4.4	1.5-9.6	12	5.8	3.0-10.2
1992	17	4.0	2.3-6.4	8	3.8	1.6-7.8	9	4.1	1.9-7.9
1993	17	3.9	2.2-6.2	9	4.2	1.9-8.3	8	3.5	1.5-7.1
1994	15	3.9	2.2-6.4	10	5.7	2.7-10.7	5	2.4	0.8-5.6
1995	13	3.3	1.8-5.6	6	3.2	1.2-7.1	7	3.4	1.3-7.0
1996	12	2.9	1.5-5.1	5	2.4	0.8-5.9	7	3.2	1.3-6.7
1997	23	5.3	3.4-8.0	17	8.1	4.7-13.1	6	2.8	1.0-6.0
1998	12	2.8	1.5-4.9	7	3.4	1.4-7.1	5	2.3	0.7-5.3
1999	17	3.9	2.2-6.2	6	2.9	1.0-6.3	11	4.6	2.3-8.3
2000	21	4.8	2.9-7.3	8	3.8	1.6-7.5	13	5.6	3.0-9.5
2001	9	2.0	0.9-3.9	4	2.0	0.5-5.1	5	2.1	0.7-5.0
2002	20	4.3	2.6-6.6	11	5.2	2.6-9.4	9	3.6	1.6-6.9
2003	17	3.7	2.2-6.0	5	2.5	0.8-5.8	12	5.2	2.7-9.1
2004	16	3.6	2.0-5.8	10	4.8	2.3-9.0	6	2.7	1.0-5.8
2005	16	3.4	2.0-5.6	12	5.2	2.7-9.2	4	1.8	0.5-4.5
2006	20	4.2	2.6-6.6	14	6.0	3.3-10.1	6	2.6	0.9-5.7
2007	14	3.2	1.7-5.3	8	3.9	1.7-7.6	6	2.5	0.9-5.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

Mortality data not provided due to small number of deaths per year.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Hodgkin lymphoma

Year of diagnosis	Incidence								
	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
2008	14	3.1	1.7-5.2	8	3.4	1.5-6.9	6	2.8	1.0-6.0
2009	18	3.7	2.2-5.9	10	4.1	1.9-7.6	8	3.4	1.4-6.6
2010	19	4.1	2.5-6.4	8	3.5	1.5-7.0	11	4.6	2.3-8.2
2011	19	3.9	2.3-6.1	10	4.1	1.9-7.6	9	3.8	1.7-7.2
2012	22	4.7	2.9-7.1	12	5.4	2.8-9.5	10	4.0	1.9-7.4
2013	17	3.5	2.0-5.7	6	2.6	0.9-5.7	11	4.3	2.1-7.9
2014	16	3.4	1.9-5.5	9	3.6	1.6-6.9	7	3.2	1.3-6.6
2015	21	4.3	2.6-6.6	13	5.5	2.9-9.5	8	3.3	1.4-6.6
2016	26	5.6	3.6-8.3	16	7.0	3.9-11.4	10	4.2	2.0-7.9
2017	13	2.9	1.5-5.0	12	5.3	2.7-9.3	1	0.5	0.0-2.7
2018	16	2.8	1.6-4.7	8	2.9	1.2-5.9	8	2.8	1.1-5.7
2019	10	2.0	0.9-3.8	6	2.8	1.0-6.0	4	1.4	0.3-3.7
2020	15	3.2	1.7-5.3	4	1.8	0.5-4.6	11	4.4	2.1-8.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

Mortality data not provided due to small number of deaths per year.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

**Richmond (Staten Island)
Non-Hodgkin lymphomas**

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	30	10.6	7.1-15.2	20	16.2	9.5-25.5	10	6.4	3.0-11.8	16	6.0	3.3-9.8	11	9.5	4.4-17.5	5	3.4	1.1-7.9
1977	27	10.2	6.7-14.9	15	13.2	7.1-22.2	12	8.1	4.2-14.1	13	4.8	2.5-8.3	9	7.7	3.3-14.9	4	2.8	0.8-7.1
1978	30	9.9	6.6-14.2	18	14.2	8.2-22.8	12	7.0	3.6-12.3	17	6.4	3.7-10.3	11	12.0	5.7-21.4	6	3.4	1.2-7.6
1979	29	9.6	6.3-13.8	16	11.3	6.2-18.9	13	7.7	4.1-13.2	14	4.7	2.5-7.9	8	5.7	2.4-11.4	6	3.7	1.3-8.0
1980	42	13.8	9.9-18.8	24	16.1	10.2-24.3	18	10.5	6.2-16.6	16	5.5	3.1-9.0	9	6.5	2.8-12.7	7	4.4	1.7-8.9
1981	38	10.9	7.7-15.1	22	13.8	8.4-21.4	16	8.4	4.8-13.8	22	6.6	4.1-10.1	10	6.2	2.9-11.7	12	6.5	3.3-11.4
1982	40	13.0	9.2-17.7	15	11.5	6.2-19.4	25	14.4	9.3-21.3	24	7.7	4.9-11.5	10	7.1	3.3-13.3	14	7.9	4.3-13.3
1983	47	14.9	10.9-19.9	24	15.7	10.0-23.7	23	12.9	8.1-19.3	28	9.0	5.9-13.0	13	9.4	4.8-16.5	15	8.2	4.6-13.6
1984	48	14.7	10.8-19.6	24	15.8	10.0-23.9	24	13.6	8.6-20.3	15	4.8	2.6-7.9	6	4.0	1.4-8.8	9	4.8	2.2-9.2
1985	49	15.3	11.2-20.2	25	17.0	10.9-25.4	24	13.3	8.5-19.9	31	9.8	6.6-14.0	15	11.5	6.1-19.5	16	8.8	5.0-14.4
1986	62	18.9	14.4-24.3	24	15.2	9.6-23.0	38	20.4	14.4-28.1	20	6.1	3.7-9.4	12	6.9	3.5-12.3	8	4.4	1.9-8.7
1987	64	18.7	14.4-24.0	32	21.3	14.4-30.3	32	16.6	11.3-23.5	19	5.6	3.3-8.8	6	4.2	1.5-9.3	13	6.4	3.4-11.0
1988	57	16.9	12.7-21.9	29	21.3	13.8-31.3	28	14.9	9.8-21.5	24	7.2	4.5-10.7	16	11.2	6.0-18.8	8	4.1	1.8-8.2
1989	59	17.9	13.5-23.1	38	27.6	18.9-38.6	21	10.6	6.5-16.2	22	6.8	4.2-10.4	13	9.9	4.9-17.5	9	5.0	2.3-9.4
1990	57	16.5	12.4-21.4	29	20.2	13.2-29.4	28	14.1	9.3-20.4	23	6.8	4.3-10.2	14	9.8	5.1-16.8	9	4.5	2.1-8.6
1991	87	24.3	19.4-30.0	53	32.8	24.3-43.4	34	17.5	12.1-24.5	25	7.1	4.6-10.5	16	10.6	6.0-17.4	9	4.3	2.0-8.2
1992	62	16.8	12.9-21.6	34	21.0	14.3-29.7	28	13.6	9.0-19.7	25	6.7	4.3-10.0	14	9.3	4.9-15.9	11	5.4	2.7-9.7
1993	76	19.8	15.5-24.8	40	22.6	15.9-31.1	36	16.8	11.8-23.3	30	8.0	5.4-11.5	17	10.1	5.8-16.3	13	6.3	3.3-10.7
1994	78	20.7	16.3-25.9	34	20.7	14.1-29.4	44	20.7	15.0-27.9	33	8.8	6.1-12.4	18	11.8	6.7-18.9	15	6.9	3.9-11.5
1995	86	22.5	17.9-27.8	39	23.2	16.2-32.0	47	21.7	15.9-28.8	33	8.5	5.9-12.0	17	9.2	5.3-14.9	16	7.3	4.2-12.0
1996	77	19.4	15.3-24.2	35	18.5	12.8-25.9	42	19.0	13.7-25.7	38	10.0	7.1-13.7	19	11.9	7.0-18.9	19	8.5	5.1-13.2
1997	101	25.2	20.5-30.6	58	33.0	24.8-42.9	43	18.9	13.7-25.5	29	7.6	5.1-10.9	12	7.6	3.8-13.4	17	7.4	4.3-11.9
1998	105	25.3	20.6-30.6	55	30.7	23.0-40.2	50	21.4	15.9-28.3	29	7.2	4.8-10.3	14	7.9	4.2-13.4	15	6.1	3.4-10.2
1999	114	26.9	22.2-32.3	57	32.2	24.1-42.0	57	24.2	18.3-31.4	39	9.5	6.7-12.9	25	15.0	9.5-22.4	14	5.7	3.1-9.6
2000	77	17.5	13.8-21.8	47	23.5	17.2-31.4	30	12.1	8.1-17.3	33	7.7	5.3-10.8	21	11.0	6.7-16.9	12	5.0	2.6-8.7
2001	111	24.9	20.5-30.0	63	31.6	24.1-40.6	48	19.3	14.2-25.7	32	7.2	4.9-10.2	18	9.8	5.7-15.6	14	5.4	2.9-9.1
2002	105	23.3	19.0-28.2	61	30.7	23.4-39.6	44	17.2	12.5-23.2	28	6.3	4.2-9.2	13	7.0	3.7-12.1	15	6.0	3.3-9.9
2003	102	22.1	18.0-26.9	57	28.8	21.6-37.5	45	17.3	12.6-23.2	30	6.8	4.6-9.7	14	7.1	3.8-12.1	16	6.2	3.5-10.1
2004	106	22.5	18.4-27.3	54	25.8	19.2-33.8	52	20.1	15.0-26.4	36	7.9	5.6-11.0	20	10.9	6.5-16.9	16	6.2	3.5-10.1
2005	118	25.2	20.8-30.2	53	25.9	19.2-34.2	65	25.5	19.7-32.6	40	8.3	5.9-11.4	17	7.8	4.4-12.7	23	8.6	5.4-13.0
2006	117	24.8	20.5-29.8	72	33.5	26.1-42.4	45	17.2	12.5-23.2	35	7.6	5.3-10.6	19	9.5	5.6-14.9	16	5.6	3.2-9.2
2007	89	18.5	14.8-22.8	40	18.5	13.1-25.4	49	18.6	13.7-24.7	31	6.9	4.7-9.8	16	8.1	4.5-13.2	15	6.0	3.3-9.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Non-Hodgkin lymphomas

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	121	24.5	20.3-29.4	63	29.7	22.6-38.2	58	21.4	16.2-27.8	22	4.6	2.9-7.0	12	5.9	3.0-10.5	10	4.0	1.9-7.4
2009	113	23.2	19.1-28.0	59	28.1	21.2-36.6	54	19.8	14.8-26.0	29	5.9	3.9-8.5	18	8.4	4.8-13.4	11	4.0	2.0-7.2
2010	128	26.1	21.7-31.1	63	29.3	22.4-37.8	65	23.9	18.4-30.6	30	6.2	4.2-8.9	17	8.0	4.5-13.0	13	5.1	2.7-8.7
2011	116	22.4	18.4-26.9	62	26.5	20.1-34.3	54	19.0	14.2-25.0	24	4.8	3.1-7.3	13	6.5	3.4-11.1	11	3.8	1.9-7.0
2012	109	20.4	16.7-24.7	50	21.3	15.7-28.4	59	20.2	15.3-26.2	35	6.6	4.6-9.2	20	9.0	5.4-14.0	15	5.2	2.9-8.7
2013	122	22.4	18.6-26.9	70	28.9	22.4-36.8	52	17.3	12.8-22.9	23	4.3	2.7-6.6	14	6.3	3.4-10.7	9	3.0	1.4-6.0
2014	143	27.1	22.8-32.1	79	34.8	27.4-43.6	64	21.2	16.2-27.4	21	4.0	2.4-6.1	14	6.8	3.7-11.3	7	2.1	0.8-4.5
2015	153	28.1	23.7-33.1	80	32.9	25.9-41.3	73	24.1	18.7-30.6	23	4.3	2.7-6.5	15	6.6	3.6-11.1	8	2.5	1.1-5.2
2016	125	22.2	18.3-26.6	60	23.7	17.9-30.8	65	21.2	16.2-27.3	22	4.0	2.5-6.1	10	4.2	2.0-7.8	12	3.6	1.8-6.5
2017	141	24.9	20.8-29.5	78	30.2	23.7-38.0	63	20.4	15.5-26.3	26	4.6	3.0-6.8	18	7.3	4.2-11.6	8	2.5	1.1-5.2
2018	148	25.8	21.7-30.4	83	32.4	25.6-40.5	65	20.1	15.4-25.9	25	4.2	2.7-6.4	13	5.3	2.8-9.2	12	3.5	1.8-6.4
2019	144	24.6	20.6-29.1	72	28.0	21.8-35.5	72	21.7	16.9-27.7	23	4.1	2.6-6.3	15	6.3	3.5-10.5	8	2.3	1.0-4.9
2020	134	22.1	18.4-26.3	68	25.0	19.2-32.0	66	19.6	15.0-25.2	33	5.6	3.8-8.0	17	7.0	4.0-11.3	16	4.5	2.5-7.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Myeloma

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	14	5.2	2.8-8.8	6	5.5	1.8-12.4	8	5.2	2.2-10.2	9	3.8	1.7-7.1	5	5.6	1.7-13.0	4	2.7	0.7-6.9
1977	6	2.4	0.9-5.2	1	0.6	0.0-4.1	5	3.3	1.1-7.7	7	2.4	1.0-5.0	2	1.3	0.2-5.2	5	3.0	1.0-7.1
1978	11	3.9	1.9-7.1	7	6.1	2.3-12.8	4	2.4	0.7-6.3	3	0.9	0.2-2.6	2	1.2	0.1-4.9	1	0.5	0.0-3.2
1979	8	2.8	1.2-5.6	3	2.2	0.5-6.6	5	3.1	1.0-7.1	5	1.8	0.6-4.2	1	0.6	0.0-4.0	4	2.5	0.7-6.3
1980	17	6.1	3.5-9.8	7	6.9	2.6-14.2	10	6.1	2.9-11.1	10	3.4	1.6-6.3	3	2.0	0.4-6.1	7	4.2	1.7-8.7
1981	10	3.6	1.7-6.6	5	5.4	1.6-12.2	5	2.8	0.9-6.6	12	4.5	2.3-7.8	7	8.2	3.1-16.4	5	2.8	0.9-6.6
1982	11	4.0	2.0-7.1	5	5.1	1.6-11.4	6	3.6	1.3-7.9	7	2.4	0.9-4.9	4	3.1	0.8-8.0	3	1.7	0.3-5.0
1983	14	4.2	2.3-7.1	11	7.2	3.6-13.0	3	1.6	0.3-4.7	10	3.1	1.5-5.7	7	5.4	2.1-11.2	3	1.6	0.3-4.8
1984	14	4.8	2.6-8.1	9	8.7	3.8-16.5	5	2.7	0.9-6.3	13	4.4	2.3-7.5	3	2.1	0.4-6.2	10	5.5	2.6-10.1
1985	20	6.5	3.9-10.0	10	7.3	3.4-13.5	10	5.6	2.7-10.2	13	4.0	2.1-6.9	7	4.7	1.8-9.9	6	3.5	1.3-7.6
1986	16	4.8	2.7-7.8	5	2.9	0.9-7.1	11	5.9	2.9-10.6	12	3.8	1.9-6.6	5	4.3	1.2-10.2	7	3.7	1.5-7.7
1987	16	5.0	2.8-8.1	8	5.8	2.3-11.7	8	4.3	1.8-8.5	11	3.6	1.8-6.4	5	4.0	1.1-9.6	6	3.3	1.2-7.1
1988	16	4.6	2.6-7.5	7	4.7	1.8-9.7	9	4.5	2.0-8.5	10	2.8	1.4-5.3	7	4.6	1.8-9.7	3	1.4	0.3-4.3
1989	24	7.2	4.6-10.7	10	6.6	3.1-12.3	14	7.6	4.1-12.8	9	2.9	1.3-5.4	2	1.7	0.2-5.7	7	3.7	1.5-7.7
1990	17	4.8	2.8-7.6	9	5.8	2.6-11.2	8	4.1	1.7-8.1	8	2.3	1.0-4.5	4	2.4	0.6-6.2	4	2.0	0.5-5.2
1991	19	5.3	3.2-8.3	8	5.9	2.5-11.7	11	5.3	2.6-9.6	12	3.7	1.9-6.4	6	5.0	1.8-10.7	6	3.2	1.2-6.9
1992	12	3.5	1.8-6.1	8	6.3	2.5-12.7	4	2.0	0.5-5.1	14	3.9	2.1-6.5	6	4.0	1.4-8.6	8	3.7	1.6-7.3
1993	16	4.5	2.6-7.3	5	4.1	1.2-9.4	11	5.2	2.6-9.3	8	2.4	1.0-4.7	3	2.8	0.5-7.8	5	2.3	0.8-5.5
1994	13	3.5	1.9-6.0	5	3.4	1.1-7.9	8	3.8	1.6-7.5	12	3.1	1.6-5.5	4	2.5	0.7-6.4	8	3.7	1.6-7.4
1995	15	3.9	2.2-6.5	9	5.0	2.3-9.6	6	2.7	1.0-5.9	10	2.7	1.3-5.0	4	2.5	0.7-6.4	6	2.8	1.0-6.0
1996	30	7.6	5.1-10.8	19	11.1	6.6-17.4	11	5.1	2.5-9.1	8	2.0	0.9-3.9	4	2.5	0.6-6.3	4	1.8	0.5-4.6
1997	23	5.8	3.7-8.7	10	6.4	3.0-11.8	13	5.6	3.0-9.6	16	4.0	2.3-6.5	10	6.3	3.0-11.5	6	2.6	1.0-5.8
1998	33	8.1	5.6-11.4	14	9.2	4.8-15.5	19	8.0	4.8-12.6	11	2.8	1.4-4.9	5	3.3	1.0-7.7	6	2.5	0.9-5.6
1999	24	5.7	3.6-8.5	10	5.6	2.6-10.5	14	5.9	3.2-9.9	8	2.0	0.8-3.9	1	1.0	0.0-4.3	7	2.9	1.1-6.0
2000	20	4.7	2.8-7.2	14	7.4	4.0-12.4	6	2.4	0.9-5.2	13	3.1	1.7-5.3	4	2.4	0.6-6.1	9	3.5	1.6-6.8
2001	37	8.6	6.0-11.8	20	11.2	6.7-17.3	17	6.8	3.9-10.9	9	2.1	0.9-3.9	6	3.3	1.1-7.3	3	1.2	0.2-3.6
2002	37	8.3	5.8-11.4	17	9.5	5.5-15.2	20	7.5	4.6-11.7	16	3.7	2.1-5.9	5	2.8	0.9-6.5	11	4.2	2.1-7.6
2003	27	6.0	3.9-8.7	14	7.7	4.1-13.1	13	5.0	2.7-8.6	16	3.6	2.1-5.9	6	3.0	1.1-6.7	10	3.8	1.8-7.1
2004	26	5.7	3.7-8.3	14	7.9	4.2-13.3	12	4.6	2.4-8.1	19	4.2	2.5-6.5	10	5.8	2.7-10.6	9	3.1	1.4-6.1
2005	35	7.2	5.0-10.1	18	8.5	5.0-13.6	17	6.0	3.5-9.8	8	1.6	0.7-3.3	4	2.3	0.6-5.8	4	1.3	0.4-3.6
2006	37	8.0	5.6-11.0	14	7.2	3.9-12.2	23	8.8	5.5-13.3	11	2.2	1.1-4.0	7	3.9	1.5-8.1	4	1.4	0.4-3.7
2007	46	9.5	7.0-12.8	25	13.2	8.5-19.6	21	7.2	4.4-11.2	11	2.3	1.2-4.2	6	3.4	1.2-7.4	5	1.9	0.6-4.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Myeloma

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	33	6.9	4.7-9.7	19	9.4	5.5-14.8	14	5.3	2.9-9.1	11	2.4	1.2-4.2	3	1.6	0.3-4.5	8	3.0	1.3-6.1
2009	49	10.0	7.4-13.3	29	13.5	9.0-19.6	20	7.0	4.2-11.0	9	1.8	0.8-3.5	6	2.8	1.0-6.2	3	1.0	0.2-3.1
2010	40	7.6	5.4-10.5	21	8.2	5.0-12.9	19	7.0	4.2-11.0	13	2.7	1.4-4.7	8	3.7	1.5-7.5	5	2.0	0.6-4.7
2011	61	11.8	8.9-15.2	31	14.6	9.8-20.9	30	10.0	6.7-14.5	5	1.0	0.3-2.3	3	1.4	0.3-4.3	2	0.7	0.1-2.7
2012	37	6.9	4.8-9.6	23	10.0	6.2-15.2	14	4.5	2.4-7.8	10	1.8	0.8-3.4	6	2.5	0.9-5.5	4	1.2	0.3-3.4
2013	51	8.8	6.5-11.7	33	13.4	9.1-19.0	18	5.5	3.2-8.9	17	3.1	1.8-5.0	6	2.5	0.9-5.6	11	3.7	1.8-6.8
2014	60	10.9	8.2-14.1	38	15.8	11.0-21.9	22	6.9	4.3-10.6	13	2.6	1.4-4.6	6	2.9	1.0-6.3	7	2.4	0.9-5.1
2015	38	6.9	4.8-9.5	23	9.6	6.0-14.5	15	4.8	2.7-8.1	17	3.2	1.8-5.2	9	4.0	1.8-7.7	8	2.6	1.1-5.4
2016	68	11.6	9.0-14.9	44	16.9	12.1-22.9	24	7.2	4.6-11.0	17	2.7	1.6-4.4	12	4.7	2.3-8.3	5	1.3	0.4-3.4
2017	50	8.6	6.4-11.5	22	9.1	5.6-13.9	28	8.3	5.5-12.2	12	2.1	1.1-3.8	7	2.7	1.1-5.7	5	1.7	0.5-4.2
2018	53	8.5	6.4-11.3	26	9.4	6.1-14.0	27	7.9	5.2-11.8	20	3.2	1.9-5.1	13	4.7	2.4-8.2	7	1.9	0.8-4.3
2019	60	9.8	7.4-12.7	34	12.8	8.8-18.1	26	7.4	4.8-11.1	12	2.0	1.0-3.6	3	1.1	0.2-3.4	9	2.6	1.2-5.1
2020	44	7.7	5.5-10.5	21	8.1	4.9-12.5	23	7.7	4.7-11.8	13	2.1	1.1-3.8	8	2.9	1.2-6.0	5	1.4	0.4-3.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020
Richmond (Staten Island)
Leukemias

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	27	10.9	7.1-16.0	15	16.0	8.4-26.9	12	8.2	4.2-14.4	19	8.1	4.8-12.7	10	12.6	5.8-23.0	9	5.9	2.6-11.2
1977	27	10.7	6.9-15.6	18	18.8	10.6-30.2	9	6.0	2.7-11.3	25	9.4	6.0-14.0	15	14.3	7.7-24.0	10	6.3	3.0-11.6
1978	33	11.5	7.8-16.3	21	16.7	9.9-26.3	12	7.7	3.9-13.5	17	6.5	3.7-10.4	12	10.3	5.0-18.3	5	3.5	1.1-8.0
1979	31	9.9	6.7-14.2	17	10.6	6.1-17.3	14	8.8	4.8-14.7	9	3.2	1.5-6.1	5	3.6	1.1-8.5	4	2.7	0.7-6.9
1980	35	11.3	7.8-15.8	14	11.8	6.1-20.2	21	11.6	7.1-17.8	22	7.4	4.5-11.2	13	12.5	6.1-21.9	9	5.0	2.3-9.6
1981	37	12.7	8.9-17.6	22	19.1	11.6-29.3	15	8.7	4.8-14.4	26	8.6	5.5-12.8	15	12.4	6.5-21.1	11	6.3	3.1-11.3
1982	31	9.8	6.6-13.9	16	11.2	6.2-18.5	15	8.2	4.6-13.6	25	8.3	5.3-12.2	14	11.1	5.9-18.9	11	6.0	3.0-10.8
1983	39	12.8	9.0-17.6	20	15.6	9.1-24.7	19	11.5	6.9-18.0	21	7.2	4.4-11.1	9	8.2	3.4-15.8	12	7.1	3.6-12.3
1984	33	10.9	7.5-15.4	21	18.0	10.7-28.0	12	6.7	3.4-11.7	34	10.9	7.5-15.2	18	13.2	7.6-21.3	16	9.0	5.1-14.7
1985	39	12.9	9.1-17.7	26	24.2	15.2-36.1	13	7.4	3.9-12.6	23	7.8	4.9-11.7	13	13.4	6.9-22.9	10	5.6	2.6-10.3
1986	40	12.2	8.7-16.7	22	15.2	9.3-23.3	18	10.0	5.9-15.8	25	7.7	4.9-11.4	9	5.2	2.4-10.2	16	8.9	5.1-14.6
1987	32	9.4	6.4-13.3	17	10.4	5.9-16.9	15	8.1	4.5-13.4	17	4.8	2.8-7.8	9	5.5	2.4-10.7	8	4.1	1.8-8.2
1988	44	13.5	9.8-18.2	17	12.1	6.9-19.6	27	14.2	9.3-20.6	27	8.5	5.6-12.4	13	10.2	5.1-18.0	14	7.5	4.1-12.6
1989	56	17.3	13.0-22.6	37	28.9	19.6-40.6	19	9.9	5.9-15.4	25	7.5	4.9-11.1	14	10.9	5.7-18.5	11	5.6	2.8-10.1
1990	47	13.5	9.9-18.1	27	16.3	10.6-24.0	20	10.2	6.2-15.7	26	8.1	5.2-11.9	15	11.5	6.1-19.3	11	5.6	2.8-10.1
1991	54	16.1	12.0-21.0	37	26.9	18.5-37.5	17	8.6	5.0-13.8	28	8.4	5.5-12.1	18	15.1	8.8-23.8	10	5.1	2.4-9.4
1992	55	15.4	11.6-20.1	34	22.0	15.0-31.1	21	10.2	6.3-15.6	31	9.0	6.1-12.8	16	11.4	6.3-18.7	15	7.4	4.1-12.2
1993	41	10.9	7.8-14.8	24	15.2	9.5-23.0	17	8.0	4.6-12.8	26	7.0	4.6-10.3	17	11.1	6.3-18.0	9	4.1	1.9-7.8
1994	53	14.1	10.5-18.5	29	18.4	12.1-26.6	24	10.8	6.9-16.1	36	9.8	6.9-13.6	22	14.7	9.1-22.5	14	6.4	3.5-10.8
1995	58	14.8	11.2-19.1	34	19.9	13.7-28.0	24	10.8	6.9-16.1	19	5.2	3.1-8.1	9	5.8	2.6-11.1	10	4.7	2.2-8.6
1996	45	11.4	8.3-15.3	23	12.8	8.0-19.3	22	9.8	6.1-14.9	29	7.6	5.1-10.9	21	12.7	7.8-19.5	8	3.6	1.5-7.1
1997	47	11.7	8.6-15.6	33	20.7	14.0-29.4	14	6.1	3.3-10.2	28	7.1	4.7-10.2	19	11.5	6.8-18.2	9	3.8	1.7-7.3
1998	58	14.0	10.6-18.1	28	15.4	10.2-22.4	30	13.2	8.9-18.8	32	7.9	5.4-11.2	22	12.6	7.8-19.1	10	4.2	2.0-7.8
1999	63	15.0	11.5-19.1	31	18.2	12.1-26.1	32	13.4	9.2-19.0	36	8.6	6.0-11.9	22	12.9	7.9-19.7	14	5.7	3.1-9.6
2000	66	15.4	11.9-19.6	36	18.9	13.1-26.4	30	12.4	8.4-17.8	22	5.2	3.3-7.9	8	4.7	2.0-9.2	14	5.5	3.0-9.3
2001	56	12.9	9.7-16.8	28	15.4	10.1-22.4	28	11.3	7.5-16.4	18	4.2	2.5-6.6	8	4.5	1.9-8.8	10	4.0	1.9-7.5
2002	65	14.5	11.2-18.6	39	19.7	13.9-27.0	26	9.9	6.4-14.6	36	8.0	5.6-11.1	23	11.6	7.3-17.5	13	4.8	2.6-8.3
2003	55	12.2	9.2-16.0	24	11.4	7.3-17.0	31	12.1	8.2-17.3	23	5.2	3.3-7.9	8	4.9	2.1-9.6	15	5.8	3.3-9.7
2004	52	11.7	8.7-15.3	21	11.1	6.8-17.1	31	12.3	8.3-17.5	26	5.9	3.8-8.6	11	5.5	2.7-9.8	15	5.4	3.0-9.0
2005	64	14.1	10.8-18.0	36	17.9	12.4-24.9	28	11.4	7.5-16.5	31	6.8	4.6-9.6	18	9.3	5.4-14.8	13	5.0	2.7-8.7
2006	72	15.6	12.2-19.7	37	18.9	13.2-26.3	35	14.1	9.8-19.7	29	6.1	4.1-8.8	16	8.4	4.7-13.9	13	5.0	2.6-8.6
2007	82	17.0	13.5-21.2	41	19.6	13.9-26.8	41	15.5	11.0-21.1	35	7.8	5.4-10.8	14	7.6	4.1-12.8	21	8.0	4.9-12.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Sex, 1976-2020**

Richmond (Staten Island)

Leukemias

Year of diagnosis/death	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	76	15.9	12.5-20.0	43	21.2	15.2-28.7	33	11.8	8.1-16.7	34	7.2	5.0-10.1	16	8.4	4.7-13.6	18	6.3	3.7-10.1
2009	89	18.1	14.5-22.4	49	22.3	16.3-29.7	40	14.5	10.3-19.9	26	5.6	3.6-8.3	12	6.6	3.3-11.4	14	4.9	2.7-8.4
2010	93	18.4	14.8-22.6	58	26.2	19.7-34.1	35	12.4	8.6-17.4	35	6.9	4.8-9.6	23	11.3	7.0-17.0	12	4.2	2.1-7.5
2011	104	20.2	16.4-24.6	51	22.5	16.6-29.9	53	18.9	14.0-24.9	22	4.5	2.8-6.8	13	6.0	3.1-10.4	9	3.4	1.5-6.6
2012	83	15.9	12.6-19.9	53	23.4	17.4-30.9	30	10.5	7.0-15.2	26	5.4	3.5-8.0	14	7.0	3.8-11.7	12	4.4	2.2-7.8
2013	102	18.8	15.3-23.0	54	22.2	16.5-29.3	48	17.2	12.6-23.1	42	7.6	5.4-10.3	26	10.8	6.9-16.0	16	5.3	3.0-8.8
2014	97	18.9	15.2-23.1	46	19.9	14.4-26.8	51	18.6	13.7-24.6	31	5.8	3.9-8.3	15	6.1	3.3-10.3	16	5.4	3.1-9.0
2015	112	20.6	16.9-25.0	72	29.6	23.0-37.6	40	13.9	9.8-19.2	23	4.3	2.7-6.6	10	5.0	2.4-9.1	13	4.3	2.2-7.6
2016	105	18.9	15.4-23.0	57	22.6	17.0-29.6	48	16.2	11.8-21.7	39	7.3	5.2-10.1	20	9.3	5.6-14.4	19	6.0	3.5-9.7
2017	104	18.0	14.6-22.0	57	22.1	16.6-28.9	47	15.4	11.2-20.8	38	6.7	4.7-9.3	17	7.7	4.4-12.4	21	6.0	3.7-9.4
2018	97	17.4	14.1-21.4	66	25.6	19.6-32.8	31	9.9	6.7-14.3	39	6.6	4.7-9.2	22	8.5	5.3-13.1	17	5.6	3.2-9.2
2019	98	17.0	13.7-20.9	57	21.6	16.3-28.3	41	13.1	9.3-18.1	37	6.4	4.5-8.9	17	6.7	3.9-10.9	20	6.0	3.7-9.5
2020	78	14.5	11.4-18.3	42	16.9	12.0-23.0	36	12.1	8.3-17.1	33	5.3	3.6-7.6	18	6.6	3.9-10.7	15	4.1	2.2-7.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic.

Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.