

**COMMUNICABLE DISEASE IN NEW YORK STATE EXCLUSIVE OF NEW YORK CITY
RATE PER 100,000 OF SELECTED DISEASES
1984 - 1994**

| | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 |
|-------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| AIDS (1) | 1.1 | 0.6 | 5.2 | 4.8 | 8.6 | 10 | 9.3 | 10.1 | 11.9 | 28.4 | 16.4 |
| AMEBIASIS | 0.4 | 0.5 | 0.5 | 0.6 | 0.6 | 0.6 | 0.7 | 0.6 | 1.1 | 1.1 | 1.1 |
| BABESIOSIS (2) | | | 0.2 | 0.1 | 0.1 | 0.2 | 0.1 | 0.3 | 0.3 | 0.3 | 0.3 |
| CAMPYLOBACTERIOSIS (2) | | | 16.3 | 17.2 | 18.3 | 18.1 | 18.1 | 18.8 | 23.7 | 21.9 | 23 |
| CONJUNC NEWBORN | 2.5 | 1.8 | 3.6 | 2.7 | 1.8 | 0.2 | 0.2 | 0.2 | 0.9 | 0.2 | 0.1 |
| CRYPTOSPORIDIOSIS (6) | | | | | | | | | | | 0.3 |
| E. COLI (6) | | | | | | | | | | | 1.4 |
| ENCEPHALITIS VIRAL | 0.3 | 0.4 | 0.3 | 0.5 | 0.3 | 0.3 | 0.4 | 0.3 | 0.2 | 0.3 | 0.3 |
| ENCEPHALITIS OTHER | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0 | 0.1 | 0.1 | 0 |
| FOOD POISONING | 1.4 | 1.4 | 1.6 | 2.5 | 1.6 | 1.1 | 0.1 | 0.4 | 0.1 | 0.2 | 0.5 |
| GIARDIASIS (2) | | | 9.1 | 11.1 | 14.6 | 18.9 | 16.3 | 18.2 | 18.5 | 17.3 | 18.2 |
| GONORRHEA | 181.2 | 174.7 | 182.9 | 167.9 | 168.7 | 164.9 | 150.9 | 136.8 | 111.8 | 100.9 | 106.4 |
| HEMOLYTIC UREMIC SYNDROME (6) | | | | | | | | | | | 0.1 |
| HEMOPHILUS INFLU (2) (4) | | | 2.3 | 2.5 | 2.6 | 2.2 | 1.7 | 0.7 | 0.5 | 0.4 | 0.7 |
| HEPATITIS A | 3.7 | 5.9 | 4.9 | 7.7 | 7.9 | 9.8 | 11.9 | 8.6 | 3.3 | 4.3 | 5 |
| HEPATITIS B | 7.8 | 8.7 | 7.6 | 7.6 | 7.2 | 6.7 | 6.5 | 5.6 | 4.8 | 4.3 | 3.7 |
| HEPATITIS C (3) | | | 0.8 | 0.8 | 0.8 | 0.7 | 0.9 | 2.2 | 1.8 | 2.4 | 2.1 |
| HEPATITIS UNSPECIFIED (3) | 1.6 | 2 | 0.4 | 0.3 | 0.2 | 0.2 | 0.2 | 0.1 | 0.1 | 0 | 0.1 |
| KAWASAKI SYNDROME (2) | | | 0.3 | 0.6 | 0.6 | 0.5 | 0.4 | 0.4 | 0.5 | 0.4 | 0.4 |
| LEGIONELLOSIS (2) | | | 0.8 | 0.6 | 0.8 | 1 | 1.5 | 1.2 | 1 | 0.8 | 0.6 |
| LISTERIOSIS (6) | | | | | | | | | | | 0.4 |
| LYME DISEASE (2) | | | 4 | 7.3 | 22.7 | 30.5 | 12.9* | 35.7* | 31.2* | 25.6* | 47.1* |
| MALARIA | 0.3 | 0.5 | 0.5 | 0.3 | 0.4 | 0.4 | 0.5 | 0.6 | 0.4 | 1.1 | 0.6 |
| MEASLES | 0.5 | 0.8 | 1 | 0.4 | 0.4 | 1.9 | 5.4 | 3.7* | 1* | 0.1* | 0.3* |
| MENINGITIS VIRAL | 4.9 | 6.9 | 5.2 | 6.4 | 3.8 | 5.8 | 5.5 | 13 | 4.6 | 5.3 | 4.3 |
| MENINGOCOCCAL | 1.4 | 1.5 | 1.3 | 1.3 | 1.4 | 1.3 | 1.3 | 1.1 | 1.1 | 1.2 | 1 |
| MENINGITIS OTHER | 3.9 | 4.5 | 2.4 | 2.5 | 2.5 | 3 | 2.8 | 2.9 | 2.7 | 2.8 | 1.4 |
| PNEUMOCOCCAL MENINGITIS (6) | | | | | | | | | | | 0.9 |
| MUMPS | 1 | 1.9 | 0.8 | 1.3 | 1 | 1.7 | 1.5 | 1 | 0.8 | 0.4 | 0.3 |
| PERTUSSIS | 1 | 1.3 | 1.5 | 1.7 | 2.3 | 1.7 | 3.2 | 1.7 | 1.6 | 3.5 | 2.3 |
| RMSF (5) | 0.1 | 0.1 | 0.2 | 0.1 | 0.1 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| RUBELLA | 1 | 0.2 | 0.2 | 0.1 | 0 | 0.1 | 0.2* | 5.1* | 0.1* | 0.2* | 0.1* |
| SALMONELLOSIS | 14.5 | 18.8 | 19.3 | 19.2 | 20.2 | 20.5 | 19 | 20 | 18.7 | 22.1 | 18.2 |
| SHIGELLOSIS | 2.9 | 4.4 | 2.6 | 9.3 | 4.6 | 8.5 | 9.9 | 7.1 | 4.2 | 7.3 | 10.3 |
| SYPHILIS EARLY | 5.8 | 6.2 | 4.6 | 5.9 | 11.3 | 18.6 | 21.2 | 14.9 | 10.2 | 7.1 | 4.6 |
| TOXIC SHOCK SYNDROME (2) | | | 0.2 | 0.2 | 0.2 | 0.1 | 0.1 | 0.2 | 0.1 | 0.2 | 0.2 |
| TUBERCULOSIS | 5.9 | 6.1 | 5.8 | 5.8 | 6.5 | 6.2 | 6.1* | 7* | 7.1* | 6.6* | 6* |
| TYPHOID FEVER | 0.1 | 0.1 | 0.1 | 0.1 | 0.2 | 0.5 | 0.2 | 0.3 | 0.2 | 0.2 | 0.1 |
| YERSINIOSIS (2) | | | 0.8 | 0.9 | 1.2 | 1.2 | 1.1 | 1.2 | 1.1 | 1.1 | 1.3 |

(1) Excludes prison inmates reported 1984 through 1994

(2) First made reportable in 1986

(3) Unspecified Hepatitis split into (I) Hepatitis C and (II) Unspecified in 1986

(4) Haemophilus Influenzae Type B Invasive Disease

(5) Rocky Mountain Spotted Fever

(6) Reportable as of 3/94 for Upstate New York

* Confirmed