

**COMMUNICABLE DISEASE IN NEW YORK CITY  
RATE PER 100,000 OF SELECTED DISEASES BY DIAGNOSIS DATE  
1987 - 1996**

	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
AIDS	71.2	87.3	91.8	101.8	118.7	139.4	159.4	131.6	89.5	139.9
AMEBIASIS	2.8	1.4	3.6	5.6	6.0	6.4	9.7	9.6	13.1	11.6
BABESIOSIS	0.0	0.1	0.0	0.0	0.1	0.0	0.0	0.1	0.1	0.1
CAMPYLOBACTERIOSIS	3.7	4.2	9.1	10.8	12.3	15.9	17.6	18.3	17	16.7
ENCEPHALITIS OTHER	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.2	0.1	0.0
GIARDIASIS	2.9	4.0	6.8	11.1	12.9	14.9	26.4	33.5	33.4	30.6
GONORRHEA	918.2	743.9	555.5	481.2	395.3	296.5	252.3	262.8	219.8	173.8
HAEMOPHILUS INFLUENZA	1.1	0.8	2.0	1.7	0.8	0.5	0.5	0.5	0.5	0.8
HEPATITIS A	2.4	5.1	6.9	10.8	17.5	12.1	14.0	12.9	13.5	8.1
HEPATITIS B	16.7	18.0	19.4	9.2	6.0	6.0	6.4	7.4	7.0	6.6
KAWASAKI SYNDROME	0.3	0.1	0.2	0.2	0.2	0.2	0.1	0.1	0.1	0.1
LEGIONELLOSIS (2)	0.4	0.7	0.9	1.3	1.1	0.1	0.1	0.2	0.1	0.3
LYME DISEASE	1.5	2.2	1.5	1.3	1.9	1.4	2.9	1.3	6.1	5.4
MALARIA	0.4	1.3	1.4	1.6	1.8	2.3	2.0	1.4	3.0	3.6
MEASLES	6.5	0.8	1.9	15.1	26.1	0.9	0.3	0.2	0.1	0.1
MENINGITIS VIRAL	2.6	2.0	2.7	2.3	5.8	2.4	3.7	2.0	2.1	1.7
MENINGOCOCCAL (1)	0.8	0.9	1.2	1.0	0.4	0.4	0.5	0.5	0.7	0.7
MENINGITIS OTHER	4.1	1.8	4.4	2.7	2.7	2.0	2.7	2.8	3.4	3.1
MUMPS	0.6	1.4	0.3	0.9	0.2	0.2	0.0	0.2	0.2	0.2
PERTUSSIS	0.2	0.2	0.3	0.3	0.3	0.3	1.6	3.0	0.9	0.8
RMSF	0.1	0.0	0.0	0.1	0.0	0.1	0.0	0.0	0.1	0.3
RUBELLA	0.0	0.1	0.2	0.1	0.0	0	0.3	0.0	0.1	0.1
SALMONELLOSIS (3)	24.9	23.2	27.7	23.0	23.3	25.0	27.6	25.9	29.1	25.7
SHIGELLOSIS	24.4	6.6	4.9	9.2	5.8	10.3	9.2	13.7	11.4	8.4
SYPHILIS EARLY	111.0	131.4	140.0	160.4	135.2	104.1	66.6	40.7	31.0	16.2
TOXIC SHOCK SYNDROME	0.1	0.1	0.1	0.1	0.0	0	0.0	0	0.1	0.1
TUBERCULOSIS (4)	30.3	31.9	34.9	48.1	50.2	52	44.2	40.9	33.4	27.5
TYPHOID FEVER	0.4	0.8	0.8	1.1	1.1	0.7	1.4	1.1	0.9	0.9
YERSINIOSIS	0.2	0.7	0.6	0.5	0.6	0.7	0.6	0.7	0.5	0.6

(1) Case definition require positive culture from blood, spinal, or other sterile sites

(2) Beginning 1991 only confirmed cases reported

(3) Salmonellosis paratyphi A and B plus Salmonellosis nontyphoidal

(4) Revised in 1978 to include persons with verified disease in past or lost to supervision for more than 12 months and had disease again